Research article

Chromatin profiling identifies transcriptional readthrough as a conserved mechanism for

piRNA cluster biogenesis in mosquitoes

Jieqiong Qu[1], Valerie Betting[1], Ruben van Iterson[1], Florence M. Kwaschik[1], Ronald P. van Rij[1*]

[1] Department of Medical Microbiology, Radboud Institute for Molecular Life Sciences, Radboud University Medical

Center, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands

Jieqiong Qu: Jieqiong.Qu@radboudumc.nl

Valerie Betting: Valerie.Betting@radboudumc.nl

Ruben van Iterson: Ruben.vanIterson@radboudumc.nl

Florence Kwaschik: kwaschik.florence@virology.uzh.ch

[*] **Corresponding author:**

Ronald van Rij: Ronald.vanRij@radboudumc.nl

Abstract

Background

The PIWI-interacting RNA (piRNA) pathway in culicine mosquitoes differs substantially from other model organisms in terms of an expanded PIWI gene family and ubiquitously expressed piRNAs in somatic tissues.

Results

Here, through comprehensive analyses of endogenous piRNAs in *Aedes aegypti*, we defined core piRNA clusters as small RNA source loci that showed ubiquitous expression in both somatic and germline tissues. These core piRNA clusters produced piRNAs from a single genomic strand and were enriched for non-retroviral endogenous viral elements (nrEVEs) in antisense orientation. Biogenesis of cluster-derived piRNAs depended on key factors, the mRNA exporter Nxf1, three Tudor proteins, Veneno, Tejas, and Yb, as well as the evolutionarily conserved cochaperone Shutdown. Combined transcriptional and chromatin state analyses showed that these core piRNA clusters do not have independent promoters, but instead depended on readthrough transcription from upstream genes for piRNA biogenesis. This biogenesis mechanism was evolutionarily conserved in other vector mosquito species, *Aedes albopictus*, *Culex quinquefasciatus*, and *Anopheles gambiae*. Interestingly, the upstream genes of many core piRNA clusters of the two *Aedes* mosquitoes were orthologous, but the downstream piRNA clusters differed both in size and sequence, especially evident by distinct nrEVE insertions from different virus families. We propose a model in which a conserved set of genes determine the loci for piRNA biogenesis, while the downstream piRNA clusters function as traps for nrEVEs, allowing adaptation to environmental challenges such as virus infection.

Conclusions

Our systematic analyses of piRNA cluster expression and chromatin state lay a solid foundation for future studies of piRNA and genome evolution in these important vector species.

Background

Aedes mosquitoes transmit arthropod-borne (arbo) viruses including dengue, Zika, and chikungunya virus [1], posing significant health and economic impact [2-4]. A female mosquito acquires arboviruses through an infectious blood meal, after which viruses establish a systemic infection. Albeit permissive to systemic infection, mosquitoes have evolved efficient antiviral strategies to limit viral replication to non-pathogenic levels [5, 6], such as the small interfering (si)RNA pathway [7, 8]. More recent studies have revealed that another class of small RNAs, PIWI-interacting RNAs (piRNAs), play a role in antiviral defense as well [9-13].

The piRNA pathway is evolutionarily conserved in animals but has been most extensively studied in the ovary of *Drosophila melanogaster*. In fruit flies, the majority of piRNAs originate from discrete genomic loci, termed piRNA clusters, that are especially rich in transposable elements (TEs) [14]. The resulting piRNAs silence TEs at the transcriptional and post-transcriptional levels to maintain genome integrity [15-17]. Two types of piRNA clusters exist in *D. melanogaster*: uni-strand and dual-strand clusters, which produce piRNAs from one or both genomic strands, respectively [14]. In somatic follicle cells, piRNAs are exclusively produced from uni-strand piRNA clusters, such as the *flamenco* (*flam*) locus [14, 18, 19]. Similar to protein-coding genes, *flam* is transcribed by RNA polymerase II and the resulting transcripts are regularly spliced, capped, and polyadenylated. The piRNA precursor transcript is subsequently exported by Nxf1/Nxt1 to Yb-bodies where primary piRNA biogenesis occurs [17, 19-21]. In germline cells, piRNAs are predominantly produced from dual-strand piRNA clusters that are marked by the repressive histone mark H3K9me3. Dual-strand clusters require a heterochromatin-dependent transcription machinery involving the Rhino-Deadlock-Cutoff (RDC) complex and Moonshiner for transcription while inhibiting splicing, capping, and polyadenylation [22-29]. A specialized nuclear export complex, consisting of an ovary-specific heterodimeric mRNA export receptor Nxf3-Nxt1, Bootlegger, and the TREX component UAP56, is recruited for export of the piRNA cluster transcript to the nuage where piRNA biogenesis factors accumulate [24, 30-33]. In the nuage, piRNA biogenesis occurs in a feed-forward loop, called ping-pong amplification, involving two PIWI proteins, Aubergine and Ago3 [14-16, 34].

While hallmark features such as the ping-pong amplification are conserved [35-39], the piRNA pathway in *Aedes* mosquitoes differs significantly from *D. melanogaster* in terms of an expanded PIWI gene family and ubiquitously expressed piRNAs in somatic tissues [40-44]. The majority of piRNAs in *D. melanogaster* are generated from TE-rich regions despite an overall low genomic TE content [14, 45, 46], whereas it is the opposite for *Aedes* mosquitoes [47]. Moreover, the mosquito piRNA pathway processes a broader repertoire of substrates including transcripts from

protein-coding genes and satellite repeats as well as cytoplasmic viral RNA, and, thus, has important functions beyond

transposon silencing [10, 35, 36, 38, 39, 48-50].

The genomes of main arbovirus vectors, *Aedes aegypti* and *Aedes albopictus*, contain sequences of non-retroviral RNA

viruses (non-retroviral endogenous elements, nrEVEs) from several families including the *Flaviviridae* and

Rhabdoviridae families. Importantly, piRNA clusters of *Aedes* mosquitoes are enriched for nrEVE sequences [12, 51-

56], suggesting that they may serve as the source of immunological memory against exogenous viruses. Indeed, a

recent study showed that piRNAs produced from a naturally occurring nrEVE protected the *Ae. aegypti* host against

its cognate virus, thus demonstrating a functional link between nrEVEs and antiviral immunity [13].

Given the rapid evolution of the piRNA pathway and the prominent differences between flies and mosquitoes [57, 58],

we explored the molecular mechanisms underlying piRNA cluster biogenesis in *Aedes* mosquitoes. We combined

transcriptome and chromatin state analyses in *Ae. aegypti* and uncovered readthrough transcription as a major

mechanism for piRNA cluster biogenesis. Further comparative analyses between two *Aedes* mosquito species revealed

an intriguing model for piRNA cluster evolution in which a conserved set of genes determine the loci for piRNA

biogenesis, while piRNA clusters function as traps for nrEVEs, allowing species-specific adaptation to viral infection

challenges.

Results

Systematic analyses of piRNA clusters in *Ae. aegypti*

As in most arthropods, the piRNA pathway of the vector mosquito *Ae. aegypti* is active in both somatic and germline

tissues [40-42, 59]. However, an important question is whether piRNA expression differs across cell lines and tissues

in *Ae. aegypti*. To answer this question, we collected piRNA sequencing data from two cell lines (Aag2 [50] and

CCL125 [42]), four somatic tissues (fat body [60], head [61], midgut [62], and thorax [59]), as well as two germline

tissues (ovary [59] and testes), and compared their piRNA cluster expression in a systematic manner (Fig. 1a).

Fig. 1 Systematic annotation of piRNA clusters in *Ae. aegypti* mosquitoes.

a. Comparison of piRNA expression from piRNA clusters between cell lines, somatic and germline tissues of *Ae. aegypti*. piRNA clusters were independently annotated in each dataset and represented as dots, with color representing the origin and dot size representing cluster size. The numbers of clusters are listed. Log scaled piRNA read counts were used to represent piRNA expression levels within a piRNA cluster. For detailed information, see Supplementary Table 1.
b. Cumulative distribution of piRNA cluster coverage on the plus strand of the *Ae. aegypti* genome. Colors are as in **a**.

c. Representative genome browser tracks showing examples of a core piRNA cluster and a germline-specific piRNA cluster. piRNA-
sized reads (25 to 30 nt) from small RNA-seq were mapped and normalized to RPKM (y-axis), with F/R indicating forward/reverse
strand. The boundaries of each cluster are indicated with dashed vertical lines. The boundaries of core piRNA clusters are based on
Aag2 piRNA clusters, and germline-specific clusters are based on ovary piRNA clusters.
d. Enrichment of the indicated repeat classes in core and germline-specific piRNA clusters, relative to the entire genome.
e. Distribution of 25 core piRNA clusters in cell lines and tissues. Within each dataset, piRNA clusters are ranked according to log
scaled cluster size (bp). The 25 core piRNA clusters are shown as dark grey bars; other piRNA clusters are shown as light grey bars.
piRNA density (RPKM) was plotted using the same colors as in **a**.
f. piRNA length distribution, PIWI protein association [39], 1U/10A bias, and the genomic distribution of the 25 core piRNA
clusters, which are ranked according to the cluster size in Aag2 cells.
118

We first annotated piRNA clusters in each cell line and tissue, using two independent pipelines (Fig. S1) [55, 56, 63].

We observed largely comparable piRNA expression between cell lines and somatic tissues, whereas the majority of

piRNA clusters in germline tissues showed tissue-dominant expression, which, however, correlated well between

ovary and testes (Fig. 1a and Fig. S1c). Uni-strand clusters were more often observed in cell lines and somatic tissues,

whereas dual-strand clusters were predominant in germline tissues (Fig. 1b). About 90% of the clusters from cell lines

and somatic tissues are uni-strand, whereas 45% and 28% of the clusters are uni-strand in ovary and testes, respectively

(Fig. S1d). Given these clear differences, we defined two classes of piRNA clusters based on their expression pattern:

25 core piRNA clusters that are universally expressed across cell lines and tissues, and 228 germline-specific piRNA

clusters that are uniquely expressed in germline tissues, ovary and testes (Fig. 1c).

Unlike piRNA clusters in *D. melanogaster* that are highly enriched for TEs [14], the 25 core piRNA clusters were

depleted of TEs except for a minor enrichment of LTR-retrotransposons, while germline-specific piRNA clusters were

enriched for Rolling-Circle (RC) transposons when compared to the whole genome (Fig. 1d). We also explored the

relationship between piRNA cluster size and piRNA density but found no correlation (Fig. 1e and Fig. S1e). Yet, the

25 core piRNA clusters produced 68% of all cluster-derived piRNAs in Aag2 cells, mainly because they are among

the largest piRNA clusters, with the largest cluster spanning ~180 kb (Fig. 1e).

We next characterized the 25 core piRNA clusters in terms of the piRNA length distribution, PIWI-protein association,

nucleotide bias, and their genomic distribution (Fig. 1f). We found that most mapped small RNAs were 25-30 nt in

length and were almost exclusively associated with Piwi5, except for cluster 23 and cluster 25 showing a strong

association with Piwi4. Moreover, core cluster-derived piRNAs showed a strong 1U but not 10A bias, suggesting that

the core piRNA clusters produce Piwi5-associated primary piRNAs. Remarkably, only three core piRNA clusters are

embedded within expressed genes in the same orientation, which raises an interesting question of how the remaining

majority of core piRNA clusters that are located in intergenic regions are generated (Fig. 1f).

Comprehensive chromatin profiles of the *Ae. aegypti* genome

The intriguing observation that most core piRNA clusters are located in intergenic regions and give rise to Piwi5-associated primary piRNAs prompted us to test whether these clusters are similar to *flam*[14, 18, 19, 64]. To gain precise insights into transcriptional regulation and chromatin states, we performed ChIP-seq of four histone marks in Aag2 cells, including H3K27ac, associated with active enhancers and promoters; H3K4me3, associated with active promoters; H3K36me3, associated with actively transcribed regions; and H3K9me3, associated with heterochromatin regions (Fig. 2a) [65-69]. For all four histone marks, two biological replicates showed strong correlations (Pearson's r > 0.98), indicating the high reproducibility of these datasets (Fig. S2b). Using ChromHMM[70], we annotated five chromatin states based on combinations of four histone marks to define heterochromatin, promoter, enhancer domains, as well as regions associated with active transcription and regions without discernible signals (Fig. 2a).

To validate the annotated chromatin states, we performed ATAC-seq to assay open chromatin regions that are biologically active, such as promoters and enhancers [71] (Fig. 2b). Distinct chromatin accessibility profiles were evident among chromatin states: the promoter and enhancer states showed the highest average levels of chromatin accessibility, followed by regions flanking the transcription state, while the heterochromatin state was relatively inaccessible, in agreement with previous observations in ENCODE datasets [72]. We next assessed the current genome annotation with chromatin states (Fig. 2c). In total, 17.4 % of the *Ae. aegypti* genome was assigned with one of the four chromatin states, with most of them located in intron and intergenic regions (Fig. S2c). 5'UTRs were enriched for promoter states, while exon and 3'UTR sequences were enriched for the transcription state, and intergenic regions for heterochromatin (Fig. 2c).

In addition to histone modifications, eukaryotic transcription units are characterized by conserved 5′ to 3′ profiles of specific RNA polymerase II (pol II) Carboxy-Terminal Domain (CTD) phospho-isoforms [73]. Phosphorylation of CTD on Ser5 is normally higher at 5′ ends of actively transcribed genes where it facilitates mRNA capping [74], whereas Ser2 phosphorylation is higher at 3′ ends where it is important for mRNA 3′ end formation [75]. We, therefore, performed a series of RNA pol II ChIP-seq experiments, using antibodies either recognizing both phospho-isoforms or one of the two phosphor-isoforms (Fig. S2d). We observed that actively expressed genes showed enrichment of active histone marks as well as RNA pol II occupancy at promoter regions, *e.g. LAP* and *eEF1a1*, whereas no signals were found for silenced genes, *e.g. AAEL020926* (Fig. 2d). Besides, this dataset provides evidence of differentially

expressed isoforms (*Vasa* and *Piwi6*) by marking the corresponding active promoter region. Overall, we observed

transcription-dependent signals of active histone marks H3K4me3, H3K27ac, and H3K36me3, as well as RNA pol II,

starting sharp at transcription start sites (TSS) and ending at transcription end sites (TES) (Fig. 2e), which has been

reported in other organisms [76-79]. Our genome-wide maps of chromatin accessibility, histone modifications, and

RNA pol II occupancy, provide a comprehensive functional annotation of the *Ae. aegypti* genome which serves as a

useful resource to understand piRNA cluster biogenesis.

Fig. 2 Chromatin profiling in *Ae. aegypti* Aag2 cells.

a. Chromatin state annotation using ChIP-seq data of four histone marks with ChromHMM [70]. Each chromatin state (row) is defined by a combinatorial pattern of specific histone marks (columns). For instance, the promoter state is distinguished by enrichment of H3K4me3 while the enhancer state is distinguished by enrichment of H3K27ac. A five-state ChromHMM was chosen because it represented biologically interpretable patterns that were not repetitive or ambiguous. Six-state and seven-state models are shown in Fig. S2a.
b. Average chromatin accessibility marked by ATAC-seq signals at each chromatin state from **a**.
c. Chromatin state coverage within each genomic feature. When a genomic region was annotated with several chromatin states, only the dominant chromatin state (> 50% coverage of the whole region) is shown.
d. Representative genome browser tracks showing chromatin profiles at expressed genes (*LAP* and *eEF1a1*), a silent gene (*AAEL020926*), genes with different isoforms (*Vasa* and *Piwi6*), as well as genes that are not annotated as such in the reference genome (downstream region of *LAP*). Chromatin state colors are as in **a**. ChIP-seq and RNA-seq signals were normalized to RPKM (y-axis).
e. Band plots (upper panel) and heatmaps (lower panel) showing average ChIP-seq signals of active histone marks, H3K4me3, H3K27ac, and H3K36me3, as well as RNA pol II, RNA pol II ser5p, and RNA pol II ser2p. Published RNA-seq datasets were used to define gene expression quartiles (Q1-Q4) and non-expressed genes (silent) (Fig. S2e) [39]. All genes were scaled to 5kb. TSS, transcription start site; TES, transcription end site. ChIP-seq signals were normalized to RPKM (y-axis).

Readthrough transcription of core piRNA clusters from upstream genes

In contrast to protein-coding genes, we did not detect signals of active histone marks and RNA pol II occupancy at the start of core piRNA clusters (Fig. 3a). Instead, active histone marks and RNA pol II signals were detected already in the upstream regions, suggesting that piRNA cluster transcription is initiated from upstream regions and that core piRNA clusters lack their own independent promoters. Besides, we did not detect enrichment of H3K9me3 signals which mark dual-strand clusters in *Drosophila* [24], neither for core piRNA clusters nor for other piRNA clusters in Aag2 cells (Fig. S3a). Only a few piRNA clusters showed H3K9me3 signals, which, however, coincided with ChIP input signals, and are likely due to the background artifacts from genomic DNA [80]. Together, our analyses revealed that most core piRNA clusters are located in intergenic regions and likely depend on readthrough transcription.

To test this hypothesis, we quantified the distance (d) between the 5' end of core piRNA clusters to the TES of its nearest gene expressed in the same orientation (Fig. 3b). Compared with scrambled controls, d of 25 core piRNA clusters was significantly shorter. In fact, most core piRNA clusters start directly after the TES of their corresponding upstream genes ($d = 0$), with the exceptions of cluster 24 and cluster 25, which are distant from any expressed genes, which we will refer to as 'stand-alone' piRNA clusters (Fig. S3b). We noticed that cluster 2 and cluster 20 were mapped to the same upstream gene, *AAEL000053* (Fig. 3b). After careful inspection, we propose that these two clusters are part of a large single element, > 220 kb in size, which was annotated as two separate clusters due to a large gap in small RNA coverage (Fig. S3b). Moreover, we experimentally validated readthrough transcription using PCR primers

210   spanning the upstream gene and piRNA clusters using cDNA as a template (Fig. 3b), whereas properly terminated

211   genes failed to amplify any products (*e.g.*, *AAEL001359* and *AAEL024736*).

Fig. 3 Confirmation of readthrough transcription

a. Average ChIP-seq signals of active histone marks, H3K4me3, H3K27ac, and H3K36me3, as well as RNA pol II in 25 core
piRNA clusters. All clusters were scaled to 5kb, with dashed lines indicating the 5' ends of the clusters. Scrambled control regions
were generated by shuffling piRNA clusters through the whole *Ae. aegypti* genome with bedtools shuffle [81].
b. The distance (*d*) between the 5' end of core piRNA clusters to TES of its nearest expressed gene in the same orientation was
quantified. Scrambled control regions were generated by shuffling piRNA clusters through the whole genome with bedtools shuffle
[81]. Statistical differences were examined by unpaired t-test. Lower panel, PCR was performed to analyze readthrough transcription
from the upstream gene into the piRNA cluster, using genomic DNA, cDNA, or mRNA as templates. piRNA clusters were ranked
according to the cluster size. For detailed primer information sees Supplementary Table 4.

c. Band plots of average ChIP-seq signals (RPKM) of four histone marks, H3K4me3, H3K27ac, H3K36me3, H3K9me3 (left), and
RNA pol II (right) of 19 upstream genes of readthrough piRNA clusters. All gene body regions were scaled to 5kb.
d. Representative genome browser tracks showing readthrough core piRNA clusters. ATAC-seq, ChIP-seq, and RNA-seq signals,
as well as piRNA-sized reads from small RNA-seq, were normalized to RPKM. The boundaries of each cluster are indicated with
dashed lines and the direction of piRNA clusters is indicated with arrows, with upstream genes labeled with gene names.

We then analyzed chromatin profiles of upstream genes, which showed canonical features of actively transcribed genes,

with the enrichment of active histone marks as well as RNA pol II ChIP-seq signals at the TSS (Fig. 3c). Expression

of several upstream genes, *AAEL006523*, *AAEL005080*, and *AAEL004586*, was confirmed by a recently published

proteomic study with > 98% hit identity [82], suggesting these genes are not only transcribed but also translated into

proteins. Hence, it is likely that these upstream genes can both deliver properly terminated gene products as well as

readthrough transcripts that are substrates for piRNA biogenesis. Notably, most readthrough core piRNA clusters

showed higher RNA-seq signals at the end of clusters (Fig. 3d), which was also true for genic piRNA clusters (Fig.

S3b). This observation is likely due to the oligo (dT)-primed RNA-seq library preparation, which in turn suggests that

these piRNA cluster transcripts are polyadenylated. Together, our analyses indicate that most core piRNA cluster

transcripts are produced by readthrough transcription from upstream genes.

We further dissected the connection between the upstream genes and the downstream piRNA production. Our previous

analyses indicated that most core piRNA clusters produce Piwi5-bound primary piRNAs (Fig. 1f). We, therefore, used

Piwi5 knockdown as a positive control to establish Stem-Loop (SL)-RT-qPCR assays to quantify the piRNA

abundance from readthrough core piRNA clusters (Fig. 4a, top panels). Using four piRNAs from two clusters (clusters

7 and 10) as readouts, we analyzed whether silencing of the upstream gene would affect its downstream piRNA

production. Upon knockdown of the upstream gene of cluster 7, *AAEL010451*, we did not observe an effect on piRNA

expression, probably due to the relatively low knockdown efficiency (Fig. 4a, middle panels). We also tried another

set of dsRNA targeting *AAEL010451* and did not achieve a better knockdown (Fig. S4a). Importantly, upon knockdown

of the upstream gene of cluster 10, *AAEL023582*, we observed a significant reduction in piRNA production from its

downstream piRNA cluster (Fig. 4a, lower panels). This effect was specific, as *AAEL023582* knockdown did not affect

the expression of piRNAs from cluster 7. These results provide a direct genetic link between upstream gene

transcription and downstream piRNA cluster expression.

Key factors of piRNA cluster biogenesis

To define factors involved in the biogenesis of core cluster-derived piRNAs, we analyzed our previously generated

small RNA-seq data from Aag2 cell upon knockdown of PIWI genes [35], genes encoding PIWI interactors [36, 83],

Veneno, *Pasilla*, *Atari*, as well as *Zucchini* (Zuc), involved in piRNA 3'end formation [37]. Among them, we observed

a clear reduction in the expression of cluster-derived piRNAs with knockdown of *Piwi5* and *Veneno*, but not with the

others (Fig. 4b). As most core piRNA clusters are *Piwi5*-dependent (Fig. 1f), we further tested three Tudor proteins

that were among the top enriched hits from a Piwi5-IP experiment [83], Vreteno (*AAEL020359*), Tejas (*AAEL022773*),

and Yb (*AAEL001939*). We also included Shutdown (*AAEL011632*), which is required for the biogenesis of all piRNA

populations in *D. melanogaster* [84]. With small RNA-seq, we observed a consistent downregulation of core piRNA

clusters upon knockdown of *Yb* and *Shutdown*; in comparison, the effect of Tejas was minor and Vreteno barely

affected piRNA expression from these core piRNA clusters (Fig. 4c).

Fig. 4 Key determinants of piRNA cluster biogenesis

a. Knockdown of upstream genes from two readthrough core piRNA clusters. *AAEL010451* is the upstream gene of cluster 7 and
AAEL023582 is the upstream gene of cluster 10. RT-qPCR was performed to confirm gene knockdown, while SL-RT-qPCR was
performed to quantify piRNA expression. All experiments were performed in biological triplicates (shown as dots) unless otherwise
specified. The interquartile range and the median are shown in box plots. Statistical differences were examined by unpaired t-test,
when compared to the non-targeting control knockdown of luciferase (dsLuc). Symbols denote statistical significance (* $p < 0.05$,
** $p < 0.01$, *** $p < 0.001$). For detailed primer information, see Supplementary Table 4.
b. Changes in piRNA levels from core piRNA clusters upon knockdown of published piRNA biogenesis factors, quantified from
small RNA-seq data and compared to the non-targeting control knockdown of luciferase (dsLuc). piRNAs were first normalized to
per million mapped miRNAs and then log scaled for comparison. The Pearson correlation coefficient is shown in the upper left
corner.
c. Changes in piRNA levels from core piRNA clusters upon knockdown of newly Piwi5 co-factors. Knockdown of target genes was
confirmed with RT-qPCR (Fig. S4b).
d. Changes in piRNA levels from core piRNA clusters upon knockdown of newly screened piRNA biogenesis factors. Knockdown
of target genes was confirmed with RT-qPCR (Fig. S4b).
e. A schematic model of piRNA biogenesis from core piRNA clusters in *Ae. aegypti*.
278

To search for key factors that are upstream in the piRNA pathway, we performed a large-scale RNAi screen, including

orthologs of known piRNA biogenesis factors in other organisms, chromatin modifiers, 16 genes of mRNA 3'end

processing, 18 genes that mediate RNA export, 26 genes of general RNA processing, as well as 29 RNA helicases (Fig.

S4). With the screen, we found that *Nxf1* knockdown significantly downregulated one of the two tested piRNAs (Fig.

S4e). We then confirmed the effect of *Nxf1* knockdown using another set of dsRNA and observed a significant

reduction of two piRNAs from two piRNA clusters. Next, we performed small RNA-seq for a more comprehensive

evaluation of all core piRNA clusters (Fig. 4d). In addition, we included UAP56 which plays an important role in

piRNA cluster export in *D. melanogaster* [30]. The effect of *Nxf1* knockdown seemed to be more heterogeneous,

however, not correlated to whether the cluster is readthrough, genic or stand-alone; whereas, knockdown of *UAP56*

barely affected piRNA expression from these core piRNA clusters, in line with results from the SL-RT-qPCR screen

(Fig S4e).

Together, these data suggest the following model for cluster-derived piRNA biogenesis: 1) piRNA cluster transcripts

are produced by readthrough transcription from an upstream coding gene; 2) key factors involved in processing and

maturation include the mRNA exporter Nxf1, three Tudor proteins, Veneno, Tejas, and Yb, the evolutionarily

conserved cochaperone Shutdown, as well as the PIWI protein Piwi5; 3) the resulting mature piRNAs are mainly

associated Piwi5 and show a strong 1U bias (Fig. 4e).

Conserved readthrough transcription for piRNA cluster biogenesis among mosquitoes

To analyze whether readthrough transcription is a shared mechanism for cluster piRNA biogenesis, we extended our

analyses to other mosquito species, including the closely related *Aedes albopictus*, and the more distant *Culex*

quinquefasciatus and *Anopheles gambiae*, (Fig. 5a).

Fig. 5 Readthrough transcription is an evolutionarily conserved mechanism for piRNA cluster biogenesis in mosquitoes

301 **a**. Phylogeny and divergence time estimation by molecular clock analysis of four mosquito genomes [85], with the genome assembly
302 (Vectorbase release 56) and the corresponding genome size annotated.
303 **b-d**. Analyses of piRNA clusters in *Ae. albopictus* (**b**), *Cx. quinquefasciatus* (**c**), and *An. gambiae* (**d**), including enrichment of
304 repeat classes in piRNA clusters (i), the distance (*d*) as illustrated in Fig. 3b (ii), and representative readthrough core piRNA clusters
305 (iii).
306

With the same strategy used for *Ae. aegypti* (Fig. 1), we annotated piRNA clusters from cell lines and tissues from

these mosquito species (Fig. S5-7). Like in *Ae. aegypti*, most piRNA clusters were uni-strand in cell lines and somatic

tissues, whereas in germline tissues piRNA clusters are more dual-stranded. About 70% of the clusters from cell lines

and somatic tissues were uni-strand, whereas the percentage was below 52% for germline tissues in all three species.

Next, we defined core piRNA clusters in *Ae. albopictus* (n=37), in *Cu. quinquefasciatus* (n=32), and in *An. gambiae*

(n=19), based on their ubiquitous expression across different cell lines and tissues. In parallel, we defined germline-

specific piRNA clusters in *Ae. albopictus* (n=244), in *Cu. quinquefasciatus* (n=587), and in *An. gambiae* (n=539)

(Supplementary Tables 5-7).

Analogous to *Ae. aegypti*, core piRNA clusters in these three species were only modestly enriched for LTR

retrotransposons (Fig. 5b-d), whereas germline-specific piRNA clusters in *Ae. albopictus* and *Cx. quinquefasciatus*

mainly showed slight enrichment of LTR retrotransposons and RC transposons (Fig. 5b and c). In striking contrast,

piRNA clusters in *An. gambiae* showed an overall enrichment of all TE classes, except for SINEs (Fig. 5d). Core

piRNA clusters in *Ae. albopictus* were also among the top-sized piRNA clusters (Fig. S5d), whereas core piRNA

clusters in *Cx. quinquefasciatus* and *An. gambiae* were more dispersed in size (Fig. S6d and S7d).

We next examined whether core piRNA clusters follow the pattern of readthrough transcription, using d (illustrated in

Fig. 3b) as an indicator. Compared to scrambled controls, d of core piRNA clusters was significantly shorter in all

three species (Fig. 5b-d). To define whether a core piRNA cluster follows readthrough transcription, we set a cutoff of

d based on the d in *Ae. aegypti* (5 kb), scaled by the genome size of each mosquito species (Fig. 5a) to account for the

higher probability to classify a piRNA cluster as a readthrough cluster in smaller genomes ($d < 10$ kb for *Ae. albopictus*;

2.5 kb for *Cx. quinquefasciatus*, and 1.0 kb for *An. gambiae*). In summary, 14 of 37 core piRNA clusters in *Ae.*

albopictus (Fig. S5e), 14 of 32 core piRNA clusters in *Cx. quinquefasciatus* (Fig. S6e), and 10 of 19 core piRNA

clusters in *An. gambiae* (Fig. S7e), fulfilled the criteria and are thus likely transcribed by readthrough transcription

from upstream genes. Indeed, many of these core piRNA clusters started directly after the upstream genes (Fig. 5b-d).

In conclusion, these analyses suggest that readthrough transcription is a common mechanism for piRNA cluster

biogenesis shared among these four vector mosquitoes.

A conserved pool of genes drives piRNA cluster expression in *Aedes* mosquitoes

Our observation that readthrough transcription is a conserved mechanism for piRNA biogenesis raises the question of

whether specific genes are selected to drive cluster expression. For the two *Aedes* mosquitoes, the expression of

upstream genes was significantly lower when compared to all expressed genes, whereas the gene size seemed to be

larger, however, the difference was not significant (Fig. 6a). In contrast, the upstream genes of *Cx. quinquefasciatus*

and *An. gambiae* were expressed at higher levels. With Gene Ontology (GO) analysis, the upstream genes in *Cx.*

quinquefasciatus were significantly enriched for genes encoding ribosomal proteins, whereas no GO term enrichment

was found in any other species.

Notably, many upstream genes are orthologs between *Ae. aegypti* and *Ae. albopictus*, which is also true for a few genic

piRNA clusters (Fig. 6b). This allowed us to define homologous piRNA clusters and compare their sequence

composition in a pair-wise manner (Fig. 6c). Despite the highly conserved upstream genes (cDNA identity > 75%),

the downstream piRNA clusters differed both in size and sequence identity. The latter was especially evident with

varied numbers of nrEVE insertions from different virus families into orthologous clusters (Fig. 6b and c, summarized

in Supplementary Table 8). Gene duplication is a common phenomenon in the genome of *Ae. albopictus* Foshan strain

[85] and we observed an interesting example where in *Ae. aegypti*, *AAEL005473* was associated with a genic piRNA

cluster (cluster 21) and depleted of nrEVE insertions, whereas its two orthologous genes in *Ae. albopictus* were

associated with a readthrough piRNA cluster (cluster 15) and a genic piRNA cluster (cluster 20), both with nrEVE

insertions (Fig. S8b).

Several studies have shown that piRNA clusters are enriched for nrEVEs in *Aedes* mosquitoes [51, 52, 55, 56, 86]. We

revisited the annotated nrEVEs in *Ae. aegypti* [56] and found that 133 of 252 nrEVEs (53%) are located within 215

piRNA clusters of Aag2 cells, and 126 of these 133 nrEVEs (95%) are located within the 25 core piRNA clusters (X^2

test, $p < 0.0001$) (Fig. 6b). The same is true for *Ae. albopictus* [55], in which 119 of 456 nrEVEs (26%) are located

within 285 piRNA clusters, and among them, 82 of these 119 nrEVEs (69%) are located within the 37 core piRNA

clusters (X^2 test, $p < 0.0001$). Besides, the vast majority of nrEVEs were inserted in antisense orientation to core piRNA

clusters in both species and thus serve as a pool of primary piRNAs that can potentially target cognate virus sequences

(Fig. 6d and e). In *Ae. albopictus*, there are two exceptions where nrEVEs insertions were in the same orientation as

core piRNA clusters, such as the nrEVE derived from a member of the *Rhabdoviridae* within cluster 20 (Fig. S8b), the

resulting piRNAs were sense to piRNA clusters, yet also showed a strong 1U bias (Fig. 6e).

Fig. 6 Systematic comparison of core piRNA clusters between mosquitoes

a. Expression level and the gene body length of upstream genes of core piRNA clusters (Core) compared to all expressed genes (Exp) in each species. Expressed genes were determined for each species (Fig. S8a). Both the distribution of the raw data and the box plot displaying the interquartile range and the median are shown. Statistical differences were examined by unpaired t-test (* p < 0.05, *** p < 0.001, **** p < 0.0001).

b. Comparison between core piRNA clusters in two *Aedes* mosquitoes. In each species, piRNA clusters were ranked according to their size and colored based on their genomic origin, with another column showing the number of nrEVE insertions. Ortholog genes were connected. 'Unknown' refers to the situation when a piRNA cluster is located in a scaffold without any annotated gene nearby.

c. Representative genome browser tracks showing two pairs of core piRNA clusters with the upstream genes being orthologs in two *Aedes* mosquitoes. The boundaries of each cluster are indicated with dashed vertical lines and the direction of the piRNA cluster is indicated with an arrow on top. nrEVEs were shown as arrows, with colors representing viral families.

d. Read size distribution and the nucleotide bias of piRNAs mapped to nrEVEs within 25 core piRNA clusters of *Ae. aegypti*.

e. Read size distribution and the nucleotide bias of piRNAs mapped to nrEVEs within 37 core piRNA clusters of *Ae. albopictus*.

375 Together, these analyses revealed that a conserved pool of genes was selected to define the genomic loci for piRNA

376 cluster biogenesis in *Aedes* mosquitoes; however, the sequence composition of piRNA clusters is not conserved and

377 core clusters have evolved as 'traps' for nrEVE insertions in a species-specific manner. Although the mechanism of

378 readthrough transcription is conserved, the identity of upstream genes is not conserved in evolutionary more distant

379 mosquito species.

Core piRNA clusters function as traps for nrEVEs

We further explored why nrEVEs are preferentially inserted into core piRNA clusters in *Aedes* mosquitoes. As nrEVE

insertions and piRNA cluster evolution are highly intertwined processes in both *Aedes* mosquitoes, we included *Cx.*

quinquefasciatus and *An. gambiae* in this analysis, because their genomes contain few nrEVE insertions and, therefore,

their core piRNA clusters likely represent the naïve state of core piRNA clusters devoid of nrEVE insertions [86]. We

compared core piRNA clusters with the remaining piRNA clusters in terms of cluster size and piRNA density. We

found core piRNA clusters are larger in cell lines and tissues of all four species, except for two germline tissues of *An.*

gambiae (Fig. 7a). The differences in cluster size are more drastic in the two *Aedes* mosquitoes than in *Cx.*

quinquefasciatus and *An. gambiae*. We observed a general pattern of higher piRNA density in core piRNA clusters of

all four species. Thus, a possible scenario is that core piRNA clusters are intrinsically larger in size and high in piRNA

density which favors insertions of nrEVEs; as a consequence, nrEVE insertions enlarge core piRNA clusters without

adversely affecting piRNA production and drive core piRNA cluster expansion.

The acquisition and loss of nrEVEs from the host genome are highly dynamic and ongoing processes. Recent studies

identified nrEVEs that were absent from the reference genome but present in wild-collected mosquitoes from different

geographical populations [13, 56]. Among these newly identified nrEVEs, three nrEVEs have definite chromosomal

integration sites, one of which was previously been shown to be in a piRNA cluster [56], which is one of the core

piRNA clusters (cluster 6, Fig. 7b). With the limited number of defined integration sites for new nrEVEs, it is

impossible to perform statistics, but it is worth noting that the 25 core piRNA clusters in total span ~1.9 Mb, occupying

only 0.15% of the *Ae. aegypti* genome. We, therefore, propose that the insertion and maintenance nrEVEs in core

piRNA clusters are unlikely to be stochastic, but rather a result of positive selection.

Discussion

Here, we comprehensively analyzed piRNA expression in four mosquito species and uncovered readthrough

transcription as a conserved mechanism for cluster piRNA biogenesis (Fig. 7c). Within *Aedes* mosquitoes, the upstream

genes are relatively lowly expressed, but evolutionary conserved to drive piRNA cluster transcription. The downstream

piRNA clusters are unidirectional, ubiquitously expressed, and function as traps for nrEVEs in a species-specific

manner. The piRNA cluster transcripts are processed by key factors, including the RNA export factor Nxf1, three

Tudor proteins, Veneno, Tejas, and Yb, as well as the cochaperone Shutdown. Eventually, mature piRNAs antisense

to nrEVEs are loaded into Piwi5 and hold the potential to target cognate viral mRNA. Our study proposes a model for

piRNA cluster biogenesis in important vector mosquitoes and provides insights into the adaptive evolution of mosquito

genomes via the endogenization of viral sequences.

Characterization of mosquito piRNA clusters

We defined two different categories of piRNA clusters: core piRNA clusters that are shared between somatic and

germline tissues; and germline-specific piRNA clusters that show restricted expression only in germline tissues. Core

piRNA clusters are predominantly uni-strand clusters and produce Piwi5-bound primary piRNAs independent of the

ping-pong amplification loop, thus resembling the *flam* locus in *D. melanogaster* [14, 18]. However, there are essential

differences. First, *flam* serves as a piRNA cluster only in somatic follicular cells of *Drosophila* ovaries [14], while in

somatic tissues outside of the ovary *flam* acts as a source of endogenous siRNAs [87]. In contrast, core piRNA clusters

in mosquitoes show ubiquitous expression in both somatic and germline tissues. We cannot exclude that the observed

expression of core piRNA clusters in germline tissues might result from the residential somatic cell populations, rather

than from the actual germline cells, which deserves further investigation. Second, unlike *flam*, which is transcribed by

RNA pol II from its own promoter similar to protein-coding genes [19], core piRNA clusters do not have their own

promoters, but depend on readthrough transcription from their upstream genes (Fig. 3). Third, *flam* is enriched for

gypsy family LTR retrotransposons, whereas core piRNA clusters are enriched for nrEVEs in *Aedes* mosquitoes (Fig.

6) [55, 56]. Thus, *flam* represses transposons by preventing the expression of *gypsy* family TEs having complementary

sequences in this locus, while core piRNA clusters with nrEVEs trapped in the antisense orientation encode a pool of

piRNAs that have the potential to target their cognate viruses. A recent study provided direct evidence of nrEVEs

mediated antiviral immunity by showing that a naturally occurring nrEVE produces piRNAs to protect the host against

its cognate virus, the cell-fusing agent virus [13]. Third, the enrichment of the *gypsy* TE family is conserved in *flam* in

three closely related species, *D. melanogaster*, *D. yakuba*, and *D. erecta* [18]. Moreover, *flam* acquires new TEs via

horizontal transfer, with related elements showing higher sequence identity between *D. simulans*, *D. sechellia*, *D.*

yakuba, and *D. erecta* than expected if TEs would be acquired through vertical transmission [18, 88]. In contrast,

nrEVE insertions vary both in numbers and virus families between *Ae. aegypti* and *Ae. albopictus* (Fig. 6). Therefore,

it is likely that core piRNA clusters acquire new nrEVEs via direct endogenization of viral sequences, showing

adaption to local viral challenges.

Germline-specific piRNA clusters produce piRNAs from both genomic strands and show restricted expression only in

germline tissues. Hence, it is logical to hypothesize that germline-specific piRNA clusters in mosquitoes may resemble

dual-strand piRNA clusters expressed in the ovary of *D. melanogaster* that are marked by heterochromatic H3K9me3

[22, 24, 25, 28]. However, no orthologs have been found in mosquitoes for those key factors that promote dual-stranded

piRNA cluster transcription, such as the RDC complex [26] and Moonshiner [28]. Besides, there is a clear depletion

of most TEs in germline-specific piRNA clusters in mosquitoes with the exception of *An. gambiae* (Fig. 5d). Therefore,

chromatin profiles in germline tissues are needed to study the regulation of germline-specific piRNA clusters. Recent

studies in *D. melanogaster* male germline revealed a sex-specific piRNA program [89, 90], which is different from

our observations in mosquitoes where piRNA clusters correlated well between ovaries and testes, which was especially

obvious in *An. gambiae* (Fig. S7a). Future studies of germline-specific piRNA clusters are required to gain more

insights into the involved key factors and the underlying mechanism of cluster transcription.

In addition to core piRNA clusters and germline-specific piRNA clusters, we observed an interesting tissue-specific

expression pattern that fewer piRNA clusters are expressed in the midgut. Such restricted piRNA expression in the

midgut from our analysis is supported by a recent study of dengue virus-infected *Ae. albopictus*, in which it was shown

that piRNAs were reduced in the midgut [91]. Interestingly, another recent study showed that the siRNA pathway fails

to efficiently silence dengue virus replication in the midgut of *Ae. aegypti* [62]. As the midgut is one of the main tissue

barriers for arboviruses to cross to establish systematic infection [92], it will be interesting to investigate the interplay

between these two small RNA pathways in the midgut and to address their role in antiviral immunity.

Readthrough transcription as a major mechanism for piRNA biogenesis in mosquitoes

Through combined transcriptome and chromatin state analyses of the *Ae. aegypti* genome, we uncovered readthrough

transcription as a major mechanism for piRNA cluster biogenesis. Knockdown of one of the upstream genes reduced

the corresponding downstream piRNA production. Moreover, many upstream genes are actually orthologs between

the two *Aedes* mosquitoes; therefore, readthrough transcription from this subset of genes is definitely not a stochastic

event, but rather a tightly regulated and evolutionarily conserved process.

Two connected questions remain to be answered: how these upstream genes are regulated to allow readthrough

transcription and how are readthrough transcripts funneled into the piRNA pathway. In germline cells of *D.*

melanogaster, the dual-stranded piRNA cluster *hsp70* locus has been proposed to follow readthrough transcription,

however, a specialized RDC complex is engaged to inhibit termination by recruiting the cleavage and polyadenylation

specificity factor (CPSF) complex in a chromatin-dependent but sequence-independent manner [24, 26]. We screened

16 genes that are involved in canonical mRNA 3'end processing including all key components of the CPSF complex

and did not observe a clear effect. Moreover, core piRNA clusters in mosquitoes are not marked by any repressive

histone mark, thus devoid of recognizable chromatin features. Therefore, it is likely that a specialized non-canonical

3'end processing pathway is recruited in a sequence-dependent manner for readthrough transcription of piRNA cluster

transcripts.

After being transcribed, the *flam* piRNA precursor is spliced, capped, polyadenylated, and exported by Nxf1/Nxt1 to

Yb-bodies where primary piRNA biogenesis occurs [17, 19-21]. We screened 18 genes of the RNA export pathway

and identified Nxf1, but not Nxt1, as an important factor for piRNA clusters. This is rather unexpected because the

heterodimeric mRNA export receptor Nxf1-Nxt1 is conserved from budding yeast to humans to mediate mRNA

translocation through nuclear pore complexes into the cytoplasm [93]. Instead, this may indicate that a specialized

non-canonical mRNA export pathway is employed to convey these readthrough transcripts to the piRNA pathway,

which involves Nxf1 but not its canonical co-factors, such as Nxt1. So far, it is not clear at which stage piRNA cluster

precursors are funneled into the piRNA biogenesis pathway. It is possible that, directly after readthrough transcription,

piRNA cluster precursors are uncoupled from the upstream gene transcript and are separately exported by Nxf1.

Another possibility is that readthrough transcripts are exported as a whole, and afterwards, the piRNA cluster

precursors are recognized by the piRNA pathway for further processing.

Our analyses identified three Tudor proteins, Veneno, Tejas, Yb, and an evolutionarily conserved cochaperone

Shutdown, as key factors for piRNA cluster biogenesis. Among them, Tejas and Yb are Piwi5 interactors, Veneno and

Shutdown are Ago3 interactors, whereas Veneno also interacts with Yb which in turn interacts with Piwi5 [36, 83].

Such a Yb-centered protein network suggests that Yb bodies are the place for primary piRNA biogenesis in mosquitoes.

However, Vreteno, another Tudor protein, did not affect piRNA production, although it is an important component of

Yb bodies in *D. melanogaster* [94]. Therefore, further characterization is needed to dissect the protein composition of

Yb bodies in *Aedes* mosquitoes.

Core piRNA clusters function as traps for nrEVEs

Our analysis revealed that core piRNA clusters function as traps for nrEVEs in the two *Aedes* mosquitoes and that they

are evolving in a species-specific manner. The different numbers and identities of acquired nrEVEs likely reflect their

exposure to virus challenges, which is supported by the finding of wild-caught mosquito populations from epidemic

regions showing a CFAV-derived nrEVE trapped in the core piRNA cluster 6 [56]. Importantly, a recent study provided

direct evidence of nrEVEs mediated antiviral immunity by showing that a naturally occurring CFAV-derived nrEVE

produces piRNAs to protect the host against its cognate virus [13]. Thus, nrEVEs that have been trapped in core piRNA

clusters can be regarded as a sequence-specific record of virus exposure that may function in concert with the siRNA

machinery which is a broadly active antiviral response.

Core piRNA clusters are intrinsically larger in size and high in piRNA density which, somehow, favors insertions of

nrEVEs. Such direct integration, transcription, and production of nrEVE-derived piRNAs constitutes a mechanism of

adaptive genome evolution of *Aedes* mosquitoes. Therefore, one possibility is to artificially mimic this heritable

immune memory, for example, by inserting sequences of pathogenic arboviruses such as Zika and dengue virus into

core piRNA clusters to generate mosquitoes resistant to infection as an intervention strategy.

Conclusion

Our study provides a systematic evaluation of piRNA profiles in four mosquito species and uncovered a conserved

readthrough transcription mechanism for piRNA biogenesis. We propose a model for piRNA cluster evolution and

highlight the function of piRNA clusters as traps for nrEVEs to shape the antiviral immunity of *Aedes* mosquitoes. Our

study lays a solid foundation for future piRNA studies to understand how the piRNA pathway affects the vector

competence of mosquito populations via nrEVE insertions, and how nrEVE-derived piRNAs influence the evolution

of viruses. Moreover, our genome-wide maps of chromatin state, genome accessibility, and pol II occupancies will be

a useful resource for understanding gene regulation in the important vector mosquito *Ae. aegypti*.

Acknowledgements

521   We thank Chet H. Loh, Rebecca Halbach, Pascal Miesen, Ezgi Taşköprü, and members of the laboratory for fruitful

522   discussion. This work is supported by a VICI grant from the Dutch Research Council (grant number 016.VICI.170.090).

Contributions

525   J.Q. and R.P.v.R designed the experiments and analyzed the data. J.Q. performed computational analyses and most of

526   the experiments, with the help from V.B. and R.v.I. for the RNAi screen and F.M.K. for PCR validation of readthrough

527   transcription. J.Q. and R.P.v.R. wrote the paper. All authors read and contributed to the manuscript.

Competing interests

530   The authors declare no competing interests.

Methods

piRNA cluster annotation

The genome assemblies of four mosquito species, *Ae. aegypti* LVP, *Ae. albopictus* Foshan FPA, *Cx. quinquefasciatus*

Johannesburg, *An. gambiae* PEST, were downloaded from Vectorbase (release 56). Publicly available small RNA-seq

datasets were collected for piRNA cluster annotation. For *Ae. aegypti*, Aag2 cell line (SRR2633298, SRR2633457,

SRR2633591, SRR3157917) [50], CCL125 cell line (SRR11252286) [42], fat body (SRR5163617, SRR5163618,

SRR5163619) [60], head (SRR950413, SRR950414) [61], midgut (SRR7430939) [62], thorax (SRR5961503,

SRR5961505) [59], Ovary (SRR5961504, SRR5961506) [59], and testes (SRR11300467). For *Ae. albopictus*, U4.4

cell line (SRR2182528) [95], whole mosquitoes (SRR2182437, SRR2182496) [95], female mosquito carcasses

(SRR11095782, SRR11095783), ovary (SRR11095792, SRR11095793), and testes (SRR11252300) [42]. For *Cx.*

quinquefasciatus, Hsu cell line (SRR11252314) [42], whole mosquitoes (SRR5457467) [96], salivary glands

(SRR8936292, SRR8936293, SRR8936298) [97], midgut (SRR8936296, SRR8936297, SRR8936299) [97], ovary

(SRR11252316) [42], and testes (SRR11252317) [42]. For *An. gambiae*, Mos55 cell line (SRR11252306) [42], head

and thorax (SRR11713504, SRR11713505) [98], midgut (SRR11713500, SRR11713501, SRR11713502) [98], fat

body (SRR11713503, SRR11713506, SRR11713507) [98], ovary (SRR11713497, SRR11713498, SRR11713499)

[98], and testes (SRR11252311) [42]. Different dataset from the same cell line or tissue were merged and used as a

single library for downstream analysis.

After trimming of adaptors with Cutadapt [99], multi-mapping reads from the small RNA-seq libraries were randomly

distributed over all possible mapping positions (--best –strata -M1) with Bowtie [100]. Annotation of piRNA clusters

was performed as described [55, 56], with several parameters optimized (Fig. S1a): 1) the minimum amount of piRNA-

sized reads (23-33 nt) mapped to per 5-kb non-overlapping window, when normalized per million mapped piRNAs

(ppm); 2) the uniquely mapped piRNA per cluster. A minimum of 20 ppm per 5-kb window and a minimum of 3

uniquely mapped piRNAs per cluster were used to annotate piRNA clusters from each cell line or tissue, separately.

This piRNA cluster annotation pipeline was solely guided by piRNA coverage of the respective genomic regions, and

did not include assumptions on nucleotide biases or strand asymmetry. In parallel, proTRAC was used to confirm

piRNA cluster annotation, which requires strand asymmetry and enrichment for ping-pong signature (1U/10A) (Fig.

S1b) [63].

Classification of core piRNA clusters and germline-specific piRNA clusters

Core piRNA clusters were defined as piRNA clusters that are universally expressed across cell lines and tissues. Within

each mosquito species, we annotated piRNA clusters individually from each cell line/ tissue, and piRNA clusters that

overlapped between at least six datasets are considered as core piRNA clusters. Bedtools intersect was used to retrieve

overlapping piRNA clusters, and the boundaries of core piRNA clusters were based on cell line piRNA clusters.

Besides, core piRNA clusters were also supported by clusters predicted by proTRAC [63], exception *Cx.*

Quinquefasciatus, due to the highly fragmented piRNA clusters predicted by proTRAC.

Germline-specific piRNA clusters were defined as piRNA clusters that are only expressed in germline tissues, ovary,

and testes, but not in any other somatic tissues. Bedtools intersect was used to get overlapped piRNA clusters between

ovary and testes, and with the option -v to remove piRNA clusters that overlapped with any cell line or somatic tissues

[81]. The boundaries of germline-specific piRNA clusters were based on ovary piRNA clusters. Detailed information

of piRNA clusters of *Ae. aegypti* was summarized in Supplementary Table 1, *Ae. albopictus* in Supplementary Table

5, *Cx. quinquefasciatus* in Supplementary Table 6, and *An. gambiae* in Supplementary Table 7.

Characterization of piRNA clusters

piRNA cluster expression level was determined by mapping piRNA-sized reads (23-33 nt) to each piRNA cluster,

using bamCoverage from deepTools [101]. Pair-wise comparison of piRNA cluster expression was performed with

log10 transformed piRNA cluster expression level using the R package ggplot2 [102]. Genome browser tracks of small

RNA-seq datasets were generated with pyGenomeTracks from deepTools [101].

For transposable elements analyses, the repeat annotation files were downloaded for *Ae. aegypti* L5 (Vectorbase,

Aedes-aegypti-LVP_AGWG_REPEATFEATURES_AaegL5.gff3), *Ae. albopictus* (NCBI, GCF_006496715.1), *Cx.*

quinquefasciatus (Vectorbase, Culex-quinquefasciatus-Johannesburg_REPEATFEATURES_CpipJ2.gff3), and for *An.*

gambiae (Vectorbase, Anopheles-gambiae-PEST_REPEATFEATURES_AgamP4.gff3). Bedtools were used to exact

TEs that overlapped with either core piRNA clusters or germline-specific piRNA clusters.

Based on genomic distribution, a piRNA cluster is defined as a genic piRNA cluster, if the piRNA cluster is 80%

covered in length by an expressed gene in the same orientation. Besides, a piRNA cluster is defined as a readthrough

piRNA cluster, if d is below the threshold. We set a cutoff of d based on the genome size of each mosquito species, 5

kb for *Ae. aegypti*, 10 kb for *Ae. albopictus*, 2.5 kb for *Cx. quinquefasciatus*, and 1.0 kb for *An. gambiae*.

ChIP-seq and analysis pipeline

Aag2 cells were grown in Leibovitz's L-15 medium (Invitrogen) supplemented with 10% fetal bovine serum (Gibco),

50 U/ml Penicillin, 50 μg/mL Streptomycin (Invitrogen), 1x Non-essential Amino Acids (Invitrogen) and 2% Tryptose

phosphate broth solution (Sigma) at 28°C in a humidified atmosphere without CO_2. Cell lines were maintained by

splitting twice weekly according to confluency.

Aag2 cells were cross-linked with 1% formaldehyde for 10 min at room temperature with gentle shaking. Cross-linking

was quenched with the addition of 1/10 volume 1.25 M glycine. Cells were washed with 1× PBS, then harvested by

scraping in Buffer B (20 mM HEPES, 0.25% Triton X-100, 10 mM EDTA, and 0.5 mM EGTA). Cells were then

pelleted by spinning at $600 \times g$ for 5 min at 4 °C. Cell pellet was resuspended in Buffer C (150 mM NaCl, 50 mM

HEPES, 1 mM EDTA, and 0.5 mM EGTA) and rotated for 10 min at 4 °C. Cells were again pelleted by spinning at

$600 \times g$ for 5 min at 4 °C. The cell pellet was resuspended in 1× incubation buffer (0.15% SDS, 1% Triton X-100,

150 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, and 20 mM HEPES) at 15 million cells/mL. Cells were sheared with a

Bioruptor Pico sonicator (Diagenode) at 4 °C using 6 cycles of 30 s on, 30 s off. Sonicated chromatin was spun at

$18,000 \times g$ for 5 min at 4 °C. Chromatin in the upper supernatant was aliquoted, immediately frozen in liquid nitrogen,

and stored at -80°C.

Sonicated chromatin from 2 million cells was used for each ChIP experiment. Chromatin was incubated in 1×

incubation buffer (0.15% SDS, 1% Triton X-100, 150 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, and 20 mM HEPES)

supplemented with 1× protease inhibitors, 0.1% BSA, and a 50:50 mix of Protein A and G Dynabeads (Invitrogen),

with agitation overnight at 4 °C. Generally, 1 µg antibody was used for each ChIP (H3K4me3, Diagenode, C15410003;

H3K27ac, Diagenode, C15410196; H3K36me3, Diagenode, C15410058; H3K9me3, Diagenode, C15410193; RNA

pol II, Santa cruz, sc-56767; RNA pol II ser5p, Abcam, ab5131; RNA pol II ser2p, Chromotek, 3E10). As a control,

no antibody was added to the ChIP input sample (sonicated chromatin). The beads were washed twice with wash buffer

1 (0.1% SDS, 0.1% sodium deoxycholate, 1% Triton X-100, 150 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, and 20

mM HEPES), once with wash buffer 2 (wash buffer 1 with 500 mM NaCl), once with wash buffer 3 (250 mM LiCl,

0.5% sodium deoxycholate, 0.5% NP-50, 1 mM EDTA, 0.5 mM EGTA, and 20 mM HEPES), and twice with wash

buffer 4 (1 mM EDTA, 0.5 mM EGTA, and 20 mM HEPES). After the washing steps, beads were rotated for 20 min

at room temperature in elution buffer (1% SDS and 0.1 M NaHCO$_3$). The supernatant was de-cross-linked with 200

mM NaCl and 100 µg/mL proteinase K for 4 h at 65 °C. De-cross-linked DNA was purified with MinElute PCR

Purification columns (Qiagen). DNA concentrations were determined by Qubit fluorometric quantification

(ThermoFisher Scientific). As a quality check step, qPCR analysis of positive control and negative control regions

using 10× diluted ChIP DNA was performed with the GoTaq qPCR system (Promega) according to the manufacturers'

recommendations. Detailed primer information of positive and negative controls is list in Supplementary Table 2.

Libraries were prepared with the Kapa Hyper Prep Kit for Illumina sequencing (Kapa Biosystems) according to the

manufacturer's protocol with the following modifications. 3 ng DNA was used as input, with NEXTflex adapters (Bioo

Scientific) and ten cycles of PCR amplification. Post-amplification cleanup and size selection was performed with dual

AMPure XP beads selection (0.6× and 0.8×). Size-selected samples were analyzed for purity with a High Sensitivity

DNA Chip on a Bioanalyzer 2100 system (Agilent), with an average size between 300 and 500 bp. Samples were

sequenced in a paired-ended manner using the NextSeq 500 (Illumina) according to standard Illumina protocols.

After an initial quality check with FastQC [103], sequencing reads were aligned to the *Ae. aegypti* reference genome

(*Ae. aegypti* LVP, VectorBase release 56) using BWA [104]. Duplicated reads were removed for further analysis.

Mapped reads from two biological duplicates were merged and used for peak calling, because of good correlation

(Pearson's r > 0.98). Peak calling was performed with the MACS2 against an input sample with standard settings and

a q value of 0.05 [105], with the option of broad peak calling for H3K36me3 and H3K9me3. The peak files from four

histone marks (H3K4me3, H3K27ac, H3K36me3, and H3K9me3, detailed in Supplementary Table 2) were used as

input for chromatin state analysis using ChromHMM [70]. The five-emission state model was chosen, as it represented

biologically interpretable combinatorial patterns that were not repetitive or ambiguous and showed general agreement

with previously published models [72, 106, 107] (Fig. S2a). The deepTools package was used to generate bigwig files

that are normalized to RPKM [101], and example track profiles were plotted with fluff [108]. Band plots and heatmaps

were generated with computeMatrix and plotProfile from deepTools [101].

ATAC-seq and analysis pipeline

ATAC libraries were prepared essentially according to a protocol for mammalian cells with minor changes [109]. In

brief, Aag2 cells were trypsinized on plates and resuspended into single cells. Cells were pelleted by spinning at $500 \times g$

for 5 min at 4°C and then washed twice with ice-cold $1\times$ PBS buffer. 50,000 cells were counted and suspended in ice-

cold $1\times$ PBS. The same volume of $2\times$ lysis buffer (20 mM Tris-HCl, 20 mM NaCl, 30 mM $MgCl_2$, 0.5% IGEPAL)

was added to 50,000 cells and pipetted up and down 20 times to lyse the cells. Cells were pelleted at $500 \times g$ for 30 min

at 4°C and suspended in 25 µL of digestion buffer (10 mM Tris pH 7.6, 5 mM $MgCl_2$, 10% DMF, 1U Tn5 transposase),

followed by tagmentation in the 37°C incubator for 1 h at 650 rpm. After tagmentation, 9 µL of clean-up buffer (0.5

M NaCl, 150 mM EDTA, 1% SDS, 2 mg/mL Protease K) was added for incubation at 40°C for 30 min with 650 rpm

shaking. Normal phase $2\times$ AMPure XP beads (68 µL) purification was performed to remove small fragments (<100

bp). Purified DNA was used for a first PCR amplification (Nextera Index Kit) with KAPA HiFi HotStart ReadyMix

(Roche) for seven cycles. Reverse phase $0.55\times$ SPRI bead (27.5 µL) purification was performed to remove large

fragments (>300 bp), followed by purification with QIAquick PCR Purification Kit (Qiagen). Purified DNA was used

for a second PCR amplification (Nextera Index Kit) with KAPA HiFi HotStart ReadyMix (Roche) for another seven

cycles. The PCR product was purified again with QIAquick PCR Purification Kit (Qiagen). DNA amounts were

determined with Qubit fluorometric quantification (ThermoFisher Scientific) and analyzed for fragment size

distribution with a DNA 1000 Chip on a Bioanalyzer 2100 system (Agilent).

Samples were sequenced in a paired-ended manner using the NextSeq 500 (Illumina) according to standard Illumina

protocols. After an initial quality check with FastQC [103], sequencing reads were aligned to *Ae. aegypti* genome

assembly (*Ae. aegypti* LVP, VectorBase release 56) using BWA [104]. Potential PCR duplicates were removed using

the Picard MarkDuplicates option. The filtered BAM files were used as input in MACS2 for peak calling with standard

settings and a q value of 0.05 [105] (Supplementary Table 2). The deepTools package was used to generate, shift

(alignmentSieve function), and normalize bigwig track files to RPKM [101], and example track profiles were plotted

with fluff [108]. Band plots were generated with computeMatrix and plotProfile from deepTools [101].

Small RNA-seq and analysis pipeline

Small RNA deep sequencing libraries were generated using the NEBNext Small RNA Library Prep Set for Illumina

(E7560, New England Biolabs), using 1 µg RNA as input. Library preparation was performed following the

manufacturer's instructions except for 3' adapter ligation, which was performed for 18 h at 16°C to enhance the ligation

efficiency of 2'-O-methylated small RNAs. Libraries were sequenced on an Illumina Hiseq 4000 machine by the

GenomEast Platform (Strasbourg, France).

After an initial quality check with FastQC [103], raw sequence reads were first trimmed with Cutadapt [99] and then

mapped with Bowtie [100]. Reads in the size range from 21 to 24 bp were mapped to the pre-miRNA sequences of *Ae.*

aegypti from miRBase database and annotated as miRNA [110]. Reads in the size range from 25 to 30 bp were regarded

as piRNAs. For PIWI-protein IP followed by small RNA-seq (PRJNA594491) [39], piRNA read counts within a

certain piRNA cluster were normalized to the corresponding IP input sample for enrichment. For analyzing the changes

in piRNA expression after knockdown of genes of interest [35-37, 83], piRNA reads counts were compared to the

luciferase control knockdown after normalization to the total mapped miRNAs. Published small RNA-seq dataset used

in this analysis include: knockdown of PIWI proteins (PRJNA266885) [35], Veneno (SRR6395463- SRR6395468)

[36], Pasilla and Atari (PRJNA664195) [83], and Zuc (SRR12231665, SRR12231683, SRR12231685, SRR12231686,

SRR12231693, SRR12231694) [37].

To analyze piRNAs mapped to nrEVEs, the latest nrEVEs annotations of *Ae. aegypti* [56] and *Ae. albopictus* [55] were

used. The length distribution was plotted in R based on the mapped reads ranging from 23 to 33 nt. For the analysis of

nucleotide biases, piRNA reads were first separated based on the strand with SAMtools [111] and then trimmed to 20

nt for sequence logo representation with WebLogo [112]. Example tracks with nrEVE annotations were generated

using pyGenomeTracks from the deepTools package[101].

RNA-seq analysis pipeline

Publicly available RNA-seq datasets were used to determine expressed genes in each mosquito species, including *Ae.*

aegypti (SRR7938851, SRR7938854, SRR7938865) [39], *Ae. albopictus* (SRR7907921, SRR7907923, SRR7907924),

Cx. quinquefasciatus (SRR16931550, SRR16931551, SRR16931552), and for *An. gambiae* (SRR1525402,

SRR1525403, SRR6127580, SRR6127581 [113]). After initial quality control by FastQC [103], raw sequence reads

were aligned to the corresponding genome, *Ae. aegypti* LVP, *Ae. albopictus* Foshan FPA, *Cx. quinquefasciatus*

Johannesburg, *An. gambiae* PEST (Vectorbase, release 56) using STAR [114] with default settings. Quantification of

gene expression (FPKM) was performed with Cuffdiff [115]. Expressed genes were determined based on the

distribution of FPKM per library (detailed in Supplementary Table 3).

mRNA purification

Two rounds of mRNA purification were performed with NEB magnetic d(T)25 beads (S1419S, New England Biolabs),

followed by DNaseI (Ambion)-treatment and reverse transcription, before cDNA PCR. PCR primers were listed in

Supplementary Table 4.

Stem-loop qPCR

Stem-loop qPCR (SL-qPCR) was performed essentially according to a protocol for miRNA quantification with minor

changes [116]. A total of 100 ng of RNA was reverse transcribed using Stem-loop reverse transcription primers

(Supplementary Table 4) and SuperScript II Reverse Transcriptase mix (Invitrogen). Stem-loop RT reaction conditions

were as follows: 30 min at 16°C, 30 min at 42°C, and 5 min at 85°C. SYBR-green qPCR was performed as following:

initial denaturation step for 5 min at 95°C, followed by 40 cycles of denaturation for 10s at 95°C, annealing for 20s at

60°C, and extension for 10s at 72°C. piRNA gene expression levels were normalized to the expression of the

housekeeping miRNA Bantam and fold changes were calculated using the 2(-ΔΔCt) method [117]. All SL-qPCR

primers were listed in Supplementary Table 4.

RNAi screen in Aag2 cells

PCR products flanked by T7 promoter sequences at both ends were *in vitro* transcribed by T7 polymerase to produce

dsRNA. Aag2 cells were seeded at a density of $1.5x10^5$ cells/well in a 24-well plate, 24 h before the dsRNA transfection

in triplicates. For each target gene, a transfection mix containing 300 μl non-supplemented L-15 medium, 500 ng

dsRNA, and 1.3 μl X-treme GENE HP DNA transfection reagent (Sigma) was prepared according to the

manufacturer's instructions. Per 24-well, 100 μl of the transfection mix was added in a dropwise manner. Three days

post-transfection, Aag2 cells were lysed in 1 mL RNA-Solv reagent (Omega Bio-tek), followed by RNA extraction

through phase separation and isopropanol precipitation. RNA integrity was evaluated on EtBr stained agarose gels,

and RNA concentration was determined using the Nanodrop ND-1000.

For RT-qPCR analyses, DNaseI (Ambion)-treated RNA was reverse transcribed with random hexamers using the

Taqman reverse transcriptase kit (Life Technologies) and PCR amplified using the GoTaq qPCR system (Promega) on

a LightCycler 480 (Roche). In RNAi experiments, expression levels of target genes were normalized to the expression

of the housekeeping gene lysosomal aspartic protease (*LAP*, *AAEL006169*), and fold changes were calculated using

the 2-ΔΔCT method [117]. All primers were listed in Supplementary Table 4.

Data availability

All generated data were deposited in the GEO database (GSE210270).

References

1.   Souza-Neto JA, Powell JR, Bonizzoni M: **Aedes aegypti vector competence studies: A review.** *Infection,*
Genetics and Evolution 2019, **67:**191-209.
2.   Diagne C, Leroy B, Vaissière A-C, Gozlan RE, Roiz D, Jarić I, Salles J-M, Bradshaw CJ, Courchamp F:
High and rising economic costs of biological invasions worldwide. *Nature* 2021:1-6.
3.   Weaver SC, Reisen WK: **Present and future arboviral threats.** *Antiviral research* 2010, **85:**328-345.
4.   Chibueze EC, Tirado V, da Silva Lopes K, Balogun OO, Takemoto Y, Swa T, Dagvadorj A, Nagata C,
Morisaki N, Menendez C: **Zika virus infection in pregnancy: a systematic review of disease course and**
complications. *Reproductive health* 2017, **14:**1-14.
5.   Cheng G, Liu Y, Wang P, Xiao X: **Mosquito defense strategies against viral infection.** *Trends in*
parasitology 2016, **32:**177-186.
6.   Lee W-S, Webster JA, Madzokere ET, Stephenson EB, Herrero LJ: **Mosquito antiviral defense**
mechanisms: a delicate balance between innate immunity and persistent viral infection. *Parasites &*
vectors 2019, **12:**1-12.
7.   Bronkhorst AW, van Rij RP: **The long and short of antiviral defense: small RNA-based immunity in**
insects. *Current opinion in virology* 2014, **7:**19-28.
8.   Blair CD: **Mosquito RNAi is the major innate immune pathway controlling arbovirus infection and**
transmission. *Future microbiology* 2011, **6:**265-277.
9.   Schnettler E, Donald CL, Human S, Watson M, Siu RW, McFarlane M, Fazakerley JK, Kohl A, Fragkoudis
R: **Knockdown of piRNA pathway proteins results in enhanced Semliki Forest virus production in**
mosquito cells. *The Journal of general virology* 2013, **94:**1680.
10. Morazzani EM, Wiley MR, Murreddu MG, Adelman ZN, Myles KM: **Production of virus-derived ping-**
pong-dependent piRNA-like small RNAs in the mosquito soma. *PLoS pathogens* 2012, **8:**e1002470.
11. Varjak M, Maringer K, Watson M, Sreenu VB, Fredericks AC, Pondeville E, Donald CL, Sterk J, Kean J,
Vazeille M: **Aedes aegypti Piwi4 is a noncanonical PIWI protein involved in antiviral responses.**
MSphere 2017, **2:**e00144-00117.
12. Tassetto M, Kunitomi M, Whitfield ZJ, Dolan PT, Sánchez-Vargas I, Garcia-Knight M, Ribiero I, Chen T,
Olson KE, Andino R: **Control of RNA viruses in mosquito cells through the acquisition of vDNA and**
endogenous viral elements. *Elife* 2019, **8:**e41244.
13. Suzuki Y, Baidaliuk A, Miesen P, Frangeul L, Crist AB, Merkling SH, Fontaine A, Lequime S, Moltini-
Conclois I, Blanc H, et al: **Non-retroviral Endogenous Viral Element Limits Cognate Virus Replication**
in Aedes aegypti Ovaries. *Current Biology* 2020, **30:**3495-3506.e3496.
14. Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ: **Discrete small RNA-**
generating loci as master regulators of transposon activity in Drosophila. *Cell* 2007, **128:**1089-1103.
15. Vagin VV, Sigova A, Li C, Seitz H, Gvozdev V, Zamore PD: **A distinct small RNA pathway silences**
selfish genetic elements in the germline. *Science* 2006, **313:**320-324.
16. Saito K, Nishida KM, Mori T, Kawamura Y, Miyoshi K, Nagami T, Siomi H, Siomi MC: **Specific**
association of Piwi with rasiRNAs derived from retrotransposon and heterochromatic regions in the
Drosophila genome. *Genes & development* 2006, **20:**2214-2222.
17. Li C, Vagin VV, Lee S, Xu J, Ma S, Xi H, Seitz H, Horwich MD, Syrzycka M, Honda BM: **Collapse of**
germline piRNAs in the absence of Argonaute3 reveals somatic piRNAs in flies. *Cell* 2009, **137:**509-
521.
18. Malone CD, Brennecke J, Dus M, Stark A, McCombie WR, Sachidanandam R, Hannon GJ: **Specialized**
piRNA pathways act in germline and somatic tissues of the Drosophila ovary. *Cell* 2009, **137:**522-535.
19. Goriaux C, Desset S, Renaud Y, Vaury C, Brasset E: **Transcriptional properties and splicing of the**
flamenco pi RNA cluster. *EMBO reports* 2014, **15:**411-418.
20. Dennis C, Brasset E, Sarkar A, Vaury C: **Export of piRNA precursors by EJC triggers assembly of**
cytoplasmic Yb-body in Drosophila. *Nature communications* 2016, **7:**1-12.
21. Qi H, Watanabe T, Ku H-Y, Liu N, Zhong M, Lin H: **The Yb body, a major site for Piwi-associated**
RNA biogenesis and a gateway for Piwi expression and transport to the nucleus in somatic cells.
Journal of Biological Chemistry 2011, **286:**3789-3797.
22. Klattenhoff C, Xi H, Li C, Lee S, Xu J, Khurana JS, Zhang F, Schultz N, Koppetsch BS, Nowosielska A:
The Drosophila HP1 homolog Rhino is required for transposon silencing and piRNA production by
dual-strand clusters. *Cell* 2009, **138:**1137-1149.

23. Pane A, Jiang P, Zhao DY, Singh M, Schüpbach T: **The Cutoff protein regulates piRNA cluster**
expression and piRNA production in the Drosophila germline. *The EMBO journal* 2011, **30:**4601-4615.
24. Mohn F, Sienski G, Handler D, Brennecke J: **The rhino-deadlock-cutoff complex licenses noncanonical**
transcription of dual-strand piRNA clusters in Drosophila. *Cell* 2014, **157:**1364-1379.
25. Zhang Z, Wang J, Schultz N, Zhang F, Parhad SS, Tu S, Vreven T, Zamore PD, Weng Z, Theurkauf WE:
The HP1 homolog rhino anchors a nuclear complex that suppresses piRNA precursor splicing. *Cell*
2014, **157:**1353-1363.
26. Chen Y-CA, Stuwe E, Luo Y, Ninova M, Le Thomas A, Rozhavskaya E, Li S, Vempati S, Laver JD, Patel
DJ: **Cutoff suppresses RNA polymerase II termination to ensure expression of piRNA precursors.**
Molecular cell 2016, **63:**97-109.
27. Le Thomas A, Stuwe E, Li S, Du J, Marinov G, Rozhkov N, Chen Y-CA, Luo Y, Sachidanandam R, Toth
KF: **Transgenerationally inherited piRNAs trigger piRNA biogenesis by changing the chromatin of**
piRNA clusters and inducing precursor processing. *Genes & development* 2014, **28:**1667-1680.
28. Andersen PR, Tirian L, Vunjak M, Brennecke J: **A heterochromatin-dependent transcription machinery**
drives piRNA expression. *Nature* 2017, **549:**54-59.
29. Chang TH, Mattei E, Gainetdinov I, Colpan C, Weng Z, Zamore PD: **Maelstrom represses canonical**
polymerase II transcription within bi-directional piRNA clusters in Drosophila melanogaster.
Molecular cell 2019, **73:**291-303. e296.
30. ElMaghraby MF, Andersen PR, Pühringer F, Hohmann U, Meixner K, Lendl T, Tirian L, Brennecke J: **A**
heterochromatin-specific RNA export pathway facilitates piRNA production. *Cell* 2019, **178:**964-979.
e920.
31. Kneuss E, Munafò M, Eastwood EL, Deumer U-S, Preall JB, Hannon GJ, Czech B: **Specialization of the**
Drosophila nuclear export family protein Nxf3 for piRNA precursor export. *Genes & development*
2019, **33:**1208-1220.
32. Pek JW, Patil VS, Kai T: **piRNA pathway and the potential processing site, the nuage, in the**
Drosophila germline. *Development, growth & differentiation* 2012, **54:**66-77.
33. Zhang F, Wang J, Xu J, Zhang Z, Koppetsch BS, Schultz N, Vreven T, Meignin C, Davis I, Zamore PD:
UAP56 couples piRNA clusters to the perinuclear transposon silencing machinery. *Cell* 2012,
151:871-884.
34. Lim AK, Kai T: **Unique germ-line organelle, nuage, functions to repress selfish genetic elements in**
Drosophila melanogaster. *Proceedings of the National Academy of Sciences* 2007, **104:**6714-6719.
35. Miesen P, Girardi E, van Rij RP: **Distinct sets of PIWI proteins produce arbovirus and transposon-**
derived piRNAs in Aedes aegypti mosquito cells. *Nucleic acids research* 2015, **43:**6545-6556.
36. Joosten J, Miesen P, Taşköprü E, Pennings B, Jansen PW, Huynen MA, Vermeulen M, Van Rij RP: **The**
Tudor protein Veneno assembles the ping-pong amplification complex that produces viral piRNAs in
Aedes mosquitoes. *Nucleic acids research* 2019, **47:**2546-2559.
37. Joosten J, Overheul GJ, Van Rij RP, Miesen P: **Endogenous piRNA-guided slicing triggers responder**
and trailer piRNA production from viral RNA in Aedes aegypti mosquitoes. *Nucleic acids research*
2021, **49:**8886-8899.
38. Betting V, Joosten J, Halbach R, Thaler M, Miesen P, Van Rij RP: **A piRNA-lncRNA regulatory network**
initiates responder and trailer piRNA formation during mosquito embryonic development. *RNA* 2021,
27:1155-1172.
39. Halbach R, Miesen P, Joosten J, Taşköprü E, Rondeel I, Pennings B, Vogels CB, Merkling SH, Koenraadt
CJ, Lambrechts L: **A satellite repeat-derived piRNA controls embryonic development of Aedes.** *Nature*
2020, **580:**274-277.
40. Miesen P, Joosten J, van Rij RP: **PIWIs go viral: arbovirus-derived piRNAs in vector mosquitoes.** *PLoS*
pathogens 2016, **12:**e1006017.
41. Akbari OS, Antoshechkin I, Amrhein H, Williams B, Diloreto R, Sandler J, Hay BA: **The developmental**
transcriptome of the mosquito Aedes aegypti, an invasive species and major arbovirus vector. *G3:*
Genes, Genomes, Genetics 2013, **3:**1493-1509.
42. Ma Q, Srivastav SP, Gamez S, Dayama G, Feitosa-Suntheimer F, Patterson EI, Johnson RM, Matson EM,
Gold AS, Brackney DE: **A mosquito small RNA genomics resource reveals dynamic evolution and host**
responses to viruses and transposons. *Genome research* 2021, **31:**512-528.
43. Campbell CL, Black WCt, Hess AM, Foy BD: **Comparative genomics of small RNA regulatory pathway**
components in vector mosquitoes. *BMC Genomics* 2008, **9:**425.

44.   Marconcini M, Hernandez L, Iovino G, Houé V, Valerio F, Palatini U, Pischedda E, Crawford JE, White
BJ, Lin T, et al: **Polymorphism analyses and protein modelling inform on functional specialization of**
Piwi clade genes in the arboviral vector Aedes albopictus. *PLOS Neglected Tropical Diseases* 2019,
13:e0007919.

45.   Mérel V, Boulesteix M, Fablet M, Vieira C: **Transposable elements in Drosophila.** *Mobile DNA* 2020,
11:1-20.

46.   Bergman CM, Quesneville H, Anxolabehere D, Ashburner M: **Recurrent insertion and duplication**
generate networks of transposable element sequences in the Drosophila melanogaster genome.
Genome Biol 2006, **7:**R112.

47.   Arensburger P, Hice RH, Wright JA, Craig NL, Atkinson PW: **The mosquito Aedes aegypti has a large**
genome size and high transposable element load but contains a low proportion of transposon-specific
piRNAs. *BMC genomics* 2011, **12:**1-23.

48.   Girardi E, Miesen P, Pennings B, Frangeul L, Saleh M-C, Ronald P: **Histone-derived piRNA biogenesis**
depends on the ping-pong partners Piwi5 and Ago3 inAedes aegypti. *Nucleic Acids Research*
2017**:**gkw1368.

49.   Vodovar N, Bronkhorst AW, Van Cleef KWR, Miesen P, Blanc H, Van Rij RP, Saleh M-C: **Arbovirus-**
Derived piRNAs Exhibit a Ping-Pong Signature in Mosquito Cells. *PLoS ONE* 2012, **7:**e30861.

50.   Miesen P, Ivens A, Buck AH, van Rij RP: **Small RNA profiling in dengue virus 2-infected Aedes**
mosquito cells reveals viral piRNAs and novel host miRNAs. *PLoS neglected tropical diseases* 2016,
10:e0004452.

51.   Whitfield ZJ, Dolan PT, Kunitomi M, Tassetto M, Seetin MG, Oh S, Heiner C, Paxinos E, Andino R: **The**
diversity, structure, and function of heritable adaptive immunity sequences in the Aedes aegypti
genome. *Current Biology* 2017, **27:**3511-3519. e3517.

52.   Horst AMt, Nigg JC, Dekker FM, Falk BW, Pfeiffer JK: **Endogenous Viral Elements Are Widespread in**
Arthropod Genomes and Commonly Give Rise to PIWI-Interacting RNAs. *Journal of Virology* 2019,
93:e02124-02118.

53.   Suzuki Y, Frangeul L, Dickson LB, Blanc H, Verdier Y, Vinh J, Lambrechts L, Saleh M-C: **Uncovering**
the Repertoire of Endogenous Flaviviral Elements in Aedes Mosquito Genomes. *Journal of Virology*
2017, **91:**JVI.00571-00517.

54.   Aguiar ERGR, Almeida JPP, Queiroz LR, Oliveira LS, Olmo RP, Faria IJS, Imler J-L, Gruber A, Matthews
BJ, Marques JT: **A single unidirectional piRNA cluster similar to the flamenco locus is the major**
source of EVE-derived transcription and small RNAs in Aedes aegypti mosquitoes. *RNA*
2020**:**rna.073965.073119.

55.   Palatini U, Masri RA, Cosme LV, Koren S, Thibaud-Nissen F, Biedler JK, Krsticevic F, Johnston JS,
Halbach R, Crawford JE: **Improved reference genome of the arboviral vector Aedes albopictus.**
Genome biology 2020, **21:**1-29.

56.   Crava CM, Varghese FS, Pischedda E, Halbach R, Palatini U, Marconcini M, Gasmi L, Redmond S, Afrane
Y, Ayala D: **Population genomics in the arboviral vector Aedes aegypti reveals the genomic**
architecture and evolution of endogenous viral elements. *Molecular ecology* 2021, **30:**1594-1611.

57.   Wierzbicki F, Kofler R, Signor S: **Evolutionary dynamics of piRNA clusters in Drosophila.** *Molecular*
ecology 2021.

58.   Assis R, Kondrashov AS: **Rapid repetitive element-mediated expansion of piRNA clusters in**
mammalian evolution. *Proceedings of the National Academy of Sciences* 2009, **106:**7079-7082.

59.   Lewis SH, Quarles KA, Yang Y, Tanguy M, Frézal L, Smith SA, Sharma PP, Cordaux R, Gilbert C, Giraud
I: **Pan-arthropod analysis reveals somatic piRNAs as an ancestral defence against transposable**
elements. *Nature ecology & evolution* 2018, **2:**174-181.

60.   Zhang X, Aksoy E, Girke T, Raikhel AS, Karginov FV: **Transcriptome-wide microRNA and target**
dynamics in the fat body during the gonadotrophic cycle of Aedes aegypti. *Proceedings of the National*
Academy of Sciences 2017, **114:**E1895-E1903.

61.   Adelman ZN, Anderson MA, Wiley MR, Murreddu MG, Samuel GH, Morazzani EM, Myles KM: **Cooler**
temperatures destabilize RNA interference and increase susceptibility of disease vector mosquitoes to
viral infection. *PLoS neglected tropical diseases* 2013, **7:**e2239.

62.   Olmo RP, Ferreira AG, Izidoro-Toledo TC, Aguiar ER, de Faria IJ, de Souza KP, Osório KP, Kuhn L,
Hammann P, de Andrade EG: **Control of dengue virus in the midgut of Aedes aegypti by ectopic**
expression of the dsRNA-binding protein Loqs2. *Nature Microbiology* 2018, **3:**1385-1393.

63.   Rosenkranz D, Zischler H: **proTRAC-a software for probabilistic piRNA cluster detection,**
899   **visualization and analysis.** *BMC bioinformatics* 2012, **13:**1-10.

900   64.   Prud'Homme N, Gans M, Masson M, Terzian C, Bucheton A: **Flamenco, a gene controlling the gypsy**
901   **retrovirus of Drosophila melanogaster.** *Genetics* 1995, **139:**697-711.

902   65.   Atlasi Y, Stunnenberg HG: **The interplay of epigenetic marks during stem cell differentiation and**
903   **development.** *Nature Reviews Genetics* 2017, **18:**643-658.

904   66.   Huang X, Gao X, Li W, Jiang S, Li R, Hong H, Zhao C, Zhou P, Chen H, Bo X: **Stable H3K4me3 is**
905   **associated with transcription initiation during early embryo development.** *Bioinformatics* 2019,
906   **35:**3931-3936.

907   67.   Ernst J, Kheradpour P, Mikkelsen TS, Shoresh N, Ward LD, Epstein CB, Zhang X, Wang L, Issner R,
908   Coyne M: **Mapping and analysis of chromatin state dynamics in nine human cell types.** *Nature* 2011,
909   **473:**43-49.

910   68.   Bell O, Wirbelauer C, Hild M, Scharf AN, Schwaiger M, MacAlpine DM, Zilbermann F, Van Leeuwen F,
911   Bell SP, Imhof A: **Localized H3K36 methylation states define histone H4K16 acetylation during**
912   **transcriptional elongation in Drosophila.** *The EMBO journal* 2007, **26:**4974-4984.

913   69.   Narlikar GJ, Sundaramoorthy R, Owen-Hughes T: **Mechanisms and functions of ATP-dependent**
914   **chromatin-remodeling enzymes.** *Cell* 2013, **154:**490-503.

915   70.   Ernst J, Kellis M: **Chromatin-state discovery and genome annotation with ChromHMM.** *Nature*
916   *protocols* 2017, **12:**2478-2492.

917   71.   Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ: **Transposition of native chromatin for**
918   **fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome**
919   **position.** *Nature methods* 2013, **10:**1213-1218.

920   72.   Gorkin DU, Barozzi I, Zhao Y, Zhang Y, Huang H, Lee AY, Li B, Chiou J, Wildberg A, Ding B: **An atlas**
921   **of dynamic chromatin landscapes in mouse fetal development.** *Nature* 2020, **583:**744-751.

922   73.   Fong N, Saldi T, Sheridan RM, Cortazar MA, Bentley DL: **RNA Pol II dynamics modulate co-**
923   **transcriptional chromatin modification, CTD phosphorylation, and transcriptional direction.**
924   *Molecular cell* 2017, **66:**546-557. e543.

925   74.   Schroeder SC, Schwer B, Shuman S, Bentley D: **Dynamic association of capping enzymes with**
926   **transcribing RNA polymerase II.** *Genes & development* 2000, **14:**2435-2440.

927   75.   Lunde BM, Reichow SL, Kim M, Suh H, Leeper TC, Yang F, Mutschler H, Buratowski S, Meinhart A,
928   Varani G: **Cooperative interaction of transcription termination factors with the RNA polymerase II C-**
929   **terminal domain.** *Nature structural & molecular biology* 2010, **17:**1195-1201.

930   76.   Yan W, Chen D, Schumacher J, Durantini D, Engelhorn J, Chen M, Carles CC, Kaufmann K: **Dynamic**
931   **control of enhancer activity drives stage-specific gene expression during flower morphogenesis.**
932   *Nature communications* 2019, **10:**1-16.

933   77.   Lopez-Atalaya JP, Ito S, Valor LM, Benito E, Barco A: **Genomic targets, and histone acetylation and**
934   **gene expression profiling of neural HDAC inhibition.** *Nucleic acids research* 2013, **41:**8072-8084.

935   78.   Ando M, Saito Y, Xu G, Bui NQ, Medetgul-Ernar K, Pu M, Fisch K, Ren S, Sakai A, Fukusumi T:
936   **Chromatin dysregulation and DNA methylation at transcription start sites associated with**
937   **transcriptional repression in cancers.** *Nature communications* 2019, **10:**1-15.

938   79.   Morselli M, Pastor WA, Montanini B, Nee K, Ferrari R, Fu K, Bonora G, Rubbi L, Clark AT, Ottonello S:
939   **In vivo targeting of de novo DNA methylation by histone modifications in yeast and mouse.** *Elife* 2015,
940   **4:**e06205.

941   80.   Amemiya HM, Kundaje A, Boyle AP: **The ENCODE blacklist: identification of problematic regions of**
942   **the genome.** *Scientific reports* 2019, **9:**1-5.

943   81.   Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features.**
944   *Bioinformatics* 2010, **26:**841-842.

945   82.   Maringer K, Yousuf A, Heesom KJ, Fan J, Lee D, Fernandez-Sesma A, Bessant C, Matthews DA,
946   Davidson AD: **Proteomics informed by transcriptomics for characterising active transposable**
947   **elements and genome annotation in Aedes aegypti.** *BMC genomics* 2017, **18:**1-18.

948   83.   Joosten J, Taşköprü E, Jansen PW, Pennings B, Vermeulen M, Van Rij RP: **PIWI proteomics identifies**
949   **Atari and Pasilla as piRNA biogenesis factors in Aedes mosquitoes.** *Cell Reports* 2021, **35:**109073.

950   84.   Olivieri D, Senti K-A, Subramanian S, Sachidanandam R, Brennecke J: **The cochaperone shutdown**
951   **defines a group of biogenesis factors essential for all piRNA populations in Drosophila.** *Molecular cell*
952   2012, **47:**954-969.

85.   Chen X-G, Jiang X, Gu J, Xu M, Wu Y, Deng Y, Zhang C, Bonizzoni M, Dermauw W, Vontas J: **Genome**
sequence of the Asian Tiger mosquito, Aedes albopictus, reveals insights into its biology, genetics, and
evolution. *Proceedings of the National Academy of Sciences* 2015, **112:**E5907-E5915.

86.   Palatini U, Miesen P, Carballar-Lejarazu R, Ometto L, Rizzo E, Tu Z, van Rij RP, Bonizzoni M:
Comparative genomics shows that viral integrations are abundant and express piRNAs in the
arboviral vectors Aedes aegypti and Aedes albopictus. *BMC genomics* 2017, **18:**1-15.

87.   Ghildiyal M, Seitz H, Horwich MD, Li C, Du T, Lee S, Xu J, Kittler EL, Zapp ML, Weng Z: **Endogenous**
siRNAs derived from transposons and mRNAs in Drosophila somatic cells. *Science* 2008, **320:**1077-
1081.

88.   Zanni V, Eymery A, Coiffet M, Zytnicki M, Luyten I, Quesneville H, Vaury C, Jensen S: **Distribution,**
evolution, and diversity of retrotransposons at the flamenco locus reflect the regulatory properties of
piRNA clusters. *Proceedings of the National Academy of Sciences* 2013, **110:**19842-19847.

89.   Chen P, Luo Y, Aravin AA: **RDC complex executes a dynamic piRNA program during Drosophila**
spermatogenesis to safeguard male fertility. *PLoS genetics* 2021, **17:**e1009591.

90.   Chen P, Kotov AA, Godneeva BK, Bazylev SS, Olenina LV, Aravin AA: **piRNA-mediated gene**
regulation and adaptation to sex-specific transposon expression in D. melanogaster male germline.
Genes & development 2021, **35:**914-935.

91.   Wang Y, Jin B, Liu P, Li J, Chen X, Gu J: **piRNA profiling of dengue virus type 2-infected Asian tiger**
mosquito and midgut tissues. *Viruses* 2018, **10:**213.

92.   Franz AW, Kantor AM, Passarelli AL, Clem RJ: **Tissue barriers to arbovirus infection in mosquitoes.**
Viruses 2015, **7:**3741-3767.

93.   Katahira J, Sträßer K, Podtelejnikov A, Mann M, Jung JU, Hurt E: **The Mex67p-mediated nuclear mRNA**
export pathway is conserved from yeast to human. *The EMBO journal* 1999, **18:**2593-2609.

94.   Handler D, Olivieri D, Novatchkova M, Gruber FS, Meixner K, Mechtler K, Stark A, Sachidanandam R,
Brennecke J: **A systematic analysis of Drosophila TUDOR domain-containing proteins identifies**
Vreteno and the Tdrd12 family as essential primary piRNA pathway factors. *The EMBO journal* 2011,
30:3977-3993.

95.   Goic B, Stapleford KA, Frangeul L, Doucet AJ, Gausson V, Blanc H, Schemmel-Jofre N, Cristofari G,
Lambrechts L, Vignuzzi M: **Virus-derived DNA drives mosquito vector tolerance to arboviral**
infection. *Nature communications* 2016, **7:**1-10.

96.   Dietrich I, Jansen S, Fall G, Lorenzen S, Rudolf M, Huber K, Heitmann A, Schicht S, Ndiaye EH, Watson
M: **RNA interference restricts Rift Valley Fever virus in multiple insect systems.** *Msphere* 2017,
2:e00090-00017.

97.   Rückert C, Prasad AN, Garcia-Luna SM, Robison A, Grubaugh ND, Weger-Lucarelli J, Ebel GD: **Small**
RNA responses of Culex mosquitoes and cell lines during acute and persistent virus infection. *Insect*
biochemistry and molecular biology 2019, **109:**13-23.

98.   Bryant WB, Ray S, Mills MK: **Global analysis of small non-coding RNA populations across tissues in**
the malaria vector, Anopheles gambiae. *Insects* 2020, **11:**406.

99.   Martin M: **Cutadapt removes adapter sequences from high-throughput sequencing reads.** *EMBnet*
journal 2011, **17:**10-12.

100.   Langmead B, Trapnell C, Pop M, Salzberg SL: **Ultrafast and memory-efficient alignment of short DNA**
sequences to the human genome. *Genome biology* 2009, **10:**1-10.

101.   Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T: **deepTools: a flexible platform for exploring deep-**
sequencing data. *Nucleic acids research* 2014, **42:**W187-W191.

102.   Wickham H: **Package 'ggplot2': elegant graphics for data analysis.** *Springer-Verlag New York doi* 2016,
10:978-970.

103.   Andrews S: **FastQC: a quality control tool for high throughput sequence data.** Babraham
Bioinformatics, Babraham Institute, Cambridge, United Kingdom; 2010.

104.   Li H, Durbin R: **Fast and accurate short read alignment with Burrows–Wheeler transform.**
bioinformatics 2009, **25:**1754-1760.

105.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li
W: **Model-based analysis of ChIP-Seq (MACS).** *Genome biology* 2008, **9:**1-9.

106.   Yue F, Cheng Y, Breschi A, Vierstra J, Wu W, Ryba T, Sandstrom R, Ma Z, Davis C, Pope BD: **A**
comparative encyclopedia of DNA elements in the mouse genome. *Nature* 2014, **515:**355-364.

107.   Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, Kheradpour P, Zhang Z, Wang
J, Ziller MJ: **Integrative analysis of 111 reference human epigenomes.** *Nature* 2015, **518:**317-330.

108. Georgiou G, van Heeringen SJ: **Fluff: exploratory analysis and visualization of high-throughput**
sequencing data. *PeerJ* 2016, **4:**e2209.
109. Qu J, Yi G, Zhou H: **p63 cooperates with CTCF to modulate chromatin architecture in skin**
keratinocytes. *Epigenetics & chromatin* 2019, **12:**1-13.
110. Kozomara A, Birgaoanu M, Griffiths-Jones S: **miRBase: from microRNA sequences to function.** *Nucleic*
acids research 2019, **47:**D155-D162.
111. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R: **The**
sequence alignment/map format and SAMtools. *Bioinformatics* 2009, **25:**2078-2079.
112. Crooks GE, Hon G, Chandonia J-M, Brenner SE: **WebLogo: a sequence logo generator.** *Genome research*
2004, **14:**1188-1190.
113. Pitts RJ, Liu C, Zhou X, Malpartida JC, Zwiebel LJ: **Odorant receptor-mediated sperm activation in**
disease vector mosquitoes. *Proceedings of the National Academy of Sciences* 2014, **111:**2566-2571.
114. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR:
STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013, **29:**15-21.
115. Trapnell C, Hendrickson DG, Sauvageau M, Goff L, Rinn JL, Pachter L: **Differential analysis of gene**
regulation at transcript resolution with RNA-seq. *Nature biotechnology* 2013, **31:**46-53.
116. Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, Nguyen JT, Barbisin M, Xu NL, Mahuvakar VR,
Andersen MR: **Real-time quantification of microRNAs by stem–loop RT–PCR.** *Nucleic acids research*
2005, **33:**e179-e179.
117. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR**
and the 2− ΔΔCT method. *methods* 2001, **25:**402-408.
118. Li XZ, Roy CK, Dong X, Bolcun-Filas E, Wang J, Han BW, Xu J, Moore MJ, Schimenti JC, Weng Z: **An**
ancient transcription factor initiates the burst of piRNA production during early meiosis in mouse
testes. *Molecular cell* 2013, **50:**67-81.
119. Sun X-J, Wei J, Wu X-Y, Hu M, Wang L, Wang H-H, Zhang Q-H, Chen S-J, Huang Q-H, Chen Z:
Identification and characterization of a novel human histone H3 lysine 36-specific methyltransferase.
Journal of Biological Chemistry 2005, **280:**35261-35271.
120. Huang H, Weng H, Zhou K, Wu T, Zhao BS, Sun M, Chen Z, Deng X, Xiao G, Auer F: **Histone H3**
trimethylation at lysine 36 guides m6A RNA modification co-transcriptionally. *Nature* 2019, **567:**414-
419.
121. Machado SR, Qu J, Koopman WJH, Miesen P: **The DEAD-box RNA helicase Dhx15 controls glycolysis**
and arbovirus replication in Aedes aegypti mosquito cells. *bioRxiv*
2022**:**2022.06.2022.497281.

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

Extended data figures and tables

1056

Fig. S1 Annotation of piRNA clusters in *Ae. aegypti* mosquitoes.

a. Optimization of parameters for piRNA cluster prediction with an established pipeline that did not include assumptions on nucleotide biases or strand asymmetry [55, 56]. Dot colors as in Fig. 1a.

Fig. S2 Quality check of the chromatin profiling dataset in Aag2 cells.

a. Chromatin state annotation using ChromHMM with different numbers of states based on ChIP-seq data in Aag2 cells.
b. Density profiles of ChIP-seq data of the indicated histone marks. Log scaled read counts within ChIP-seq peaks between two biological replicates were plotted. A level bar reflecting the number of ChIP-seq peaks is shown in the lower right corner. For detailed ChIP-seq peaks, see Supplementary Table 2.
c. Genomic distribution of five chromatin states. Chromatin state colors are as in Fig. 2a and the *Ae. aegypti* genome background distribution was shown as the 1st gray bar. For instance, 54.9% of the promoter state was located in intron regions. The category 'others' consists of non-coding RNAs and pseudogenes.
d. Density profiles of RNA pol II, RNA pol II ser5p, and RNA pol II ser2p ChIP-seq in duplicates. For detailed information about peaks enriched for the three RNA pol II, see Supplementary Table 2.
e. Published RNA-seq datasets were used to define expressed genes [39]. A cutoff of $\log_{10}$FPKM=-0.6 was chosen to distinguish expressed genes from non-expressed genes (Supplementary Table 3).

Fig. S3 Chromatin signatures of Aag2 piRNA clusters

a. Band plots (upper panel) and heat maps (lower panel) showing average signals of ChIP-seq data across 215 piRNA clusters in Aag2 cells (left) and 158 piRNA clusters (right) after removal of 57 piRNA clusters that overlapped with strong ChIP input signals (regions from MACS2 peak calling, described in M&M) to exclude background noise from genomic DNA. ChIP-seq signals were normalized to RPKM (y-axis). In the heatmap, all piRNA clusters were scaled to 5 kb, and the upstream/downstream 5 kb regions were shown.

b. Representative genome browser tracks showing examples of a readthrough piRNA cluster, a genic piRNA cluster, and a stand-alone piRNA cluster. ATAC-seq, ChIP-seq, and RNA-seq signals, as well as piRNA-sized reads from small RNA-seq, were normalized to RPKM. The boundaries of each cluster are indicated with dashed lines and the direction of piRNA clusters is indicated with arrows.

1094

Fig. S4 A large-scale RNAi screen to identify key factors involved in cluster-derived piRNA biogenesis

a. Knockdown of *AAEL010451* using another dsRNA. RT-qPCR was performed to confirm gene knockdown, while SL-RT-qPCR was performed to quantify piRNA expression. All experiments were performed in biological triplicates (shown as dots) unless otherwise specified. The interquartile range and the median are shown in box plots. Statistical differences were examined by unpaired t-test, when compared to the non-targeting control knockdown of luciferase (dsLuc). Symbols denote statistical significance (* $p < 0.05$). For detailed primer information, see Supplementary Table 4.

b. RT-qPCR was performed to confirm knockdown of target genes while SL-RT-qPCR was performed to quantify piRNA
abundance. All experiments were performed in biological duplicates or triplicates (shown as dots). The interquartile range and the
median were shown in box plots. Statistical differences were examined by unpaired t-test, when compared to the non-targeting
control knockdown of luciferase (dsLuc). Symbols denote statistical significance (* $p < 0.05$, *** $p < 0.001$). For detailed primer
information, see Supplementary Table 4.
c-f. RNAi screen was performed with orthologs of known piRNA biogenesis factors in other organisms and chromatin modifiers.
In other model systems, transcription factors such as A-MYB and Ci are important for piRNA cluster transcription [19, 118]. We
also tested the H3K36me3 writer, Setd2 [119], due to the observed readthrough transcription that was marked by rather a strong
continuation of H3K36me3 signals (Fig. 3a and d); and subsequently, two m^{6}A writers, Mellt3 and Mellt14, as recent studies
suggested H3K36me3 guided the deposition of m^{6}A at 3'UTRs [120], which may serve as a signal triggering readthrough
transcription (**c**), 16 mRNA 3'end processing genes (**d**), 18 RNA export genes and confirmation of *Nxf1* knockdown (dsNxf1) with
another set of dsRNA (lower panel) (**e**), 26 general RNA processing genes [39] and 29 RNA helicase genes [121] (**f**).
1113

1114

Fig. S5 Systematic analysis of piRNA clusters in *Ae. albopictus* mosquitoes.

a. Comparison of piRNA expression within piRNA clusters between the U4.4 cell line, somatic tissue (carcass), and germline tissues
(ovary and testes) of *Ae. albopictus*. piRNA clusters were independently annotated in each dataset and represented as dots, with
color representing the origin and dot size representing cluster size. The numbers of clusters are listed. Log scaled piRNA read counts
were used to represent piRNA expression levels within a piRNA cluster. For detailed piRNA cluster information, see Supplementary
Table 5.

b. Cumulative distribution of piRNA cluster coverage on the plus strand of the *Ae. albopictus* genome.
c. Quantification of uni-strand and dual-strand clusters in the indicated cell lines and tissues, with an arbitrary threshold of 80% or
90% piRNAs produced from one genomic strand defined as uni-strand piRNA clusters.
d. Distribution of 37 core piRNA clusters in cell lines and tissues. Within each dataset, piRNA clusters were ranked according to
the cluster size. The 37 core piRNA clusters are shown as dark grey bars; other piRNA clusters are shown as light grey bars. piRNA
density (RPKM) was plotted using the same colors as in **a**.
e. Summary of the origin of core piRNA clusters, the distribution of nrEVEs, and the overlapping gene (for genic clusters) or the
nearest expressed gene in the same orientation for readthrough piRNA clusters and other clusters. 'Unknown' refers to the situation
when a piRNA cluster is located in a scaffold without any annotated gene nearby.
1130

1131

Fig. S6 Systematic analysis of piRNA clusters in *Cx. quinquefasciatus* mosquitoes.

a. Comparison of piRNA expression within piRNA clusters between the Hsu cell line, somatic tissues (salivary glands and midgut),
and germline tissues (ovary and testes) of *Cx. quinquefasciatus*. piRNA clusters were independently annotated in each dataset and
represented as dots, with color representing the origin and dot size representing cluster size. The numbers of clusters are listed. Log
scaled piRNA read counts were used to represent piRNA expression levels within a piRNA cluster. For detailed piRNA cluster
information, see Supplementary Table 6.
b. Cumulative distribution of piRNA cluster coverage on the plus strand of the *Cx. quinquefasciatus* genome.
c. Quantification of uni-strand and dual-strand clusters in the indicated cell lines and tissues, with an arbitrary threshold of 80% or
90% piRNAs produced from one genomic strand defined as uni-strand piRNA clusters.
d. Distribution of 32 core piRNA clusters in cell lines and tissues. Within each dataset, piRNA clusters were ranked according to
the cluster size. The 32 core piRNA clusters are shown as dark grey bars; other piRNA clusters are shown as light grey bars. piRNA
density (RPKM) was plotted using the same colors as in **a**.
e. Summary of the origin of core piRNA clusters and the overlapping gene (for genic clusters) or the nearest expressed gene in the
same orientation for readthrough piRNA clusters and other clusters. 'Unknown' refers to the situation when a piRNA cluster is
located in a scaffold without any annotated gene nearby.
1147

1148

Fig. S7 Systematic analysis of piRNA clusters in *An. gambiae* mosquitoes.

a. Comparison of piRNA expression within piRNA clusters between the Mos55 cell line, somatic tissues (head thorax, midgut and
fat body), and germline tissues (ovary and testes) of *An. gambiae*. piRNA clusters were independently annotated in each dataset and
represented as dots, with color representing the origin and dot size representing cluster size. The numbers of clusters are listed. Log
scaled piRNA read counts were used to represent piRNA expression levels within a piRNA cluster. For detailed piRNA cluster
information, see Supplementary Table 7.
b. Cumulative distribution of piRNA cluster coverage on the plus strand of the *An. gambiae* genome.
c. Quantification of uni-strand and dual-strand clusters in the indicated cell lines and tissues, with an arbitrary threshold of 80% or
90% piRNAs produced from one genomic strand defined as uni-strand piRNA clusters.
d. Distribution of 19 core piRNA clusters in cell lines and tissues. Within each dataset, piRNA clusters were ranked according to
the cluster size. The 19 core piRNA clusters are shown as dark grey bars; other piRNA clusters are shown as light grey bars. piRNA
density (RPKM) is plotted using the same colors as in **a**.
e. Summary of the origin of core piRNA clusters and the overlapping gene (for genic clusters) or the nearest expressed gene in the
same orientation for readthrough piRNA clusters and other clusters. 'Unknown' refers to the situation when a piRNA cluster is
located in a scaffold without any annotated gene nearby.
1164

1165

Fig. S8 Comparison of upstream genes across different mosquito species.

a. Determination of the expressed genes in each species. A cutoff was chosen at the dip of the distribution plot to separate expressed
genes from non-expressed genes, which are detailed in Supplementary Table 3.
b. In *Ae. aegypti*, *AAEL005473* is associated with a genic piRNA cluster (cluster 21). Its two ortholog genes in *Ae. albopictus* are
associated with a readthrough piRNA cluster (*AALFPA_048291*, cluster 15) and a genic piRNA cluster (*AALFPA_079151*, cluster
20). The boundaries of each cluster are indicated with dashed lines and the direction of the piRNA cluster is indicated with an arrow
on top. nrEVEs were shown as arrows, with colors representing viral families.
1173

1174

Supplementary Information

Additional file 1: Supplementary Figures S1-S8

Additional file 2: Table S1. piRNA clusters in Ae. aegypti.xlsx

Additional file 3: Table S2. chromatin profiles in Ae. aegypti Aag2 cells.xlsx

Additional file 4: Table S3. Expressed genes in four mosquito species.xlsx

Additional file 5: Table S4. A summary of primers.xlsx

Additional file 6: Table S5. piRNA clusters in Ae. albopictus.xlsx

Additional file 7: Table S6. piRNA clusters in Cx. quinquefasciatus.xlsx

Additional file 8: Table S7. piRNA clusters in An. gambiae.xlsx

Additional file 9: Table S8. Summary of nrEVEs in core piRNA clusters of Aedes mosquitoes.xlsx