# Can growth in captivity alter the calcaneal microanatomy of a wild ungulate?

**Romain cottereau[1,#], Katia Ortiz[2,3], Yann[2-4] Locatelli,**

**Alexandra Houssaye[1,\*], Thomas Cucchi[5,\*]**

[1] CNRS, UMR 7179 Mécanismes Adaptatifs et Evolution, Muséum d'Histoire Naturelle de Paris, France

[2] Réserve Zoologique de la Haute-Touche, Muséum National d'Histoire Naturelle, Obterre, France

[3] Institut de Systématique, Evolution, Biodiversité, UMR 7205, Muséum National d'Histoire Naturelle CNRS

UPMC EPHE, UA, Paris, France

[4] Physiologie de la Reproduction et des Comportements, UMR 7247, INRAE CNRS Université de Tours IFCE,

Nouzilly, France

[5] UMR 7209 Archéozoologie, Archéobotanique : Sociétés, Pratiques et Environnements, Muséum d'Histoire

Naturelle de Paris, France

*: co last authors

#: corresponding author

## Abstract

Reduced mobility associated with captivity induces changes in biomechanical stress on the
skeleton of domesticated animals. Due to bone plasticity, bone's morphology and internal
structure can respond to these new biomechanical stresses over individuals' lifetime. In a
context where documenting early process of animal domestication is a real zooarchaeological
challenge. This study will test the hypothesis that change in mobility patterns during a wild
ungulate's life will alter the internal structure of its limb bones and provide a proof of concept
for the application of this knowledge in Zooarchaeology. Using the calcaneus as a phenotypic
marker through qualitative and quantitative 3D microanatomical analyses, we relied on a
comparative study across wild boars (*Sus scrofa*) populations from controlled experimental
conditions with different mobility patterns (natural habitat, large pen, and stall) and
archaeological specimens collected from middle and late Mesolithic as surrogate for the norm
of reaction in European wild boar phenotype before the spread of agriculture and domestic pigs.
Results provide evidence for compressive and tensile forces as the main elements affecting the
variation in the cortical thickness along the calcaneus. Furthermore, changes in the internal
structure of the calcaneus between mobility patterns are observed but their intensity is not
directly associated with the degree of mobility restriction and only weakly impacted by the size
or weight of the individuals. Despite having greater bone volume, the calcaneus of the

Mesolithic wild boars displays a very similar microanatomy compared to the present-day hunted or captive wild boars. These results suggest that calcaneal microanatomy is more affected by population differences than by locomotor variation. For all these reasons, this preliminary study doesn't support the use of microanatomy of the calcaneus as an indicator of change in locomotor behaviour induced by captivity in the archaeological record.

Keywords: calcaneus; Bone structure; captivity; domestication; *Sus scrofa*; functional morphology

## Introduction

The bones that make up the vertebrate skeleton are plastic organs whose morphology, both external and internal, adapt in response to physical stresses (Roux, 1881; Wolff, 1986; Hall, 1983, 2005; Ruff et al., 2006; Du et al., 2020), beyond the predominant influence of heredity (Hall, 1989; Cubo et al., 2005). Muscle mobilization applies stresses that influence bone growth, development, and remodeling (Marcus, 2002). Numerous studies on human skeletons have shown that intensive physical activity has characteristic consequences on skeletal microanatomy (Zanker & Swaine, 2000; Modlesky et al., 2008; Maïmoun & Sultan 2011; Maïmoun et al., 2013). Conversely, the bone resorption observed in astronauts (Lang et al., 2004) and bedridden individuals (Krølner & Toft., 1983) illustrates bone accommodation to the absence of gravity (Carmeliet & Bouillon, 2001) and inactivity. Therefore, differences in mobility between individuals, which engenders differences in biomechanical stresses, affect the bones' structure.

Captive animals may grow in limited areas, which does not involve the same range of movement and forces applied to their bones compared to free-ranging individuals. Analysing the biomechanical bone plasticity is a major asset for paleoanthropologists and archaeozoologists trying to decipher individual-scale lifestyle changes from bones anatomy in order to document changes in activity patterns of past humans and other animals (Trinkaus et al., 1994, Agarwal, 2016). In paleoanthropology, this biomechanical component has been used to better understand the evolution of bipedalism in hominids (Ruff, 2018) and to observe the morphological consequences of the transition from a nomadic hunter-gatherer to a farmer-herder lifestyle (Pinhasi & Stock, 2011). In archaeozoology, changes between a domestic and a wild lifestyle related to locomotion and gait changes, have been estimated as a prevalent factor over load-carrying on structural changes of domestic donkeys limb bones (Shackelford et al.,

2013). Despite this, archaeozoologists have always considered that the first observable

morphological changes occurred only late in the domestication process and could not perceive

its initial stages since they would not involve genetic isolation or selective reproduction (Vigne

et al., 2011; Colledge et al., 2016). To develop new phenotypic proxy of the first steps of animal

domestication, an experimental project (DOMEXP) has been performed to test whether growth

in captivity of a wild ungulate would leave a morpho-functional imprint distinguishable from

its natural habitat norm of reaction. Hence, early process of cultural control (Hecker 1982) could

be traced from bones anatomy long before the so-called "domestication syndrome". To test

this hypothesis, the DOMEXP project used a genetically homogeneous wild boar (*Sus scrofa*)

population to control the genetic and environmental factors influencing skeletal variation. From this populations, 6 months old piglets were captured after weaning to be reared until the age of two years under two mobility reduction regimes. Geometric morphometrics' analyses have provided proof of concept that this mobility reduction in a wild ungulate population could leave morphometric prints on skulls (Neaux et al., 2021) and calcaneus (Harbers et al.,

Fig. 1. Lateral view of the tarsal region of a skeletally mature artiodactyl, showing the calcaneus with other associated bones, ligaments and tendons. The large arrow in the dorsal direction indicates the orientation of the force exerted by the Achilles tendon during paw extension, loading the dorsal cortex in compression ("C") and the plantar cortex in tension ("T"). Modified from Su et al. (1999).

2020a). Microanatomical investigations have also revealed changes in the 3D topography of

the cortical thickness in the humeral shaft (Harbers et al., 2020b). However, the impact of

mobility reduction on the micraoanatomical features of the skeleton remained to be explored.

For this reason, this study investigates the microanatomical variation of the calcaneus from the

experimental specimens of the DOMEXP project. Tarsal bones like the calcaneum are

particularly considered as more informative than others in investigating the mechanical forces

that apply to individuals during locomotion (Lovejoy et al., 1999; Carter & Beaupré, 2001;

Pearson & Lieberman, 2004). Proximally articulated to the fibula and distally to the cuboid

bone, the calcaneus is intensely stressed during locomotion in terrestrial tetrapods. It acts as a lever arm for the ankle extensor muscles, and is subject to high tensile, flexural, and compressive forces (Fig. 1; Hussain, 1975; Carrano, 1997; Bleefeld & Bock, 2002; Bassarova et al., 2009; Barone, 2017). The calcaneus is therefore a particularly interesting object of study to investigate differences in mobility between individuals (Harbers et al., 2020a). Moreover, the high compactness of the calcaneus makes it resistant to taphonomic alterations, which gives it a good general preservation in archaeological contexts (Binford, 1978).

This study compares the calcaneus microanatomy variation between adult captive-bred and free-ranging wild boars to assess if captivity can induce microanatomical variation in a wild ungulate. We also include wild boar calcanei from a Mesolithic (ca. 8000 years old) context of hunter-gatherers in northern France to observe how much the variation in wild boar microanatomy has changed since the neolithisation of Europe and whether this proxy can be used to infer change in locomotor behaviour from archaeological calcanei.

## Material and method

### Experimental protocol

24 wild boars aged six months were sampled from a control population living in a fenced forest of about 100,000 $m^2$ in northern France. These specimens were then divided into two equal samples with the same sex ratio and reared until 24 month in two different contexts of reduced mobility: (1) a stall of 100 $m^2$, where males and females were separated (each group had 45 $m^2$) and (2) a wooded pen of 3000 $m^2$. These two contexts represent respectively 99.9% and 97% of range reduction compared to the control population, which prevents the average daily distances measured in free boar populations (Palencia et al. 2019; Russo et al. 1997). The stall presented no opportunity for foraging. In the wooded enclosure, this opportunity was limited due to space limitations. Dry feed pellets suitable for feeding domestic pigs were provided to both groups.

### Specimens studied

Our dataset includes calcaneus bones from 47 specimens: 26 from the DOMEXP experimental farm - 11 stall specimens (stall), 10 pen specimens (pen), five from the control population (control) - 15 hunted wild boars from two Northern France forests: eight from the Chambord forest and seven from the Compiègne forest (Natural habitat) and six archaeological specimens

from the Mesolithic contexts (Meso) of Noyen sur Seine (Mordant et al., 2013, Marinval-Vigne

et al., 1989) in Northern France (Table 1).The six archaeological specimens from Mesolithic

deposits have been accumulated by hunter-gatherer living in Western Europe before the

Neolithic dispersal via the Mediterranean and Danubian pathways, which introduced domestic

pigs from the Near East (Larson 2006) that interbred with local populations of wild boars

(Frantz et al. 2020). Two specimens are radiocarbon dated from the Middle Mesolithic (between

-8000 and -7300 cal. BP) and four specimens to the Final Mesolithic dated between -7000 and

-6200 cal. BP (Mordant et al., 2013). All the specimens bar the archaeological samples have

associated age and sex informations. Three of the archaeological calcanei belong to adult

individuals since their proximal epiphysis is fused which is known to happen around 2 years in

wild boars (Bridault et al., 2000).

## Data acquisition

147    All the calcanei have been scanned with high-resolution microtomography (EasyTom 40-150

148    scanner, RX Solutions) at the MRI platform, hosted at ISEM, University of Montpellier (UMR

149    5554); reconstructions were then performed using X-Act (RX Solutions).

## Virtual thin sections

For qualitative comparisons, three virtual sections (Fig. 2, Appendix 1,2 and 3) were made for

each calcaneus, following Barone (2017) for terminology and orientations. The bones were

oriented as follows: in dorsal view (Fig. 2a), bone's axis is vertical and the fibular trochlea is

oriented toward the observer, its dorsal part aligns with the contour of the bone's medial border;

in medial view (Fig. 2b) the *sustentaculum tali* is directed towards the observer, the observation

angle is fixed when the anterior edge of the fibular trochlea is no longer visible upon rotation

from the anterior view to the medial view. Sagittal sections (SS) run in dorsal view from the

distal end tip and the midpoint of the thickness at the proximal epiphysis base (Fig. 2). The

frontal sections (FS) extend from the distal end tip to the midpoint of the proximal epiphysis

base (see purple arrows on Fig. 2b). Cross-sections (TS) are taken perpendicular to the FS plane

at 1/3 of the total length of the bone (from the proximal epiphysis tip to the distal tip) (Fig. 2).

These sectional planes were chosen to depict large portions of the bone in order to analyze the

microanatomical structure (e.g., trabecular network, cortical thickness) while being easily

created with good reproducibility for all specimens. Virtual sections were created using

VGSTUDIO MAX, versions 2.2 (Volume Graphics Inc.).

## Calcaneus 3D mapping

To observe and measure the thickness variations of the compact cortex along the calcaneus,
bone tissue was segmented (excluding soft tissue and cavities) using image data reconstructed
with Avizo 9.4 (VSG, Burlington, MA, USA). Then, the outer cortex has been isolated from
the trabecular bone, limited by the inner surface of the cortex for each bone. This segmentation
step was done manually with a combination of Avizo's "remove islands" (to eliminate isolated
volumes that are too small) and "smooth labels" (to connect the slices selection more
realistically) functions to optimize the segmentation repeatability and consistency. Then, the
distances between the inner and outer surfaces of the cortex were calculated in Avizo 9.4 using
the "surface distance" function. Finally, a distance isosurface was obtained with a colour
gradient that appears on the external surface of the bone (Fig. 4; Appendix 4). This colour
gradient showing the relative variation of cortical thickness within each bone, is specific to each
specimen since it varies between the minimum and maximum cortical thicknesses, warmer
colours being used for higher thicknesses, and colder colours for lower thicknesses. Therefore,
two specimens with similar colorimetry may have different absolute cortical thicknesses.

## Quantitative parameters

185 Quantitative parameters used to characterize the internal structure of the bones are: (1)
186 cortex/medullary area volumes, (2) overall bone and trabecular tissue compactness, and average

Fig. 2 Calcaneus of *Sus scrofa,* specimen 2017-578 illustrating the planes of the virtual sections: SS, Sagital section; FS, Frontal section; TS, Transverse section. Db, Dorsal border; Pb, Plantar border. Purple arrows indicate the midpoint of the proximal epiphysis base.

and (3) maximum cortical thicknesses following Houssaye et al. (2021). Most are ratios produced using the volume values obtained from the "material statistics" function in Avizo after the segmentation and cortex isolation steps. The parameters used in the statistical analyses in this study are:

1) whole bone volume in mm$^3$ (WBV), as an indicator of size; 2) bone compactness (C= bone tissue volume*100/WBV); 3) relative trabecular bone tissue fraction (%Trab = trabecular bone tissue volume*100/bone tissue volume); 4) trabecular compactness (Tc= trabecular bone tissue*100/trabecular volume). From the mean

208  (MeanT) and maximum (MaxT) cortical thicknesses that were obtained directly in Avizo 9.4

209  using the 'distance' function were calculated 5) RMaxT and 6) RMeanT, relative maximum and

210  mean thicknesses, by dividing MaxT and MeanT by a mean radius r, obtained from the whole

211  bone volume and considering that the calcanea are cylinders (as v=πr² h, r= √(v /hπ)).

Table. 1 List of material and different parameters used in this study where each row corresponds to a specimen. Ctxt, Context; Prov, Provenance; WBV, total bone volume mm3; C, compactness ratio; %Trab, percentage of trabecular bone volume to cortical bone volume; TC, trabecular compactness; RMeanT, mean relative cortex thickness; RMaxT, maximum relative cortex thickness. Nh, Natural habitat; St, Experimental stall; Co, Experimental control; M, Mesolithic; P, Experimental pen; Comp, Compiègne; Cham, Chambord; Noy2/3, Noyen-sur-seine 2/3; Urc, Urcier.

| Ctxt | Prov | Sex | Age (month) | Weight (kg) | WBV (mm3) | C | %Trab | TC | RMeanT | RMaxT |
|------|------|-----|-------------|-------------|-----------|-----|-------|-----|--------|-------|
| Nh | Comp | F | 10 | 77 | 15252 | 81 | 28 | 57 | 0,08 | 0,34 |
| Nh | Comp | M | 13 | 64 | 18441 | 73 | 43 | 56 | 0,05 | 0,19 |
| Nh | Comp | M | 18 | 113 | 20011 | 83 | 27 | 61 | 0,10 | 0,44 |
| Nh | Comp | F | 12 | 61 | 16112 | 85 | 31 | 67 | 0,09 | 0,39 |
| Nh | Comp | M | 17 | 86 | 20892 | 82 | 35 | 65 | 0,08 | 0,32 |
| Nh | Cham | F | 36 | 52 | 11552 | 81 | 40 | 64 | 0,07 | 0,19 |
| Nh | Cham | M | 72 | 123 | 19932 | 87 | 30 | 68 | 0,10 | 0,31 |
| Nh | Comp | F | 18 | 61 | 14069 | 86 | 35 | 69 | 0,08 | 0,34 |
| Nh | Comp | . | 2 | . | 866 | 82 | 56 | 72 | 0,04 | 0,13 |
| Nh | Cham | F | 17 | . | 8541 | 65 | 51 | 51 | 0,07 | 0,21 |
| Nh | Cham | F | 18 | . | 13125 | 70 | 35 | 49 | 0,07 | 0,28 |
| Nh | Cham | F | 20 | . | 11478 | 78 | 36 | 59 | 0,07 | 0,22 |
| Nh | Cham | F | 18 | . | 13698 | 71 | 41 | 53 | 0,05 | 0,15 |
| Nh | Cham | F | 19 | . | 12480 | 78 | 26 | 50 | 0,09 | 0,17 |
| Nh | Cham | F | 19 | NA | 10870 | 77 | 31 | 55 | 0,08 | 0,29 |
| St | LHT | F | 25 | 91 | 13301 | 73 | 59 | 78 | 0,06 | 0,19 |
| St | LHT | F | 25 | 62 | 9179 | 75 | 42 | 60 | 0,06 | 0,20 |
| St | LHT | F | 25 | 67 | 13416 | 78 | 42 | 62 | 0,06 | 0,18 |
| St | LHT | F | 25 | 89 | 15853 | 73 | 39 | 53 | 0,05 | 0,19 |
| St | LHT | F | 25 | 5 | 15059 | 73 | 39 | 54 | 0,07 | 0,17 |
| St | LHT | F | 25 | 91 | 12971 | 79 | 34 | 59 | 0,08 | 0,30 |
| P | LHT | M | 25 | 84 | 14344 | 75 | 34 | 53 | 0,07 | 0,22 |
| P | LHT | F | 25 | 92 | 15051 | 84 | 36 | 68 | 0,09 | 0,26 |
| P | LHT | F | 25 | 54 | 16201 | 85 | 33 | 69 | 0,09 | 0,36 |
| P | LHT | M | 25 | 67 | 14028 | 76 | 34 | 54 | 0,08 | 0,22 |
| St | LHT | M | 25 | 84 | 20258 | 79 | 49 | 66 | 0,06 | 0,18 |
| St | LHT | M | 25 | 89 | 22432 | 73 | 37 | 57 | 0,07 | 0,22 |
| St | LHT | M | 25 | 78 | 20155 | 81 | 44 | 70 | 0,07 | 0,21 |
| St | LHT | M | 25 | 96 | 22497 | 74 | 41 | 58 | 0,07 | 0,21 |
| St | LHT | M | 25 | 73 | 19239 | 72 | 42 | 55 | 0,06 | 0,18 |
| P | LHT | M | 25 | 91 | 19452 | 82 | 37 | 66 | 0,07 | 0,24 |
| P | LHT | F | 25 | 67 | 15750 | 77 | 33 | 54 | 0,07 | 0,22 |
| P | LHT | M | 25 | 84 | 21659 | 74 | 37 | 55 | 0,08 | 0,27 |
| P | LHT | M | 25 | 90 | 18859 | 71 | 31 | 47 | 0,07 | 0,25 |
| P | LHT | M | 25 | 87 | 19942 | 75 | 37 | 54 | 0,07 | 0,22 |
| P | LHT | M | 25 | 92 | 20783 | 77 | 33 | 56 | 0,08 | 0,26 |
| Co | Urc | M | 20 | 53 | 13733 | 76 | 46 | 61 | 0,05 | 0,19 |
| Co | Urc | M | 8 | 35 | 13711 | 71 | 55 | 60 | 0,04 | 0,17 |
| Co | Urc | M | 20 | 52 | 14446 | 77 | 50 | 64 | 0,05 | 0,22 |
| Co | Urc | F | 84 | 110 | 24561 | 65 | 64 | 56 | 0,04 | 0,14 |
| Co | Urc | F | 96 | 60 | 16299 | 79 | 49 | 66 | 0,06 | 0,19 |
| M | Noy3 | . | . | . | 15154 | 73 | 43 | 55 | 0,04 | 0,27 |
| M | Noy3 | . | . | . | 16109 | 67 | 50 | 54 | 0,05 | 0,23 |
| M | Noy3 | . | . | . | 16144 | 76 | 54 | 79 | 0,07 | 0,26 |
| M | Noy3 | . | . | . | 34372 | 75 | 36 | 55 | 0,06 | 0,26 |
| M | Noy2 | . | . | . | 23577 | 88 | 20 | 64 | 0,13 | 0,48 |
| M | Noy2 | . | . | . | 25740 | 76 | 30 | 51 | 0,08 | 0,26 |

## Statistical analysis

Statistical tests and graphical representations were performed in R (R Core Team. 2017) using the Rstudio software (see appendix 5 and 6 for the script and the data Table). A linear regression model ("lm" function of the "stats" package) as well as a regression coefficient ("cor" function of the "stats" package) were used to assess the linear relationships between the variables (1) whole bone volume (WBV; considered as an estimate of bone size), (2) weight, and (3) age of each individual with all the other quantitative parameters used (Table 2). To explore the distribution of specimens based on their quantitative microanatomical parameters and the variation patterns, we performed a standardized Principal Component Analyses (PCA; "dudi.pca" function of the ade4 package; David and Jacobs 2014).

To estimate the role of factors such as Sex (male/female/indeterminate), origin (Compiègne/ Chambord/ Urcier/ La Haute Touche/ Noyen sur Seine) and mobility status (Natural habitat /Experimental control/ Experimental pen/ Experimental stall/Mesolithic wild boars) in the variation of the quantitative microanatomical parameters, we used analyses of variance (ANOVA; function "anova_test" of the "rstatix" package) after checking the conditions of normality (function "shapiro_test" of the "rstatix" package) and homogeneity of variances (function "levene_test" of the "rstatix" package). When overall difference is significant, we computed pairwise comparison tests using Tukey hsd tests ("tukey_hsd" function of the "rstatix" package). When the variables did not meet the conditions of homogeneity of variances and/or normality, we used the kruskal-Wallis test (function "Kruskal_test" of the "rstatix" package), a non-parametric alternative to ANOVA. When this test is significant, a Dunn's test ("dunn_test (p.adjust.method="bonferroni")" function of the "rstatix" package) is used to compare pairwise differences between the groups concerned.

MANOVA ("res.man" function) was also used to test the overall difference in microanatomical variables between mobility contexts.

## **Results**

### Qualitative descriptions
#### *General microanatomical pattern*

The general microanatomical pattern of the wild boar calcaneus is close to the structure of long bones of terrestrial mammals, with the cortex forming a tubular diaphysis but with a very short diaphyseal part (sections Fig. 3). The proximal epiphysis is not fused to the rest of the bone for the most part of the specimens, except for the oldest. The cortex surrounds cancellous bone (trabecular bone and intertrabecular spaces), which is essentially quite dense, and a small open

medullary cavity, about 1cm (Pradat187; Fig. 3c,h) to 2 cm long (2013-1286; Fig. 3d,i) and 1 cm wide, at the level of the *Sustentaculum tali* (FigXa-j). The thickness of the trabeculae is relatively homogeneous along the calcaneus, except around the medullary cavity, where they are generally thicker (Fig. 3a-j). Also, the trabecular density is heterogeneous, with some specimens having many trabeculae (2017-570; Fig. 3b,g,l) while others have twice as less but thicker trabeculae (2013-1286; Fig. 3 d,I,n). Finally, the bone density in the distal part strongly varies between individuals from compact (2017-570; Fig. 3b,g) to spongious (Calc2139; Fig. 3e,j).

*Sagittal sections*

In all specimens, the dorsal and plantar margins (Db and Pb) show a relatively high cortical thickness, especially at mid-diaphysis. Spongious bone shows anisotropic trabeculae (i.e. with a preferential direction) above the medullary cavity in the bone center (Fig. 3a-e). Anisotropic trabeculae follow the two main directions that are represented with green/outline arrows in the figure 3a and schematized with intersecting lines in figure 1. The cortical thicknesses of the plantar and dorsal margins vary from thick (1 cm in specimen 2017-570; Fig. 3b) to proportionally twice as thin (0.5 cm in Pradat187; Fig. 3c). A variation in the cortical thickness of the plantar border of the proximal epiphysis is also observed, it is very thin (1mm) in some specimens like Pradat 187 (Fig. 3c) while others, such as 2017-555 (Fig. 3a), show a clear thickening of the cortex (2mm) in this area. Similarly, bone is highly compact next to the fibular trochlea for numerous specimens (Calc2139; Fig. 3e) or rather spongious for some individuals (2013-1286; Fig. 3d).

*Frontal sections*

All individuals have a cortex that varies relatively little in thickness (about 2-3mm) on the medial and lateral sides (Fig. 3f-e). In contrast, the cortical thickness in the proximal epiphysis varies from thin (0.5 cm in specimen 2013-1286; Fig. 3i) to twice thicker (0.3 cm in specimen 2017-555; Fig. 3f). Similarly, the compactness and the cortical thickness of the *sustentaculum tali* varies between individuals, some of which show compact bone and thick cortex (Padat187; Fig. 3h) while others show spongious bone and thin cortex (2013-1286; Fig. 3i).

*Transverse sections*

The transverse sections' shapes are generally oval and elongate (2017-555; Fig. 3k) but several specimens present a rounder section (2017-570; Fig. 3l). The cortical thickness is fairly constant

280    across the sections, but some specimens show a cortical thickening at the plantar border (2013-
281    1286; Fig. 3n).

283    None of the variation observed between specimens is clearly qualitatively associated to any
284    main parameter of the study, namely the context, provenance, sex, size or weight.

Fig. 3 Virtual thin sections of the calcaneus of boars, a,f,k, 2017-555 (Experimental stall); b,g,l 2017-570 (Experimental pen); c,h,m Pradat187 (Experimental control); d,i,n, 2013-1286 (Natural habitat); e,j,o, Calc2139 (Mesolithic). Db, Dorsal border; De, Distal end; Ft, Fibular trochlea; Pb, Plantar border; Pe, Proximal epiphysis; St, *Sustentaculu tali*. Scale bars of sagittal and frontal sections equal 2 cm; Scale bars of transverse sections equal 1 cm.

## 3D mapping of the cortical thickness

The 3D mappings of the cortical thickness and its variation across each bone, are quite similar between specimens (Appendix 1). In agreement with the observation of the virtual sections, there is a fairly extensive area with greater cortical thickness at the plantar border of the calcaneus (Pb; Fig. 4A). Although generally less extensive, thickenings of the cortex can also be noted on the dorsal margin (Db) and on the dorsal base of the *sustentaculum tali* (Sb; Fig. 4B). There is little thickening of the cortex on the proximal end (Pe). Finally, there is no lateral or medial thickening noted.

Twenty-five of the 47 specimens show strong cortical thickening at the plantar margin (Pb; Appendix 1), like in 2017-568 (Fig. 4A); this is less visible in the 22 others, like Pradat 185 and 2013-1264 (Fig. 4D; Fig. 4G). Twenty-three specimens have thick cortical bone at the dorsal margin (Db; Table 1; Appendix 1), like in Pradat 185 and 2017-568 boars (Fig. 4B and Fig. 4E), whereas the others do not show such thickening like the calcaneus of boar 2013-1264 (Fig. 4H). Twenty-height specimens show a relative thickening at the base of the sustentaculum (Sb; Table 1), this widening is particularly noticeable in boar 2013-1264 (Fig. 4H) and absent in others like Pradat 185 (Fig. 4E). A slight recurrent cortical thickening on the distal end is observed in part of the specimens like 2017-568 (Fig. 4a). Twenty-seven specimens have a slightly thicker cortex on the proximal end (Pe) such as 2017-568 (Fig. 4C), 15 show no or very little thickening, such as Pradat185 and 2013-1264, respectively (Fig. 4F and Fig. 4I). Four of the proximal ends of the Mesolithic specimens were not found because they were not fused to the rest of the bone.

Fig. 4 3D mappings of the *Sus scrofa* calcaneus relative cortical thickness. A-C specimen 2017-568; D-F Pradat 185; G-I 2013-1264. Anatomical abbreviations: Db, dorsal border; Ls, lateral side; Ms, medial side; Pb, plantar border; Pe, proximal epiphysis; Sb, Sustentaculum base; St, *Sustentaculum tali*. D, A and G are in plantar view; E, B and H are in dorsal view; F and I and C are in posterior view. Abbreviations for orientations, prox, proximal; lat, lateral; med, medial; post, posterior; ant, anterior.

## Quantitative analyses

### *Microanatomical covariation with Weight, age size and sex.*

The Whole bone volume (WBV) is expectedly only correlated to the body weight and the PCA axes (Table. 2). Only %Trab is significantly correlated with specimen age with a slight increase in the proportion of trabecular bone over cortical bone as boars get older (r=0.32). Variables C, %Trab, RMeanT, RMaxT, weight, and TC did not differ between males and females. Only whole calcaneus volume (WBV) variation differs with sex (Kruskal Wallis: $p<0.01$) with males larger than females (Fig. 6). Mesolithic specimens (sex unknown) have larger calcanei than present-day males (Dunn's test: $p<0.01$).

Table. 2 Values obtained for tests of the effect of total volume, age, and weight on the different parameters and the PCA axes. r: correlation coefficient; p: p-value of the linear regression model; WBV: whole bone volume; RMeanT, mean relative cortical thickness; RMaxT, maximum relative cortical thickness; PC1, PC2: position of individuals on the first two axes of the PCA.

| | WBV | Age | Weight | C | %Trab | TC | RMeanT | RMaxT | PC1 | PC2 |
|---|---|---|---|---|---|---|---|---|---|---|
| WBV | X | X | X | r=0,07 | r=-0,13 | r=-0,05 | r=0,13 | r=0,20 | **r=-0,46** | **r=0,36** |
| n=46 | | | | p=0,63 | p=0,38 | p=0,75 | p=0,37 | p=0,18 | **P<0,01** | **p=0,01** |
| Age | r=0,30 | X | X | r=0,47 | **r=0,32** | r=0,17 | r=-0,19 | r=-0,20 | r=0,07 | r=0,25 |
| n=40 | P=0,06 | | | p=0,92 | **p=0,04** | p=0,29 | p=0,23 | p=0,21 | p=0,65 | p=0,12 |
| Weight | **r=0,56** | r=0,27 | X | r=0,08 | r=-0,16 | r=0,02 | r=0,30 | r=0,26 | r=0,34 | r=0,31 |
| n=34 | **P<0,01** | P=0,12 | | p=0,65 | p=0,35 | p=0,93 | p=0,08 | p=0,14 | p=0,05 | p=0,07 |

### *Patterns of calcanei microanatomical variations and mobility contexts*

Specimen 2013-1287, corresponding to the youngest individual (2 months), was removed before performing PCA (n=46; Fig. 5) and other quantitative analyses. Axes 1 and 2 of the PCA explain 75.3% of the total variance. Furthermore, contribution of the variables to the axes (Fig. 5), show that RMeanT and RMaxT covary while WBV and %Trab vary in opposite ways. The first axis is influenced by the cortical thickness (TC), %Trab and C, while only TC greatly structures the second axis. WBV is correlated with the PCA axes whereas age, weight and the microanatomical parameters are not (table 2). The different mobility contexts induce significant microanatomical differences (MANOVA: p-value<0.01). On the PCA (figure 5), experimental penned, stalled, and control wild boars are quite distinct along PC1, whereas the Mesolithic specimens of Noyen-sur-Seine covers almost all variation along PC1. Differences across the locomotor contexts are observed for RMeanT (Kruskal Wallis: p<0.01), RMaxT (Kruskal Wallis: p<0.01) and %Trab (Kruskal Wallis: p<0.01). RMeanT differences are significant

Fig. 5 Distribution of the 46 specimens and their associated context groups on the first two axes of the PCA computed on: WBV, total bone volume; RMeanT, relative mean bone thickness; RMaxT, relative maximum bone thickness; C, compactness ratio; TC, trabecular compactness ratio; %Trab, percentage of trabecular bone volume to cortical bone volume. meso, Mesolithic; nat hab, Natural habitat.

between wild boars from the Natural habitat and the Experimental control groups (Dunn test:

p<0.01) and between Experimental control and Experimental pen groups (Dunn's test: p<0.01).

RMaxT is significantly different between Experimental control and Mesolithic wild boars

(Dunn's test: p<0.01). %Trab is significant different between Experimental control and natural

habitat (Dunn's test: p<0.01) and between Experimental pen and control (Dunn's test: p<0.01).

Thus, the pen-reared boars have a higher average cortex thickness (RMeanT) than the control

group, for which, conversely, it is the percentage of trabecular bone (%Trab) that is higher (Fig.

Fig. 6 Calcaneus microanatomical variation in *Sus scrofa* from experimental populations and Mesolithic contexts. WBV, whole bone volume; RMeanT, relative mean cortical thickness; RMaxT, relative maximum cortical thickness; C, compactness ratio; TC, trabecular compactness ratio; %Trab, percentage of trabecular bone volume to cortical bone volume. meso, Mesolithic; nat hab, Natural habitat.

6). The stall group had intermediate RMeanT, RMaxT and %Trab values compared to the pen

and control groups. In addition, there is a generally greater standard deviation in the Natural

habitat and Mesolithic groups for all parameters, except TC, for which individuals in the stall

group are slightly more dispersed than those in the Natural habitat group. On the other hand, no

group is distinguished along PC2, the variables C and TC did not vary according to context

(ANOVA; p-value C =0.39; p-value TC =0.82; Fig. 6); Finally, the whole volume (WBV) of

the calcaneus differed slightly between groups (ANOVA: p=0.04) but only the difference

between hunted Mesolithic and modern wild boars in their natural habitat was significant

(Tukey hsd test: p=0.02; Fig. 6), with Mesolithic individuals being larger.

## Discussion

### 1.Overall calcaneal microanatomy in wild and captive wild boars

The qualitative description of the 3D maps and virtual thin sections identified a strong cortical

thickening on the plantar and dorsal borders (Pb and Db; Fig. 4). These observations are

consistent with the main constraints identified in the calcaneus of artiodactyls with significant

compression, bending, and tension forces on the plantar and dorsal borders (Lanyon 1973; Fig.

1). Thickenings of the plantar and dorsal margins have also been identified for wild deers

(Skedros et al., 2001) and pasture-raised domestic sheeps (Skedros et al., 2007). Moreover, the

anisotropric properties of the boar specimens trabeculae to follow an antero-posterior

orientations (outline arrows Fig. 3a) are congruent with the distribution of constraints

mentioned above, as internal bone structure organized itself to better respond to stress (Wolff,

1986; Ruff et al., 2006; Van der Meulen et al., 2006).

Although the entheses (i.e., areas of ligament and tendon insertion (Djukic et al., 2015)) are

regions of stress concentration, these areas show little or no effect on the microanatomy of boar

calcaneus. The loads applied to the calcaneus of artiodactyls are primarily shared between the

long plantar ligament and the Achilles tendon, which connects the calcaneus to the

gastrocnemius and soleus muscles, the latters forming the sural triceps (Lanyon,1973; Woo et

al., 1981; Skedros et al., 2001; Skedros et al., 2007; Barone, 2020). However, while the long

plantar ligament attaches all along the plantar border, only a part of this edge is thickened in

the boar calcaneus. Furthermore, an important thickening of similar proportion is also observed

on the dorsal edge, whereas this bone side shows no enthesis. The Achilles tendon insertion at

the proximal end of the calcaneus is itself covered by a tendinous structure, the calcaneus cap

of the superficial flexor digitorum muscle (*m. flexor superficialis*), taking an attachment point

from the tip of the proximal end of the calcaneus to phalanges (Su et al., 1999; Barone, 2017;

Fig. 1).  A slight cortical thickening on the proximal epiphysis of our specimens coincides with

the insertion of the tendons of the *m. flexor superficialis* muscles of the toes and of the *m.*

*gastrocnemius* muscles (Bénévent & Bressot, 1968; Barone 2017).

Regarding the articular surfaces, the *sustentaculum tali* is a strong protuberance on the medial

side forming an articular surface with the talus, the fibular trochlea is an articular surface for

the malleolar bone and the end of the distal part articulate with the cuboid bone (Barone, 2017),

recurrent slight cortical thickening is also observed in those regions.

Overall, the microanatomy seems to reflect the tension and compression forces with a strong

cortical thickening on Pb and Db as well as the anisotropy of the trabeculae. The areas of contact

with other bones are also represented with important bone density on the fibular trochlea, the

*sustentaculum tali* and the end of the distal part.

## 2.Intra individual variation in calcaneal Microanatomy

Beyond the general pattern, no clear relationship is observed between the variability of the

microanatomical parameters and the factors explored in this study (context, sex, weight, size).

However, we found important inter-individual differences, notably in the extent and depth of

the cortical thickness of the plantar and dorsal edges, and on the proximal epiphysis; the length

of the medullary cavity; the number and thickness of the trabeculae; the bone tissue density of

the *sustentaculum tali*, at the distal part of the bone and next to the fibular trochlea; and the

transverse sectional shape. This suggests that other factors influencing bone development

during growth must be explored to further understand inter-individual disparity.

Only a few correlations (positive or negative) of the microanatomical variables with bone size,

weight, and age of the individuals (Table 2), are significant. These three parameters, therefore,

have a limited impact on the microanatomical organisation. Body weight is only correlated with

bone size (WBV). WBV influences the PCA axes (Table 2) while, paradoxically, bone size is

not directly correlated with any of the microanatomic variables in isolation, which is congruent

with the observations of the sections and 3D maps that have not found any link between

specimens' size and their microanatomy. Thus, it is the covariation between the variables that

are themselves weakly correlated with whole volume that makes the relationship between whole

volume and all variables significant. Nevertheless, although age is not one of the parameters on

which this study focuses, a weak correlation was found between age and the trabecular

percentage (%Trab; r=0.32), but trabecular percentage is not significantly correlated with bone

size since age and bone size are neither significantly correlated. The increase in the proportion

of trabecular bone tissue with age is not related to an increase in trabecular compactness (TC)

because this parameter is not significantly correlated with age (p=0.29); the volume of the

medullary zone tends to increase since the cortex becomes proportionally thinner. This result is

surprising because the opposite phenomenon occurs in the calcaneus of deer (Skedros et al.,

2001) and sheep (Skedros et al., 2007), where a thickening of the cortex is observed with size
in relation to the medullary zone. Conversely, while it is surprising that age does not correlate
with whole bone volume (WBV; Table2), this shows that age and volume do not follow a linear
relationship or that intraspecific variability in calcaneus size between specimens exceeds the
effect of growth. However, our sample does not adequately test the relationship between the
variables and ontogeny because of the large proportion of individuals of the same age (25
months).

### 3.Change in mobility regime and calcaneal Microanatomy

Despite the lack of directly observable influence of the mobility context over the calcaneal
microanatomy, we found quantitative effect of mobility differences in the cortical thickness
(RmeanT and RmaxT), trabecular percentage (%Trab) and in the overall variation patterns
(PCA), indicating that difference in mobility context influences the microanatomical
characteristics of the calcaneus, although not in a strongly discriminatory pattern. However, we
didn't find the expected microanatomical proximity of wild boars living in their natural habitat
and their dissimilarity from wild boars kept in captivity, as seen in previous studies on the
calcaneus 3D externalshape and form (Harbers et al.2020). The variations in relative mean
cortical thickness (RMeanT) illustrates the general trend of variations related to the mobility
regime (Fig. 6). While we expected to observe similar average thicknesses between animals
that had similar mobility conditions (e.g. Natural habitat and Experimental control), we found
that control individuals have a much lower cortical thickness than wild individuals from natural
habitat. Thus, modern and Mesolithic wild boars hunted in their natural habitat display similar
microanatomy with the wild boars which grew in a very small living space (enclosure of 4m2
in a hangar of 100m2.). The most divergent microanatomy from the wild boar norm of reaction
have been observed in the control populations from the wild boar farm of Urcier, which have a
much lower cortical thickness than wild boars in their natural habitat. All these results suggest
that the microanatomy of the *Sus scrofa* calcaneus does not strongly reflect the mobility regime,
contradicting the strong microanatomical signal associated with locomotor restriction
evidenced in battery chickens, including osteoporosis related to inactivity (Rath et al., 2000).
However, the locomotor restrictions of these reared animals are generally greater than those
imposed on the animals of this study.

## 4. Greater microanatomical variability in modern and ancient wild boars hunted in their natural habitat

The six Mesolithic calcanei from the archaeological site of Noyen-sur-Seine show similarity for all parameters and variables with the other groups (Fig. 6). Thus, the Mesolithic individuals share the same micronatomy with their modern relative. We also found greater microanatomical variability in wild boars populations both modern and Mesolithic. External factors were much less controlled than for the DOMEXP groups, thus resulting in more elements to affect bone plasticity. In addition, the animals from Compiègne and Chambord (Natural habitat) and NoyenEns2 and NoyenEns3 (archaeology) are different populations so they have greater genetic variability in comparison to the DOMEXP group. Furthemore, the Mesolithic specimens belong to individuals from multiple generations separated by several centuries. This genetic variability is much a substantial factor that has a greater impact on the observed phenotype than the intrapopulation variability related to motricity in this study. Consequently, when several populations are included in the same group (Natural habitat and Mesolithic), their variability exceeds that observed between the same population placed in different locomotor contexts (stall, pen, control). Thus, the microanatomy of the calcaneus appears to be more affected by population differences than by the locomotor context in which the animals grew.

In addition to this explanation, the wider locomotor regime of wild boars in their natural habitat would foster greater ecophenotypic variation compared to captive specimens with reduced mobility and more stereotypical locomotor behaviour. One the one hand, wild boars in nature have a locomotor repertoire that must respond to several problems that are not encountered in captivity, such as foraging or escape (Spitz & Janeau, 1990, 1995). Their daily travel is generally less than 10km but it could be up to 80 km in one night (Keuling et al., 2009), peak speeds of 40 km/h and high jumps up to 1.5 m have also been observed (Baskin & Danell, 2003). One the other hand, there is no study describing the skeletal repercussion of stereotypic behaviours. However, because these types of behaviours performed to compensate for lack of activity are induced during significant psychological depressions in individuals (Rushen, 1993; Andre, 2007), other experimental approaches involving living specimens are not desirable.

quantifying

Also, diet plays a role in the development of the skeleton (Randoin & Causeret, 1945). Experimental specimens were fed nutritionally balanced pellets (15% protein) intended for pig breeding to ensure consistent growth and bone formation. In their natural habitat, food availability, seasonal, and geographical variations are major factors influencing food selection

by wild boars (Ballari, & Barrios-García, 2014), Thus, the diet factor possibly also contributes to the high variability observed in wild (free-ranging) individuals. Furthemore, it is likely that there are differences in substrates between the forests and areas in which these boars lived; this may have important implications for the autopod (Kappelman, 1988). Indeed, the substrate factor seems potentially important for the stall-reared group where the flat ground, was covered with moss and straw mats, whereas the more irregular natural terrain of wild boars (like control group) implies variable and multidirectional soil reaction forces (Hanot et al., 2017). The group of pens were raised on a flat terrain covered with grass and a few trees.

The overview of the different studies related to the DOMEXP project show that bone plasticity associated with domestication varies between bones. The most surprising result of the present study is that the bone plasticity of the external morphology of wild boar calcaneus' is more variable than its microanatomy, although the latter is considered to be more plastic than bone external morphology (Dumont et al., 2013; Kivell, 2016; Vlachopoulos et al., 2017). Since the results of the study show that boars living in different contexts can have the same microanatomical pattern, calcaneal microanatomy cannot be used to infer a captive lifestyle. However, a strong diversity is noted in the microanatomy between wild and Mesolithic specimens. A better understanding of the factors that regulate calcaneal variability would possibly allow inferences related to habitat (type of soil, open or closed environment) or diet.

## Aknowledgements

We thank the reviewers Ignacio Aguilar Lazagabaster and Max Price as well as our recommender Nimrod Marom for their helpful and positive comments that have helped improve this work. Thank to Florian Bogaert, who helped us to realize the segmentation of a part of specimens. Preprint version X of this article has been peer-reviewed and recommended by Peer Community In Archaeology (https://XXX link to the article).

## Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article. Alexandra Houssaye is recommender of PCI Paleontolgy.

## Funding

This research has been funded by the project Emergence SU-19-3-EMRG-02. This research has also benefited from financial supports of the Muséum national d'Histoire naturelle (Paris) and the CNRS INEE (Institut écologie et environnement).
This research was funded by ANR through the DOMEXP project (ANR\u000213-JSH3-0003-01), the LabEx ANR-10-LABX-0003-BCDiv, in the programme 'Investissements d'avenir' ANR-11-IDEX-0004-02, programme Emergence SU-19-3-EMRG-02.

## Bibliography

Agarwal, S.C., 2016. Bone morphologies and histories: Life course approaches in bioarchaeology. American Journal of Physical Anthropology. Am. J. Phys. Anthropol., 159: 130-149. URL https://onlinelibrary.wiley.com/doi/full/10.1002/ajpa.22905

Andre, J.P., 2007. An introduction to normal and pathological behaviour in psittacine. Bulletin de l'Académie Vétérinaire de France, 2007 vol.160, N°3 https://doi.org/10.4267/2042/47885

Barone, R., 2000. Anatomie comparée des mammifères domestiques Tome 2 -Arthologie et myologie- (4eéd.). Association centrale d'entraide vétérinaire.

Barone, R., 2017. Anatomie comparée des mammifères domestiques Tome 1 - ostéologie – (5e éd.). Vigot.

Baskin, L., Danell K., 2003 Ecology of ungulates: a handbook of species in Eastern Europe and Northern and Central Asia. Heidelberg, Germany:Springer Science & Business Media.

Bassarova, M., Janis, C.M., Archer, M., 2009. The Calcaneum—On the Heels of Marsupial Locomotion. J Mammal Evol 16, 1–23. https://doi.org/10.1007/s10914-008-9093-7

Bénévent, M., Bressot, C., 1968. Guide pour la dissection et l'identification des principaux muscles de la carcasse chez le mouton. Annales de biologie animale, biochimie, biophysique 8, 147–193.

Binford, L.R., 1978. Nunamiut: ethnoarchaeology. New York, NY, Academic Press.

Bleefeld, A., Bock, W., 2002. Unique anatomy of lagomorph calcaneus. Acta Palaeontologica Polonica, 47, 181–183

Carmeliet, G., Bouillon, R., 2001. Space flight: a challenge for normal bone homeostasis. Crit. Rev. Eukar. Gene 11, 1–3.

Carrano, M.T., 1997. Morphological indicators of foot posture in mammals: a statistical and biomechanical analysis. Zool. J. Linn. Soc. 121, 77–104. doi:10.1111/j.1096-3642.1997.tb00148.x

Carter, D.R., Beaupré, G.S., 2001. Skeletal Function and Form. Mechanobiology of Skeletal Development, Aging and Regeneration. Cambridge University Press, Cambridge.

554     Cowin, S.C., Hegedus, D.H., 1976. Bone remodeling I : a theory of adaptive elasticity. J. Elasticity 6,
555         313–326.
556
557     Cubo, J., Ponton, F., Laurin, M., De Margerie, E., Castanet, J., 2005. Phylogenetic Signal in Bone
558         Microstructure of Sauropsids. Systematic Biology 54, 562–574.
559         https://doi.org/10.1080/10635150591003461
560
561     Cucchi, T., Domont, A., Harbers, H., Leduc, C., Guidez, A., Bridault, A., Hongo, H., Price, M., Peters,
562         J., Briois, F., Guilaine, J., Vigne, J.D., 2021. Bones geometric morphometrics illustrate 10th
563         millennium cal. BP domestication of autochthonous Cypriot wild boar (Sus scrofa circeus nov.
564         ssp). Sci Rep 11, 11435. https://doi.org/10.1038/s41598-021-90933-w
565
566     Curran, S.C., 2012. Expanding ecomorphological methods: geometric morphometric analysis of
567         Cervidae post-crania. J. Archaeol. Sci. 39, 181-184
568
569     David, C.C., Jacobs, D.J., 2014. Principal Component Analysis: a method for determining the essential
570         dynamics of proteins. In: Livesay DR (ed) Protein Dynamics: Methods and Protocols. Humana
571         Press, Totowa, pp 193–226
572
573     Divé, J., Eisenmann, V., 1991. Identification and discrimination of first phalanges from Pleistocene
574         and modern Equus, wild and domestic. In: Meadow, R.,Uerpmann, H.-P. (Eds.), 1991. Equids in
575         the Ancient World, vol. II. Ludwig Reichert Verlag, Wiesbaden, pp. 278-333
576
577     Djukic, K., Milovanovic, P., Hahn, M., Busse, B., Amling, M., Djuric, M., 2015. Bone
578         microarchitecture at muscle attachment sites: The relationship between macroscopic scores of
579         entheses and their cortical and trabecular microstructural design. American Journal of Physical
580         Anthropology 157, 81–93. https://doi.org/10.1002/ajpa.22691
581
582     Du, T.Y., Standen, E.M., 2020. Terrestrial acclimation and exercise lead to bone functional response
583         in Polypterus senegalus pectoral fins. Journal of Experimental Biology 223.
584         https://doi.org/10.1242/jeb.217554
585
586     Dumont, M., Laurin, M., Jacques, F., Pellé, E., Dabin, W., de Buffrénil, V., 2013. Inner
587         architecture of vertebral centra in terrestrial and aquatic mammals: a two-dimensional
588         comparative study. Journal of Morphology 274, 570–584.
589
590     Eisenmann, V., 1986. Comparative osteology of modern and fossil horses, half-asses, and asses. In:
591         Meadow, R.H., Uerpmann, H.-P. (eds.), Equids in the Ancient World. Ludwig Reichert Verlag,
592         Wiesbaden, pp. 68-116.
593
594     Eisenmann, V., Beckouche, S., 1986. Identification and discrimination of metapodials from Pleistocene
595         and modern Equus, wild and domestic. In: Meadow, R.H.,Uerpmann, H.-P. (Eds.), Equids in the
596         Ancient World. Ludwig Reichert Verlag, Wiesbaden, pp. 117-163.
597
598     Eisenmann, V., 1995. What metapodial morphometry has to say about some Miocene Hipparions. In:
599         Vrba, E.S., Denton, G.H., Partridge, T.C., Burckle, L.H. (Eds.), Paleoclimate and Evolution, with
600         Emphasis on Human Origins. Yale University Press, New Haver, pp. 148-164.
601
602     Frost, H.M., 1983. A determinant of bone architecture. The minimum effective strain. Clin Orthop
603         Relat Res 286–292.
604

Frost, H.M., 1994. Wolff's Law and bone's structural adaptations to mechanical usage: an overview for clinicians. The Angle Orthodontist 64, 175–188. https://doi.org/10.1043/0003-3219(1994)064<0175:WLABSA>2.0.CO;2

Fyhrie, D.P., Carter, D.R., 1986. A unifying principle relating stress to trabecular bone morphology. Journal of Orthopaedic Research 4, 304–317. https://doi.org/10.1002/jor.1100040307

Hall, B. K., 1983. Epigenetic control in development and evolution, p. 353–379. In: Development and Evolution. B. C. Goodwin, N. Holder, and C. C. Wylie (eds.). Cambridge University Press, Cambridge.

Hall, B. K., 1989. Genetic and epigenetic control of vertebrate embryonic development. Netherlands Journal of Zoology 40:352–361.

Hall, R.L., Shereff, M.J., 1993 Anatomy of the calcaneus. Clin. Orthop. Relat. Res. 290, 27–35.

Hall, B.K., 2005. Bones And Cartilage: Developmental And Evolutionary Biology. Academic Press Inc, Australia ; San Diego, Calif.

Hanot, P., Herrel, A., Claude, G., Cornette, R., 2017. Morphological integration in the appendicular skeleton of two domestic taxa: The horse and donkey. Proceedings of the Royal Society B: Biological Sciences 284, 20171241. https://doi.org/10.1098/rspb.2017.1241

Harbers, H., Neaux, D., Ortiz, K., Blanc, B., Laurens, F., Baly, I., Callou, C., Schafberg, R., Haruda, A., Lecompte, F., Casabianca, F., Studer, J., Renaud, S., Cornette, R., Locatelli, Y., Vigne, J.D., Herrel, A., Cucchi, T., 2020a. The mark of captivity: plastic responses in the ankle bone of a wild ungulate (Sus scrofa). R Soc Open Sci 7:192039. https://doi.org/10.1098/rsos.192039

Harbers, H., Zanolli, C., Cazenave, M., Theil, J.-C., Ortiz, K., Blanc, B., Locatelli, Y., Schafberg, R., Lecompte, F., Baly, I., Laurens, F., Callou, C., Herrel, A., Puymerail, L., Cucchi, T., 2020b. Investigating the impact of captivity and domestication on limb bone cortical morphology: an experimental approach using a wild boar model. Scientific Reports 10, 19070. https://doi.org/10.1038/s41598-020-75496-6

Hegedus, D.H., Cowin, S.C., 1976. Bone remodeling II: small strain adaptive elasticity. J. Elasticity 6, 337–352

Houssaye, A., Taverne, M., Cornette, R., 2018. 3D quantitative comparative analysis of long bone diaphysis variations in microanatomy and cross-sectional geometry. J Anat 232:836–849

Huiskes, R., Ruimerman, R., van Lenthe, G.H., Janssen, J.D., 2000. Effects of mechanical forces on maintenance and adaptation of form in trabecular bone. Nature 405, 704–706. https://doi.org/10.1038/35015116

Hussain, S., 1975. Evolutionary and Functional Anatomy of the Pelvic Limb in Fossil and Recent Equidae (Perissodactyla, Mammalia). Anatomia, Histologia, Embryologia: Journal of Veterinary Medicine Series C, 4, 193–222. doi:10.1111/j.1439-0264.1975.tb00637.x

Kappelman, J., 1988. Morphology and locomotor adaptations of the bovid femur in relation to habitat. Journal of Morphology, 198, 119–130. https://doi.org/10.1002/jmor.1051980111

Kelly, S.A., Czech, P.P., Wight, J.T., Blank, K.M., Garland, T., 2006. Experimental evolution and phenotypic plasticity of hindlimb bones in high-activity house mice. Journal of Morphology 267, 360–374. https://doi.org/10.1002/jmor.10407

Keuling, O., Stier, N., Roth, M., 2009. Commuting, shifting or remaining?: Different spatial utilisation
patterns of wild boar Sus scrofa L. in forest and field crops during summer. Mamm. Biol. 74,
145–152. (doi:10.1016/j.mambio.2008.05.007)

Kivell, T.L., 2016. A review of trabecular bone functional adaptation: what have we learned from
trabecular analyses in extant hominoids and what can we apply to fossils? Journal of Anatomy
228, 569–594. https://doi.org/10.1111/joa.12446

Krølner, B., Toft, B., 1983. Vertebral Bone Loss: An Unheeded Side Effect of Therapeutic Bed Rest.
Clinical Science 64, 537–540. https://doi.org/10.1042/cs0640537

Lang, T., LeBlanc, A., Evans, H., Lu, Y., Genant, H., Yu, A., 2004. Cortical and Trabecular Bone
Mineral Loss From the Spine and Hip in Long-Duration Spaceflight. Journal of Bone and Mineral
Research 19, 1006–1012. https://doi.org/10.1359/JBMR.040307

Lanyon, L.E., 1973. Analysis of surface bone strain in the calcaneus of sheep during normal
locomotion: Strain analysis of the calcaneus. Journal of Biomechanics 6, 41–49.
https://doi.org/10.1016/0021-9290(73)90036-5

Lanyon, L.E., 1992. Control of bone architecture by functional load bearing. Journal of Bone and
Mineral Research 7, S369–S375. https://doi.org/10.1002/jbmr.5650071403

Laros, G.S., Tipton, C.M., Cooper, R.R., Stimmel, P., Davis, R., Matthes, R.D., 1971. Influence of
Physical Activity on Ligament Insertions in the Knees of Dogs. JBJS 53, 275–286.

Lieberman, D.E., Devlin, M.J., Pearson, O.M., 2001. Articular area responses to mechanical loading:
effects of exercise, age, and skeletal location. Am. J. Phys. Anthropol. 116, 266-277.

Lovejoy, C.O., Cohn, M.J., White, T.D., 1999. Morphological analysis of the mammalian postcranium:
a developmental perspective. Proc. Nat. Acad. Sci. U.S. A. 96, 13247e13252

Maïmoun, L., Sultan, C., 2011. Effects of physical activity on bone remodeling. Metabolism 60, 373–
388. https://doi.org/10.1016/j.metabol.2010.03.001

Maïmoun, L., Coste, O., Philibert, P., Briot, K., Mura, T., Galtier, F., Mariano-Goulart, D., Paris, F.,
Sultan, C., 2013. Peripubertal female athletes in high-impact sports show improved bone mass
acquisition    and    bone    geometry.    Metabolism    62,    1088–1098.
https://doi.org/10.1016/j.metabol.2012.11.010

Marcus R., 2002. Chapter 86 - Mechanisms of Exercise Effects on Bone. In Principles of bone biology
(second edition) (eds JP Bilezikian, LG Raisz, GA Rodan), pp. 1477–1488. San Diego, CA:
Academic Press

Marinval-Vigne, M.C., Mordant, D., Auboire, G., Augereau, A., Bailon, S., Dauphin, C., Delibrias, G.,
Krier, V., Leclerc, A.S., Leroyer, C., Marinval, P., Mordant, C., Rodriquez, P., Vilette, P., Vigne,
J.-D., 1989. Noyen-sur-Seince, site stratifié en milieu fluviatile: Une étude multidisciplinaire
intégrée. Bulletin de la Société préhistorique française 86, 370–379.

Martin,   R.B.,   2000.   Toward   a   unifying   theory   of   bone   remodeling.   Bone   26,   1–6.
https://doi.org/10.1016/S8756-3282(99)00241-0

Modlesky, C.M., Majumdar, S., Dudley, G.A., 2008. Trabecular bone microarchitecture in female
collegiate gymnasts. Osteoporos Int 19, 1011–1018. https://doi.org/10.1007/s00198-007-0522-x

713
714 Mordant, D., Boris, V., Vigne, J.D., 2013. Noyen-sur-Seine, vingt-cinq ans après. - Société
715     Préhistorique française, Séance 2, p. 37-50.
716
717 Neaux, D., Blanc, B., Ortiz, K., Locatelli, Y., Laurens, F., Baly, I., Callou, C., Lecompte, F., Cornette,
718     R., Sansalone, G., Haruda, A., Schafberg, R., Vigne, J.D., Debat, V., Herrel, A., Cucchi, T., 2021.
719     How Changes in Functional Demands Associated with Captivity Affect the Skull Shape of a Wild
720     Boar (Sus scrofa). Evol Biol 48, 27–40. https://doi.org/10.1007/s11692-020-09521-x
721
722 Newman, S., Leeson, S., 1999. The effect of dietary supplementation with 1,25-
723     dihydroxycholecalciferol or vitamin C on the characteristics of the tibia of older laying hens.
724     Poultry Science 78, 85–90. https://doi.org/10.1093/ps/78.1.85
725
726 Noyes, M.D., Frank, DELUCAS, J., Torvik, P., 1974. Biomechanics of Anterior Cruciate Ligament
727     Failure: An Analysis of Strain-Rate Sensitivity and Mechanisms of Failure in Primates. The
728     Journal of bone and joint surgery. American volume 56, 236–53.
729     https://doi.org/10.2106/00004623-197456020-00002
730
731 Palencia, P., Vicente, J., Barroso, P., Barasona, J., Soriguer, R.C. and Acevedo, P., 2019. Estimating
732     day range from camera-trap data: the animals' behaviour as a key parameter. J Zool, 309: 182-
733     190. https://doi.org/10.1111/jzo.12710
734
735 Parfitt, A.M., 2002. Targeted and nontargeted bone remodeling: Relationship to basic multicellular unit
736     origination and progression. Bone 30, 5–7.
737
738 Pearson, O.M., Lieberman, D.E., 2004. The aging of Wolff's 'law': Ontogeny and responses to
739     mechanical loading in cortical bone. Am. J. Phys. Anthropol. 125, 63–99.
740     (doi:10.1002/ajpa.20155)
741
742 Pelletier, M., Kotiaho, A., Niinimäki, S., Salmi, A.-K., 2020. Identifying early stages of reindeer
743     domestication in the archaeological record: a 3D morphological investigation on forelimb bones
744     of modern populations from Fennoscandia. Archaeol Anthropol Sci 12, 169.
745     https://doi.org/10.1007/s12520-020-01123-0
746
747 Pinhasi, R., Stock, J.T., 2011. Human Bioarchaeology of the Transition to Agriculture. John Wiley &
748     Sons.
749
750 R Core Team. 2017 R: A language and environment for statistical computing. Vienna, Austria: R
751     Foundation for Statistical Computing. https://www.R-project.org.
752
753 Randoin, L., Causeret, J., 1945. Experiments on the effect of the phosphorus, calcium, magnesium and
754     vitamin D contents of the diet on bone development in the white rat. Bull: Soc. sci. Hyg. aliment.
755     33, 134–143.
756
757 Rath, N.C., Huff, G.R., Huff, W.E., Balog, J.M., 2000. Factors Regulating Bone Maturity and Strength
758     in Poultry1. Poultry Science 79, 1024–1032. https://doi.org/10.1093/ps/79.7.1024
759
760 Robling, A.G., Castillo, A.B., Turner, C.H., 2006. Biomechanical and molecular regulation of bone
761     remodeling. *Annual Review of Biomedical Engineering*, *8*, 455-498.
762     https://doi.org/10.1146/annurev.bioeng.8.061505.095721
763
764 Rossel, S., Marshall, F., Peters, J., Pilgram, T., Adams, M.D., O'Connor, D., 2008. Domestication of
765     the donkey: timing, processes, and indicators. Proc. Natl. Acad. Sci. U. S. A. 105 (10), 3715e3720.
766

Roux, W., 1881. Der Kampf der Teile des Organismus. Ein Beitrag zur Vervollständigung der mechanischen Zweckmäßigkeitslehre. Wilhelm Engelmann, Leipzig

Ruff, C., B. Holt, and E. Trinkaus., 2006. Who's afraid of the big bad Wolff ?: "Wolff's Law" and bone functional adaptation. American Journal of Physical Anthropology 129:484–498

Ruff, C. B., Puymerail, L., Macchiarelli, R., Sipla, J., Ciochon, R. L., 2015. Structure and composition of the Trinil femora : Functional and taxonomic implications. Journal of Human Evolution, 80, 147-158. https://doi.org/10.1016/j.jhevol.2014.12.004

Ruff, C.B., 2018 Chapitre 6 Biomechanical analyses of archeological human skeletons. Dans M.A. Katzenberg, & A.L. Grauer, Biological Anthropology of the Human Skeleton. John Wiley & Sons.

Ruimerman, R., Hilbers, P., van Rietbergen, B., Huiskes, R. 2005. A theoretical framework for strain-related trabecular bone maintenance and adaptation. Journal of Biomechanics, 38, 931-941. https://doi.org/10.1016/j.jbiomech.2004.03.037

Rushen, J., 1993. The "coping" hypothesis of stereotypic behaviour. *Animal Behaviour, 45*(3), 613–615. https://doi.org/10.1006/anbe.1993.1071

Russo, L., Massei, G., Genov, P.V., 1997. Daily home range and activity of wild boar in a Mediterranean area free from hunting. Ethology Ecology & Evolution 9, 287–294. https://doi.org/10.1080/08927014.1997.9522888

Schriefer, J.L., Warden, S.J., Saxon, L.K., Robling, A.G., Turner, C.H., 2005. Cellular accommodation and the response of bone to mechanical loading. Journal of Biomechanics 38, 1838–1845. https://doi.org/10.1016/j.jbiomech.2004.08.017

Shackelford, L., Marshall, F., Peters, J., 2013. Identifying donkey domestication through changes in cross-sectional geometry of long bones. Journal of Archaeological Science 40, 4170–4179. https://doi.org/10.1016/j.jas.2013.06.006

Skedros, J.G., Mason, M.W., Bloebaum, R.D., 2001. Modeling and remodeling in a developing artiodactyl calcaneus: A model for evaluating Frost's Mechanostat hypothesis and its corollaries. The Anatomical Record 263, 167–185. https://doi.org/10.1002/ar.1094

Skedros, J.G., Sorenson, S.M., Hunt, K.J., Holyoak, J.D., 2007. Ontogenetic structural and material variations in ovine calcanei: A model for interpreting bone adaptation. The Anatomical Record 290, 284–300. https://doi.org/10.1002/ar.20423

Spitz, F., Janeau, G., 1990. Spatial strategies: an attempt to classify daily movements of wild boar. Acta Theriol. 35, 129–149. doi:10.4098/AT.arch.90-14

Spitz, F., Janeau, G., 1995 Daily selection of habitat in wild boar (Sus scrofa). J. Zool. 237, 423–434.

Su, S.C., Skedros, J.G., Bachus, K.N., Bloebaum, R.D., 1999. Loading conditions and cortical bone construction of an artiodactyl calcaneus. Journal of Experimental Biology 202, 3239–3254. https://doi.org/10.1242/jzeb.202.22.3239

Taylor, D., 1997. Bone maintenance and remodeling: A control system based on fatigue damage. Journal of Orthopaedic Research 15, 601–606. https://doi.org/10.1002/jor.1100150417

Trinkaus, E., Churchill, S.E., Ruff, C.B., 1994. Postcranial robusticity in Homo. II: Humeral bilateral
asymmetry and bone plasticity. American Journal of Physical Anthropology 93, 1–34.
https://doi.org/10.1002/ajpa.1330930102

Turner, C.H., Owan, I., Takano, Y., 1995. Mechanotransduction in bone : Role of strain rate. American
Journal   of   Physiology-Endocrinology   and   Metabolism,   269(3),   E438-E442.
https://doi.org/10.1152/ajpendo.1995.269.3.E438

Turner, C.H., 1998. Three rules for bone adaptation to mechanical stimuli. Bone, 23, 399-407.
https://doi.org/10.1016/S8756-3282(98)00118-5

Turner, C.H., 1999. Toward a mathematical description of bone biology: the principle of cellular
accommodation. Calcif Tissue Int 65, 466–471. https://doi.org/10.1007/s002239900734

Van der Meulen, M.C.H., Morgan, T.G., Yang, X., Baldini, T.H., Myers, E.R., Wright, T.M., Bostrom,
M.P.G., 2006. Cancellous bone adaptation to in vivo loading in a rabbit model. Bone 38, 871–
877. https://doi.org/10.1016/j.bone.2005.11.026

Vlachopoulos, D., Barker, A.R., Ubago-Guisado, E., Fatouros, I.G., Knapp, K.M., Williams, C.A.,
Gracia-Marco, L., 2017. Longitudinal Adaptations of Bone Mass, Geometry, and Metabolism in
Adolescent Male Athletes: The PRO-BONE Study. Journal of Bone and Mineral Research 32,
2269–2277.

Vigne, J.D., Carrere, I., Briois, F., Guilaine, J., 2011. The early process of mammal domestication in
the Near East: New evidence from the Pre-Neolithic and Pre-Pottery Neolithic in Cyprus. Current
Anthropology 52, S255–S271.

Wolff, J., 1986. "The Law of Bone Remodeling" (translation of the German 1892 edition by P. Maquet
and R. Furlong). Berlin Heidelberg New York: Springer

Woo, S.L., Kuei, S.C., Amiel, D., Gomez, M.A., Hayes, W.C., White, F.C., Akeson, W.H., 1981. The
effect of prolonged physical training on the properties of long bone: a study of Wolff's Law. J
Bone Joint Surg Am 63, 780–787.

Zanker, C.L., Swaine, I.L., 2000. Responses of bone turnover markers to repeated endurance running
in humans under conditions of energy balance or energy restriction. Eur J Appl Physiol 83, 434–
440. https://doi.org/10.1007/s004210000293