Title: Co-fluctuations of neural activity define intra-V1 networks related to
perceptual organization

Authors:

Mitchell Valdes-Sosa*[1,2], Marlis Ontivero-Ortega*[1,3], Jorge Iglesias-Fuster[1], Agustin Lage-

Castellanos[1], Lidice Galan-Garcia[1], Pedro Valdes-Sosa[2,1].

Affiliations:

[1] Cuban Center for Neuroscience, Havana, Cuba

[2] The Clinical Hospital of Chengdu Brain Sciences, University of Electronic Sciences and

Technology of China, Chengdu, China

[3] Psychology Faculty, Ghent University, Belgium

* Both authors contributed equally

Email: mitchell@cneuro.edu.cu; luna@cneuro.edu.cu

Abstract

Using functional resonance imaging (fMRI), we studied the relationship between

perceptual organization and network topology within the primary visual cortex (V1). Twenty-six

humans (male and female) were recorded during active observation of two Global and two Local

Navon letters. Correlations between fMRI fluctuations from different V1 sites were measured

(after removing stimulus-evoked signals) in windows specific to each condition. Intra-V1, like

brain-wide networks, presented an overall decrease of correlations during stimulation compared

to baseline and increased statistical dimensionality. Massive edgewise testing and network

based-statistics (both corrected by FDR) identified differences between conditions of connection

strengths that were mapped to the visual field. Global letters elicited long links mainly

connecting V1 sites mapping the lower left/right visual quadrants. Shorter links were associated

with Local letters, primarily mapped within the lower-left visual quadrant. Frequently link

lengths exceeded V1 population receptive field sizes. These connections were not observed in

the time-locked (feedforward) responses shared across participants. Thus, these networks reflect

activity idiosyncratic to each participant, possibly generated by interactions within or feedback to

V1. Perception would sculpt V1 connectivity, with specific increases in link strengths (in a

background of decreases). These findings could help shed light on V1 as a "cognitive

blackboard".

Keywords: V1, perceptual organization, connectivity, networks, fMRI

Running title: Within-V1 networks and perceptual organization
38
39

Introduction

The primary visual cortex (V1) is the main recipient of visual input from the retina, and the initial station of specialized -feedforward- (Lamme & Roelfsema, 2000) routes enlisting higher-order visual areas (Goodale & Milner, 2018; Haak & Beckmann, 2016). Animal studies indicate that V1 neurons have small receptive fields (RF) narrowly tuned for different stimulus features (van Essen et al., 1984). The largest sizes of human V1 population RFs (pRFs) (Wandell & Winawer, 2015), estimated with functional magnetic resonance imaging (fMRI), do not exceed 5° of visual angle in V1. These results indicate that V1 is a bank of local feature detectors that provide a basic description of visual scenes (DeValois & DeValois, 1991).

Visual perceptual organization is a critical stage that groups different stimulus elements, separates figure from background, establishes whole/parts relations, and enables object-based attention (Kimchi et al., 2016; Pomerantz & Portillo, 2015; Wagemans et al., 2012). These processes frequently encompass regions in the visual field (VF) spanning distances bigger than pRF sizes in V1. Thus, perceptual organization needs information integration across sizeable visual field sectors, usually thought to be achieved in higher-order visual areas with larger RFs (Klink et al., 2021; Wandell & Winawer, 2015).

Interestingly, V1 contains information about global stimulus properties that extend across distances bigger than its small RFs. Multivariate pattern analysis (MVPA) of human fMRI activity provides evidence for this. Scene identity can be decoded within regions of V1 that map large but occluded parts of the VF (Smith & Muckli, 2010). Object shapes presented in the VF periphery can be decoded from the (unstimulated) foveal V1 (Williams et al., 2008). MVPA correctly predicts visual working-memory content from V1 activity without exogenous stimulation (Harrison & Tong, 2009), reflecting purely top-down information. This enriched V1

activity, not explainable by its classical RFs, could play an essential role in processes such as

attention, working memory, and imagery, possibly acting like a cognitive blackboard (Roelfsema

& de Lange, 2016).

Interactions between neuronal populations are needed to represent large-scale -global-

properties in V1. Lateral connections between V1 neurons having similar edge-orientation could

play a role, facilitating the detection of extended contours (Chen et al., 2014; Kapadia et al.,

1995; Liang et al., 2017; Schwarz & Bolz, 1991). But input from higher-order visual areas

(Roelfsema & de Lange, 2016) is required to make more complex features (e.g., object form or

scene identity) and broader spatial coverage available to V1. Massive fiber connections from the

visual higher-order regions to V1 could provide an anatomical basis for this feedback (Markov et

al., 2011; Markov et al., 2014).

To understand perceptual organization, we must measure cooperation between neurons

scattered within V1, higher-order visual cortices, and beyond. Doing so is a daunting task.

Luckily, this cooperation may imply the synchronization of neural activity (Singer, 2021), which

could perhaps be measured in V1. Brain sites with synchronized slow fMRI signals are parts of

the same large-scale brain circuits (Bijsterbosch et al., 2017, Bijsterbosch et al., 2020).

Encouragingly for our purposes, fine-grained patterns of visual neural circuitry are also reflected

by this type of synchronization. Connective receptive field (cRF) modelling uses the co-

fluctuations of voxels in V1 with those in other cortical areas to estimate the equivalent of pRFs

for the latter (Bock et al., 2015; Gravel et al., 2014; Haak et al., 2013). Estimated cRFs

correspond well with pRFs obtained with standard techniques. Voxel-to-voxel multivariate

regressions (Baldassano et al., 2012) between V1 and target cortical areas have been used to

estimate retinotopic mappings in the latter. The estimated maps fit closely with results from

classical retinotopic mappings (Baldassano et al., 2016).

Therefore, the neural cooperation underlying perceptual organization could be reflected

in co-fluctuations of fMRI activity between V1 sites. This idea was tested recently by Nasr and

co-workers (Nasr et al., 2021) in human subjects. Larger correlations were found between V1

regions of interest (ROIs) in opposite hemispheres when these areas mapped parts of the same

compared to parts of different objects. This effect was stronger for stimuli in the lower visual

field, congruent with a similar bias for global perception in behavioral data (Nasr & Tootell,

2020; Previc, 1990). These results suggest that changes in perceptual organization are associated

with modifications of functional connectivity within V1.

Here we follow up on the Nasr et al. study, using different stimuli and a finer anatomical

scale. Global vs. Local compound (i.e., Navon) letters were used (Figure 1A), which induced

attention to different spatial scales. We hypothesized that intra-V1 network topology would

depend on the perceptual organization of observed scenes. Vertexwise correlation of fMRI

activity (between all cortical V1 nodes) was measured in time-windows corresponding to each

stimulus. Furthermore, a novel pipeline for intra-V1 connectivity analysis is proposed (Figure 2)

to identify reliable differences in network topology across conditions. Both edge-based and

network-based tests were employed (Chung et al., 2021; Noble et al., 2022). We mapped links

that differed consistently between stimulus classes onto the VF by exploiting the close

correspondence between V1 anatomy and retinotopy (Benson et al., 2012; Benson & Winawer,

2018).

Figure 1. Stimuli and Global>Local activation maps. A) Stimuli used in the experiment.

From left to right, Global 'E' (EG) and 'U' (UG), Local 'E' (EL) and 'U' (UL). Black/white are

reversed. Second-level activation maps of our data (FDR threshold q=0.05), in B) lateral and

medial views and in C) posterior views. On the right in C), green lines show the borders of

retinotopic areas from the Wang et al. (2015) atlas. The colour bar corresponds to t-values.

We first verified if the intraV1-networks identified here presented three previously

reported network properties. One is the dependence of correlations between sites in V1 on their

separation along the cortical surface (Dawson et al., 2016). The second property is a reduction of

correlations during tasks compared to rest (Ito et al., 2020). The third is that statistical

dimensionality increases during tasks compared to rest for brain-wide networks (Ito et al., 2020),

reflecting less information redundancy. We then tested our central hypothesis with an edgewise

analysis predicting the lengths and spatial distribution of V1-links would correspond with the

respective perceptual windows of Global and Local letters (Han et al., 1999). Furthermore, it was

predicted that asymmetries in the VF mapping of significant links and perceptual abilities would

mirror each other (Barbot et al., 2021; Kupers et al., 2019). Specifically, we expected stronger

interhemispheric connections in the lower visual field for Global compared to Local stimuli

(consistent with Nasr et al., 2021) and more VF links for Local compared to Global stimuli (see

Robertson & Ivry, 2000).

Our final question concerned the relative involvement of different fMRI sources in

network generation. The V1 networks could be driven by feedforward responses shared across

participants or activity idiosyncratic to each participant. We modelled task fMRI activity as a

sum of components representing different sources (Nastase et al., 2019), the weight of which

could be modified in the fMRI signal by appropriate preprocessing. This procedure allowed us to

assess the role of these sources in intra-V1 connectivity by using a network-based statistical

approach (Noble & Scheinost, 2020) while also confirming findings from the edgewise method.

Results

The intra-V1-network topologies associated with Global/Local Navon letters were

compared. We used modified Navon figures that allowed temporal separation of the two levels,

thus unmixing their fMRI responses. These stimuli (alternating with background masks) were

presented in 20 s blocks, preceded by a Base mask during 20 s, within an MRI scanner for

twenty-six participants. Local and Global versions of 'E' and 'U' were used, and the task was to

count minor -and infrequent- deviations in letter shape (see Valdés-Sosa et al., 2020).

A second-level analysis (FDR corrected, q=0.05) found increased activity in the foveal

V1 for Local -compared to Global- letters (Figures 1B and C). Previous work has explained this

localization by a more restricted spatial focus of attention for local attention (Sasaki et al., 2001).

Increased activity was also found for Global -compared to Local- letters in a small region of the

para-hippocampal cortex.

<u>V1 network analysis pipeline</u>

We developed the analysis pipeline depicted in Figure 2 to estimate V1 connectivity

based on fMRI activity and segregated by stimuli. In each participant, the fMRI time series were

projected from their native volume to their cortical surface and subsequently registered to the

FreeSurfer average surface. Nuisance variables were regressed out of the fMRI time series to

eliminate the effects of noise, artifacts, and physiological contaminants (Ciric et al., 2017). The

primary analysis was carried out on background activity, defined as the residual time series after

regressing out the stimulus-evoked activity. The stimulus-evoked activity was modelled by

convolving the SPM canonical hemodynamic response function (HRF) with the timing of each

stimulus separately. As described in another section, additional variants of Step B from Figure 2,

no-deconvolution and deconvolution with finite impulse responses (FIR), were also tested and

used to explore the neural sources of V1 networks. Time segments related to the same condition

were identified (after accounting for the HRF delay) and concatenated. The fMRI time series

from 863 vertices from the V1 area mapping of the central 8° of the visual field as defined in

(Benson & Winawer, 2018) were extracted. The Pearson correlations between these time series

were calculated.

The second-level connectivity analysis consisted of contrasts between the matrices from

different stimulus conditions. Subtracting correlations minimizes the effect of between-site

geodesic and retinotopic-eccentricity separation, which otherwise would dominate V1

connectivity estimates. Two within-subject statistical approaches were used. The first consisted

of massive univariate t-tests on the connection (edge) strengths, with correction for multiple

testing through false discovery rate (FDR, q=0.05) thresholds. Edgewise analysis has the

advantage of greater spatial precision but also less statistical power (Noble et al., 2022).

Descriptors related to topology were derived from the networks sparsified with the FDR

thresholds, using ten geometric definitions of connectivity subspaces or connSubsp (e.g., links

crossing the vertical meridian in the lower visual field, see Figure 7 for cartoon depictions).

We used constrained network-based statistics -cNBS- (Noble and Scheinost, 2020), which is

somewhat less precise spatially than edgewise analysis but is more powerful. This method inputs

the edgewise tests, which are grouped into "a priori" subnetworks, for which summary statistics

are generated and submitted to permutation tests. Here the connSubsp partition was used for

cNBS.

Figure 2. Primary data analysis pipeline. A) The first five principal components of white

matter and ventricular (red and blue respectively in the image) time series (and derived

measures) were extracted from the volume fMRI data. B) A general linear model on cortical

surface data was used to regress out the contributions of these noise parameters, motion

correction parameters (and derived measures), and stimulus-evoked activity. C) The resulting

background activity was windowed to correspond with stimulus effects (considering the

hemodynamic response lag). Correlation matrices between all V1 cortical vertices were obtained

per subject, Fisher transformed and subtracted to produce difference matrices. D) A massive

univariate t-test was performed on each entry in the difference matrices across subjects. After

this step, network-based statistics (cNBS) were calculated in addition to edgewise statistics, as

described in the following steps. E) Mixture modelling of the t-values was used to obtain false

discovery rate thresholds, illustrated here by a Gaussian null (blue) and two gamma functions

(red). F) Connections on the cortex that survived the FDR thresholds were mapped to visual field

coordinates (see red arrows), using population values for the pRF centers of each V1 cortical

vertex (blue dots). An example of a between-vertex link is shown in green on the inflated cortical

surface and the visual field. G) In the edgewise analysis, descriptors of the connections were

measured.

Plausibility check of V1-connectivity matrices

The compatibility of our intra-V1 networks with previous findings in the literature was

tested. First, intra-V1 correlations between sites decrease with their separation on the surface

(Dawson et al., 2016). As seen in Figure 3 A, the same result was obtained for intra-V1

correlations. Power functions (similar across conditions, $y=3.25*x^{-0.75}$ with R2=0.85 for the

Local, and $y=3.23*x^{-0.74}$ with R2=0.82 for the Global letters) explained a large proportion of the

relationship between correlations and geodesic distance. This finding validates our strategy of

subtracting measures between conditions to eliminate this effect, which would swamp any other

influence on the correlations.

V1 networks and perceptual organization

Figure 3. Intra-V1 connectivity plausibility check results. A) Between-site correlations (black dots) in the left V1 as a function of their geodesic separation over the cortical surface. A first-order polynomial regression fit to the data is shown as a red curve for each condition. B) The 2D histograms of binned correlations for pairs of stimuli across all edges in V1. The black diagonal line indicates identical correlation values. Base is the pre-stimulus baseline. C) Dimensionality for different V1-covariance matrices.

Second, correlations in interregional and intra-V1functional networks are smaller for task fMRI than in the resting state (Ito et al., 2020). This reduction was also valid here for our intra-V1 connectivity (Figure 3B). Correlations obtained associated with the baseline (our substitute for resting state) were larger than for the Global and Local segments at 76% and 70 % of the edges, respectively. These effects were significant in paired t-tests on the z-transformed V1 correlations (t(349029)=271.3, p<0.00001 for Global, and t(349029)=429.5, p<0.00001 for Local letters). There was also a significant (t(349029)=,271, p<0.0001), albeit less pronounced,

advantage for Global over Local letters (in 24% of edges). Measures of Cohen's d confirmed this

pattern. The effect size was moderate (0.72) for the baseline>Global, strong (1.6) for

baseline>Local, and small 0.45) for Global>Local comparisons.

Third, the statistical dimensionality of brain-wide networks increases during tasks

compared to the resting state. The V1-covariance matrix dimensionality associated with each

stimulus and preceding Base segments (segregated by the level of letters following them) are

shown in Figure 3C. Larger dimensionality was found for Global ($F_{(2, 50)}=13.0$,$p=0.0013$) and

Local ($F_{(2, 50)}=16.5$, $p=0.0004$) letters compared to both types of Base. Effects related to letter

identity, letter level, or their interaction were absent. Dimensionality for Global relative to Local

was also reduced, a non-significant effect that should be verified in a properly powered study.

Therefore, our intra-V1 networks exhibited network properties consistent with those reported

previously.

Edgewise analysis of intra-V1 networks associated with Global/Local Letters

Our main goal was to test if V1 networks associated with Global and Local stimuli

differed in topology. The results of the edgewise connectivity analyses (with canonical-HRF

deconvolution) are shown in Figure 4. FDR thresholds were found at both tails of the distribution

of t-values (Figure 4A), probably because distinct network topologies were present for each

condition. The link maps in the VF (Figure 4B) show these topological differences. It is evident

that for the Global condition, numerous links connected the left and right visual hemifields,

especially for the lower visual quadrants. These horizontal –left/right- links were absent for the

Local condition, for which most edges remained within one quadrant, especially the inferior right

quadrant. Power is reduced in edgewise analyses due to the correction for multiple comparisons.

Figure 5 displays links with Cohen's d values above 0.5. This figure confirms the results of the t-

tests and suggests that some links with moderate/large effect sizes did not survive the FDR

threshold (which tends to be conservative). FDR threshold estimation failed for the edgewise

analyses with the no-deconvolution, FIR deconvolution, and intersubject correlation schemes.

Figure 4. A) Distribution of t-values and mixture models fits. Bayesian information

criterion (BIC) values are displayed above each mixture model. B) Plot in the visual field of

connections surviving the FDR threshold (q=0.05) for Local>Global contrast, and C) for the

Global>Local. Blue dots are pRFs centers for each V1 vertex. FDR threshold in insets. The lines

connect the two centers for each significant link, with a color code indicating their absolute t-

value. D and E) Approximate size of stimuli in the visual field. As shown in the method section,

a narrow sector surrounding the lower half of the vertical meridian had missing data.

Figure 5. Effect sizes (Cohen's d) for the V1 links for each contrast (no vertex-wise FDR
thresholding). The maps are sorted according to effect size ranges indicated in the insets. Only d
> 0.5 are shown. A) Global>local. B) Local>Global.

Figure 6. Histograms of link t-values and lengths in the visual field. A) 2-D histogram of link lengths and t value counts for elements surviving the Global>Local FDR threshold; B) 2-D histogram of link lengths and t value counts for elements surviving the Local>Global FDR threshold. C) Normalized cumulative histogram of distances for links in A) and B). The vertical black dashed line corresponds to maximum pRF sizes in V1 for the central 8 degrees of eccentricity. The horizontal line corresponds to the 50% cumulative distribution level (medians).

As predicted, significant connections tended to be longer in visual field space for the Global>Local contrast (median=4.9º) than for the Local>Global contrast (median=3.6º) (Figure

6). This difference in distributions was highly significant in a Kruskal-Wallis test (df=1, Chi-

sq=3.6471e03, p<0.0001). Note that the t-values are larger also for the Global compared to the

Local conditions. Furthermore, the median length for both types of stimuli was greater than 3º of

the visual field, which is the maximum pRF size in central 8º of the visual field (Wandell &

Winawer, 2015) studied here. To conclude, the edgewise analysis upheld our main prediction

that intra-V1 network topologies were adjusted to the perceptual windows required by Local and

by Global stimuli, with links that did not fit within single V1 pRFs.

Visual field asymmetries in intra-V1 networks

We examined the strength and VF distribution of intra-V1 links that were significant in

the Global vs. Local contrasts by measuring the proportion that crossed from one visual quadrant

to another or stayed in the same quadrant (Figure 7). For this, we used the connSubsp

subnetwork partition. Connections for the Global>Local case mainly crossed the vertical

meridian, linking the left and right visual hemifields (i.e., V1 from the two cerebral

hemispheres), mainly in the lower quadrants. Diagonal and antidiagonal connections were also

present for this condition.

Figure 7. The proportion of connections between/within visual quadrants. Insets schematize the types of connections (connSubsp groups, also used in the network-based analysis). LRup and LRlow, left-right crossings in the upper and lower hemifields. UDleft and UDright, up-down crossings in the left and right hemifields. Main Diag and Antidiag are diagonal links. URquad, ULquad, LLquad, and LRquad, are the links limited to the upper-right, upper-left, lower-left and lower-right visual quadrants.

In the Local>Global case, few connections crossed the vertical meridian, and most links stayed within the lower right quadrant (with a smaller number in the lower left quadrant). In this case, links between the upper and lower quadrants on the right side of the VF, something not seen for the Global>Local contrast.

Using Fisher exact tests, we found significant differences between the Global and Local conditions in the proportion of crossings for the following subspaces: lower visual field (p<0.0001), up-down crossings in the left hemifield (p<0.023), up-down in the right (p<0.0001), crossings staying within the lower left (p<0.0001) and lower right quadrants (p<0.0001). Thus, there were evident asymmetries -exclusive for each condition- in link distribution across the VF. As will be discussed below, these effects parallel psychophysical asymmetries.

<u>Enhancement and suppression of fMRI sources</u>

Different fMRI sources could drive intra-V1 networks, whose relative contributions can

be selectively modulated at the GLM step in our pipeline. The stimulus sequence and timing

were identical for all participants. Hence, it was possible to model the fMRI signal at each

cortical vertex as the sum of four sources, according to the following equation (modified from

Nastase et al., 2019):

$Y(stim, sub) = C(stim) + IdPl(stim, sub) + IdNpl(stim, sub) + N(sub)$ (1)

Where *stim* is the stimulus, *sub* is the participant, *Y* is the fMRI time series, *C* is the response

shared across participants, *IdPl* is the idiosyncratic phase-locked response to the stimuli, *IdNpl* is

the idiosyncratic non-phase-locked response, and *N* represents noise components. Since *N* was

reduced in all analyses by regressing out nuisance covariates, the used signals would mainly

contain mixtures of the first three terms of equation 1. Table 1 shows how different fMRI

preprocessing schemes can theoretically suppress specific components.

Table 1. Effects of preprocessing on fMRI sources for intra-V networks.

Components contributing to correlations after each preprocessing strategy			
	$C(stim)$	$IdPl(stim, sub)$	$IdNpl(stim, sub)$
ISC	Keep	-	-
No-deconvolution	Keep	Keep	Keep
Deconvolve with Canonical HRF	-	Keep	Keep
Deconvolve with FIR	-	-	Keep

Intersubject correlations (ISCs) were estimated between the average fMRI time series at

each V1 vertex across n-1 participants and the time series of the other vertices from the left-out

participant (these correlation matrices were subsequently averaged). Averaging will isolate the

shared response *C*. In the no-deconvolution scheme, task-events responses were not regressed

out (preserving *C, IdPl* and *IdNp*). Deconvolving task effects with the canonical HRF (as used

in previous studies, Keller et al., 2022; Tran et al., 2018) suppresses *C*, leaving

IdPl and dNpl. Here the canonical HRF captured *C* variance well, with significant correlations

in V1 sites (0.8- 1.0) between the grand averages and the time course modelled with this HRF.

Deconvolving responses to task-events with an FIR basis suppresses the phase-locked

components (Goutte et al., 2000; Kay et al., 2008), leaving only *IdNpl*.

Our characterization of the effects of preprocessing schemes (Table 1) is sustained by the

power spectra of the resulting fMRI time series (Figure 8). Base + stimulus blocks lasted 44 s

implying a fundamental frequency of 0.023 Hz. The Grand average (dominated by the *C*

component) presents prominent peaks at this frequency, its harmonics, and the main

subharmonic (probably generated by signal aliasing). These peaks are smaller in the raw data,

lesser after canonical HRF deconvolution, and smallest after FIR deconvolving.

Figure 8. Relative fMRI power spectra. Each curve represents results from a different

preprocessing strategy. All spectra were calculated after denoising. The vertical gray lines

represent the fundamental frequency of stimulation (1/44 s), the first seven harmonics, and the

0.5 sub-harmonic. The grand average is the spectrum of the time series after averaging across

subjects. Raw is the average of spectra across participants when the stimulus-response was not

deconvolved. Canonical HRF and FIR are the averages of spectra across participants after the

stimulus-response was regressed out using the respective basis functions.

<u>Network-based analysis (cNBS)</u>

We confirmed results from the edge-based analysis with a constrained network-based

statistic (cNBS), which has greater statistical power (Noble et al., 2022). We employed the "a

priori" connSubsp partition to avoid circularity. In this analysis, we also compared the effects of

different preprocessing schemes. The cNBS in each connSubsp group was defined as the mean t-

values of its links Global Vs. Local contrast (restricted to the vertices in the central 4° of

eccentricity). A null distribution for the cNBS of each subnetwork was created by randomly

flipping the signs of the Global Vs. Local contrast across participants and then recalculating the

cNBS 10,000 times. The probability of the observed cNBS in the null distribution was

Bonferroni corrected for multiple tests (shown in Table 2).

Row	ISC. G>L	ISC. L>G	No deconv. G>L	No deconv. L>G	Canonical-HRF deconvol. G>L	Canonical-HRF deconvol. L>G	FIR deconvolv G>L	FIR deconvolv L>G
LRu	0.3485	0.6516	**0.0001**	1.0000	**0.0001**	1.0000	**0.0001**	1.0000
LRlo	0.8858	0.1143	**0.0001**	1.0000	**0.0001**	1.0000	**0.0001**	1.0000
UDle	0.6369	0.3632	1.0000	**0.0001**	0.9994	**0.0007**	0.9997	**0.0004**
UDr	0.1718	0.8283	0.9995	**0.0006**	0.9978	**0.0023**	0.9998	**0.0003**
ULeq	0.6667	0.3334	1.0000	**0.0001**	0.9983	**0.0018**	0.9996	**0.0005**
URq	0.0047	0.9954	0.9988	**0.0013**	0.9998	**0.0003**	0.9999	**0.0002**
LoLeq	0.5521	0.4480	0.9999	**0.0002**	1.0000	**0.0001**	0.9998	**0.0003**
LoRq	0.1116	0.8885	0.9949	0.0052	0.9748	0.0253	0.9991	**0.0010**
DiagP	0.6262	0.3739	**0.0001**	1.0000	**0.0001**	1.0000	**0.0001**	1.0000
DiagS	0.9178	0.0823	**0.0001**	1.0000	**0.0001**	1.0000	**0.0001**	1.0000

Table 2. cNBS permutation test results as a function of preprocessing type. Entries are

the probability of belonging to the null distribution. With Bonferroni correction, p=0.05 is 0.005,

significant results in bold font.

Results from the cNBS analysis are consistent with those from the edgewise tests for the

time series deconvolved with the classic HRF (compare with Figures 5 and 6). Subnetworks

crossing the vertical meridian (LRu and LRlo) or that were diagonal (DiagP and DiagS) were

significantly above chance for the Global>Local contrast. Conversely, subnetworks crossing the

horizontal meridian or those restricted to one quadrant were significantly above chance for the

Local>Global contrast. These findings were also valid with no deconvolution or with FIR

deconvolution. In contrast, none of the tests using ISC (representing activity common to all

subjects) were significant. This analysis was confirmed by bootstrap (across participants)

estimates of the cNBS.

Figure 9. Violin plot of bootstrap cNBS replicates in the Global-Local contrast as a function of

preprocessing type. cNBS was calculated for the ten connSubsp by resampling participants with

replacement (n=1000). Positive and negative values indicate larger cNBS values for the Global

and Local conditions, respectively. Black dots correspond to means, the black line to 99%

confidence intervals, and the violin plot colour to preprocessing type.

The 95% Confidence intervals (CIs) excluded zero for all subnetworks with no-

deconvolution, canonical HRF- and FIR- deconvolution. In sharp contrast, all CIs for ISC

included zero. There was a trend to larger cNBS in the following order: no-deconvolution,

HRFF- deconvolution and FIR-deconvolution. However, this was confirmed by bootstrapping

the differences in cNBS between preprocessing variants only with uncorrected p-values ($p<0.01$)

and for the Global>Local contrast. These results demonstrate that the intra-V1 networks

described in this article do not originate from the *C* component -common to all participants-

relying instead on contributions to the fMRI that are characteristic of each individual.

Discussion

Intra-V1, like brain-wide networks, presented an overall decrease of correlations during

stimulation compared to baseline but with increased statistical dimensionality. Patterns of

synchrony in neural activity between V1 cortical nodes varied systematically with re-

configurations in perceptual organization. Compared with Local letters, Global Navon letters

were associated with longer connections in the visual field that frequently crossed the vertical

meridian, especially in the lower visual field. In contrast, links related to Local letters were

shorter and restricted mainly to the lower right visual quadrant and only infrequently crossed the

vertical meridian. These effects were not seen in the intersubject correlations.

Our intra-V1 networks had characteristics previously reported for brain networks. Firstly,

intra-V1 correlations decreased with greater separation between cortical nodes, as reported in

other studies (Dawson et al., 2016; Raemaekers et al., 2014). We circumvented this significant

effect due to geodesic distance by subtracting the correlation matrices associated with different

conditions. Secondly, we found that correlations within V1 dropped with Navon letter

stimulation (compared to the Base), whereas statistical dimensionality increased. The same

outcome has been reported for brain-wide networks using large datasets (Ito et al., 2020),

suggesting less information redundancy between brain regions during tasks, possibly boosting

information coding capacity. Finally, the peaks of fMRI power spectra in our V1 data exhibited

the typical 1/f slope and changed as predicted with stimulus deconvolution.

The edgewise and network-based methods produced consistent results, with all links that

were significant in the first analysis belonging to significant sub-networks of the latter. The

edgewise analysis was more conservative, with FDR threshold estimation failing in some

analyses. Consequently, edgewise testing, followed by correction for multiple comparisons, only

allows visualization of 'tip of the iceberg' effects (Chen et al., 2022). This idea is reinforced by

the many links with large Cohen d values that did not survive FDR thresholds. However, t-maps

(especially with canonical HRF deconvolution) allowed more precise visualization of network

topology changes in the VF.

Our results are congruent with Nasr et al., 2021. In both studies, the synchrony of fMRI

activity between the left and right V1 sites increased when they mapped parts of the same object

compared to parts of different objects. In our case, Global letters straddled the vertical meridian

of the VF, requiring inter-hemispheric cooperation to perceive their overall shape. Like the

fragmented condition in Nasr et al., most of our local letters were circumscribed to either the left

or right visual VFs, precluding the need for interhemispheric integration. The segregation of

most local letters to one VF would explain this case's lack of left/right links. Attention to local

letters in compound figures implies perceptual segregation into smaller elements (Han &

Humphreys, 2002), consistent with the short V1 links confined to single quadrants.

Moreover, in both studies, the significant interhemispheric connections were more robust

in dorsal V1 (i.e., the lower VF). Visual perception is thought to be more Global in the lower

than the upper VF (Christman, 1993; Levine & McAnany, 2005; Previc, 1990), perhaps related

to greater sensitivity for lower spatial frequency (SF) components in this region compared to the

upper VF. Lower SF components are critical for extracting Global shapes in Navon figures,

whereas higher SF components are more important for perceiving Local shapes (Flevaris &

Robertson, 2016). The neural basis of this vertical asymmetry has been recently examined (Nasr

& Tootell, 2020).

Links related to Local letters were mainly restricted to the lower right VF (projecting to

the left hemisphere). When different Navon letters are briefly flashed to each VF, this

competition favours right-sided stimuli when attending the local level and prefers left-sided

stimuli when attending the global level(Flevaris et al., 2010). Patients with left hemisphere

lesions perform worse on recognizing Local Navon letters than those with right hemisphere

lesions (Robertson & Ivry, 2000), who fail more on Global letters. Furthermore, attention to

Local Navon letters elicits larger event-related potentials over the left side of the scalp (Han et

al., 2002; Iglesias-Fuster et al., 2014; Jiang & Han, 2005), whereas Global Navon letters elicit

larger responses over the right side. The left/right cerebral hemispheres are thought to be

specialized for Local/Global processing (Flevaris & Robertson, 2016; Han & Humphreys, 2002),

an idea which fits the concentration of links associated with Local letters in the right VF.

In contrast to Nasr and co-workers, we found for the Local>Global contrast significant

correlations between the dorsal and ventral V1 (corresponding to lower/upper VFs) from the

same hemisphere. These connections crossing the horizontal meridian perhaps signify

augmented vertical neural binding. The reasons for this discrepancy are unclear, but we list two

hypotheses to explore. First, we used a more extensive set of nuisance covariates (as standard in

functional connectivity work, e.g., Cole et al., 2021) than Nasr et al., which could have increased

sensitivity by lowering noise. Second, we used actively attended foveal stimuli, whereas

unattended peripheral stimuli were used in Nasr et al. The former approach could perhaps

strengthen links crossing the horizontal meridian.

The feedforward response to the stimuli (*C*) shared among participants (Nastase et al.,

2019) does not contribute to the intra-V1 networks described here. Although the leave-one-

subject-out procedure used to calculate ISC can inflate positive findings (Chen et al., 2016), we

found only null results with this method. Conversely, the networks are present after

deconvolution with the canonical HRF or FIR (procedures that suppress *C*). These findings

implicate fMRI activity idiosyncratic to the participants as the network source. Idiosyncratic

activity can be phase locked to the stimuli (e.g., fluctuations in evoked activity amplitude or

duration) or non-phased locked (e.g., spontaneous oscillations, also known as background

activity).

A second clue to the neural sources of the networks is the lengths of significant links in

the VF, which exceed the largest pRF sizes in the V1 sectors mapping our stimuli (Benson &

Winawer, 2018; Wandell & Winawer, 2015). Connection lengths implicate either long-range

lateral interactions within V1 (Kapadia et al., 1995; Liang et al., 2017) or feedback from higher-

order visual areas with large RFs (Briggs, 2020; Williams et al., 2008; Petro & Muckli, 2017). It

is thought (Field & Hayes, 2004; Kuai et al., 2017; Schwarz & Bolz, 1991) that enhanced co-

activation of neurons via lateral connections in the superficial layers of V1 can facilitate the

detection of extended contours. Microelectrode recordings in monkeys suggest that contour

detection emerges first in V4 and then is fed back to V1 (Chen et al., 2014; Liang et al., 2017).

Layer-specific fMRI studies in humans have found effects of global stimulus configuration in the

superficial layers of V1 (Lawrence et al., 2019; Marquardt et al., 2019; Petro & Muckli, 2016),

possibly originating from feedback originating in other cortical areas.

Joint fMRI and electrophysiological recordings are required to understand the V1

networks described here. Ultra-slow spontaneous fMRI fluctuations are reported to be associated

with the envelope of gamma-band neural activity (Nir et al., 2007; Scheeringa et al., 2011;

Mateo et al., 2017). Gamma-band activity could affect vasomotor oscillations in cortical surface

pial arterioles, which in turn would influence the arrival of oxyhemoglobin to cortical tissue,

consequently changing the fMRI signal (Mateo et al., 2017; Drew et al., 2020). A seminal

hypothesis (Gray et al., 1989) is that perceptual binding of contours is produced by the

synchronous neuronal activity of neurons representing different aspects of the same object

(Singer, 2021), although not all agree (Riesenhuber & Poggio, 1999). Neuronal synchronization

in beta-band neural activity has also been proposed as a player in contour integration (Kermani

et al., 2020).

One limitation of this study is the lack of eye-movement control, which would allow

stimuli to stray in the VF, blurring the V1 retinotopic mapping. Another limitation is that we

used a population prior map (Benson & Winawer, 2018) to plot V1 vertices onto the VF when

individualized retinotopic mapping would be more precise. To register cortical anatomy across

participants, we used the default FreeSurfer inter-subject surface registration (Fischl, 2012)

instead of more accurate methods (e.g., MSMSULC and MSMALL, Coalson et al., 2018). Due

to missing values, the significance of link effects in a narrow sector of the VF may be

underestimated. A caveat for follow-up studies is that our methods in their present form are

optimal for block designs for which clear temporal separation between experimental conditions

is guaranteed. Furthermore, it would be wise only to compare networks associated with stimuli

differing minimally in retinal layout.

Ideas and speculations

V1 networks, estimated from background fMRI activity, seem to reflect the perceptual

organization of observed visual scenes, with stable topologies across individuals. Although the

spatial resolution of these networks is coarse compared to microelectrode recordings in monkeys

(and blind to faster neuronal interactions), they speak to a finer scale than ROI-based networks.

Furthermore, they are easy to estimate in human experiments, and their visualization in the VF

benefits from the close correspondence between retinotopy and anatomy found in V1. They

could occupy an intermediate level of analysis of V1 circuitry, perhaps analogous to fMRI-based

pRFs relative to microelectrode-measured RFs. Several testable predictions of this idea

immediately come to mind.

Intra-V1 networks could facilitate object recognition in noisy backgrounds, help

circumvent camouflage, and guide object-based attention, with success in these tasks co-varying

alongside network efficiency. The operation of Gestalt laws and different interpretations of

ambiguous (Wyatte et al., 2014), or multi-stable figures (Li et al., 2017), may be reflected in V1

networks. Their topology may also vary as attention shifts within the same stimuli (Flevaris et

al., 2011; Luck et al., 1997). Canonical and non-canonical orientations stimulus orientations

(e.g., upright vs. inverted faces, Strother et al., 2011) could also be associated with different V1

network configurations. Successful and aborted induction of illusory surfaces (e.g., Kanisza

figures) (Chen et al., 2020; Keane et al., 2021) could also generate different network topologies.

The physiological basis of intra-V1 networks could be confirmed in ultrahigh-field fMRI

studies with cortical-layer specific recordings (Huber et al., 2020; Marquardt et al., 2019; Petro

& Muckli, 2017). Together with granger causality, or mediation analysis, layer-specific

recordings could be used to dissect the roles of lateral interaction in V1 from feedback influence

from higher-order areas in the association of network topology with perceptual organization.

<u>Conclusions</u>

Correlations in intra-V1 networks are decreased during stimulation compared to baseline

for most links, with specific increases in some connections. The result is larger statistical

dimensionality. Thus, perception organization could sculpt V1 connectivity. Global and Local

letters elicited different V1 topologies that partially mirror VF perceptual asymmetries. Time-

locked responses shared across participants do not contribute to these networks, whose link

lengths in the VF frequently exceeded V1 pRF sizes. These networks possibly reflect interactions

within V1 or feedback to this area from higher-order areas, which would modulate fMRI activity

idiosyncratic to each participant. These findings, and further studies with the methods we

developed, could help shed light on V1 as a "cognitive blackboard". The main appeal of studying

intra-V1 networks is that it may allow indirect but precise visualization of cooperative effects

among higher-order visual areas, otherwise hidden in voxels scattered across the cortex.

Materials and Methods

<u>Participants</u>

Twenty-six human volunteers aged from 23 to 28 years (9 females) participated in the

study. All had normal, or corrected-to-normal, vision did not present any medical condition and

were right-handed except for two cases. The ethics committee of the University for Electronic

Science and Technology of China (UESTC) approved the procedures, and participants gave

written informed consent following the Helsinki declaration.

<u>Task and stimuli</u>

This experiment is described in more detail in Valdés-Sosa et al., 2020. The stimuli were

projected on a screen at the subject's feet, viewed through an angled mirror fixed to the MRI

head-coil, and were generated using the Cogent Matlab toolbox

(http://www.vislab.ucl.ac.uk/cogent.php). Modified Global and Local Navon letters, made from

white lines on a black background (Figure 1A), alternated with a baseline matrix of (about 2.0°

wide and 5.3° high). This matrix was built out of smaller placeholder elements shaped like '8's

(with visual angles about 40' wide and 1° 3' high). Two letters, 'E' and 'U', were presented at both

levels. Blocks consisted of ten 1 sec repetitions of the same letter (within a fixed level) that

alternated with 1 sec baseline periods, thus lasting 20 s. Blocks were initiated by a 1 s cue

('Global' or 'Local'), followed by a 19 s baseline, and ended with a 4 s wait period where counts

of minor deviations in letter shape were reported. Four or five runs were presented, each

consisting of 12 blocks.

<u>Data acquisition and basic image preprocessing</u>

Data acquisition and initial image preprocessing are described in detail in Valdés-Sosa et

al., 2020. Recordings were carried out with a GE Discovery MR750 3T scanner (General

Electric Medical Systems, Milwaukee, WI, USA) using an eight-channel receiver head coil. A

T1-weighted image was obtained with 1 x 1 x 0.5 mm resolution. A mid-gray surface was

reconstructed from the T1 image for each subject using Freesurfer

(http://surfer.nmr.mgh.harvard.edu), registered to the FsAverage template, and subsampled to

81924 vertices. Discs with 5mm radii were defined over the FsAverage surface using the Surfing

toolbox (http://surfing.sourceforget.net). The T1 image was also segmented and normalized to

MNI space using SPM12. For each subject, masks of white matter and of cerebrospinal fluid

(CSF) restricted to the ventricles (using a template in MNI space)

(https://sites.google.com/site/mrilateralventricle/template were created using a threshold of tissue

probability greater than 0.9.

Functional images were obtained with a T2*- weighted echo planar imaging sequence

(TR=2.5s; TE =40 ms; flip angle=90°). Spatial resolution was 1.875 x 1.875 x 2.9, with 135

images per run. The initial five volumes in all runs were discarded to stabilize T1 magnetization.

Pre-preprocessing of functional data included artifact correction (ArtRepair toolbox,

http://cibsr.stanford.edu/tools/ArtRepair/ArtRepair.htm), followed by slice-timing, head motion

correction (with the extraction of motion parameters) and unwarping using SPM8

(http://www.fil.ion.ucl.ac.uk/spm/). The fMRI time series were then projected to each

individual's mid-gray cortical surface and high-pass filtered with a time constant of 128 s.

Preprocessing of fMRI time series and intra-V1 connectivity estimates

Background activity was defined as the residual time series of each surface vertex after

regressing out the effects of the stimuli (evoked response) and 64 nuisance parameters using the

general linear model (GLM). The nuisance regressors included the six primary motion

parameters (obtained from head motion correction), their derivatives, and the quadratics of both

these sets (24 motion regressors in total). Physiologic noise was modelled using the aCompCor

method (Behzadi et al., 2007) on the time series extracted separately from the white matter and

the ventricles masks described above. The first five principal components from each set of time

series, the derivatives of these components, and the quadratics of all these parameters were

obtained (40 regressors in total). Therefore, the nuisance regressors totaled 64 parameters. We

applied a univariate general linear model (GLM) with the vertex-wise fMRI time series as

dependent variables in three variants. The first variant modelled the stimulation block as a square

wave convolved with the canonical hemodynamic function from the SPM-12 toolbox (canonical-

HRF). In a second variant (no-deconvolve), only nuisance regressors were included in the design

matrix. In the last variant, a FIR basis (18 stick functions covering 45 s: the approximate duration

of a stimulus cycle) was used. Intersubject correlations were also calculated, as described in the

results section. All four variants were used in the network-based statistical analysis. We verified

with a Fourier analysis the effects of these preprocessing schemes on the periodic stimulus-

related activity. The power spectra of complete time series were extracted for each individual

and averaged over participants. We used Bartlett's method to calculate the spectra, which

involves averaging the periodograms of consecutive and non-overlapping segments.

Theoretically, this method does not have the best statistical properties. However, after

comparisons with multi-taper spectrum estimation, little difference was found in the spectral

estimates and much less over-smoothing of peaks with the simpler method.

After applying the GLM, the time series were segmented into blocks considering the time

shift introduced by the hemodynamic function. These segments were linearly detrended.

Segments corresponding to the same stimulus were concatenated. The activity from surface

vertices was smoothed by averaging with the time series of its neighbours in the 5 mm discs

mentioned above. For this article, we used only vertices corresponding to the central 8° of

eccentricity in a population prior retinotopic map of V1 (Benson & Winawer, 2018). Different

vertices could be missing due to noise or BOLD signal dropout. These bad vertices were

concentrated in a small subset of the central fovea and a narrow circular sector about 10 degrees

wide, surrounding the lower portion of the vertical visual field meridian (see Figure 10). Data

from all participants was obtained in 560 V1 vertices, whereas missing data for one case was

found in 85 and for two participants in 68 vertices. The maximum number of missing

participants was 7 (in 3 vertices). Thus, caution should be exercised in interpreting results from

the visual field sectors with the most missing data. Still, the conclusions about the overall

differences between connectivity for Global and Local stimuli are not undercut by this fact.

Figure 10. Missing V1 vertices. The left side displays a plot of how many participants
had missing data at each measured location within the visual field (axis conventions for this
graph are the same as in Figure 1). On the right side, a histogram of how many vertices (y-axis)
had missing data as a function of the number of participants in which this happened (x-axis).

Covariance and Pearson correlation matrices in V1 were calculated based on these time

series for each participant and stimulus condition and transformed to z values using the Fisher

transformation. Negative correlations were set to zero (Sporns & Betzel, 2016), and missing

values in each subject's V1-correlation matrix were substituted by the median value of the other

subjects

<u>Plausibility check of V1-connectivity matrices</u>

As a plausibility check of the V1-connectivity matrices, three previously described

properties of fMRI networks were tested. First, we tested the relationship between the correlation

and geodesic distance between V1. The geodesic distance between V1 cortical sites for each

cortical hemisphere (based on the FsAverage surface) was calculated with the Fast Marching

toolbox (https://es.mathworks.com/matlabcentral/fileexchange/6110-toolbox-fast-marching). The

Fisher-transformed correlations were fit as power functions of geodesic distance. MATLAB

curve-fit toolbox. Second, the size of correlations from resting state (here represented by a pre-

stimulation Base period) and task fMRI were compared. Correlations for the two stimulus

conditions were also compared. The values from the group average of the V1-correlation

matrices for each pair of conditions were used to calculate 2-dimensional histogram density

plots. Edgewise differences between conditions were submitted to groupwise t-tests. Finally, the

dimensionality of resting state and task fMRI data was also verified. Statistical dimensionality

for fMRI data was estimated by the 'participation ratio' (Litwin-Kumar et al., 2017), as used by

Ito et al. (Ito et al., 2020). The V1-covariance matrices were obtained for each condition and

their preceding baseline, and their eigenvalues were extracted. Roughly explained, the

participation ratio finds the number of eigenvalues needed to explain variance greater than a

given threshold, with more components indicating a higher dimensionality of the data.

Dimensionality values were submitted to a one-way repeated-measures analysis of variance

across stimulus conditions.

Edgewise analysis

Within-subjects, two-tailed, group t-tests were calculated based on the differences

between Global and Local connectivity matrices. Multiple comparison corrections were based

False Discovery Rates (FDR) with q=0.05. FDRs thresholds were estimated for both tails of the

t-value distribution by using mixture modelling (Bielczyk et al., 2018; Gorgolewski et al., 2012)

that estimated a Null (H0) and alternative distribution (H1s) from the histogram of observed t-

values. Four mixture models were fit using expectation-maximization: a) Only a Gaussian-H0; b)

The Gaussian H0 plus a negative gamma H1 distribution; c) the Gaussian-H0 plus a positive

gamma H1 distribution; d) And the Gaussian-H0 plus negative gamma and positive gamma H1

distributions. The model with the lowest Bayesian Information Criterion (BIC) was the best fit

(Gorgolewski et al., 2012). The FDR was calculated for each t-value as the quotient of the area

under the H0 distribution above the observed t-value relative to the number of counts of the

empirical histogram (which has contributions from H0 and H1). The FDR was calculated for

both tails of the histogram. The minimum/maximum t-value in the right/left tails associated with

an FDR less than q was defined as the threshold. For some preprocessing variants, this threshold

estimation failed.

Constrained network-based statistics (cNBS)

Ten sub-networks (connSubsp) of the intra-V1 connectivity matrix were predefined

(illustrated in the top row of Figure 7). The number of connections falling into each sub-network

was counted and displayed in histograms, and Fisher's exact test was carried out to contrast the

conditions. The cNBS methodology (Noble & Scheinost, 2020) was used for statistical tests of

the differences between Global and Local conditions. The cNBS was calculated by averaging the

edgewise t-values within each connSubsp sub-network. The null distribution of the cNBS in each

subnetwork was obtained by permutation resampling (flipping the signs of the observed t-values

over participants, n=10,000) and recalculating the statistic. The p-value for the observed cNBS in

each group was calculated from the null distribution. Moreover, confidence intervals for the

subnetwork cNBSs were obtained by bootstrapping over participants 1000 times.

Projection of V1 connectivity onto the visual space

The mean polar angle and eccentricity were extracted for V1 cortical vertices from a

population prior map (Benson & Winawer, 2018). Graph plots depicted significant edges from

the edgewise tests in visual field coordinates. The colour of lines in the graph plot represented

either the t-tests surviving the FDR thresholds or Cohen's d binned according to effect strength.

The length of these lines was measured in visual degrees, and the difference in distributions

between experimental conditions was assessed with a Kruskal-Wallis test.

Bibliography

Baldassano, C., Fei-Fei, L., & Beck, D. M. (2016). Pinpointing the peripheral bias in neural

scene processing networks during natural viewing. *Journal of Vision*, *16*(2), 9.

https://doi.org/10.1167/16.2.9.doi

Baldassano, C., Iordan, M. C., Beck, D. M., & Fei-Fei, L. (2012). Voxel-level functional

connectivity using spatial regularization. *NeuroImage*, *63*(3), 1099–1106.

https://doi.org/10.1016/j.neuroimage.2012.07.046

Barbot, A., Xue, S., & Carrasco, M. (2021). Asymmetries in visual acuity around the visual field.

Journal of Vision, *21*(1), 1–23. https://doi.org/10.1167/JOV.21.1.2

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction

method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1), 90–101.

https://doi.org/10.1016/j.neuroimage.2007.04.042

Benson, N. C., Butt, O. H., Datta, R., Radoeva, P. D., Brainard, D. H., & Aguirre, G. K. (2012).

The retinotopic organization of striate cortex is well predicted by surface topology. *Current*

Biology : CB, *22*(21), 2081–2085. https://doi.org/10.1016/j.cub.2012.09.014

Benson, N. C., & Winawer, J. (2018). Bayesian analysis of retinotopic maps. *ELife*, *7*, 1–29.

https://doi.org/10.7554/elife.40224

Bielczyk, N. Z., Walocha, F., Ebel, P. W., Haak, K. v., Llera, A., Buitelaar, J. K., Glennon, J. C.,

& Beckmann, C. F. (2018). Thresholding functional connectomes by means of mixture

modeling. *NeuroImage*, *171*(October 2017), 402–414.

https://doi.org/10.1016/j.neuroimage.2018.01.003

Bijsterbosch, J., Harrison, S. J., Jbabdi, S., Woolrich, M., Beckmann, C., Smith, S., & Duff, E. P.

(2020). Challenges and future directions for representations of functional brain

organization. *Nature Neuroscience*, *23*(12), 1484–1495. https://doi.org/10.1038/s41593-

020-00726-z

Bijsterbosch, J., Smith, S., & Beckmann, C. (2017). *Introduction to resting state fMRI functional*

connectivity. Oxford University Press.

Bock, A. S., Binda, P., Benson, N. C., Bridge, H., Watkins, K. E., & Fine, I. (2015). Resting-

State Retinotopic Organization in the Absence of Retinal Input and Visual Experience.

Journal of Neuroscience, *35*(36), 12366–12382. https://doi.org/10.1523/JNEUROSCI.4715-

14.2015

Briggs, F. (2020). Role of Feedback Connections in Central Visual Processing. *Annual Review of*

Vision Science, *6*, 313–334. https://doi.org/10.1146/annurev-vision-121219-081716

Chen, G., Shin, Y. W., Taylor, P. A., Glen, D. R., Reynolds, R. C., Israel, R. B., & Cox, R. W.

(2016). Untangling the relatedness among correlations, part I: Nonparametric approaches to

inter-subject correlation analysis at the group level. *NeuroImage*, *142*, 248–259.

https://doi.org/10.1016/j.neuroimage.2016.05.023

Chen, G., Taylor, P. A., Stoddard, J., Cox, R. W., Bandettini, P. A., & Pessoa, L. (2022). Sources

of Information Waste in Neuroimaging: Mishandling Structures, Thinking Dichotomously,

and Over-Reducing Data. *Aperture Neuro*, *2021*(5), 1–22.

https://doi.org/10.52294/2e179dbf-5e37-4338-a639-9ceb92b055ea

Chen, M., Yan, Y., Gong, X., Gilbert, C. D., Liang, H., & Li, W. (2014). Incremental Integration

of Global Contours through Interplay between Visual Cortical Areas. *Neuron*, *82*(3), 682–

694. https://doi.org/10.1016/j.neuron.2014.03.023

Chen, S., Weidner, R., Zeng, H., Fink, G. R., Müller, H. J., & Conci, M. (2020). Tracking the

completion of parts into whole objects: Retinotopic activation in response to illusory figures

in the lateral occipital complex. *NeuroImage*, *207*(June 2019), 116426.

https://doi.org/10.1016/j.neuroimage.2019.116426

Klink, P.C., Chen, X., Vanduffel, W., & Roelfsema, P. R. (2021). Population receptive fields in

non-human primates from whole-brain fmri and large-scale neurophysiology in visual

cortex. *ELife*, *10*, 1–35. https://doi.org/10.7554/eLife.67304

Christman, S. D. (1993). Local-global processing in the upper versus lower visual fields. *Bulletin*

of the Psychonomic Society, *31*(4), 275–278. https://doi.org/10.3758/BF03334927

Chung, J., Bridgeford, E., Arroyo, J., Pedigo, B. D., Saad-Eldin, A., Gopalakrishnan, V., Xiang,

L., Priebe, C. E., & Vogelstein, J. T. (2021). Statistical connectomics. *Annual Review of*

Statistics and Its Application, *8*, 463–492. https://doi.org/10.1146/annurev-statistics-

042720-023234

Ciric, R., Wolf, D. H., Power, J. D., Roalf, D. R., Baum, G. L., Ruparel, K., Shinohara, R. T.,

Elliott, M. A., Eickhoff, S. B., Davatzikos, C., Gur, R. C., Gur, R. E., Bassett, D. S., &

Satterthwaite, T. D. (2017). Benchmarking of participant-level confound regression

strategies for the control of motion artifact in studies of functional connectivity.

NeuroImage, *154*(March), 174–187. https://doi.org/10.1016/j.neuroimage.2017.03.020

Coalson, T. S., van Essen, D. C., & Glasser, M. F. (2018). The impact of traditional

neuroimaging methods on the spatial localization of cortical areas. *Proceedings of the*

National Academy of Sciences of the United States of America, *115*(27), E6356–E6365.

https://doi.org/10.1073/pnas.1801582115

Cole, M. W., Ito, T., Cocuzza, C., & Sanchez-Romero, R. (2021). The functional relevance of

task-state functional connectivity. *Journal of Neuroscience*, *41*(12), 2684–2702.

https://doi.org/10.1523/JNEUROSCI.1713-20.2021

Dawson, D. A., Lam, J., Lewis, L. B., Carbonell, F. M., Mendola, J. D., & Shmuel, A. (2016).

Partial Correlation-Based Retinotopically Organized Resting-State Functional Connectivity

Within and Between Areas of the Visual Cortex Reflects More Than Cortical Distance.

Brain Connectivity, *6*(1), 57–75. https://doi.org/10.1089/brain.2014.0331

DeValois, R. L., & DeValois, K. K. (1991). Spatial Vision. In *Oxford Psychology Series*.

https://doi.org/10.1093/acprof:oso/9780195066579.001.0001

Drew, P. J., Mateo, C., Turner, K. L., Yu, X., & Kleinfeld, D. (2020). Ultra-slow oscillations in

fMRI and resting-state connectivity: neuronal and vascular contributions and technical

confounds. Neuron, 107(5), 782-804.

Field, D. J., & Hayes, A. (2004). Contour Integration and the Lateral Connections of V1

Neurons. In *The Visual Neurosciences* (pp. 1096--1079).

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Flevaris, A. v, Bentin, S., & Robertson, L. C. (2010). Local or global? Attentional selection of

spatial frequencies binds shapes to hierarchical levels. *Psychological Science*, *21*(3), 424–

431. https://doi.org/10.1177/0956797609359909

Flevaris, A. v., Bentin, S., & Robertson, L. C. (2011). Attentional selection of relative SF

mediates global versus local processing: Evidence from EEG. *Journal of Vision*, *11*(7), 1–

12. https://doi.org/10.1167/11.7.1

Flevaris, A. v., & Robertson, L. C. (2016). Spatial frequency selection and integration of global

and local information in visual processing: A selective review and tribute to Shlomo Bentin.

Neuropsychologia, *83*, 192–200. https://doi.org/10.1016/j.neuropsychologia.2015.10.024

Goodale, M. A., & Milner, A. D. (2018). Two visual pathways – Where have they taken us and

where will they lead in future? *Cortex*, *98*, 283–292.

https://doi.org/10.1016/j.cortex.2017.12.002

Gorgolewski, K. J., Storkey, A. J., Bastin, M. E., & Pernet, C. R. (2012). Adaptive thresholding

for reliable topological inference in single subject fMRI analysis. *Frontiers in Human

Neuroscience*, *6*(AUGUST), 1–14. https://doi.org/10.3389/fnhum.2012.00245

Goutte, C., Nielsen, F. Å., & Hansen, L. K. (2000). Modeling the haemodynamic response in

fMRI using smooth FIR filters. *IEEE Transactions on Medical Imaging*, *19*(12), 1188–

1201. https://doi.org/10.1109/42.897811

Gravel, N., Harvey, B., Nordhjem, B., Haak, K. V., Dumoulin, S. O., Renken, R., Ćurčić-Blake,

B., & Cornelissen, F. W. (2014). Cortical connective field estimates from resting state fMRI

activity. *Frontiers in Neuroscience*, *8*(OCT), 1–10.

https://doi.org/10.3389/fnins.2014.00339

Gray, C. M., König, P., Engel, A. K., & Singer, W. (1989). Oscillatory responses in cat visual

cortex exhibit inter-columnar synchronization which reflects global stimulus properties.

Nature, *338*(6213), 334–337. https://doi.org/10.1038/338334a0

Haak, K. v., & Beckmann, C. F. (2016). Objective analysis of the topological organization of the

human cortical visual connectome suggests three visual pathways. *Cortex*, 1–11.

https://doi.org/10.1016/j.cortex.2017.03.020

Haak, K. v., Winawer, J., Harvey, B. M., Renken, R., Dumoulin, S. O., Wandell, B. A., &

Cornelissen, F. W. (2013). Connective field modeling. *NeuroImage*, *66*, 376–384.

https://doi.org/10.1016/j.neuroimage.2012.10.037

Han, S., & Humphreys, G. W. (2002). Segmentation and selection contribute to local processing

in hierarchical analysis. *Quarterly Journal of Experimental Psychology Section A: Human

Experimental Psychology*, *55*(1), 5–21. https://doi.org/10.1080/02724980143000127

Han, S., Humphreys, G. W., & Chen, L. (1999). Parallel and competitive processes in

hierarchical analysis: perceptual grouping and encoding of closure. *Journal of Experimental

Psychology. Human Perception and Performance*, *25*(5), 1411–1432.

https://doi.org/10.1037/0096-1523.25.5.1411

Han, S., Weaver, J. A., Murray, S. O., Kang, X., Yund, E. W., & Woods, D. L. (2002).

Hemispheric asymmetry in global/local processing: Effects of stimulus position and spatial

frequency. *NeuroImage*, *17*(3), 1290–1299. https://doi.org/10.1006/nimg.2002.1255

Harrison, S. A., & Tong, F. (2009). Decoding reveals the contents of visual working memory in

early visual areas. *Nature*, *458*(7238), 632–635. https://doi.org/10.1038/nature07832

Huber, L., Finn, E. S., Chai, Y., Goebel, R., Stirnberg, R., Stöcker, T., Marrett, S., Uludag, K.,

Kim, S. G., Han, S. H., Bandettini, P. A., & Poser, B. A. (2020). Layer-dependent

functional connectivity methods. *Progress in Neurobiology*, *May*, 101835.

https://doi.org/10.1016/j.pneurobio.2020.101835

Iglesias-Fuster, J., Santos-Rodríguez, Y., Trujillo-Barreto, N., & Valdés-Sosa, M. J. (2014).

Asynchronous presentation of global and local information reveals effects of attention on

brain electrical activity specific to each level. *Frontiers in Psychology*, *5*(OCT), 1–14.

https://doi.org/10.3389/fpsyg.2014.01570

Ito, T., Brincat, S. L., Siegel, M., Mill, R. D., He, B. J., Miller, E. K., Rotstein, H. G., & Cole, M.

W. (2020). Task-evoked activity quenches neural correlations and variability across cortical

areas. *PLoS Computational Biology*, *16*(8 August), e1007983.

https://doi.org/10.1371/JOURNAL.PCBI.1007983

Jiang, Y., & Han, S. (2005). Neural mechanisms of global/local processing of bilateral visual

inputs: an ERP study. *Clinical Neurophysiology : Official Journal of the International*

Federation of Clinical Neurophysiology, *116*(6), 1444–1454.

https://doi.org/10.1016/j.clinph.2005.02.014

Kapadia, M. K., Ito, M., Gilbert, C. D., & Westheimer, G. (1995). Improvement in visual

sensitivity by changes in local context: Parallel studies in human observers and in V1 of

alert monkeys. *Neuron*, *15*(4), 843–856. https://doi.org/10.1016/0896-6273(95)90175-2

Kay, K. N., David, S. V., Prenger, R. J., Hansen, K. A., & Gallant, J. L. (2008). Modeling low-

frequency fluctuation and hemodynamic response timecourse in event-related fMRI. *Human*

Brain Mapping, *29*(2), 142–156. https://doi.org/10.1002/hbm.20379

Keane, B. P., Barch, D. M., Mill, R. D., Silverstein, S. M., Krekelberg, B., & Cole, M. W.

(2021). Brain network mechanisms of visual shape completion. *NeuroImage*, *236*.

https://doi.org/10.1016/j.neuroimage.2021.118069

Keller, A. S., Jagadeesh, A. v., Bugatus, L., Williams, L. M., & Grill-Spector, K. (2022).

Attention enhances category representations across the brain with strengthened residual

correlations to ventral temporal cortex. *NeuroImage*, *249*(January), 118900.

https://doi.org/10.1016/j.neuroimage.2022.118900

Kermani, M., Zavitz, E., Oakley, B., Price, N. S. C., Hagan, M. A., & Wong, Y. T. (2020). Long-

range neural coherence encodes stimulus information in primate visual cortex. *BioRxiv*.

https://doi.org/10.1101/2020.06.22.164269

Kimchi, R., Yeshurun, Y., Spehar, B., & Pirkner, Y. (2016). Perceptual organization, visual

attention, and objecthood. *Vision Research*, *126*, 34–51.

https://doi.org/10.1016/j.visres.2015.07.008

Kuai, S. G., Li, W., Yu, C., & Kourtzi, Z. (2017). Contour Integration over Time:

Psychophysical and fMRI Evidence. *Cerebral Cortex*, *27*(5), 3042–3051.

https://doi.org/10.1093/cercor/bhw147

Kupers, E. R., Carrasco, M., & Winawer, J. (2019). Modeling visual performance differences

'around' the visual field: A computational observer approach. *PLoS Computational Biology*,

15(5), 1–29. https://doi.org/10.1371/journal.pcbi.1007063

Lamme, V. A. F., & Roelfsema, P. R. (2000). The distinct modes of vision offered by

feedforward and recurrent processing. *Trends in Neurosciences*, *23*(11), 571–579.

https://doi.org/10.1016/S0166-2236(00)01657-X

Lawrence, S. J. D., Formisano, E., Muckli, L., & de Lange, F. P. (2019). Laminar fMRI:

Applications for cognitive neuroscience. *NeuroImage*, *197*(March), 785–791.

https://doi.org/10.1016/j.neuroimage.2017.07.004

Levine, M. W., & McAnany, J. J. (2005). The relative capabilities of the upper and lower visual

hemifields. *Vision Research*, *45*(21), 2820–2830.

https://doi.org/10.1016/j.visres.2005.04.001

Li, H.-H., Rankin, J., Rinzel, J., Carrasco, M., & Heeger, D. J. (2017). Attention model of

binocular rivalry. *Proceedings of the National Academy of Sciences*, *114*(30), E6192–

E6201. https://doi.org/10.1073/pnas.1620475114

Liang, H., Gong, X., Chen, M., Yan, Y., Li, W., & Gilbert, C. D. (2017). Interactions between

feedback and lateral connections in the primary visual cortex. *Proceedings of the National

Academy of Sciences of the United States of America*, *114*(32), 8637–8642.

https://doi.org/10.1073/pnas.1706183114

Litwin-Kumar, A., Harris, K. D., Axel, R., Sompolinsky, H., & Abbott, L. F. (2017). Optimal

Degrees of Synaptic Connectivity. *Neuron*, *93*(5), 1153-1164.e7.

https://doi.org/10.1016/j.neuron.2017.01.030

Luck, S. J., Chelazzi, L., Hillyard, S. a, & Desimone, R. (1997). Neural mechanisms of spatial

selective attention in areas V1, V2, and V4 of macaque visual cortex. *Journal of*

Neurophysiology, *77*(1), 24–42.

Markov, N. T., Misery, P., Falchier, A., Lamy, C., Vezoli, J., Quilodran, R., Gariel, M. A.,

Giroud, P., Ercsey-Ravasz, M., Pilaz, L. J., Huissoud, C., Barone, P., Dehay, C., Toroczkai,

Z., Van Essen, D. C., Kennedy, H., & Knoblauch, K. (2011). Weight consistency specifies

regularities of macaque cortical networks. *Cerebral Cortex*, *21*(6), 1254–1272.

https://doi.org/10.1093/cercor/bhq201

Markov, N. T., Vezoli, J., Chameau, P., Falchier, A., Quilodran, R., Huissoud, C., Lamy, C.,

Misery, P., Giroud, P., Ullman, S., Barone, P., Dehay, C., Knoblauch, K., & Kennedy, H.

(2014). Anatomy of hierarchy: Feedforward and feedback pathways in macaque visual

cortex. *Journal of Comparative Neurology*, *522*(1), 225–259.

https://doi.org/10.1002/cne.23458

Marquardt, I., de Weerd, P., Schneider, M., Gulban, O. F., Ivanov, D., & Uludağ, K. (2019).

Depth-resolved ultra-high field fMRI reveals feedback contributions to surface motion

perception. *BioRxiv*, 1–28. https://doi.org/10.1101/653626

Mateo, C., Knutsen, P.M., Tsai, P.S., Shih, A.Y. & Kleinfeld, D. (2017). Entrainment of arteriole

vasomotor fluctuations by neural activity is a basis of blood-oxygenation-level-dependent

"resting state" connectivity. Neuron, 96, 1–13.

Nasr, S., Kleinfeld, D., & Polimeni, J. R. (2021). The Global Configuration of Visual Stimuli

Alters Co-Fluctuations of Cross-Hemispheric Human Brain Activity. *The Journal of*

Neuroscience, *41*(47), 9756–9766. https://doi.org/10.1523/jneurosci.3214-20.2021

Nasr, S., & Tootell, R. B. H. (2020). Asymmetries in global perception are represented in near-

versus far-preferring clusters in human visual cortex. *Journal of Neuroscience*, *40*(2), 355–

368. https://doi.org/10.1523/JNEUROSCI.2124-19.2019

Nastase, S. A., Gazzola, V., Hasson, U., & Keysers, C. (2019). Measuring shared responses

across subjects using intersubject correlation. *Social Cognitive and Affective Neuroscience*,

14(6), 669–687. https://doi.org/10.1093/scan/nsz037

Nir, Y., Fisch, L., Mukamel, R., Gelbard-Sagiv, H., Arieli, A., Fried, I., & Malach, R. (2007).

Coupling between neuronal firing rate, gamma LFP, and BOLD fMRI is related to

interneuronal correlations. Current biology, 17(15), 1275-1285.

Noble, S., Mejia, A. F., Zalesky, A., & Scheinost, D. (2022). Improving power in functional

magnetic resonance imaging by moving beyond cluster-level inference. *Proceedings of the*

National Academy of Sciences, *119*(32), e2203020119.

https://doi.org/10.1073/PNAS.2203020119

Noble, S., & Scheinost, D. (2020). The Constrained Network-Based Statistic: A New Level of

Inference for Neuroimaging. In *Lecture Notes in Computer Science (including subseries*

Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics): Vol. 12267

LNCS (pp. 458–468). https://doi.org/10.1007/978-3-030-59728-3_45

Petro, L. S., & Muckli, L. (2016). The brain's predictive prowess revealed in primary visual

cortex. *Proceedings of the National Academy of Sciences*, *113*(5), 1124–1125.

https://doi.org/10.1073/pnas.1523834113

Petro, L. S., & Muckli, L. (2017). The laminar integration of sensory inputs with feedback

signals in human cortex. *Brain and Cognition*, *112*, 54–57.

https://doi.org/10.1016/j.bandc.2016.06.007

Pomerantz, J. R., & Portillo, M. C. (2015). Emergent Features, Gestalts, and Feature Integration

Theory. In J. Wolfe & L. C. Robertson (Eds.), *From Perception to Consciousness*. Oxford

University Press. https://doi.org/10.1093/acprof:osobl/9780199734337.003.0016

Previc, F. H. (1990). Functional specialization in the lower and upper visual fields in humans: Its

ecological origins and neurophysiological implications. *Behavioral and Brain Sciences*,

13(3), 519–542. https://doi.org/10.1017/S0140525X00080018

Raemaekers, M., Schellekens, W., van Wezel, R. J. A., Petridou, N., Kristo, G., & Ramsey, N. F.

(2014). Patterns of resting state connectivity in human primary visual cortical areas: A 7T

fMRI study. *NeuroImage*, *84*, 911–921. https://doi.org/10.1016/j.neuroimage.2013.09.060

Riesenhuber, M., & Poggio, T. (1999). Are cortical models really bound by the "binding

problem"? *Neuron*, *24*(1), 87–93. https://doi.org/10.1016/S0896-6273(00)80824-7

Robertson, L. C., & Ivry, R. (2000). Hemispheric asymmetries: Attention to visual and auditory

primitives. *Current Directions in Psychological Science*, *9*(2), 59–63.

https://doi.org/10.1111/1467-8721.00061

Roelfsema, P. R., & de Lange, F. P. (2016). Early Visual Cortex as a Multiscale Cognitive

Blackboard. *Annual Review of Vision Science*, *2*(1), 131–151.

https://doi.org/10.1146/annurev-vision-111815-114443

Sasaki, Y., Hadjikhani, N., Fischl, B., Liu, A. K., Marret, S., Dale, A. M., & Tootell, R. B. H.

(2001). Local and global attention are mapped retinotopically in human occipital cortex.

Proceedings of the National Academy of Sciences, *98*(4), 2077–2082.

https://doi.org/10.1073/pnas.98.4.2077

Scheeringa, R., Fries, P., Petersson, K. M., Oostenveld, R., Grothe, I., Norris, D. G., ... &

Bastiaansen, M. C. (2011). Neuronal dynamics underlying high-and low-frequency EEG

oscillations contribute independently to the human BOLD signal. Neuron, 69(3), 572-583.

Schwarz, C., & Bolz, J. (1991). Functional specificity of a long-range horizontal connection in

cat visual cortex: A cross-correlation study. *Journal of Neuroscience*, *11*(10), 2995–3007.

https://doi.org/10.1523/jneurosci.11-10-02995.1991

Singer, W. (2021). Recurrent dynamics in the cerebral cortex: Integration of sensory evidence

with stored knowledge. *Proceedings of the National Academy of Sciences*, *118*(33).

https://doi.org/10.1073/pnas.2101043118

Smith, F. W., & Muckli, L. (2010). Nonstimulated early visual areas carry information about

surrounding context. *Proceedings of the National Academy of Sciences of the United States*

of America, *107*(46), 20099–20103. https://doi.org/10.1073/pnas.1000233107

Sporns, O., & Betzel, R. F. (2016). Modular brain networks. *Annual Review of Psychology*,

67(September 2015), 613–640. https://doi.org/10.1146/annurev-psych-122414-033634

Strother, L., Mathuranath, P. S., Aldcroft, A., Lavell, C., Goodale, M. A., & Vilis, T. (2011).

Face inversion reduces the persistence of global form and its neural correlates. *PLoS ONE*,

6(4). https://doi.org/10.1371/journal.pone.0018705

Tran, S. M., McGregor, K. M., James, G. A., Gopinath, K., Krishnamurthy, V., Krishnamurthy,

L. C., & Crosson, B. (2018). Task-residual functional connectivity of language and attention

networks. *Brain and Cognition*, *122*(January), 52–58.

https://doi.org/10.1016/j.bandc.2018.02.003

Valdés-Sosa, M., Ontivero-Ortega, M., Iglesias-Fuster, J., Lage-Castellanos, A., Gong, J., Luo,

C., Castro-Laguardia, A. M., Bobes, M. A., Marinazzo, D., & Yao, D. (2020). Objects seen

as scenes: Neural circuitry for attending whole or parts. *NeuroImage*, *210*(January).

https://doi.org/10.1016/j.neuroimage.2020.116526

van Essen, D. C., Newsome, W. T., & Maunsell, J. H. R. (1984). The visual field representation

in striate cortex of the macaque monkey: Asymmetries, anisotropies, and individual

variability. *Vision Research*, *24*(5), 429–448. https://doi.org/10.1016/0042-6989(84)90041-

5

Wagemans, J., Feldman, J., Gepshtein, S., Kimchi, R., Pomerantz, J. R., van der Helm, P. a, &

van Leeuwen, C. (2012). A century of Gestalt psychology in visual perception: II.

Conceptual and theoretical foundations. *Psychological Bulletin*, *138*(6), 1218–1252.

https://doi.org/10.1037/a0029334

Wandell, B. A., & Winawer, J. (2015). Computational neuroimaging and population receptive

fields. *Trends in Cognitive Sciences*, *19*(6), 349–357.

https://doi.org/10.1016/j.tics.2015.03.009

Williams, M. A., Baker, C. I., op de Beeck, H. P., Mok Shim, W., Dang, S., Triantafyllou, C., &

Kanwisher, N. (2008). Feedback of visual object information to foveal retinotopic cortex.

Nature Neuroscience, *11*(12), 1439–1445. https://doi.org/10.1038/nn.2218

Wyatte, D., Jilk, D. J., & O'Reilly, R. C. (2014). Early recurrent feedback facilitates visual

object recognition under challenging conditions. *Frontiers in Psychology*, *5*(JUL), 1–10.

https://doi.org/10.3389/fpsyg.2014.00674

Author contributions

Designed research; MVS, MOO

Contributed analytic tools: MVS, MOO, ALC, PVS

Analyzed data: MVS, MOO, JIF

Wrote the paper: MVS, MOO, JIF, PVS

Competing Interests

The authors declare no competing interests.

Acknowledgements

We thank the University of Electronic Science and Technology of China (UESTC) for funding and providing special equipment and facilities for this project. This work was supported by the VLIR-UOS project "A Cuban National School of Neurotechnology for Cognitive Aging", the National Fund for Science and Innovation of Cuba, the National Natural Science Foundation of China (#81861128001) and the 1110 Project (B12027) of China. The authors are grateful to K. Uludag for his suggestions on the manuscript.