Non-canonical Telomerase Reverse Transcriptase Controls Osteogenic Reprogramming

of Aortic Valve Cells Through STAT5

Rolando A. Cuevas[1], Luis Hortells[1,2], Claire C. Chu[1], Ryan Wong[1], Alex Crane[1], Camille

Boufford[1], Cailyn Regan[1], William J. Moorhead III[1], Michael J. Bashline[1], Aneesha Parwal[1],

Angelina M. Parise[1], Parya Behzadi[1], Mark J. Brown[1], Aditi Gurkar[3], Dennis Bruemmer[4], John

Sembrat[5], Ibrahim Sultan[6], Thomas G. Gleason[6], Marie Billaud[6], and Cynthia St. Hilaire[1,8*]

[1] Division of Cardiology, Department of Medicine, Pittsburgh Heart, Lung, Blood and Vascular

Medicine Institute, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

[2] current address - Institute of Experimental and Clinical Pharmacology and Toxicology,

University of Freiburg, 79110 Freiburg, Germany.

[3] Aging Institute, Division of Geriatrics, Department of Medicine, University of Pittsburgh,

Pittsburgh, Pennsylvania, USA.

[4] Department of Cardiovascular Medicine, Cleveland Clinic, Cleveland, OH, USA.

[5] Division of Pulmonary, Allergy, and Critical Care Medicine, University of Pittsburgh School of

Medicine, Pittsburgh, PA, USA.

[6] Division of Cardiac Surgery, Department of Cardiothoracic Surgery, Heart and Vascular

Institute, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA.

[7] Department of Surgery, Division of Cardiac Surgery, Brigham and Women's Hospital &

Harvard Medical School, Boston, Maryland, USA.

[8] Department of Bioengineering, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

Running Title: TERT/STAT5 is required for valvular calcification

*Corresponding Author Contact Info:

Cynthia St. Hilaire, PhD, FAHA

University of Pittsburgh

BSTWR 1744.1

200 Lothrop Street, Pittsburgh, PA 15261

+1 412-648-9441 sthilair@pitt.edu

TOTAL WORD COUNT: 6617

ABSTRACT WORD COUNT: 268

Background: Calcific aortic valve disease (CAVD) is the pathological remodeling of valve

leaflets. The initial steps in the osteogenic reprogramming of the leaflet are not fully understood.

Studies have shown that TERT overexpression primes mesenchymal stem cells to differentiate

into osteoblasts, and we investigated whether TERT contributes to the osteogenic

reprogramming of valve interstitial cells (VICs).

Methods: Human control and CAVD aortic valve leaflets and patient-specific hVICs were used

in in vivo and in vitro calcification assays. Loss of function experiments in hVICs and cells

isolated from $Tert^{-/-}$ and $Terc^{-/-}$ mice were used for mechanistic studies. In silico modeling,

proximity ligation and co-immunoprecipitation assays defined novel TERT interacting partners.

Chromatin immunoprecipitation and CUT&TAG sequencing defined protein-DNA interactions.

Results: TERT protein was highly expressed in calcified valve leaflets without changes in

telomere length, DNA damage, or senescence markers, and these features were retained in

isolated primary hVICs. *TERT* expression increased with osteogenic stimuli, while knock-down

or genetic deletion of *TERT* prevented calcification. Mechanistically, TERT was required to

initiate osteogenic reprogramming independent of its canonical telomere-extending activity and

the lncRNA *TERC*. TERT exerts its osteogenic functions via binding with Signal Transducer and

Activator of Transcription 5 (STAT5). Depletion and inhibition of STAT5 prevented calcification.

STAT5 was found to bind the promoter region of Runt-Related Transcription Factor 2 (*RUNX2*),

the master regulator of osteogenic reprogramming. Finally, we demonstrate that TERT and

STAT5 are upregulated and colocalized in CAVD tissue.

Conclusions: TERT's non-canonical activity is required to initiate VIC calcification. TERT

partners with STAT5 to bind the *RUNX2* gene promoter. These data identify a novel mechanism

and potential therapeutic target to decrease vascular calcification.

ABBREVIATIONS:

αSMA, alpha 2 smooth muscle actin; αTUB, alpha tubulin; hVICs, human aortic valve interstitial

cells; CAVD, calcific aortic valve disease; Ch-IP, chromatin immunoprecipitation; ECM,

extracellular matrix; γ-H2AX, phosphorylated gamma histone 2AX; LPS, lipopolysaccharide;

BMMSCs, bone marrow mesenchymal stem cells; MSCs, Mesenchymal stem cells; NT, no

treatment; mVICs, mouse aortic valve interstitial cells OPN, osteopontin; OST, osteogenic

treatment; PCNA, proliferating cell nuclear antigen; RUNX2, Runt-elated transcription factor 2;

STAT5, Signal Transducer and Activator of Transcription 5A/B; TERC, telomerase RNA

component; TERT, telomerase reverse transcriptase catalytic subunit; TL, telomere length.

Key Words:

aortic stenosis, calcific aortic valve disease, vascular calcification, osteogenic reprogramming,

TERT, STAT5

Novelty and Significance

What Is Known?

• Calcific aortic valve disease (CAVD) is the most prevalent form of aortic valve pathology.

CAVD strongly correlates with age and leads to heart failure and a high risk of stroke.

Currently, the only therapeutic option is valve replacement, which comes with significant

healthcare costs and additional risks to patients.

• Runt-related transcription factor 2 (RUNX2) is the master transcription factor required for

osteogenic differentiation of stem cells to osteoblasts and osteogenic reprogramming of

cardiovascular cells. Yet, the early events driving its transcription in aortic valve cells are

poorly defined.

• In addition to its reverse transcriptase enzymatic activity, TERT exhibits non-canonical

transcriptional regulatory functions and overexpression of TERT primes mesenchymal stem

cells to differentiate down the osteoblast lineage.

What New Information Does This Article Contribute?

• TERT protein levels in calcified aortic leaflets and valve interstitial cells, and its non-

canonical osteogenic activity are independent of changes in telomere length and cell

senescence.

• Genetic loss or depletion of TERT blocks calcification in valve interstitial cells, coronary

smooth muscle cells, and mesenchymal stem cells.

• Early in the osteogenic reprogramming of VICs TERT co-localizes with STAT5, and this

TERT-tethered STAT5 binds to the *RUNX2* gene promoter, the master regulator of

osteogenic transcriptional programs.

• STAT5 depletion and pharmacological inhibition prevent calcification of human valve

interstitial cells, coronary smooth muscle cells, and mesenchymal stem cells.

What are the clinical implications?

• We have identified TERT-STAT5 as a novel signaling axis that orchestrates the early steps

in the osteogenic reprogramming of aortic valve cells. Inhibiting TERT and STAT5 activity, or

their interaction, may be leveraged to develop pharmacological or biological therapeutic

strategies to halt or prevent calcification in the aortic valve, bioprosthetic valves, and

perhaps other cardiovascular tissues.

• Invasive and expensive surgical procedures are currently the only treatment option for

patients with CAVD. The discovery and defining of the early events driving vascular

calcification identifies novel and druggable targets for developing non-surgical therapies.

INTRODUCTION

Calcific aortic valve disease (CAVD) encompasses a spectrum of pathological remodeling of the

valve leaflet, ranging from mild valve thickening that displays microcalcifications to

macrocalcifications that predominate in more advanced disease stages.[1,2] The advanced state,

aortic stenosis, results from fibrotic remodeling and the formation of calcified nodules on the

valve leaflet that impede and disrupt blood flow, causing excessive strain on the cardiac tissue,

increasing the risk of stroke and leading to heart failure.[3] CAVD severity and incidence increase

with age, with CAVD prevalence >1,000 per 100,000 individuals ≥75 years of age.[4,5] Further,

roughly 0.4-0.6% of the population harbors bicuspid aortic valve malformations, and these

malformed valves are prone to calcify and develop CAVD.[4] Currently, no therapeutic treatment

halts or reverses pathologic calcification that occurs in the aortic valve leaflets or other

cardiovascular tissues. The only therapeutic option to treat aortic stenosis is valve replacement

via surgical or transcatheter procedures, incurring substantial medical costs and health care

burdens to the patient.[6]

Aortic valve interstitial cells (VICs) are the primary cell type within the valve leaflet and are

responsible for maintaining leaflet integrity. VICs reside in a quiescent state; however, adaptive

or maladaptive responses to environmental factors such as inflammation, extracellular matrix

(ECM) breakdown, and mechanical stress may disrupt VIC homeostasis.[7,8] With alterations in

homeostasis, VICs acquire a myofibroblast-like phenotype, termed "activated VICs," capable of

proliferation, contraction, and secretion of proteins that further remodel the extracellular milieu.[9]

Activated VICs can transition into calcifying cells; however, the initial steps regulating this

process are not fully understood. It is well established that the osteogenic reprogramming of

cardiovascular cells such as VICs and vascular smooth muscle cells (SMCs) is analogous to the

differentiation of mesenchymal stem cells (MSCs) into bone-forming osteoblasts; osteogenic

differentiation of MSCs and the reprogramming of cardiovascular cells is primarily orchestrated

by Runt-related transcription factor 2 (*RUNX2*), the master regulator of osteogenic

transcriptional programs.[10-12] Other osteogenic signature markers, such as osteocalcin, alkaline

phosphatase, and osteopontin, are upregulated in the differentiation of stem cells to osteoblasts

and the calcifying of VICs.[9,13] While it is accepted that VICs upregulate the expression of

osteogenic genes in response to stresses such as inflammation, ECM remodeling, and

mechanical stress, the specific mechanistic steps between those stresses and the initial

reprogramming events in the activation of osteogenic transcriptional reprogramming remain ill-

defined.

The telomerase complex extends and maintains the ends of chromosomes after cell division.

The core complex consists of the telomerase RNA component (*TERC*), telomerase reverse

transcriptase (TERT) protein, and other scaffolding proteins.[14] In addition to its role in telomere

maintenance, TERT also exhibits non-canonical functions independent of telomerase activity,

such as regulating gene expression and orchestrating chromatin remodeling.[15-17] For example,

TERT acts as a cofactor in Wnt-dependent promoter activation by interacting with the chromatin

remodeling protein SWI/SNF related, matrix associated, actin dependent regulator of chromatin,

subfamily a, member 4 (SMARCA4), facilitating the expression of Wnt/β-catenin target genes in

stem cells.[18] In a murine model of atherosclerosis, TERT induced chromatin remodeling and

SMC proliferation by chaperoning the retinoblastoma-associated transcription factor 1 (E2F1)

binding to S-phase gene promoters.[19,20] Linking together the themes of calcification and non-

canonical TERT, two co-published studies identified that the overexpression of *TERT* primed

human MSCs to differentiate down the osteogenic lineage and develop bone-like structures.[21,22]

The mechanism by which TERT drives the osteogenic differentiation of human MSCs is

unknown. Collectively, these studies provide strong evidence of telomerase-independent

functions of TERT in altering chromatin states and regulating gene expression via various

protein-protein interactions.

Signal transducers and activators of transcription (STAT) proteins play essential roles in

inflammatory, proliferative, and apoptotic programs, mediating the signal transduction pathways

of various cytokines and growth factors.[23] STATs function as transcription factors and recruit

chromatin remodeling and histone-modifying proteins to upregulate gene transcription.[24]

STAT5A and STAT5B are encoded by two highly conserved genes in humans and mice; 96% of

their amino acid sequence is shared, and many of their activities are redundant.[25] Herein,

unless specific tools utilized allow a distinction, we will refer to both generally as STAT5. Upon

phosphorylation, STAT5 dimerizes and translocates into the nucleus to bind the consensus

motif (TTCNNNGAA) and activates the promoters of various target genes by recruiting

transcriptional co-activators.[26,27] Furthermore, STAT5 was shown to promote the osteogenic

differentiation of murine MSCs.[28]

In this study, we tested the hypothesis that TERT contributes to the osteogenic reprogramming

of VICs. We utilized primary human aortic valve tissues from control donors and CAVD patients

and generated patient-specific VIC lines for in vitro disease modeling. We established baseline

patterns and osteogenic induction of *TERT* and calcification markers in CAVD tissue and VICs.

We assessed the consequences of genetic deletion or depletion of *TERT* during osteogenic

reprogramming in several cell types and in in vivo and ex vivo calcification models. We further

defined the underlying mechanism by which TERT and STAT5 participate in the osteogenic

reprogramming of VICs.

METHODS

Data Availability

The Materials, Methods, and Major Resources Table are found in the Supplemental Material.

The raw data, experimental materials, and analytic methods are available from the

corresponding author upon reasonable request.

RESULTS

TERT is elevated in CAVD tissues.

Human aortic valves were collected after surgical aortic valve replacement procedures or from

cadaveric tissue obtained via the Center for Organ Recovery and Education (CORE) of the

University of Pittsburgh and processed as previously described.[29] Macroscopic examination and

histological staining with Von Kossa were used to determine and confirm whether valves could

be classified as a non-calcified control or as having calcific aortic valve disease (CAVD) (**Figure**

1A, Supplemental Figure 1A, Supplemental Table 1). RUNX2 is the master transcription

factor required to initiate stem cells to differentiate into osteoblasts and is critical for the

pathological osteogenic reprogramming of vascular cells.[30-33] Immunofluorescent staining

showed RUNX2, and the late calcification marker OPN were significantly upregulated in CAVD

tissue compared to control tissues, indicating activation of osteogenic transcriptional programs

in CAVD tissues (**Supplemental Figure 1B**). Previous studies found that overexpression of

TERT primes human MSCs to differentiate down the osteogenic lineage.[21,22] We found that

TERT transcript levels were significantly upregulated in CAVD tissue relative to control samples

(**Figure 1B**), and *TERT* expression levels showed no correlation with the donor's age

(**Supplemental Figure 1C**). Markers of senescence (Cyclin Dependent Kinase Inhibitor 1A,

CDKN1A, Galactosidase Beta 1, *GLB1*)*,* proliferation (Proliferating Cell Nuclear Antigen,

PCNA), DNA damage (Tumor Protein 53, *TP53*), and cytoskeletal markers (actin Alpha 2

Smooth Muscle/αSMA, *ACTA2,* Vimentin, *VIM,* interleukin 6, *IL6*) showed no statistically

significant differences, while the inflammatory marker and tumor necrosis factor-alpha (*TNF)*

was significantly elevated in CAVD samples (**Supplemental Figure 1D**), supporting

observations by others that indicate a role for inflammatory signaling in aortic valve calcification

pathogenesis.[34,35]

We detected elevated TERT protein signal in CAVD valves compared to non-calcified controls,

with the signal localized to areas of calcification. Similarly, the number of cells expressing TERT

was significantly elevated in our CAVD tissues compared to non-calcified control tissues

(**Figure 1C**). No statistically significant differences were observed in the staining levels of

proliferation marker proliferating cell nuclear antigen (PCNA) or the DNA damage indicator

phosphorylated gamma histone 2AX (γ-H2AX) between CAVD and control tissues

(**Supplemental Figure 1E**). We also found no statistically significant differences in the relative

mean telomere length between the control and CAVD tissues in our study (**Figure 1D**),

indicating no overt influence of the canonical telomere-extending functions of TERT. The mean

age of CAVD patients in this study was slightly elevated compared to the control patients, 68

and 54 years old, respectively (**Figure 1E**). Together, these data show that TERT, osteogenic

markers, and inflammatory signatures are upregulated in CAVD tissues compared to controls,

while markers indicating alterations in proliferation, DNA damage, and senescence are not

altered. Despite the elevated TERT expression and protein levels, telomere length remains

unaltered between CAVD and control tissue, suggesting non-canonical TERT activity.

CAVD human VICs at baseline recapitulate observations in CAVD tissues.

We created patient-specific valve interstitial cell (VIC) lines from CAVD and control valves.[29] At

baseline, freshly isolated hVICs from CAVD and control patients exhibited no morphological or

statistical differences in their proliferative or migratory capacity (**Figure 2, A, B, C, and**

Supplemental Figure 1F). TERT and RUNX2 proteins were significantly elevated at baseline in

CAVD hVICs compared to control hVICs (**Figure 2D**). We observed no statistical differences in

the protein levels of the hVIC activation markers SM22 or alpha SMA between CAVD and

control hVICs but observed a significant increase of the mineralizing enzyme tissue non-specific

alkaline phosphatase (TNAP) in CAVD hVICs (**Supplemental 1G**). Like in the leaflets, there

was no statistical difference in the relative telomere length between CAVD and control hVICs

(**Figure 2E**). These data show that hVICs isolated from CAVD patients display elevated TERT

and RUNX2 expression compared to non-calcified control hVICs, recapitulating the osteogenic

phenotype observed in CAVD leaflet tissue.

TERT expression is upregulated in human VICs in an in vitro osteogenic disease model.

To model CAVD in vitro and establish a role for TERT during hVIC osteogenic reprogramming,

we cultured CAVD and control hVICs in osteogenic media conditions (OST), a media widely

utilized to induce osteogenic differentiation of stem cells to osteoblasts.[36] Alizarin Red staining

for calcification revealed that both OST-treated CAVD and control hVICs lay down calcified

matrix as early as 14 days (d14), CAVD hVICs calcified earlier relative to control hVICs (**Figure**

2F). Importantly, we observed that CAVD hVICs calcify de novo under no treatment (NT)

conditions, suggesting that CAVD hVICs are primed for osteogenic reprogramming (**Figure 2F**).

TERT protein levels were significantly increased in control hVICs after d14 of OST treatment,

while in CAVD hVICs, TERT was elevated even under NT conditions (**Figure 2G**). The

calcification markers RUNX2 and OPN were significantly upregulated, indicating active

osteogenic reprogramming (**Figure 2G**). TERT protein signal was elevated in CAVD hVICs

relative to control hVICs at d14 under NT conditions (**Figure 2H**, left panels). At d14 of OST, the

TERT protein staining was elevated and localized near areas of calcification in both control and

CAVD hVICs; however, CAVD hVICs formed larger calcified nodules (**Figure 2H**, right panels).

We observed no significant differences in cell numbers at the end of the 14-day assay under NT

or OST treatment conditions (**Supplemental Figure 2A**). As CAVD hVICs exhibit elevated

osteogenic markers at baseline (**Supplemental Figure 1G**) and are prone to calcify (**Figure 2H**)

de novo, we assessed transcripts in control hVICs under OST. Osteogenic markers *BMP2*,

THBS1, *ALPL,* and myofibroblast markers *CNN1*, *TAGLN*, and *SMTN* were significantly

upregulated in control hVICs at d14 of OST treatment (**Figure 2I**). Likewise, the calcification-

associated gene *FOXO1* and the CAVD-associated metalloproteinase *MMP13* were also

significantly upregulated in OST conditions, as shown previously.[37,38] Lastly, the pro-

inflammatory transcription factor Signal Transducer and Activator of Transcription 5A *(STAT5A)*

was significantly upregulated, while the expression of the isoform *STAT5B* remained

unchanged. These results show that hVICs undergo osteogenic reprogramming instead of de

novo mineral nucleation.[39]

In cancer and stem cells, elevated expression of TERT is associated with long telomeres and

enables unlimited proliferation, while short telomeres can trigger cellular senescence.[40,41]

Although we observed no differences in telomere length in our tissues and cells in this study

(Figures 1D and 2E), we assessed cellular senescence and observed no significant differences

in the expression of senescence-associated β-galactosidase (SA-β-gal, *GLB1*) between CAVD

and control hVICs at baseline **(Supplemental Figure 2B)** nor increased in control hVICs in OST

treatment (**Figure 2I**). SA-β-gal activity staining indicated that cellular senescence was not

engaged even after 28 days of OST treatment (**Supplemental Figure 2C**). These data

confirmed that cellular senescence is not operative during hVICs osteogenic reprogramming.

TERT is required for the osteogenic differentiation of human MSCs.

Previous studies have shown that the overexpression of TERT primes human mesenchymal

stem cells (hMSCs) to differentiate into osteoblasts.[21,22] To assess the role of endogenous

TERT in osteogenic differentiation, we subjected hMSCs to NT and OST treatment. OST

treatment induced robust calcification as early as day 14, while cells in NT conditions did not

calcify (**Supplemental Figure 3A**). Protein analysis showed that the TERT protein levels

significantly increased on day 3 of the OST treatment, coinciding with increased *RUNX2*

expression and RUNX2 protein levels (**Supplemental Figures 3B and 3C**). hMSCs in OST

treatment exhibited intense TERT staining, and TERT-positive cells clustered around calcified

nodules (**Supplemental Figure 3D**). Using lentiviral-mediated transduction of expressing short-

hairpin RNA to deplete TERT (shTERT) or scrambled control (shControl), we found that

knockdown of *TERT* significantly inhibited the calcification of hMSCs (**Supplemental Figures**

3E and 3F). While previous studies show that overexpression of *TERT* primed hMSCs to

undergo osteogenic differentiation, our data show that TERT is required for the osteogenic

differentiation of hMSCs.[21,22]

TERT is required for calcification in vitro and in vivo.

Control hVICs were transduced with shControl and shTERT lentivirus, and like hMSCs,

knockdown of *TERT* significantly inhibited calcification compared to the control group and

downregulated RUNX2 protein levels (**Figures 3A and 3B**). We assessed TERT's role in

calcifying human coronary artery smooth muscle cells (hCASMCs), which calcify in

atherosclerotic plaques.[42] Knocking down TERT significantly inhibited hCASMCs calcification

(**Figure 3C**). Together, these data indicate that TERT is essential for the osteogenic

reprogramming of hVICs and hCASMCs. Next, mice VIC (mVICs) and bone marrow MSCs

(mBMMSCs) lines were generated from *Tert*$^{+/+}$ and *Tert*$^{-/-}$ mice.[43] In line with our observations in

human cells with depleted TERT, we observed significant inhibition of calcification in *Tert*$^{-/-}$

mVICs (**Figure 3D**) and *Tert*$^{-/-}$ mBMMSCs (**Supplemental Figure 4A**) compared to *Tert*$^{+/+}$ cells.

No statistically significant differences were observed in senescence-associated β-galactosidase

activity in mVICs or mBMMSCs (**Supplemental Figures 4B** and **4C**).

To determine whether TERT is required to initiate and/or maintain osteogenic reprogramming,

we cultured control hVICs in OST and transduced shTERT lentivirus 7 or 14 days later, and

calcification was assessed at day 21. Control hVICs transduced with shTERT on day 7 showed

a significant reduction in calcification compared to the OST alone, while transduction of shTERT

on day 14 did not show any decrease in calcification (**Supplemental Figure 4D**), indicating that

TERT is required to initiate reprogramming to the osteogenic phenotype.

Next, we used *Tert*$^{+/+}$ and *Tert*$^{-/-}$ mice to test whether Tert is required for diet-induced non-

atherosclerotic medial arterial calcification in vivo. Of note, *Tert*$^{-/-}$ mice were generated from

heterozygous breeding pairs (F1 generation) and thus did not exhibit shortened telomeres.[44]

Tert$^{+/+}$ and *Tert*$^{-/-}$ mice were fed the A6P6 diet, which was previously shown to induce medial

arterial calcification in the vasculature of mice.[45] At the end of the 12-week regime, we observed

significantly higher levels of calcification in the aorta of *Tert*$^{+/+}$ compared to *Tert*$^{-/-}$ mice (**Figure**

3E). Staining for Runx2 protein mirrored calcification presentation and was significantly higher in

the *Tert*$^{+/+}$ compared to *Tert*$^{-/-}$ tissue (**Figure 3F**). These findings were validated in an ex vivo

calcification assay. Thoracic aortic rings obtained from the *Tert*$^{+/+}$ and *Tert*$^{-/-}$ mice were cultured

in ex vivo OST or control NT medium for 14 days. *Tert*$^{+/+}$ aortic rings exhibited significantly

elevated calcification compared to aortic rings from *Tert*$^{-/-}$ mice (**Supplemental Figure 4E**).

Tert$^{+/+}$ aortic rings also exhibited prominent staining for Runx2 protein, which was absent in

aortic rings from *Tert*[-/-] mice (**Supplemental Figure 4F**), and western blot analysis confirmed

elevated Runx2 protein levels in *Tert*[+/+] aortic rings compared to *Tert*[-/-] (**Supplemental Figure**

4G). Neither calcification nor Runx2 was detected in the NT groups. Together, these data show

that TERT is required for osteogenic reprogramming of cells in vitro and in vivo.

TERT's canonical telomere-extending activity requires the telomerase RNA component (*Terc*) to

maintain chromosome telomere length and genome integrity.[14] *Terc* exerts a dual function in the

telomerase complex; it guides the telomerase complex to the telomere ends and serves as the

primer for TERTs reverse transcriptase activity, *Terc* also is a scaffold that tethers the additional

proteins of the telomerase complex (i.e., Dyskerin, NOP10, GAR1 and NHP2).[46] *Terc* is

essential for the canonical telomere-extending activity of TERT.[47] To ensure that TERT's role in

calcification is independent of its telomere-extending function, we isolated mVICs from *Terc*[-/-]

mice. While *Tert*[-/-] mVICs did not calcify under OST conditions, *Terc*[-/-] mVICs were significantly

calcified (**Figure 3G**). Similarly, in hVICs, BIBR1532, which blocks the reverse transcriptase

activity of TERT,[48] did not prevent calcification nor decrease the expression of *RUNX2*

(**Supplemental Figure 4H and 4I**). Together, these data illustrate that the canonical telomere-

extending function of TERT does not contribute to osteogenic reprogramming.

TERT/STAT5 interact and bind to the *RUNX2* gene promoter to initiate osteogenic

reprogramming.

Several studies have shown that TERT participates in gene transcription by physically

interacting with transcription factors.[17,20,49,50] RUNX2 is the master regulator of osteogenic

differentiation of stem cells.[30,31] Increased *RUNX2* expression and transcriptional activity are the

hallmarks of osteoblast differentiation and cardiovascular calcification and are required for

osteogenic reprogramming of vascular cells.[32,33] As our data show that RUNX2 is significantly

downregulated in *TERT*-knockdown and deficient cells (Figure 3), we used the LASAGNA[51] and

TRANSFAC/MATCH[52] programs to identify putative transcription factor binding sites in the 5 kb

region upstream of the human *RUNX2* gene promoter (NM_001015051.3) and identified eight

potential STAT5 binding sites (**Supplemental Table 2**). We sought to determine whether

STAT5 is bound to the *RUNX2* gene promoter and focused on the two most proximal predicted

STAT5 sites located at -1371 bp and -193 bp upstream of the transcriptional start site (TSS), as

both have high homology to the tetrameric consensus STAT5 binding site and show an elevated

likelihood of being functional sites (**Figures 4A** and **4B**). As inflammation and inflammatory

signaling pathways are known to contribute to CAVD pathogenesis[1,53], we used bacterial

lipopolysaccharide (LPS) as a broad method to activate inflammatory response pathways in

hVICs. We found that LPS was sufficient to induce calcification of hVICs and that TERT-positive

cells accumulated near calcified nodules (**Figure 4C**). Protein analysis showed that OST and

LPS significantly upregulate TERT, RUNX2, and STAT5 proteins (**Figure 4D**). TERT protein

levels were elevated in both the cytosol and nucleus of cells upon OST stimulation

(**Supplemental Figure 5A**). Under NT conditions, STAT5 is observed in the membrane and

cytosol but translocates and accumulates in the nucleus of hVICs in OST (**Supplemental**

Figure 5B). Next, we used genetic and pharmacological approaches to determine the

requirement of STAT5 for the calcification of hVICs. The small molecules StafiA-1 and StafiB-1

specifically inhibit STAT5A and STAT5B, respectively.[54,55] We found that StafiA-1 significantly

inhibited calcification of control hVICs during a 21-day OST assay and that this inhibition was of

a greater extent than the inhibition observed when control hVICs were treated with StafiB-1,

suggesting that STAT5A is the main participant during the osteogenic reprogramming of hVICs

(**Figure 4E**). StafiA-1 significantly reduced calcification in hCASMCs and hMSCs

(**Supplemental Figures 5C** and **5D**). Lentiviral-mediated delivery of shControl or shSTAT5

constructs showed that knocking down STAT5 significantly attenuated calcification (**Figure 4F**),

confirming an essential role for STAT5 in the osteogenic reprogramming of hVICs. Knocking

down STAT5 after 7 or 14 days of OST treatment did not reduce calcification (**Figure 4G**),

indicating that STAT5 activity is required during the initial steps of osteogenic reprogramming.

Others have established that TERT can interact with transcription factors to regulate gene

expression.[17,20,49,50] We assessed whether TERT and STAT5 directly interact. Three methods

were utilized: proximity ligation assay (PLA), co-immunoprecipitation (IP), and exogenously

expressing TERT and STAT5 reporters. PLA can detect protein-protein proximity <40 nm

apart.[56] PLA revealed that OST significantly increased TERT and STAT5 colocalization in

control hVICs after 21 days of OST treatment relative to NT condition (**Figure 4H**). Focusing

only on nuclear co-localization, TERT/STAT5 puncta are significantly elevated in the nucleus of

calcifying hVICs relative to the NT condition (**Supplementary Figure 5E**). Protein IP also shows

that TERT and STAT5 physically interact in hVICs upon OST treatment relative to NT condition

(**Figure 4I**). These data show that TERT and STAT5 interact and translocate into the nucleus

during the osteogenic reprogramming of hVICs. Supporting a role for STAT5 in osteogenic

reprogramming, STAT5 protein levels were significantly lower in the aorta of $Tert^{-/-}$ mice fed the

A6P6 diet compared to the aorta from $Tert^{+/+}$ mice (**Supplemental Figures 5F**), and these

findings were replicated in the ex vivo calcification assay (**Supplemental Figure 5G**). While the

domains of TERT were initially defined by their role in the telomere-extending functions of

TERT, protein partners required for the known non-canonical functions of TERT have been

found in several domains throughout the protein.[57] Further, sites for protein binding partners to

STAT5, such as epigenetic modifiers and transcriptional regulators, have been found throughout

the protein.[58] We used AI-assisted technology to generate tertiary structures of TERT and

STAT5 dimers and predict how they interact. AlphaFold[59] mapped the interaction between

human TERT and dimerized human STAT5a, and these binding data were visualized using

HDOCK[60] (**Supplemental Figure 5H and 5I**). Next, we tested TERT and STAT5 interaction by

expressing them exogenously. We fused the human CDS of TERT and STAT5 to NFAST and

CFAST splitFAST reporters, respectively, to generate TERT-NFAST and STAT5A-CFAST

fusion proteins. This approach allows monitoring of the formation and dissociation of a protein

assembly.[61] Control hVICs were co-transfected with TERT-NFAST and STAT5A-CFAST

plasmids and labeled with 5 µM HBR-3,5DOM (4-hydroxy-3,5- dimethoxy benzylidene

rhodamine) at d3, d5, and d7 of OST treatment or control NT. We found that the TERT/STAT5

splitFAST signal was highly elevated in calcifying control hVICs relative to the NT condition as

early as d3 throughout d7, supporting the early formation of the TERT/STAT5 complex in

calcifying hVICs (**Figure 4J**). Together, these approaches confirm TERT and STAT5 interaction

during osteogenic reprogramming.

To assess the function of TERT-STAT5 interaction, we performed chromatin

immunoprecipitation analysis (Ch-IP). We found that STAT5 is significantly enriched on both

putative STAT5 binding sites in the *RUNX2* promoter during osteogenic reprogramming (**Figure**

4K). *RUNX2* transcription is initiated from two distinct promoters: the distal promoter region P1,

located upstream of exon 1, and the proximal P2 promoter, located upstream of exon 2.[62,63] The

P1 promoter is highly active and required for osteoblast progenitor expansion, osteoblast

maturation, and bone formation, while the P2 promoter-derived isoform is expressed in

osteoblastic and non-osteoblastic mesenchymal cells.[64-66] Genome-wide CUT&Tag analysis of

the *Tert*[+/+] mVICs showed that STAT5 binds to the *Runx2* P1 and P2 promoters during

osteogenic differentiation and that STAT5 was absent in the same region of *Runx2* promoters in

Tert[-/-] mVICs, highlighting the Tert requirement for Stat5 binding to *Runx2* gene promoters

(**Figure 4L**, GEO accession GSE274226). Next, by comparing *Tert*[+/+] and *Tert*[-/-] CUT&Tag

sequencing of *Tert*[+/+] and *Tert*[-/-] mVICs, we found that Stat5 occupies promoter, intragenic, and

intergenic regions throughout the genome (**Figure 4M**). Gene Ontology (GO) pathway analysis

revealed that genes with high Stat5 binding participate in biological processes related to protein

phosphorylation, JAK/STAT signaling, inflammation, osteoblast differentiation, SMC function,

and chromatin remodeling (**Figure 4N**). Together, these data identify that TERT's effects are

mediated via STAT5 binding on the *Runx2* gene promoter and the promoters involved in several

pathways known to be operative during osteogenic reprogramming of cardiovascular cells.

TERT and STAT5 are upregulated in calcified tissue.

Confirming our in vitro and murine in vivo data above, in human CAVD tissue, we found that

STAT5 protein levels are significantly elevated and localized to areas of calcification in the

calcified leaflet, mirroring TERT distribution. Additionally, the number of STAT5-positive cells

was significantly higher in CAVD tissue compared to control tissue (**Figure 5A**). Furthermore,

the number of TERT/STAT5 co-positive nuclei was significantly greater in CAVD valves

compared to control valves (**Figure 5B**). These data support our in vitro investigations and

indicate that TERT/STAT5 interaction contributes to human CAVD pathogenesis.

TERT interacts with SMARCA4 during the osteogenic reprogramming of VICs.

Studies by others have shown that TERT facilitates chromatin accessibility and transcription by

recruiting SMARCA4 (previously referred to as BRG1), the catalytic subunit of the SWI/SNF

chromatin remodeling complex, to specific DNA loci.[18,67,68] We investigated whether SMARCA4

plays a role in the osteogenic reprogramming of hVICs via interacting with TERT. There was no

difference in SMARCA4 protein staining in the aorta of *Tert*[+/+] and *Tert*[-/-] mice fed the A6P6 diet,

nor in aortic rings ex vivo (**Supplemental Figures 6A and 6B**). However, in hVICs, expression

analysis revealed significant upregulation of *SMARCA4* transcripts as early as day five of OST

stimulation (**Supplementary Figure 6C**). Immunoprecipitation showed that TERT and

SMARCA4 physically interact in hVICs upon OST treatment (**Supplementary Figure 6D**).

Immunofluorescence staining showed that TERT and SMARCA4 are detected and colocalized

in cells in the calcified human leaflet (**Supplementary Figure 6E**), mirroring TERT and STAT5

distribution in human CAVD tissue. These findings suggest that SMARCA4 plays an active role

in the osteogenic reprogramming of VICs.

DISCUSSION

This study investigated the early steps required for the osteogenic reprogramming of human

valve interstitial cells in CAVD (**Figure 6**). This study establishes a critical role for TERT in this

process. Specifically, our data show that 1) TERT is upregulated in CAVD tissue and elevated in

primary VIC lines stimulated with osteogenic media. While TERT is elevated in these calcified

tissues and cells, telomere length is not impacted, nor is there evidence of cell senescence; 2)

TERT is required for the calcification of VICs, CASMCs, and MSCs. The requirement for TERT

occurs early in the process of osteogenic reprogramming, as knocking down TERT protein

levels after seven days of osteogenic stimulation did not deter calcification. These osteogenic

reprogramming activities of TERT are independent of its telomerase enzymatic activity, as $Terc^-$

$^{/-}$ cells or the use of the TERT enzymatic inhibitor BIBR1532 did not deter calcification.

Further, TERT co-immunoprecipitated with the chromatin remodeling enzyme SMARCA4, and

these two proteins were also found in the nucleus of CAVD tissues; 3) Inflammatory signals are

known to contribute to CAVD pathogenesis, and we found such signaling can increase TERT

levels, as well as the transcription factor STAT5. In silico modeling predicted TERT/STAT5

interaction, which was proven experimentally with immunoprecipitation, proximity ligation assay,

and a fluorescence complementation system developed for the visualization of transient protein-

protein interactions, and like TERT, genetic knockdown of STAT5, or specific inhibition of

STAT5 activity, prevented calcification. Lastly, the requirement for TERT was established in in

vivo and ex vivo calcification models. Collectively, these experiments support a novel role for

TERT and STAT5 in driving CAVD pathogenesis and cardiovascular reprogramming and

identify a unique, druggable target for cardiovascular calcification.

TERT is well-known for its canonical telomere-extending activity, yet TERT also regulates gene

expression and chromatin remodeling independent of affecting telomere integrity. TERT

interacts physically with several transcription factors to promote the expression of several

genes, including β-catenin, to stimulate genes involved in bone development.[15,16,49] In a wire-

injury restenosis model, TERT binds to the transcription factors SP1 at the *VEGF* promoter and

E2F1 at *MCM7*, *CCNA1*, and *PCNA* promoters, stimulating SMC proliferation and neointima

formation.[20,43,50,69] We did not detect proliferative or migratory differences between control and

CAVD VICs despite elevated levels of TERT at baseline (Figures 2B and 2C), indicating cell-

specific and stress-specific functions of TERT.

In addition to supporting transcription factor function, TERT can act as a cofactor to orchestrate

chromatin remodeling machinery. Studies have shown that TERT serves as a chaperone to

facilitate the recruitment of SMARCA4 and histone acetyltransferases to stimulate chromatin

accessibility and transcription or to maintain heterochromatin structure,[18,67,68] and loss of TERT

abrogates the cellular response to DNA double-strand breaks and alters the overall chromatin

configuration without affecting telomere integrity.[17] We now show that TERT is upregulated in

calcified valve tissues and VICs under osteogenic and inflammatory stimuli. TERT interacts with

STAT5, which binds to the *RUNX2* gene promoter and drives osteogenic reprogramming of the

aortic valve cells. TERT depletion reduced *RUNX2* expression and calcification in several cell

types, and TERT, STAT5, and the chromatin remodeling enzyme SMARCA4 co-

immunoprecipitated in OST-treated VICs (Supplementary Figure 6). Further, TERT and

SMARCA4 co-localized in the nucleus of CAVD tissues. These discoveries, along with the

findings of others showing that *TERT* overexpression drives osteogenic differentiation in human

MSCs,[21,22] support a broad role for TERT serving as a co-factor bridging chromatin remodeling

complexes and transcription factors to facilitate euchromatin formation to modulating cellular

phenotypic transition and osteogenic reprogramming.

Telomere length is considered a biomarker of aging, yet it varies widely between people of the

same age, even twins.[70] Several studies have utilized circulating leukocytes to investigate

whether shortened telomeres correlate with various vascular disease states.[71,72] One study

correlated telomere length in circulating leukocytes with aortic stenosis and found that leukocyte

telomere length was slightly shorter in individuals with aortic stenosis than in age-matched

controls.[73] However, telomere length in leukocytes does not indicate global telomere length in

an individual, telomerase complex activity, or TERT protein function in all cell types. Further,

somatic mosaicism and clonal hematopoiesis discoveries underscore the importance of not

ascribing telomere length in one cell type to telomere length and/or TERT activities in another.[74]

Short telomeres in circulating leukocytes may reflect increased cell turnover due to systemic

inflammation, as leukocytes are highly proliferative, and telomere attrition is highly likely to be a

tissue- and disease-dependent process. In this regard, Huzen et al. compared the telomere

length of the leukocytes of age-matched controls and patients with atherosclerosis and the

telomere length of the cells in the atherosclerotic plaques of the patients.[75] They found that

while telomere length in leukocytes from the atherosclerosis patient group was significantly

shorter than in the control group, telomere length in the atherosclerotic plaques was significantly

longer, indicative of increased telomerase activity. Thus, total leukocyte telomere length does

not represent telomere length in the diseased tissue. Our study investigated TERT protein levels

directly in the valve tissue and patient-specific VICs. We determined that the expression levels

of TERT and osteogenic markers in CAVD tissue and cells were elevated, without changes in

telomere length, proliferation, and DNA damage markers (Figures 1-2, Supplemental Figures 1-

4), suggesting non-canonical TERT activity in these cells.

VIC calcification shares aspects of the transcriptional program observed during the

differentiation of MSCs into osteoblasts, including the upregulation of *RUNX2, BGLAP, TNAP*,

and the secretion of bone-forming proteins and accumulation of calcium minerals.[10,12,76]

Likewise, signals such as mechanical stress, disturbed flow, and inflammation promote VIC

calcification and the osteogenic differentiation of MSCs.[77] These observations suggest that VICs

switch from their quiescent state and undergo an osteogenic reprogramming akin to the

differentiation of MSCs into osteoblasts, acquiring a calcifying phenotype. The mechanisms

coordinating the early molecular events in these cell transitions have yet to be fully understood.

We have hypothesized that fully differentiated cells can de-differentiate into a transitory,

multipotent stem cell-like state and then acquire an osteogenic phenotype.[78] VICs exhibit the

potential to differentiate into osteogenic, chondrogenic, or adipogenic lineages, properties

comparable to mesenchymal stem cells.[79] We chose to investigate TERT in the osteogenic

reprogramming of VICs as TERT is highly expressed in stem cells, and its overexpression

primes mesenchymal stem cells into the osteogenic lineage.[21,22] Our data supports the role of

TERT in the osteogenic reprogramming of VICs, whereby TERT serves as a bridge between

DNA-binding transcription factors and chromatin remodeling enzymes.

The osteogenic differentiation of stem cells is orchestrated by RUNX2, which is also detected in

the osteogenic reprogramming of valvular and vascular cells.[32,33] In a recent transcriptomic

study, Xu et al. identified 14 cell populations in aortic valve tissue samples from two control and

four CAVD valves.[80] The severity of CAVD was assessed via ultrasound imaging, which detects

leaflet mobility; however, examining the macroscopic pictures and histological staining of valve

leaflets used in this analysis shows that while these valves are indeed stiff, they do not exhibit

robust nodules of calcification as do the valves in our data set (Figures 1A, 1C, 5, and

Supplemental Figure 1A). Their analysis identified three VIC populations defined by expression

of *FOS, HSPA6, and SPARC.* However, cells from six valves were pooled, with 10.4% of cells

from control tissue and the remaining 89.6% from CAVD tissues. Bulk transcriptomic data from

these tissues show only a small 1.49-fold increase in *RUNX2* gene expression in CAVD tissues,

in stark contrast with our more stringent analysis of *RUNX2* expression by quantitative PCR

(Figure 2, Supplemental Figure 4) and what is currently accepted.[10-12] Inspection of the publicly

available bulk RNA-sequencing data (PRJNA552159) showed no increase in *TERT* transcript

expression. While the hemodynamic data showed those valves had undergone fibrotic

remodeling, few calcifying cells were present, as seen by their histological staining. This lack of

calcification may explain why TERT expression was undetected in their study. In the general

population, aortic sclerosis commonly goes undetected until a patient develops symptoms such

as dyspnea, angina, and syncope, as the stiffness and remodeling of the leaflet obstructs

ventricular outflow. A population-based study assessed aortic valve disease in 552 randomly

selected men and women aged 55 to 86. Using imaging and Doppler echocardiography, mild

calcification was observed in 40% of the individuals, severe calcification in 13%, and

symptomatic stenosis in only 2.2%.[81] This population-based study illustrates how common aortic

valve remodeling is and highlights the importance of using multiple modalities to accurately

determine valve pathology (i.e., fibrotic vs calcified). In our study, a portion of each patient valve

was used for cell isolation while the rest was utilized for calcification assessment by histological

staining, following a workflow that we developed that guarantees that cells isolated from

surgically removed and postmortem tissues retain their proliferative capacity and endothelial

and interstitial phenotypes in culture.[29] While unbiased sequencing approaches can help

uncover novel mechanisms contributing to disease states, they are skewed towards expression

in a snapshot and can miss short-lived transitory events.

The function of STAT5 in bone formation is multifaceted. While it was shown that STAT5 could

promote the differentiation of MSCs into osteoblasts by upregulating the osteogenic genes in a

Jak2-dependent manner, a Stat5a global knockout mouse exhibited elevated bone mass and

mineral density.[82,83] These conflicting reports indicate that the role of STAT5 in bone formation

and remodeling is complex, and more studies are required. STAT5 has diverse functions as a

transcription factor: it interacts with histone acetyltransferases and transcription factors such as

the glucocorticoid receptor, SP1, YY1, and C/EBPβ to stimulate gene expression,[84-86] and

studies in cancer cells identified that STAT5 can induce *TERT* expression.[87,88] In silico modeling

found multiple putative STAT5 binding sites in the *RUNX2* P1 promoter region, which were

confirmed with Ch-IP assays. This confirms that STAT5/TERT acts to initiate the osteogenic

reprogramming of VICS, establishing a novel mechanism in CAVD pathogenesis whereby

TERT/STAT5 bind to each other and activate osteogenic gene transcription, which drives the

early events in cellular osteogenic reprogramming. Cell phenotype reprogramming requires

chromatin restructuring to allow for the transcription of repressed genes. As observed in other

cell types, we found TERT co-immunoprecipitated with SMARCA4, as well as STAT5, and

genetic depletion of TERT or STAT5 midway through the OST assay (i.e., day 14) did not

perturb the calcification of cells. This observation strongly supports that TERT complexes with

SMARCA4 to remodel chromatin and with the STAT5 transcription factor to bind and activate

RUNX2 gene transcription. We explored the potential physical direct interaction between TERT

and STAT5, and we propose that the RNA binding domain of TERT is the most likely candidate

to interact with STAT5 (Supplemental Figure 5). Studies identifying critical domains for

interaction are currently ongoing.

The strengths of this study lie in the utilization of human and mouse tissues and primary lines

generated from valve leaflets. While our results are robust, it is important to address the

limitations of our study. Firstly, we worked with a limited number of human samples obtained

from cadaveric donors who matched sex and age. Due to this limited sample size, we limited

our statistical analyses to non-parametric analysis, comparing two to four groups, and no mixed

models were included. Regardless, one strength of our study is the multiprong approach to

determining the role of TERT and STAT5 in vascular calcification. Secondly, STAT5A and

STAT5B are two different genes, with 96% of their amino acid sequence shared and their

activities redundant.[25] We have determined that VICs express both variants; however, our

antibodies and shRNA constructs could not differentiate between the two variants. Thus, the two

isoforms may interact with TERT to bind the *RUNX2* gene promoter. STAT5 is activated by

multiple cytokines, including IL-2, IL-6, and TNF-α, and it was shown that IL-2 specifically

activates STAT5 to induce TERT expression in cancer cells.[87,89] Whether STAT5 first induces

TERT expression and then TERT and STAT5 cooperate to control the expression of *RUNX2*

needs to be determined. Thirdly, there is redundancy regarding STAT1/3/5 functions and

signaling.[90] Our study was not designed to address all the complexities of STAT signaling per se

but rather to investigate the role of TERT as a key protein that orchestrates the osteogenic

reprogramming of aortic valve cells; our future studies will explore the role of STAT5 further.

Our work highlights the importance of primary human tissue-based studies. Murine models are

not ideal for studying non-genetic drivers of CAVD, as mice do not develop CAVD de novo.[91,92]

Mouse leaflets' structure differs from human leaflets as they lack a trilaminar organization and

present a melanocyte cell population.[93-95] Recent elegant work from the Giachelli group showed

that in mice, calcification happens in the sinus wall and, to a lesser extent, at the leaflet hinge,

the region between the leaflet VICs and SMCs of the aorta, with no detectable calcification in

the aortic valve leaflets after 26 weeks of a high-fat diet in a *Ldlr* knockout mice model.[96] This

distribution of aortic root calcification is also observed in a mouse model described by a Honda

et al which detected calcification in the sinus wall 16 weeks post wire-induced mechanical

injury.[91] Due to the lack of aortic valve leaflet calcification in atherosclerotic or mechanical injury

murine models, we utilized a diet-induced medial arterial calcification (MAC) model.[45] MAC is

not associated with lipid deposition, fibrous cap formation, or intimal hyperplasia; instead, MAC

stems from the progressive accumulation of calcium and phosphate within the arterial walls.[97]

These structural features distinguish MAC from atherosclerosis and are more representative of

the pathogenesis of valve leaflet calcification.[2,97]

This study has uncovered the early molecular events that enable the osteogenic reprogramming

of the interstitial cells of the aortic valve. Current CAVD therapies are surgical, limited to

mechanical or bioprosthetic valve replacement, and performed only when the disease has

progressed to the advanced point of affecting blood flow and heart function. Our data show that

perturbing TERT or STAT5 can inhibit calcification; however, broadly inhibiting one or the other

of these proteins may have deleterious off-target effects. Understanding the specific binding

domains and other protein binding partners of TERT and STAT5 will enable the development of

small molecules to block their interaction, which is a more feasible therapeutic strategy for aortic

valve calcification than global inhibition. CAVD affects over 9.5 million people in the U.S. and

has an associated 56-67% 5-year mortality rate; hence, the need for early, non-surgical

therapies is great.[4] This study uncovers a novel and targetable pathway that may be leveraged

to halt or prevent CAVD progression.

ACKNOWLEDGEMENTS

We want to acknowledge and thank the Center for Organ Research and Education for their help

and support, as well as tissue donors and their families for making this study possible. We thank

Dr. Alison Tebo for her guidance and assistance in constructing the SplitFAST plasmids tool

used in this study. We would also like to thank Kristin Konopka, tissue bank coordinator at The

Department of Cardiothoracic Surgery, UPMC. We thank Jason Dobbins for his critical reading

of this manuscript.

SOURCES OF FUNDING

This study was supported by funds from the National Institutes of Health (R01 HL142932, R56

HL168657, St. Hilaire), the American Heart Association (20IPA35260111, St. Hilaire; 902641,

Cuevas; 24POST1186619, Behzadi), the McKamish Family Foundation (St. Hilaire and Sultan)

and the Samuel and Emma Winters Foundation (St. Hilaire).

DISCLOSURES

The University of Pittsburgh has filed a provisional patent application regarding technology on

behalf of CSH and RAC based on the findings presented in this manuscript. IS receives

institutional research support from Atricure and Medtronic and is a consultant for Medtronic

Vascular. None of these conflicts of interest are related to this work. All other authors have nothing

to disclose.

FIGURE LEGENDS

Figure 1. TERT is elevated in CAVD tissues. (A) Examples of non-calcified control (top) and

calcified CAVD (bottom) human valves used in this study. Arrowheads indicate calcified nodules

Scalebar. 1 cm. (B), TERT mRNA transcript quantification in control and CAVD valve tissues. n

= 8 control, n= 9 CAVD. (C), Representative serial sections of Von Kossa (dark precipitation)

and TERT immunofluorescent staining in control and CAVD valve tissues. Scalebar 100 µm.

Graphs show the quantification of TERT intensity and the number of TERT-positive cells. n = 9

control, n = 9 CAVD. (D), Telomere length measurements in valve tissues. n = 19 control, n =

12 CAVD. (E), Age of the patients utilized in this study. n = 25 control, n = 16 CAVD. Data are

shown as means ± SD. P values were calculated using the Mann-Whitney U test and shown on

each graph. L = lumen.

Figure 2. TERT is elevated in CAVD hVICs at baseline and upregulation during

osteogenic reprogramming. (A), Representative images showing cell morphology of control

and CAVD hVICs. Scalebar 50 µm. (B), Proliferation of control and CAVD hVICs at baseline. n

= 3 both groups. (C) Relative migration distances of control and CAVD VICs. n = 10 both

groups. (D) Western blot staining of control and CAVD hVICs at baseline. Quantification of

protein levels is shown on the right panels. n = 4 both groups. (E) Telomere length

measurements in VICs. n = 8, control, n = 5 CAVD. (F) Representative images of early

calcification in hVICs growing in normal conditions (no treatment NT) or stimulated with

osteogenic media (OST) for 14 days. Calcium deposition was visualized by Alizarin Red

staining. Scalebar 50 µm. Quantification of calcification on the right graphs. n = 7 both groups.

(G), Western blot staining of samples from control and CAVD hVICs after 14 days of NT or

OST. Quantification of protein levels is shown on the bottom graphs. n = 4 both groups. (H),

Representative immunofluorescent staining images of TERT in control and CAVD hVICs after

14 days of NT or OST. Calcium deposition was visualized with OsteoImage. Scalebar 100 µm.

(I) Expression profile control VICs after 14 days of NT and OST treatment. n = 4 both groups.

Data are shown as means ± SD. P values were calculated by 2-way ANOVA (Figure C), the

Mann-Whitney U test (Figures B, D, E, G, and I), and the Kruskal-Wallis H test with Dunn post

hoc test Figure F) and shown on each graph. * P = 0.0286 in Figure I.

Figure 3. TERT is required to initiate osteogenic transition and calcification. (A) Control

hVICs were transduced lentivirus containing shControl or shTERT and followed by OST

stimulation for up to 28 days. Calcium deposition was visualized by Alizarin Red staining. n = 6

shControl, n = 6 shTERT at 21 days and n = 9 each group at day 28. Scalebar 400 µm. (B)

Representative western blot staining of control hVICs transduced with lentivirus containing

either shControl or shTERT followed by 7 days of OST. (C) hCASMCs were transduced with

shControl or shTERT and stimulated with OST for 14 and 21 days. n = 4 for each group at both

time points. Scalebar 400 µm. (D) Representative images of mVIC isolated from $Tert^{+/+}$ and $Tert^{-}$

$^{/-}$ mice were stimulated with OST for 28 days. Scalebar 400 µm. (E) Representative Von Kossa

staining images of $Tert^{+/+}$ and $Tert^{-/-}$ mice aortas (dark precipitation). Scale bar 400 µm.

Quantification of calcification is shown on the right panel. (F) Representative immunofluorescent

images of Runx2 expression indicative of calcification (red, insert) on $Tert^{+/+}$ and $Tert^{-/-}$ mice

aorta. Scale bar 400 μm. Runx2 quantification of calcification is shown on the right panel. (G)

Representative images of mVICs isolated from F1 $Tert^{-/-}$ or $Terc^{-/-}$ mice were stimulated with

OST for up to 21 days. Scalebar 100 µm. Data are shown as means ± SD. P values were

calculated using the Kruskal-Wallis H test, Dunn post hoc test (Figures A, C, and G), and Mann-

Whitney U test (Figures D, E, and F).

Figure 4. STAT5 binds to TERT and is required for the calcification of human VICs. (A)

Graphic of in silico analysis identifying the consensus binding site for STAT5. (B), Diagram

depicting STAT5 binding to RUNX2 promoter and the positions of the predicted STAT5 binding

sites (purple boxes). (C) Representative immunofluorescence images of control hVICs after 14

days of LPS treatment or OST stimulation. Scalebar 50 μm. (D) Representative western blot

staining of hVICs after 14 days of OST stimulation. (E) Representative images of control hVICs

treated with 10 μM of the STAT5 inhibitors StafiA-1 or StafiB-1 throughout 28 days of

osteogenic stimulation. Scalebar 400 μm. n = 8 each group. The quantification of calcification is

shown on the right graph. (F) Control hVICs were transduced with lentivirus containing

shControl or shSTAT5 constructs and treated with OST for 21 days. Calcium deposition was

visualized by Alizarin Red staining. n = 9 for each group. Scalebar 400 μm. (G) Representative

images of hVIC (left panels) treated with 10 μM of the STAT5 inhibitors StafiA-1 starting at d7 or

d14 of OST treatment. Samples were collected at d21. Scalebar 400 μm. n = 6 for each group.

(H) Representative images of TERT/STAT5 complex (red foci) in control hVICs cultured for 21

days in osteogenic medium and detected by PLA. Scalebar 100 μm. n = 4 for each group. (I)

TERT/STAT5 interaction was detected by co-immunoprecipitation in control hVICs under OST

for 21 days. (J) Control hVIC co-expressing TERT-NFAST and STAT5A-CFAST were labeled

with 5 μM HBR-3,5DOM (4-hydroxy-3,5- dimethoxybenzylidene rhodamine, coral) and imaged

at d3, d5, and d7 of osteogenic treatment (OST) or control no-treatment (NT). Scale bars 50

μm. (K) Chromatin immunoprecipitation shows enrichment of STAT5 on RUNX2 promoter OST

conditions compared to NT. The base position is indicated in the blue boxes. n=4 for each group

as in B. (L) STAT5 CUT&Tag sequencing tracks in genomic regions of Runx2 in no-treatment

control and osteogenic condition-treated mice VICs. (M) Stacked chart representing genome-

wide STAT5 occupancy in promoters, genes, and intergenic regions frequency in mice VICs. (N)

Top 20 GO pathways enriched in genes with significant loss of STAT5 in $Tert^{+/+}$vs $Tert^{-/-}$ mice

VICs. Data are shown as means ± SD. P values were calculated by the Mann-Whitney U test

(Figures F, H, and J) and the Kruskal-Wallis H test (Figures E and G) and indicated on top of

each graph.

Figure 5. TERT and STAT5 are upregulated and colocalized in CAVD tissue. (A)

Representative serial sections of Von Kossa (dark precipitation) and TERT and STAT5

immunofluorescent staining in control and CAVD tissues. Scalebar 100 µm. STAT5 intensity

and STAT5-positive cells (STAT5+) are quantified on the right panels. (B) Quantification of

STAT5-TERT-positive cells. Data are shown as means ± SD, n = 5 control, n = 6 CAVD. P

values were calculated with the Mann-Whitney U test and indicated on each graph. L = lumen.

Figure 6. TERT/STAT5 promotes osteogenic reprogramming. Inflammatory signals promote

the upregulation and interaction of TERT and STAT5A. The TERT/STAT5 complex translocates

into the nucleus, where TERT interacts with SMARCA4 (also known as BRG1), which facilitates

a permissive chromatin configuration, and STAT5A binds specific sites on the RUNX2 promoter,

leading to the osteogenic reprogramming of valve interstitial cells during the early stages of

CAVD pathogenesis.

Supplemental Figure 1. Characterization of human CAVD valves and patient-isolated

VICs at baseline. (A) Representative section of Von Kossa staining in control and valve CAVD

leaflets tissue. Scalebar 1 mm. (B) Representative serial sections of RUNX2 immunofluorescent

staining and OPN in control and CAVD valve tissues. Scalebar 100 µm. Quantification of

RUNX2-positive cells and OPN-positive cells on the leaflet specimen is shown on the right

graphs. n = 5 control, n = 6 CAVD. (C) Correlation between age of the donor and TERT

expression in valve tissues. n = 8 control, n = 9 CAVD. (D) Gene expression profile in the valve

leaflets. n = 7 each group. (E) Representative images of immunofluorescent staining of PCNA

(top panels) and γ-H2AX (bottom panels) in control and CAVD valve tissues. Scalebar 100 µm.

Quantification of PCNA-positive cells and γ-H2AX-positive cells are shown on the right graphs.

n = 4 control, n = 3 CAVD. (F) Representative images of a scratch assay in control and CAVD

hVICs. Time points are indicated on top of each panel. Blue lines denote the cell migration front.

Scalebar 1 mm. n = 10 of each group. (G) Representative images of αSMA, CNN, SM22, TNAP,

and f-Actin in control and CAVD hVICs. Scalebar 50 μm. Signal quantification is shown on the

right. n = 4 each group. Data are shown as means ± SD. P values were calculated with the

Mann-Whitney U test (Figures A, B, D, F, and G), Pearson correlation test (Figure C), and two-

way ANOVA with the Sidak post hoc test (Figure D and shown on top of each graph). L =

lumen.

Supplemental Figure 2. Inhibition of reverse transcriptase activity of TERT does not

prevent osteogenic reprogramming of hVICs. (A) Cell counts of control and CAVD hVICs

after 14 days of NT or OST stimulation. n = 4 each group. (B) β-galactosidase (BLG1) mRNA

transcript quantification in control and CAVD hVICs at baseline. n = 5 control, n= 4 CAVD. (C)

Representative images of senescence-associated β-galactosidase activity staining of hVICs

after 28 days of NT or OST stimulation. Blue cells were considered positive for senescence.

Quantification of senescence is shown on the right graph. n = 4 each group. Data are shown as

means ± SD. P values were calculated with the Mann-Whitney U test (Figure B) and the

Kruskal-Wallis H test with Dunn pairwise comparison post hoc test (Figures A, C) and shown on

top of each graph. Scalebar 400 μm. n = 6 for each group.

Supplemental Figure 3. TERT is upregulated and required for the osteogenic

differentiation of human mesenchymal stem cells. (A) Representative images of hMSCs

stimulated with OST for 14 days. Calcium deposition was visualized by Alizarin Red staining.

Scalebar 50 μm. (B) Western blot staining of samples from hMSCs collected during the 14 days

of OST treatment. (C) RUNX2 expression profile of differentiating hMSCs stimulated for 14 days

with OST media. n = 3. (D) Representative TERT immunofluorescent staining images of hMSCs

stimulated with OST for 14 days. Scalebar 100 µm. (E) Representative images of hMSCs

infected with shTERT or shControl virus and stimulated with OST for 14 days. Calcium

deposition was visualized by Alizarin Red staining. The quantification of calcification is shown

on the right graph. n = 3. (F) Western blot staining of samples from hMSCs collected during the

14 days of OST treatment. Data are shown as means ± SD. P values were calculated with the

Kruskal-Wallis H test with Dunn pairwise comparison post hoc test (Figure C) and the Mann-

Whitney U test (Figure E) and shown on each graph.

Supplemental Figure 4. Inhibition of reverse transcriptase activity of TERT does not

prevent osteogenic reprogramming of VICs. (A) Representative images of mBMMSCs

isolated from $Tert^{+/+}$ or $Tert^{-/-}$ mice and stimulated with OST for 21 days. Scalebar 100 µm.

Quantification of calcification is presented as average absorbance on the right graphs, n = 5 for

each group. (B, C) Representative images of a senescence-associated β-galactosidase (SA-β-

gal) staining of mVICs and mBMMSCs isolated from $Tert^{+/+}$ or F1 $Tert^{-/-}$ mice after 21 days of

OST stimulation. In each group, blue cells were considered positive for senescence, n = 5. (D)

Schematic of OST treatment with delayed shTERT transduction (top panel). VICs were

transduced with shTERT, and calcification was assessed with Alizarin Red S stain at day 21.

Scalebar 400 µm. n = 6 for each group. (E) Representative sections of Von Kossa staining in

thoracic aortic rings isolated from $Tert^{+/+}$ and $Tert^{-/-}$ mice after 14 days of OST treatment relative

to NT control treatment. Quantification is shown on the right. Scale bar 400 $µ$m (F)

Immunofluorescence staining of thoracic aortic rings after OST treatment as in B. Quantification

of Runx2 is shown on the right graph. Scale bar 400 $µ$m (G) Western blot staining of aortic ring

specimens collected and treated in B. Quantification of Runx2 is shown on the right graph. (H)

Human VICs were treated with the TERT inhibitor BiBR1532 and stimulated with OST, and

calcification was assessed on day 28, with n = 5 in each group. (I) TERT, RUNX2, and STAT5A

expression were evaluated on day 14, n = 6 DMSO, n = 5 BIBR1532. Data are shown as means

± SD. P values were calculated with the Kruskal-Wallis H test with Dunn pairwise comparison

post hoc test (Figures A, B, C, E, H) and the Mann-Whitney U test (Figures D, F, G, I) and are

shown on top of each graph.

Supplemental Figure 5. STAT5 is upregulated, binds to TERT, and is required for

osteogenic reprogramming of VICs. (A) Western blot analysis of cytoplasmic and nuclear

fractions of TERT from control VICs obtained after seven days of OST stimulation. Fractions

purity was assessed by staining for markers of cytoplasm (TUB) and nucleoplasm (Histone H3).

Cyt, cytoplasmic fraction; Nuc, nuclear fraction. (B) Representative immunofluorescence image

of STAT5 shows subcellular distribution after 14 days of NT and OST stimulation. Scalebar 40

μm. (C) Representative images of CASMCs treated with 10 μM of the STAT5 inhibitors StafiA-1

or StafiB-1 during the 14 days of OST stimulation. Scalebar 400 μm. n = 4 for each group. (D)

Representative images of MSCs treated with 10 μM of the STAT5 inhibitors StafiA-1 or StafiB-1

during the 21 days of OST stimulation. Scalebar 400 μm. n = 4 for each group. (E)

Representative images of TERT/STAT5 complex (red foci) detected by proximity ligation assay

(PLA) in the nucleus of control VICs. Scalebar 25 μm. The PLA foci quantification and intensity

are shown on the right graph. n = 4 for each group. (H) Mol*3D Viewer rendering of the TERT-

STAT5 dimer interaction predicted by the HDock. Red boxes indicate the position of the

interaction between TERT and STAT5. (F) Representative immunofluorescent images of Stat5

expression after 12 weeks of high adenine high phosphate diet on $Tert^{-/-}$ and $Tert^{-/-}$ mice aorta.

Scale bar 400 μm. Quantification of Stat5 expression is shown on the right panel. (G)

Representative images of Stat5 immunofluorescence staining of thoracic aortic rings isolated

from $Tert^{+/+}$ and $Tert^{-/-}$ mice after 14 days of OST treatment relative to NT control treatment.

Quantification of Stat5 is shown on the right graph. Scale bar 400 μm (I) The diagram shows the

regions involved in the predicted interaction as indicated in A, and the docking and confidence

scores predicted by HDock are indicated. TEN: DNA anchor domain for processive telomeric

repeat addition, RBD: Telomerase RNA binding domain, RT: Finger and palm region of the

reverse transcriptase domain, CTE: C-terminal extension or putative thumb domain of the

reverse transcriptase, CCD: Coiled-coil domain, required for nuclear localization, DBD: DNA

binding domain, HLK: Helical linker and DNA binding domain, SH2: Domain required for

activation and dimerization, TD: Transactivation domain. Data are shown as means ± SD. P

values were calculated with the Mann-Whitney U test (Figures A and G) and the Kruskal-Wallis

H test with Dunn pairwise comparison post-test (Figures B, E, and F) and indicated in each

graph.

Supplemental Figure 6. Chromatin remodeling enzyme SMARCA4 is upregulated and

interacts with TERT during the osteogenic reprogramming of VICs. (A) Representative

immunofluorescent images of Smarca4 expression after 12 weeks of high adenine high

phosphate diet on $Tert^{-/-}$ and $Tert^{-/-}$ mice aorta. Scale bar 400 μm. Quantification of Smarca4

expression is shown on the right panel. (B) Representative images of Smarca4

immunofluorescence staining of thoracic aortic rings isolated from $Tert^{+/+}$ and $Tert^{-/-}$ mice after

14 days of OST treatment relative to NT control treatment. Quantification of Smarca4 is shown

on the right graph. (C) SMARCA4 expression was evaluated on days 5 and 10 on VICs

stimulated with OST. n = 5 each group. (E) TERT/STAT and TERT/SMARCA4 interaction was

detected by co-immunoprecipitation upon osteogenic stimulation on VIC. (E) Representative

TERT and SMARCA4 immunofluorescent staining in control and CAVD tissues. The number of

positive cells is shown on the right graphs. n = 9 control, n = 9 CAVD. Scalebar 100 μm. Data

are shown as means ± SD, and P values were calculated with the Mann-Whitney U test and

indicated on each graph in Figures A and C. L = lumen.

REFERENCES

1. Kraler S, Blaser MC, Aikawa E, Camici GG, Luscher TF. Calcific aortic valve disease:
from molecular and cellular mechanisms to medical therapy. *Eur Heart J.* 2022;43:683-
697. doi: 10.1093/eurheartj/ehab757

2. Small AM, Yutzey KE, Binstadt BA, Voigts Key K, Bouatia-Naji N, Milan D, Aikawa E,
Otto CM, St Hilaire C, American Heart Association Council on G, et al. Unraveling the
Mechanisms of Valvular Heart Disease to Identify Medical Therapy Targets: A Scientific
Statement From the American Heart Association. *Circulation.* 2024;150:e109-e128. doi:
10.1161/CIR.0000000000001254

3. Buttner P, Feistner L, Lurz P, Thiele H, Hutcheson JD, Schlotter F. Dissecting Calcific
Aortic Valve Disease-The Role, Etiology, and Drivers of Valvular Fibrosis. *Front*
Cardiovasc Med. 2021;8:660797. doi: 10.3389/fcvm.2021.660797

4. Coffey S, Roberts-Thomson R, Brown A, Carapetis J, Chen M, Enriquez-Sarano M,
Zuhlke L, Prendergast BD. Global epidemiology of valvular heart disease. *Nat Rev*
Cardiol. 2021;18:853-864. doi: 10.1038/s41569-021-00570-z

5. Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM, Barengo
NC, Beaton AZ, Benjamin EJ, Benziger CP, et al. Global Burden of Cardiovascular
Diseases and Risk Factors, 1990-2019: Update From the GBD 2019 Study. *J Am Coll*
Cardiol. 2020;76:2982-3021. doi: 10.1016/j.jacc.2020.11.010

6. Virani SS, Alonso A, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP,
Chamberlain AM, Chang AR, Cheng S, Delling FN, et al. Heart Disease and Stroke
Statistics-2020 Update: A Report From the American Heart Association. *Circulation.*
2020;141:e139-e596. doi: 10.1161/CIR.0000000000000757

7. Rajamannan NM, Evans FJ, Aikawa E, Grande-Allen KJ, Demer LL, Heistad DD,
Simmons CA, Masters KS, Mathieu P, O'Brien KD, et al. Calcific aortic valve disease:
not simply a degenerative process: A review and agenda for research from the National
Heart and Lung and Blood Institute Aortic Stenosis Working Group. Executive summary:
Calcific aortic valve disease-2011 update. *Circulation.* 2011;124:1783-1791. doi:
10.1161/CIRCULATIONAHA.110.006767

8.   Chen JH, Simmons CA. Cell-matrix interactions in the pathobiology of calcific aortic
910   valve disease: critical roles for matricellular, matricrine, and matrix mechanics cues. *Circ*
911   *Res*. 2011;108:1510-1524. doi: 10.1161/CIRCRESAHA.110.234237

912   9.   Liu AC, Joag VR, Gotlieb AI. The emerging role of valve interstitial cell phenotypes in
913   regulating heart valve pathobiology. *Am J Pathol*. 2007;171:1407-1418. doi:
914   10.2353/ajpath.2007.070251

915   10.   Rajamannan NM, Subramaniam M, Rickard D, Stock SR, Donovan J, Springett M,
916   Orszulak T, Fullerton DA, Tajik AJ, Bonow RO, et al. Human aortic valve calcification is
917   associated with an osteoblast phenotype. *Circulation*. 2003;107:2181-2184. doi:
918   10.1161/01.CIR.0000070591.21548.69

919   11.   Rajamannan NM. Calcific aortic valve disease: cellular origins of valve calcification.
920   *Arterioscler Thromb Vasc Biol*. 2011;31:2777-2778. doi: 10.1161/ATVBAHA.111.237610

921   12.   Aikawa E, Libby P. A Rock and a Hard Place: Chiseling Away at the Multiple
922   Mechanisms of Aortic Stenosis. *Circulation*. 2017;135:1951-1955. doi:
923   10.1161/CIRCULATIONAHA.117.027776

924   13.   Wirrig EE, Yutzey KE. Conserved transcriptional regulatory mechanisms in aortic valve
925   development and disease. *Arterioscler Thromb Vasc Biol*. 2014;34:737-741. doi:
926   10.1161/ATVBAHA.113.302071

927   14.   Calado RT, Young NS. Telomere diseases. *N Engl J Med*. 2009;361:2353-2365. doi:
928   10.1056/NEJMra0903373

929   15.   Stewart SA, Hahn WC, O'Connor BF, Banner EN, Lundberg AS, Modha P, Mizuno H,
930   Brooks MW, Fleming M, Zimonjic DB, et al. Telomerase contributes to tumorigenesis by
931   a telomere length-independent mechanism. *Proc Natl Acad Sci U S A*. 2002;99:12606-
932   12611. doi: 10.1073/pnas.182407599

933   16.   Choi J, Southworth LK, Sarin KY, Venteicher AS, Ma W, Chang W, Cheung P, Jun S,
934   Artandi MK, Shah N, et al. TERT promotes epithelial proliferation through transcriptional
935   control of a Myc- and Wnt-related developmental program. *PLoS Genet*. 2008;4:e10.
936   doi: 10.1371/journal.pgen.0040010

937   17.   Masutomi K, Possemato R, Wong JM, Currier JL, Tothova Z, Manola JB, Ganesan S,
938   Lansdorp PM, Collins K, Hahn WC. The telomerase reverse transcriptase regulates
939   chromatin state and DNA damage responses. *Proc Natl Acad Sci U S A*.
940   2005;102:8222-8227. doi: 10.1073/pnas.0503095102

18.   Park JI, Venteicher AS, Hong JY, Choi J, Jun S, Shkreli M, Chang W, Meng Z, Cheung
P, Ji H, et al. Telomerase modulates Wnt signalling by association with target gene
chromatin. *Nature*. 2009;460:66-72. doi: 10.1038/nature08137

19.   Aono J, Ruiz-Rodriguez E, Qing H, Findeisen HM, Jones KL, Heywood EB, Bruemmer
D. Telomerase Inhibition by Everolimus Suppresses Smooth Muscle Cell Proliferation
and Neointima Formation Through Epigenetic Gene Silencing. *JACC Basic Transl Sci*.
2016;1:49-60. doi: 10.1016/j.jacbts.2016.01.002

20.   Endorf EB, Qing H, Aono J, Terami N, Doyon G, Hyzny E, Jones KL, Findeisen HM,
Bruemmer D. Telomerase Reverse Transcriptase Deficiency Prevents Neointima
Formation Through Chromatin Silencing of E2F1 Target Genes. *Arterioscler Thromb*
Vasc Biol. 2017;37:301-311. doi: 10.1161/ATVBAHA.116.308717

21.   Shi S, Gronthos S, Chen S, Reddi A, Counter CM, Robey PG, Wang CY. Bone
formation by human postnatal bone marrow stromal stem cells is enhanced by
telomerase expression. *Nat Biotechnol*. 2002;20:587-591. doi: 10.1038/nbt0602-587

22.   Simonsen JL, Rosada C, Serakinci N, Justesen J, Stenderup K, Rattan SI, Jensen TG,
Kassem M. Telomerase expression extends the proliferative life-span and maintains the
osteogenic potential of human bone marrow stromal cells. *Nat Biotechnol*. 2002;20:592-
596. doi: 10.1038/nbt0602-592

23.   Leonard WJ, O'Shea JJ. Jaks and STATs: biological implications. *Annu Rev Immunol*.
1998;16:293-322. doi: 10.1146/annurev.immunol.16.1.293

24.   Hu X, Li J, Fu M, Zhao X, Wang W. The JAK/STAT signaling pathway: from bench to
clinic. *Signal Transduct Target Ther*. 2021;6:402. doi: 10.1038/s41392-021-00791-1

25.   Okutani Y, Kitanaka A, Tanaka T, Kamano H, Ohnishi H, Kubota Y, Ishida T, Takahara
J. Src directly tyrosine-phosphorylates STAT5 on its activation site and is involved in
erythropoietin-induced signaling pathway. *Oncogene*. 2001;20:6643-6650. doi:
10.1038/sj.onc.1204807

26.   Wingelhofer B, Neubauer HA, Valent P, Han X, Constantinescu SN, Gunning PT, Muller
M, Moriggl R. Implications of STAT3 and STAT5 signaling on gene regulation and
chromatin remodeling in hematopoietic cancer. *Leukemia*. 2018;32:1713-1726. doi:
10.1038/s41375-018-0117-x

27.   Gu L, Liao Z, Hoang DT, Dagvadorj A, Gupta S, Blackmon S, Ellsworth E, Talati P, Leiby
B, Zinda M, et al. Pharmacologic inhibition of Jak2-Stat5 signaling By Jak2 inhibitor

AZD1480 potently suppresses growth of both primary and castrate-resistant prostate
cancer. *Clin Cancer Res*. 2013;19:5658-5674. doi: 10.1158/1078-0432.CCR-13-0422

28.   Dieudonne FX, Severe N, Biosse-Duplan M, Weng JJ, Su Y, Marie PJ. Promotion of
osteoblast differentiation in mesenchymal cells through Cbl-mediated control of STAT5
activity. *Stem Cells*. 2013;31:1340-1349. doi: 10.1002/stem.1380

29.   Cuevas RA, Chu CC, Moorhead WJ, 3rd, Wong R, Sultan I, St Hilaire C. Isolation of
Human Primary Valve Cells for In vitro Disease Modeling. *J Vis Exp*. 2021:e62439. doi:
10.3791/62439

30.   Ducy P, Zhang R, Geoffroy V, Ridall AL, Karsenty G. Osf2/Cbfa1: a transcriptional
activator of osteoblast differentiation. *Cell*. 1997;89:747-754. doi: 10.1016/s0092-
8674(00)80257-3

31.   Otto F, Thornell AP, Crompton T, Denzel A, Gilmour KC, Rosewell IR, Stamp GW,
Beddington RS, Mundlos S, Olsen BR, et al. Cbfa1, a candidate gene for cleidocranial
dysplasia syndrome, is essential for osteoblast differentiation and bone development.
Cell. 1997;89:765-771. doi: 10.1016/s0092-8674(00)80259-7

32.   Sun Y, Byon CH, Yuan K, Chen J, Mao X, Heath JM, Javed A, Zhang K, Anderson PG,
Chen Y. Smooth muscle cell-specific runx2 deficiency inhibits vascular calcification. *Circ*
Res. 2012;111:543-552. doi: 10.1161/CIRCRESAHA.112.267237

33.   Lin ME, Chen T, Leaf EM, Speer MY, Giachelli CM. Runx2 Expression in Smooth
Muscle Cells Is Required for Arterial Medial Calcification in Mice. *Am J Pathol*.
2015;185:1958-1969. doi: 10.1016/j.ajpath.2015.03.020

34.   Parra-Izquierdo I, Castanos-Mollor I, Lopez J, Gomez C, San Roman JA, Sanchez
Crespo M, Garcia-Rodriguez C. Calcification Induced by Type I Interferon in Human
Aortic Valve Interstitial Cells Is Larger in Males and Blunted by a Janus Kinase Inhibitor.
Arterioscler Thromb Vasc Biol. 2018;38:2148-2159. doi: 10.1161/ATVBAHA.118.311504

35.   Parra-Izquierdo I, Castanos-Mollor I, Lopez J, Gomez C, San Roman JA, Sanchez
Crespo M, Garcia-Rodriguez C. Lipopolysaccharide and interferon-gamma team up to
activate HIF-1alpha via STAT1 in normoxia and exhibit sex differences in human aortic
valve interstitial cells. *Biochim Biophys Acta Mol Basis Dis*. 2019;1865:2168-2179. doi:
10.1016/j.bbadis.2019.04.014

36.   Jin H, St Hilaire C, Huang Y, Yang D, Dmitrieva NI, Negro A, Schwartzbeck R, Liu Y, Yu
Z, Walts A, et al. Increased activity of TNAP compensates for reduced adenosine

production and promotes ectopic calcification in the genetic disease ACDC. *Sci Signal*.

2016;9:ra121. doi: 10.1126/scisignal.aaf9109

37.   Moorhead WJ, 3rd, Chu CC, Cuevas RA, Callahan Jt, Wong R, Regan C, Boufford CK,

Sur S, Liu M, Gomez D, et al. Dysregulation of FOXO1 (Forkhead Box O1 Protein)

Drives Calcification in Arterial Calcification due to Deficiency of CD73 and Is Present in

Peripheral Artery Disease. *Arterioscler Thromb Vasc Biol*. 2020;40:1680-1694. doi:

10.1161/ATVBAHA.119.313765

38.   Jung JJ, Razavian M, Kim HY, Ye Y, Golestani R, Toczek J, Zhang J, Sadeghi MM.

Matrix metalloproteinase inhibitor, doxycycline and progression of calcific aortic valve

disease in hyperlipidemic mice. *Sci Rep*. 2016;6:32659. doi: 10.1038/srep32659

39.   Millan A, Lanzer P, Sorribas V. The Thermodynamics of Medial Vascular Calcification.

Front Cell Dev Biol. 2021;9:633465. doi: 10.3389/fcell.2021.633465

40.   Kim NW, Piatyszek MA, Prowse KR, Harley CB, West MD, Ho PL, Coviello GM, Wright

WE, Weinrich SL, Shay JW. Specific association of human telomerase activity with

immortal cells and cancer. *Science*. 1994;266:2011-2015. doi: 10.1126/science.7605428

41.   Hao LY, Armanios M, Strong MA, Karim B, Feldser DM, Huso D, Greider CW. Short

telomeres, even in the presence of telomerase, limit tissue renewal capacity. *Cell*.

2005;123:1121-1131. doi: 10.1016/j.cell.2005.11.020

42.   Jin HY, Weir-McCall JR, Leipsic JA, Son JW, Sellers SL, Shao M, Blanke P, Ahmadi A,

Hadamitzky M, Kim YJ, et al. The Relationship Between Coronary Calcification and the

Natural History of Coronary Artery Disease. *JACC Cardiovasc Imaging*. 2021;14:233-

242. doi: 10.1016/j.jcmg.2020.08.036

43.   Ogawa D, Nomiyama T, Nakamachi T, Heywood EB, Stone JF, Berger JP, Law RE,

Bruemmer D. Activation of peroxisome proliferator-activated receptor gamma

suppresses telomerase activity in vascular smooth muscle cells. *Circ Res*. 2006;98:e50-

59. doi: 10.1161/01.RES.0000218271.93076.c3

44.   Chiang YJ, Hemann MT, Hathcock KS, Tessarollo L, Feigenbaum L, Hahn WC, Hodes

RJ. Expression of telomerase RNA template, but not telomerase reverse transcriptase,

is limiting for telomere length maintenance in vivo. *Mol Cell Biol*. 2004;24:7024-7031.

doi: 10.1128/MCB.24.16.7024-7031.2004

45.   Tani T, Orimo H, Shimizu A, Tsuruoka S. Development of a novel chronic kidney

disease mouse model to evaluate the progression of hyperphosphatemia and associated

mineral bone disease. *Sci Rep*. 2017;7:2233. doi: 10.1038/s41598-017-02351-6

46.   Roake CM, Artandi SE. Regulation of human telomerase in homeostasis and disease.
1039   *Nat Rev Mol Cell Biol*. 2020;21:384-397. doi: 10.1038/s41580-020-0234-z
1040   47.   Feng J, Funk WD, Wang SS, Weinrich SL, Avilion AA, Chiu CP, Adams RR, Chang E,
1041   Allsopp RC, Yu J, et al. The RNA component of human telomerase. *Science*.
1042   1995;269:1236-1241. doi: 10.1126/science.7544491
1043   48.   Pascolo E, Wenz C, Lingner J, Hauel N, Priepke H, Kauffmann I, Garin-Chesa P, Rettig
1044   WJ, Damm K, Schnapp A. Mechanism of human telomerase inhibition by BIBR1532, a
1045   synthetic, non-nucleosidic drug candidate. *J Biol Chem*. 2002;277:15566-15572. doi:
1046   10.1074/jbc.M201266200
1047   49.   Ghosh A, Saginc G, Leow SC, Khattar E, Shin EM, Yan TD, Wong M, Zhang Z, Li G,
1048   Sung WK, et al. Telomerase directly regulates NF-kappaB-dependent transcription. *Nat
1049   Cell Biol*. 2012;14:1270-1281. doi: 10.1038/ncb2621
1050   50.   Liu N, Ding D, Hao W, Yang F, Wu X, Wang M, Xu X, Ju Z, Liu JP, Song Z, et al. hTERT
1051   promotes tumor angiogenesis by activating VEGF via interactions with the Sp1
1052   transcription factor. *Nucleic Acids Res*. 2016;44:8693-8703. doi: 10.1093/nar/gkw549
1053   51.   Lee C, Huang CH. LASAGNA: a novel algorithm for transcription factor binding site
1054   alignment. *BMC Bioinformatics*. 2013;14:108. doi: 10.1186/1471-2105-14-108
1055   52.   Wingender E, Dietze P, Karas H, Knuppel R. TRANSFAC: a database on transcription
1056   factors and their DNA binding sites. *Nucleic Acids Res*. 1996;24:238-241. doi:
1057   10.1093/nar/24.1.238
1058   53.   Babu AN, Meng X, Zou N, Yang X, Wang M, Song Y, Cleveland JC, Weyant M,
1059   Banerjee A, Fullerton DA. Lipopolysaccharide stimulation of human aortic valve
1060   interstitial cells activates inflammation and osteogenesis. *Ann Thorac Surg*. 2008;86:71-
1061   76. doi: 10.1016/j.athoracsur.2008.03.008
1062   54.   Natarajan K, Muller-Klieser D, Rubner S, Berg T. Stafia-1: a STAT5a-Selective Inhibitor
1063   Developed via Docking-Based Screening of in Silico O-Phosphorylated Fragments.
1064   *Chemistry*. 2020;26:148-154. doi: 10.1002/chem.201904147
1065   55.   Elumalai N, Berg A, Natarajan K, Scharow A, Berg T. Nanomolar inhibitors of the
1066   transcription factor STAT5b with high selectivity over STAT5a. *Angew Chem Int Ed Engl*.
1067   2015;54:4758-4763. doi: 10.1002/anie.201410672
1068   56.   Fredriksson S, Gullberg M, Jarvius J, Olsson C, Pietras K, Gustafsdottir SM, Ostman A,
1069   Landegren U. Protein detection using proximity-dependent DNA ligation assays. *Nat
1070   Biotechnol*. 2002;20:473-477. doi: 10.1038/nbt0502-473

57.   Gillis AJ, Schuller AP, Skordalakes E. Structure of the Tribolium castaneum telomerase
catalytic subunit TERT. *Nature*. 2008;455:633-637. doi: 10.1038/nature07283

58.   Neculai D, Neculai AM, Verrier S, Straub K, Klumpp K, Pfitzner E, Becker S. Structure of
the unphosphorylated STAT5a dimer. *J Biol Chem*. 2005;280:40782-40787. doi:
10.1074/jbc.M507682200

59.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Zidek A, Potapenko A, et al. Highly accurate protein structure prediction with
AlphaFold. *Nature*. 2021;596:583-589. doi: 10.1038/s41586-021-03819-2

60.   Yan Y, Tao H, He J, Huang SY. The HDOCK server for integrated protein-protein
docking. *Nat Protoc*. 2020;15:1829-1852. doi: 10.1038/s41596-020-0312-x

61.   Tebo AG, Gautier A. A split fluorescent reporter with rapid and reversible
complementation. *Nat Commun*. 2019;10:2822. doi: 10.1038/s41467-019-10855-0

62.   Liu JC, Lengner CJ, Gaur T, Lou Y, Hussain S, Jones MD, Borodic B, Colby JL,
Steinman HA, van Wijnen AJ, et al. Runx2 protein expression utilizes the Runx2 P1
promoter to establish osteoprogenitor cell number for normal bone formation. *J Biol
Chem*. 2011;286:30057-30070. doi: 10.1074/jbc.M111.241505

63.   Stock M, Otto F. Control of RUNX2 isoform expression: the role of promoters and
enhancers. *J Cell Biochem*. 2005;95:506-517. doi: 10.1002/jcb.20471

64.   Zhang S, Xiao Z, Luo J, He N, Mahlios J, Quarles LD. Dose-dependent effects of Runx2
on bone development. *J Bone Miner Res*. 2009;24:1889-1904. doi: 10.1359/jbmr.090502

65.   Okura H, Sato S, Kishikawa S, Kaneto S, Nakashima T, Yoshida N, Takayanagi H,
Kiyono H. Runx2-I isoform contributes to fetal bone formation even in the absence of
specific N-terminal amino acids. *PLoS One*. 2014;9:e108294. doi:
10.1371/journal.pone.0108294

66.   Banerjee C, Javed A, Choi JY, Green J, Rosen V, van Wijnen AJ, Stein JL, Lian JB,
Stein GS. Differential regulation of the two principal Runx2/Cbfa1 n-terminal isoforms in
response to bone morphogenetic protein-2 during development of the osteoblast
phenotype. *Endocrinology*. 2001;142:4026-4039. doi: 10.1210/endo.142.9.8367

67.   Maida Y, Yasukawa M, Okamoto N, Ohka S, Kinoshita K, Totoki Y, Ito TK, Minamino T,
Nakamura H, Yamaguchi S, et al. Involvement of telomerase reverse transcriptase in
heterochromatin maintenance. *Mol Cell Biol*. 2014;34:1576-1593. doi:
10.1128/MCB.00093-14

68.   Okamoto N, Yasukawa M, Nguyen C, Kasim V, Maida Y, Possemato R, Shibata T, Ligon
1104   KL, Fukami K, Hahn WC, et al. Maintenance of tumor initiating cells of defined genetic
1105   composition by nucleostemin. *Proc Natl Acad Sci U S A*. 2011;108:20388-20393. doi:
1106   10.1073/pnas.1015171108

1107   69.   Gizard F, Nomiyama T, Zhao Y, Findeisen HM, Heywood EB, Jones KL, Staels B,
1108   Bruemmer D. The PPARalpha/p16INK4a pathway inhibits vascular smooth muscle cell
1109   proliferation by repressing cell cycle-dependent telomerase activation. *Circ Res*.
1110   2008;103:1155-1163. doi: 10.1161/CIRCRESAHA.108.186205

1111   70.   Slagboom PE, Droog S, Boomsma DI. Genetic determination of telomere size in
1112   humans: a twin study of three age groups. *Am J Hum Genet*. 1994;55:876-882.

1113   71.   Brouilette S, Singh RK, Thompson JR, Goodall AH, Samani NJ. White cell telomere
1114   length and risk of premature myocardial infarction. *Arterioscler Thromb Vasc Biol*.
1115   2003;23:842-846. doi: 10.1161/01.ATV.0000067426.96344.32

1116   72.   Samani NJ, Boultby R, Butler R, Thompson JR, Goodall AH. Telomere shortening in
1117   atherosclerosis. *Lancet*. 2001;358:472-473. doi: 10.1016/S0140-6736(01)05633-1

1118   73.   Kurz DJ, Kloeckener-Gruissem B, Akhmedov A, Eberli FR, Buhler I, Berger W, Bertel O,
1119   Luscher TF. Degenerative aortic valve stenosis, but not coronary disease, is associated
1120   with shorter telomere length in the elderly. *Arterioscler Thromb Vasc Biol*. 2006;26:e114-
1121   117. doi: 10.1161/01.ATV.0000222961.24912.69

1122   74.   Evans MA, Walsh K. Clonal hematopoiesis, somatic mosaicism, and age-associated
1123   disease. *Physiol Rev*. 2023;103:649-716. doi: 10.1152/physrev.00004.2022

1124   75.   Huzen J, Peeters W, de Boer RA, Moll FL, Wong LS, Codd V, de Kleijn DP, de Smet BJ,
1125   van Veldhuisen DJ, Samani NJ, et al. Circulating leukocyte and carotid atherosclerotic
1126   plaque telomere length: interrelation, association with plaque characteristics, and
1127   restenosis after endarterectomy. *Arterioscler Thromb Vasc Biol*. 2011;31:1219-1225. doi:
1128   10.1161/ATVBAHA.110.217158

1129   76.   Yutzey KE, Demer LL, Body SC, Huggins GS, Towler DA, Giachelli CM, Hofmann-
1130   Bowman MA, Mortlock DP, Rogers MB, Sadeghi MM, et al. Calcific aortic valve disease:
1131   a consensus summary from the Alliance of Investigators on Calcific Aortic Valve
1132   Disease. *Arterioscler Thromb Vasc Biol*. 2014;34:2387-2393. doi:
1133   10.1161/ATVBAHA.114.302523

1134   77.   Hoerstrup SP, Kadner A, Melnitchouk S, Trojan A, Eid K, Tracy J, Sodian R, Visjager JF,

1135   Kolb SA, Grunenfelder J, et al. Tissue engineering of functional trileaflet heart valves

1136   from human marrow stromal cells. *Circulation*. 2002;106:I143-150.

1137   78.   Hortells L, Sur S, St Hilaire C. Cell Phenotype Transitions in Cardiovascular

1138   Calcification. *Front Cardiovasc Med*. 2018;5:27. doi: 10.3389/fcvm.2018.00027

1139   79.   Bogdanova M, Zabirnyk A, Malashicheva A, Enayati KZ, Karlsen TA, Kaljusto ML,

1140   Kvitting JP, Dissen E, Sullivan GJ, Kostareva A, et al. Interstitial cells in calcified aortic

1141   valves have reduced differentiation potential and stem cell-like properties. *Sci Rep*.

1142   2019;9:12934. doi: 10.1038/s41598-019-49016-0

1143   80.   Xu K, Xie S, Huang Y, Zhou T, Liu M, Zhu P, Wang C, Shi J, Li F, Sellke FW, et al. Cell-

1144   Type Transcriptome Atlas of Human Aortic Valves Reveal Cell Heterogeneity and

1145   Endothelial to Mesenchymal Transition Involved in Calcific Aortic Valve Disease.

1146   *Arterioscler Thromb Vasc Biol*. 2020;40:2910-2921. doi: 10.1161/ATVBAHA.120.314789

1147   81.   Lindroos M, Kupari M, Heikkila J, Tilvis R. Prevalence of aortic valve abnormalities in the

1148   elderly: an echocardiographic study of a random population sample. *J Am Coll Cardiol*.

1149   1993;21:1220-1225. doi: 10.1016/0735-1097(93)90249-z

1150   82.   Joung YH, Lim EJ, Darvin P, Chung SC, Jang JW, Do Park K, Lee HK, Kim HS, Park T,

1151   Yang YM. MSM enhances GH signaling via the Jak2/STAT5b pathway in osteoblast-like

1152   cells and osteoblast differentiation through the activation of STAT5b in MSCs. *PLoS*

1153   *One*. 2012;7:e47477. doi: 10.1371/journal.pone.0047477

1154   83.   Lee KM, Park KH, Hwang JS, Lee M, Yoon DS, Ryu HA, Jung HS, Park KW, Kim J,

1155   Park SW, et al. Inhibition of STAT5A promotes osteogenesis by DLX5 regulation. *Cell*

1156   *Death Dis*. 2018;9:1136. doi: 10.1038/s41419-018-1184-7

1157   84.   Paulson M, Pisharody S, Pan L, Guadagno S, Mui AL, Levy DE. Stat protein

1158   transactivation domains recruit p300/CBP through widely divergent sequences. *J Biol*

1159   *Chem*. 1999;274:25343-25349. doi: 10.1074/jbc.274.36.25343

1160   85.   Martino A, Holmes JHt, Lord JD, Moon JJ, Nelson BH. Stat5 and Sp1 regulate

1161   transcription of the cyclin D2 gene in response to IL-2. *J Immunol*. 2001;166:1723-1729.

1162   doi: 10.4049/jimmunol.166.3.1723

1163   86.   Bergad PL, Towle HC, Berry SA. Yin-yang 1 and glucocorticoid receptor participate in

1164   the Stat5-mediated growth hormone response of the serine protease inhibitor 2.1 gene.

1165   *J Biol Chem*. 2000;275:8114-8120. doi: 10.1074/jbc.275.11.8114

87.   Yamada O, Ozaki K, Furukawa T, Machida M, Wang YH, Motoji T, Mitsuishi T, Akiyama
M, Yamada H, Kawauchi K, et al. Activation of STAT5 confers imatinib resistance on
leukemic cells through the transcription of TERT and MDR1. *Cell Signal*. 2011;23:1119-
1127. doi: 10.1016/j.cellsig.2011.02.005

88.   Yamada O, Ozaki K, Akiyama M, Kawauchi K. JAK-STAT and JAK-PI3K-mTORC1
pathways regulate telomerase transcriptionally and posttranslationally in ATL cells. *Mol*
Cancer Ther. 2012;11:1112-1121. doi: 10.1158/1535-7163.MCT-11-0850

89.   Tormo AJ, Letellier MC, Sharma M, Elson G, Crabe S, Gauchat JF. IL-6 activates
STAT5 in T cells. *Cytokine*. 2012;60:575-582. doi: 10.1016/j.cyto.2012.07.002

90.   Tanabe Y, Nishibori T, Su L, Arduini RM, Baker DP, David M. Cutting edge: role of
STAT1, STAT3, and STAT5 in IFN-alpha beta responses in T lymphocytes. *J Immunol*.
2005;174:609-613. doi: 10.4049/jimmunol.174.2.609

91.   Honda S, Miyamoto T, Watanabe T, Narumi T, Kadowaki S, Honda Y, Otaki Y,
Hasegawa H, Netsu S, Funayama A, et al. A novel mouse model of aortic valve stenosis
induced by direct wire injury. *Arterioscler Thromb Vasc Biol*. 2014;34:270-278. doi:
10.1161/ATVBAHA.113.302610

92.   Sider KL, Blaser MC, Simmons CA. Animal models of calcific aortic valve disease. *Int J*
Inflam. 2011;2011:364310. doi: 10.4061/2011/364310

93.   Balani K, Brito FC, Kos L, Agarwal A. Melanocyte pigmentation stiffens murine cardiac
tricuspid valve leaflet. *J R Soc Interface*. 2009;6:1097-1102. doi: 10.1098/rsif.2009.0174

94.   Carneiro F, Kruithof BP, Balani K, Agarwal A, Gaussin V, Kos L. Relationships between
melanocytes, mechanical properties and extracellular matrix composition in mouse heart
valves. *J Long Term Eff Med Implants*. 2015;25:17-26. doi:
10.1615/jlongtermeffmedimplants.2015011748

95.   Mjaatvedt CH, Kern CB, Norris RA, Fairey S, Cave CL. Normal distribution of
melanocytes in the mouse heart. *Anat Rec A Discov Mol Cell Evol Biol*. 2005;285:748-
757. doi: 10.1002/ar.a.20210

96.   Dharmarajan S, Speer MY, Pierce K, Lally J, Leaf EM, Lin ME, Scatena M, Giachelli CM.
Role of Runx2 in Calcific Aortic Valve Disease in Mouse Models. *Front Cardiovasc Med*.
2021;8:687210. doi: 10.3389/fcvm.2021.687210

97.   Lanzer P, Hannan FM, Lanzer JD, Janzen J, Raggi P, Furniss D, Schuchardt M,
Thakker R, Fok PW, Saez-Rodriguez J, et al. Medial Arterial Calcification: JACC State-

of-the-Art Review. *J Am Coll Cardiol*. 2021;78:1145-1165. doi:

10.1016/j.jacc.2021.06.049

1200

Figure 1. TERT is elevated in CAVD tissues. (**A**) Examples of non-calcified control (top) and calcified CAVD (bottom) human valves used in this study. Arrowheads indicate calcified nodules Scalebar. 1 cm. (**B**), *TERT* mRNA transcript quantification in control and CAVD valve tissues. n = 8 control, n= 9 CAVD. (**C**), Representative serial sections of Von Kossa (dark precipitation) and TERT immunofluorescent staining in control and CAVD valve tissues. Scalebar 100 μm. Graphs show the quantification of TERT intensity and the number of TERT-positive cells. n = 9 control, n = 9 CAVD. (**D**), Telomere length measurements in valve tissues. n = 19 control, n = 12 CAVD. (**E**), Age of the patients utilized in this study. n = 25 control, n = 16 CAVD. Data are shown as means ± SD. P values were calculated using the Mann-Whitney U test and shown on each graph. L = lumen.

Figure 2. TERT is elevated in CAVD hVICs at baseline and upregulation during osteogenic reprogramming. (A), Representative images showing cell morphology of control and CAVD hVICs. Scalebar 50 μm. (**B**), Proliferation of control and CAVD hVICs at baseline. n = 3 both groups. (**C**) Relative migration distances of control and CAVD VICs. n = 10 both groups. (**D**) Western blot staining of control and CAVD hVICs at baseline. Quantification of protein levels is shown on the right panels. n = 4 both groups. (**E**) Telomere length measurements in VICs. n = 8, control, n = 5 CAVD. (**F**) Representative images of early calcification in hVICs growing in normal conditions (no treatment NT) or stimulated with osteogenic media (OST) for 14 days. Calcium deposition was visualized by Alizarin Red staining. Scalebar 50 μm. Quantification of calcification on the right graphs. n = 7 both groups. (**G**), Western blot staining of samples from control and CAVD hVICs after 14 days of NT or OST. Quantification of protein levels is shown on the bottom graphs. n = 4 both groups. (**H**), Representative immunofluorescent staining images of TERT in control and CAVD hVICs after 14 days of NT or OST. Calcium deposition was visualized with OsteoImage. Scalebar 100 μm. (**I**) Expression profile control VICs after 14 days of NT and OST treatment. n = 4 both groups. Data are shown as means ± SD. P values were calculated by 2-way ANOVA (Figure C), the Mann-Whitney U test (Figures B, D, E, G, and I), and the Kruskal-Wallis H test with Dunn post hoc test Figure F) and shown on each graph. * P = 0.0286 in Figure I.

Figure 3. TERT is required to initiate osteogenic transition and calcify. (**A**) Control hVICs were transduced lentivirus containing shControl or shTERT and followed by OST stimulation for up to 28 days. Calcium deposition was visualized by Alizarin Red staining. n = 6 shControl, n = 6 shTERT at 21 days and n = 9 each group at day 28. Scalebar 400 μm. (**B**) Representative western blot staining of control hVICs transduced with lentivirus containing either shControl or shTERT followed by 7 days of OST. (**C**) hCASMCs were transduced with shControl or shTERT and stimulated with OST for 14 and 21 days. n = 4 for each group at both time points. Scalebar 400 μm. (**D**) Representative images of mVIC isolated from *Tert*^{+/+} and *Tert*^{-/-} mice were stimulated with OST for 28 days. Scalebar 400 μm. (**E**) Representative Von Kossa staining images of *Tert*^{+/+} and *Tert*^{-/-} mice aortas (dark precipitation). Scale bar 400 μm. Quantification of calcification is shown on the right panel. (**F**) Representative immunofluorescent images of Runx2 expression indicative of calcification (red, insert) on *Tert*^{+/+} and *Tert*^{-/-} mice aorta. Scale bar 400 μm. Runx2 quantification of calcification is shown on the right panel. (**G**) Representative images of mVICs isolated from F1 *Tert*^{-/-} or *Terc*^{-/-} mice were stimulated with OST for up to 21 days. Scalebar 100 μm. Data are shown as means ± SD. P values were calculated using the Kruskal-Wallis H test, Dunn post hoc test (Figures A, C, and G), and Mann-Whitney U test (Figures D, E, and F).

Figure 4. STAT5 binds to TERT and is required for the calcification of human VICs. (A) Graphic of in silico analysis identifying the consensus binding site for STAT5. (**B**), Diagram depicting STAT5 binding to RUNX2 promoter and the positions of the predicted STAT5 binding sites (purple boxes). (**C**) Representative immunofluorescence images of control hVICs after 14 days of LPS treatment or OST stimulation. Scalebar 50 μm. (**D**) Representative western blot staining of hVICs after 14 days of OST stimulation. (**E**) Representative images of control hVICs treated with 10 μM of the STAT5 inhibitors StafiA-1 or StafiB-1 throughout 28 days of osteogenic stimulation. Scalebar 400 μm. n = 8 each group. The quantification of calcification is shown on the right graph. (**F**) Control hVICs were transduced with lentivirus containing shControl or shSTAT5 constructs and treated with OST for 21 days. Calcium deposition was visualized by Alizarin Red staining. n = 9 for each group. Scalebar 400 μm. (**G**) Representative images of hVIC (left panels) treated with 10 μM of the STAT5 inhibitors StafiA-1 starting at d7 or d14 of OST treatment. Samples were collected at d21. Scalebar 400 μm. n = 6 for each group. (**H**) Representative images of TERT/STAT5 complex (red foci) in control hVICs cultured for 21 days in osteogenic medium and detected by PLA. Scalebar 100 μm. n = 4 for each group. (**I**) TERT/STAT5 interaction was detected by co-immunoprecipitation in control hVICs under OST for 21 days. (**J**) Control hVIC co-expressing TERT-NFAST and STAT5A-CFAST were labeled with 5 μM HBR-3,5DOM (4-hydroxy-3,5-dimethoxybenzylidene rhodamine, coral) and imaged at d3, d5, and d7 of osteogenic treatment (OST) or control no-treatment (NT). Scale bars 50 μm. (**K**) Chromatin immunoprecipitation shows enrichment of STAT5 on RUNX2 promoter OST conditions compared to NT. The base position is indicated in the blue boxes. n=4 for each group as in B. (**L**) STAT5 CUT&Tag sequencing tracks in genomic regions of Runx2 in no-treatment control and osteogenic condition-treated mice VICs. (**M**) Stacked chart representing genome-wide STAT5 occupancy in promoters, genes, and intergenic regions frequency in mice VICs. (**N**) Top 20 GO pathways enriched in genes with significant loss of STAT5 in *Tert*[+/+] vs *Tert*[-/-] mice VICs. Data are shown as means ± SD. P values were calculated by the Mann-Whitney U test (Figures F, H, and J) and the Kruskal-Wallis H test (Figures E and G) and indicated on top of each graph.

Figure 5. TERT and STAT5 are upregulated and colocalized in CAVD tissue. (**A**) Representative serial sections of Von Kossa (dark precipitation) and TERT and STAT5 immunofluorescent staining in control and CAVD tissues. Scalebar 100 μm. STAT5 intensity and STAT5-positive cells (STAT5+) are quantified on the right panels. (**B**) Quantification of STAT5-TERT-positive cells. Data are shown as means ± SD, n = 5 control, n = 6 CAVD. P values were calculated with the Mann-Whitney U test and indicated on each graph. L = lumen.

Figure 6. TERT/STAT5 promotes osteogenic reprogramming. Inflammatory signals promote the upregulation and interaction of TERT and STAT5A. The TERT/STAT5 complex translocates into the nucleus, where TERT interacts with SMARCA4 (also known as BRG1), which facilitates a permissive chromatin configuration, and STAT5A binds specific sites on the RUNX2 promoter, leading to the osteogenic reprogramming of valve interstitial cells during the early stages of CAVD pathogenesis.

Supplemental Figure 1. Characterization of human CAVD valves and patient-isolated VICs at base-line. (**A**) Representative section of Von Kossa staining in control and valve CAVD leaflets tissue. Scalebar 1 mm. (**B**) Representative serial sections of RUNX2 immunofluorescent staining and OPN in control and CAVD valve tissues. Scalebar 100 μm. Quantification of RUNX2-positive cells and OPN-positive cells on the leaflet specimen is shown on the right graphs. n = 5 control, n = 6 CAVD. (**C**) Correlation between age of the donor and TERT expression in valve tissues. n = 8 control, n = 9 CAVD. (**D**) Gene expression profile in the valve leaflets. n = 7 each group. (**E**) Representative images of immunofluorescent staining of PCNA (top panels) and γ-H2AX (bottom panels) in control and CAVD valve tissues. Scalebar 100 μm. Quantification of PCNA-positive cells and γ-H2AX-positive cells are shown on the right graphs. n = 4 control, n = 3 CAVD. (**F**) Representative images of a scratch assay in control and CAVD hVICs. Time points are indicated on top of each panel. Blue lines denote the cell migration front. Scalebar 1 mm. n = 10 of each group. (**G**) Representative images of αSMA, CNN, SM22, TNAP, and f-Actin in control and CAVD hVICs. Scalebar 50 μm. Signal quantification is shown on the right. n = 4 each group. Data are shown as means ± SD. P values were calculated with the Mann-Whitney U test (Figures A, B, D, F, and G), Pearson correlation test (Figure C), and two-way ANOVA with the Sidak post hoc test (Figure D and shown on top of each graph). L = lumen.

Supplemental Figure 2. Inhibition of reverse transcriptase activity of TERT does not prevent osteogenic reprogramming of hVICs. (**A**) Cell counts of control and CAVD hVICs after 14 days of NT or OST stimulation. n = 4 each group. (**B**) β-galactosidase (*BLG1*) mRNA transcript quantification in control and CAVD hVICs at baseline. n = 5 control, n= 4 CAVD. (**C**) Representative images of senescence-associated β-galactosidase activity staining of hVICs after 28 days of NT or OST stimulation. Blue cells were considered positive for senescence. Quantification of senescence is shown on the right graph. n = 4 each group. Data are shown as means ± SD. P values were calculated with the Mann-Whitney U test (Figure B) and the Kruskal-Wallis H test with Dunn pairwise comparison post hoc test (Figures A, C) and shown on top of each graph. Scalebar 400 µm. n = 6 for each group.

Supplemental Figure 3. TERT is upregulated and required for the osteogenic differentiation of human mesenchymal stem cells. (**A**) Representative images of hMSCs stimulated with OST for 14 days. Calcium deposition was visualized by Alizarin Red staining. Scalebar 50 μm. (**B**) Western blot staining of samples from hMSCs collected during the 14 days of OST treatment. (**C**) *RUNX2* expression profile of differentiating hMSCs stimulated for 14 days with OST media. n = 3. (**D**) Representative TERT immunofluorescent staining images of hMSCs stimulated with OST for 14 days. Scalebar 100 μm. (**E**) Representative images of hMSCs infected with shTERT or shControl virus and stimulated with OST for 14 days. Calcium deposition was visualized by Alizarin Red staining. The quantification of calcification is shown on the right graph. n = 3. (**F**) Western blot staining of samples from hMSCs collected during the 14 days of OST treatment. Data are shown as means ± SD. P values were calculated with the Kruskal-Wallis H test with Dunn pairwise comparison post hoc test (Figure C) and the Mann-Whitney U test (Figure E) and shown on each graph.

Supplemental Figure 4. Inhibition of reverse transcriptase activity of TERT does not prevent osteo-genic reprogramming of VICs. (**A**) Representative images of mBMMSCs isolated from *Tert*[+/+] or *Tert*[-/-] mice and stimulated with OST for 21 days. Scalebar 100 μm. Quantification of calcification is presented as average absorbance on the right graphs, n = 5 for each group. (**B, C**) Representative images of a senescence-associated β-galactosidase (SA-β-gal) staining of mVICs and mBMMSCs isolated from *Tert*[+/+] or F1 *Tert*[-/-] mice after 21 days of OST stimulation. In each group, blue cells were considered positive for senescence, n = 5. (**D**) Schematic of OST treatment with delayed shTERT transduction (top panel). VICs were transduced with shTERT, and calcification was assessed with Alizarin Red S stain at day 21. Scalebar 400 μm. n = 6 for each group. (**E**) Representative sections of Von Kossa staining in thoracic aortic rings isolated from *Tert*[+/+] and *Tert*[-/-] mice after 14 days of OST treatment relative to NT control treatment. Quantification is shown on the right. Scale bar 400 μm (**F**) Immunofluorescence staining of thoracic aortic rings after OST treatment as in B. Quantification of Runx2 is shown on the right graph. Scale bar 400 μm (**G**) Western blot staining of aortic ring specimens collected and treated in B. Quantification of Runx2 is shown on the right graph. (**H**) Human VICs were treated with the TERT inhibitor BiBR1532 and stimulated with OST, and calcification was assessed on day 28, with n = 5 in each group. (**I**) *TERT*, *RUNX2*, and *STAT5A* expression were evaluated on day 14, n = 6 DMSO, n = 5 BIBR1532. Data are shown as means ± SD. P values were calculated with the Kruskal-Wallis H test with Dunn pairwise comparison post hoc test (Figures A, B, C, E, H) and the Mann-Whitney U test (Figures D, F, G, I) and are shown on top of each graph.

Supplemental Figure 5. STAT5 is upregulated, binds to TERT, and is required for osteogenic reprogramming of VICs. (**A**) Western blot analysis of cytoplasmic and nuclear fractions of TERT from control VICs obtained after seven days of OST stimulation. Fractions purity was assessed by staining for markers of cytoplasm (TUB) and nucleoplasm (Histone H3). Cyt, cytoplasmic fraction; Nuc, nuclear fraction. (**B**) Representative immunofluorescence image of STAT5 shows subcellular distribution after 14 days of NT and OST stimulation. Scalebar 40 µm. (**C**) Representative images of CASMCs treated with 10 µM of the STAT5 inhibitors StafiA-1 or StafiB-1 during the 14 days of OST stimulation. Scalebar 400 µm. n = 4 for each group. (**D**) Representative images of MSCs treated with 10 µM of the STAT5 inhibitors StafiA-1 or StafiB-1 during the 21 days of OST stimulation. Scalebar 400 µm. n = 4 for each group. (**E**) Representative images of TERT/STAT5 complex (red foci) detected by proximity ligation assay (PLA) in the nucleus of control VICs. Scalebar 25 µm. The PLA foci quantification and intensity are shown on the right graph. n = 4 for each group. (**H**) Mol*3D Viewer rendering of the TERT-STAT5 dimer interaction predicted by the HDock. Red boxes indicate the position of the interaction between TERT and STAT5. (**F**) Representative immunofluorescent images of Stat5 expression after 12 weeks of high adenine high phosphate diet on *Tert⁻/⁻* and *Tert⁻/⁻* mice aorta. Scale bar 400 µm. Quantification of Stat5 expression is shown on the right panel. (**G**) Representative images of Stat5 immunofluorescence staining of thoracic aortic rings isolated from *Tert⁺/⁺* and *Tert⁻/⁻* mice after 14 days of OST treatment relative to NT control treatment. Quantification of Stat5 is shown on the right graph. Scale bar 400 µm (**I**) The diagram shows the regions involved in the predicted interaction as indicated in A, and the docking and confidence scores predicted by HDock are indicated. TEN: DNA anchor domain for processive telomeric repeat addition, RBD: Telomerase RNA binding domain, RT: Finger and palm region of the reverse transcriptase domain, CTE: C-terminal extension or putative thumb domain of the reverse transcriptase, CCD: Coiled-coil domain, required for nuclear localization, DBD: DNA binding domain, HLK: Helical linker and DNA binding domain, SH2: Domain required for activation and dimerization, TD: Transactivation domain. Data are shown as means ± SD. P values were calculated with the Mann-Whitney U test (Figures A and G) and the Kruskal-Wallis H test with Dunn pairwise comparison post-test (Figures B, E, and F) and indicated in each graph.

Supplemental Figure 6. Chromatin remodeling enzyme SMARCA4 is upregulated and interacts with TERT during the osteogenic reprogramming of VICs. (A) Representative immunofluorescent images of Smarca4 expression after 12 weeks of high adenine high phosphate diet on *Tert*[-/-] and *Tert*[-/-] mice aorta. Scale bar 400 µm. Quantification of Smarca4 expression is shown on the right panel. (B) Representative images of Smarca4 immunofluorescence staining of thoracic aortic rings isolated from *Tert*[+/+] and *Tert*[-/-] mice after 14 days of OST treatment relative to NT control treatment. Quantification of Smarca4 is shown on the right graph. (C) *SMARCA4* expression was evaluated on days 5 and 10 on VICs stimulated with OST. n = 5 each group. (E) TERT/STAT and TERT/SMARCA4 interaction was detected by co-immunoprecipitation upon osteogenic stimulation on VIC. (E) Representative TERT and SMARCA4 immunofluorescent staining in control and CAVD tissues. The number of positive cells is shown on the right graphs. n = 9 control, n = 9 CAVD. Scalebar 100 µm. Data are shown as means ± SD, and P values were calculated with the Mann-Whitney U test and indicated on each graph in Figures A and C. L = lumen.