**Pathogen Clearance-based Correlates of Immunoprotection**

**Reveal Insightful Features of Vaccine Immunity**

Xianbin Tian[1], Juanjuan Wang[1,2]*, Haoze Chen[1], Ming Ding[1], Qian Jin[1], and Jing-Ren Zhang[1,2]*

[1]Center for Infectious Disease Research, Department of Basic Medical Science, School of

Medicine, Tsinghua University, Beijing, China;

[2]Tsinghua-Peking Center for Life Sciences, Tsinghua University, Beijing, China;

**\*Correspondence** to: wangjj15@tsinghua.org.cn or zhanglab@tsinghua.edu.cn

**ABSTRACT**

Vaccination has significantly reduced the incidence of invasive infections caused by several bacterial pathogens, including *Streptococcus pneumoniae, Haemophilus influenzae* and *Neisseria meningitidis*. However, no vaccines are available for many other invasive pathogens. A major hurdle in vaccine development is the lack of functional markers to quantify vaccine immunity in eliminating pathogens in the process of infection. Based on our recent discovery of the liver as the major organ of vaccine-induced clearance of blood-borne virulent bacteria, we here describe a new vaccine evaluation system that quantitatively characterize important properties of effective vaccines in shuffling virulent bacteria from the blood circulation to the liver resident macrophage Kupffer cells (KCs) and sinusoidal endothelial cells (LSECs) in animal models. This system consists of three related correlates or assays: pathogen clearance from the bloodstream, pathogen trapping in the liver, and pathogen capture by KCs/LSECs. The readouts of these correlates were consistently associated with the serotype-specific immunoprotection levels of the 13-valent pneumococcal polysaccharide conjugate vaccine (PCV13) against lethal infections of *S. pneumoniae,* a major invasive pathogen of community-acquired infections in humans. Furthermore, these assays were able to identify a set of functional differences between two licensed PCV13 brands. Finally, the reliability and sensitivity of these correlates in reflecting vaccine efficacy were verified with whole cell vaccines of *Klebsiella pneumoniae* and *Escherichia coli*, two major pathogens in hospital-acquired invasive infections. This system may be used as cost effective readouts to evaluate the immunoprotective potential of vaccine candidates in the preclinical phase by filling the current technical gap in vaccine evaluation between the conventional *in vitro* approaches (e.g., antibody production and pathogen neutralization/opsonophagocytosis) and survival of immunized animals.

**Key words:** vaccine, immunoprotection, blood infection, *Streptococcus pneumoniae*, liver, Kupffer cell, and liver sinusoidal endothelial cell.

## INTRODUCTION

Sepsis caused by microbial infections of the blood circulation represents an important global health problem and economic burden[1,2]. Approximately 49 million sepsis cases and 11 million sepsis-associated deaths were reported worldwide in 2017, representing 19.7% of all global deaths[2]. Sepsis pathogenesis is initiated by the entry and proliferation of microbial pathogens in the blood circulation, subsequent induction of excessive inflammatory responses, and life-threatening multi-organ dysfunctions[3]. Pathogenic bacteria are the common pathogens of septic infections, which consist of both the Gram-positive bacteria (*Streptococcus pneumoniae*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Enterococcus faecalis*) and Gram-negative bacteria (*Acinetobacter baumannii*, *Escherichia coli*, *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa*)[4].

*S. pneumoniae* (pneumococcus), an encapsulated Gram-positive bacterium residing in the nasopharynx of humans, is an opportunistic pathogen of community-acquired pneumonia, bacteremia, and meningitis, particularly in children under 5 years old and the elderly[5]. There are forms of capsular polysaccharide (CPS)-based vaccines globally licensed to prevent invasive pneumococcal diseases (IPDs) (e.g. bacteremia and meningitis): pneumococcal polysaccharide vaccine (PPV) and pneumococcal conjugate vaccine (PCV)[6]. PPV is the first generation of pneumococcal vaccine that contains T cell-independent CPS antigens with poor immunogenicity in children under two years of ages[7]. PCV represents the next generation of CPS-based vaccine, in which CPSs are chemically linked to carrier proteins (e.g. diphtheria and tetanus toxoids) to activate both B and T lymphocytes to produce IgG antibodies[8]. The global recommendation of PCV has significantly reduced the pneumococcus-associated death worldwide[9]. Since the first introduction of PCV7 (Prevnar by Wyeth) that contains the CPSs of 7 IPD-associated serotypes, several PCVs with additional serotypes have been introduced, such as PCV10 (Synflorix by GlaxoSmithKline), PCV13 (Prevnar 13 by Pfizer, Weuphoria by Walvax, Weiminfeibao by Minhai), PCV15 (Vaxneuvance by Merck), PCV20 (Prevnar 20 by Pfizer)[6,10]. The three abovementioned PCV13 brands are all licensed in China.

Vaccination has greatly contributed to the control of many infectious diseases. While programmed immunization has led to global eradication (e.g., smallpox) or nearly eradication (e.g., polio) of a few viral diseases, a number of bacterial diseases have become preventable by vaccination, such as diphtheria (*Corynebacterium diphtheria*), whooping cough (*Bordetella pertussis*), tetanus (*Clostridium tetani*), meningitis (*Neisseria meningitidis*), bacterial pneumonia and sepsis (*S. pneumoniae*, *Haemophilus influenzae*). Except for *S. pneumoniae*, there are no available vaccines for any of the major blood infection pathogens after many

unsuccessful efforts in developing vaccines for each of these pathogens. One of the major

hurdles in the process of the current vaccine development is the lack of reliable and sensitive

immunoprotection correlates to identify any problems in the early steps of vaccine

development. Protection efficacy of vaccines relies on two functional aspects of immunization:

1) the features of vaccine-induced antibody and/or T-cell responses, and 2) the potency of

vaccine-elicited immunity of pathogen elimination upon pathogen entry[11]. Beyond protection

against infections in animal models (e.g., survival rate), the current vaccine evaluation system

mostly depends on *in vitro* assessment of vaccine-induced immune responses, such as antibody

titer, antibody-mediated neutralization and opsonophagocytosis of pathogen. As an example,

the current pneumococcal vaccines are based on measuring antibody level by enzyme-linked

immunosorbent assay (ELISA) and antibody-mediated phagocytosis of neutrophil by

opsonophagocytic killing assay (OPKA)[12,13]. These parameters don't inform how vaccine-

elicited immunity enable hosts to recognize and clear pathogens in the process of infection.

Therefore, there is a great need for reliable and sensitive *in vivo* techniques for functional

evaluation of vaccine immunity in the recognition and elimination of invading pathogens in

the process of infection.

The liver represents a major organ to capture and eliminate invading microbes from the

blood circulation, which are executed by the liver resident macrophage Kupffer cells (KCs)

and liver sinusoidal endothelial cells (LSECs)[14]. KCs and LSECs represent approximately 20%

and 50% of the liver non-parenchymal cells (NPCs), respectively[15]. KCs are the most dominant

resident macrophages in mammals, representing approximately 90% of total resident

macrophages in the body[16]. While KCs are specialized for the clearance of large particles (e.g.

bacteria)[17], LSECs preferentially clear small particles (e.g. viruses)[18]. Our recent studies have

discovered that the liver is the major immune organ to execute vaccine-activated capture and

killing of pathogenic bacteria in the bloodstream of mice[19,20]. In particular, PCV13 activates

both KCs and LSECs to rapidly recognize and capture IgG-coated pneumococci in the

bloodstream[20].

This study aims to establish an *in vivo* vaccine evaluation system to quantify the functional

features of vaccine-activated pathogen clearance from the blood circulation to the liver, based

on our recent findings of liver-based vaccine immunity[19,20]. We have developed three related

new correlates or assays: pathogen clearance from the bloodstream, pathogen trapping in the

liver, and pathogen capture by KCs/LSECs in the liver sinusoids. The reliability and sensitivity

of these assays have been tested in three pathogens and four vaccines. Potential applications of

this new system in vaccine evaluation and development are discussed.

 **RESULTS**

 **The clearance rate of blood-borne pneumococci predicts the protection efficacy of PCV13**

 Our recent studies have shown that effective clearance of bacteria from the bloodstream in

 the early phase of septic infections is essential for host survival[21], and efficient vaccination

 enhances such the pathogen clearance of certain encapsulated high-virulence bacteria that

 otherwise persist in the blood circulation[19,20]. To assess the direct relationship between the

 vaccine-elicited pathogen clearance rate and host survival, we compared the PCV13-induced

 bacterial clearance of different vaccine-covered serotypes of pneumococci from the

 bloodstream. In particular, we focused on the five high-virulence (HV) serotypes (1, 3, 4, 5 and

 6A) covered by PCV13 because the clearance of these serotypes highly depends on PCV13

 immunization in mice[19,20], and the rest of the eight low-virulence (LV) serotypes (6B, 7F, 9V,

 14, 18C, 19A, 19F and 23F) are effectively cleared in naïve mice[21]. We first measured CPS-

 specific IgG antibodies in PCV13-immunized and control mice by enzyme-linked

 immunosorbent assay (ELISA) because antibody titer has been used as an efficacy marker of

 pneumococcal polysaccharide conjugate vaccines[22,23]. Though PCV13 immunization

 generated antibodies to each of the five serotypes, the ELISA results revealed substantial

 variations in antibody level, with serotypes 4 and 6A as the highest and lowest in the antibody

 range, respectively (**Fig. 1A**). We next monitored the dynamic of bacteremia in immunized and

 control mice during the first 30 min post intravenous (i.v.) infection of pneumococci. In the

 adjuvant group, all of the five representative strains retained abundantly in blood circulation

 within 30 min post i.v. infection of $10^6$ CFU (**Fig. 1B**). In contrast, the circulating pneumococci

 were cleared quickly from the bloodstream even to undetectable level in PCV13-immunized

 mice at 30 min post infection (**Fig. 1C**). Consistent with our previous findings of serotype-3

 and -4 pneumococci[20], this result demonstrates that PCV13 induces fast pneumococcal

 clearance of all the five HV serotypes from the bloodstream.

 Previous studies have indicated that pneumococcal conjugate vaccines PCV7/PCV13

 confer relatively poor protection against invasive disease caused by serotype 3 in humans[24-26].

 We next tested if pathogen clearance rate could identify this functional difference in mice using

 higher challenge doses. PCV13-immunized mice rapidly cleared $10^7$ CFU of serotype-4, -5 and

 -6A pneumococci from the circulation with $CT_{50}$ of 1.5, 0.7, and 0.8 min, respectively. In sharp

 contrast, serotype-1 and -3 pneumococci were still largely circulating in the blood of the

 immunized mice with marginal reduction in bacteremia level post i.v. inoculation with $10^7$ CFU

 (**Fig. 1D**). With a much higher infection dose ($10^8$ CFU), the immunized mice still rapidly

 cleared 99% inoculum of serotype-5 and -4 pneumococci with $CT_{50}$ of 0.7 and 1.5 min,

respectively (**Fig. 1E**), but were incapable of effectively clearing the other 3 serotypes at this

infection dose, with 58%, 74%, and 32% of serotype-1, -3 and -6A inoculum being still detected

in the circulation at 30 min, respectively. Overall, these experiments have ranked the functional

capacity of PCV13 immunization in clearing the five HV serotypes in the following order: 5,

4, 6A, 1 and 3. The uneven clearance of various serotypes in PCV13-immunized mice strongly

suggests that the different antigens in PCV13 elicit variable levels of functional immunity.

To determine if the differences of PCV13-elicited clearance of various pneumococcal

serotypes reflect protection efficacy against different serotypes of *S. pneumoniae*, we measured

the survival of PCV13-immunized mice post intraperitoneal (i.p.) infection. Compared with

the full mortality of adjuvant groups post i.p. infection with $10^6$ CFU, PCV13-immunized mice

conferred a full protection against serotype 1 (**Fig. 1F**), 4 (**Fig. 1H**), 5 (**Fig. 1I**) and 6A (**Fig.**

**1J**), and 80% protection against serotype 3 (**Fig. 1G**). We further determined the upper limits

of PCV13 immunization against each serotype with higher levels of inoculum. In line with the

different impacts of PCV13 on blood clearance of the five serotypes (**Figs. 1D-E**), the

vaccination conferred the strongest protection against lethal infection of serotype-4, -5 and -

6A pneumococci, with a 50% lethal dose ($LD_{50}$) of $7.1 \times 10^7$ CFU (**Fig. 1H**), $1.2 \times 10^8$ CFU

(**Fig. 1I**) and $5.6 \times 10^7$ CFU (**Fig. 1J**), respectively. By comparison, the immunized mice were

much more susceptible to infection of serotype-3 pneumococci with a $LD_{50}$ of $2.2 \times 10^6$ CFU

(**Fig. 1G**), and, to a less extent, to that of serotype-1 pneumococci ($LD_{50} = 3.2 \times 10^7$ CFU) (**Fig.**

**1F**). Overall comparison revealed a dramatic 54.5-fold difference in $LD_{50}$ between the best

(serotype 5) and worst (serotype 3) protected categories. The order of protection efficacy

induced by PCV13 immunization against infection of the HV serotypes (5, 4, 6A, 1 and 3) is

the same as the order of blood pathogen clearance rate. Taken together, these results

demonstrate that the PCV13-elicited pathogen clearance rate in the bloodstream is highly

predictive of the protection efficacy.

**Hepatic trapping of blood-borne pneumococci correlates the protection efficacy of**

**PCV13**

Our recent studies have found that the liver is the major executive organ for vaccine-

induced trapping of high-virulence encapsulated *S. pneumoniae*[20] and *Klebsiella pneumoniae*

serotypes[19] from the blood circulation. To determine the potential of vaccine-activated hepatic

pathogen trapping as a functional correlate for vaccine protection efficacy, we detected the

bacterial CFU count in five major organs and blood as previously described[20]. Consistent with

the poor clearance of the five HV serotypes in mice of adjuvant group, these animals showed

marginal pathogen trapping in the liver for serotypes 1, 3, 4, 5, and 6A post i.v. infection with

$10^6$ CFU (**Fig. 2A**). In contrast, virtually all of the viable pneumococci in PCV13-immunized

mice were detected in the liver, with near sterilization of the blood circulation, at 30 min post

i.v. infection with $10^6$ CFU (**Fig. 2B**). Comparison of the total CFU counts between PCV13

and adjuvant groups revealed significantly lower levels of bacterial burden in vaccinated mice,

which reflected vaccine-activated pathogen killing in the liver[20], and the full protection of

vaccinated mice against infection with $10^6$ CFU of the five HV serotypes (**Fig. 1**). However,

PCV13-immunized mice displayed a serotype-specific pattern of vaccine-activated pathogen

trapping in the liver at higher levels of infection dose. Although the majority of viable serotype-

1 (77%), -3 (72%), -4 (84%), -5 (91%) and -6A (86%) pneumococci were still found in the

livers of PCV13-immunized mice at 30 min post i.v. infection with $10^7$ CFU, serotype-1 (18%)

and -3 (20%) pneumococci escaped hepatic capture and remained in the bloodstream (**Fig. 2C**).

The vaccine-induced hepatic trapping of the five serotypes is ranked from the highest to the

lowest as: 5, 6A, 4, 1 and 3. Relatively poorer hepatic trapping of serotypes 1 and 3 in the

immunized mice was also reflected by relatively higher levels of total bacteria detected for

these two serotypes at 30 min. When infection dose was increased to $10^8$ CFU, relative

proportion of hepatic pathogen trapping was in the order as: 5 (97%), 4 (96%), 6A (21%), 3

(19%) and 1 (3%) and serotype-specific variation in vaccine-induced hepatic pathogen trapping

became more dramatic (**Fig. 2D**). These results show that vaccine-activated hepatic trapping

of blood-borne pneumococci strongly correlates with the protection efficacy of PCV13.

**Pathogen capture by liver immune cells indicates the protection efficacy of PCV13**

The vaccine-activated dual-track bacterial capture by liver resident macrophages

Kupffer cells (KCs) and sinusoidal endothelial cells (LSECs) has been found as the cellular

basis of vaccine-elicited pathogen trapping in the liver[20,21]. To assess the potential of cellular

capture of circulating pathogens as a functional correlate for vaccine protection efficacy, time-

lapse intravital microscopy (IVM) imaging was used to visualize and quantify cellular capture

of serotype-1, -3, -4, -5 and -6A pneumococci post i.v. infection with $5 \times 10^7$ CFU in the liver

sinusoids of PCV13-immunized and control mice. While IVM showed that the HV

pneumococci were mostly passing the vasculature of control mice (**Fig. 3A, Movies 1-5, left**

**panel**), but were rapidly tethered mainly onto LSECs and less on KCs (**Fig. 3A, Movies 1-5,**

**right panel**). In agreement with the serotype-specific protection efficacies of PCV13 (**Fig. 1E**),

quantification of immobilized bacteria per field of view (FOV) revealed the highest numbers

of total captured bacteria for serotypes 4 (110) and 5 (87) as compared with fewer bacteria for

serotypes 1 (64), 3 (61) and 6A (74) (**Fig. 3B**). The real-time imaging of vaccine-activated

pathogen capture by the liver immune cells in live mice has provided immune cell-based data

that are consistent with the protection efficacy of PCV13.

**The *in vivo* immunoprotection correlates distinguish the potency levels of different**

**PCV13 brands**

The abovementioned data have established an *in vivo* system to evaluate vaccine protection

efficacy against bacterial septic infection, including the quantitation of pathogen clearance

from blood, trapping in the liver, and cellular capture by LSECs and KCs. Since these

parameters uncovered remarkable differences of PCV13-induced immunity in protecting lethal

infection against different vaccine-covered HV pneumococcal serotypes, we wondered

whether PCV13 vaccines have brand-specific differences in protecting against lethal infection

of the same serotypes of pneumococci, which could reveal potential differences of vaccine

efficacy. We thus compared abovementioned PCV13 (referred to as PCV13) with another brand

of PCV13 (referred to as PCV13X) using the three parameters, which are both licensed in

China. When i.v. challenged with $10^7$ CFU of bacteria, mice immunized by PCV13 showed

significant faster clearance of serotype-1 pneumococci ($CT_{50}$, 3.7 min) than the PCV13X-

immunized counterparts ($CT_{50}$, >30 min) (**Fig. 4A, left panel**). Accordingly, PCV13

immunization yielded more potent hepatic trapping as evidenced by the relative higher

proportion of liver-trapped bacteria out of the total viable bacteria in PCV13-immunized mice

(77%) than that in PCV13X-immunized counterparts (26%) (**Fig. 4A**, **middle panel**). Although

there was a modest difference in the total bacteria captured by the liver immune cells between

PCV13- and PCV13X-immunized mice, it is worth to mention that PCV13 induced a

significantly more potent bacterial capture by LSECs than PCV13X (**Fig. 4A**, **right panel,**

**Movie 6**), which may explain the difference in protection efficacy between the two vaccines

(**Fig. 5**). In a similar pattern, PCV13 immunization also yielded a more effective clearance of

serotype-3 pneumococci ($CT_{50}$, 7.1 min) than PCV13X ($CT_{50}$, >30 min), which is mirrored by

more effective hepatic trapping and cellular capture of the pathogen in PCV13-immunized

mice (**Figs. 4B** and **4F**, **Movie 7**). To a less extent, PCV13 also exhibited a similar pattern of

functional advantage over the other brand in the three aspects of pneumococcal clearance of

serotype 6A, with more efficient levels of blood pathogen clearance ($CT_{50}$ of 0.8 min for

PCV13 vs. 2.0 min for PCV13X), hepatic trapping (86% for PCV13 vs. 88% for PCV13X) and

total cellular capture in the liver sinusoids (74 for PCV13 vs. 47 for PCV13X) (**Figs. 4C** and

**4F**; **Movie 8**).

Potential brand-specific vaccine effects for serotype-4 and -5 pneumococci were evaluated

with an infection dose of $10^8$ CFU because this was the most revealing dose for the serotype-

specific vaccine effects of these two types (**Fig. 1**). While infection with serotype-4

pneumococci revealed no substantial differences between the two vaccines in the three

parameters (**Figs. 4D** and **4F**, **Movie 9**), PCV13 showed the most dramatic advantage over the

other brand in the vaccine effects against serotype-5 pneumococci. PCV13-immunized mice

displayed a highly efficient blood clearance ($CT_{50}$ of 0.7 min for PCV13 vs. >30 min for

PCV13X), hepatic trapping (97% for PCV13 vs. 15% for PCV13X) and total cellular capture

(87 for PCV13 vs. 36 for PCV13X) in the liver sinusoids (**Figs. 4E** and **4F**, **Movie 10**). Together,

these data show that the three *in vivo* immunoprotection correlates are capable of identifying

quantitative differences in vaccine-induced pathogen clearance functions.

We finally verified the PCV13 brand-specific variations in CPS-specific IgG antibodies

and protection against lethal infections. ELISA test revealed that PCV13 generally induced

more CPS-specific antibodies than PCV13X across the five serotypes tested thus far (**Fig. 5A**).

In a similar manner, PCV13 immunization led to a consistently more potent protection against

septic infections with all the five HV serotypes. With the doses that were fully protective by

PCV13, PCV13X immunization generated partial protection for serotype-1 (40%), -4 (60%),

and -5 (40%) pneumococci (**Figs. 5B, 5D** and **5E**). For the infection of both serotype-3 and -

6A pneumococci, PCV13 immunization conferred 80% protection of mice while PCV13X

immunization only provided 20% protection (**Figs. 5C** and **5F**). Overall, the superior protection

of PCV13 against septic infection of the HV pneumococcal serotypes than PCV13X agrees

with the quantitative differences of vaccine-activated pathogen clearance in the three novel *in*

*vivo* immunoprotection correlates. In summary, the novel *in vivo* vaccine evaluation system

with three immunoprotection correlates can accurately reveal the potency levels of

pneumococcal vaccines.

**The *in vivo* immunoprotection correlates reveal limited efficacy of whole cell vaccines**

We finally determined whether our PCV13-based *in vivo* vaccine evaluation system could

characterize the functional features of other vaccines beyond *S. pneumoniae.* This potential

was first tested with an inactivated whole cell vaccine (WCV) for a hypervirulent *Klebsiella*

*pneumoniae* (capsular serotype K2), a Gram-negative bacterium and a major pathogen of blood

infections in humans[27]. Our initial trial showed that the WCV confers a full protection against

lethal infection of the same *K. pneumoniae* strain (**Fig. 6A**). While all of naïve mice succumbed

to infection in the first 36 hr post i.p inoculation with $10^6$ CFU of bacteria, the group of

immunized mice completely survived the same challenge. Further analysis revealed that the vaccinated mice gained the ability to clear *K. pneumoniae* from the blood circulation in the first 30 min post i.v. inoculation, with a $CT_{50}$ of 0.6 min, whereas the pathogen were mostly circulating in the bloodstream of naïve mice (**Fig. 6B**). Quantitation of bacteria in the liver and other major organs uncovered 92% of detectable *K. pneumoniae* in the livers of vaccinated mice as compared with 28% hepatic bacteria in naïve mice at 30 min post i.v. infection (**Fig. 6C**). The vaccine-activated clearance of blood-borne *K. pneumoniae* was also manifested by virtual sterilization of the bloodstream in vaccinated mice at 30 min as compared with severe bacteremia in naïve mice (43% of total bacteria in blood). IVM imaging revealed significantly enhanced capture of circulating bacteria by KCs but not LSECs in the sinusoids of vaccinated mice (**Fig. 6D** and **6I**), which is consistent with relatively low protection capacity of the WCV (full mortality at higher infection dose of $10^7$ CFU, **Fig. 6A**). We further confirmed the reliability of the *in vivo* immunoprotection correlates with inactivated WCV of a virulent *Escherichia coli* K1 strain, a capsular serotype commonly causing human blood infection and neonate meningitis[27]. The i.p. infection with a lethal dose of the same *E. coli* strain led to 100% mortality in naïve mice but a complete protection in the vaccinated mice (**Fig. 6E**). In line with the immunoprotection capacity, WCV immunized mice also showed rapid clearance of the pathogen from the circulation (**Fig. 6F**), effective pathogen capture in the liver (**Fig. 6G**) by KCs (**Fig. 6H** and **6J**), as evidenced by dominant proportion of the detectable bacteria in the liver (81%) and virtual blood sterilization of each immunized mouse. It should be mentioned that the KCs of *E. coli* WCV-immunized mice were much more capable of pathogen capture (94 bacteria per FOV) than the counterparts of *K. pneumoniae*-WCV (13 bacteria per FOV), suggesting that the former is more effective. This notion is supported by more significant pathogen killing in the *E. coli* WCV-immunized mice, as indicated by more significant reduction in the total bacteria in *E. coli* WCV-immunized mice (53% of the total bacteria) than the *K. pneumoniae* counterparts (104% of the total bacteria). Lastly, IVM imaging did not revealed any significant enhancement of pathogen capture by LSECs by either *K. pneumoniae* WCV (**Fig. 6I**) or *E. coli* WCV (**Fig. 6J**), which is in sharp contrast to the dominant role of LSECs in PCV13-immunized mice (**Fig. 3**). Based on the extreme importance of LSECs in upgrading vaccine protection potency[20], the failure of WCV in activating pathogen capture of LSECs indicated the limited efficacy of the whole cell vaccines as compared with the licensed PCV13. In conclusion, these data have confirmed the reliability of our *in vivo* immunoprotection correlates in quantifying the functional features and potency of vaccine-activated immunity.

## DISCCUSSION

Immunoprotection efficacy is the core function of all vaccines, regardless the natures of pathogenesis and immunization. However, the current vaccine evaluation system almost entirely depends on *in vitro* features of vaccine-induced responses beyond impact on host disease severity and/or survival, such as antibody titer and antibody-mediated neutralization and/or opsonophagocytosis[22,28,29]. The dynamics of vaccine-induced *in vivo* actions toward pathogen elimination in the process of infection have remained as a "black hole". In this study, we have used four different vaccines (two PCV13s and two WCVs) for three pathogens (*S. pneumoniae*, *K. pneumoniae* and *E. coli*) to characterize an *in vivo* vaccine evaluation system, based on our recent discoveries of the liver as the major immune organ to capture and kill highly virulent bacteria from the blood circulation in vaccinated mice[19,20]. This system includes three related vaccine-activated functions in the defense against invasive infection: pathogen clearance from the bloodstream, trapping in the liver, and cellular capture by KCs and/or LSECs. These correlates have thus bridged our knowledge gap between vaccine-induced antibody response and immunoprotection. Despite that many effective vaccines have been developed by conventional approaches, many important infectious diseases remain unpreventable by vaccination. These functional features of vaccine immunity will not only upgrade our understanding of vaccine biology, but may also empower future vaccine development, particularly for those invasive pathogens that are able to survive in the blood circulation by circumventing innate capture by Kupffer cells (encapsulated bacteria, fungi, and protozoa) and LSECs (viruses).

We have clearly shown that the kinetics of pathogen decay in the bloodstream of vaccinated mice accurately reflect the immunoprotection level of vaccines (survival rate). The pneumococcal serotypes with faster clearance rates or lower $CT_{50}$ values at high infection doses (serotypes 4, 5 and 6A) in PCV13-immunized mice consistently exhibited stronger immunity to lethal challenge or higher $LD_{50}$ values (**Fig. 1**). Conversely, the serotypes with higher $CT_{50}$ counts also displayed lower $LD_{50}$ values (serotypes 1 and 3). This functional association is further supported by PCV13 brand-specific correlation between blood pathogen clearance and immunoprotection (**Figs. 4** and **5**). Lastly, *E. coli* WCV-immunized mice that displayed faster pathogen clearance in the bloodstream than the *K. pneumoniae* counterparts also showed more potent immunoprotection (**Fig. 6**). Vaccine-activated removal of invading bacteria from bloodstream has been commonly used as a parameter for vaccine evaluation[20,30,31]. However, to the best of our knowledge, this is the first report of demonstrating quantitative relationship between vaccine-activated blood pathogen clearance and immunoprotection. An advantage of

this parameter is its suitability in repetitive sampling of the blood by retroorbital bleeding without sacrificing the animals. Along with the dynamic detection of blood bacteria at multiple time points post i.v. infection, typically within 30 min, both the clearance proportion and rate of circulating bacteria can be measured (**Fig. 1**), which makes it more sensitive and quantitative.

Our data have shown that the vaccine-activated pathogen trapping in the liver is well suited as a functional correlate for immunoprotection potency of vaccines against blood-borne bacteria. Serotypes 4, 5 and 6A with the superior immunoprotection or high $LD_{50}$ values in PCV13-immunzed mice displayed higher percentages of total bacteria in the liver than serotypes 1 and 3, which exhibited the inferior immunoprotection and hepatic pathogen trapping (**Fig. 2**). This finding was verified by functional differences of two PCV13 brands in immunoprotection and hepatic trapping of five HV pneumococcal serotypes (**Figs. 4** and **5**). Moreover, we have used two WCVs to confirm that vaccine-activated hepatic pathogen trapping can be used to evaluate the immunoprotection potency of vaccines beyond the polysaccharide conjugate vaccines. *E. coli* WCV was superior to the *K. pneumoniae* WCV in both hepatic pathogen trapping and immunoprotection efficacy (**Fig. 6**). The liver has been recognized as a major organ to clear blood-borne microbes for over six decades[32-37]. Our recent studies have uncovered the essential role of the liver as the organ for the execution of vaccine-activated immunity against blood infections of virulent bacteria[19,20], which has changed our conventional view that vaccine-activated immunity eliminates blood-borne bacteria by circulating phagocytes[6,38].

Consistent with the abovementioned causal relationship between hepatic pathogen trapping and immunoprotection efficacy, this study has demonstrated that cellular pathogen capture by KCs and LSECs in the liver sinusoids is a reliable and sensitive indicator of vaccine immunoprotection. This relies on the state-of-art non-invasive intravital imaging technology to visualize the process of pathogen capture by KCs and LSECs in the liver sinusoidal vasculatures. This technique can provide multiple facets of functional information on vaccine immunity: overall pathogen capture and ratio of KC- and LSEC-captured bacteria. As manifested by significant difference of two WCVs in FOV value (**Fig. 6**), the vaccine activating stronger pathogen capture by KCs confers more potent immunoprotection. However, as indicated in our recent work[20], this study shows that LSEC-mediated pathogen capture is an accurate correlate of the elite vaccine immunity. LSEC-dominant pathogen capture in the liver accompanies the potent immunoprotection in PCV13-immunized mice. It is worth mentioning that the levels of LSEC-captured bacteria distinguish the subtle difference of PCV13 to PCV13X in immunoprotection against serotype-4 pneumococci (**Fig. 4D**), which could not be

differentiated by either blood pathogen clearance or hepatic pathogen trapping. In sharp

contrast, the mice immunized with two WCVs showed only pathogen capture by KCs and

relatively weaker immunoprotection efficacies. Along this line, effective activation of LSEC's

pathogen capture will be the direction for the development of potent vaccines for *E. coli*, *K.*

*pneumoniae* and other common pathogens of blood infections in humans.

This study has uncovered important insights into the spatial and technical venues for future

improvement of polysaccharide conjugate vaccines. While previous epidemiological surveys

have reported relatively poor protection of pneumococcal conjugate vaccines against serotype-

3 pneumococci in humans[24-26,39], the mechanisms behind the functional differences between

serotype 3 and other types have remained undefined. Our data in mouse blood infection model

have shown that, despite comparable levels of PCV13-induced serotype-specific IgG

antibodies for the five serotypes, PCV13-induced immunoprotection against serotype-3 is

strikingly inferior to other serotypes (e.g., $LD_{50}$ values). Accordingly, our *in vivo* vaccine

evaluation system reveals that PCV13 induces much poorer levels of blood pathogen clearance,

hepatic pathogen trapping and cellular pathogen capture in the liver. Because vaccine-induced

IgG antibodies, particularly IgG1 subtype, define the pathogen binding from KCs to LSECs in

the liver[20], our data indicate that PCV13 induces a much lower level of protective antibodies

to the serotype-3 capsule, which cannot be identified by measuring the total antibodies.

It is highly intriguing to observe substantial differences between two PCV13 brands in

various immunoprotective correlates. While there may be differences in manufactural

procedures, such as the purity of capsular polysaccharides and polysaccharide-protein

conjugation, it is noticeable that the two brands use different number of carrier proteins. While

PCV13X uses a single carrier protein for all the 13 serotypes, PCV13 contains the same carrier

protein as PCV13X and an additional carrier, in which 7 serotypes are conjugated to one carrier

while the other 6 serotypes are linked to the other carrier. The carrier protein is essential for

PCV to activate helper T cell to generate high-affinity antibodies and long-term memory

compared with polysaccharide vaccines[7]. Since both the vaccines share one carrier protein, our

data strongly suggest that the functional superiority of PCV13 is rooted in the extra carrier.

These immunological differences may be caused by excessive occupation of immunological

capacity by a single carrier protein in PCV13X, a phenomenon called carrier protein-induced

epitope suppression (CIES)[40]. The competition for helper T cells to recognize and present the

same carrier protein conjugated with different haptens (polysaccharides in PCV13) is the

molecular basis for CIES[41,42]. In other words, PCV13 may benefit from potential attenuation

of the CIES effect by using dividing the labor between the two carriers.

In summary, our *in vivo* vaccine evaluation system can not only reliably and sensitively

distinguish the functional features of vaccines, but also provide functional guidance to future

vaccine design and evaluation. The three assays of this new vaccine evaluation system can be

directly applicable to the evaluation and development of new vaccines for bacteria, fungi and

parasites that cause systemic infections, particularly the encapsulated bacteria that are

refractory to phagocytic clearance in the liver (e.g., *E. coli* and *K. pneumoniae*). With certain

technical modifications, our findings may also be used for evaluation of vaccine candidates for

viral pathogens that depend on viremia for dissemination and vector-borne transmission.

 **METHOD AND MATERIAL**

**Bacteria strains and cultivation**

All the strains of *S. pneumoniae*, *E. coli* and *K. pneumoniae* used in this study are listed in **Table S1**. Pneumococcal strains were cultured in Todd-Hewitt broth (THB) (Oxoid) with 0.5% yeast extract or on tryptic soy agar (TSA) (BD Difco) plates with 5% defibrinated sheep blood at 37℃ with 5% $CO_2$ as described[21]. *E. coli* and *K. pneumoniae* were cultured in Luria-Bertani (LB) broth at 37℃ with shaking as described[19].

**Vaccines and immunoprotection**

PCV13 and PCV13X used in this study have been approved by the Chinese National Medical Products Administration (NMPA). The two vaccines contain the same set of 13 capsular polysaccharide antigens or serotypes (1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F), with an apparent difference in the carrier protein based on the information at the manufacturer's websites. All the 13 CPSs are linked to a single carrier protein in PCV13X; in PCV13, 7 serotypes are conjugated to the common carrier with PCV13X, whereas the other 6 serotypes to the second carrier.

C57BL/6J mice and CD1 mice were obtained from Beijing HFK Bioscience (Beijing, China) and Vital River (Beijing, China), respectively, and kept in SPF facilities at Laboratory Animal Resources Center of Tsinghua University. All animal experiments were conducted according to animal protocols approved by the Institutional Animal Care and Use Committee at Tsinghua University. Immunization with PCV13 was carried out in 5-week-old C57BL/6J mice as previously described[20]. 50 μl (1/10 dose of human) of PCV13 or adjuvant (aluminum phosphate) was intramuscularly (i.m.) injected. Two boost immunizations were carried out by the same procedure 14 and 28 days later. Immunization with the whole cell vaccines of *E. coli* and *K. pneumoniae* was carried out in 5-week-old CD1 mice as previously described[43,44]. Briefly, $10^8$ colony forming unit (CFU) of *E. coli* strain TH14512 (K1) or *K. pneumoniae* strain TH13044 (K2) were treated with 0.5% formalin in phosphate-buffered saline (PBS) at 4°C overnight, washed with PBS by centrifugation, resuspended in 475 μl PBS, and mixed with 25 μl of Adju-Phos® adjuvant (InvivoGen) before subcutaneous (s.c.) immunization. The same procedure was repeated once 7 days later. Immune sera were collected by retro-orbital bleeding 2 weeks after final immunization.

Immunoprotection was assessed by intraperitoneal (i.p.) infection with various doses of bacteria in 100 μl PBS 3 weeks post final immunization as described[20]. The animal survival

was observed every 12 hrs within 7 days post infection. The 50% lethal dose ($LD_{50}$) was calculated by $LD_{50}$ calculator (Quest GraphTM, AAT Bioquest, Inc).

**Bacterial clearance from the bloodstream**

The clearance of *S. pneumoniae*, *E. coli* and *K. pneumoniae* from the bloodstream was evaluated by assessing kinetics of bacteremia retro-orbital bleeding and CFU plating on TSA blood plates at various time points in the first 30 min post intravenous (i.v.) inoculation of bacteria in 100 μl PBS as described previously[20]. Clearance time for half of the inoculum ($CT_{50}$) values were obtained as described[21].

**Bacterial trapping in the liver**

Bacteria trapped in the liver and other major organs (heart, kidney, lung and spleen) were determined as described[20]. Briefly, bacterial burdens in the blood, heart, kidney, liver, lung and spleen of each mouse were measured by CFU plating at 30 min post infection and presented as CFU count for each organ. Bacteria trapped in the liver are presented as CFU count or percentage of bacteria detected in the liver out of the total viable bacteria detected in the blood and five major organs.

**Cellular capture of bacteria by liver immune cells**

Bacterial capture by KCs and LSECs was assessed by real-time intravital microscopy (IVM) imaging of mouse liver with inverted confocal laser scanning microscopy as described[45]. The LSECs and KCs were labelled by i.v. injection of 2.5 μg AF594 anti-CD31 (BioLegend) and AF647 anti-F4/80 (BioLegend) antibodies, respectively, 30 min before infection. $5 \times 10^7$ CFU of FITC-labeled bacteria were i.v. inoculated to track the immobile bacteria captured in the liver sinusoids. FITC labeling was accomplished by incubating $10^8$ CFU of bacteria with 100 μl PBS containing 1 μg FITC (Sigma-Aldrich) for 30 min at room temperature in dark. Images were acquired with Leica TCS SP8 confocal microscope using 10×/0.45 NA and 20×/0.80 NA HC PL APO objectives. The microscope was equipped with Acousto Optics without filters. Fluorescence signals were detected by photomultiplier tubes and hybrid photo detectors ($600 \times 600$ pixels for time-lapse series and $1,024 \times 1,024$ pixels for photographs). Three laser excitation wavelengths (488, 585, and 635 nm) were employed by white light laser (1.5 mw, Laser kit WLL2, 470–670 nm). Real-time imaging was monitored for at least 1 min post infection. More than five random fields of view (FOV) at 10 min post infection were selected to quantify the bacteria captured by LSECs and KCs.

**Enzyme-linked immunosorbent assay (ELISA)**

CPS-specific IgG was quantified by enzyme-linked immunosorbent assay (ELISA) as described[20,46]. Briefly, 96-well plates were coated with capsular polysaccharides and then blocked by 5% non-fat dry milk (BD Difco). The mouse serum was serially diluted with PBS and added into CPS-coated wells for two-hour incubation at 37°C. Bound antibodies were detected by incubation with goat anti-mouse HRP-conjugated IgG (1:2,000, EasyBio, Beijing, China) for 1 h. The optical absorbance at wavelength of 450 nm was measured after 5-minute incubation of TMB substrates (TIANGEN, China) and stopped by the addition of 1 M phosphoric acid.

## REFERENCES

1. Cecconi M, Evans L, Levy M, Rhodes A. Sepsis and septic shock. *Lancet* 2018; **392**(10141): 75-87.

2. Rudd KE, Johnson SC, Agesa KM, et al. Global, regional, and national sepsis incidence and mortality, 1990-2017: analysis for the Global Burden of Disease Study. *Lancet* 2020; **395**(10219): 200-11.

3. Munford RS, Suffredini AF. Sepsis, severe sepsis and septic shock. In: Bennett JE, Dolin RD, Blaser MJ, eds. Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases. 8 ed. Philadelphia: Elsevier Health Sciences; 2014: 914-34.

4. Mayr FB, Yende S, Angus DC. Epidemiology of severe sepsis. *Virulence* 2014; **5**(1): 4-11.

5. Bogaert D, de Groot R, Hermans PWM. *Streptococcus pneumoniae* colonisation: the key to pneumococcal disease. *Lancet Infect Dis* 2004; **4**(3): 144-54.

6. Briles DE, Paton JC, Mukerji R, Swiatlo E, Crain MJ. Pneumococcal vaccines. *Microbiol Spectr* 2019; **7(6)**: GPP3-0028-2018.

7. Rappuoli R, De Gregorio E. A sweet T cell response. *Nat Med* 2011; **17**(12): 1551-2.

8. Rappuoli R. Glycoconjugate vaccines: Principles and mechanisms. *Sci Transl Med* 2018; **10**(456).

9. Wahl B, O'Brien KL, Greenbaum A, et al. Burden of *Streptococcus pneumoniae* and *Haemophilus influenzae* type b disease in children in the era of conjugate vaccines: global, regional, and national estimates for 2000-15. *Lancet Glob Health* 2018; **6**(7): e744-e57.

10. Two new pneumococcal vaccines-Prevnar 20 and Vaxneuvance. *JAMA* 2021; **326**(24): 2521-2.

11. De Gregorio E, D'Oro U, Bertholet S, Rappuoli R. Vaccines. In: Paul WE, ed. Fundamental Immunology. Philadelphia: Wolters Kluwer; 2012: 1032-68.

12. Fleck RA, Romero-Steiner S, Nahm MH. Use of HL-60 cell line to measure opsonic capacity of pneumococcal antibodies. *Clin Diagn Lab Immunol* 2005; **12**(1): 19-27.

13. Jackson LA, Gurtman A, Rice K, et al. Immunogenicity and safety of a 13-valent pneumococcal conjugate vaccine in adults 70 years of age and older previously vaccinated with 23-valent pneumococcal polysaccharide vaccine. *Vaccine* 2013; **31**(35): 3585-93.

14. Kubes P, Jenne C. Immune responses in the liver. *Annu Rev Immunol* 2018; **36**: 247-77.

15. Racanelli V, Rehermann B. The liver as an immunological organ. *Hepatology* 2006; **43**(2 Suppl 1): S54-62.

16. Bilzer M, Roggel F, Gerbes AL. Role of Kupffer cells in host defense and liver disease. *Liver Int* 2006; **26**(10): 1175-86.

17. Krenkel O, Tacke F. Liver macrophages in tissue homeostasis and disease. *Nat Rev Immunol* 2017; **17**(5): 306-21.

18. Sorensen KK, Simon-Santamaria J, McCuskey RS, Smedsrod B. Liver sinusoidal endothelial cells. *Compr Physiol* 2015; **5**(4): 1751-74.

19. Huang X, Li X, An H, et al. Capsule type defines the capability of *Klebsiella pneumoniae* in evading Kupffer cell capture in the liver. *PLoS Pathog* 2022; **18**(8): e1010693.

20. Wang J, An H, Ding M, et al. Potent bacterial vaccines require FcγRIIB-mediated pathogen capture by liver sinusoidal endothelium. *bioRxiv* 2022; doi: 2022.07.19.500551.

21. An H, Qian C, Huang Y, et al. Functional vulnerability of liver macrophages to capsules defines virulence of blood-borne bacteria. *J Exp Med* 2022; **219**(4): e20212032.

22. LaFon DC, Nahm MH. Measuring immune responses to pneumococcal vaccines. *J Immunol Methods* 2018; **461**: 37-43.

23. Recommendations to assure the quality, safety and efficacy of pneumococcal conjugate vaccines, Annex 3, TRS No 977. 2013.

24. Slotved HC, Dalby T, Harboe ZB, et al. The incidence of invasive pneumococcal serotype 3 disease in the Danish population is not reduced by PCV-13 vaccination. *Heliyon* 2016; **2**(11):

e00198.

25. Lapidot R, Shea KM, Yildirim I, Cabral HJ, Pelton SI, Department Of Public Health TM.
Characteristics of serotype 3 invasive pneumococcal disease before and after universal
childhood immunization with PCV13 in Massachusetts. *Pathogens* 2020; **9**(5).

26. Menzies R, Stein AN, Booy R, Van Buynder PG, Litt J, Cripps AW. The impact of the
changing pneumococcal national immunisation program among older Australians. *Vaccine*
2021; **39**(4): 720-8.

27. Rhee C, Wang R, Zhang Z, et al. Epidemiology of hospital-onset versus community-onset
sepsis in U.S. hospitals and association with mortality: a retrospective analysis using electronic
clinical data. *Crit Care Med* 2019; **47**(9): 1169-76.

28. Paschall AV, Middleton DR, Avci FY. Opsonophagocytic killing assay to assess
immunological responses against bacterial pathogens. *J Vis Exp* 2019; (146).

29. Plotkin SA. Correlates of protection induced by vaccination. *Clin Vaccine Immunol* 2010;
**17**(7): 1055-65.

30. Brown EJ, Hosea SW, Frank MM. The role of antibody and complement in the
reticuloendothelial clearance of pneumococci from the bloodstream. *Rev Infect Dis* 1983; **5
Suppl 4**: S797-805.

31. Li Y, Hill A, Beitelshees M, et al. Directed vaccination against pneumococcal disease. *Proc
Natl Acad Sci U S A* 2016; **113**(25): 6898-903.

32. Martin SP, Kerby GP, Holland BC. A method for measuring removal of bacteria from the
blood by the various organs of the intact animal. *Proc Soc Exp Biol Med* 1949; **72**(1): 63-8.

33. Rogers DE. Studies on bacteriemia. I. Mechanisms relating to the persistence of
bacteriemia in rabbits following the intravenous injection of *Staphylococci*. *J Exp Med* 1956;
**103**(6): 713-42.

34. Benacerraf B, Sebestyen MM, Schlossman S. A quantitative study of the kinetics of blood
clearance of P$^{32}$-labelled *Escherichia coli* and *Staphylococci* by the reticuloendothelial system.
*J Exp Med* 1959; **110**(1): 27-48.

35. Mackaness GB. Cellular resistance to infection. *J Exp Med* 1962; **116**: 381-406.

36. Brown EJ, Hosea SW, Frank MM. The role of complement in the localization of
pneumococci in the splanchnic reticuloendothelial system during experimental bacteremia. *J
Immunol* 1981; **126**(6): 2230-5.

37. Gregory SH, Sagnimeni AJ, Wing EJ. Bacteria in the bloodstream are trapped in the liver
and killed by immigrating neutrophils. *J Immunol* 1996; **157**(6): 2514-20.

38. Nahm MH, Katz J. Immunity to extracellular bacteria. In: E.P. W, ed. Fundamental
Immunology. 6th ed. Philadephia: Lippincott-Raven Publishers; 2012: 1001-15.

39. Andrews NJ, Waight PA, Burbidge P, et al. Serotype-specific effectiveness and correlates
of protection for the 13-valent pneumococcal conjugate vaccine: a postlicensure indirect cohort
study. *Lancet Infect Dis* 2014; **14**(9): 839-46.

40. Pobre K, Tashani M, Ridda I, Rashid H, Wong M, Booy R. Carrier priming or suppression:
understanding carrier priming enhancement of anti-polysaccharide antibody response to
conjugate vaccines. *Vaccine* 2014; **32**(13): 1423-30.

41. Herzenberg LA, Tokuhisa T. Epitope-specific regulation. I. Carrier-specific induction of
suppression for IgG anti-hapten antibody responses. *J Exp Med* 1982; **155**(6): 1730-40.

42. Dagan R, Goldblatt D, Maleckar JR, Yaich M, Eskola J. Reduction of antibody response to
an 11-valent pneumococcal vaccine coadministered with a vaccine containing acellular
pertussis components. *Infect Immun* 2004; **72**(9): 5383-91.

43. Serushago BA, Mitsuyama M, Handa T, Koga T, Nomoto K. Role of antibodies against
outer-membrane proteins in murine resistance to infection with encapsulated *Klebsiella
pneumoniae*. *J Gen Microbiol* 1989; **135**(8): 2259-68.

44. Gohar A, Abdeltawab NF, Fahmy A, Amin MA. Development of safe, effective and

immunogenic vaccine candidate for diarrheagenic *Escherichia coli* main pathotypes in a mouse
model. *BMC Res Notes* 2016; **9**: 80.
45. Wang J, Kubes P. A reservoir of mature cavity macrophages that can rapidly invade visceral
organs to affect tissue repair. *Cell* 2016; **165**(3): 668-78.
46. Wang Y, Wen Z, Pan X, Briles DE, He Y, Zhang JR. Novel immunoprotective proteins of
*Streptococcus pneumoniae* identified by opsonophagocytosis killing screen. *Infect Immun* 2018;
**86**(9) e00423-18.
604
605

**Acknowledgments:** We thank Tsinghua research platforms for assistance in animal
experimentation (Laboratory Animal Research Center) and IVM imaging (Center for
Cell Biology).

**Funding:** This work was supported by grants from National Natural Science Foundation of
China 31820103001, 31530082, 81671972, 31728002 (J.-R.Z.) and 3210010245 (J.W.);
China Postdoctoral Science Foundation 2020M680518 (J.W.); Tsinghua-Peking Joint
Center for Life Sciences Postdoctoral Foundation (J.W.).

**Author contributions:**

Conceptualization: X.T., J.W., J.-R.Z.;

Experimentation: X.T., J.W., H.C., M.D., Q.J., J.-R.Z.;

Methodology: X.T., J.W., H.C., M.D., Q.J.;

Investigation: J X.T., J.W., J.-R.Z.;

Funding: J.W., J.-R.Z.;

Project administration: J.W., J.-R.Z.;

Supervision: X.T., J.W., J.-R.Z.;

Writing original draft: X.T., J.W., J.-R.Z.;

**Competing interests:**

Patent applications have been filed on the basis of this work.

**Data and materials availability:**

All data are available in the main text or the supplementary materials.
626

# FIGURES AND LEGENDS

**Figure 1. Vaccine-activated pathogen clearance from the circulation in PCV13-imminzed**

**mice**.

**(A)** CPS-specific IgG titers of serotype 1, 3, 4, 5, 6A in sera of adjuvant-treated (control) and

PCV13-immunized mice as detected by ELISA. n = 3-5.

**(B)** Bacteremia levels of adjuvant-treated mice i.v. infected with $10^6$ CFU of serotype-1, -3, -

4, -5, -6A pneumococci in the first 30 min. n = 3.

**(C-E)** Bacteremia level of PCV13-immunized mice i.v. infected with $10^6$ **(C)**, $10^7$ **(D)** or $10^8$

**(E)** CFU of serotype-1, -3, -4, -5, -6A pneumococci in the first 30 min. n = 3.

**(F-J)** Survival rate of adjuvant-treated and PCV13-immunized mice i.p. infection with various

doses of serotype-1 **(F)**, -3 **(G)**, -4 **(H)**, -5 **(I)** or -6A **(J)** pneumococci. $LD_{50}$ is presented at the

top. n = 5.

Data presented as the mean values ± standard error of mean (SEM). Dotted line in (B-E)

indicates the limit of detection.

**Figure 2. Vaccine-activated hepatic trapping of blood-borne pneumococci in PCV13-immunized mice.**

**(A)** Proportional bacterial burden in the blood and each of the five major organs of adjuvant-treated mice at 30 min post i.v. infection with $10^6$ CFU of serotype-1, -3, -4, -5 and -6A pneumococci. Relative proportion of hepatic bacteria is presented in or at the top of each bar as a percentage of total bacteria detected in each mouse. n = 3.

**(B-D)** Proportional bacterial burden in the blood and five major organs of PCV13-immunized mice at 30 min post i.v. infection with $10^6$ **(B)**, $10^7$ **(C)** or $10^8$ **(D)** CFU of serotype-1, -3, -4, -5 and -6A pneumococci. Relative proportion of hepatic bacteria is presented in each bar as in **(A)**.

Data presented as the mean values ± standard error of mean (SEM).

**Figure 3. Vaccine-activated pathogen capture by LSECs and KCs in PCV13-immunized mice.**

**(A)** Representative IVM images of pneumococci (green), Kupffer cells (KCs, red) and liver sinusoid endothelial cells (LSECs, cyan) in the liver sinusoids of adjuvant-treated (control, left panel) or PCV13-immunized (right panel) mice 10 min after i.v. infection with $5 \times 10^7$ CFU of serotype-1, -3, -4, -5 and -6A pneumococci. n = 2. The process of bacterial capture is demonstrated in Movies 1-5.

**(B)** Quantitation of bacteria immobilized on LSECs and KCs of adjuvant-treated (control) or PCV13-immunized mice in at least five random fields of view (FOV).

Data presented as the mean values ± standard error of mean (SEM).

**Figure 4. Brand-specific functional variations of PCV13 revealed by the *in vivo***

**immunoprotection correlates.**

**(A-E)** Bacteremia in the first 30 min (left panel), bacterial burden in the blood and major organs

at 30 min (middle panel), and bacterial capture by LSECs and KCs (right panel) in PCV13- or

PCV13X-immunized mice post i.v. infection with serotype-1 (**A**), -3 (**B**), -6A (**C**), -4 (**D**) and

-5 (**E**) pneumococci. Relative proportion of hepatic bacteria is indicated in or above each bar

as in Fig. 2A. The data for the first two panels for each serotype were derived from the same

experiment with the following infection dose: $10^7$ CFU for serotypes 1, 3 and 6A; $10^8$ CFU for

serotypes 4 and 5. Bacterial capture (n = 2) was quantitated by IVM imaging as in Fig. 3B.

**(F)** Representative IVM images of the liver sinusoids in PCV13- (left panel) or PCV13X-

immunized (right panel) mice 10 min after i.v. infection with $5 \times 10^7$ CFU of serotype-1, -3, -

4, -5 and -6A pneumococci as in Fig. 3A. The process of bacterial capture is demonstrated in

Movie 6 (serotype 1), Movie 7 (serotype 3), Movie 8 (serotype 6A), Movie 9 (serotype 4), and

Movie 10 (serotype 5).

**Figure 5. Antibody titer and immunoprotection of efficacy in PCV13- and PCV13X-immunized mice.**

**(A)** CPS-specific IgG titers of serotype 1, 3, 4, 5 and 6A in sera of PCV13- or PCV13X-immunized mice. n = 3-5.

**(B-F)** The survival of PCV13- or PCV13X-immunized mice post i.p. infection with $2 \times 10^7$ CFU serotype-1 (**B**), $1 \times 10^6$ CFU serotype-3 (**C**), $5 \times 10^7$ CFU serotype-4 (**D**), $1 \times 10^8$ CFU serotype-5 (**E**) or $5 \times 10^7$ CFU serotype-6A (**F**) pneumococci. n = 5.

Data presented as the mean values ± standard error of mean (SEM).

**Figure 6. Vaccine-activated pathogen clearance, trapping in the liver and capture by Kupffer cells in whole cell vaccine (WCV)-immunized mice.**

**(A-D)** Functional features of *K. pneumoniae* WCV. Host survival (**A**), bacteremia in the first 30 min (**B**), bacterial burden in blood and major organs at 30 min (**C**), and bacterial capture by LSECs and KCs (**D**) post infection of *K. pneumoniae* strain TH13044 (serotype K2) in naïve or WCV-immunized mice. Infections were performed with the following doses and routes: $10^6$ or $10^7$ CFU (i.p.) for (A), $5 \times 10^6$ CFU (i.v.) for (B-C), $5 \times 10^7$ CFU (i.v.) for (D). n = 5 for (A), 3 for (B-C), and 2 for (D).

**(E-H)** Functional features of *K. pneumoniae* WCV. Host survival (**E**), bacteremia (**F**), bacterial burden in blood and major organs (**G**), and bacterial capture (**H**) post infection of *E. coli* strain TH14512 (serotype K1) in naïve or WCV-immunized mice. Infections were performed with

the following doses and routes: $5 \times 10^7$ CFU (i.p.) for (E), $10^8$ CFU (i.v.) for (F-G), $5 \times 10^7$

CFU (i.v.) for (H). n = 5 for (E), 3 for (F-G), and 2 for (H).

**(I-J)** Representative IVM images of the liver sinusoids in naive (left panel) or WCV-

immunized (right panel) mice 10 min after i.v. infection with $5 \times 10^7$ CFU of *K. pneumoniae*

**(I)** and *E. coli* **(J)** as in Fig. 3A. The process of bacterial capture is demonstrated in Movies 11-

12.

Data presented as the mean values ± standard error of mean (SEM).

     **SUPLMENATRY MATERIALS**

712     **Table S1. Bacterial strains used in this study**

| Strain ID | Species | Serotype | Source | Region | Reference |
|-----------|---------|----------|--------|--------|-----------|
| TH2866 | *S. pneumoniae* | 1 | Blood | China | [21] |
| TH865 | *S. pneumoniae* | 3 | Cerebrospinal Fluid | USA | [21] |
| TIGR4 | *S. pneumoniae* | 4 | Blood | Norway | [21] |
| TH2940 | *S. pneumoniae* | 5 | Blood | China | [21] |
| TH870 | *S. pneumoniae* | 6A | Blood | USA | [21] |
| TH13044 | *K. pneumoniae* | K2 | Blood | China | [19] |
| TH14512 | *E. coli* | K1 | Blood | China | [21] |

**Movie 1-5. PCV13-elicited pneumococcal capture of high-virulence serotypes in liver**

**sinusoid.** Capture of i.v. inoculated serotype-1 (Movie 1), -3 (Movie 2), -4 (Movie 3), -5

(Movie 4) and -6A (Movie 5) pneumococci ($5 \times 10^7$ CFU, FITC-labelled, green) from the

bloodstream by liver sinusoid endothelial cells (LSECs, cyan) and resident macrophage

Kupffer cells (KCs, red) in adjuvant-treated (control) or PCV13-immunized mice. Real-time

imaging was visualized for at least 1 minute by intravital microscopy immediately post

infection. The playback rate is eight frames per second. The quantitative data were shown in

Fig. 3B; selected images in Fig. 3A.

**Movie 6-10. More efficient pneumococcal capture in liver sinusoids elicited by PCV13**

**than PCV13X.** $5 \times 10^7$ CFU of FITC-labelled serotype-1 (Movie 6), -3 (Movie 7), -6A (Movie

8), -4 (Movie 9) and -5 (Movie 10) pneumococci in PCV13-or PCV13X-immunized mice were

visualized as in Movie 1. The quantitative data were shown in Figs. 4A-E; selected images in

Fig. 4F.

**Movie 11-12. Whole cell vaccine induced capture of *K. pneumoniae* and *E. coli* by KCs.** 5

$\times 10^7$ CFU of FITC-labelled *K. pneumoniae* strain TH13044 (Movie 11) and *E. coli* strain

TH14512 (Movie 12) in naive or WCV-immunized mice were visualized as in Movie 1. The

quantitative data were shown in Fig. 6D and 6H; selected images in Figs. 6I-J.