The *Vibrio* Type III Secretion System 2 is not restricted to the *Vibrionaceae* and

encodes differentially distributed repertoires of effector proteins

Jerez SA[1, ‡], Plaza N [2, ‡], Bravo V[3, ‡], Urrutia I[2, ‡], Blondel CJ[1]*

[1] Instituto de Ciencias Biomédicas, Facultad de Medicina y Facultad de Ciencias de la

Vida, Universidad Andrés Bello, Santiago, Chile.

[2] Instituto de Ciencias Biomédicas, Facultad de Ciencias de la Salud, Universidad

Autónoma de Chile, Santiago, Chile.

[3] Programa Centro de Investigación Biomédica y Aplicada, (CIBAP), Escuela de Medicina,

Facultad de Ciencias Médicas, Universidad de Santiago de Chile, Santiago, Chile.

‡ These authors contributed equally to this work.

*Corresponding author: carlos.blondel@unab.cl

Mailing address, Instituto de Ciencias Biomédicas, Universidad Andrés Bello. Echaurren

183, Santiago, 8320000, Chile.

ABSTRACT

Vibrio parahaemolyticus is the leading cause of seafood-borne gastroenteritis worldwide. A distinctive feature of the O3:K6 pandemic clone, and its derivatives, is the presence of a second, phylogenetically distinct, Type III Secretion System (T3SS2) encoded within the genomic island VPaI-7. The T3SS2 allows the delivery of effector proteins directly into the cytosol of infected eukaryotic cells to subvert key host cell processes, critical for *V. parahaemolyticus* to colonize and cause disease. Furthermore, the T3SS2 also increases the environmental fitness of *V. parahaemolyticus* in its interaction with bacterivorous protists; hence it has been proposed that it contributed to the global oceanic spread of the pandemic clone. Several reports have identified T3SS2-related genes in *Vibrio* and non-*Vibrio* species, suggesting that the T3SS2 gene cluster is not restricted to the *Vibrionaceae* and can mobilize through horizontal gene transfer events. In this work, we performed a large-scale genomic analysis to determine the phylogenetic distribution of the T3SS2 gene cluster and its repertoire of effector proteins. We identified putative T3SS2 gene clusters in 1130 bacterial genomes from eight bacterial genera and five bacterial families. A hierarchical clustering analysis allowed us to define six T3SS2 subgroups (I-VI) with different repertoires of T3SS2 effector proteins, redefining the concepts of T3SS2 core and accessory effector proteins. Finally, we identified a subset of T3SS2 gene clusters that lack most T3SS2 effector proteins described to date and provided a list of 10 novel effector candidates for this subgroup through bioinformatic analysis. Collectively, our findings indicate that the T3SS2 extends beyond the *Vibrionaceae* family and suggest that different effector protein repertories could have a differential impact on the pathogenic potential and environmental fitness of all the bacteria that have acquired the *Vibrio* T3SS2 gene cluster.

DATA SUMMARY

1. All genome sequences used in this study were downloaded from the National Center for Biotechnology Information (NCBI) RefSeq or GenBank databases (See **Table S1** for accession numbers).

2. Files for the T3SS2 reconstructed phylogenetic tree (Newick tree and MSA fasta file), hierarchical clustering data analysis file from MORPHEUS, **Table S1** with genome accession numbers and all the data of the absence/presence of T3SS2-related components, and **Table S2** with the prediction of novel effector proteins are available as part of the online Supporting Dataset at the Zenodo data repository (https://doi.org/10.5281/zenodo.7016552). The T3SS2 phylogenetic tree can be interactively visualized in https://itol.embl.de/tree/190161901253747116626959067#

INTRODUCTION

Vibrio parahaemolyticus is a Gram-negative bacterium that resides in marine and estuarine environments [1, 2]. In 1996, a pandemic clone of the O3:K6 serotype emerged and spread around the globe, becoming the leading cause of seafood-borne gastroenteritis worldwide [3]. Whole-genome sequencing of the pandemic strain RIMD2210633 revealed the presence of a novel Type III Secretion System (T3SS2) encoded within an 80-kb genomic island on chromosome 2, known as the *V. parahaemolyticus* pathogenicity island 7 (VPaI-7) [4]. While all *V. parahaemolyticus* strains encode a T3SS on chromosome 1 (T3SS1), only *V. parahaemolyticus* strains derived from the original pandemic clone harbor the phylogenetically distinct T3SS2 [4].

T3SS are complex nanomachines that enable Gram-negative bacteria to deliver proteins, known as effectors, directly from the bacterial cytosol into the cytosol of eukaryotic cells [5, 6]. The translocation of effectors into target cells enables bacteria to subvert a wide variety of host cell functions [7–9]. In addition, the specific repertoire of proteins delivered to infected cells determines which cellular networks are subverted by the bacterium and ultimately influence the outcome of the infection [5, 6, 10]. In this context, even closely related T3SSs can have different repertoires of effector proteins due to the due to horizontal gene transfer (HGT) and/or genomic rearrangement events [11, 12]. A recent global survey described that T3SS are widely distributed among Gram-negative bacteria, identifying 174 nonredundant T3SS from 109 distinct bacterial genera [13]. T3SS have been classified in 13 (I-XIII) major categories based on sequence divergence of T3SS core components [13]. These T3SS have been classified in 13 (I-XIII) major categories based on sequence divergence of T3SS core components [13]. According to this study, the *Vibrio* T3SS1 is part of category Ib, and the T3SS2 is the founding member of the T3SS phylogenetic category X [13].

Currently, the T3SS2 is considered the main virulence factor of *V.*

parahaemolyticus. Studies in animal models of infection revealed that, while the T3SS1

plays a minor role during infection, the T3SS2 is essential for the enterotoxicity and the

development of the clinical signs of the disease [14–17]. Therefore, the T3SS2 is

considered the major virulence factor of pandemic *V. parahaemolyticus* [18]. The gene

expression of the T3SS2 gene cluster is induced by bile salts [19, 20] through a regulatory

network involving the VtrA, VtrC, and VtrB proteins [19, 21, 22] and the T3SS2 function

requires host cell-surface fucosylation for efficient delivery of effector protein into infected

cells [23]. Notably, the importance of the T3SS2 goes beyond increasing the pathogenic

potential of pandemic strains, as the T3SS2 increases the environmental fitness of *V.*

parahaemolyticus by promoting bacterial survival against predatory amoeba [24].

Phylogenetic analysis based on the amino acid sequence divergence of T3SS

structural components has identified two distinct lineages of the T3SS2, known as

phylotypes T3SS2α and T3SS2β [25, 26]. T3SS2α and T3SS2β gene clusters share a

similar genetic organization and a subset of effector proteins (**Fig. 1**) [18]. A more recent

study has proposed an additional category within T3SS2β, known as T3SS2γ [27]. The

T3SS2γ is very similar to the T3SS2β gene cluster but has some gene content differences,

such as the presence of a *tdh* gene inserted within the T3SS gene cluster [27] (**Fig. 1**).

Most of our knowledge of the T3SS2 comes from studies of the T3SS2α of *V.*

parahaemolyticus strain RIMD2210633 and T3SS2α *V. cholerae* strain AM-19226

(reviewed in [18, 28]). In this context, 15 different T3SS2 effector proteins have been

identified, of which only 5 are shared between *V. parahaemolyticus* RIMD2210633 and *V.*

cholerae AM-19226 [29–42]. These 5 effectors are therefore called core effector proteins,

while the remaining 10 are known as accessory effector proteins [18]. While useful, this

classification has the limitation of comparing the distribution of effector of only 2 *Vibrio*

strains of the same T3SS2 phylotype.

The T3SS2 gene cluster is not restricted to *V. parahaemolyticus* as similar T3SSs

have been found in other species such as non-O1/non-O139 *V. cholerae*, *V. mimicus* and

V. anguillarum [25, 43–46]. Furthermore, several reports have identified the presence of

one or more T3SS2-related genes in strains of bacterial genera other than *Vibrio*, including

Aeromonas [47], *Grimontia* [20], *Photorhabdus* [48, 49], *Providencia* [50] and *Shewanella*

[49, 51]. Altogether, these reports suggest that the presence of T3SS2 gene clusters

extends beyond the *Vibrionaceae* family and could have been acquired through HGT

events. Nevertheless, these correspond to isolated observations performed in a limited

number of strains and bacterial genomes. Therefore, the overall taxonomic and

phylogenetic distribution of the T3SS2 and its effector proteins in bacterial genomes is

currently unknown.

To address this issue, in this study, we performed a large-scale bioinformatic

analysis to gain insight into the phylogenetic distribution of the T3SS2 gene clusters,

including effector proteins. Our analysis identified 1030 bacterial genomes encoding

putative T3SS2 gene clusters distributed over 43 different bacterial species from 8

bacterial genera. In addition, we classified these T3SS2s into 6 different subgroups in

terms of the repertoire of effector proteins they possess. Our work provides evidence that

the T3SS2 extends beyond the *Vibrionaceae* family and that acquisition of both the T3SS2

and different repertoires of effector proteins could have an impact on the pathogenic

potential and environmental fitness of these bacteria.

Impact Statement

One of the distinctive features of the O3:K6 pandemic clone of *Vibrio parahaemolyticus* is the presence of a phylogenetically distinct Type III Secretion System (T3SS) known as T3SS2. T3SS are complex nanomachines that allow the delivery of effector proteins directly into the cytosol of eukaryotic cells to manipulate different cellular functions. The *Vibrio* T3SS2 is the major virulence factor of the pandemic clone and a pivotal contributor to the pathogenesis and environmental fitness of the bacterium. Several studies have identified the T3SS2 gene cluster in other bacterial species beyond the *Vibrionaceae* family, suggesting that this T3SS can be mobilized through horizontal gene transfer events. In this work, we performed a phylogenetic distribution analysis of the T3SS2 gene cluster in publicly available bacterial genome sequences. We identified putative T3SS2 gene clusters in 1130 bacterial genomes from 43 different bacterial species from 8 genera and 5 bacterial families. Furthermore, we classified these T3SS2s into 6 different subgroups (I-VI) in terms of their repertoire of effector proteins. Our findings indicate that the T3SS2 is not restricted to the *Vibrionaceae* family and suggest that different effector protein repertories could have a differential impact on the pathogenic potential and environmental fitness of all the identified bacteria that have acquired this T3SS2.

    METHODS

In silico identification and analyzes of T3SS2 loci and components

To identify putative T3SS2 loci, the amino acid sequence of the VPA1338 (SctN)

and VPA1339 (SctC) proteins of *V. parahaemolyticus* RIMD 2210633 were used as

queries in BLASTp, to perform analyses [52] using publicly available bacterial genome

sequences of the National Center for Biotechnology Information (NCBI) database

(February 2021). A cut-off of 70% identity and 80% sequence coverage threshold were

used to select for positive matches. All genome sequence assemblies with positive hits

were downloaded from the NCBI RefSeq or GenBank databases (See **Table S1** for

accession numbers and metadata) and used to build local BLAST databases. To identify

the remaining 37 T3SS2-related components in these 1130 genomes, the nucleotide and

aminoacid sequence of each component (**Table S2**) was used as queries for either

tBLASTx or BLASTp analyzes, respectively. A 50% identity and 80% sequence coverage

threshold were used to select for positive matches. For comparative genomic analysis,

nucleotide sequences were visualized and analyzed by the Artemis version 18.1 [53], the

Artemis Comparison Tool (ACT) release 6 [54], the multiple aligner Mauve [55] and

EasyFig v2.2.2 [56]. For the hierarchical clustering analysis, a presence/absence matrix of

each T3SS2 component was constructed for each bacterial genome (**Table S1**) and

uploaded as a *csv* file to the online server of the Versatile matrix visualization and analysis

software MORPHEUS (one minus Pearson correlation, average linkage method)

(https://software.broadinstitute.org/morpheus). The hierarchical clustering data analysis file

from MORPHEUS is available as part of the online Supporting Data at the Zenodo data

repository (https://doi.org/10.5281/zenodo.7016552). The presence and absence of each

identified component were visualized as a colored matrix generated in GraphPad Prism

version 9 (GraphPad Software, San Diego, California, USA).

T3SS2 phylogenetic analyses

The 1130 concatenated SctN and SctC amino acid sequences were aligned with

ClustalW using the Molecular Evolutionary Genetics Analysis (MEGA) software version 7.0

[57]. The phylogenetic tree was built from the alignments obtained from MEGA by

performing a bootstrap test of phylogeny (1000 replications) using the Maximum-

Likelihood (ML) method with a Jones-Taylor-Thornton (JTT) + CAT correction model using

FastTree2 (Galaxy Version 2.1.10+galaxy1) [58] in the Galaxy platform

(https://usegalaxy.org/) [59]. The phylogenetic tree was further visualized and edited to

include important metadata with iTOL [60]. Files for the T3SS2 reconstructed phylogenetic

tree (Newick tree) is available as part of the online Supporting Data at the Zenodo data

repository (https://doi.org/10.5281/zenodo.7016552). The T3SS2 phylogenetic tree can be

interactively visualized in https://itol.embl.de/tree/190161901253747116626959067#

Identification of novel T3SS2 effector candidates

Each Open Reading Frame (ORF) within the predicted T3SS2 gene clusters of

Subgroup VI (including 1 kb upstream and downstream of the first and last predicted

component, respectively) was analyzed by means of the EffectiveDB [61], Bastion3 [62]

and BEAN2.0 [63] pipelines to identify potential T3S signal sequences. ORFs with positive

hits in at least one of these pipelines was further analyzed to predict putative functional

domains with the InterProScan [64] and NCBI-CDD [65]. Finally, each ORF was also

analyzed by the we used the structure-based homology tool HHpred [66]. Multiple

sequence alignments were performed by MAFFT [67] and T-Coffee Expresso [68] and

visualized by ESPript 3.0 [69].

RESULTS AND DISCUSSION

T3SS2 gene clusters extend beyond the *Vibrionaceae* family

We screened all publicly available bacterial genomes in the NCBI database to identify the full breadth of bacterial species harboring putative T3SS2 gene clusters. First, we performed BLASTP analysis in search for homologs of the VPA1338 (SctN) and VPA1339 (SctC) proteins of *V. parahaemolyticus* RIMD 2210633. SctN and SctC were chosen as they are two of the most conserved T3SS structural proteins and are often used for phylogenetic clustering and classification of T3SSs [13, 70]. We used stringent cut-offs to avoid identifying homolog proteins from phylogenetically close T3SSs (≥70□% of identity and over ≥80□% of coverage). Our analysis identified the presence of VPA1338 and VPA1339 homologs in 1130 bacterial genomes (**Fig. 2, Table S1**).

To determine if these 1130 bacterial genomes could encode complete T3SS2 gene clusters, we further screened each genome for the presence of the 37 T3SS2-related components previously identified in *V. parahaemolyticus* RIMD 2210633 (T3SS2α), *V. parahaemolyticus* MAVP-R (T3SS2β) and *V. cholerae* AM-19226 T3SS2α (**Table S1**). These include ORFs encoding 16 structural proteins, 15 bacterial effector proteins, 3 regulatory proteins (VtrA, VtrB, and VtrC), and each of the 3 hemolysins linked to T3SS2 gene clusters (TDH$_{Vp}$, TRH$_{Vc}$ and TRH$_{Vp}$). Most of the 1130 bacterial genomes (1106) encoded each of the T3SS structural genes suggesting that they most likely harbor complete T3SS2 gene clusters (**Fig. 3**). The exception was 24 *Vibrio anguilarum* genomes that only possess 8 of the 16 T3SS2 structural components (**Table S1**). This was not unexpected as it has been previously reported that some *V. anguillarum* strains lack most of the T3SS2 structural components due to a large internal deletion of the T3SS2 gene cluster [71], suggesting that these strains do not encode functional T3SS2s.

Taxonomic distribution analysis identified putative T3SS2 gene clusters in 43

bacterial species from 8 bacterial genera (*Vibrio*, *Photobacterium*, *Grimontia*,

Photorhabdus, *Yersinia*, *Shewanella,* and *Aeromonas*) and 5 distinct families

(*Aeromonadaceae*, *Morganellaea*, *Shewanellaceae*, *Vibrionaceae,* and *Yersiniaceae*) (**Fig.**

2A). In some genus, we identified multiple species harboring putative T3SS2 gene clusters

while in other genus we could identify as little as one bacterial species harboring T3SS2

gene components (**Fig. 2A**). In terms of bacterial species, the genus *Vibrio* contained the

largest number of different species harboring putative T3SS2 gene clusters (29 species),

followed by the *Photorhabdus*, *Shewanella* and *Providencia* (each with 3 species), the

Yersinia genus (2 species) and the *Photobacterium*, *Grimontia,* and *Aeromonas* each with

a single species. Finally, our analysis did not identify bacterial strains harboring multiple

T3SS2 gene clusters within their genomes.

Interestingly, most of the different bacterial species identified to harbor putative

T3SS2 gene clusters can be found in aquatic habitats, suggesting that acquisition and

transfer of the T3SS2 actively occur in this environment. Among the exceptions was the

identification of T3SS2 gene clusters in genomes of strains of the *Photorhabdus* genus

(**Fig. 2A** and **Table S1**). Bacteria from the genus *Photorhabdus* comprise different

bacterial species that are important insect pathogens, including some that can also cause

human disease [72]. Wilkinson et al.[48], previously identified a T3SS2 gene cluster in the

genome of *Photorhabdus asymbiotica* strain ATCC43949. Our analysis expanded this

finding by identifying complete T3SS2 gene clusters in different strains of *P. australis*, *P.*

heterorhabitis and *P. asymbiotica* species (**Fig. S2** and **Table S1**). Every T3SS2 gene

cluster identified is inserted in the exact genetic location compared to the *P. laumondii*

strain TT01 genome, as previously reported for *P. asymbiotica* strain ATCC43949 [48]. We

also identified T3SS2 gene clusters in 1 strain of *Yersinia kristensenii* and 2 strains of

Yersinia enterocolitica (**Fig. 2A** and **Table S1**). Interestingly, while *Yersinia enterocolitica*

can be found in aquatic environments, the 2 strains with putative T3SS2 gene clusters

were isolated from wild ungulate carcasses and belong to the highly virulent biotype 1B

[73].

Altogether, the taxonomic distribution of T3SS2 gene clusters in bacterial genomes

indicates that this secretion system is not restricted to the *Vibrionaceae* family and

supports the notion that T3SS2 gene clusters can mobilize through horizontal gene

transfer events among environmental bacteria.

The T3SS2β phylotype is most abundant and divergent than the phylotype T3SS2α

The T3SS2 is divided into two major phylotypes known as T3SS2α and T3SS2β

[18], and an additional subcategory within T3SS2β known as T3SS2γ [27]. To determine

the distribution of T3SS2α, T3SS2β and T3SS2γ, among the 1130 T3SS2s identified, we

performed a phylogenetic analysis using concatenated SctN and SctC aminoacidic

sequences. We also included the concatenated SctN and SctC sequences of the

Agrobacterium radiobacter K84 and *Rhizobium etli* CIAT 652 T3SSs, which are

representatives of T3SS category XI, the closest phylogenetic category to the *Vibrio*

T3SS2 [13]. The phylogenetic clustering analysis showed that all the 1030 putative

T3SS2s grouped closely together while the T3SS of *Agrobacterium radiobacter* K84 and

Rhizobium etli CIAT 652 cluster in a more distant branch (**Fig. 2B**), supporting the notion

that all the 1130 T3SS identified in this work correspond to bona fide T3SS2s. Our

analysis also revealed that 36,6% (n= 414) of the identified T3SS2s belonged to phylotype

T3SS2α and 61% (n= 689) to phylotype T3SS2β. The phylogenetic analysis also revealed

that the greatest sequence divergence was observed within the T3SS2β phylotype (**Fig.**

2C).

As mentioned above, recent reports have created a subcategory within T3SS2β of

V. parahaemolyticus known as T3SS2γ (based mainly on the presence of both the *tdh* and

trh genes) [27]. To determine the percentage of T3SS2γ gene clusters within the T3SS2β

family, we performed a gene content analysis based on the definition of T3SS2γ gene

clusters proposed by Xu et al., [13]. The analysis identified that 40,5% (n= 280) of T3SS2β

gene clusters could be classified as T3SS2γ. Notably, the presence of T3SS2γ did not

cluster to any major clade of the T3SS2 phylogenetic tree (**Fig. 2B**), supporting the notion

that T3SS2γ is not a third T3SS2 phylotype but rather a subcategory within T3SS2β.

Taxonomic distribution analysis identified the presence of T3SS2α and T3SS2β in

strains of each of the 5 bacterial families (**Fig. 2A and 2B**), with no clear

overrepresentation of any phylotype within a particular bacterial family. Nevertheless,

T3SS2β was present in a more significant number of different bacterial species (37 out of

44) in comparison to T3SS2α (19 out of 44 species) (**Fig. 2A**). Finally, the T3SS2γ

subcategory was identified solely in *V. parahaemolyticus* strains which agree with previous

reports [27]. Since many *V. parahaemolyticus* strains which harbor T3SS2 gene clusters

are related to the RIMD 2210633 pandemic clone, we analyzed the distribution of T3SS2

phylotypes among different *V. parahaemolyticus* Sequence Types (STs). Our analysis

showed that T3SS2α was overrepresented in *V. parahaemolyticus* strains from ST3 (44%),

ST120 (11,5%), ST189 (7%) and ST332 (16,4%) while the T3SS2β was overrepresented

in strains from ST3 (11,5%), ST36 (25,2%) and ST631 (7,5%) (**Fig. S1**).

T3SS2 gene clusters share a high degree of synteny

To gain insight into the genomic context of the identified T3SS2 gene clusters, we

performed a comparative genomic analysis in a subset of bacterial strains with complete or

almost complete bacterial genomes (large enough contigs to allow a comparative

analysis). Notably, there was a high gene cluster synteny between the T3SS2 gene of *V.*

parahaemolyticus RIMD2210633 and the gene clusters of bacterial species of different

genera (e.g., *Providencia, Photobacterium, Grimontia* and *Photorhabdus*) (**Fig. 3 and Fig.**

S2). A high gene cluster synteny was also observed between the T3SS2 gene clusters

with a lower sequence identity, such as the gene clusters of *Aeromonas* sp. DNP9 and

Yersinia enterocolitica Y1Wildboar1B (**Fig. 3C**).

In terms of genome location, in addition to the described chromosomal location of

T3SS2 within VPAI-7 of *V. parahamolyticus* RIMD2210633 [4], we identified a T3SS2 gene

cluster within the plasmid of a recently sequenced *V. meditteraniae* strain 117-T6

(CP033579) [74] (**Fig. 4A**). Another interesting observation was the presence of a T3SS2

gene cluster inserted right next to a T3SS1 gene cluster in *V. campbelli* strain UMTGB204.

In this strain both T3SSs are inserted within the vicinity of the same serine tRNA (**Fig. 4B**),

in which the T3SS1 of *V. parahaemolyticus* RIMD 2210633 is located. Finally, in each of

the T3SS2 gene clusters identified in *Grimontia hollisae* strains, a large gene encoding a

putative adhesin (SiiE-like adhesin) is encoded in the middle of the T3SS2 gene cluster

(**Fig. 4A**).

Hierarchical clustering analysis identifies 6 subgroups with different repertoires of

effector proteins

While phylogenetic analysis of T3SSs using core structural components provides

insight into the evolution of T3SSs, the acquisition, transfer, or loss of accessory and

effector encoding genes provides an additional layer of complexity to the evolution of these

systems [11, 13, 70]. To gain further insight into this diversity, we performed a hierarchical

clustering analysis of 37 T3SS2-related (**Fig. 5**). For the analysis of the effector proteins,

we included each T3SS2 effector described for either *V. parahaemolyticus* RIMD 2210633

or *V. cholerae* AM 19226 [18]. The hierarchical clustering analysis allowed us to identify 6

major subgroups (I to VI) with similar gene content profiles (**Fig 5**, **Table S1**).

Most structural components were highly conserved in all subgroups, except for the

hydrophilic translocator VopW of subgroups I and VI and the VopB2 and VopD2 translocon

proteins of subgroup VI. The *vtrA*, *vtrB,* and *vtrC* genes, which encode the regulatory

complex that induces the expression of T3SS2 gene clusters in response to bile [19, 22],

were also highly conserved, except for subgroups III and VI, where many bacterial strains

harbored divergent homologs of VtrA and VtrC (**Fig. 5**).

Since most structural genes were highly conserved across all genomes, the

hierarchical clustering analysis mainly classified each T3SS2 regarding the repertoire of

effector proteins identified. As shown in **Fig. 6A**, each T3SS2 subgroup was enriched in a

distinct set of effector proteins and associated hemolysins. While most subgroups were

enriched for a combination of 10-12 different T3SS2 effectors (subgroups I to V), subgroup

VI was vastly underrepresented in terms of the presence of known effector proteins. This

subgroup was primarily enriched in the VgpA effector protein, lacking most other known

T3SS2 effector proteins. The presence of VgpA homologs in each subgroup supports this

effector´s critical role in T3SS2 function [39, 75]. As expected, the current phylotype

classification (T3SS2α versus T3SS2β) was a poor predictor of the presence/absence of

individual effector proteins (**Fig. 6A and B**), highlighting the utility of a gene-content

classification scheme.

Interestingly, most T3SS2 subgroups were enriched for genomes of particular

bacterial species (**Fig. 6B**): i) Subgroup I included strains from 4 different bacterial species

(*P. damselae*, *V. cholerae*, *V. mimicus,* and *Vibrio sp.*); ii) Subgroup II included strains only

from *Grimontia hollisae* species; iii) Subgroup III included *V. parahaemolyticus* strains

exclusively; iv) Subgroup IV included strains from 3 species (*V. cholerae*, *V. mimicus* and

V. parahaemolyticus); v) Subgroup V included strains from 5 species (*V.*

parahaemolyticus, *V. cholerae*, *V. diabolicus*, *V. mimicus* and *Vibrio sp.*); and subgroup VI

harbored the most diverse set of bacterial species with a total of 39 bacterial species (**Fig.**

6B). Subgroup VI was also the only subgroup with bacterial strains beyond the

Vibrionaceae family. Finally, while no T3SS2 subgroup was exclusive to a particular

T3SS2 phylotype some subgroups were indeed overrepresented by different phylotypes.

Subgroups I, III and IV were overrepresented in strains from the T3SS2α phylotype, while

subgroups II, V and VI were overrepresented in strains from the T3SS2β phylotype.

Finally, every *V. parahaemolyticus* strain of the T3SS2γ subcategory clustered within

subgroup V (**Table S1**).

Altogether, the hierarchical clustering analysis suggests that T3SS2s have

acquired a diverse set of effector proteins which could ultimately impact on the contribution

of each T3SS2 to the pathogenic potential and environmental fitness of different bacterial

strains.

T3SS2 subgroup VI encodes 10 novel effector candidates

As mentioned above, the T3SS2 gene clusters of subgroup VI stand out as they

are the most taxonomically diverse (**Fig. 6B**) and lack most of the known T3SS2 effector

proteins (**Fig. 5, 6A**). The high degree of synteny of these gene clusters to the T3SS2

gene cluster of *V. parahaemolyticus* RIMD2210633 (**Fig. 3 and Fig. S2**) and their

distribution in the T3SS2 phylogenetic tree (**Fig 2**) suggests that these are not part of a

phylogenetically distinct T3SS2, but rather have a different set of effector proteins. To test

this hypothesis, we extracted the T3SS2 gene cluster from selected bacterial genomes of

Subgroup VI (strains with closed genomes or sufficiently large contigs to include complete

T3SS2 gene clusters) and performed bioinformatic analysis to identify putative T3SS2

effector candidates. We analyzed each ORF of these T3SS2 clusters based on 4 criteria,

including: i) identification of putative T3SS secretion signals by T3SS effector prediction

pipelines (BEAN2.0, Bastion3 and EffectiveDB); ii) bioinformatic analysis to identify

conserved functional domains (InterProScan, NCBI-CDD); iii) sequence identity to known

T3SS effector proteins of other bacterial species (BLAST); and iv) functional predictions

via remote homology searches using the HHpred pipeline.

382   Our analysis identified a set of 10 novel effector candidates that harbored putative

T3SS secretion signal sequences and putative functional domains present in known T3SS

effector proteins. These included putative ADP-ribosyltransferases, tyrosine

phosphatases, glycosyltransferases and Rho-activating proteins (**Table 1 and Fig. 7A**). In

addition, 5 of these ORFs encode homologs of known T3SS effector proteins or toxins

such as the effector proteins OspC1, AoPH and VopC of *S. flexneri*, *A. hydrophila* and *V.*

parahaemolyticus respectively (**Table 1 and Fig. 7A**).

389   Interestingly, one candidate harbored a predicted GDSL lipase domain

(IPR001087) and a 22,4% sequence identity to the VPA0226 protein of *V.*

parahaemolyticus (**Fig. S4**). VPA0226 is a secreted lipase required for *V.*

parahaemolyticus RIMD2210633 to egress from the infected cell. Notably, in *V.*

parahaemolyticus VPA0226 is not secreted by the T3SS2 but rather by the T2SS [76]. The

10 effector candidates identified were present in diverse bacterial species of the T3SS2

subgroup VI, except for the putative tyrosine phosphatases (DYE18_RS08865) that also

was identified in *Grimontia hollisae* strains of T3SS2 subgroup II (**Fig. 7B and Fig. 7C**).

397   Overall, our analysis suggests that subgroup VI encodes a unique set of novel

398   T3SS2 effector proteins not present in any other T3SS2 subgroup. Further work is needed

399   to confirm that each candidate corresponds to T3SS2 effector proteins.

401   **Revisiting the current classification of core and accessory T3SS2 effector proteins**

402   Recently, Hu et al.[13], proposed a classification of T3SS2 effector proteins based

on the presence of these effectors in *V. parahaemolyticus* and *V. cholerae.* Effectors

present in both *V. cholerae* and *V. parahaemolyticus* are known as core effectors, and

those present in either *V. cholerae* or *V. parahaemolyticus* have been called accessory

effector proteins. While useful, this classification relies on analyzing only two strains of two

bacterial species.

To revisit this classification, we analyzed the presence/absence of each T3SS2

effector protein in the 43 bacterial species in which we have identified putative T3SS2

gene clusters. As shown in **Fig. S3**, only VgpA was present in at least one strain of most

bacterial species (88%). The high prevalence of this effector protein could be explained by

its additional structural role [39]. Another effector, VopT, was identified exclusively in *V.*

parahaemolyticus strains. The remaining 13 T3SS2 effector proteins were present at

various degrees with no overrepresentation in any bacterial species. Altogether, our data

show a wide distribution of T3SS2 effector proteins in different bacterial species and that

there is no clear correlation between the number of bacterial species that harbor a

particular effector protein, which is the basis of the current classification scheme. In this

context, we propose that a better alternative is to classify T3SS2s based on the repertoire

of their effector proteins in the context of all identified T3SS2 loci in bacterial genomes

(subgroups I-VI). One of the limitations of this classification scheme is that it is dynamic, as

it may change over time if novel effector encoding genes are acquired through HGT or if

some are lost due to genomic rearrangements.

CONCLUSION

The acquisition and transfer of T3SS gene clusters can potentially increase the virulence and the environmental fitness of the recipient bacteria. In this context, it has been proposed that the acquisition of the T3SS2 gene cluster by *V. parahaemolyticus* was a crucial factor in the global spread and increased pathogenic potential of the O3:K6 pandemic clone and its derivatives. In this study, we performed a genome-wide analysis to determine the phylogenetic distribution of the *Vibrio* T3SS2 gene cluster in public genome databases. Our analysis revealed that T3SS2 gene clusters extend beyond the *Vibrionaceae* family, share a high degree of synteny and that it is possible to categorize T3SS2 in six different subgroups (I-VI) in terms of their repertoire of effector proteins. Notably, it is plausible to think that acquisition of the T3SS2 gene cluster by HGT events in each of the bacteria identified in this study could increase their human pathogenic potential and environmental fitness, just as it has been proposed for the *V. parahaemolyticus* pandemic clone. Finally, since the repertoire of T3SS effector proteins delivered to the target cells ultimately defines the cellular outcome of an infection, we believe that future studies should focus on testing the contribution of each repertoire of T3SS2 effector proteins to the pathogenic potential and environmental fitness of these bacterial species.

CONFLICT OF INTERESTS

The authors declare that there are no conflicts of interest.

FUNDING INFORMATION

This work was funded by the Howard Hughes Medical Institute (HHMI)-Gulbenkian

International Research Scholar Grant #55008749 and FONDECYT Grant 1201805 (ANID)

from CJB. SJ is supported by the national PhD grant 21210879 (ANID). IMU is supported

by FONDECYT Grant 3200874 (ANID). VB is supported by DICYT grant 022001BZ

(USACH).

AUTHOR CONTRIBUTIONS

Sebastian Jerez: Investigation, Visualization, review and editing final manuscript.

Nicolas Plaza: Investigation, Visualization, review and editing final manuscript.

Italo M Urrutia: Investigation, Visualization, review and editing final manuscript.

Veronica Bravo: Investigation, Visualization, review and editing final manuscript.

Carlos J Blondel: Conceptualization, Methodology, Formal analysis, Investigation,

Visualization, Resources, Data curation, Writing-Original Draft and review and editing.,

Funding acquisition, Supervision and Project administration.

ACKNOWLEDGMENTS

We thank members of the Blondel Laboratory for helpful discussions on all aspects of this

project and for their comments on the manuscript.

 FIGURE LEGENDS

Fig. 1. The T3SS2 is divided into two major phylotypes T3SS2α and T3SS2β, and a

subcategory within T3SS2β known as T3SS2γ. Schematic depiction of a comparison of

the T3SS2 gene clusters in *V. cholerae* AM-19226, *V. parahaemolyticus* RIMD2210633, *V.*

parahaemolyticus MAVP-R, and MAVP-Q. BLASTn alignment was performed and

visualized using Easyfig. Genes encoding different components are highlighted in color.

Fig. 2. The T3SS2 extends beyond the *Vibrionaceae* family. (A) Identification and

taxonomic distribution of T3SS2 gene clusters in bacterial genomes. The presence of

T3SS2 gene clusters from phylotypes T3SS2α (red) and T3SS2β (blue) are shown for

each taxonomic category. The numbers correspond to the total numbers of bacterial

genomes within each taxonomic category that harbored a putative T3SS2 gene cluster. (B)

Maximum-likelihood phylogenetic tree built from the alignments of concatenated SctN and

SctC amino acid sequences with a bootstrap test of phylogeny (1000 replications) with a

Jones-Taylor-Thornton (JTT) correction model and visualized by iTOL. The presence of

hemolysis and subgroups are shown as colored strips. T3SS2α and T3SS2β are shown in

red and blue branches, respectively. (C) Same tree as B but with branch length

information.

Fig. 3. T3SS2 gene clusters share a high degree of synteny. Schematic depiction of a

comparison of the T3SS2 gene clusters in *V. parahaemolyticus* RIMD2210633 in

comparison to the gene clusters of strains of *Providencia* (A), *Photobacterium* (B) and

Aeromonas, and *Yersinia enterocolitica* strains (C). BLASTn alignments were performed

and visualized using Easyfig. Genes encoding T3SS-related genes are highlighted in blue.

Fig. 4. Distinctive genomic locations of T3SS2 gene clusters. (A) Comparison of the

T3SS2 gene cluster in *V. parahaemolyticus* RIMD2210633 with the T3SS2 gene cluster

encoded in *V. mediterranei* 116-T6 plasmid and the T3SS2 gene cluster of *Grimontia*

hollisae NCTC11640. (B) Schematic depiction of a comparison of the T3SS1 and T3SS2

gene clusters inserted in the vicinity of the serine tRNA in *Vibrio campbellii* strain

UMTGB204. BLASTn alignments were performed and visualized using Easyfig. Genes

encoding T3SS-related genes are highlighted in blue.

Fig. 5. Hierarchical clustering analysis of T3SS2-related genes identifies 6 distinct

subgroups. Clustering analysis was performed using Versatile matrix visualization and

analysis software MORPHEUS (one minus Pearson correlation, average linkage method).

Boxes representing genes not detected, detected, or detected but divergent (less than

70% identity and 80% sequence coverage) are colored in grey, blue or red, respectively.

An asterisk highlights the group of *V. anguillarum* genomes that only harbored a limited

number of T3SS structural components.

Fig. 6. Differential distribution of T3SS2 effector proteins. (A) Frequency of detection

of each T3SS2-related gene in each T3SS2 phylotype and Subgroup. Frequencies are

color-coded and expressed as percentages. (B) Taxonomic distribution of the T3SS2

Subgroups (I-VI) in comparison to the T3SS2 phylotypes.

Fig. 7. T3SS2 Subgroup VI encodes 10 novel effector candidates. (A) Schematic

representation of novel candidate highlighting the presence of predicted functional

domains in different colors. (B) Taxonomic distribution of each predicted effector protein

identified in bacterial genomes. Red presence, white absence. (C) Frequency of the

presence of each predicted novel effector protein in the different T3SS2 subgroups (I-VI).

Frequencies are color-coded and expressed as percentages.

Table S1. Table with information regarding the presence/absence and hierarchical

clustering analysis of T3SS2 components in the bacterial genome sequences (including

accession numbers).

Table S2. Prediction of T3SS effector candidates within the T3SS2 subgroup VI gene

clusters. Representative ORFs for each prediction are shown with the results of different

prediction software, the identification of putative functional domains (InterproScan), and

the amino acid sequence identity to known T3SS effector proteins from other organisms.

Fig. S1. Distribution of *Vibrio parahaemolyticus* Sequence Types (STs) in terms of the

presence of T3SS2α or T3SS2β gene clusters.

Fig. S2. Gene synteny between the T3SS2 gene clusters of *V. parahaemolyticus*

RIMD2210633 and the identified T3SS2 gene clusters of *Photorhabdus* species. BLASTn

alignments were performed and visualized using Easyfig. Genes encoding T3SS-related

genes are highlighted in blue.

Fig. S3. Only VgpA is widely conserved among bacterial species harboring T3SS2 gene

clusters. The distribution matrix shows the detection of each effector protein in at least one

bacterial strain in red. The total number of bacterial species in which each effector was

detected is shown at the bottom of the matrix.

Fig. S4. Multiple sequence alignment of the amino acid sequence of the predicted effector

BCT80_RS20435 and the secreted lipase VPA0226 and the *Salmonella* T3SS effector

protein SseJ. The alignment was performed using T-Coffee Expresso and visualized by

ESPript 3.0. Amino acids within a red background correspond to positions with 100%

identity; amino acids with a yellow background correspond to positions with >70% identity.

REFERENCES

1. Brumfield KD, Usmani M, Chen KM, Gangwar M, Jutla AS, *et al.* Environmental parameters associated with incidence and transmission of pathogenic Vibrio spp. *Environ Microbiol* 2021;23:7314–7340.

2. Letchumanan V, Chan K-G, Lee L-H. Vibrio parahaemolyticus: a review on the pathogenesis, prevalence, and advance molecular identification techniques. *Frontiers in microbiology* 2014;5:705.

3. Velazquez-Roman J, León-Sicairos N, Hernández-Díaz L de J, Canizalez-Roman A. Pandemic Vibrio parahaemolyticus O3:K6 on the American continent. *Frontiers in cellular and infection microbiology* 2013;3:110.

4. Makino K, Oshima K, Kurokawa K, Yokoyama K, Uda T, *et al.* Genome sequence of Vibrio parahaemolyticus: a pathogenic mechanism distinct from that of V cholerae. *Lancet (London, England)* 2003;361:743–749.

5. Galán JE, Lara-Tejero M, Marlovits TC, Wagner S. Bacterial Type III Secretion Systems: Specialized Nanomachines for Protein Delivery into Target Cells. *Annu Rev Microbiol* 2014;68:1–24.

6. Lara-Tejero M, Galán JE. Protein Secretion in Bacteria. *Ecosal Plus* 2019;8:245–259.

7. Portaliou AG, Tsolis KC, Loos MS, Zorzini V, Economou A. Type III Secretion: Building and Operating a Remarkable Nanomachine. *Trends in biochemical sciences* 2016;41:175–189.

8. Dean P. Functional domains and motifs of bacterial type III effector proteins and their roles in infection. *FEMS Microbiology Reviews* 2011;35:1100–1125.

9. Scott NE, Hartland EL. Post-translational Mechanisms of Host Subversion by Bacterial Effectors. *Trends Mol Med* 2017;23:1088–1102.

10. Ruano-Gallego D, Sanchez-Garrido J, Kozik Z, Núñez-Berrueco E, Cepeda-Molero M, *et al.* Type III secretion system effectors form robust and flexible intracellular virulence networks. *Science*;371. Epub ahead of print 2021. DOI: 10.1126/science.abc9531.

11. Brown NF, Finlay B. Potential origins and horizontal transfer of type III secretion systems and effectors. *Mob Genetic Elements* 2011;1:118–121.

12. Stavrinides J, Ma W, Guttman DS. Terminal Reassortment Drives the Quantum Evolution of Type III Effectors in Bacterial Pathogens. *Plos Pathog* 2006;2:e104.

13. Hu Y, Huang H, Cheng X, Shu X, White AP, *et al.* A global survey of bacterial type III secretion systems and their effectors: A global survey of bacterial type III secretion systems. *Environ Microbiol* 2017;19:3879–3895.

14. Yang H, Santos M de S, Lee J, Law HT, Chimalapati S, *et al.* A Novel Mouse Model of Enteric
Vibrio parahaemolyticus Infection Reveals that the Type III Secretion System 2 Effector VopC
Plays a Key Role in Tissue Invasion and Gastroenteritis. *Mbio* 2019;10:e02608-19.

15. Ritchie JM, Rui H, Zhou X, Iida T, Kodoma T, *et al.* Inflammation and disintegration of
intestinal villi in an experimental model for Vibrio parahaemolyticus-induced diarrhea. *PLoS*
pathogens 2012;8:e1002593.

16. Piñeyro P, Zhou X, Orfe LH, Friel PJ, Lahmers K, *et al.* Development of two animal models to
study the function of Vibrio parahaemolyticus type III secretion systems. *Infection and immunity*
2010;78:4551–4559.

17. Hubbard TP, Chao MC, Abel S, Blondel CJ, Wiesch PAZ, *et al.* Genetic analysis of Vibrio
parahaemolyticus intestinal colonization. *Proc National Acad Sci* 2016;113:201601718.

18. Matsuda S, Hiyoshi H, Tandhavanant S, Kodama T. Advances on Vibrio parahaemolyticus
research in the postgenomic era. *Microbiol Immunol* 2020;64:167–181.

19. Gotoh K, Kodama T, Hiyoshi H, Izutsu K, Park K-S, *et al.* Bile acid-induced virulence gene
expression of Vibrio parahaemolyticus reveals a novel therapeutic potential for bile acid
sequestrants. *PloS one* 2010;5:e13365.

20. Li P, Rivera-Cancel G, Kinch LN, Salomon D, Tomchick DR, *et al.* Bile salt receptor complex
activates a pathogenic type III secretion system. *Elife* 2016;5:e15718.

21. Kodama T, Gotoh K, Hiyoshi H, Morita M, Izutsu K, *et al.* Two regulators of Vibrio
parahaemolyticus play important roles in enterotoxicity by controlling the expression of genes in
the Vp-PAI region. *PloS one* 2010;5:e8678.

22. Li P, Rivera-Cancel G, Kinch LN, Salomon D, Tomchick DR, *et al.* Bile salt receptor complex
activates a pathogenic type III secretion system. *Elife* 2016;5:e15718.

23. Blondel CJ, Park JS, Hubbard TP, Pacheco AR, Kuehl CJ, *et al.* CRISPR/Cas9 Screens Reveal
Requirements for Host Cell Sulfation and Fucosylation in Bacterial Type III Secretion System-
Mediated Cytotoxicity. *Cell Host Microbe* 2016;20:226–237.

24. Matz C, Nouri B, McCarter L, Martinez-Urtaza J. Acquired type III secretion system determines
environmental fitness of epidemic Vibrio parahaemolyticus in the interaction with bacterivorous
protists. *Plos One* 2011;6:e20275.

25. Okada N, Iida T, Park K-S, Goto N, Yasunaga T, *et al.* Identification and characterization of a
novel type III secretion system in trh-positive Vibrio parahaemolyticus strain TH3996 reveal
genetic lineage and diversity of pathogenic machinery beyond the species level. *Infection and*
immunity 2009;77:904–913.

26. Okada N, Matsuda S, Matsuyama J, Park K-S, Reyes C de los, *et al.* Presence of genes for type
III secretion system 2 in Vibrio mimicus strains. *BMC microbiology* 2010;10:302.

27. Xu F, Gonzalez-Escalona N, Drees KP, Sebra RP, Cooper VS, *et al.* Parallel Evolution of Two
Clades of an Atlantic-Endemic Pathogenic Lineage of Vibrio parahaemolyticus by Independent
Acquisition of Related Pathogenicity Islands. *Appl Environ Microb* 2017;83:e01168-17.

28. Miller KA, Tomberlin KF, Dziejman M. Vibrio variations on a type three theme. *Curr Opin*
Microbiol 2019;47:66–73.

29. Plaza N, Urrutia IM, Garcia K, Waldor MK, Blondel CJ. Identification of a Family of Vibrio
Type III Secretion System Effectors That Contain a Conserved Serine/Threonine Kinase Domain.
Msphere 2021;6:e00599-21.

30. Zhang L, Krachler AM, Broberg CA, Li Y, Mirzaei H, *et al.* Type III effector VopC mediates
invasion for Vibrio species. *Cell reports* 2012;1:453–460.

31. Kodama T, Rokuda M, Park K-S, Cantarelli VV, Matsuda S, *et al.* Identification and
characterization of VopT, a novel ADP-ribosyltransferase effector protein secreted via the Vibrio
parahaemolyticus type III secretion system 2. *Cellular Microbiology* 2007;9:2598–2609.

32. Hiyoshi H, Okada R, Matsuda S, Gotoh K, Akeda Y, *et al.* Interaction between the Type III
Effector VopO and GEF-H1 Activates the RhoA-ROCK Pathway. *Plos Pathog* 2015;11:e1004694.

33. Zhou X, Gewurz BE, Ritchie JM, Takasaki K, Greenfeld H, *et al.* A Vibrio parahaemolyticus
T3SS effector mediates pathogenesis by independently enabling intestinal colonization and
inhibiting TAK1 activation. *Cell reports* 2013;3:1690–1702.

34. Trosky JE, Mukherjee S, Burdette DL, Roberts M, McCarter L, *et al.* Inhibition of MAPK
Signaling Pathways by VopA from Vibrio parahaemolyticus. *J Biol Chem* 2004;279:51953–51957.

35. Hiyoshi H, Kodama T, Saito K, Gotoh K, Matsuda S, *et al.* VopV, an F-Actin-Binding Type III
Secretion Effector, Is Required for Vibrio parahaemolyticus-Induced Enterotoxicity. *Cell Host*
Microbe 2011;10:401–409.

36. Liverman ADB, Cheng H-C, Trosky JE, Leung DW, Yarbrough ML, *et al.* Arp2/3-independent
assembly of actin by Vibrio type III effector VopL. *Proc National Acad Sci* 2007;104:17117–
17122.

37. Calder T, Kinch LN, Fernandez J, Salomon D, Grishin NV, *et al.* Vibrio Type III Effector
VPA1380 Is Related to the Cysteine Protease Domain of Large Bacterial Toxins. *Plos One*
2014;9:e104387.

38. Hu M, Zhang Y, Gu D, Chen X, Waldor MK, *et al.* Nucleolar c□Myc recruitment by a Vibrio
T3SS effector promotes host cell proliferation and bacterial virulence. *Embo J* 2020;e105699.

39. Tandhavanant S, Matsuda S, Hiyoshi H, Iida T, Kodama T. Vibrio parahaemolyticus Senses
Intracellular K+ To Translocate Type III Secretion System 2 Effectors Effectively. *Mbio*
2018;9:e01366-18.

40. Suzuki M, Danilchanka O, Mekalanos JJ. Vibrio cholerae T3SS Effector VopE Modulates
Mitochondrial Dynamics and Innate Immune Signaling by Targeting Miro GTPases. *Cell Host*
Microbe 2014;16:581–591.

41. Chaand M, Miller KA, Sofia MK, Schlesener C, Weaver JWA, *et al.* Type Three Secretion
System Island-Encoded Proteins Required for Colonization by Non-O1/Non-O139 Serogroup
Vibrio cholerae. *Infect Immun* 2015;83:2862–2869.

42. Alam A, Miller KA, Chaand M, Butler JS, Dziejman M. Identification of Vibrio cholerae Type
III Secretion System Effector Proteins. *Infect Immun* 2011;79:1728–1740.

43. Dziejman M, Serruto D, Tam VC, Sturtevant D, Diraphat P, *et al.* Genomic characterization of
non-O1, non-O139 Vibrio cholerae reveals genes for a type III secretion system. *P Natl Acad Sci*
Usa 2005;102:3465–3470.

44. Okada N, Matsuda S, Matsuyama J, Park K-S, Reyes C de los, *et al.* Presence of genes for type
III secretion system 2 in Vibriomimicus strains. *Bmc Microbiol* 2010;10:302.

45. Naka H, Dias GM, Thompson CC, Dubay C, Thompson FL, *et al.* Complete Genome Sequence
of the Marine Fish Pathogen Vibrio anguillarum Harboring the pJM1 Virulence Plasmid and
Genomic Comparison with Other Virulent Strains of V. anguillarum and V. ordalii. *Infect Immun*
2011;79:2889–2900.

46. Arteaga M, Velasco J, Rodriguez S, Vidal M, Arellano C, *et al.* Genomic characterization of the
non-O1/non-O139 Vibrio cholerae strain that caused a gastroenteritis outbreak in Santiago, Chile,
2018. *Microb Genom* 2020;6:e000340.

47. Vázquez-Rosas-Landa M, Ponce-Soto GY, Eguiarte LE, Souza V. Comparative genomics of
free-living Gammaproteobacteria: pathogenesis-related genes or interaction-related genes? *Pathog*
Dis 2017;75:ftx059.

48. Wilkinson P, Waterfield NR, Crossman L, Corton C, Sanchez-Contreras M, *et al.* Comparative
genomics of the emerging human pathogen Photorhabdus asymbiotica with the insect pathogen
Photorhabdus luminescens. *Bmc Genomics* 2009;10:302.

49. Akeda Y, Kodama T, Saito K, Iida T, Oishi K, *et al.* Identification of the Vibrio
parahaemolyticus type III secretion system 2□associated chaperone VocC for the T3SS2□specific
effector VopC. *Fems Microbiol Lett* 2011;324:156–164.

50. Hu Y, Huang H, Cheng X, Shu X, White AP, *et al.* A global survey of bacterial type III
secretion systems and their effectors. *Environ Microbiol* 2017;19:3879–3895.

51. Plaza N, Urrutia IM, Garcia K, Waldor MK, Blondel CJ. Identification of a Family of Vibrio
Type III Secretion System Effectors That Contain a Conserved Serine/Threonine Kinase Domain.
Msphere 2021;6:e00599-21.

52. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J*
Mol Biol 1990;215:403–410.

53. Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. Artemis: an integrated platform for
visualization and analysis of high-throughput sequence-based experimental data. *Bioinformatics*
2012;28:464–469.

54. Carver TJ, Rutherford KM, Berriman M, Rajandream M-A, Barrell BG, *et al.* ACT: the Artemis
comparison tool. *Bioinformatics* 2005;21:3422–3423.

55. Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: Multiple Alignment of Conserved
Genomic Sequence With Rearrangements. *Genome Res* 2004;14:1394–1403.

56. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. *Bioinformatics*
2011;27:1009–1010.

57. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version
7.0 for Bigger Datasets. *Mol Biol Evol* 2016;33:1870–1874.

58. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees for
Large Alignments. *Plos One* 2010;5:e9490.

59. Afgan E, Baker D, van den Beek M, Blankenberg D, Bouvier D, *et al.* The Galaxy platform for
accessible, reproducible and collaborative biomedical analyses: 2016 update. *Nucleic Acids Res*
2016;44:W3–W10.

60. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic Acids Res 2019;47:gkz239-.

61. Eichinger V, Nussbaumer T, Platzer A, Jehl M-A, Arnold R, *et al.* EffectiveDB—updates and
novel features for a better annotation of bacterial secreted proteins and Type III, IV, VI secretion
systems. *Nucleic Acids Res* 2016;44:D669–D674.

62. Wang J, Li J, Yang B, Xie R, Marquez-Lago TT, *et al.* Bastion3: a two-layer ensemble predictor
of type III secreted effectors. *Bioinformatics* 2018;35:2017–2028.

63. Dong X, Lu X, Zhang Z. BEAN 2.0: an integrated web resource for the identification and
functional analysis of type III secreted effectors. *Database* 2015;2015:bav064.

64. Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, *et al.* The InterPro protein
families and domains database: 20 years on. *Nucleic Acids Res* 2020;49:gkaa977-.

65. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, *et al.* CDD/SPARCLE: the conserved
domain database in 2020. *Nucleic Acids Res* 2019;48:D265–D268.

66. Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, *et al.* A Completely Reimplemented
MPI Bioinformatics Toolkit with a New HHpred Server at its Core. *J Mol Biol* 2018;430:2237–
2243.

67. Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. *Brief Bioinform* 2017;20:bbx108-.

68. Notredame C, Higgins DG, Heringa J. T-Coffee: A Novel Method for Fast and Accurate
Multiple Sequence Alignment. *JMB* 2000;302:205–217.

69. Robert X, Gouet P. Deciphering key features in protein structures with the new ENDscript
server. *Nucleic Acids Res* 2014;42:W320–W324.

70. Abby SS, Rocha EPC. The non-flagellar type III secretion system evolved from the bacterial
flagellum and diversified into host-cell adapted systems. *PLoS genetics* 2012;8:e1002983.

71. Naka H, Dias GM, Thompson CC, Dubay C, Thompson FL, *et al.* Complete genome sequence
of the marine fish pathogen Vibrio anguillarum harboring the pJM1 virulence plasmid and genomic
comparison with other virulent strains of V. anguillarum and V. ordalii. *Infection and immunity*
2011;79:2889–2900.

72. Gerrard JG, McNevin S, Alfredson D, Forgan-Smith R, Fraser N. Photorhabdus Species:
Bioluminescent Bacteria as Human Pathogens? *Emerg Infect Dis* 2003;9:251–254.

73. Macori G, Romano A, Adriano D, Razzuoli E, Bianchi DM, *et al.* Draft Genome Sequences of
Four Yersinia enterocolitica Strains, Isolated from Wild Ungulate Carcasses. *Genome Announc*
2017;5:e00192-17.

74. Liu Q, Xu M, He Y, Tao Z, Yang R, *et al.* Complete Genome Sequence of Vibrio mediterranei
117-T6, a Potentially Pathogenic Bacterium Isolated from the Conchocelis of Pyropia spp.
Microbiol Resour Announc 2019;8:e01569-18.

75. Hu M, Zhang Y, Gu D, Chen X, Waldor MK, *et al.* Nucleolar c☐Myc recruitment by a Vibrio
T3SS effector promotes host cell proliferation and bacterial virulence. *Embo J* 2020;e105699.

76. Chimalapati S, Santos M de S, Lafrance AE, Ray A, Lee W-R, *et al.* Vibrio deploys type 2
secreted lipase to esterify cholesterol with host fatty acids and mediate cell egress. *Elife*
2020;9:e58057.

T3SS2 gene cluster

V. cholerae AM-19226 T3SS2α

V. parahaemolyticus RIMD2210633 T3SS2α

V. parahaemolyticus MAVP-R T3SS2β

V. parahaemolyticus MAVP-Q T3SS2γ

100%

40%

| Structural components | Hypothetical proteins | Urease gene cluster |
| Effector proteins | TDH / TRH | Integrase/Transposase |

A

Family	Genus	Species	T3SS2 Phylotype α	T3SS2 Phylotype β	
Aeromonadaceae	Aeromonas	*Aeromonas sp.*		■	1
Morganallaceae	Photorhabdus	*P. asymbiotica*	■	■	2
		P. australis	■		2
	Providencia	*P. heterorhabditis*	■		3
		P. burhodogranariea		■	1
		P. rettgeri		■	1
		P. sneebia		■	1
Shewanellaceae	Shewanella	*S. baltica*		■	3
		S. psychrophila		■	1
	Grimontia	*Shewanella sp.*		■	7
Photobacterium		*G. hollisae*		■	10
		P. damselae	■		1
Vibrionaceae	Vibrio	*V. aestuarianus*	■		3
		V. agarivorans	■		1
		V. alginolyticus		■	1
		V. anguillarum	■	■	30
		V. aquaticus	■		1
		V. aquimaris		■	1
		V. azureus		■	2
		V. campbellii		■	3
		V. cholerae	■	■	162
		V. coralliilyticus		■	1
		V. corolliirubri	■		1
		V. diabolicus		■	2
		V. harveyi	■	■	4
		V. hyugaensis		■	1
		V. injenensis		■	1
		V. lentus	■	■	33
		V. mediterranei		■	1
		V. metschnikovii		■	1
		V. mimicus	■	■	3
		V. ordalii		■	2
		V. ostreicida	■		2
		V. parahaemolyticus	■	■	810
		V. pectenicida		■	2
		V. qinghaiensis		■	1
		V. sagamiensis		■	1
		Vibrio sp.	■	■	16
		V. splendidus	■	■	3
		V. tasmaniensis		■	4
		V. tetraodonis		■	1
Yersiniaceae	Yersinia	*Y. enterocolitica*		■	2
		Y. kristensenii		■	1
		# Total genomes			1130

B

T3SS2β

T3SS2α

Hemolysins
- ■ TDH
- ■ TRH Vc
- ■ TRH Vp

Subgroups (Hierarchical clustering)
- ■ I
- ■ II
- ■ III
- ■ IV
- ■ V
- ■ VI

T3SSβ subgroup
- ■ T3SS2γ

C

T3SS2α   T3SS2β

Category X

Category XI

Tree scale: 0.1

A

Vibrio mediterranei
117-T6 plasmid

Vibrio parahaemolyticus
RIMD2210633

Grimontia hollisae
NCTC11640

SiiE-like adhesin

100%

21%

B

Vibrio campbellii
DS40M4

Ser-tRNA

Vibrio campbellii
UMTGB204

Ser-tRNA   T3SS2 gene cluster   T3SS1 gene cluster

100%

66%