Lipid metabolism dysfunction following symbiont elimination is linked to altered

Kennedy pathway homeostasis

Geoffrey M. Attardo[1,2†], Joshua B. Benoit[1,3†], Veronika Michalkova[1,4], Alekhya

Kondragunta[3], Aaron A. Baumann[1,5], Brian L. Weiss[1], Anna Malacrida[6], Francesca

Scolari[6,7], and Serap Aksoy[1]

[1]Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New

Haven, Connecticut, United States of America

[2]Department of Entomology and Nematology, Division of Agriculture and Natural

Resources, University of California Davis, Davis, CA 95616

[3]Department of Biological Sciences, University of Cincinnati, Cincinnati, OH, 45221

[4]Section of Molecular and Applied Zoology, Institute of Zoology, Slovak Academy of

Sciences, Bratislava, Slovakia

[5]Center for Agricultural Synthetic Biology, University of Tennessee, Knoxville, TN,

37996

[6]Department of Biology and Biotechnology, University of Pavia, 27100 Pavia, Italy

[7]Institute of Molecular Genetics (IGM), Italian National Research Council (CNR), Via

Abbiategrasso 207, 27100 Pavia, Italy

For correspondence: gmattardo@ucdavis.edu, joshua.benoit@uc.edu

†these authors contributed equally to this work

Abstract

Lipid metabolism is critical for insect reproduction, especially for species that invest

heavily into early developmental stages of their offspring. The role of symbiotic bacteria

during this process is unknown but likely essential, especially in the case of obligate

microbes that fulfill key biological functions in the host. Using a combined lipidomics,

functional genomics and biochemical strategy, we examined the role of lipid metabolism

in the interaction between the viviparous tsetse fly (*Glossina morsitans morsitans*) and

its obligate endosymbiotic bacteria (*Wigglesworthia glossinidia*) during tsetse

pregnancy. We observed increased *CTP:phosphocholine cytidylyltransferase* (*cct1*)

expression during pregnancy. This gene codes for the enzyme that functions as the rate

limiting step in phosphatidylcholine biosynthesis in the Kennedy pathway which is

critical for stored lipid metabolism and progeny development. Experimental removal of

Wigglesworthia impaired lipid metabolism via disruption of the Kennedy pathway,

yielding obese mothers whose developing progeny ultimately starve. Functional

validation via experimental *cct1* suppression revealed a phenotype similar to females

lacking obligate *Wigglesworthia* symbionts. These results indicate that, in *Glossina*,

symbiont-derived factors, likely B vitamins, are critical for proper function of both lipid

biosynthesis and lipolysis. Loss of the symbiosis has a dramatic impact on *Glossina*

fecundity, and may be broadly applicable to other insect systems, particularly to species

that require symbiotic partners to maximize lipolysis and reproductive output.

Keywords: B vitamin, reproduction, *Wigglesworthia*, tsetse fly, phospholipids

Introduction

Reproduction represents an evolutionarily imperative and metabolically taxing

physiological process for all animals (Flatt, 2011; Perrin and Sibly, 1993). This may be

especially true for insects, which maintain a high reproductive rate, even though

progeny quality declines throughout adulthood (Barreaux et al., 2022; Michalkova et al.,

2014a). One such insect is the tsetse fly, *Glossina spp*., which reproduces via a process

called 'adenotrophic viviparity' (gland fed, live birth (Benoit et al., 2015)). Female tsetse

ovulate one egg per gonotrophic cycle and following fertilization and embryogenesis,

larvigenesis proceeds within the maternal uterus. *In utero,* larvae are nourished

exclusively by milk secretions produced by uniquely adapted maternal accessory

glands, called the milk glands. Thus, female tsetse flies supply all the nutrients

necessary to support embryonic and larval development of their progeny. Tsetse milk

consists of 20-30 mg of carbohydrates, lipids and proteins in an aqueous base (Benoit

et al., 2015, 2014; Cmelik et al., 1969; Tobe, 1978). The rapid incorporation of nutrients

into milk products during lactation reduces total maternal lipid and protein content by

nearly 50% and 25%, respectively (Attardo et al., 2012). Following birth, the milk glands

rapidly involute and lipid reserves accumulate to support a subsequent offspring

(Attardo et al., 2012; Baumann et al., 2013; Benoit et al., 2018). Tsetse mothers endure

this metabolically intense K-selected reproductive strategy continuously through their

adulthood, ultimately birthing up to 8-12 larvae per lifetime (Barreaux et al., 2022;

Michalkova et al., 2014a).

Tsetse flies have evolved to feed exclusively on vertebrate blood. While blood is

nutrient-rich, it lacks in sufficient quantity several of the vitamins and cofactors required

to support the metabolically intense process of milk production. To overcome this,

tsetse flies have developed intimate, long-term associations with symbiotic bacteria that

supplement nutrients missing from host blood. One of these bacteria is the maternally

transmitted obligate endosymbiont *Wigglesworthia glossinidia*, which has associated

exclusively with tsetse species for 50-80 million years (Chen et al., 1999). As a result,

Wigglesworthia has experienced massive genome erosion (Akman et al., 2002; Hall et

al., 2019). However, the bacterium retains genes required to produce a variety of B

vitamins that are largely absent from vertebrate blood (Akman et al., 2002; Bing et al.,

2017). Metabolic processing of proline, which circulates in tsetse hemolymph and

replaces carbohydrates as the flies' prominent energy source, is fueled by B vitamins

generated by *Wigglesworthia* (Michalkova et al., 2014b). Accordingly, antibiotic-

mediated clearance of *Wigglesworthia* results in a dramatic reduction in multiple B

vitamin-associated compounds that serve as essential cofactors in fundamental

metabolic pathways, including the pentose phosphate pathway, nucleotide biosynthesis,

methionine/cysteine metabolism, and amino acid and lipid metabolism. These

deficiencies yield adult female flies incapable of milk production which consequently

results in abortion of early instar larvae (Benoit et al., 2017; Bing et al., 2017; Nogge,

1976). However, the mechanistic basis for this phenotype remains unclear.

Here we examined the interplay among obligate symbionts, lipid metabolism, and

reproductive output of tsetse flies. Maternal transfer of nutrients to developing offspring

is an energetically demanding process for all insects. However, tsetse flies, similar to

other viviparous systems, invest even more into their progeny by providing resources for

embryonic and larval development. This study demonstrates that critical symbiont-

derived factors promote phospholipid synthesis, facilitating rapid lipolysis of stored lipid

reserves in order to nourish the developing intrauterine progeny. Other insects that rely

on symbiont-derived B vitamins, including bed bugs and lice (Benoit et al., 2016; Boyd

et al., 2017; Hosokawa et al., 2010; Kirkness et al., 2010), are likely to experience

impaired lipolysis and reduced fecundity in the absences of their symbiotic partner(s),

as the molecular mechanisms underlying this process are not unique to *Glossina*

(Thomas et al., 2020). Furthermore, similarities in live birth across animal systems

indicate that micronutrient deprivation is likely to have significant impacts on metabolic

processes with similar negative consequences that extend from invertebrates to

vertebrates (Fouks et al., 2022).

Results

<u>Aposymbiotic flies are obese and fail to reproduce</u>

Antibiotic-treated, endosymbiont-free (aposymbiotic) tsetse flies failed to reproduce

successfully, as previously shown (Benoit et al., 2017; Michalkova et al., 2014b; Nogge,

1976). When early instar larvae from aposymbiotic mothers were removed before

abortion and compared to control larvae, their lipid levels were 25-30% lower, indicating

substantially reduced lipid transfer from mother to larvae in the aposymbiotic state (Fig.

1). Aposymbiotic females show higher total lipid levels, with an average of 5.66 $\pm$ 0.44

lipids per mg dry mass relative to 4.85 $\pm$ 0.34 in symbiotic females. This phenotype

was associated with increased lipid droplet (LD) size in the fat body (FB), suggesting

defect(s) in lipolysis. Additionally, Nile blue staining of FB tissue revealed a reduction in

charged phospholipids relative to neutral storage lipid species, including tri- and di-

glycerides, in aposymbiotic flies relative to control flies (Fig. 1). These initial studies

suggest a major dysfunction in lipid metabolism following symbiont clearance.

<u>Lipidomics reveals massive shifts in metabolic lipid processing</u>

To understand the biochemical landscape facilitating the accumulation of

triacylglycerols in endosymbiont-free flies, we generated aposymbiotic females via

tetracycline application, and analyzed pooled FB and milk gland (MG) tissues,

comparing aposymbiotic and untreated (control), age-matched females. Analysis of the

differential biochemical composition revealed substantial alterations in the composition

of multiple biochemical categories in aposymbiotic flies, including B vitamins,

creatine/creatinine, amino acids, diacylglycerols (DAG), phospholipids, sphingolipids,

lysolipids and polyunsaturated fatty acids (Suppl. Figs. 1-6). These changes yield an

overall composition difference in the lipid profiles between control and aposymbiotic flies

(Suppl. Fig. 7, ANOSIM, R = 0.879, P = 1.03E-06).

Many of the observed changes are via either direct or indirect effects of

dysregulation of phospholipid (PL) biosynthesis via the Kennedy pathway (Fig. 2).

Notably, aposymbiotic flies were deficient in key precursors to PL synthesis including

choline, phosphocholine, and the majority of the detected DAGs, and

phosphatidylcholines (PCs). Many DAG variants are present in adult female tsetse, and

they differ in acyl chain lengths and desaturation state. The most abundant DAGs in

control flies contain 16/18 carbon unsaturated or mono-unsaturated side chains (Fig. 2

and Suppl. Fig. 2), while aposymbiotic flies show a broad reduction in abundance of

these DAGs. Among phosphatidylcholines, aposymbiotic flies generally lack moieties

containing 16/18 carbon unsaturated or mono-unsaturated side chains, while those

containing 18/20 carbon polyunsaturated fatty acids are abundant (Suppl. Fig. 1). This

may reflect the observed overabundance of polyunsaturated free fatty acids in

aposymbiotic flies (Suppl. Fig. 4). The substantial deficiencies in the precursor

molecules required for phosphatidylcholine production suggest that *Wigglesworthia*-

derived factors either directly or indirectly facilitate the production of phospholipid

precursors.

Aposymbiotic flies also demonstrate a profound deficiency in creatine and

creatinine (Suppl. Fig. 6). Creatine/creatinine biosynthesis and the urea cycle are

dependent on folate, which is provided to tsetse flies by *Wigglesworthia* (Snyder and

Rio, 2015). Components of the urea cycle are also reduced in abundance in

aposymbiotic flies. The urea cycle is also dependent on folate for proper function

(Brosnan and Brosnan, 2010; Fatterpaker et al., 1951). Creatine/creatinine are key

regulators of lipid homeostasis for adrenergic and diet-induced thermogenesis in brown

and beige adipose tissue in vertebrates. Induced deficiencies in in creatine/creatinine in

mice results in diet induced obesity, which can be rescued by dietary creatine

supplementation (Kazak et al., 2017). The mechanism behind how these compounds

regulate lipolysis remains to be elucidated but may also be involved in maintenance of

lipid homeostasis throughout the reproductive cycle.

<u>Choline-phosphate cytidylyltransferase 1 represents a critical enzyme during tsetse fly</u>

<u>reproductive cycles</u>

Prior RNA-seq analyses of gene expression during tsetse pregnancy and milk

production revealed a suite of highly-expressed milk protein-encoding genes (Attardo et

al., 2019; Benoit et al., 2014; International Glossina Genome Initiative, 2014). Notably,

transcripts for the enzyme choline-phosphate cytidylyltransferase 1 (*cct1*) were

upregulated in a similar manner to milk proteins in pregnant flies (Fig. 3). This enzyme

participates in phosphatidylcholine synthesis. Phosphatidyl cholines are critical for

regulation of LD size and formation to facilitate lipolysis during energy-intensive

processes (Krahmer et al., 2011). The expression of *cct1* increases during milk gland

involution following parturition (Fig. 3). Tissue-specific expression analysis confirmed

cct1 transcripts were abundant in the FB and MG. In support of this finding, *in situ*

hybridization revealed *cct1* expression in both of these tissues (Fig. 3C). These data

suggest that *cct1* is critical for lipid metabolism during tsetse pregnancy, likely

functioning in the FB and MG, to break down lipids critical as energy for milk production,

or necessary to produce milk LDs.

<u>Suppression of cct1 phenocopies the aposymbiotic state</u>

To confirm if impaired *cct1* function and altered phospholipid metabolism perturbs lipid

metabolism during viviparity, we suppressed *cct1* expression using RNA interference

(RNAi) (Fig. 4). Similar to aposymbiotic flies, *cct1* knockdown flies produce few progeny

(Fig. 4). We noted a massive decrease in levels of phospholipids and PC in developing

larvae, indicating that *cct1* is necessary to ensure sufficient lipid transfer to the

developing larva. This decrease likely accounts for the 40-45% reduction in total lipids

measured from larvae deposited from *cct1* knockdown flies (0.20 $\pm$ 0.03 mg lipid/ mg

dry mass for *cct1* RNAi vs 0.29 $\pm$ 0.04 for control). In aposymbiotic females, *cct1*

expression increases substantially, perhaps as a compensatory response to reduced

phospholipid levels, which are required for lipid transfer during pregnancy.

Tetracycline treatment impairs mitochondrial function (Moullan et al., 2015),

which could represent an off-target outcome that significantly impacts tsetse lipid

metabolism. As such, we treated flies with glyphosate, which specifically inhibits

Wigglesworthia folate production (Rio et al., 2019). This yielded flies that presented

obesity measures similar to their aposymbiotic counterparts (Fig. 4G). Lastly,

aposymbiotic flies were fed on blood with a cocktail of B vitamins and yeast extract,

which reduced the lipid levels within the flies, providing a direct link between B vitamins

and obesity levels (Fig. 4H). This finding confirmed that *Wigglesworthia*-derived B

vitamins acts as an essential factor for tsetse lipid homeostasis.

Discussion

This study defines a novel feature of host-symbiont dynamics in tsetse showing

dysfunctional PL metabolism as a resultant phenotype of symbiont elimination (Fig. 5).

Tsetse fly pregnancy requires a rapid transfer of substantial nutritional resources from

mother to offspring. A key mechanism in this process is the production of PC by the

Kennedy biosynthesis pathway. The *cct1* gene encodes a rate-limiting enzyme in this

pathway that prepares the choline headgroup for attachment to cytidyl triphosphate to

generate CDP-choline. CDP-choline is then combined with diacylglycerol to produce the

final phosphatidyl choline product. The observed increase in *cct1* transcripts during

pregnancy suggests that this enzyme/pathway is critical to provision of PC for LD

production/function for lipolysis in the FB and for incorporation into products secreted

from the MG (Benoit et al., 2015; International Glossina Genome Initiative, 2014).

Production of PC is required to coat the outer surface of LDs produced by the

endoplasmic reticulum, and is thus critical for lipid metabolism (Olzmann and Carvalho,

2019). The amphipathic nature of PC prevents aberrant LD fusion and provides the

biochemical conditions to recruit LD-associated proteins required for lipolysis (Pol et al.,

2014).

In mammals, disruption of appropriate PC to PE ratios in LDs causes formation

of large droplets with reduced surface area and altered surface protein composition.

The end result is abnormal fat retention within a cell or tissue/organ (steatosis;

(Listenberger et al., 2018). In tsetse flies, symbiont clearance produces obesity, similar

to that observed in mammals where individuals see increased lipid levels in conjunction

with symbiont dysfunction. Flies initially fed blood spiked with tetracycline are cleared of

their *Wigglesworthia*, with any lingering bacterial-derived factors clearing through

subsequent feedings. Importantly, multiple feeding events following tetracycline

treatment over the course of two weeks reduces the direct impact of this treatment,

indicating that the phenotypes are associated with the loss of the symbiont. Symbiont

absence results in PC deficiency that inhibits metabolism, mobilization and integration

of stored lipids during lactation. Reproduction is terminated due to lipid retention in the

FB and inefficient lipid transfer to the intrauterine larva. This is likely due to the

combined effect of the mothers inability to process FB lipids due to dysfunction in lipid

droplet formation in the FB and MG as *cct1* is abundantly expressed in both locations.

The biochemical observations associated with this phenotype suggest at least

two points of failure in phosphatidylcholine biosynthesis via the Kennedy (CDP-choline)

pathway. The first limiting factor appears to be deficiencies in choline and choline

phosphate levels, which form the phosphatidylcholine headgroup. The production of

choline is dependent on methylation reactions driven by enzymes that utilize S-adenosyl

methionine (SAM) as a cofactor. Prior work in aposymbiotic *Glossina* demonstrates a

severe deficiency of SAM in aposymbiotic flies (Bing et al., 2017). Production of SAM is

dependent on the cysteine-methionine pathway, which is dependent on the B vitamins

pyridoxal and folate as enzymatic cofactors (Loenen, 2006; Nguyen et al., 2001).

Methylation reactions require SAM as a cofactor for the production of choline and

conversion of phosphatidylethanolamine (PE) to PC (Gibellini and Smith, 2010; Mazen

Noureddin et al., 2015; Snyder and Rio, 2015). In accordance with the observations in

mammalian systems, the mechanistic disruption occurring in the aposymbiotic flies is

most likely due to the deficiency in symbiont-derived B vitamins.

Wigglesworthia produces an array of B vitamins (B1, B2, B3, B5, B6, B7 and B9)

(Akman et al., 2002; Bing et al., 2017; Rio et al., 2019; Snyder and Rio, 2015).

Deficiencies in B6, folate and, consequently, SAM have a negative impact on

phospholipid biosynthesis in mammalian systems. Issues related to the production of

phosphatidylcholine have been linked to folate restriction (Abratte et al., 2008). The

aposymbiotic tsetse phenotype bears resemblance to the pathology observed in non-

alcoholic fatty liver disease (NAFLD) in mammals as PC deficiency is a key dysfunction

(Bianca M. Arendt et al., 2013; Jacobs et al., 2013; Ville Männistö et al., 2019). Rats

exhibiting a NAFLD phenotype have been rescued by dietary supplementation with

phosphatidylcholine (Kitagawa et al., 2015). The phenotype observed in aposymbiotic

flies (this study and in prior studies (Bing et al., 2017; Snyder and Rio, 2015)) is

phenotypically equivalent to that observed when *cct1* is suppressed. Folate deficiency is

likely a major underlying factor for the phenotype of dysfunctional metabolism of lipids

we observe in flies lacking symbionts. Induction of folate deficiency in tsetse through

treatment with glyphosate impacts vector competency and reproduction (Rio et al.,

2019; Snyder and Rio, 2015), highlighting that folate provision by *Wigglesworthia*

represents a critical lynchpin in tsetse metabolism. While female tsetse are able to

survive B vitamin deficiency, the metabolic deficiencies prevent them from performing

the mass mobilization of lipids from FB stores to the MG required for milk production

during pregnancy. Our supplementation of B vitamins to aposymbiotic flies reduces

obesity and has been previously shown to recover fecundity (Michalkova et al., 2014b),

providing a direct link between B vitamins provided by *Wigglesworthia* and obesity in

Glossina. This is not surprising as *cct* is overexpressed in aposymbiotic lines, indicating

once specific B vitamins are available lipolysis can occur more rapidly to support

reproduction.

269   Another disruption in phosphatidylcholine biosynthesis appears to derive from

DAG deficiency. The lack of DAGs results in inadequate substrate availability for fusion

with cytidine 5'-diphosphocholine by DAG cholinephosphotransferase. DAGs can be

synthesized *de novo* during the production of triglycerides or they can be produced

during lipolysis of triglycerides upon removal of a fatty acid side chain by a lipase.

Triglycerides are overabundant in aposymbiotic flies, suggesting that lipolysis of the

existing triglycerides is compromised under these conditions. In insects, the lipase

responsible for this activity is called *brummer lipase* (*Drosophila*, CG5295), and is

orthologous to the Adipose triglyceride lipase (ATGL) in mammals (Alexander Mildner et

al., 2005; Grönke et al., 2007). Brummer and ATGL are both regulated by the

phosphorylation state of other lipid droplet bound proteins, called perilipins (Sztalryd

and Brasaemle, 2017). Perilipins sequester a co-activator of ATGL in their non-

phosphorylated state called CGI-58, which inhibits lipolysis (Itabe et al., 2017). Recent

work demonstrates that LDs lacking proper phosphatidylcholine levels show increased

levels of associated perilipins and a steatosis (lipid accumulation) phenotype

(Listenberger et al., 2018). The phosphatidylcholine deficiencies in aposymbiotic

Glossina could be functioning via the same paradigm, resulting in perilipin accumulation

and inhibition of *brummer*-mediated lipolysis resulting in DAG deficiency and TAG

accumulation.

The symbiosis between tsetse flies and *Wigglesworthia* is essential for this

complex system and both are dependent on one another for their survival. These

studies demonstrate that lipid metabolism, a foundational process in *Glossina* biology, is

impaired in the absence of their obligate symbionts. This is likely a model for other

symbiotic systems, particularly those compensating for dietary nutritional deficiencies

such as blood feeding invertebrates that are supplied with B vitamins from bacterial

symbionts. This includes ticks (Duron et al., 2018), bed bugs (Nikoh et al., 2014), and

lice (Boyd et al., 2017), which are all epidemiologically important blood-feeding disease

vectors and require obligate symbionts to provide B vitamins. Beyond blood-feeding

arthropods, specific bacteria associated with plant feeding insects also provide B

vitamins (Salem et al., 2014), suggesting that lipid metabolism and B vitamin

dependency issues likely occur in many systems. Lastly, B vitamins can be provided to

insect species that have transient bacterial symbionts, such as in mosquitoes (Wang et

al., 2021), suggesting that the interdependence of B vitamins and lipid metabolism may

be a widespread phenomenon among arthropod species. As a key example,

mosquitoes display altered lipid metabolism related to microbiome elimination and B

vitamin deficiency (Didion et al., 2021; Romoli et al., 2021; Valzania et al., 2018; Wang

et al., 2021). Specifically, the mosquitoes fail to accumulate lipids at critical time points,

such as early in development and when lipids are accumulated for dormancy (Didion et

al., 2021; Romoli et al., 2021; Valzania et al., 2018; Wang et al., 2021), indicating a

critical role for B vitamins, lipid metabolism, and symbiont presence for many biological

processes, including the critical contribution to reproduction we observed in this study.

Along with the role among arthropod systems, genomic studies have identified

significant functional convergence among live-bearing systems in invertebrate and

vertebrate lineages (Fouks et al., 2022). This includes aspects that range from

metabolism to immune function (Fouks et al., 2022). In the vertebrate systems, fetal

development requires essential fatty acids and long chain polyunsaturated fatty acids

(Herrera, 2002; Knopp et al., 1998; Ryckman et al., 2015). As in the insect system, the

accumulation of lipids early in vertebrate pregnancy is a critical feature that builds up

stores providing lipids to the developing progeny (Herrera, 2002). As pregnancy

progresses, the mother transitions from lipogenesis to lipolysis to increase the lipid

supply to the fetus (Herrera et al., 1994, 1988). An imbalance in lipid levels provided to

the neonates, either through malnutrition of the mother or impaired transfers, can yield

many negative consequences ranging from delayed growth/development to death

(Furness et al., 2012; Herrera, 2002). Similar to this study, B vitamin levels have been

documented to alter lipid metabolism in mammalian systems with most focus on human

and murine systems (Castaño-Moreno et al., 2020; da Silva et al., 2020; Herrera, 2002).

Low B vitamin levels during pregnancy have been directly associated with a reduction in

circulating lipid levels, increased obesity in the mother, and a reduction in health of the

neonate (Adaikalakoteswari et al., 2017; Castaño-Moreno et al., 2020; McNeil et al.,

2009), which is functionally similar to the obesity and reduced larval growth we

observed in tsetse flies lacking B vitamins produced by *Wigglesworthia*. As observed in

this study for tsetse flies, the induction of obesity acts through impaired PC synthesis

that requires specific B vitamins, which is also critical for vertebrate lipid metabolism (da

Silva et al., 2014). Thus, a distinct commonality of viviparity among animals is the

requirements of specific B vitamins for lipid metabolism to meet the energetic demand

of internal progeny development. This expands on genomic similarities (Fouks et al.

2022) to other physiological processes that are similar between invertebrate and

vertebrate viviparity.

Methods

Flies

Glossina morsitans morsitans flies were maintained on defibrinated bovine blood

(Lampire Biologicals, Pipersville, PA, USA) at 25°C and 50-60% RH using an artificial

membrane feeding system (Moloo, 1971) at the insectaries at Yale School of Public

Health and the Slovak Academy of Sciences. Females were mated 3-5 days after

emergence. Flies were collected according to established developmental markers

based on oocyte, embryo and larva presence (Attardo et al., 2006; Strickler-Dinglasan

et al., 2006; Yang et al., 2010). Aposymbiotic tsetse larvae were derived from females

fed a diet supplemented with tetracycline (20 µg per ml of blood) to clear their

indigenous microbiota and yeast extract (1%) to provide vitamins in the absence of

obligate *Wigglesworthia* (Pais et al., 2008). A subset of the progeny were allowed to

feed of blood supplemented with B vitamins and yeast extract according to Michalkova

et al. (2014b), which has been shown to partially alleviate B vitamin deficiencies

following *Wigglesworthia* elimination.

Milk extraction

Milk products were extracted from the guts of developing 3rd instar larvae based on our

previous studies (Benoit et al., 2014, 2012). Briefly, actively-feeding individuals were

removed from the uterus and their digestive tracts were dissected. A pulled glass

capillary needle was utilized to pierce into the digestive tract and the contents were

removed using reverse pressure. Each sample was stored at -80ºC until utilization in the

experiments.

Lipidomics

Tissues (milk glands and fat body) were dissected from aged-matched WT flies that

received normal blood meals for five weeks (control - symbiotic) and aposymbiotic

females and prepared according to our previously established protocol (Bing et al.,

2017). Briefly, tissues were added 100 μl of ice cold 1X phosphate buffered saline

(PBS), homogenized with a pestle and stored at -80° C. For both aposymbiotic and

control flies, samples consisted of four replicates of 20 individuals each. Samples were

shipped to Metabolon (Morrisville, NC, USA) and analysis was performed using the

proprietary Metabolon DiscoveryHD4 global lipidomics platform. Each sample was

screened for the presence and quantity of lipids by mass spectrometry using a panel

specific lipids. Relative differences between samples were determined by comparison of

mean metabolite abundance across the biological replicates. Significance of the

difference between symbiotic and aposymbiotic samples was determined by Welch's

two-sample t-test: p-value cut off of (<0.05). False discovery rate (FDR) was tested for

by q-value with a cut off of (<0.1). Non-Metric Multidimensional Scaling (NMDS) plots

were created using the metaMDS function from the R package vegan (Dixon, 2003;

Oksanen, 2007) to determine differences in the general lipid composition between

control and aposymbiotic individuals. Analysis of similarities (ANOSIM, using the

anosim from the R package vegan) tests was performed to test for significance in the

general differences.

RNA extraction and qPCR

RNA was extracted from flies with TRizol reagent (Invitrogen, Carlsbad, CA, USA) using

the manufacturer's recommended protocol. RNA was treated with TURBO DNA Free kit

(Ambion, Austin, TX, USA). RNA was precipitated with alcohol to remove residual salt,

purified with a RNeasy kit (Qiagen, Maryland, USA) and stored at -70ºC until use. cDNA

was prepared using the Superscript III reverse transcriptase (Invitrogen).

Transcript abundance was determined utilizing quantitative PCR (qPCR) on a Bio-Rad

CFX detection system (Hercules). Primer sequences used were those from Table S1.

Results were analyzed with CFX manager software version 3.1 (Bio-Rad). Ct values for

genes of interest were standardized by Ct values for the control gene (tubulin) and

relative to the average value for the control treatment or newly emerged flies, yielding a

relative fold change in gene expression.

RNA interference of *cct1*

Suppression of *cct1* was performed utilizing double stranded RNA (dsRNA) according to

our previous studies on tsetse gene suppression (Benoit et al., 2017, 2012). The T7

promoter sequence was added to the 5′ end of the primer sequences and PCR

amplification conditions are described (Table S1). The PCR products were purified

using QIAquick PCR purification kit (Qiagen, Valencia, CA) and cloned into pGEM T-

Easy vector (Promega, Madison, WI) and verified by sequencing (Keck DNA

sequencing facility, Yale University). dsRNAs were synthesized using the MEGAscript

RNAi Kit (Ambion, Austin, TX), purified using a RNeasy Mini Kit (Qiagen, Valencia, CA)

and siRNAs were generated by using the Block-iT Dicer RNAi kit (Invitrogen, Carlsbad,

CA). The siRNA concentration was adjusted to 600–800 ng/µl in PBS and each fly was

injected with 1.5 µl siRNA using a pulled glass capillary needle into their thorax.

Concentration of siRNAs were kept at 600–800 ng/µl within the 1.5 µl injected.

Knockdown efficiency was determined 5d after injection by qPCR and normalized to

tubulin levels. Fecundity following *cct1* was assessed according to previously

established protocols (Benoit et al., 2014, 2012).

<u>Glyphosate treatment</u>

Groups of *G. morsitans* flies were maintained on a diet containing 100 µM glyphosate

100 µM glyphosate combined with 500 nM folic acid [N-(phosphonomethyl)glycine;

Sigma-Aldrich, St. Louis, MO, USA] or (Sigma-Aldrich). This treatment has been

previously documented to impact folate synthesis (Rio et al., 2019; Snyder and Rio,

2015). Following six bloodmeals, the flies were examined to determine if lipid and

phosphotydilcholine levels were increased or decreased. These treatments were

conducted to eliminate off-target effects of tetracycline supplementation validate

Wigglesworthia's role in tsetse lipid metabolism provide a secondary validation of

impaired lipid metabolism without treatment with antibiotics.

<u>Total lipid assay</u>

Total lipids present within biological samples were determined with a standard vanillin

assay (Attardo et al., 2012; Rosendale et al., 2019; Van Handel, 1985). Whole flies and

individual tissues were dried at 0% RH at 60ºC and weighed to determine the dry mass.

The flies were homogenized in 2 ml of chloroform:methanol (2:1). The supernatant was

removed and placed into a glass tube and the solvent was evaporated at 90ºC. The

dried lipids were treated with 0.4 ml of concentrated sulfuric acid at 90ºC for 10 min. The

acid/lipid mixture (40 µl) was combined with 4 ml vanillin reagent. Samples were

measured spectrophotometrically at 525 nm and the total lipid content was calculated

against a lipid standard.

In situ hybridization for *cct1*

Milk gland tubules and fat body were removed from tsetse fly females at six-seven days

after the previous birth. The combined milk gland/fat body was placed into Carnoy's

fixative for a five-six day fixation period (Attardo et al., 2008; Michalkova et al., 2014a).

Antisense/sense digoxigenin-labeled RNA probes for *cct1* were generated using the

MAXIscript T7 transcription kit following manufacturer's protocol (Ambion, Austin, TX,

USA) using a primer set with a T7 primer (Table S1). Antibody solutions were made

using α-Digoxigenin-rhodamine, Fab fragments (Roche Applied Science, Penzberg,

Germany) for FISH probe detection (1:200 dilution) and rabbit α-GmmMGP (1:2500)

antibodies. Alexa Fluor 488 goat α-rabbit IgG (Invitrogen) at a dilution of 1:500 was

added as a secondary antibody for immunohistochemistry (Attardo et al., 2008;

Michalkova et al., 2014a). Slides were mounted in VECTASHIELD Mounting Medium

with DAPI (Vector laboratories Inc. Burlingame, CA, USA). Samples were observed

using a Zeiss Axioskop2 microscope (Zeiss, Thornwood, NY, USA) equipped with a

fluorescent filter and viewed and imaged at 400x magnification. Images were captured

using an Infinity1 USB 2.0 camera and software (Lumenera Corporation, Ottawa,

Ontario, Canada) and merged in Adobe Photoshop.

Phosphatidylcholine assay

Phosphatidylcholine levels were assessed through the use of a colorimetric assay

(Caymen Chemicals, Ann Arbor, MI, USA). Extracted milk from the guts of 3rd instar

larvae (10 µl) was placed within the sample well of a 96 well plate. The colorimetric

reaction was initiated by adding 100 µl of the reaction mixture. Following mixing for a

few seconds, the plate was incubated at 37°C for 60 min and concentration determined

by comparison to a phosphatidylcholine standard.

<u>Nile Blue staining</u>

Lipid composition of fat body cells from control and aposymbiotic *Glossina* was

analyzed microscopically using the lipophilic Nile Blue stain. Nile Blue is a differential

stain, which results in charged phospholipids (fatty acids, chromolipids and

phospholipids) being stained blue and neutral lipids (triglycerides, cholesterol esters,

steroids) being stained red/pink. The method used here is a modified version of the

protocol described by Canete and colleagues (Canete et al., 1983). Fat body tissues

were dissected from flies reared as described in the Lipidomics section above.

Dissected tissues were immediately transferred to ice cold 4% paraformaldehyde in

PBS and allowed to fix overnight at 4°C. Fixed tissues were then rinsed in PBS followed

by immersion in 0.1 mg/ml aqueous Nile Blue sulfate (Sigma-Aldrich) for 30 min. After

staining, tissues were rinsed three times with PBS for 1 min per rinse to remove excess

stain. Tissues were then mounted in glycerol on single depression concave microscope

slides. Stained fat bodies were visualized using a Zeiss Axio Vert.A1 scope and images

captured using the Zeiss Zen imaging software. Color differences were quantified by

measuring the average amount of red and blue color in five 100 pixel squares within the

center of each lipid droplet with the use of Adobe Photoshop.

<u>Statistical analyses</u>

Lipidomics data were analyzed using the Metaboanalyst software package

(https://www.metaboanalyst.ca/) (Pang et al., 2021). Metabolite abundance values

represent peak area counts measured from compound peaks. Samples were

normalized across samples using the sum of raw area counts for each sample. Missing

values were imputed by replacement with minimum detected values for each

metabolite. Data were rescaled by median centering to a value of 1 for each sample.

Significant differences between control and aposymbiotic samples were measured by t-

test followed by p-value correction using false discovery rate (FDR)/Benjamini and

Hochberg analysis. Differences in relative abundance were considered significant below

an FDR corrected P-value of 0.05. Differences in other biological attributes were

assessed through the use of t-test or ANOVA.

Funding

This study was supported by awards from National Institutes of Health USA (AI109263,

AI128523, F32AI093023, and AI08177405), FAO/IAEA Coordinated Research Program

"Improving SIT For Tsetse Flies through Research on their Symbionts" and

"Improvement of Colony Management in Insect Mass-rearing for SIT Applications", and

the Cariplo-Regione Lombardia research grant IMPROVE (F.S.).

Figure legends

Figure 1: Reproduction and lipid metabolism is dysfunctional in aposymbiotic flies.
Proportion of females giving birth (A), N = 4, and lipid content in the surviving larvae (B)
are significantly reduced in aposymbiotic flies (t-test, P-value 0.007 and 0.0003,
respectively). N = 8-10. C-H) Nile Blue staining reveals significant differences in the
neutral and charged lipid composition in aposymbiotic and control flies. I) Volume of the
fat body is increased significantly in aposymbiotic flies (t-test, P-value 0.008). N = 12-14.
J) Image quantification reveals significantly higher charged lipids in the control
compared to the aposymbiotic flies (t-test, P-value 0.002). N = 12-14.

Figure 2: Lipidomics reveals a significant shift in lipid moetities following symbiont
removal. Compounds showing a significantly different fold change in their abundance in
control (yellow) and aposymbiotic (green) flies are shown. Red denotes increase in
compounds of the Kennedy pathway in aposymbiotic flies, while blue indicates lipids
showing higher fold change in control individuals. Specifically, there is a major
dysfunction in the Kennedy pathway for phosphatidylcholine synthesis. N = 5, *
indicates significance based on a t-test.

Figure 3: Increased expression of *choline-phosphate cytidylyltransferase 1* (*cct1*) is
associated with tsetse fly pregnancy. A) Previous RNA-seq studies (Attardo et al., 2019;
Benoit et al., 2014) reveal that *cct1* is expressed in pregnant/lactating flies. B) qPCR
indicated that the expression pattern for *cct1* is correlated with the pregnancy cycle,
similar to a milk protein critical to feed the developing larvae (*milk gland protein 1*,

mgp1). N = 4. C) Expression of *cct1* is localized in both the milk glands and fat body,

suggesting critical roles in lipid breakdown and milk production. N = 4, *, denotes

significance based on a Kal's proportion test or t-test.

Figure 4: Suppression of *cct1* produces aposymbiotic phenotypes of altered lipid

metabolism during pregnancy. A) RNA interference reduces the expression of *cct1*. N =

5, *, denotes significance based on a t-test. *Mgp1, milk gland protein 1.* B-C) Total

progeny production and progeny generated per day are reduced following suppression

of *cct1*. N = 3 groups of 10. Different letters denote significance based on an ANOVA.

Acid smase1, *asmase1*. D-E) Milk phospholipids and phosphatidylcholine are reduced

when *cct1* is reduced. N = 12. *, denotes significance based on a t-test. F) Aposymbiotic

flies show increased expression of *cct1* as a potential compensatory mechanism to

allow for lipolysis during pregnancy. N = 10. *, denotes significance based on a t-test. G)

Glyphosate treatment increased lipid levels within flies. N = 8. *, denotes significance

based on a t-test. H) Blood supplementation with B vitamins or yeast extract reduced

obesity in flies. N = 10. *, denotes significance based on ANOVA.

Figure 5: Summary of the dynamics between symbiont loss and reproduction during

tsetse fly pregnancy in relation to altered lipolysis. The percentage annotations in the

panel depicting the Kennedy Pathway represent the relative amount of the specific lipid

metabolite in the aposymbiotic tsetse fly relative to the untreated, wild-type control

which would be equivalent to 100%.

Supplemental Figure 1: Absolute abundance of detected fat body/milk gland

phospholipids and associated compounds in symbiotic and aposymbiotic *G. morsitans.*

Abundance values represent normalized area under compound peaks. Significance

determined by Welch's Two-Sample t-test followed by FDR adjustment of resulting p-

values.

Supplemental Figure 2: Absolute abundance of detected fat body/milk gland

diacylglycerols in symbiotic and aposymbiotic *G. morsitans.* Abundance values

represent normalized area under compound peaks. Significance determined by Welch's

Two-Sample t-test followed by FDR adjustment of resulting p-values.

Supplemental Figure 3: Absolute abundance of detected fat body/milk gland

sphingolipids and associated compounds in symbiotic and aposymbiotic *G. morsitans.*

Abundance values represent normalized area under compound peaks. Significance

determined by Welch's Two-Sample t-test followed by FDR adjustment of resulting p-

values.

Supplemental Figure 4: Absolute abundance of detected fat body/milk gland

polyunsaturated fatty acids (PUFAs) in symbiotic and aposymbiotic *G. morsitans.*

Abundance values represent normalized area under compound peaks. Significance

determined by Welch's Two-Sample t-test followed by FDR adjustment of resulting p-

values.

Supplemental Figure 5: Absolute abundance of detected fat body/milk gland lysolipids

in symbiotic and aposymbiotic *G. morsitans.* Abundance values represent normalized

area under compound peaks. Significance determined by Welch's Two-Sample t-test

followed by FDR adjustment of resulting p-values.

Supplemental Figure 6: Absolute abundance of detected fat body/milk gland amino

acid and vitamin B associated metabolic pathways undergoing significant changes in

abundance between in symbiotic and aposymbiotic *G. morsitans.* Abundance values

represent normalized area under compound peaks. Significance determined by Welch's

Two-Sample t-test followed by FDR adjustment of resulting p-values.

Supplemental Figure 7: NMDS analysis comparing the lipid profile differences

between aposymbiotic and control flies. Significant variation between aposymbiotic and

control samples (ANOSIM, R = 0.879, P = 1.03E-06).

Table S1: Primers used for qPCR, generation of dsRNA, and *in situ* hybridization.

References

Abratte CM, Wang W, Li R, Moriarty DJ, Caudill MA. 2008. Folate intake and the MTHFR C677T genotype influence choline status in young Mexican American women. *J Nutr Biochem* **19**:158–165.

Adaikalakoteswari A, Vatish M, Alam MT, Ott S, Kumar S, Saravanan P. 2017. Low Vitamin B12 in Pregnancy Is Associated With Adipose-Derived Circulating miRs Targeting PPARγ and Insulin Resistance. *J Clin Endocrinol Metab* **102**:4200–4209.

Akman L, Yamashita A, Watanabe H, Oshima K, Shiba T, Hattori M, Aksoy S. 2002. Genome sequence of the endocellular obligate symbiont of tsetse flies, Wigglesworthia glossinidia. *Nat Genet* **32**:402–407.

Alexander Mildner, Sonja Fellert, Norbert Tennagels, Günter Müller, Herbert Jäckle, Sebastian Grönke, Stefan Petry, Ronald P. Kühnlein. 2005. Brummer lipase is an evolutionary conserved fat storage regulator in Drosophila. *Cell Metab*. doi:10.1016/j.cmet.2005.04.003

Attardo GM, Abd-Alla AMM, Acosta-Serrano A, Allen JE, Bateta R, Benoit JB, Bourtzis K, Caers J, Caljon G, Christensen MB, Farrow DW, Friedrich M, Hua-Van A, Jennings EC, Larkin DM, Lawson D, Lehane MJ, Lenis VP, Lowy-Gallego E, Macharia RW, Malacrida AR, Marco HG, Masiga D, Maslen GL, Matetovici I, Meisel RP, Meki I, Michalkova V, Miller WJ, Minx P, Mireji PO, Ometto L, Parker AG, Rio R, Rose C, Rosendale AJ, Rota-Stabelli O, Savini G, Schoofs L, Scolari F, Swain MT, Takáč P, Tomlinson C, Tsiamis G, Van Den Abbeele J, Vigneron A, Wang J, Warren WC, Waterhouse RM, Weirauch MT, Weiss BL, Wilson RK, Zhao X, Aksoy S. 2019. Comparative genomic analysis of six Glossina genomes, vectors of African trypanosomes. *Genome Biol* **20**:187.

Attardo GM, Benoit JB, Michalkova V, Yang G, Roller L, Bohova J, Takáč P, Aksoy S. 2012. Analysis of lipolysis underlying lactation in the tsetse fly, Glossina morsitans. *Insect Biochem Mol Biol* **42**:360–370.

Attardo GM, Guz N, Strickler-Dinglasan P, Aksoy S. 2006. Molecular aspects of viviparous reproductive biology of the tsetse fly (Glossina morsitans morsitans): Regulation of yolk and milk gland protein synthesis. *Journal of Insect Physiology*. doi:10.1016/j.jinsphys.2006.07.007

Attardo GM, Lohs C, Heddi A, Alam UH, Yildirim S, Aksoy S. 2008. Analysis of milk gland structure and function in Glossina morsitans: milk protein production, symbiont populations and fecundity. *J Insect Physiol* **54**:1236–1242.

Barreaux AMG, Higginson AD, Bonsall MB, English S. 2022. Incorporating effects of age on energy dynamics predicts nonlinear maternal allocation patterns in iteroparous animals. *Proc Biol Sci* **289**:20211884.

Baumann AA, Benoit JB, Michalkova V, Mireji P, Attardo GM, Moulton JK, Wilson TG, Aksoy S. 2013. Juvenile hormone and insulin signaling pathways regulate lipid levels during lactation and dry periods of tsetse fly pregnancy. *Mol Cell Endocrinol* **372**:30–41.

Benoit JB, Adelman ZN, Reinhardt K, Dolan A, Poelchau M, Jennings EC, Szuter EM, Hagan RW, Gujar H, Shukla JN, Zhu F, Mohan M, Nelson DR, Rosendale AJ, Derst C, Resnik V, Wernig S, Menegazzi P, Wegener C, Peschel N, Hendershot JM, Blenau W, Predel R, Johnston PR, Ioannidis P, Waterhouse RM, Nauen R,

638 Schorn C, Ott M-C, Maiwald F, Johnston JS, Gondhalekar AD, Scharf ME,
639 Peterson BF, Raje KR, Hottel BA, Armisén D, Crumière AJJ, Refki PN, Santos ME,
640 Sghaier E, Viala S, Khila A, Ahn S-J, Childers C, Lee C-Y, Lin H, Hughes DST,
641 Duncan EJ, Murali SC, Qu J, Dugan S, Lee SL, Chao H, Dinh H, Han Y,
642 Doddapaneni H, Worley KC, Muzny DM, Wheeler D, Panfilio KA, Vargas Jentzsch
643 IM, Vargo EL, Booth W, Friedrich M, Weirauch MT, Anderson MAE, Jones JW,
644 Mittapalli O, Zhao C, Zhou J-J, Evans JD, Attardo GM, Robertson HM, Zdobnov
645 EM, Ribeiro JMC, Gibbs RA, Werren JH, Palli SR, Schal C, Richards S. 2016.
646 Unique features of a global human ectoparasite identified through sequencing of
647 the bed bug genome. *Nat Commun* **7**:10165.

648 Benoit JB, Attardo GM, Baumann AA, Michalkova V, Aksoy S. 2015. Adenotrophic
649 Viviparity in Tsetse Flies: Potential for Population Control and as an Insect Model
650 for Lactation. *Annual Review of Entomology*. doi:10.1146/annurev-ento-010814-
651 020834

652 Benoit JB, Attardo GM, Michalkova V, Krause TB, Bohova J, Zhang Q, Baumann AA,
653 Mireji PO, Takáč P, Denlinger DL, Ribeiro JM, Aksoy S. 2014. A novel highly
654 divergent protein family identified from a viviparous insect by RNA-seq analysis: a
655 potential target for tsetse fly-specific abortifacients. *PLoS Genet* **10**:e1003874.

656 Benoit JB, Attardo GM, Michalkova V, Takác P, Bohova J, Aksoy S. 2012.
657 Sphingomyelinase activity in mother's milk is essential for juvenile development: a
658 case from lactating tsetse flies. *Biol Reprod* **87**:17, 1–10.

659 Benoit JB, Michalkova V, Didion EM, Xiao Y, Baumann AA, Attardo GM, Aksoy S. 2018.
660 Rapid autophagic regression of the milk gland during involution is critical for
661 maximizing tsetse viviparous reproductive output. *PLoS Negl Trop Dis*
662 **12**:e0006204.

663 Benoit JB, Vigneron A, Broderick NA, Wu Y, Sun JS, Carlson JR, Aksoy S, Weiss BL.
664 2017. Symbiont-induced odorant binding proteins mediate insect host
665 hematopoiesis. *Elife* **6**. doi:10.7554/eLife.19535

666 Bianca M. Arendt, David W.L. Ma, Brigitte Simons, Seham Noureldin, George
667 Therapondos, Maha Guindi, Morris Sherman, Johane P. Allard. 2013. Nonalcoholic
668 fatty liver disease is associated with lower hepatic and erythrocyte ratios of
669 phosphatidylcholine to phosphatidylethanolamine. *Appl Physiol Nutr Metab*.
670 doi:10.1139/apnm-2012-0261

671 Bing X, Attardo GM, Vigneron A, Aksoy E, Scolari F, Malacrida A, Weiss BL, Aksoy S.
672 2017. Unravelling the relationship between the tsetse fly and its obligate symbiont
673 Wigglesworthia: transcriptomic and metabolomic landscapes reveal highly
674 integrated physiological networks. *Proceedings of the Royal Society B: Biological*
675 *Sciences* **284**:20170360.

676 Boyd BM, Allen JM, Nguyen N-P, Vachaspati P, Quicksall ZS, Warnow T, Mugisha L,
677 Johnson KP, Reed DL. 2017. Primates, Lice and Bacteria: Speciation and Genome
678 Evolution in the Symbionts of Hominid Lice. *Mol Biol Evol* **34**:1743–1757.

679 Brosnan JT, Brosnan ME. 2010. Creatine metabolism and the urea cycle. *Mol Genet*
680 *Metab* **100, Supplement**:S49–S52.

681 Canete M, Hazen MJ, Stockert JC. 1983. NILE BLUE SULFATE STAINING FOR
682 DEMONSTRATION OF LIPIDS IN FLUORESCENCE MICROSCOPY. *Acta*
683 *Histochem Cytochem* **16**:286–288.

Castaño-Moreno E, Castillo V, Peñailillo R, Llanos MN, Valenzuela R, Ronco AM. 2020.
Fatty acid and lipid metabolism in liver of pregnant mice and their offspring is
influenced by unbalanced folates/vitamin B12 diets. *Prostaglandins Leukot Essent*
Fatty Acids **154**:102057.
Chen X, Li S, Aksoy S. 1999. Concordant evolution of a symbiont with its host insect
species: molecular phylogeny of genus Glossina and its bacteriome-associated
endosymbiont, Wigglesworthia glossinidia. *J Mol Evol* **48**:49–58.
Cmelik SHW, Bursell E, Slack E. 1969. Composition of the gut contents of third-instar
tsetse larvae (Glossina morsitans westwood). *Comparative Biochemistry and*
Physiology. doi:10.1016/0010-406x(69)91764-2
da Silva MT, Mujica-Coopman MF, Figueiredo ACC, Hampel D, Vieira LS, Farias DR,
Shahab-Ferdows S, Allen LH, Brito A, Lamers Y, Kac G, S Vaz J. 2020. Maternal
plasma folate concentration is positively associated with serum total cholesterol
and low-density lipoprotein across the three trimesters of pregnancy. *Sci Rep*
10:20141.
da Silva RP, Kelly KB, Al Rajabi A, Jacobs RL. 2014. Novel insights on interactions
between folate and lipid metabolism. *Biofactors* **40**:277–283.
Didion EM, Sabree ZL, Kenyon L, Nine G, Hagan RW, Osman S, Benoit JB. 2021.
Microbiome reduction prevents lipid accumulation during early diapause in the
northern house mosquito, Culex pipiens pipiens. *J Insect Physiol* **134**:104295.
Dixon P. 2003. VEGAN, a package of R functions for community ecology. *J Veg Sci*
14:927–930.
Duron O, Morel O, Noël V, Buysse M, Binetruy F, Lancelot R, Loire E, Ménard C,
Bouchez O, Vavre F, Vial L. 2018. Tick-Bacteria Mutualism Depends on B Vitamin
Synthesis Pathways. *Curr Biol* **28**:1896–1902.e5.
Fatterpaker P, Marfatia U, Sreenivasan A. 1951. Influence of folic acid and vitamin B12
on formation of creatine in vitro and in vivo. *Nature* **167**:1067–1068.
Flatt T. 2011. Survival costs of reproduction in Drosophila. *Exp Gerontol* **46**:369–375.
Fouks B, Harrison MC, Mikhailova AA, Marchal E, English S, Carruthers M, Jennings
EC, Pippel M, Attardo GM, Benoit JB, Bornberg-Bauer E, Tobe SS. 2022. Live-
bearing cockroach genome reveals convergent evolutionary mechanisms linked to
viviparity in insects and beyond. *bioRxiv*. doi:10.1101/2022.02.03.478960
Furness DLF, Yasin N, Dekker GA, Thompson SD, Roberts CT. 2012. Maternal red
blood cell folate concentration at 10–12 weeks gestation and pregnancy outcome. *J*
Matern Fetal Neonatal Med **25**:1423–1427.
Gibellini F, Smith TK. 2010. The Kennedy pathway--De novo synthesis of
phosphatidylethanolamine and phosphatidylcholine. *IUBMB Life* **62**:414–428.
Grönke S, Müller G, Hirsch J, Fellert S, Andreou A, Haase T, Jäckle H, Kühnlein RP.
2007. Dual lipolytic control of body fat storage and mobilization in Drosophila. *PLoS*
Biol **5**:e137.
Hall RJ, Flanagan LA, Bottery MJ, Springthorpe V, Thorpe S, Darby AC, Wood AJ,
Thomas GH. 2019. A Tale of Three Species: Adaptation of Sodalis glossinidius to
Tsetse Biology, Wigglesworthia Metabolism, and Host Diet. *MBio* **10**.
doi:10.1128/mBio.02106-18
Herrera E. 2002. Lipid metabolism in pregnancy and its consequences in the fetus and
newborn. *Endocrine* **19**:43–55.

Herrera E, Lasunción MA, Gomez-Coronado D, Aranda P, López-Luna P, Maier I. 1988.
Role of lipoprotein lipase activity on lipoprotein metabolism and the fate of
circulating triglycerides in pregnancy. *American Journal of Obstetrics and*
Gynecology. doi:10.1016/0002-9378(88)90193-7
Herrera E, Muñoz C, López-Luna P, Ramos P. 1994. Carbohydrate-lipid interactions
during gestation and their control by insulin. *Braz J Med Biol Res* **27**:2499–2519.
Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010. Wolbachia as a
bacteriocyte-associated nutritional mutualist. *Proc Natl Acad Sci U S A* **107**:769–
774.
International Glossina Genome Initiative. 2014. Genome sequence of the tsetse fly
(Glossina morsitans): vector of African trypanosomiasis. *Science* **344**:380–386.
Itabe H, Yamaguchi T, Nimura S, Sasabe N. 2017. Perilipins: a diversity of intracellular
lipid droplet proteins. *Lipids Health Dis* **16**:83.
Jacobs RL, van der Veen JN, Vance DE. 2013. Finding the balance: the role of S-
adenosylmethionine and phosphatidylcholine metabolism in development of
nonalcoholic fatty liver disease. *Hepatology* **58**:1207–1209.
Kazak L, Chouchani ET, Lu GZ, Jedrychowski MP, Bare CJ, Mina AI, Kumari M, Zhang
S, Vuckovic I, Laznik-Bogoslavski D, Dzeja P, Banks AS, Rosen ED, Spiegelman
BM. 2017. Genetic Depletion of Adipocyte Creatine Metabolism Inhibits Diet-
Induced Thermogenesis and Drives Obesity. *Cell Metab* **26**:660–671.e3.
Kirkness EF, Haas BJ, Sun W, Braig HR, Perotti MA, Clark JM, Lee SH, Robertson HM,
Kennedy RC, Elhaik E, Gerlach D, Kriventseva EV, Elsik CG, Graur D, Hill CA,
Veenstra JA, Walenz B, Tubío JMC, Ribeiro JMC, Rozas J, Johnston JS, Reese
JT, Popadic A, Tojo M, Raoult D, Reed DL, Tomoyasu Y, Kraus E, Mittapalli O,
Margam VM, Li H-M, Meyer JM, Johnson RM, Romero-Severson J, Vanzee JP,
Alvarez-Ponce D, Vieira FG, Aguadé M, Guirao-Rico S, Anzola JM, Yoon KS,
Strycharz JP, Unger MF, Christley S, Lobo NF, Seufferheld MJ, Wang N, Dasch
GA, Struchiner CJ, Madey G, Hannick LI, Bidwell S, Joardar V, Caler E, Shao R,
Barker SC, Cameron S, Bruggner RV, Regier A, Johnson J, Viswanathan L,
Utterback TR, Sutton GG, Lawson D, Waterhouse RM, Venter JC, Strausberg RL,
Berenbaum MR, Collins FH, Zdobnov EM, Pittendrigh BR. 2010. Genome
sequences of the human body louse and its primary endosymbiont provide insights
into the permanent parasitic lifestyle. *Proc Natl Acad Sci U S A* **107**:12168–12173.
Kitagawa E, Yamamoto T, Yamamoto K, Nakagawa T, Hayakawa T. 2015.
Accumulation of lipid in rat liver was induced by vitamin B6 deficiency and was
ameliorated by supplemental phosphatidylcholine in the diet. *Biosci Biotechnol*
Biochem **79**:1320–1326.
Knopp RH, Bonet B, Zhu X. 1998. Lipid Metabolism in Pregnancy In: Cowett RM, editor.
Principles of Perinatal—Neonatal Metabolism. New York, NY: Springer New York.
pp. 221–258.
Krahmer N, Guo Y, Wilfling F, Hilger M, Lingrell S, Heger K, Newman HW, Schmidt-
Supprian M, Vance DE, Mann M, Farese RV Jr, Walther TC. 2011.
Phosphatidylcholine synthesis for lipid droplet expansion is mediated by localized
activation of CTP:phosphocholine cytidylyltransferase. *Cell Metab* **14**:504–515.
Listenberger L, Townsend E, Rickertsen C, Hains A, Brown E, Inwards EG, Stoeckman
AK, Matis MP, Sampathkumar RS, Osna NA, Kharbanda KK. 2018. Decreasing

Phosphatidylcholine on the Surface of the Lipid Droplet Correlates with Altered
Protein Binding and Steatosis. *Cells* **7**. doi:10.3390/cells7120230

Loenen WA. 2006. S-adenosylmethionine: jack of all trades and master of everything?
Biochem Soc Trans **34**:330–333.

Mazen Noureddin, José M. Mato, Shelly C. Lu. 2015. Nonalcoholic fatty liver disease:
update on pathogenesis, diagnosis, treatment and the role of S-
adenosylmethionine. *Exp Biol Med*. doi:10.1177/1535370215579161

McNeil CJ, Hay SM, Rucklidge GJ, Reid MD, Duncan GJ, Rees WD. 2009. Maternal
diets deficient in folic acid and related methyl donors modify mechanisms
associated with lipid metabolism in the fetal liver of the rat. *British Journal of*
Nutrition. doi:10.1017/s0007114509990389

Michalkova V, Benoit JB, Attardo GM, Medlock J, Aksoy S. 2014a. Amelioration of
reproduction-associated oxidative stress in a viviparous insect is critical to prevent
reproductive senescence. *PLoS One* **9**:e87554.

Michalkova V, Benoit JB, Weiss BL, Attardo GM, Aksoy S. 2014b. Vitamin B6 generated
by obligate symbionts is critical for maintaining proline homeostasis and fecundity
in tsetse flies. *Appl Environ Microbiol* **80**:5844–5853.

Moloo SK. 1971. An artificial feeding technique for Glossina. *Parasitology* **63**:507–512.

Moullan N, Mouchiroud L, Wang X, Ryu D, Williams EG, Mottis A, Jovaisaite V,
Frochaux MV, Quiros PM, Deplancke B, Houtkooper RH, Auwerx J. 2015.
Tetracyclines Disturb Mitochondrial Function across Eukaryotic Models: A Call for
Caution in Biomedical Research. *Cell Rep* **10**:1681–1691.

Nguyen TT, Hayakawa T, Tsuge H. 2001. Effect of vitamin B6 deficiency on the
synthesis and accumulation of S-adenosylhomocysteine and S-adenosylmethionine
in rat tissues. *J Nutr Sci Vitaminol* **47**:188–194.

Nikoh N, Hosokawa T, Moriyama M, Oshima K, Hattori M, Fukatsu T. 2014.
Evolutionary origin of insect-Wolbachia nutritional mutualism. *Proceedings of the*
National Academy of Sciences. doi:10.1073/pnas.1409284111

Nogge G. 1976. Sterility in tsetse flies (Glossina morsitans Westwood) caused by loss
of symbionts. *Experientia* **32**:995–996.

Oksanen J. 2007. vegan : Community Ecology Package. R package version 1.8-5.
http://www.cran.r-project.org.

Olzmann JA, Carvalho P. 2019. Dynamics and functions of lipid droplets. *Nat Rev Mol*
Cell Biol **20**:137–155.

Pais R, Lohs C, Wu Y, Wang J, Aksoy S. 2008. The obligate mutualist Wigglesworthia
glossinidia influences reproduction, digestion, and immunity processes of its host,
the tsetse fly. *Appl Environ Microbiol* **74**:5965–5974.

Pang Z, Chong J, Zhou G, de Lima Morais DA, Chang L, Barrette M, Gauthier C,
Jacques P-É, Li S, Xia J. 2021. MetaboAnalyst 5.0: narrowing the gap between raw
spectra and functional insights. *Nucleic Acids Res* **49**:W388–W396.

Perrin N, Sibly RM. 1993. Dynamic models of energy allocation and investment. *Annu*
Rev Ecol Syst **24**:379–410.

Pol A, Gross SP, Parton RG. 2014. Biogenesis of the multifunctional lipid droplet:
Lipids, proteins, and sites. *J Cell Biol* **204**:635–646.

Rio RVM, Jozwick AKS, Savage AF, Sabet A, Vigneron A, Wu Y, Aksoy S, Weiss BL.
2019. Mutualist-Provisioned Resources Impact Vector Competency. *MBio* **10**.

doi:10.1128/mBio.00018-19

Romoli O, Schönbeck JC, Hapfelmeier S, Gendrin M. 2021. Production of germ-free
mosquitoes via transient colonisation allows stage-specific investigation of host–
microbiota interactions. *Nat Commun* **12**:1–16.

Rosendale AJ, Dunlevy ME, McCue MD, Benoit JB. 2019. Progressive behavioural,
physiological and transcriptomic shifts over the course of prolonged starvation in
ticks. *Mol Ecol* **28**:49–65.

Ryckman KK, Spracklen CN, Smith CJ, Robinson JG, Saftlas AF. 2015. Maternal lipid
levels during pregnancy and gestational diabetes: a systematic review and meta-
analysis. *BJOG: An International Journal of Obstetrics & Gynaecology*.
doi:10.1111/1471-0528.13261

Salem H, Bauer E, Strauss AS, Vogel H, Marz M, Kaltenpoth M. 2014. Vitamin
supplementation by gut symbionts ensures metabolic homeostasis in an insect
host. *Proceedings of the Royal Society B: Biological Sciences*.
doi:10.1098/rspb.2014.1838

Snyder AK, Rio RVM. 2015. "Wigglesworthia morsitans" Folate (Vitamin B9)
Biosynthesis Contributes to Tsetse Host Fitness. *Appl Environ Microbiol* **81**:5375–
5386.

Strickler-Dinglasan PM, Guz N, Attardo G, Aksoy S. 2006. Molecular characterization of
iron binding proteins from Glossina morsitans morsitans (Diptera: Glossinidae).
Insect Biochemistry and Molecular Biology. doi:10.1016/j.ibmb.2006.09.003

Sztalryd C, Brasaemle DL. 2017. The perilipin family of lipid droplet proteins:
Gatekeepers of intracellular lipolysis. *Biochim Biophys Acta Mol Cell Biol Lipids*
1862:1221–1232.

Thomas GWC, Dohmen E, Hughes DST, Murali SC, Poelchau M, Glastad K, Anstead
CA, Ayoub NA, Batterham P, Bellair M, Others. 2020. Gene content evolution in the
arthropods. *Genome Biol* **21**:1–14.

Tobe SS. 1978. Reproductive physiology of Glossina. *Annu Rev Entomol* **23**:283–307.

Valzania L, Coon KL, Vogel KJ, Brown MR, Strand MR. 2018. Hypoxia-induced
transcription factor signaling is essential for larval growth of the mosquito Aedes
aegypti. *Proc Natl Acad Sci U S A* **115**:457–465.

Van Handel E. 1985. Rapid determination of glycogen and sugars in mosquitoes. *J Am*
Mosq Control Assoc **1**:299–301.

Ville Männistö, Dorota Kaminska, Vesa Kärjä, Mika Tiainen, Vanessa D. de Mello, Kati
Hanhineva, Pasi Soininen, Mika Ala-Korpela, Jussi Pihlajamäki. 2019. Total liver
phosphatidylcholine content associates with non-alcoholic steatohepatitis and
glycine N-methyltransferase expression. *Liver Int*. doi:10.1111/liv.14174

Wang Y, Eum JH, Harrison RE, Valzania L, Yang X, Johnson JA, Huck DT, Brown MR,
Strand MR. 2021. Riboflavin instability is a key factor underlying the requirement of
a gut microbiota for mosquito development. *Proc Natl Acad Sci U S A* **118**.
doi:10.1073/pnas.2101080118

Yang G, Attardo GM, Lohs C, Aksoy S. 2010. Molecular characterization of two novel
milk proteins in the tsetse fly (Glossina morsitans morsitans). *Insect Mol Biol*
19:253–262.

Figure 1

Figure 2

A.

Kennedy pathway

Choline

Choline kinase

Phosphocholine

CTP: Phosphocholine
Cytidylyltransferase (CCT)

Cytidine-diphosphocholine

DAG:CDP-Choline
Cholinephosphotransferase

Phosphatidylcholine

RNA-seq results from lactating and dry flies

cho

cct2

cct1 *

cpt1

cpt2

cpt3

Relative expression level

B.

Oogenesis — — → Dry (non-lactating)

Embryogenesis - - → Lactating

Larvigenesis

mgp1

ctt1

Relative transcript levels

Time after adult emergence (d)

Right ovary
Uterus
Left ovary

Adult emergence

Parturiton

Lactation state

C.

Relative transcript levels

Before lactation

During lactation

Milk gland

Fat body

Milk gland

FB/MG   Ov   MT   RT   B   MG   HG   SG

Figure 5

Lipid Metabolism Dysfunction in Aposymbiotic Tsetse Flies

Symbiotic

Aposymbiotic

The Kennedy Pathway
Phospholipid biosynthesis