Oxidative reactions catalyzed by hydrogen peroxide produced by *Streptococcus pneumoniae* and

other Streptococci Cause the Release and Degradation of Heme from Hemoglobin

**Babek Alibayov[1], Anna Scasny[1], Faidad Khan[1], Aidan Creel[1,2], Perriann Smith[1,3], Ana G. Jop Vidal[1],**

**Fa'alataitaua M. Fitisemanu[4], Teresita Padilla-Benavides[4], Jeffrey Weiser[5], and Jorge E. Vidal[1,#],**

[1]Department of Cell and Molecular Biology, School of Medicine, and [2]Summer Undergraduate

Research Experience Program (SURE), School of Graduate Studies in the Health Sciences, University

of Mississippi Medical Center, Jackson MS; [3]Mississippi INBRE Research Scholar, University of

Southern Mississippi, Hattiesburg MS; [4]Department of Molecular Biology and Biochemistry,

Wesleyan University, Middletown CT; [5]Department of Microbiology, NYU Langone Health, New

York NY,

Running title: Spn-$H_2O_2$ oxidizes hemoglobin degrading heme

[#]Address correspondence to:

Jorge E. Vidal, PhD

2500 North State Street, Jackson Mississippi 39216

Phone: (601) 984-1711, Fax: (601) 984-1501

Email: jvidal@umc.edu

**Abstract**

Streptococcus pneumoniae (Spn) strains cause pneumonia that kills millions every year worldwide. Spn produces Ply, a hemolysin that lyses erythrocytes releasing hemoglobin and also produces the pro-oxidant hydrogen peroxide (Spn-$H_2O_2$) during growth. The hallmark of the pathophysiology of hemolytic diseases is the oxidation of hemoglobin but oxidative reactions catalyzed by Spn-$H_2O_2$ has been poorly studied. We characterized the oxidation of hemoglobin by Spn-$H_2O_2$. We prepared a series of single ($\Delta spxB$, or $\Delta lctO$), double mutant ($\Delta spxB\Delta lctO$) and complemented strains in TIGR4, D39 and EF3030. We then utilized an *in vitro* model with oxy-hemoglobin to demonstrate that oxy-hemoglobin was oxidized rapidly, within 30 min of incubation, by Spn-$H_2O_2$ to met-hemoglobin and that the main source of Spn-$H_2O_2$ was pyruvate oxidase (SpxB). Moreover, extended incubation caused the release and the degradation of heme. We then assessed oxidation of hemoglobin and heme degradation by other bacteria inhabitants of the respiratory tract. All hydrogen peroxide-producing streptococci tested caused the oxidation of hemoglobin and heme degradation whereas those bacterial species that produce <1 µM $H_2O_2$, neither oxidized hemoglobin nor degraded heme. An *ex vivo* bacteremia model confirmed that oxidation of hemoglobin and heme degradation occurred concurrently with hemoglobin that was released from erythrocytes by Ply. Finally, gene expression studies demonstrated that heme, but not red blood cells or hemoglobin induced an upregulated transcription of the *spxB* gene. Oxidation of hemoglobin may be important for pathogenesis and for the symbiosis of hydrogen peroxide-producing bacteria with other species by providing nutrients such as iron.

**Introduction.**

Streptococcus pneumoniae (Spn) colonizes the lower respiratory epithelium, causing pneumonia that can progress to invasive pneumococcal disease (IPD)(1-4). Spn mainly affects the most vulnerable populations: young children, immunocompromised patients, and the elderly (1, 4, 5). Globally, Spn causes ~15 million cases of illness each year, leading to more than a million deaths due to pneumonia (6-8). Pneumonia is an acute infection of the pulmonary parenchyma that presents with pulmonary hemorrhage, inflammatory congestion, hepatization, suppurative infiltration and lung parenchymal injury (4, 9-13). If left untreated, Spn invades the bloodstream, producing septic shock, multiorgan failure, cardiotoxicity, and death within days from onset (4, 9-12, 14, 15).

Spn strains produce the toxin pneumolysin (Ply) that has cytotoxic activity against lung epithelial and endothelial cells (16-18). Ply is a 53-kDa surface located protein (19, 20) that can also be released through autolysis (21, 22). Besides its toxicity against cells, Ply has hemolytic activity against erythrocytes of several mammalian species and hemolysis is observed whether Ply is released into the supernatant or located on the bacterial membrane (20). Under standard culture conditions, i.e., growing bacteria at 37°C in a 5% $CO_2$ atmosphere, Ply is responsible for almost all hemolytic activity in cultures of Spn strains (11), causing the release of a large amount of hemoglobin into the supernatant (20, 21, 23). Transcription of the ply gene encoding this pneumolysin is upregulated when Spn is cultured under conditions mimicking lung infection (24) and therefore, Ply-induced release of hemoglobin is expected to occur during pneumococcal lung infection.

Hemoglobin is a tetrameric protein (64 kDa) made of four subunits, two alpha and two beta chains. Each subunit contains a heme center that reversibly binds oxygen through a penta-

coordinate heme molecule containing ferrous iron ($Fe^{+2}$), known as oxy-hemoglobin (25, 26). When

hemoglobin is released from erythrocytes, heme-hemoglobin can be observed by spectroscopy at

~415 nm (25, 27, 28). This region is known as the Soret region peak and represents heme-

hemoglobin, while the alpha and beta chains are characterized by two absorption peaks of ~540

and ~570 nm (27, 28). Cell-free heme is toxic to cells because of its potential to generate toxic

radicals (29).

Release of hemoglobin during hemolytic disease such as sickle cell disease and other

hemorrhagic conditions, causes oxidation of hemoglobin (30). Oxidized hemoglobin releases heme,

which in turn induces adverse effects including leukocyte activation, cytokine up-regulation, and

production of oxidants leading to lung epithelial injury and damage to the vascular system (29-31).

Reactive oxygen species (ROS), such as nitric oxide (NO) (32) hypochlorous acid conjugate base

(OCl-)(24), and $H_2O_2$ (33, 34), drive a catalytic cycle including the oxidation of oxy-hemoglobin to

ferryl hemoglobin (Hb-$Fe^{+4}$), and auto - reduction of the ferryl intermediate to met-hemoglobin (Hb-

$Fe^{+3}$) that is detected by spectroscopy at 405 nm (35, 36). Additional pro-oxidant molecules, such as

$H_2O_2$, produce more ferryl hemoglobin and cause heme degradation (28, 36). Thus, the oxidation of

hemoglobin results in highly oxidizing ferryl hemoglobin (Hb-$Fe^{4+}$), met-hemoglobin (Hb-$Fe^{3+}$) and

the release of toxic heme that disrupts the mitochondria membrane potential of lung epithelial cells

and endothelial cells, and caused lipid peroxidation (28, 30, 36-38).

Because Spn causes hemorrhage in the lung and produces a large amount of highly reactive

$H_2O_2$ (39, 40), oxidation of cell-free hemoglobin might occur during pneumococcal disease. $H_2O_2$ is a

by-product of the oxidation of pyruvate, end product of the glycolysis, produced by the enzyme

pyruvate oxidase (SpxB) (41, 42). This reaction consumes oxygen and produces $H_2O_2$. In the absence

of oxygen, pyruvate is converted to L-lactate while in the presence of molecular $O_2$, L-lactate feeds

back to pyruvate, via lactate oxidase (LctO), in a reaction that also produces $H_2O_2$ (41, 42). It has

been calculated that ~85% of $H_2O_2$ released in cultures of Spn occurs through the reaction catalyzed

by SpxB, whereas the remaining ~15% is attributed to LctO (43, 44). Experiments using animal

models have demonstrated that Spn-$H_2O_2$ plays a role in lung colonization, and in the translocation

of Spn from the lungs to the bloodstream (45-48). Details of the mechanism(s) are largely unknown.

We recently discovered that Spn-$H_2O_2$ oxidizes hemoglobin to met-hemoglobin, a non-oxygen

binding form of hemoglobin (23). We also demonstrated that the oxidation of hemoglobin causes

the characteristic greenish halo around pneumococcal colonies grown on blood agar plates, known

as alpha hemolysis (23). Thus, the phenotype known as alpha hemolysis is not hemolysis but the

oxidation of hemoglobin to met-hemoglobin (23). Another alpha-hemolytic streptococci, *S. gordonii*,

oxidizes hemoglobin to met-hemoglobin through catalytic reactions driven by $H_2O_2$ (49).

Oxidation of cell-free hemoglobin induces lung injury in patients with acute respiratory distress

syndrome (31), and in children with SCD, oxidation of hemoglobin (Hb)S to Hb-$Fe^{4+}$ and then its

reduction to met-hemoglobin results in an acute hemolytic vascular inflammatory process causing

acute lung injury (30, 50). We hypothesize that the oxidation of hemoglobin during pneumococcal

disease can exacerbates cytotoxicity in the lungs but details of these oxidative reactions in cultures

of Spn strains are not available. Therefore, in this study we comprehensively investigated the

oxidation of hemoglobin by Spn cultures. We discovered that the oxidative capacity of Spn cultures,

under the culture conditions utilized here, was attributed to $H_2O_2$ produced through catalytic

reactions of SpxB but not to that of LctO. We also found that Spn-$H_2O_2$ caused the degradation of

heme and that a number of $H_2O_2$-producing, alpha-hemolytic, Streptococci oxidized hemoglobin and

degraded heme from hemoglobin. An *ex vivo* model of pneumococcal bacteremia further

demonstrated that oxidation of hemoglobin and heme degradation occurred rapidly and that heme,

but not hemoglobin or red blood cells, caused an upregulated transcription of *spxB*. Oxidative

reactions driven by Spn-$H_2O_2$ might contribute to the pathogenesis of pneumococcal disease and

can also be important for the symbiosis with other non-hydrogen peroxide producing bacteria by

providing nutrients such as heme-iron.

**Results**

***Streptococcus pneumoniae*-produced hydrogen peroxide oxidizes heme-hemoglobin, causing the**

**loss of the Soret peak.** We demonstrated that the alpha and beta chains of oxy-hemoglobin are

deoxygenated by *S. pneumoniae* strains but further oxidation of heme-$Fe^{+2}$ was not investigated

(23). To assess this, Todd-Hewitt broth (THY) supplemented with sheep oxy-hemoglobin (oxyHb),

which contains heme-$Fe^{2+}$, was inoculated with TIGR4 and incubated at 37°C in a 5% $CO_2$ and ~20%

$O_2$ atmosphere. As a control, uninfected THY-oxyHb was incubated under the same conditions. After

30 min of incubation, TIGR4 cultures caused the shift of the Soret curve from heme-$Fe^{2+}$, that peaks

at 415 nm, to a Soret curve of heme-$Fe^{3+}$ at 405 nm (Fig. 1A). As expected, this hemoglobin

preparation was deoxygenated producing met-hemoglobin (Fig. 1B). Similar oxidation of heme-$Fe^{2+}$

to heme-$Fe^{3+}$ was observed when oxy-hemoglobin was purified from horse erythrocytes and

incubated with Spn strains TIGR4, or D39 (not shown). Cultures of THY-oxyHb inoculated with

hydrogen peroxide-deficient strain TIGR4Δ*spxB*Δ*lctO,* or D39Δ*spxB*Δ*lctO*, did not cause the shift of

the Soret curve (Fig. 1A and not shown). Hemoglobin in uninfected cultures was not auto-oxidized

after this 30 min incubation period (Fig. 1A).

Extended incubation of TIGR4 in THY-oxyHb revealed that the Soret curve was lost in a time-

dependent manner (Fig. 1C). To assess if Spn-$H_2O_2$ had caused the loss of the Soret peak, we

inoculated THY-oxyHb cultures with TIGR4Δ*spxB*Δ*lctO* and the culture was incubated for 6 h. A Soret

curve was observed in this culture but the heme-$Fe^{2+}$ peak (415 nm), observed after 30 min post-

inoculation (Fig. 1A), had shifted to 405 nm representing heme-$Fe^{3+}$ (Fig. 1D). The deoxygenation of

hemoglobin in THY-oxyHb cultures of TIGR4$\Delta$spxB$\Delta$lctO was, however, due to autoxidation because

uninfected cultures incubated under the same conditions at 37°C, and with a 5% $CO_2$ and ~20% $O_2$

atmosphere, had shifted to heme-$Fe^{3+}$ as well (Fig. 1C).

To further confirm that the loss of the Soret peak was due to Spn-$H_2O_2$, we prepared a

complemented strain by cloning back spxB and lctO ($\Omega$spxB$\Omega$lctO) between SP_1113 and SP_1114

(Fig. 2). We also complemented TIGR4$\Delta$spxB with a single copy of spxB to be utilized later (Fig. 2).

This chromosomal location is unaffected by exogenous regulatory elements, such that

complementing genes were controlled by their own promoters and insertion of genes within

SP_1113 and SP_1114 did not affect virulence (51-54). Compared to wt and TIGR4$\Delta$spxB$\Delta$lctO,

complemented strains showed a slight but not significant growth delay (Fig. 2B); however $H_2O_2$

production was restored to wt levels, as assessed with the Prussian blue-forming reaction and the

Amplex Red kit (Fig. 2C and 2D) (55). In THY-oxyHb cultures of $\Omega$spxB$\Omega$lctO incubated for 6 h, the

Soret curve had flattened (Fig. 1D). These results suggest that Spn-$H_2O_2$ causes heme degradation.

Given that heme-$Fe^{2+}$ in THY-oxyHb cultures was oxidized to heme-$Fe^{2+}$ (Fig. 1A) prior to

losing the Soret curve (Fig. 1C), we next assessed if the oxidation state of heme was important for

the subsequent loss of the Soret curve. To investigate this, we supplemented THY with deoxy-

hemoglobin (i.e., met-hemoglobin) containing heme-$Fe^{3+}$ (THY-Hb) and the culture was incubated

with Spn. A similar loss of the Soret curve was observed with TIGR4 wt and $\Omega$spxB$\Omega$lctO (Fig. 1E),

but an intact curve was obtained in cultures of TIGR4$\Delta$spxB$\Delta$lctO (Fig. 1E), indicating that heme-$Fe^{3+}$

was further degraded. These results demonstrated that Spn-$H_2O_2$ converts oxyHb to metHb.

**Exogenous hydrogen peroxide supplementation at levels produced in *S. pneumoniae* cultures recapitulates hemoglobin oxidation.** We then assessed whether a similar concentration of hydrogen peroxide to that observed in wt cultures of Spn (<1 mM) [Fig. 2D and references (39, 40)] can alter the heme-$Fe^{3+}$ curve. For easy comparison of the multiple concentrations utilized in this experiment, we plotted the maximum absorbance obtained by spectroscopy for THY-Hb (405 nm) and we compared against that of THY-Hb treated with an increasing concentration of hydrogen peroxide. As shown in Fig. 1F, 10 µM hemoglobin in THY-Hb peaks at an absorbance of ~0.8. After 4 h of incubation with hydrogen peroxide, the absorbance was significantly reduced with a concentration of 88 µM and above, including the concentration of hydrogen peroxide observed in Spn cultures (~880 µM) (Fig. 1F). Altogether, these results indicate that the heme moiety in oxidized hemoglobin is degraded by Spn-$H_2O_2$.

**Oxidation of hemoglobin in *S. pneumoniae* cultures is mainly caused by hydrogen peroxide catalyzed through pyruvate oxidase SpxB.** Under aerobic conditions, hydrogen peroxide in pneumococcal cultures is produced through catalytic reactions of pyruvate oxidase (SpxB) and lactate oxidase (LctO) (43, 44). To assess the contribution of each enzyme, single Δ*spxB* or Δ*lctO* prepared in the TIGR4, or EF3030, background were assessed in THY-Hb cultures. In cultures of wt strains TIGR4, and EF3030, isogenic mutant TIGR4Δ*lctO*, and EF3030Δ*lctO*, Ω*spxB*Ω*lctO* and TIGR4Δ*spxB*Ω*spxB* complemented strain the Soret Heme-Hb$^{3+}$ curve was flattened, causing a statistically significant reduction of the 405 nm peak (Fig. 3A). In contrast, in THY-Hb cultures of TIGR4Δ*spxB* or EF3030Δ*spxB,* the Soret curve remained intact, similar to THY-Hb cultures inoculated with the corresponding Δ*spxB*Δ*lctO* mutant (Fig. 3A and 3B). Additional D39Δ*spxB*Δ*lctO*, or D39Δ*spxB* strain (P878), hydrogen peroxide-deficient mutants did not alter the Soret curve while

D39Δ*spxB*/*spxB* complemented strain, P1221, caused the loss of the Soret curve (Fig. 3C). These

experiments demonstrated that oxidation of hemoglobin in cultures of Spn is mainly caused by

hydrogen peroxide sourced from reactions catalyzed by SpxB.

**Spn-produced $H_2O_2$ degrades heme-$Fe^{3+}$.** Heme degradation occurs through enzymatic reactions or

by nonselective destruction of heme double bonds caused by reactive oxygen species, such as $H_2O_2$

(35, 56). To investigate whether Spn-$H_2O_2$ caused degradation of heme, THY-Hb medium was

inoculated with TIGR4, EF3030, their hydrogen peroxide mutant derivatives, or complemented

strains and heme was assessed in supernatants using an in-gel heme assay (57, 58). Signal of heme,

heme-bound to hemoglobin monomers or dimers, was detected in uninfected THY-Hb cultures

incubated for 6 h (Fig. 4A). Heme signal in the supernatant of THY-Hb cultures incubated with TIGR4

for 2 h was similar to that of the uninoculated control but it decreased after 4 h post-inoculation

and was not detected 6 h post-inoculation (Fig. 4A). In THY-Hb cultures of TIGR4, TIGR4Δ*lctO*,

TIGR4Ω*spxB*Ω*lctO*, TIGR4Δ*spxB*Ω*spxB,* EF3030, or EF3030Δ*lctO* the signal of heme had already

disappeared after 6 h of incubation (Fig. 4B and 4C). In contrast, in THY-Hb cultures of

TIGR4Δ*spxB*Δ*lctO,* TIGR4Δ*spxB*, EF3030Δ*spxB*Δ*lctO*, or EF3030Δ*spxB*, signal of free heme, and heme-

bound to hemoglobin was detected, thus confirming that degradation of heme had occurred in

these cultures through hydrogen peroxide produced by SpxB.

As a loading control and to additionally investigate if hemoglobin had been degraded,

supernatants were probed by Western blot with a polyclonal anti-hemoglobin antibody. Our results

revealed a similar concentration of hemoglobin in those supernatants, confirming that the apo-

protein was not degraded after the 2, 4 or 6 h incubation period with Spn (Fig. 4A, bottom panel). As

expected since heme was degraded by Spn-$H_2O_2$, the intracellular concentration of iron was 79.19,

18.21, or 108.79 µM of Fe/µg of protein in TIGR4, TIGR4$\Delta$*spxB*$\Delta$*lctO* or TIGR4$\Omega$*spxB*$\Omega$*lctO*,

respectively, confirming that iron was released and taken by wt and complemented strain.

Altogether, these experiments demonstrated that Spn-$H_2O_2$ degrades free-heme and heme bound

to hemoglobin.

**Streptococci that produces hydrogen peroxide, but not other respiratory bacteria, causes the**

**degradation of heme.** Non-encapsulated Spn (NESpn) and other alpha-hemolytic streptococci

produce hydrogen peroxide (59, 60). We therefore assessed a collection of NESpn and oral

streptococci for heme degradation. We included *S. mutans* as an additional control since strains lack

production of $H_2O_2$ (61, 62). As shown in Fig. 5, degradation of heme-$Fe^{3+}$ was observed in THY-Hb

cultures of NESpn strains MNZ41, MNZ67 and *S. salivarius*, *S. mitis*, *S. pseudopneumonanie* and *S.*

*oralis* but not in cultures of *S. mutans* (Fig. 5A). We also tested cultures of other bacterial species

that cohabit the upper airways along with oral streptococci, such as *Haemophilus influenzae*,

*Staphylococcus aureus* and assessed cultures of *Klebsiella pneumoniae*. The incubation of THY-Hb

with any of these bacterial species did not cause heme degradation (Fig. 5A). Streptococcal species

that caused heme degradation produced hydrogen peroxide, while those unable to affect the heme-

hemoglobin Soret curve did not (Fig. 5B). Altogether, these experiments demonstrated that Spn and

other alpha hemolytic streptococci that produce abundant hydrogen peroxide, oxidizes hemoglobin

causing the degradation of heme.

**Heme is oxidized and degraded in an *ex-vivo* model of pneumococcal bacteremia.** During the

course of pneumococcal bacteremia erythrocytes are lysed by the pneumolysin (Ply), released

through autolysis (63). Ply is also located on the bacterial membrane where it lyses red blood cells

through contact (19, 20). Abundant Spn-$H_2O_2$ is produce as early as 2 h post-inoculation (Fig. 2D)

thereby we sought to assess whether oxidation of hemoglobin and heme degradation occur in an

*ex-vivo* model of pneumococcal bacteremia (Fig. 6A). To assess this, we supplemented THY broth

with a suspension of red blood cells (THY-RBC), and this medium was infected with TIGR4,

TIGR4Δ*ply*, or TIGR4Δ*spxB*Δ*lctO*. The oxidation and concentration of hemoglobin as well as heme

degradation were investigated.

As expected, the Soret curve was not observed in supernatants from non-lysed THY-RBC

control and it was also absent in THY-RBC supernatants infected with TIGR4Δ*ply,* because it does

not lyse erythrocytes (Fig. 6B). Given that RBCs contains oxy-hemoglobin (Hb-$Fe^{2+}$), the Soret curve

of the positive control, where RBC had been lysed with saponin, peaked at 415 nm with an average

absorbance of 1.38 (Fig. 6B). The concentration of cell-free hemoglobin in this control supernatant

was 17.35 µM as measured by the Quantichrom assay (Table 1). The Soret curve from THY-RBC

cultures infected with TIGR4 peaked at 405 nm with an average absorbance of 1.39 (Fig. 6B) and

concentration of hemoglobin of 12.8 µM (Table 1). These results indicate that maximum TIGR4-

induced release of hemoglobin is achieved within 30 min of incubation, compared with the saponin-

induced released control, and that the oxidation of heme-$Fe^{2+}$ to heme-$Fe^{3+}$ had already occurred

(Fig. 6B).  Similar to the wt and because TIGR4Δ*spxB*Δ*lctO* produces Ply (21, 23), a Soret curve was

observed in THY-RBC cultures infected with this hydrogen peroxide deficient mutant as soon as 30

min post-inoculation (Fig. 6B). This Soret curve peaked at 415 nm with an average absorbance of

1.21 and a concentration of hemoglobin of 16.4 µM (Table 1), indicating that most hemoglobin had

been released but, unlike the wt strain, oxidation of heme-$Fe^{2+}$ did not occur because of the lack of

hydrogen peroxide. Accordingly, oxidation of hemoglobin did not occur in TIGR4 cultures treated

with catalase but it was oxidized to met-hemoglobin in TIGR4 cultures treated with protease

inhibitors (Fig. 6B).

After 6 h of incubation, autoxidation of uninoculated, saponin-released hemoglobin in THY-

RBC cultures (Fig. 6C, Hb-$Fe^{3+}$) and those infected with TIGR4 treated with catalase (Fig. 6C) were

oxidized. THY-RBC medium infected with TIGR4 had already flattened the heme-$Fe^{3+}$ curve and

therefore heme had been degraded but not in those of TIGR4 treated with catalase (Fig. 6C).

Degradation of heme was not inhibited by supplementing THY-RBC before the infection with

protease inhibitors (Fig. 6C). We next treated saponin-released oxy-hemoglobin with increasing

amounts of $H_2O_2$ and the oxidation to met-hemoglobin and then degradation of heme was

recapitulated (Fig. 6D). Altogether, our results indicate that oxidation of hemoglobin and

degradation of heme by Spn-$H_2O_2$ occur during pneumococcal bacteremia.

To further mimic bacteremia, we added THY with 10% of human serum and supplemented

this broth culture with 10 µM met-hemoglobin. In addition, non-diluted human serum was

supplemented with 10 µM met-hemoglobin. These hemoglobin-containing serum cultures were

infected with TIGR4 and incubated at different times. A time course degradation of met-hemoglobin

was observed in both culture conditions (Fig. 6E and 6F), confirming that hemoglobin is oxidized and

degraded when Spn growths in human serum.

Because the release of hemoglobin and its oxidation was observed early during the

simulated bacteremia (i.e., 30 min), to gain insights into the mechanism leading to hemoglobin

oxidation and heme degradation, we investigated expression of *ply*, *spxB*, and *lctO* during

incubation of Spn in THY-RBC, THY-Hb or in THY supplemented with heme. Compared to THY control

cultures, the expression of *ply, spxB,* and *lctO* did not significantly change when TIGR4 grew in broth

cultures added with RBCs or hemoglobin. However, transcription of *spxB* was upregulated when

TIGR4 was incubated in THY containing heme, whereas the transcription *lctO* and *ply* was

downregulated (Fig. 7). Together, these experiments indicate that under these culture conditions

heme was the inducer for Spn-$H_2O_2$ production.

**Discussion**

We have demonstrated in this study that $H_2O_2$ produced by the pneumococcus, Spn-$H_2O_2$,

and other alpha-hemolytic Streptococci, caused the oxidation of hemoglobin to met-hemoglobin

with a subsequent degradation of heme. The current study also demonstrated that the oxidation of

hemoglobin and heme degradation occurred in a simulated bacteremia infection suggesting that

these oxidative reactions may occur during pneumococcal disease.

*In vitro* oxidation of hemoglobin with hydrogen peroxide produces ferryl hemoglobin [Hb-

$Fe^{4+}$], a highly reactive form of hemoglobin that gets reduced to met-hemoglobin (64). With an

increased presence of hydrogen peroxide molecules, met-hemoglobin gets oxidized back to Hb-$Fe^{4+}$

, which in turn causes heme degradation (65).  Iron stored in heme-hemoglobin is poorly reactive

but once freed up, as a consequence of heme degradation, free iron can react with hydrogen

peroxide to produce another highly reactive hydroxyl radical (66, 67).  Indeed, we have

demonstrated that free iron present in THY medium is enough to generate hydroxyl radicals ($^{\bullet}$OH)

(40), killing *S. aureus* strains rapidly (39, 68, 69). The increase of iron in the culture medium due to

heme degradation should result in an increased generation of $^{\bullet}$OH  radicals. The potential presence

of two highly reactive radicals, Hb-$Fe^{4+}$ and $^{\bullet}$OH, for the cytotoxicity seen during pneumococcal

disease is being investigated in our laboratories.

It has been investigated that nearly 3% of oxy-hemoglobin is auto-oxidized inside

erythrocytes, and this auto-oxidation is caused by $2\times10^{-7}$ mM of hydrogen peroxide generated in the

298 cytosol of red blood cells (70). Although auto-oxidation by $CO_2$ occurred in our *in vitro* system 4 h

299 post-inoculation (i.e., auto-oxidation was not observed when incubated at 37°C with no $CO_2$), Spn

300 cultures produce $10^7$-fold more (~1 mM) hydrogen peroxide within a few hours of incubation and

301 caused the rapid oxidation of hemoglobin leading to the release of heme, a toxic molecule, with its

302 subsequent degradation (65). We demonstrated that heme degradation caused an increased on

303 free-iron in the medium that was taken by pneumococci. These novel observations might have

304 important implications for pathogenesis of hydrogen peroxide-producing Streptococci but also for

305 the symbiosis observed in bacterial communities where some species can benefit from heme and

306 iron release by hydrogen peroxide-producing bacteria.

308   In children with SCD, hemoglobin (Hb)S is oxidized to Hb-$Fe^{4+}$ then reduced to met-

309 hemoglobin and resulting in an acute hemolytic vascular inflammatory process causing acute lung

310 injury (30, 50). Both Hb-$Fe^{4+}$ and met-hemoglobin induce a drop of mitochondrial oxygen

311 consumption and mitochondrial membrane potential in epithelial lung cells (36). Acute chest

312 syndrome is an important cause of hospitalization and mortality in children with SCD and Spn is a

313 leading etiologic agent of acute chest syndrome. While the precise role of Spn is still unclear, SCD

314 children are ~100-fold more susceptible to pneumococcal infection (71). Studies of the oxidation of

315 hemoglobin by Spn-$H_2O_2$, however, have been neglected. We recently demonstrated that the

316 growth of planktonic Spn, and formation of pneumococcal biofilms, is enhanced by supplementing

317 the culture medium with human hemoglobin (72, 73). Besides iron, other nutrients may become

318 accessible to pneumococcus, since the oxidation of hemoglobin by hydrogen peroxide releases

319 globin-derived peptides (29), but also a striking transcriptome remodeling was identified that

included upregulated transcription of genes encoding transporters of glyco-conjugated molecules

(72, 73).

Besides nutrients, oxidative reactions may lead to unintended consequences. During

pneumococcal pneumonia the pulmonary parenchyma presents with hemorrhage, inflammatory

congestion, hepatization, suppurative infiltration, and lung parenchymal injury (4, 9-13). Whereas

there are a number of virulence factors implicated in the pathophysiology of lung infection (63), the

oxidation of hemoglobin through Spn-$H_2O_2$ may also be an important contributor to pneumococcal

disease. For example, mice intranasally inoculated with Spn develop lung hemorrhage, at necropsy,

and histological analysis reveals lung consolidation associated with alveolar septal edema and

pleomorphic alveolar, interstitial, and perivascular cellular inflammation (74). Spn mutants in the

*spxB* gene whether with a pleotropic capsule defect or not, were attenuated for virulence in mouse

models of pneumococcal disease (46, 47, 75).

There are other important implications as a result of the oxidation of hemoglobin by

hydrogen peroxide produced by streptococci. Heme release and/or heme degradation through,

hydrogen peroxide, may be beneficial to other bacteria which do not synthetize large amounts of

this pro-oxidant but that require iron for their metabolism and pathogenesis. For instance, it has

been recently demonstrated that dental plaque bacteria such as *S. gordonii*, which produces

hydrogen peroxide, oxidizes hemoglobin to release heme, and this free-heme facilitates the

colonization of the dental plaque by a Gram-negative strain *Porphyromonas gingivalis* (49). *P.*

*gingivalis* does not produce siderophores but it has a sophisticated mechanism for heme-iron

acquisition (76). Conversion of oxy-hemoglobin to met-hemoglobin, by hydrogen peroxide

producing bacteria, allows the release of heme that can be taken by these microorganisms.

In  summary,  we  demonstrated  that  Spn  and  other  hydrogen  peroxide-producing

Streptococci, oxidizes hemoglobin from human and other species, releasing and degrading heme.

These oxidative reactions occurred early during the lag phase of growth indicating that metabolic

adaptation  for  growth  in  the  presence  of  hemoglobin  stimulated  production  of  Spn-$H_2O_2$.  This

upregulated metabolic adaption was the result in part of the release of heme since gene expression

studies demonstrated an upregulation of $spxB$ transcription but not that of $lctO$ or the pneumolysin

gene  $ply$,  under  the  culture  conditions  tested.  Noteworthy,  incubation  with  erythrocytes  or

hemoglobin did not induced a regulation of $spxB$ in the current study. We have observed a potential

toxic radical produced during the oxidation of hemoglobin, spanning the incubation time utilized in

gene  expression  studies,  that  may  explain  the  observed  downregulation  of  $spxB$  (Scasny  et  al,

unpublished  observations).  Efforts  are  in  place  to  assess  the  specific  role  for  these  oxidative

reactions for pathogenesis and bacterial symbiosis.

**Material and Methods**

**Bacterial strains, material, and growth conditions**. All Spn wild type strains and mutant derivatives

used in this study are listed in Table 2. Bacterial stocks were prepared in medium containing skim

milk-tryptone-glucose-glycerin  (STGG)  and  stored  at  -80°C  (77).  Pneumococci  were  rotinously

cultivated at 37°C in a 5% $CO_2$ and ~20% $O_2$ atmosphere on blood agar plate (BAP). Non-typeable

*Haemophilus influenzae* (NTHI) was grown on chocolate agar plate (CAP), which was prepared with

heat-lysed  horse  blood,  a  source  of  hemin  and  NAD.  *Klebsiella  pneumoniae*  and  *Staphylococcus*

*aureus* were cultivated on tryptic soy agar (TSA). Experiments were performed using Todd-Hewitt-

broth  with  0.5%  yeast  extract  (THY),   THY  added  with  10  µM  human  met-hemoglobin  (Sigma-

Aldrich), THY added with 5 µM heme (Sigma-Aldrich), THY added with sheep erythrocytes (Quad

Five), and horse erythrocytes (Lampire). In experiments inoculated with NTHI, THY medium was

supplemented with hemin (10 µg/ml), and NAD (2 µg/ml). Brain heart infusion (BHI) broth, Bacto

agar, Bacto Tryptic Soy Broth, Bact Yeast Extract and Bacto Todd Hewitt Broth were purchased from

Becton, Dickinson and Company. Antibiotics utilized were gentamycin, erythromycin,

spectinomycin, chloramphenicol, and were all sourced from Sigma-Aldrich. Hydrogen peroxide

(Fisher Scientific), Nicotinamide adenine dinucleotide (NAD(+)) (Sigma-Aldrich), Catalase (Sigma-

Aldrich), Complete™, Mini Protease Inhibitor Cocktail (Roche), $FeCl_3 \cdot 6H_2O$ and potassium

hexacyanoferrate (III) (Sigma-Aldrich), o-dianisidine (Alfa Aesar) /sodium acetate (Sigma-Aldrich),

ethanol and methanol (Fisher Scientific). Human serum was purchased from MP Biomedicals.

**Preparation of inoculum for experiments.** The inoculum was prepared as previously described (40).

Briefly, bacteria were inoculated on blood agar plates (BAP) and incubated overnight at 37°C in a 5%

$CO_2$ atmosphere. Bacteria were then harvested from plates by PBS washes and this suspension was

used to inoculate THY with or without additives, that was brought to a final $OD_{600}$ of ~0.1. This

suspension contained ~$5.15 \times 10^8$ cfu/ml as confirmed by dilution and platting of aliquots of the

suspension.

**Bacterial broth culture medium containing oxy-hemoglobin.** Sheep or horse erythrocytes were

washed three times with sterile PBS (pH=7.4) at 300 × $g$ for 5 min in a refrigerated centrifuge

(Eppendorf). These washed erythrocytes suspension was lysed with 0.1% saponin and the

concentration of oxy-hemoglobin in this preparation was determined using the QuantiChrom™

Hemoglobin Assay Kit (BioAssay Systems); oxy-hemoglobin was then added to THY broth to a final

concentration of ~10 µM. The concentration, and redox state of oxy-hemoglobin was verified by

spectroscopy before each experiment. In some experiments, THY containing oxy-hemoglobin was

supplemented with an excess of catalase to a final concentration of 400 U/ml, or with 1X complete

protease inhibitors cocktail (Roche).

**Studies of the oxidation of oxy-hemoglobin, met-hemoglobin and heme degradation by**

**spectroscopy.** Experiments were conducted using six-well microplates (Genesee Scientific) added

with THY containing a 2% suspension of washed sheep erythrocytes, 10 µM of oxy-hemoglobin, or

10 µM of met-hemoglobin. In some experiments human serum was supplemented with 10 µM of

met-hemoglobin, or THY was supplemented with human serum to a final 10% concentration (v/v)

and added with 10 µM of met-hemoglobin. These media were inoculated with Spn or the other

bacterial species listed in each experiment, and incubated for the indicated time at 37°C in a 5% $CO_2$

and ~20% $O_2$ atmosphere. At the of the incubation time, supernatants containing planktonic

bacteria were removed from cultures and bacteria were pelleted down by centrifugation at 13,000

rpm x 5 min in a refrigerated micro-centrifuge (Eppendorf). In another set of experiments, THY

containing 10 µM of human met-hemoglobin was treated with different concentrations of hydrogen

peroxide for 6 h. The oxidation of oxy-hemoglobin to met-hemoglobin and further oxidation and

degradation of met-hemoglobin was determined by analyzing the spectra spanning 200 nm through

1000 nm in bacteria-free supernatants, using a spectrophotometer Omega BMG LabTech

(ThermoFisher).

**Quantification of hydrogen peroxide production.** Hydrogen peroxide production was quantified

from bacterial cultures inoculated in six-well microplates (Genesee Scientific) that had been added

with THY. These cultures were incubated at 37°C in a 5% $CO_2$ and ~20% $O_2$ atmosphere for the

indicated time. At the end of the incubation, the culture supernatant was collected, centrifuged at

4°C for 5 min at 15,000 rpm and then the supernatant was transferred to a new tube and kept in ice

for no more than 30 min. Some supernatants were filtered sterilized using a 0.4 µm syringe filter

(Fisher scientific). Collected supernatants were diluted with 1× reaction buffer of the Amplex Red

$H_2O_2$ assay kit (Molecular Probes), and hydrogen peroxide levels were quantified according to the

manufacturer's instructions using a spectrophotometer Omega BMG LabTech (ThermoFisher).

**Qualitative detection of hydrogen peroxide production by *S. pneumoniae* strains.** Qualitative

detection of $H_2O_2$ production by Spn was performed using the Prussian blue agar assay, which limit

of detection is 2.5 nm $H_2O_2$ (55). Prussian blue agar plates were made with 1.0 g/liter of $FeCl_3 \cdot 6H_2O$,

1.0 g/liter of potassium hexacyanoferrate III ($K_3[Fe(CN)_6]$) also known as Prussian blue, 37 g/liter of

dehydrated BHI broth, and 15 g/liter of agar. Prior to experiments, detection of $H_2O_2$ was assessed

by dropping 10 µl of a 1 mM $H_2O_2$ solution that reduced $K_3[Fe(CN)_6]$ and causing its precipitation and

then a blue halo within minutes. PBS was used as a negative control. Spn strains or culture

supernatants (10 µl) obtained as described above, were inoculated, or spotted, respectively, onto

Prussian Blue agar plates. These plates were incubated at 37 °C in a 5% $CO_2$ and ~20% $O_2$

atmosphere overnight (Spn) or for 2 h (supernatants) after which plates were photographed with a

Canon Rebel EOS T5 camera system and digital pictures were analyzed.

**Western blotting.** Bacteria were grown in THY with 10 µM hemoglobin at 37°C for 2 h, 4 h, and 6 h.

After incubation the cells were harvested by centrifuge at 15,000 rpm for 5 min and the supernatant

were transferred to new tube. The collected samples were combined with 4x reducing sample

buffer and boiled 5 minutes. The mixture was loaded into 10 well 4-12% Mini-PROTEAN® TGX™

precast gel (Biorad), and run 2 hours at 90 V in running buffer. Gels were transferred to

nitrocellulose membranes in transfer buffer supplemented with 20% ethanol with Trans-Blot Turbo

Transfer System. Membranes were blocked 1 h at room temperature in 5% non-fat dry milk in TBS

supplemented with 0.1% Tween-20 (TBST). Then a membrane was incubated with the Hb polyclonal

antibody (Invitrogen) at 1:1000 dilution in 5% non-fat dry milk in TBST overnight at 4°C. Next day,

the blot was washed 3x 5 min with TBST, and incubated with donkey anti-goat IgG-HRP (Santa Cruz)

secondary antibody 1:5000 in 5% non-fat dry milk in TBST 1 h at room temperature. Blots were

washed 3x 5 min in TBST and 2mL each SuperSignal Substrate (Thermo Fisher Scientific) reagent was

added to blot and incubated for 5 min at room temperature. Blots were imaged on a ChemiDoc MP

(BioRad) using ImageLab 5.0 software, automatic exposure settings for Chemiluminescence, high

specificity, optimizing for bright bands.

**In-gel heme staining.** Bacteria were grown in THY with 10 µM hemoglobin at 37°C for 2, 4, or 6 h,

after which supernatants were collected, spun down and combined with non-reducing loading

buffer. The mixtures were loaded into a non-denaturing 12% Mini-PROTEAN® TGX™ precast gel

(Biorad), and run for 2 h at 90 V in running buffer lacking SDS. The gel was then stained using a

published protocol (35). Briefly, after electrophoresis gels were immersed into a methanol/sodium

acetate (pH=5) solution and incubated at room temperature on rocking platform (VWR) at 1 speed

for 2 min with constant shaking. After incubation, the gel was added with a solution of o-

dianisidine/sodium acetate (pH=5)and incubated for 20 min in the dark. To visualize heme in the

gel, 3% hydrogen peroxide was added, and the reaction was stopped by washing with distilled

water. The gel was then photographed with a Canon Rebel EOS T5 camera system and digital

pictures were analyzed.

**RNA extraction and qRT-PCR analysis.** Bacteria were inoculated into THY, or THY containing 5 µM

heme, 5 µM Hb, or a 1% suspension of RBC and incubated at 37°C for 2 h. After incubation the

bacterial suspension was collected, and mixed with RNAprotect Reagent (Qiagen). Then cells were

harvested and the total RNA was extracted using the RNeasy mini kit (Qiagen) following the

manufacturer's instruction. DNA was removed using the Turbo DNase-free kit (Life Technologies).

The RNA concentration was obtained using a NanoDrop spectrophotometer (Thermo Fisher

Scientific), and 200 ng of RNA were cDNA transcribed using the iScript™ cDNA Synthesis Kit (Biorad).

Gene expression analysis was carried out using the PerfeCTa SYBR® Green SuperMix (Quantabio)

and CFX96 Touch Real-Time PCR system (Biorad). Primers used for the qRT-PCR analysis are listed in

Table 3. The following conditions were utilized: 1 cycle at 95°C for 3 min and 40 cycles of 95°C for 15

s, 60°C for 30 s and 72°C for 30 s. Melting curves were generated to confirm the absence of primer

dimers. The relative quantitation of mRNA expression was normalized to the constitutive expression

of the housekeeping 16S rRNA gene and calculated by the comparative CT ($2^{-\Delta\Delta CT}$) method (78).

**Preparation of *lctO* mutation in Spn strains EF3030 and D39.** To prepare an isogenic *lctO* mutant,

we PCR-amplified, using primers lcto-spec1 and lcto-spec2 (Table 3), an insertionally inactivated

*lctO*-spectinomycin fragment from strain TIGR4Δ*lctO* (40). This PCR product was purified using the

QIAquick PCR purification kit (Qiagen) and then transformed into EF3030 or D39 using an standard

transformation procedure (79). Transformants were harvested in BAP with spectinomycin

(100 μg/ml) and the insertional inactivation of *lctO* was confirmed by PCR with primers lcto-spec1 /

lcto-spec2 and sequencing.

**Construction of complemented strains.** To complement *spxB*, or *spxB* and *lctO,* into TIGR4Δ*spxB* or

TIGR4Δ*spxB*Δ*lctO*, respectively, we chose a 'neutral' chromosomal location within the *hplA* gene

(SP1113) in which insertions do not affect virulence (52). A *spxB* complemented fragment was

prepared by amplifying an upstream region carrying a fragment of the *hlpA* gene, using as a

template DNA purified from TIGR4, and primers hlpA-up1 and hlpA-up2 (Table 3). We then

generated a PCR fragment, using DNA purified from TIGR4 as a template, containing the *spxB* gene

and its promotor with primers spxB1 and spxB2 (Table 3). A downstream fragment containing the

*catP* gene from chloramphenicol resistance and a fragment of ORF SP1114 was amplified using as a

template DNA from strain JWV500 using primers hplA-down1 and hlpA-down2. PCR products were

purified using a QIAquick PCR purification kit (Qiagen) and ligated by splicing overlap extension PCR

with primers hlpA-up1/hlpA-down2. The PCR-ligated product was verified in a 0.8% DNA gel,

purified from the gel using QIAquick Gel Extraction Kit (Qiagen) and re-amplified by PCR with

primers hlpA-up1 and hlpA-down2. This PCR product was purified as above and used to transform

TIGR4Δ*spxB* using a standard procedure (79). Transformants were harvested in BAP plates

containing chloramphenicol (5 µg/ml) and screened by PCR using primers spxB1/spxB2 to confirm

the complementation of *spxB*.

To construct a *spxB* and *lctO* complemented strain, an upstream DNA fragment containing

*hlpA* and *spxB* was amplified using DNA purified from TIGR4Δ*spxB*Ω*spxB* and primers hlpA1 and

spxB2. Then, a DNA fragment was prepared by PCR amplification of the *lctO* gene and its putative

promotor region using TIGR4 DNA as a template and primers lctO1 and lctO2. The downstream

fragment containing the *catP* gene and a fragment of ORF SP1114 was used as prepared above.

After purification with a QIAquick PCR purification kit (Qiagen), these three DNA fragments were

ligated by splicing overlap extension PCR with primers hlpA-up1/hlpA-down2 and purified. This

construct was re-amplified with primers hlpA-up1 and hlpA-down2, purified and transformed into

TIGR4Δ*spxB*Δ*lctO*. Transformants were screened by PCR using primers hlpA-up1/hlpA-down2and

selected completed clones were further confirmed by $H_2O_2$ production.

**Metal content analysis.** Pneumococcal biofilms cultured in THY-Hb for 4 h were washed 3x with

ice-cold PBS, resuspended in 1 ml PBS and sonicated at high intensity for 5 min in 30 s on 30 s off

cycles using a Bioruptor (Diagenode, Denville, NJ, USA). Protein was determined in all samples by

the Bradford method and a known mass of sample was mineralized in concentrated $HNO_3$ trace

metal grade (Thermo Fisher Scientific) as previously described (80). Iron in each sample was

determined by atomic absorbance spectroscopy (AAS) using a flame 55 AA Absorption Spectrometer

(Agilent, Santa Clara, CA, USA). Analytical grade standards for iron (Thermo Fischer Scientific) were

diluted in 18$\Omega$V water. The iron content on each sample was normalized to the initial mass of

protein as previously reported for other metals (81, 82).

**Statistical analysis.** We performed one-way analysis of variances (ANOVA) followed by Dunnet's

multiple comparison test, when more than two groups were compared or the Student $t$ test to

compare two groups, as indicated. All statistical analysis was performed using the software Graph

Pad Prism (version 8.3.1).

**Figure legends**

**Fig. 1. Hydrogen peroxide produced in _S. pneumoniae_ cultures oxidizes human hemoglobin.**

TIGR4, TIGR4$\Delta spxB\Delta lctO$, or TIGR4$\Omega spxB\Omega lctO$ was inoculated in THY containing oxy-hemoglobin

[Hb (Fe$^{+2}$)] or met-hemoglobin [Hb (Fe$^{+3}$)], or left uninfected, and cultures were incubated for (A-B)

30 min,        (C) 2 h or 6 h, (D-F) 6 h at 37°C in a 5% $CO_2$ atmosphere. Supernatants were collected

and the spectra were obtained using a spectrophotometer Omega BMG LabTech. Panel B shows the

enclosed area in A, corresponding to the absorbance peaks of the alpha chain and the beta chain of

oxy-hemoglobin, flattened in met-hemoglobin. In (A, B and E) the dotted lines indicate the Soret

peak for oxy-hemoglobin (415 nm) or met-hemoglobin (405 nm). (F) THY supplemented with met-

hemoglobin (Hb-Fe$^{3+}$) was added with the indicated concentration of hydrogen peroxide, or left

untreated. These cultures were incubated for 6 h at 37°C in a 5% $CO_2$ atmosphere after which the

supernatants were purified and the spectra were obtained as above. The absorbance at 405 nm

obtained for each assessed concentration, was utilized to construct the graphic. Error bars in all

panels represent the standard errors of the means calculated using data from at least three

independent experiments performed with two technical replicates. In (B-F) One-way ANOVA with

Dunnet's test for multiple comparison was performed, *$p < 0.05$ compared against the Soret peak of

uninfected THY-Hb-$Fe^{2+}$ (B and D), uninfected THY-Hb-$Fe^{2+}$ and incubated 2 h (C) or uninfected THY-

Hb-$Fe^{3+}$(E and F).

**Fig. 2. Preparation of a TIGR4Δ*spxB*Δ*lctO*-derivative complemented strain carrying *spxB* and *lctO***

**or *spxB*.** (A) Schematic diagram showing the chromosomal region where *spxB* and *lctO* (top) or *spxB*

(bottom) was complemented in TIGR4Δ*spxB*Δ*lctO*, to create TIGR4Ω*spxB*Ω*lctO*, TIGR4Ω*spxB*.

Promotor regions of each gene (P) are indicated as well as terminators (†); gene IDs were included

using TIGR4 nomenclature. (B-D) TIGR4, TIGR4Δ*spxB*Δ*lctO*, TIGR4Ω*spxB*Ω*lctO*, or TIGR4Ω*spxB* was

inoculated in THY and incubated at 37°C in a 5% $CO_2$ atmosphere. (B) The $OD_{600}$ of the cultures was

obtained at the indicated time. Error bars represent the standard errors of the means calculated

using data from at least three independent experiments performed with two technical replicates.

(C) Supernatants from 4 h cultures were sterilized and an aliquot was inoculated onto Prussian-blue

agar plates (55) and the plates were incubated 2 h at room temperature after which they were

photographed. This experiment was repeated at least three times. (D) Hydrogen peroxide was

quantified from these supernatants using the Amplex Red assay. Error bars represent the standard

errors of the means calculated using data from two independent experiments performed with two

technical replicates. Limit of detection=0.56 μM.

**Fig. 3. Oxidation of hemoglobin occurs through hydrogen peroxide produced through SpxB**

**activity but not LctO.** (A) TIGR4, TIGR4Δ*spxB*Δ*lctO*, TIGR4Δ*spxB*, TIGR4Δ*lctO*, TIGR4Ω*spxB*Ω*lctO*, or

TIGR4Ω*spxB* or (B) EF3030, EF3030Δ*spxB*Δ*lctO*, EF3030Δ*spxB*, or EF3030Δ*lctO*, or (C) D39,

D39Δ*spxB*Δ*lctO*, D39 Δ*spxB*, or D39 Δ*spxB*/*spxB*  were inoculated in THY containing met-

hemoglobin [Hb ($Fe^{3+}$)] and cultures were incubated for 4 h at 37°C in a 5% $CO_2$ atmosphere.

Supernatants were collected and the spectra were obtained using a spectrophotometer Omega

BMG LabTech. The absorbance at 405 nm obtained from each assessed strain, was utilized to

construct the graphics. Error bars in all panels represent the standard errors of the means calculated

using data from at least three independent experiments performed with two technical replicates

each. One-way ANOVA with Dunnet's test for multiple comparison was performed; NS=not

significant compared to THY-Hb ($Fe^{3+}$), *$p<0.0001$ compared with untreated THY-Hb ($Fe^{3+}$), or

$\Delta spxB\Delta lctO$, or $\Delta spxB$; **$p<0.003$ compared with untreated THY-Hb ($Fe^{3+}$).

**Fig. 4. Oxidation of human hemoglobin by Spn-$H_2O_2$ leads to heme degradation.** (A-C) The samples

were stained by an in-gel staining method. (A) TIGR4 wt was inoculated in THY supplemented with

met-hemoglobin (THY-Hb) and incubated for 2, 4 or 6 h at 37°C in a 5% $CO_2$ atmosphere.

Supernatant was harvested and run in a 12% polyacrylamide gel under non-denaturing conditions

and heme was stained by an in-gel heme staining procedure. Free heme, heme-bound to

hemoglobin monomer (Hb-monomer) or to hemoglobin dimers (Hb-dimer), and the observed

molecular size in kDa, are indicated. In the bottom panel, supernatants were Western blotted using

a polyclonal anti-hemoglobin antibody. (B) TIGR4, TIGR4$\Delta spxB\Delta lctO$, TIGR4$\Delta lctO$, TIGR4$\Delta spxB$,

TIGR4$\Omega spxB$, TIGR4$\Omega spxB\Omega lctO$, or (C) EF3030 wt strains or EF3030$\Delta spxB\Delta lctO$, EF3030$\Delta spxB$, or

EF3030$\Delta lctO$, was inoculated into THY-Hb and incubated for six hours. Supernatants were harvested

and analyzed as in (A).

**Fig. 5. Oxidation of hemoglobin occurs in cultures of oral Streptococci that produce hydrogen**

**peroxide.** THY supplemented with met-hemoglobin (THY-Hb) was left uninoculated or inoculated

with the indicated strain and incubated for 4 h at 37°C in a 5% $CO_2$ atmosphere. Supernatants were

collected and the spectra were obtained using a spectrophotometer Omega BMG LabTech. The

absorbance obtained at 405 nm for each assessed strain, was utilized to construct the graphic. Error

bars represent the standard errors of the means calculated using data from at least three

independent experiments performed with two technical replicates. One-way ANOVA with Dunnet's

test for multiple comparison was performed, *$p<0.05$, or NS=not significant, compared with

untreated THY-Hb. (B) Hydrogen peroxide was quantified from 4 h culture supernatants of each

strain and using the Amplex Red assay. Error bars represent the standard errors of the means

calculated using data from two independent experiments performed with two technical replicates.

Limit of detection=0.56 µM.

**Fig. 6. Hemoglobin is oxidized by Spn-$H_2O_2$ in an *ex vivo* bacteremia model with erythrocytes.** (A)

Model of *ex vivo* pneumococcal bacteremia showing pneumococci with pneumolysin (Ply) located in

the membrane; Ply also gets released through autolysis. Membrane-permeable hydrogen peroxide

($H_2O_2$) is produced as a by-product of the oxidation of pyruvate. Bottom panel, oxy-hemoglobin (Hb-

$Fe^{2+}$) is then release from erythrocytes by Ply-induced erythrocyte lyses and once released, oxy-

hemoglobin gets oxidized by Spn-$H_2O_2$ to met-hemoglobin (Hb-$Fe^{3+}$). Alternatively, top panel, Spn-

$H_2O_2$ permeates inside erythrocytes causing the oxidation of hemoglobin that is then released from

lysed erythrocytes as met-hemoglobin (Hb-$Fe^{3+}$). (B and C) A suspension (2%) of sheep erythrocytes

in THY was inoculated with TIGR4, TIGR4 supplemented with catalase, TIGR4 supplemented with

protease inhibitors (PI), TIGR4$\Delta spxB\Delta lctO$, or TIGR4$\Delta ply$ and the inoculated cultures were incubated

for 30 min (B) or 6 h (C) at 37°C in a 5% $CO_2$ atmosphere. As a control, erythrocytes were lysed with

saponin and oxy-hemoglobin (THY-$Hb^{2+}$), that was released into the supernatant, was considered as

maximum hemoglobin release. (D) Oxy-hemoglobin was left untreated (Hb-$Fe^{2+}$) or added with

added with $H_2O_2$ and incubated for 6 h.(E and F) Human serum was supplemented with 10 µM met-

hemoglobin (HS+$Hb^{3+}$) or serum was used to supplement THY to a final 10% (v/v) and added with 10

µM met-hemoglobin (10% HS+$Hb^{3+}$). These media were inoculated with TIGR4 and incubated for 2,

4 or 6 h. Supernatants from experiments in panels (B-F) were collected and the spectra were

obtained using a spectrophotometer Omega BMG LabTech. In panels B through F, error bars

represent the standard errors of the means calculated using data from at least three independent

experiments. One-way ANOVA with Dunnet's test for multiple comparison was performed, $*p<0.05$,

or NS=not significant, compared with (B and C) TIGR4/catalase, (D) untreated Hb-$Fe^{2+}$, (E) untreated

10% HS+$Hb^{3+}$, or (F) untreated HS+$Hb^{3++}$.

**Fig. 7. Regulation of *ply*, *spxB* and *lctO* genes by incubation with erythrocytes, hemoglobin, or heme.** TIGR4

wt was inoculated in THY broth alone, in THY broth containing 1% sheep erythrocytes, THY with 10

μM hemoglobin (Hb) or THY with 5 μM heme. Cultures were incubated for 2 h at 37°C in a 5% $CO_2$

atmosphere, bacteria were then harvested, and RNA was extracted from these cultures and utilized

as template in qRT-PCR reactions with primers that amplified the *ply*, *spxB*, or *lctO* gene. Average $C_T$

values were normalized to the 16S rRNA gene, and the fold differences, relative to THY control,

were calculated using the comparative $C_T$ method ($2^{-\Delta\Delta C_T}$)(78).

**Acknowledgements**

This study was supported in part by a grant from the National Institutes of Health (NIH; 5R21AI144571-03) to JEV and Wesleyan University institutional funds to TP-B. BA was supported by a Fulbright scholarship awarded by the US Department of State. FMF is supported by the Ronald E. McNair Postbaccalaureate Fellowship program. The content is solely the responsibility of the authors and does not necessarily represent the official view of the NIH or the US Department of State. The authors thank Jaime Carrazco-Carrillo from Wesleyan University for his assistance in some laboratory assays.

**References**

1. Jedrzejas MJ. 2001. Pneumococcal virulence factors: structure and function. Microbiol Mol Biol Rev 65:187-207.

2. Kadioglu A, Weiser JN, Paton JC, Andrew PW. 2008. The role of *Streptococcus pneumoniae* virulence factors in host respiratory colonization and disease. Nat Rev Microbiol 6:288-301.

3. Tomasz A. 2000. *Streptococcus pneumoniae* : molecular biology & mechanisms of disease. Mary Ann Liebert, Inc., Larchmont, NY.

4. van der Poll T, Opal SM. 2009. Pathogenesis, treatment, and prevention of pneumococcal pneumonia. Lancet 374:1543-56.

5. Klugman KP, Madhi SA, Albrich WC. 2008. Novel approaches to the identification of *Streptococcus pneumoniae* as the cause of community-acquired pneumonia. Clin Infect Dis 47 Suppl 3:S202-6.

6.     O'Brien KL, Wolfson LJ, Watt JP, Henkle E, Deloria-Knoll M, McCall N, Lee E, Mulholland K,

Levine OS, Cherian T. 2009. Burden of disease caused by *Streptococcus pneumoniae* in

children younger than 5 years: global estimates. Lancet 374:893-902.

7.     Walker CL, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, O'Brien KL, Campbell H, Black

RE. 2013. Global burden of childhood pneumonia and diarrhoea. Lancet 381:1405-16.

8.     Holyoake LV, Hunt S, Sanguinetti G, Cook GM, Howard MJ, Rowe ML, Poole RK, Shepherd M.

2016. CydDC-mediated reductant export in Escherichia coli controls the transcriptional

wiring of energy metabolism and combats nitrosative stress. Biochem J 473:693-701.

9.     Chao Y, Marks LR, Pettigrew MM, Hakansson AP. 2014. Streptococcus pneumoniae biofilm

formation and dispersion during colonization and disease. Front Cell Infect Microbiol 4:194.

10.    Simell B, Auranen K, Kayhty H, Goldblatt D, Dagan R, O'Brien KL, Pneumococcal Carriage G.

2012. The fundamental link between pneumococcal carriage and disease. Expert Rev

Vaccines 11:841-55.

11.    Eurich DT, Marrie TJ, Minhas-Sandhu JK, Majumdar SR. 2017. Risk of heart failure after

community acquired pneumonia: prospective controlled study with 10 years of follow-up.

BMJ 356:j413.

12.    Musher DM, Rueda AM, Kaka AS, Mapara SM. 2007. The association between pneumococcal

pneumonia and acute cardiac events. Clin Infect Dis 45:158-65.

13.    Masters IB, Isles AF, Grimwood K. 2017. Necrotizing pneumonia: an emerging problem in

children? Pneumonia (Nathan) 9:11.

14.    Brown AO, Millett ER, Quint JK, Orihuela CJ. 2015. Cardiotoxicity during invasive

pneumococcal disease. Am J Respir Crit Care Med 191:739-45.

15. Brissac T, Shenoy AT, Patterson LA, Orihuela CJ. 2018. Cell Invasion and Pyruvate Oxidase-

Derived H2O2 Are Critical for Streptococcus pneumoniae-Mediated Cardiomyocyte Killing.

Infect Immun 86.

16. Nishimoto AT, Rosch JW, Tuomanen EI. 2020. Pneumolysin: Pathogenesis and Therapeutic

Target. Front Microbiol 11:1543.

17. Rubins JB, Duane PG, Charboneau D, Janoff EN. 1992. Toxicity of pneumolysin to pulmonary

endothelial cells in vitro. Infect Immun 60:1740-6.

18. Rubins JB, Duane PG, Clawson D, Charboneau D, Young J, Niewoehner DE. 1993. Toxicity of

pneumolysin to pulmonary alveolar epithelial cells. Infect Immun 61:1352-8.

19. Price KE, Greene NG, Camilli A. 2012. Export requirements of pneumolysin in Streptococcus

pneumoniae. J Bacteriol 194:3651-60.

20. Shak JR, Ludewick HP, Howery KE, Sakai F, Yi H, Harvey RM, Paton JC, Klugman KP, Vidal JE.

2013. Novel role for the Streptococcus pneumoniae toxin pneumolysin in the assembly of

biofilms. mBio 4:e00655-13.

21. Bryant JC, Dabbs RC, Oswalt KL, Brown LR, Rosch JW, Seo KS, Donaldson JR, McDaniel LS,

Thornton JA. 2016. Pyruvate oxidase of Streptococcus pneumoniae contributes to

pneumolysin release. BMC Microbiol 16:271.

22. Jacques LC, Panagiotou S, Baltazar M, Senghore M, Khandaker S, Xu R, Bricio-Moreno L, Yang

M, Dowson CG, Everett DB, Neill DR, Kadioglu A. 2020. Increased pathogenicity of

pneumococcal serotype 1 is driven by rapid autolysis and release of pneumolysin. Nat

Commun 11:1892.

23.   McDevitt E, Khan F, Scasny A, Thompson CD, Eichenbaum Z, McDaniel LS, Vidal JE. 2020.

Hydrogen Peroxide Production by Streptococcus pneumoniae Results in Alpha-hemolysis by

Oxidation of Oxy-hemoglobin to Met-hemoglobin. mSphere 5.

24.   Aprianto R, Slager J, Holsappel S, Veening JW. 2018. High-resolution analysis of the

pneumococcal transcriptome under a wide range of infection-relevant conditions. Nucleic

Acids Res 46:9990-10006.

25.   Kakar S, Hoffman FG, Storz JF, Fabian M, Hargrove MS. 2010. Structure and reactivity of

hexacoordinate hemoglobins. Biophys Chem 152:1-14.

26.   Perutz MF. 1990. Mechanisms regulating the reactions of human hemoglobin with oxygen

and carbon monoxide. Annu Rev Physiol 52:1-25.

27.   Ludlow JT, Wilkerson RG, Sahu KK, Nappe TM. 2020. Methemoglobinemia, StatPearls,

Treasure Island (FL).

28.   Balla J, Jacob HS, Balla G, Nath K, Eaton JW, Vercellotti GM. 1993. Endothelial-cell heme

uptake from heme proteins: induction of sensitization and desensitization to oxidant

damage. Proc Natl Acad Sci U S A 90:9285-9.

29.   Posta N, Csosz E, Oros M, Petho D, Potor L, Kallo G, Hendrik Z, Sikura KE, Mehes G, Toth C,

Posta J, Balla G, Balla J. 2020. Hemoglobin oxidation generates globin-derived peptides in

atherosclerotic lesions and intraventricular hemorrhage of the brain, provoking endothelial

dysfunction. Lab Invest 100:986-1002.

30.   Kato GJ, Steinberg MH, Gladwin MT. 2017. Intravascular hemolysis and the pathophysiology

of sickle cell disease. J Clin Invest 127:750-760.

31.    Shaver CM, Upchurch CP, Janz DR, Grove BS, Putz ND, Wickersham NE, Dikalov SI, Ware LB,

Bastarache JA. 2016. Cell-free hemoglobin: a novel mediator of acute lung injury. Am J

Physiol Lung Cell Mol Physiol 310:L532-41.

32.    Almeida CB, Souza LE, Leonardo FC, Costa FT, Werneck CC, Covas DT, Costa FF, Conran N.

2015. Acute hemolytic vascular inflammatory processes are prevented by nitric oxide

replacement or a single dose of hydroxyurea. Blood 126:711-20.

33.    Rice RH, Lee YM, Brown WD. 1983. Interactions of heme proteins with hydrogen peroxide:

protein crosslinking and covalent binding of benzo[a]pyrene and 17 beta-estradiol. Arch

Biochem Biophys 221:417-27.

34.    Nagababu E, Rifkind JM. 1998. Formation of fluorescent heme degradation products during

the oxidation of hemoglobin by hydrogen peroxide. Biochem Biophys Res Commun 247:592-

6.

35.    Maitra D, Byun J, Andreana PR, Abdulhamid I, Diamond MP, Saed GM, Pennathur S, Abu-

Soud HM. 2011. Reaction of hemoglobin with HOCl: mechanism of heme destruction and

free iron release. Free Radic Biol Med 51:374-86.

36.    Kassa T, Jana S, Strader MB, Meng F, Jia Y, Wilson MT, Alayash AI. 2015. Sickle Cell

Hemoglobin in the Ferryl State Promotes betaCys-93 Oxidation and Mitochondrial

Dysfunction in Epithelial Lung Cells (E10). J Biol Chem 290:27939-58.

37.    Nader E, Romana M, Connes P. 2020. The Red Blood Cell-Inflammation Vicious Circle in

Sickle Cell Disease. Front Immunol 11:454.

38.    Buehler PW, Humar R, Schaer DJ. 2020. Haptoglobin Therapeutics and Compartmentalization

of Cell-Free Hemoglobin Toxicity. Trends Mol Med 26:683-697.

39.   Pericone CD, Overweg K, Hermans PW, Weiser JN. 2000. Inhibitory and bactericidal effects of

hydrogen peroxide production by Streptococcus pneumoniae on other inhabitants of the

upper respiratory tract. Infect Immun 68:3990-7.

40.   Wu X, Gordon O, Jiang W, Antezana BS, Angulo-Zamudio UA, Del Rio C, Moller A, Brissac T,

Tierney ARP, Warncke K, Orihuela CJ, Read TD, Vidal JE. 2019. Interaction between

Streptococcus pneumoniae and Staphylococcus aureus Generates (.)OH Radicals That

Rapidly Kill Staphylococcus aureus Strains. J Bacteriol 201.

41.   Lisher JP, Tsui HT, Ramos-Montanez S, Hentchel KL, Martin JE, Trinidad JC, Winkler ME,

Giedroc DP. 2017. Biological and Chemical Adaptation to Endogenous Hydrogen Peroxide

Production in Streptococcus pneumoniae D39. mSphere 2.

42.   Taniai H, Iida K, Seki M, Saito M, Shiota S, Nakayama H, Yoshida S. 2008. Concerted action of

lactate oxidase and pyruvate oxidase in aerobic growth of Streptococcus pneumoniae: role

of lactate as an energy source. J Bacteriol 190:3572-9.

43.   Pericone CD, Park S, Imlay JA, Weiser JN. 2003. Factors contributing to hydrogen peroxide

resistance in Streptococcus pneumoniae include pyruvate oxidase (SpxB) and avoidance of

the toxic effects of the fenton reaction. J Bacteriol 185:6815-25.

44.   Ramos-Montanez S, Kazmierczak KM, Hentchel KL, Winkler ME. 2010. Instability of ackA

(acetate kinase) mutations and their effects on acetyl phosphate and ATP amounts in

Streptococcus pneumoniae D39. J Bacteriol 192:6390-400.

45.   Echlin H, Frank M, Rock C, Rosch JW. 2020. Role of the pyruvate metabolic network on

carbohydrate metabolism and virulence in Streptococcus pneumoniae. Mol Microbiol

doi:10.1111/mmi.14557.

46.   Echlin H, Frank MW, Iverson A, Chang TC, Johnson MD, Rock CO, Rosch JW. 2016. Pyruvate

Oxidase as a Critical Link between Metabolism and Capsule Biosynthesis in Streptococcus

pneumoniae. PLoS Pathog 12:e1005951.

47.   Ramos-Montanez S, Tsui HC, Wayne KJ, Morris JL, Peters LE, Zhang F, Kazmierczak KM, Sham

LT, Winkler ME. 2008. Polymorphism and regulation of the spxB (pyruvate oxidase) virulence

factor gene by a CBS-HotDog domain protein (SpxR) in serotype 2 Streptococcus

pneumoniae. Mol Microbiol 67:729-46.

48.   Rai P, Parrish M, Tay IJ, Li N, Ackerman S, He F, Kwang J, Chow VT, Engelward BP. 2015.

Streptococcus pneumoniae secretes hydrogen peroxide leading to DNA damage and

apoptosis in lung cells. Proc Natl Acad Sci U S A 112:E3421-30.

49.   Brown JL, Yates EA, Bielecki M, Olczak T, Smalley JW. 2018. Potential role for Streptococcus

gordonii-derived hydrogen peroxide in heme acquisition by Porphyromonas gingivalis. Mol

Oral Microbiol 33:322-335.

50.   van Beers EJ, Samsel L, Mendelsohn L, Saiyed R, Fertrin KY, Brantner CA, Daniels MP, Nichols

J, McCoy JP, Kato GJ. 2014. Imaging flow cytometry for automated detection of hypoxia-

induced erythrocyte shape change in sickle cell disease. Am J Hematol 89:598-603.

51.   Guiral S, Henard V, Laaberki MH, Granadel C, Prudhomme M, Martin B, Claverys JP. 2006.

Construction and evaluation of a chromosomal expression platform (CEP) for ectopic,

maltose-driven gene expression in Streptococcus pneumoniae. Microbiology (Reading)

152:343-349.

52.   Kjos M, Aprianto R, Fernandes VE, Andrew PW, van Strijp JA, Nijland R, Veening JW. 2015.

Bright fluorescent Streptococcus pneumoniae for live-cell imaging of host-pathogen

interactions. J Bacteriol 197:807-18.

53.    Gallay C, Sanselicio S, Anderson ME, Soh YM, Liu X, Stamsas GA, Pelliciari S, van Raaphorst R,

761            Denereaz J, Kjos M, Murray H, Gruber S, Grossman AD, Veening JW. 2021. CcrZ is a

762            pneumococcal spatiotemporal cell cycle regulator that interacts with FtsZ and controls DNA

763            replication by modulating the activity of DnaA. Nat Microbiol 6:1175-1187.

764    54.    Mercy C, Ducret A, Slager J, Lavergne JP, Freton C, Nagarajan SN, Garcia PS, Noirot-Gros MF,

765            Dubarry N, Nourikyan J, Veening JW, Grangeasse C. 2019. RocS drives chromosome

766            segregation and nucleoid protection in Streptococcus pneumoniae. Nat Microbiol 4:1661-

767            1670.

768    55.    Saito M, Seki M, Iida K, Nakayama H, Yoshida S. 2007. A novel agar medium to detect

769            hydrogen peroxide-producing bacteria based on the prussian blue-forming reaction.

770            Microbiol Immunol 51:889-92.

771    56.    Nagababu E, Rifkind JM. 2004. Heme degradation by reactive oxygen species. Antioxid Redox

772            Signal 6:967-78.

773    57.    Pamplona A, Ferreira A, Balla J, Jeney V, Balla G, Epiphanio S, Chora A, Rodrigues CD,

774            Gregoire IP, Cunha-Rodrigues M, Portugal S, Soares MP, Mota MM. 2007. Heme oxygenase-1

775            and carbon monoxide suppress the pathogenesis of experimental cerebral malaria. Nat Med

776            13:703-10.

777    58.    Mingone CJ, Gupte SA, Chow JL, Ahmad M, Abraham NG, Wolin MS. 2006. Protoporphyrin IX

778            generation from delta-aminolevulinic acid elicits pulmonary artery relaxation and soluble

779            guanylate cyclase activation. Am J Physiol Lung Cell Mol Physiol 291:L337-44.

780    59.    Keller LE, Robinson DA, McDaniel LS. 2016. Nonencapsulated Streptococcus pneumoniae:

781            Emergence and Pathogenesis. mBio 7:e01792.

782    60.    Facklam R. 2002. What happened to the streptococci: overview of taxonomic and

783            nomenclature changes. Clin Microbiol Rev 15:613-30.

784    61.    Kajfasz JK, Rivera-Ramos I, Abranches J, Martinez AR, Rosalen PL, Derr AM, Quivey RG,

785            Lemos JA. 2010. Two Spx proteins modulate stress tolerance, survival, and virulence in

786            Streptococcus mutans. J Bacteriol 192:2546-56.

787    62.    van der Hoeven JS, Hoogendoorn H. 1990. Uptake of oxygen, release and degradation of

788            hydrogen peroxide by Streptococcus mutans NCTC 10449. Antonie Van Leeuwenhoek 57:91-

789            5.

790    63.    Weiser JN, Ferreira DM, Paton JC. 2018. Streptococcus pneumoniae: transmission,

791            colonization and invasion. Nat Rev Microbiol 16:355-367.

792    64.    De Jesus-Bonilla W, Ramirez-Melendez E, Cerda J, Lopez-Garriga J. 2002. Evidence for

793            nonhydrogen bonded compound II in cyclic reaction of hemoglobin I from Lucina pectinata

794            with hydrogen peroxide. Biopolymers 67:178-85.

795    65.    Nagababu E, Rifkind JM. 2000. Heme degradation during autoxidation of oxyhemoglobin.

796            Biochem Biophys Res Commun 273:839-45.

797    66.    Mao GD, Thomas PD, Poznansky MJ. 1994. Oxidation of spin trap 5,5-dimethyl-1-pyrroline-1-

798            oxide in an electron paramagnetic resonance study of the reaction of methemoglobin with

799            hydrogen peroxide. Free Radic Biol Med 16:493-500.

800    67.    Iuliano L, Violi F, Pedersen JZ, Pratico D, Rotilio G, Balsano F. 1992. Free radical-mediated

801            platelet activation by hemoglobin released from red blood cells. Arch Biochem Biophys

802            299:220-4.

68.   Khan F, Wu X, Matzkin GL, Khan MA, Sakai F, Vidal JE. 2016. Streptococcus pneumoniae

Eradicates Preformed Staphylococcus aureus Biofilms through a Mechanism Requiring

Physical Contact. Front Cell Infect Microbiol 6:104.

69.   Regev-Yochay G, Trzcinski K, Thompson CM, Malley R, Lipsitch M. 2006. Interference

between Streptococcus pneumoniae and Staphylococcus aureus: In vitro hydrogen peroxide-

mediated killing by Streptococcus pneumoniae. J Bacteriol 188:4996-5001.

70.   Giulivi C, Hochstein P, Davies KJ. 1994. Hydrogen peroxide production by red blood cells.

Free Radic Biol Med 16:123-9.

71.   Assad Z, Michel M, Valtuille Z, Lazzati A, Boizeau P, Madhi F, Gaschignard J, Pham LL, Caseris

M, Cohen R, Kaguelidou F, Varon E, Alberti C, Faye A, Angoulvant F, Koehl B, Ouldali N. 2022.

Incidence of Acute Chest Syndrome in Children With Sickle Cell Disease Following

Implementation of the 13-Valent Pneumococcal Conjugate Vaccine in France. JAMA Netw

Open 5:e2225141.

72.   Akhter F, Womack E, Vidal JE, Le Breton Y, McIver KS, Pawar S, Eichenbaum Z. 2021.

Hemoglobin induces early and robust biofilm development in Streptococcus pneumoniae by

a pathway that involves comC but not the cognate comDE two-component system. Infect

Immun doi:10.1128/IAI.00779-20.

73.   Akhter F, Womack E, Vidal JE, Le Breton Y, McIver KS, Pawar S, Eichenbaum Z. 2020.

Hemoglobin stimulates vigorous growth of Streptococcus pneumoniae and shapes the

pathogen's global transcriptome. Sci Rep 10:15202.

74.   Gotts JE, Bernard O, Chun L, Croze RH, Ross JT, Nesseler N, Wu X, Abbott J, Fang X, Calfee CS,

Matthay MA. 2019. Clinically relevant model of pneumococcal pneumonia, ARDS, and

nonpulmonary organ dysfunction in mice. Am J Physiol Lung Cell Mol Physiol 317:L717-L736.

75.    Spellerberg B, Cundell DR, Sandros J, Pearce BJ, Idanpaan-Heikkila I, Rosenow C, Masure HR.

827            1996. Pyruvate oxidase, as a determinant of virulence in Streptococcus pneumoniae. Mol

828            Microbiol 19:803-13.

829    76.    Smalley JW, Olczak T. 2017. Heme acquisition mechanisms of Porphyromonas gingivalis -

830            strategies used in a polymicrobial community in a heme-limited host environment. Mol Oral

831            Microbiol 32:1-23.

832    77.    O'Brien KL, Bronsdon MA, Dagan R, Yagupsky P, Janco J, Elliott J, Whitney CG, Yang YH,

833            Robinson LG, Schwartz B, Carlone GM. 2001. Evaluation of a medium (STGG) for transport

834            and optimal recovery of Streptococcus pneumoniae from nasopharyngeal secretions

835            collected during field studies. J Clin Microbiol 39:1021-4.

836    78.    Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time

837            quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25:402-8.

838    79.    Vidal JE, Ludewick HP, Kunkel RM, Zahner D, Klugman KP. 2011. The LuxS-dependent

839            quorum-sensing system regulates early biofilm formation by Streptococcus pneumoniae

840            strain D39. Infect Immun 79:4050-60.

841    80.    Raimunda D, Padilla-Benavides T, Vogt S, Boutigny S, Tomkinson KN, Finney LA, Arguello JM.

842            2013. Periplasmic response upon disruption of transmembrane Cu transport in

843            Pseudomonas aeruginosa. Metallomics 5:144-51.

844    81.    Tavera-Montanez C, Hainer SJ, Cangussu D, Gordon SJV, Xiao Y, Reyes-Gutierrez P, Imbalzano

845            AN, Navea JG, Fazzio TG, Padilla-Benavides T. 2019. The classic metal-sensing transcription

846            factor MTF1 promotes myogenesis in response to copper. FASEB J 33:14556-14574.

82.  Gordon SJV, Xiao Y, Paskavitz AL, Navarro-Tito N, Navea JG, Padilla-Benavides T. 2019.

Atomic Absorbance Spectroscopy to Measure Intracellular Zinc Pools in Mammalian Cells. J

Vis Exp doi:10.3791/59519.

83.  Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, Heidelberg J, DeBoy RT, Haft

DH, Dodson RJ, Durkin AS, Gwinn M, Kolonay JF, Nelson WC, Peterson JD, Umayam LA, White

O, Salzberg SL, Lewis MR, Radune D, Holtzapple E, Khouri H, Wolf AM, Utterback TR, Hansen

CL, McDonald LA, Feldblyum TV, Angiuoli S, Dickinson T, Hickey EK, Holt IE, Loftus BJ, Yang F,

Smith HO, Venter JC, Dougherty BA, Morrison DA, Hollingshead SK, Fraser CM. 2001.

Complete genome sequence of a virulent isolate of Streptococcus pneumoniae. Science

293:498-506.

84.  Lanie JA, Ng WL, Kazmierczak KM, Andrzejewski TM, Davidsen TM, Wayne KJ, Tettelin H,

Glass JI, Winkler ME. 2007. Genome sequence of Avery's virulent serotype 2 strain D39 of

Streptococcus pneumoniae and comparison with that of unencapsulated laboratory strain

R6. J Bacteriol 189:38-51.

85.  Keller LE, Luo X, Thornton JA, Seo KS, Moon BY, Robinson DA, McDaniel LS. 2015.

Immunization with Pneumococcal Surface Protein K of Nonencapsulated Streptococcus

pneumoniae Provides Protection in a Mouse Model of Colonization. Clin Vaccine Immunol

22:1146-53.

86.  Junges R, Maienschein-Cline M, Morrison DA, Petersen FC. 2019. Complete Genome

Sequence of Streptococcus pneumoniae Serotype 19F Strain EF3030. Microbiol Resour

Announc 8.

87.    Keller LE, Thomas JC, Luo X, Nahm MH, McDaniel LS, Robinson DA. 2013. Draft Genome

869           Sequences of Five Multilocus Sequence Types of Nonencapsulated Streptococcus

870           pneumoniae. Genome Announc 1.

871    88.    Boake WC. 1956. Antistaphylocoagulase in experimental staphylococcal infections. J

872           Immunol 76:89-96.

888    **Table 1. Quantification of cell-free hemoglobin and the heme-hemoglobin Soret curve.**

| Strains | Cell-free hemoglobin (µM) | Absorbance at 415 nm |
|---|---|---|
| Uninfected | 17.35 | 1.38 |
| TIGR4 | 12.8 | 1.39 |
| TIGR4Δ*spxB*Δ*lctO* | 16.4 | 1.21 |

| Strain | Characteristics | Reference or source |
|---|---|---|
| TIGR4 | Reference strain, whole genome sequenced, capsular serotype 4 | (83) |
| SPJV41 (TIGR4Δ*spxB*Δ*lctO*) | TIGR4 with a deleted *spxB* and *lctO* gene by insertion-deletion with an erythromycin and spectinomycin cassette, respectively | (40) |
| SPJV29 (TIGR4Δ*spxB*) | TIGR4 with an insertionally inactivated *spxB*::*ermB* (Ery^r) | (23) |
| SPJV42 (TIGR4Δ*lctO*) | TIGR4 with an insertionally inactivated *lctO*:: *aad9* (Spc^r) | (40) |
| SPJV43 (TIGR4Δ*spxB*Ω*spxB*) | SPJV29 complemented with the gene *spxB* and its promoter | This study |
| SPJV44 (TIGR4Δ*spxB*Δ*lctO*Ω*spxB*- | SPJV41 complemented with the gene | This study |

| lctO) | spxB and lctO | | 907 |
|---|---|---|---|
| D39 | Avery strain, clinical isolate, capsular serotype 2 | (84) | 908 |
| P878 | D39 with a deletion in spxB | (39) | 909 |
| P1221 | P878 complemented with pMU1328::spxB | (43) | 910 |
| SPJV45 (D39ΔspxBΔlctO) | D39 with a deleted spxB and lctO gene by insertion-deletion with an erythromycin and spectinomycin cassette, respectively | (23) | 911 <br> 912 |
| EF3030 | Clinical isolate, caccine serotype 19F strain | (85, 86) | 913 |
| SPJV49 (EF3030ΔspxBΔlctO) | EF3030 with a deleted spxB and lctO gene by insertion-deletion with an erythromycin and spectinomycin cassette, respectively | (23) | 914 <br> 915 |
| SPJV50 (EF3030ΔspxB) | EF3030 with an insertionally inactivated spxB::ermB (Ery$^r$) | This study | 916 |
| SPJV51 (EF3030ΔlctO) | EF3030 with an insertionally inactivated lctO:: aad9 (Spc$^r$) | This study | 917 |
| MNZ41 | Non-encapsulated S. pneumoniae strain | (87) | 918 |
| MNZ67 | Non-encapsulated S. pneumoniae strain | (87) | |
| S. pseudopneumoniae | ATCC 960 | Laboratory stock | 919 |
| S. salivarius | ATCC 7073 | Laboratory stock | 920 |
| S. oralis | ATCC 35037 | Laboratory stock | 921 |
| K. pneumoniae | ATCC 700603 | Laboratory stock | 922 |
| Staphylococcus aureus strain Newman | NCTC 8178, ATCC 13420 | (88) | 923 |
| Non-typeable Haemophilus influenzae (NTHI) | M5029 | Laboratory stock | 924 |
| S. mitis | ATCC 49456 | Laboratory stock | 925 |
| S. mutans | ATCC 25175 | Laboratory stock | 926 |

927    s study

| Primer | Sequence (5' → 3') |
|---|---|
| hlpA-up1 | TCAGCAGGTTCATGAGGGAA |

| | |
|---|---|
| hlpA-up2 | TCATTTCTGTTTTATAACAAAGTCCGGATCCTTTAACAGCGT |
| hlpA-down1 | CTCGCCGAAAATCAAATATGATCACTCACGGCATGGATGA |
| hlpA-down2 | CAAAACAACATTGCCCGACG |
| spxB1 | CTTTGTTATAAAACAGAAATGA |
| spxB2 | CATATTTGATTTTCGGCGAG |
| hlpA-up3 | TTTCCCCAATTATTCACCCCACATATTTGATTTTCGGCGAG |
| lcto1 | TGGGGTGAATAATTGGGGAAA |
| lcto2 | CATTCAGTGGAGGCAATCTGT |
| lcto-spec1 | TTTCTGAGTAGGCGGAGTGG |
| lcto-spec2 | GCAATTTTCAGAGCAGCTTGG |
| JVS109L | TCTACTCAATCGACGTAGGTAACA |
| JVS110R | TATTGAACGTTTGTGATAACGTCT |
| JVS59L | TGAGACTAAGGTTACAGCTTACAG |
| JVS60R | CTAATTTTGACAGAGAGATTACGA |
| JVS105L | TTTGCAATGTAGAAAATCCAAGTA |
| JVS106R | AAATCTCTGGAAGGTCAACAGTAG |
| JVS49L | CTGATGACTGTCAATCGAGATACC |
| JVS50R | AATGGACGAATCAACTCCATAA |
| JVS35L | AACCAAGTAACTTTGAAAGAAGAC |
| JVS36R | AAATTTAGAATCGTGGAATTTTT |

**A**

THY-Hb

TIGR4
Time (h)
2  4  6

**B**

TIGR4
Δ*spxB*Δ*lcto*
Δ*lcto*
Δ*spxB*
Ω*spxB*Ω*lcto*
Ω*spxB*
THY-Hb

**C**

EF3030
Δ*spxB*Δ*lcto*
Δ*spxB*
Δ*lctO*

Hb-dimer (28)
Hb-mono (15)
Heme (0.5)

**Wb anti-Hb**