BIN1 knockdown rescues systolic dysfunction in the aging heart

Maartje Westhoff Ph.D.[1]*, Silvia G. del Villar Ph.D.[1]*, Taylor L. Voelker[1] Ph.D., Phung N. Thai Ph.D.[2,3], Nipavan Chiamvimonvat M.D.[2,3,4], Eamonn J. Dickson Ph.D.[1], and Rose E. Dixon Ph.D.[1]

These authors contributed equally to this work.

[1]Department of Physiology and Membrane Biology, University of California Davis, Davis, CA, USA
[2]Division of Cardiovascular Medicine, Department of Internal Medicine, University of California, Davis, Davis, CA, USA
[3]Department of Veterans Affairs, Northern California Health Care System, Mather, CA, USA
[4]Department of Pharmacology, University of California Davis, Davis, CA, USA

MW: https://orcid.org/0000-0002-5421-8609
SGdV: https://orcid.org/0000-0002-5612-4761
TLV: https://orcid.org/0000-0002-6779-9350
PNT: https://orcid.org/0000-0001-7344-150X
NC: https://orcid.org/0000-0001-9499-8817
EJD: https://orcid.org/0000-0001-8619-7630
RED: https://orcid.org/0000-0003-0655-690X

Corresponding author: Rose E. Dixon, redickson@ucdavis.edu Department of Physiology and Membrane Biology, 4112 Tupper Hall, One Shields Avenue, University of California Davis, Davis, CA 95616, USA

Abstract word count: 150

Main text word count (Introduction, Results, and Discussion): 5083

Abstract

Cardiac dysfunction is a hallmark of aging in humans and mice. Here we report that a two-week treatment to restore youthful Bridging Integrator 1 (BIN1) levels in the hearts of 24-month-old mice rejuvenated cardiac function and substantially reversed the aging phenotype. Our data indicate that age-associated overexpression of BIN1 occurs alongside dysregulated endosomal recycling and disrupted trafficking of cardiac $Ca_V1.2$ and type 2 ryanodine receptors. These deficiencies affect channel function at rest and their upregulation during acute stress. *In vivo* echocardiography revealed reduced systolic function in old mice. BIN1 knockdown using an adeno-associated virus serotype 9 packaged shRNA-mBIN1 restored the nanoscale distribution and clustering plasticity of ryanodine receptors and recovered Ca^{2+} transient amplitudes and cardiac systolic function toward youthful levels. Enhanced systolic function correlated with increased phosphorylation of the myofilament protein cardiac myosin binding protein-C. These results reveal BIN1 knockdown as a novel therapeutic strategy to rejuvenate the aging myocardium.

Keywords:

Calcium channels, ryanodine receptors, aging, $Ca_V1.2$, BIN1, *β*-adrenergic receptors, EC-coupling, clustering, cardiac myosin binding protein-C, cardiac troponin I

Graphical Abstract

Introduction

Old age is a major independent risk factor for cardiovascular disease which remains the leading cause of death globally[1,2]. As humans advance into old age, regardless of overall health, certain intrinsic and progressive changes in cardiac structure and function occur including enhanced atrial and ventricular fibrosis, left ventricular hypertrophy, and diastolic dysfunction[3-5]. Maximal systolic function is also impaired due to a reduced responsiveness to sympathetic nervous system activation and subsequent β-adrenergic receptor (β-AR) signaling[3-5]. The decline of this reactive stimulus can precipitate exercise intolerance in the elderly and render the heart more vulnerable to damage during acutely stressful events when hypoxia and metabolic stress ensue[4,6]. In mice, similar intrinsic cardiac changes occur over a much shorter lifespan and in the absence of other risk factors like smoking or diabetes, making mice a useful model to understand the molecular mechanisms of cardiac aging[3].

The force of myocardial contraction and the rate and extent of the subsequent relaxation are tuned by both Ca^{2+} and myofilament-dependent processes. Examination and comparison of these points of regulation and their integration in the *in vivo* function of young and old mice may identify targets to improve the cardiac aging phenotype. Weakened myocardial contractile responses to βAR-stimulation have been linked to an age-associated decrease in the ability of the βAR signaling cascade to augment intracellular Ca^{2+} transient amplitude[7,8]. This has been attributed to a reduced capacity to increase $Ca_V1.2$ activity and availability, and thus to diminished Ca^{2+} influx during the action potential (AP)[5]. Activation of cAMP-dependent protein kinase A (PKA), downstream of β-AR stimulation, leads to phosphorylation of the channel complex[9,10], removing an inhibitory brake[11,12] and resulting in increased channel open probability (P_o) due to potentiation of longer-duration 'mode 2' openings[13], and an increased number of functional channels[14,15]. We recently reported that β-AR activation also mobilizes a sub-sarcolemmal pool of $Ca_V1.2$ channels, triggering a PKA-dependent, dynamic increase in sarcolemmal $Ca_V1.2$ expression in ventricular myocytes[15,16]. Consequent enlargement of $Ca_V1.2$ channel clusters

facilitates cooperative channel interactions[16,17], and contributes to an amplification of Ca^{2+} influx

that rapidly tunes excitation-contraction (EC)-coupling to meet increased hemodynamic and

metabolic demands. We hypothesized that loss of this rheostatic mechanism in aging myocytes

could impose a narrow inotropic dynamic range on the system, limiting the response to β-AR

stimulation.

Nanoscale redistribution and augmentation of cardiac Ca^{2+} channel clustering in response

to β-AR stimulation also extends to RyR2. Accordingly, acute treatment with the βAR-agonist

isoproterenol (ISO) or a phosphorylation-inducing cocktail promotes enhanced RyR2 cluster

sizes[18,19], as recently reviewed[20]. This dynamic redistribution of RyR2 into larger clusters is

orchestrated by bridging integrator 1 (BIN1)[18] and is linked to increased Ca^{2+} spark frequency

which summate to produce larger Ca^{2+} transients[19,21]. There is widespread agreement that β-AR

stimulated Ca^{2+} transient augmentation is impaired in the aging myocardium[5,8], however the

underlying mechanisms are not understood.

Here, we present a comprehensive examination of cardiac Ca^{2+} channel function, their

nanoscale distribution, and ISO-stimulated redistribution to test the central hypothesis that age-

dependent disruption in the basal and on-demand trafficking and clustering of these channels

restricts EC-coupling plasticity during myocardial aging and contributes to the pathophysiology of

cardiac aging. We report that old myocytes exhibit basal super-clustering of both $Ca_V1.2$ and

RyR2, with no additional dynamic response to ISO. We link these deficits in cardiac Ca^{2+} channel

recycling and mobility to an age-associated upregulation in BIN1 and the development of

endosomal trafficking deficits or "endosomal traffic jams". Crucially, we find that shRNA-mediated

knockdown of BIN1 in old animals restores a young phenotype by re-establishing RyR2

organization, Ca^{2+} transient amplitude and their β-AR stimulated augmentation, and rejuvenating

myofilament protein phosphorylation levels to recover youthful systolic function.

Results

β-AR stimulated I_{Ca} augmentation is diminished in aging

We began our study with an investigation into the $Ca_V1.2$ channel response to β-AR stimulation

by recording whole-cell Ca^{2+} currents (I_{Ca}) from 3-month-old and 24-month-old (henceforth

referred to as young and old, respectively) ventricular myocytes under control and 100 nM ISO-

stimulated conditions. In young myocytes, ISO elicited a 1.59-fold increase in peak I_{Ca} density

(Fig. 1a-b, and Supplementary Table 1). However, old myocytes exhibited a blunted response

showing only a 1.19-fold increase with ISO and basal currents were significantly larger in old

myocytes than in young controls (Fig. 1a-c). The maximal ISO-stimulated current amplitude was

similar in young and old myocytes (Fig. 1c), suggesting that the "ceiling" was intact but that the

"floor" or baseline shifted with aging leaving less scope for current augmentation with acute

adrenergic stress. Given the reduced ISO-stimulated current augmentation, these data suggest

a reduced β-adrenergic responsiveness in old mice versus young.

Reduced capacity of β-AR stimulation to tune EC-coupling in aging myocytes

Influx of Ca^{2+} through $Ca_V1.2$ channels during AP-mediated depolarization stimulates Ca^{2+}-

induced Ca^{2+}-release from cross-dyad RyR2 to initiate EC-coupling. The amplitude of the

resultant Ca^{2+} transient largely dictates the magnitude of myocardial contraction and the fraction

of blood ejected, assuming constant Ca^{2+} sensitivity of the myofilaments. β-AR stimulation

enhances both $Ca_V1.2$ and RyR2-mediated Ca^{2+} influx to produce positive inotropy and a greater

ejection fraction. We thus examined Ca^{2+} transients in young and old myocytes to investigate the

effects of aging on β-AR tuning of EC-coupling. Transients evoked from Fluo-4AM-loaded

myocytes paced at 1 Hz exhibited a 4.6 ± 0.4-fold enhancement in amplitude with ISO in young

cells (Fig. 1d-f). In a parallel with the I_{Ca} results, basal Ca^{2+} transients in aging myocytes were

larger than in young myocytes (Fig. 1d and f) and the fold-increase with ISO was almost halved

(2.1 $\pm$ 0.3-fold change; Fig. 1e). These results further suggest reduced β-adrenergic

responsiveness in old hearts and diminished myocardial responses to acute stress.

Age-dependent alterations in nanoscale distribution and clustering of Ca$_V$1.2 channels

Increased sarcolemmal Ca$_V$1.2 channel expression and clustering at rest in aging cells could

explain the age-dependent increase in basal I_{Ca}. Furthermore, the reduced responsivity to ISO

could be explained by an impaired capacity to mobilize/recycle additional endosomal Ca$_V$1.2

channels in aging myocytes. In young cells, the result of ISO-stimulated channel recycling can be

observed as Ca$_V$1.2 super-clustering in single molecule localization microscopy (SMLM)[15]. To

determine whether ISO-stimulated recycling was still functional in aging myocytes, we examined

the nanoscale distribution and clustering of Ca$_V$1.2 in t-tubule and crest regions of young and old

myocytes using super-resolution SMLM with and without ISO-stimulation. These experiments

confirmed that the super-clustering response to ISO was intact in the t-tubules of young myocytes,

where Ca$_V$1.2 channel cluster areas in ISO-treated young myocytes were 28.8% larger than

unstimulated controls. However, in old myocytes, channels were already basally super-clustered,

showed no augmentation in area with ISO, and were similarly sized to clusters in ISO-stimulated

young cells (Fig. 2a-b). This phenomenon was isolated to the t-tubules as crest Ca$_V$1.2

populations showed no significant alteration in cluster area (Supplementary Fig. 1). Overall, the

age-dependent increase in basal Ca$_V$1.2 channel cluster area and failure to elicit any additional

increase with ISO suggests that a larger number of channels are already present in old myocyte

t-tubule sarcolemma and that the ISO-stimulated channel insertion response is impaired in aging.

Age-dependent alterations in nanoscale distribution and clustering of RyR2 channels

Other groups have reported a similar phosphorylation-induced cluster size expansion of the cross-

dyad RyR2s in young ventricular myocytes[18,19]. In agreement with those reports, SMLM

performed on young myocytes immunostained against RyR2, revealed that average RyR2 cluster

size grew by 32.7% in young myocytes after acute ISO treatment (Fig. 2c-d). However, mean

RyR2 cluster areas in unstimulated old myocytes were similarly sized to the expanded clusters

observed in ISO-stimulated young myocytes and did not undergo any further expansion with ISO.

These findings suggest that the ISO-stimulated nanoscale redistribution of RyR2 is also impaired

in aging.

ISO-stimulated enhancement of Ca$_V$1.2-RyR2 interactions is absent in aging

Since functional crosstalk between Ca$_V$1.2 and RyR2 is an essential requirement for cardiac EC-

coupling, we next examined the number of sites of proximity between Ca$_V$1.2 and RyR2 using a

proximity ligation assay (PLA). In this assay, a fluorescent signal is emitted at each location where

Ca$_V$1.2 and RyR2 come within 40 nm of each other[22]. In young myocytes, acute ISO treatment

triggered the emergence of more Ca$_V$1.2-RyR2 proximity sites (Fig. 2e-f). This phenomenon was

not observed in old myocytes, where the number of Ca$_V$1.2-RyR2 proximity sites appeared to

have already reached a maximal level in the absence of any exogenous stimulation.

Dynamic TIRF imaging reveals reduced ISO-stimulated Ca$_V$1.2 insertion in aging

The basal super-clustering and absence of an additional response to ISO in old myocytes may

indicate: (i) the endosomal reservoir is empty after already being recycled and inserted; and/or (ii)

the trafficking and mobilization of channels is somehow impaired by aging. To interrogate these

hypotheses, we visualized real-time dynamics of the channels utilizing a Ca$_V$1.2 "biosensor"

approach pioneered by our lab. In this technique, adeno-associated virus serotype 9 (AAV9)-

Ca$_V$β$_2$-paGFP auxiliary subunits are transduced via retro-orbital injections and then visualized in

isolated myocytes using TIRF microscopy as previously described[15,16]. The relative balance

between insertion and endocytosis dictates the overall expression of any membrane protein. An

insertion-heavy mismatch between them will lead to increased membrane expression of the

protein, while an endocytosis-heavy mismatch will favor reduced expression over time. We used

image math to calculate the relative population of channels that were inserted during the ISO

treatment, removed/endocytosed, or that remained static during the whole recording. As expected

based on our previous studies[15,16], young myocytes responded to ISO by inserting more $Ca_V1.2$

channels into the sarcolemma than they endocytosed, resulting in a net change in $Ca_V\beta_2$-paGFP

in the TIRF footprint that on average amounted to 18% (Fig. 3a, c-g). However, old myocytes

exhibited a comparatively blunted response with fewer ISO-stimulated insertions (Fig. 3b-c) and

a larger population of static channels in the TIRF footprint suggesting impaired mobility (Fig. 3e).

Consequently, the ISO-induced change in TIRF footprint $Ca_V\beta_2$-paGFP amounted to only 9.5%

(Fig. 3g). Overall, these data reveal an age-dependent deficit in ISO-triggered insertion of $Ca_V1.2$,

and a shift towards reduced channel mobility or "stranding" at the plasma membrane.

Endosomal traffic jams impede mobilization of endosomal $Ca_V1.2$ reservoirs in aging

We further investigated the effects of aging on the endosomal reservoir and capacity to mobilize

$Ca_V1.2$ channel cargo in response to ISO by examining EEA1-positive early endosome, and

Rab11-positive recycling endosome pools of $Ca_V1.2$ in young and old myocytes. Accordingly,

Airyscan super-resolution imaging on young myocytes revealed immunostained $Ca_V1.2$ on

14.89% of EEA1 positive pixels (Fig. 4a-b). As we have previously reported, *β*-AR activation

stimulates the recycling of a portion of the early endosome $Ca_V1.2$ to the plasma membrane via

the Rab4-choreographed fast endosomal recycling pathway[15]. This was evidenced by a

significant decrease in the % colocalization between EEA1 and $Ca_V1.2$ after ISO in young

myocytes. In old myocytes, there was a trending increase in the percentage of EEA1/$Ca_V1.2$

colocalization in unstimulated cells and no ISO-induced endosomal emptying, suggesting

trafficking alterations (Fig. 4b). Supporting this idea, EEA1-positive early endosomes were

discernibly and quantifiably larger in old myocytes than in young (Fig. 4a and c).

Recycling deficiencies extended into Rab11-positive recycling endosomes where a similar

age-dependent absence of ISO-induced emptying of their $Ca_V1.2$ cargo was observed in Rab11

and $Ca_V1.2$ double-stained myocytes (Fig. 4d-e). Recycling endosomes were also found to be

significantly enlarged in old myocytes compared to young (Fig. 4f). Rab7 positive late endosomes

did not display any age-dependent alterations in their size or $Ca_V1.2$ cargo suggesting that the

degradation pathway may be intact (Supplementary Fig. 2).

BIN1 is overexpressed and mislocalized in old myocytes

Endosomal enlargement has been linked to endosomal dysfunction and deficiencies in membrane

protein recycling in Alzheimer's disease (AD)[23]. Endosome swelling results from an imbalance in

cargo coming into and leaving endosomes, sometimes referred to as endosomal traffic jams[24].

BIN1/amphiphysin II has been implicated as playing a role in the development of endosomal traffic

jams in AD-stricken neurons[24]. In the heart, BIN1 is better known for its role in targeted delivery

of $Ca_V1.2$[25,26], t-tubule biogenesis[27], micro-folding[28], and maintenance[29], and in dyad formation[30].

However, in neurons, where there are no t-tubules, BIN1 is known to play a role in mediating

endosomal membrane curvature and tubule formation required for cargo exit and recycling from

endosomes[31,32]. BIN1 overexpression has been linked to aging and neurodegeneration in the

brain[33-35] and has been seen to produce endosomal expansion and traffic jams[24,31,32].

Furthermore, BIN1 expression levels are known to affect ion channel trafficking in both neurons[36]

and ventricular myocytes[25,26]. We thus examined BIN1 expression and localization in young and

old myocytes and found an age-associated upregulation (Fig. 5a-b) and redistribution of BIN1

with old myocytes exhibiting a vesicular, endosomal pattern of BIN1-staining alongside the

expected t-tubule population (Fig. 5c; *left*).

BIN1 knockdown in old animals rejuvenates RyR2 plasticity and β-AR responsiveness

Since BIN1 became upregulated and mislocalized with aging with a more endosome-like pattern

of expression, we hypothesized that knocking down BIN1 to young levels might resolve the

endosomal traffic jams and restore β-AR responsivity to aging hearts. To test this, we transduced

live mice with AAV9-GFP-mBIN1-shRNA (shRNA-mBIN1) or AAV9-GFP-scramble-shRNA (shRNA-scrmb; control) via retro-orbital injection. Two weeks post-injection, hearts were harvested, and successful shRNA-mediated knockdown of BIN1 was confirmed via both immunostaining and western blot analysis. BIN1 knockdown restored youthful localization of BIN1 to the t-tubules (Fig. 5c; *right*), and reduced BIN1 expression levels in old hearts to young heart levels (Fig. 5d-e). Control shRNA-scrmb transduced old hearts had similar BIN1 expression levels to non-transduced old hearts (Fig. 5d-e) and in all experiments, cells isolated from shRNA-scrmb transduced hearts maintained an old phenotype (Fig. 5c, 5f-m, Supplementary Fig. 3, and Supplementary Table 1). Contrary to our hypothesis, I_{Ca} recordings from shRNA-mBIN1 transduced old myocytes did not indicate restoration of ISO responsivity (Fig. 5f-g, Supplementary Fig. 3a). In line with that finding, BIN1 knockdown did not reverse the endosomal swelling seen with aging suggesting that it is not an essential contributor to endosomal dysfunction in cardiomyocytes (Supplementary Fig. 4). However, basal Ca^{2+} transient amplitude in shRNA-mBIN1 transduced old myocytes resembled that of young myocytes and *β*-adrenergic responsiveness was fully restored to young levels (Fig. 5h-i, Supplementary Fig. 3b).

The recovery of Ca^{2+} transient amplitude suggested that BIN1 knockdown may predominantly impact RyR2 function and/or distribution and not $Ca_V1.2$. To investigate that possibility, we performed SMLM on shRNA-mBIN1 transduced myocytes finding the nanoscale distribution of $Ca_V1.2$ was unaltered compared to non-transduced old myocytes in that they remained basally super-clustered and did not recover the super-clustering response to ISO (Fig. 5j-k, and Supplementary Fig. 3c). In contrast, RyR2 appeared rejuvenated and exhibited a young cell-like nanoscale distribution and response to ISO (Fig. 5l-m, and Supplementary Fig. 3d). Overall, these results suggest that knockdown of BIN1 in aging hearts recovers cardiac *β*-adrenergic responsiveness by restoring RyR2 clustering plasticity.

BIN1 knockdown in old animals rescues systolic but not diastolic function

Since all our experiments were performed in isolated cells, the question remained, would

BIN1 knockdown improve old heart contractile function *in vivo*? To address this, we performed

echocardiography in young, old, and old shRNA-transduced mice. In conscious mice fractional

shortening (FS) and ejection fractions (EF) were reduced in old mice compared to young (Fig. 6a-

c) suggesting an age-dependent systolic dysfunction. To examine the effects of shRNA-scrmb or

shRNA-mBIN1 on old heart function, we performed echocardiograms on old mice to record their

initial function prior to treatment with shRNA (Fig. 6d-e; *left*). Those mice were then retro-orbitally

injected with either shRNA-scrmb (control) or shRNA-mBIN1 and after a two-week transduction

period, repeat echocardiograms were obtained (Fig. 6d-e; *right*). In agreement with the

rejuvenating effects of BIN1 knockdown observed in the *in vitro* functional analysis of Ca^{2+}

transients, basal ventricular function was restored to a more youthful phenotype upon BIN1

knockdown, with shRNA-mBIN1 treated old mice displaying improved FS and EF (Fig. 6f-g).

Old mice also exhibited signs of left ventricular (LV) hypertrophy including increased LV

mass (Supplementary Fig. 5a). Two weeks of BIN1 knockdown did not significantly improve LV

mass although there was a downward trajectory toward a more youthful LV mass in paired old

mice pre- and post-shRNA-mBIN1 that was not apparent in the shRNA-scrmb cohort

(Supplementary Fig. 5b). LV end diastolic volumes (EDV) and end systolic volumes (ESV) were

significantly increased in old mice compared to young (Supplementary Fig. 5e and g). The

resultant increase in preload in old mice would be expected to enhance ventricular output via the

Frank-Starling mechanism to maintain stroke volume in the face of reduced contractility. shRNA-

mBIN1 transduction in old mice significantly reduced EDV and ESV to youthful levels

(Supplementary Fig. 5f and h). Accordingly, cardiac output (CO) was significantly rejuvenated by

BIN1 knockdown in old mice, increasing from a mean of 14.82 mL/min pre-injection to 17.32

mL/min two weeks post-knockdown and approaching the 18.61 mL/min seen in young mice

(Supplementary Fig. 5i-j). The restoration of youthful CO, EDV, ESV, FS and EF by BIN1

knockdown suggests this treatment rejuvenates systolic function in old mice.

We further measured the *in vivo* effects of ISO on ventricular contractile function by

performing echocardiograms on anesthetized young and old mice before and after intraperitoneal

injection of ISO. As above, repeat echocardiograms were performed on old mice 2-weeks after

retro-orbital injection of shRNA-scrmb or shRNA-mBIN1 (Supplementary Figs. 6 and 7). As has

been previously reported, resting and ISO-stimulated heart rates were significantly lower in old

mice compared to young[37], but neither the scrambled shRNA nor the BIN1 knockdown had any

significant effect on heart rate suggesting that pacemaker activity is not affected by these

treatments (Supplementary Figs. 5c-d, 6a-b, and 7a-b). In agreement with the *in vivo* data from

conscious mice, basal ventricular contractile function was rejuvenated by BIN1 knockdown with

CO, FS, EF, EDV and ESV (Supplementary Figs. 6 and 7) all improving toward youthful levels.

Importantly this effect was not observed in shRNA-scrmb transduced mice suggesting a specific

effect of BIN1 knockdown.

To assess changes in diastolic function with aging, we performed pulsed wave Doppler

imaging on unconscious mice. Old mice displayed an enhanced isovolumetric relaxation time

(IVRT) and reduced mitral valve early to late ventricular filling velocity ratio (MV E/A), indicative

of diastolic dysfunction, which was not remedied with BIN1 knockdown (Fig. 7). Altogether, these

results suggest that BIN1 knockdown may represent a new therapeutic strategy to improve age-

related deficits in systolic function.

Age-dependent decrease in myofilament Ca^{2+} sensitivity is rescued by BIN1 knockdown

Ca^{2+} sensitivity of the myofilaments can also influence systolic and diastolic function thus we

examined the expression and phosphorylation state of two cardiac myofilament proteins that are

known to impact Ca^{2+} sensitivity in a phosphorylation-dependent manner, specifically cardiac

troponin-I (cTnI)[38,39] and cardiac myosin binding protein-C (cMyBP-C)[40,41]. We probed for these

myofilament proteins in heart lysates isolated from each cohort. We began with cTnI finding its

total expression increased with aging, but relative expression of the phosphorylated form

(pS23/24) reduced (Fig. 8a-c). While shRNA-mBIN1 treatment displayed a trend toward a more

youthful pS23/24 expression, the trend did not become significant. Total expression of cMyBP-C

was unaltered with aging but its phosphorylation at S273 was significantly decreased (Fig. 8d-f).

shRNA-mBIN1 treatment restored youthful cMyBP-C pS273 expression levels. Our results

support a model where reduced phosphorylation of cMyBP-C in old hearts contributes to deficits

in contractile function while reduced cTnI phosphorylation favors longer relaxation and diastolic

dysfunction. BIN1 knockdown appears to restore systolic but not diastolic function by improving

phosphorylation of cMyBP-C but not cTnI.

Discussion

We report six novel findings that provide insight into the mechanistic basis of age-related cardiac

dysfunction and reveal BIN1 as a new therapeutic target to rejuvenate the aging heart and extend

cardiovascular health into old age: 1) patch clamp and Ca^{2+} transient recordings revealed an age-

associated loss of cardiac β-adrenergic responsiveness measured by the functional augmentation

of $Ca_V1.2$ and RyR2 channel activity; 2) super-resolution imaging and PLA revealed an age-

associated dysregulation of $Ca_V1.2$ and RyR2 clustering plasticity, and juxta-positioning at dyads;

3) examination of endosomal pools and $Ca_V1.2$ dynamics revealed ISO-stimulated Ca^{2+} channel

mobility and recycling is impaired in aging due to trafficking deficits that cause endosomal traffic

jams; 4) western blots and super-resolution imaging revealed that BIN1, important for t-tubule

biogenesis[27], and $Ca_V1.2$, RyR2[18,25,26] and sarcomere[42] organization, is overexpressed in aging

hearts; 5) shRNA-mediated BIN1 knockdown to youthful levels rejuvenated RyR2 clustering

plasticity, basal Ca^{2+} transient amplitudes and their functional augmentation by β-AR-stimulation,

and myofilament protein phosphorylation; and finally 6) *in vivo* echocardiography and Doppler

imaging demonstrated that BIN1 knockdown in old mice restored youthful systolic (but not

diastolic) function and substantially reversed the aging phenotype. Our knockdown approach

using AAV9 packaged shRNA against BIN1 is rapid, occurring within 2 weeks of retro-orbital

injection, and has translational potential as AAVs are approved for use in humans[43].

Prior studies have shown that basal $Ca_V1.2$ activity and β-AR-mediated I_{Ca} enhancement

are altered with aging[44]. However, the effects on basal $Ca_V1.2$ currents appear to vary depending

on sex, species, and possibly even strain with various groups reporting enhanced[44-46],

unchanged[47], or reduced[48-50] currents in aged models. In our hands, myocytes from 24-month-old

C57BL/6J mice had enhanced basal $Ca_V1.2$ currents compared to young (3-month-old). Old

myocytes also exhibited basally enhanced Ca^{2+} transient amplitudes. Prior findings on Ca^{2+}

transient alterations in the aging myocardium range from basal amplitude augmentation[45,51] and

faster decay kinetics[52] to no or little reported change[50], to basal reduction in CaT amplitude and

slower rate of decay[46,53]. This echoes the variability in age-associated alterations of I_{Ca} and

reflects the complexity of aging. Importantly though, and in line with our results, there is agreement

that the ability of β-AR-signaling to augment Ca^{2+} transient amplitude and tune EC-coupling and

inotropy is diminished with aging[7,8,54-59].

This study provides the first evidence linking age-associated β-AR hypo-responsivity to

alterations in Ca^{2+} channel clustering and organization. Super-resolution microscopy studies of

the nanoscale arrangement of $Ca_V1.2$ and RyR2 have fueled the emerging concept that tunable

insertion, recycling, and clustering of these channels physiologically during receptor signaling

cascades or pathologically in disease, affects Ca^{2+} release efficiency and myocardial

contractility[15,16,19,20,60-62]. Larger $Ca_V1.2$ clusters facilitate enhanced $Ca_V1.2$-$Ca_V1.2$ physical

interactions and cooperative gating that amplifies I_{Ca}[16,17,63] while larger RyR2 clusters favor

enhanced Ca^{2+} spark frequency and larger Ca^{2+} transients[21]. We previously reported that acute

β-AR activation in young ventricular myocytes promotes rapid, dynamic augmentation of

sarcolemmal $Ca_V1.2$ channel abundance as channels are mobilized from endosomal reservoirs

via fast and slow recycling pathways[15,16]. Here, super-resolution microscopy and dynamic channel

tracking experiments demonstrated that this response is impaired with aging, such that $Ca_V1.2$

are basally super-clustered to a similar extent to that seen in young ISO-stimulated myocytes.

These results, correlated with the basally elevated I_{Ca} and impaired current augmentation with

ISO, hint that the endosomal reservoir of $Ca_V1.2$ cargo has already been expended and/or

endosomal recycling pathways are impaired. Our data suggest the latter is true, revealing reduced

ISO-induced insertions in old myocytes, an aggregation of channels on swollen endosomes, and

a larger pool of static channels that appear stranded in the sarcolemma. Whether this pool of

static channels reflects increased lifetime of $Ca_V1.2$ in the membrane remains to be established.

The main conclusion from this data is that the endosomal pathway and homeostatic regulation of

$Ca_V1.2$ channel expression at the sarcolemma is impaired with aging.

Deficiencies in recycling and endosomal enlargement are familiar sights in Alzheimer's

disease (AD) where endosomal enlargement is the first neuro-cytopathological hallmark of AD,

visible even before neurofibrillary tangles or amyloid β plaques[23]. An imbalance in cargo trafficking

into and out of endosomes is thought to result in endosome swelling and recycling pathway

deficiencies termed endosomal traffic jams[24]. As mentioned earlier, alterations in BIN1 expression

are thought to contribute to endosomal dysregulation in AD neurons. Thus, when we found BIN1

expression was significantly increased in aging, and appeared to assume a more endosomal

localization pattern, we thought, since it was in the "right place at the right time" that knockdown

of BIN1 would restore youthful endosomal trafficking and improve $Ca_V1.2$ channel cargo

mobilization, recovering $β$-AR responsivity and $Ca_V1.2$ channel functional augmentation. That

hypothesis was thoroughly disproven with results showing BIN1 knockdown did not in fact

accomplish any of those things. Several other proteins have been implicated as sources or

contributors to endosomal dysfunction and swelling in AD neurons as reviewed[24]. Future studies

should investigate whether any of those proteins underlie age-associated endosomal dysfunction

and impaired $Ca_V1.2$ recycling in ventricular myocytes.

On the other side of the dyadic cleft, RyR2 also display enlarged basal clusters that fail to

undergo augmentation with ISO in aging myocytes. Other groups have reported phosphorylation-

stimulated clustering of RyR2[18,19] but this is the first report of disruption of this dynamic response

with aging. Furthermore, it is the first to find that spatial proximity between $Ca_V1.2$ and RyR2

increases with acute ISO in young myocytes. This suggests rapid dynamic formation of new dyads

or expansion of existing dyads during ISO-stimulation. The implication here is that $β$-AR activation

stimulates parallel $Ca_V1.2$ and RyR2 cluster expansion on either side of the dyadic cleft to

maximize the efficiency of Ca^{2+} release but that this additional tuning capacity or reserve is lost

with aging and is apparently already expended at rest. Cardiac RyR2 trafficking and mobility is

poorly understood and understudied but one report has linked RyR2 redistribution upon

phosphorylation to BIN1, evidenced by reduced recruitment of phosphorylated RyR2 to dyadic

regions in cardiac-specific *BIN1* knockout cardiomyocytes[18]. Interestingly, BIN1 knockdown in old

mice had a more profound effect on RyR2 distribution, Ca^{2+} transient amplitude, and inotropy than

it had on $Ca_V1.2$. To that end, RyR2 nanoscale distribution was rejuvenated by BIN1 knockdown

in old hearts but $Ca_V1.2$ still exhibited the old super-clustered phenotype and heightened basal

I_{Ca}. So, while our initial rationale for knocking down BIN1 in the aging heart BIN1 was flawed,

serendipitously we discovered that BIN1 knockdown has a rejuvenating effect on RyR2.

Echocardiography confirmed the benefits of BIN1 knockdown on the aging heart with

parameters of systolic function including EF, FS, and CO all substantially improved after just a

two-week knockdown of BIN1. EDV and ESV also declined to youthful levels reflecting the

increased contractile function. In humans, resting systolic function is generally preserved with

aging, however it becomes significantly impaired during exercise and acute stress[3,64,65]. The

resting systolic dysfunction observed in the current study may reflect the difference between

human and mouse heart's basal autonomic balance at room temperature. Mice have relatively

more sympathetic nervous system activity at rest than humans. With aging, autonomic balance

becomes even more tilted toward sympathetic activity as parasympathetic input appears

reduced[66,67]. In this way, a resting mouse heart is not necessarily a fair comparison to a resting

human heart and may instead be a better reflection of an exercising human heart. Thus, BIN1

knockdown may present a therapeutic option to enhance systolic function in aging humans during

acute exercise and stress.

Diastolic dysfunction is a more universal finding with aging in both humans and mice[4,68]

and our findings of a prolonged IVRT with aging and reduced E/A ratio agree with prior studies

reporting impaired relaxation. However, BIN1 knockdown did not remedy diastolic dysfunction.

This is perhaps expected as enhanced fibrosis[68] with aging is one of the main causes of the

enhanced stiffness that underlies diastolic issues in the aging heart, and there is no reason to

suspect that BIN1 knockdown would improve fibrosis.

Reduced Ca^{2+} sensitivity of the myofilaments can also contribute to diastolic dysfunction.

Enhanced phosphorylation of the thin myofilament protein cTnI by PKA at ser-23 and -24

accelerates myocardial relaxation by reducing myofilament Ca^{2+} sensitivity, favoring Ca^{2+}

dissociation from troponin-C, and leading to faster actin-myosin detachment[38,39]. Our results

indicate that aging hearts display significantly lower levels of cTnI phosphorylation favoring slower

relaxation. This deficit was not improved or corrected by BIN1 knockdown and diastolic

dysfunction prevailed. Given the reciprocal relationship between cTnI phosphorylation and

myofilament Ca^{2+} sensitivity, one would predict that the age-associated reduction in cTnI pS23/24

levels would result in increased Ca^{2+} sensitivity in old mice compared to young, leading to

increased Ca^{2+}-dependent force generation. However, our *in vitro* finding of larger amplitude Ca^{2+}

transients in old myocytes accompanied by reduced *in vivo* contractile function in old mice

suggests quite the opposite. Specifically, our functional results suggest an age-associated

reduction in the Ca^{2+} sensitivity of the myofilaments that is remedied by BIN1 knockdown. Thus,

we broadened our search to examine cMyBP-C, another myofilament protein. Decreased

phosphorylation of cMyBP-C causes decreases myofilament Ca^{2+} sensitivity[41] and contractile

dysfunction in heart failure patients[40]. Furthermore, in the dephosphorylated state, cMyBP-c

inhibits actin-myosin interactions leading to reduced force development[41]. We found old mouse

hearts had lower cMyBP-C pS273[69] expression levels than young hearts. A similar age-

associated decline in cMyBP-C phosphorylation has been reported by others and found to

correlate with deterioration of cardiac function[70]. Furthermore, phosphorylation-mimetic cMyBP-

C mutant mice exhibit preserved cardiac function and enhanced longevity suggesting

phosphorylation of cMyBP-C is cardioprotective[70]. In a novel finding, we report that BIN1

knockdown rescues cMyBP-C pS273 phosphorylation levels which potentially explains the

improved contractile function and Ca^{2+} sensitivity in shRNA-mBIN1 transduced mice. But how can

one link BIN1 expression levels to sarcomeric proteins? While we do not yet have the full

mechanistic picture of how this occurs, there is evidence that BIN1 binds to and influences

sarcomeric proteins. For example, in skeletal muscle, BIN1 is said to act as an essential adaptor

protein and scaffold that promotes mature sarcomere assembly[42]. Increased BIN1 expression is

also known to promote enhanced expression of myosin heavy chain[71] and to associate with and

stabilize actin filaments[72]. Future studies should examine the mechanism of BIN1 regulatory

influences over cMyBP-C but it is an intriguing idea first proposed by Fernando et al in 2009[42],

that a single protein (BIN1) could orchestrate sarcolemmal and sarcomeric architecture and

regulation.

It is important to mention that although our study focused on the effects of BIN1

upregulation with aging, prior work has revealed that BIN1 *downregulation* is associated with heart

failure[26,29] and that exogenous supplementation of a cardiac specific isoform of BIN1 can improve

cardiac function in mice with pressure overload-induced heart failure[73]. It is possible that BIN1

levels must be tightly regulated and that any deviation from a homeostatic normal level could be

damaging. In Alzheimer's disease such a model has been presented where too much or too little

BIN1 expression are both associated with enhanced risk of disease[35,74].

In summary, we find that BIN1 knockdown restores youthful Ca^{2+} transient amplitude, *β-*

AR mediated transient augmentation, and *in vivo* systolic function in the aging myocardium. This

study sets the groundwork for future work and raises the hope that a similar approach could

improve cardiac function, restore exercise tolerance, and increase capacity to endure acute stress

in elderly patients.

Acknowledgments

We extend our thanks to Dr. Sakthivel Sadayappan for the generous gift of the total cMyBP-C

antibody and the cMyBP-C Ser-273 phospho-site specific antibody[69]. We thank Dr. L. Fernando

Santana and Heather Spooner for reading and providing critical suggestions on our manuscript.

Sources of Funding

This work was supported by NIH R01AG063796 and R01HL159304 to R.E.D.; R01GM127513

and RF1NS131379 to E.J.D.; by the NIGMS funded Pharmacology Training Program T32

GM099608 and later by an AHA Predoctoral Fellowship 827909 to T.L.V., by a Postdoctoral

Fellowship from NIH T32 Training Grant in Basic & Translational Cardiovascular Science T32

HL086350 and NIH F32 HL149288 to P.N.T.; and by NIH R01HL085727, R01HL085844,

R01HL137228, S10OD010389 Core Equipment Grant, and VA Merit Review Grant I01 BX000576

and I01 CX001490 to N.C. Graphical Abstract was generated using Biorender.com.

Disclosures

None.

References

1   Roth, G. A. *et al.* Global Burden of Cardiovascular Diseases and Risk Factors, 1990-2019:
Update From the GBD 2019 Study. *Journal of the American College of Cardiology* **76**, 2982-
3021, doi:10.1016/j.jacc.2020.11.010 (2020).

2   Dhingra, R. & Vasan, R. S. Age as a risk factor. *Med Clin North Am* **96**, 87-91,
doi:10.1016/j.mcna.2011.11.003 (2012).

3   Dai, D. F., Chen, T., Johnson, S. C., Szeto, H. & Rabinovitch, P. S. Cardiac aging: from
molecular mechanisms to significance in human health and disease. *Antioxid Redox Signal*
16, 1492-1526, doi:10.1089/ars.2011.4179 (2012).

4   Strait, J. B. & Lakatta, E. G. Aging-Associated Cardiovascular Changes and Their
Relationship to Heart Failure. *Heart Failure Clinics* **8**, 143-164, doi:10.1016/j.hfc.2011.08.011
(2012).

5   Lakatta, E. G. & Sollott, S. J. Perspectives on mammalian cardiovascular aging: humans to
molecules. *Comp Biochem Physiol A Mol Integr Physiol* **132**, 699-721 (2002).

6   Ferrara, N. *et al.* beta-adrenergic receptor responsiveness in aging heart and clinical
implications. *Front Physiol* **4**, 396, doi:10.3389/fphys.2013.00396 (2014).

7   Xiao, R. P. *et al.* Age-associated reductions in cardiac beta1- and beta2-adrenergic responses
without changes in inhibitory G proteins or receptor kinases. *J Clin Invest* **101**, 1273-1282
(1998).

8   Xiao, R. P., Spurgeon, H. A., O'Connor, F. & Lakatta, E. G. Age-associated changes in beta-
adrenergic modulation on rat cardiac excitation-contraction coupling. *J Clin Invest* **94**, 2051-
2059 (1994).

9   Osterrieder, W. *et al.* Injection of subunits of cyclic AMP-dependent protein kinase into cardiac
myocytes modulates Ca2+ current. *Nature* **298**, 576-578 (1982).

10 Hartzell, H. C., Mery, P. F., Fischmeister, R. & Szabo, G. Sympathetic regulation of cardiac
calcium current is due exclusively to cAMP-dependent phosphorylation. *Nature* **351**, 573-576,
doi:10.1038/351573a0 (1991).

11 Liu, G. *et al.* Mechanism of adrenergic CaV1.2 stimulation revealed by proximity proteomics.
Nature **577**, 695-700, doi:10.1038/s41586-020-1947-z (2020).

12 Manning, J. R. *et al.* Rad GTPase deletion increases L-type calcium channel current leading
to increased cardiac contraction. *J Am Heart Assoc* **2**, e000459,
doi:10.1161/JAHA.113.000459 (2013).

13 Yue, D. T., Herzig, S. & Marban, E. Beta-adrenergic stimulation of calcium channels occurs
by potentiation of high-activity gating modes. *Proc Natl Acad Sci U S A* **87**, 753-757. (1990).

14 Bean, B. P., Nowycky, M. C. & Tsien, R. W. Beta-adrenergic modulation of calcium channels
in frog ventricular heart cells. *Nature* **307**, 371-375 (1984).

15 Del Villar, S. G. *et al.* beta-Adrenergic control of sarcolemmal CaV1.2 abundance by small
GTPase Rab proteins. *Proc Natl Acad Sci U S A* **118**, doi:10.1073/pnas.2017937118 (2021).

16 Ito, D. W. *et al.* beta-adrenergic-mediated dynamic augmentation of sarcolemmal CaV 1.2
clustering and co-operativity in ventricular myocytes. *J Physiol* **597**, 2139-2162,
doi:10.1113/JP277283 (2019).

17 Dixon, R. E., Navedo, M. F., Binder, M. D. & Santana, L. F. Mechanisms and Physiological
Implications of Cooperative Gating of Ion Channels Clusters. *Physiological reviews*,
doi:10.1152/physrev.00022.2021 (2021).

18 Fu, Y. *et al.* Isoproterenol Promotes Rapid Ryanodine Receptor Movement to Bridging
Integrator 1 (BIN1)-Organized Dyads. *Circulation* **133**, 388-397,
doi:10.1161/CIRCULATIONAHA.115.018535 (2016).

19 Asghari, P. *et al.* Cardiac ryanodine receptor distribution is dynamic and changed by auxiliary
proteins and post-translational modification. *Elife* **9**, doi:10.7554/eLife.51602 (2020).

20 Dixon, R. E. Nanoscale Organization, Regulation, and Dynamic Reorganization of Cardiac
Calcium Channels. *Front Physiol* **12**, 810408, doi:10.3389/fphys.2021.810408 (2021).

21 Galice, S., Xie, Y., Yang, Y., Sato, D. & Bers, D. M. Size Matters: Ryanodine Receptor Cluster
Size Affects Arrhythmogenic Sarcoplasmic Reticulum Calcium Release. *J Am Heart Assoc* **7**,
doi:10.1161/JAHA.118.008724 (2018).

22 Fredriksson, S. *et al.* Protein detection using proximity-dependent DNA ligation assays.
Nature biotechnology **20**, 473-477, doi:10.1038/nbt0502-473 (2002).

23 Cataldo, A. M. *et al.* Endocytic pathway abnormalities precede amyloid beta deposition in
sporadic Alzheimer's disease and Down syndrome: differential effects of APOE genotype and
presenilin mutations. *Am J Pathol* **157**, 277-286, doi:10.1016/s0002-9440(10)64538-5 (2000).

24 Small, S. A., Simoes-Spassov, S., Mayeux, R. & Petsko, G. A. Endosomal Traffic Jams
Represent a Pathogenic Hub and Therapeutic Target in Alzheimer's Disease. *Trends
Neurosci* **40**, 592-602, doi:10.1016/j.tins.2017.08.003 (2017).

25 Hong, T. T. *et al.* BIN1 localizes the L-type calcium channel to cardiac T-tubules. *PLoS Biol*
8, e1000312, doi:10.1371/journal.pbio.1000312 (2010).

26 Hong, T. T. *et al.* BIN1 is reduced and Cav1.2 trafficking is impaired in human failing
cardiomyocytes. *Heart Rhythm* **9**, 812-820, doi:10.1016/j.hrthm.2011.11.055 (2012).

27 Lee, E. *et al.* Amphiphysin 2 (Bin1) and T-tubule biogenesis in muscle. *Science* **297**, 1193-
1196, doi:10.1126/science.1071362 (2002).

28 Hong, T. *et al.* Cardiac BIN1 folds T-tubule membrane, controlling ion flux and limiting
arrhythmia. *Nature medicine* **20**, 624-632, doi:10.1038/nm.3543 (2014).

29 Caldwell, J. L. *et al.* Dependence of Cardiac Transverse Tubules on the BAR Domain Protein
Amphiphysin II (BIN-1). *Circulation Research* **115**, 986-996,
doi:10.1161/circresaha.116.303448 (2014).

30 De La Mata, A. *et al.* BIN1 Induces the Formation of T-Tubules and Adult-Like Ca(2+) Release
Units in Developing Cardiomyocytes. *Stem cells (Dayton, Ohio)* **37**, 54-64,
doi:10.1002/stem.2927 (2019).

31 Pant, S. *et al.* AMPH-1/Amphiphysin/Bin1 functions with RME-1/Ehd1 in endocytic recycling.
Nat Cell Biol **11**, 1399-1410, doi:10.1038/ncb1986 (2009).

32 Lambert, E. *et al.* The Alzheimer susceptibility gene BIN1 induces isoform-dependent
neurotoxicity through early endosome defects. *Acta Neuropathol Commun* **10**, 4,
doi:10.1186/s40478-021-01285-5 (2022).

33 Tan, M. S., Yu, J. T. & Tan, L. Bridging integrator 1 (BIN1): form, function, and Alzheimer's
disease. *Trends in molecular medicine* **19**, 594-603, doi:10.1016/j.molmed.2013.06.004
(2013).

34 Wang, H. F. *et al.* Bridging Integrator 1 (BIN1) Genotypes Mediate Alzheimer's Disease Risk
by Altering Neuronal Degeneration. *J Alzheimers Dis* **52**, 179-190, doi:10.3233/JAD-150972
(2016).

35 Chapuis, J. *et al.* Increased expression of BIN1 mediates Alzheimer genetic risk by modulating
tau pathology. *Mol Psychiatry* **18**, 1225-1234, doi:10.1038/mp.2013.1 (2013).

36 Schurmann, B. *et al.* A novel role for the late-onset Alzheimer's disease (LOAD)-associated
protein Bin1 in regulating postsynaptic trafficking and glutamatergic signaling. *Mol Psychiatry*
25, 2000-2016, doi:10.1038/s41380-019-0407-3 (2020).

37 Choi, S., Baudot, M., Vivas, O. & Moreno, C. M. Slowing down as we age: aging of the cardiac
pacemaker's neural control. *Geroscience*, doi:10.1007/s11357-021-00420-3 (2021).

38 Pi, Y., Zhang, D., Kemnitz, K. R., Wang, H. & Walker, J. W. Protein kinase C and A sites on
troponin I regulate myofilament Ca2+ sensitivity and ATPase activity in the mouse
myocardium. *J Physiol* **552**, 845-857, doi:10.1113/jphysiol.2003.045260 (2003).

39 Kentish, J. C. *et al.* Phosphorylation of troponin I by protein kinase A accelerates relaxation
and crossbridge cycle kinetics in mouse ventricular muscle. *Circ Res* **88**, 1059-1065,
doi:10.1161/hh1001.091640 (2001).

40 El-Armouche, A. *et al.* Decreased phosphorylation levels of cardiac myosin-binding protein-C
in human and experimental heart failure. *J Mol Cell Cardiol* **43**, 223-229,
doi:10.1016/j.yjmcc.2007.05.003 (2007).

41 Kumar, M., Haghighi, K., Kranias, E. G. & Sadayappan, S. Phosphorylation of cardiac myosin-
binding protein-C contributes to calcium homeostasis. *J Biol Chem* **295**, 11275-11291,
doi:10.1074/jbc.RA120.013296 (2020).

42 Fernando, P. *et al.* Bin1 SRC homology 3 domain acts as a scaffold for myofiber sarcomere
assembly. *J Biol Chem* **284**, 27674-27686, doi:10.1074/jbc.M109.029538 (2009).

43 High, K. A. & Aubourg, P. rAAV Human Trial Experience. *Adeno-Associated Virus: Methods*
and Protocols **807**, 429-457, doi:10.1007/978-1-61779-370-7_18 (2011).

44 Josephson, I. R., Guia, A., Stern, M. D. & Lakatta, E. G. Alterations in properties of L-type Ca
channels in aging rat heart. *J Mol Cell Cardiol* **34**, 297-308, doi:10.1006/jmcc.2001.1512
(2002).

45 Dibb, K. M., Rueckschloss, U., Eisner, D. A., Isenberg, G. & Trafford, A. W. Mechanisms
underlying enhanced cardiac excitation contraction coupling observed in the senescent sheep
myocardium. *J Mol Cell Cardiol* **37**, 1171-1181, doi:10.1016/j.yjmcc.2004.09.005 (2004).

46 Isenberg, G., Borschke, B. & Rueckschloss, U. Ca2+ transients of cardiomyocytes from
senescent mice peak late and decay slowly. *Cell Calcium* **34**, 271-280, doi:10.1016/s0143-
4160(03)00121-0 (2003).

47 Walker, K. E., Lakatta, E. G. & Houser, S. R. Age associated changes in membrane currents
in rat ventricular myocytes. *Cardiovascular research* **27**, 1968-1977,
doi:10.1093/cvr/27.11.1968 (1993).

48 Grandy, S. A. & Howlett, S. E. Cardiac excitation-contraction coupling is altered in myocytes
from aged male mice but not in cells from aged female mice. *Am J Physiol Heart Circ Physiol*
291, H2362-2370, doi:10.1152/ajpheart.00070.2006 (2006).

49 Liu, S. J., Wyeth, R. P., Melchert, R. B. & Kennedy, R. H. Aging-associated changes in whole
cell K(+) and L-type Ca(2+) currents in rat ventricular myocytes. *Am J Physiol Heart Circ*
Physiol **279**, H889-900, doi:10.1152/ajpheart.2000.279.3.H889 (2000).

50 Kong, C. H. T. *et al.* The Effects of Aging on the Regulation of T-Tubular ICa by Caveolin in
Mouse Ventricular Myocytes. *J Gerontol A Biol Sci Med Sci* **73**, 711-719,
doi:10.1093/gerona/glx242 (2018).

51 Rueckschloss, U., Villmow, M. & Klockner, U. NADPH oxidase-derived superoxide impairs
calcium transients and contraction in aged murine ventricular myocytes. *Exp Gerontol* **45**,
788-796, doi:10.1016/j.exger.2010.05.002 (2010).

52 Francis Stuart, S. D. *et al.* Age-related changes in cardiac electrophysiology and calcium
handling in response to sympathetic nerve stimulation. *J Physiol* **596**, 3977-3991,
doi:10.1113/JP276396 (2018).

53 Janczewski, A. M., Spurgeon, H. A. & Lakatta, E. G. Action potential prolongation in cardiac
myocytes of old rats is an adaptation to sustain youthful intracellular Ca2+ regulation. *J Mol*
Cell Cardiol **34**, 641-648, doi:10.1006/jmcc.2002.2004 (2002).

54 White, M. *et al.* Age-related changes in beta-adrenergic neuroeffector systems in the human
heart. *Circulation* **90**, 1225-1238 (1994).

55 Lakatta, E. G. Deficient Neuroendocrine Regulation of the Cardiovascular-System with
Advancing Age in Healthy Humans. *Circulation* **87**, 631-636 (1993).

56 Stratton, J. R. *et al.* Differences in cardiovascular responses to isoproterenol in relation to age
and exercise training in healthy men. *Circulation* **86**, 504-512 (1992).

57 Davies, C. H., Ferrara, N. & Harding, S. E. Beta-adrenoceptor function changes with age of
subject in myocytes from non-failing human ventricle. *Cardiovascular research* **31**, 152-156
(1996).

58 Lakatta, E. G., Gerstenblith, G., Angell, C. S., Shock, N. W. & Weisfeldt, M. L. Diminished
inotropic response of aged myocardium to catecholamines. *Circ Res* **36**, 262-269 (1975).

59 Cerbai, E. *et al.* Beta-adrenoceptor subtypes in young and old rat ventricular myocytes: a
combined patch-clamp and binding study. *Br J Pharmacol* **116**, 1835-1842 (1995).

60 MacQuaide, N. *et al.* Ryanodine receptor cluster fragmentation and redistribution in persistent
atrial fibrillation enhance calcium release. *Cardiovascular research* **108**, 387-398,
doi:10.1093/cvr/cvv231 (2015).

61 Kolstad, T. R. *et al.* Ryanodine receptor dispersion disrupts Ca(2+) release in failing cardiac
myocytes. *Elife* **7**, doi:10.7554/eLife.39427 (2018).

62 Hou, Y. *et al.* Nanoscale Organisation of Ryanodine Receptors and Junctophilin-2 in the
Failing Human Heart. *Front Physiol* **12**, 724372, doi:10.3389/fphys.2021.724372 (2021).

63 Dixon, R. E., Yuan, C., Cheng, E. P., Navedo, M. F. & Santana, L. F. Ca2+ signaling
amplification by oligomerization of L-type Cav1.2 channels. *Proc Natl Acad Sci U S A* **109**,
1749-1754, doi:10.1073/pnas.1116731109 (2012).

64 Schulman, S. P. *et al.* Age-related decline in left ventricular filling at rest and exercise. *Am J*
Physiol **263**, H1932-1938, doi:10.1152/ajpheart.1992.263.6.H1932 (1992).

65 Fleg, J. L. & Lakatta, E. G. in *Cardiovascular disease in the elderly*   21-64 (CRC Press,
2008).

66 Comelli, M. *et al.* Rhythm dynamics of the aging heart: an experimental study using conscious,
restrained   mice.   *Am   J   Physiol   Heart   Circ   Physiol*   **319**,   H893-H905,
doi:10.1152/ajpheart.00379.2020 (2020).

67 Piantoni, C. *et al.* Age-Related Changes in Cardiac Autonomic Modulation and Heart Rate
Variability in Mice. *Front Neurosci* **15**, 617698, doi:10.3389/fnins.2021.617698 (2021).

68 Dai, D. F. *et al.* Overexpression of catalase targeted to mitochondria attenuates murine
cardiac   aging.   *Circulation*   **119**,   2789-2797,   doi:CIRCULATIONAHA.108.822403
[pii]10.1161/CIRCULATIONAHA.108.822403 (2009).

69 Copeland, O. *et al.* Analysis of cardiac myosin binding protein-C phosphorylation in human
heart muscle. *J Mol Cell Cardiol* **49**, 1003-1011, doi:10.1016/j.yjmcc.2010.09.007 (2010).

70 Rosas, P. C. *et al.* Cardiac Myosin Binding Protein-C Phosphorylation Mitigates Age-Related
Cardiac Dysfunction: Hope for Better Aging? *JACC Basic Transl Sci* **4**, 817-830,
doi:10.1016/j.jacbts.2019.06.003 (2019).

71 Wechsler-Reya, R. J., Elliott, K. J. & Prendergast, G. C. A role for the putative tumor
suppressor Bin1 in muscle cell differentiation. *Molecular and cellular biology* **18**, 566-575,
doi:10.1128/MCB.18.1.566 (1998).

72 Drager, N. M. *et al.* Bin1 directly remodels actin dynamics through its BAR domain. *Embo*
Reports **18**, 2051-2066, doi:10.15252/embr.201744137 (2017).

73 Li, J., Agvanian, S., Zhou, K., Shaw, R. M. & Hong, T. Exogenous Cardiac Bridging Integrator
1 Benefits Mouse Hearts With Pre-existing Pressure Overload-Induced Heart Failure. *Front*
Physiol **11**, 708, doi:10.3389/fphys.2020.00708 (2020).

74 Calafate, S., Flavin, W., Verstreken, P. & Moechars, D. Loss of Bin1 Promotes the
Propagation of Tau Pathology. *Cell Rep* **17**, 931-940, doi:10.1016/j.celrep.2016.09.063
(2016).

Figures and Figure Legends

Figure 1. *β*-AR stimulated augmentation of I_{Ca} and Ca²⁺ transients is diminished in aging. **a**, representative whole-cell currents elicited from young and old ventricular myocytes before (control; black) and during application of ISO (blue). **b**, dot-plots showing the fold change in peak current with ISO in young (N = 6, n = 8) and old (N = 5, n = 11) myocytes. **c**, dot-plots showing peak I_{Ca} before and after ISO. **d**, representative Ca²⁺ transients recorded before and after ISO from paced young (N = 6, n = 10) and old (N = 5, n = 11) myocytes. **e**, dot-plots showing the fold-increase in Ca²⁺ transient amplitude after ISO and **f**, Ca²⁺ transient amplitude before and after ISO. Unpaired Student's t-tests were performed on data sets displayed in b and e. Two-way ANOVAs with post-hoc Tukey's multiple comparison tests were performed on data displayed in c and f.

Figure 2. Basal super-clustering, impaired β-AR responsiveness, and enhanced proximity of Ca$_V$1.2 and RyR2 in aged myocytes. **a**, SMLM localization maps showing Ca$_V$1.2 channel localization and distribution in the t-tubules of young and old ventricular myocytes with or without ISO-stimulation. Yellow boxes indicate the location of the regions of interest magnified in the top right of each image. **b**, dot-plots showing mean Ca$_V$1.2 channel cluster areas in young (control: N = 3, n = 9; ISO: N = 3, n = 9) and old (control: N = 4, n = 11; ISO: N = 4, n = 10) myocytes. **c** and **d**, show the same for RyR2 in young (control: N = 3, n = 17; ISO: N = 3, n = 11) and old (control: N = 3, n = 12; ISO: N = 3, n = 11) myocytes. **e**, Representative fluorescence PLA (red)/DAPI (blue) images of myocytes with and without ISO-stimulation. **f**, dot-plot summarizing the cellular density of PLA fluorescent puncta normalized to the density in young control cells (young control: N = 3, n = 19; young ISO: N = 3, n = 16; old control: N = 3, n = 14; old ISO: N = 3, n = 16). Statistical analyses on data summarized in b, d, and f were performed using two-way ANOVAs with post-hoc Tukey's multiple comparison tests.

Figure 3. Dynamic TIRF imaging reveals age-associated trafficking deficits of Ca$_V$1.2.

a and **b**, Representative TIRF images of transduced Ca$_V\beta_{2a}$-paGFP young (**a**) and old (**b**) ventricular myocytes before (*top*) and after ISO (*bottom*). Channel populations that were inserted (red), endocytosed (blue), or static (white) during the ISO treatment are represented to the *right*. **c-e**, dot-plots summarizing the quantification of inserted (**c**), endocytosed (**d**), and static (**e**) Ca$_V\beta_{2a}$-paGFP populations. **f**, is a compilation of the data in c, d, and e to show the relative % of the channel pool that is inserted, endocytosed, and static. **g**, is a summary plot showing the net % change of plasma membrane Ca$_V\beta_{2a}$-paGFP after ISO for each cohort of young (*N* = 3, *n* = 10) and old (*N* = 3, *n* = 10) myocytes. Data were analyzed for statistical analysis using unpaired Student's t-tests.

Figure 4. Aging impairs β-AR-stimulated $Ca_V1.2$ recycling. a, Airyscan super-resolution images of $Ca_V1.2$ and EEA1 immuno-stained myocytes with and without ISO. *Bottom:* Binary colocalization maps show pixels in which $Ca_V1.2$ and EEA1 completely overlapped. **b**, dot-plots summarizing % colocalization between EEA1 and $Ca_V1.2$, and **c**, EEA1 positive endosome areas in young (control: *N* = 3, *n* = 16; ISO: *N* = 3, *n* = 16) and old (control: *N* = 3, *n* = 14; ISO: *N* = 3, *n* = 14) myocytes. **d-f,** same layout for myocytes co-stained for $Ca_V1.2$ and Rab11 recycling endosomes in young (control: *N* = 3, *n* = 13; ISO: *N* = 3, *n* = 13) and old (control: *N* = 4, *n* =16; old ISO: *N* = 3, *n* = 10) myocytes. Data were analyzed using two-way ANOVAs with post-hoc Tukey's multiple comparison tests.

Figure 5. BIN1 knockdown restores β-AR augmentation of Ca^{2+} transients and RyR2 clustering dynamics. a, western blot of BIN1 expression in whole heart lysates from young and old mice. Ponceau was used for normalization. **b**, histogram showing normalized BIN1 levels relative to young male (N = 5, average of 1-2 replicates). **c**, representative Airyscan images of young, old, and old transduced myocytes immunostained against BIN1. **d**, western blot of BIN1 expression in whole heart lysates from: young, old, shRNA-mBIN1 or shRNA-scrmb transduced mice. Ponceau was used for normalization. **e**, histogram showing normalized BIN1 levels relative

to young (N = 3, average of 1 - 3 replicates). **f**, representative whole-cell currents from shRNA-

scrmb and shRNA-mBIN1 myocytes before and during ISO application. **g**, fold change in peak I_{Ca}

with ISO in young, old, shRNA-scrmb (N = 3, n = 7) and shRNA-mBIN1 (N = 3, n = 9) myocytes.

h, representative Ca^{2+} transients recorded from old shRNA-scrmb and shRNA-mBIN1 myocytes

before and after ISO. **i**, fold increase after ISO from young, old, shRNA-scrmb (N = 3, n = 12) and

shRNA-mBIN1 (N = 5, n = 13) myocytes. **j**, SMLM localization maps showing Ca_V1.2 channel

localization on t-tubules of myocytes from old shRNA-scrmb and shRNA-mBIN1, with or without

ISO-stimulation. Regions of interest are highlighted by yellow boxes. **h**, fold change in mean

Ca_V1.2 channel cluster area with ISO in the old shRNA-scrmb (control: N = 3, n = 14; ISO: N = 3,

n = 13) and shRNA-mBIN1 myocytes (control: N = 3, n = 12; ISO: N = 3, n = 11). **l** and **m**, show

the same layout for RyR2 immunostained old shRNA-scrmb (control: N = 3, n = 12; ISO: N = 3, n

= 13) and shRNA-mBIN1 myocytes (control: N = 3, n = 9; ISO: N = 3, n = 8). Old and young data

points in g, i, k and m are reproduced from data in Figs. 1b, 1e, 2b and 2d respectively. Statistical

analysis was performed on data in b using a two-way ANOVA, and data in e, g, i, k and m using

one-way ANOVAs.

Figure 6. BIN1 knockdown improves cardiac contractility in old mice. a, representative M-

mode echocardiogram images from conscious young and old mice. Summary dot-plots showing

b, fractional shortening and **c,** ejection fraction in young (N = 11) and old (N = 20) mice are shown.

d and **e,** representative M-mode echocardiogram images from conscious old mice before (*left*)

and two-weeks after RO-injection (*right*) of shRNA-scrmb (**d**) and shRNA-mBIN1 (**e**). Summary

dot-plots for **f,** fractional shortening and **g**, ejection fraction showing paired results before and

after RO-injection for shRNA-scrmb (N = 9) and shRNA-mBIN1 (N = 9) groups are displayed.

Unpaired Student's t-tests were performed on data displayed in b and c. Paired Student's t-tests

were performed on data displayed in f and g.

Figure 7. BIN1 knockdown does not recover age-related diastolic dysfunction. Representative pulsed-wave (PW) doppler images from unconscious young (**a**) and old mice before (left) and two-weeks after RO-injection (right) of shRNA-scrmb (**b**) and shRNA-mBIN1 (**c**). Summary dot-plots for young (N = 7) and old (N = 11) mice, and paired results before and after RO-injection of old mice with shRNA-scrmb (N = 6) and shRNA-mBIN1 (N = 5) for the following diastolic measurements are displayed: **d** and **e**, isovolumetric relaxation time (IVRT) and, **f** and **g,** mitral valve E/A ratio (MV E/A). Unpaired Student's t-tests were performed on data displayed in d and f. Paired Student's t-tests were performed on data displayed in e and g.

Figure 8. Myofilament Ca²⁺ sensitivity is altered in aging. a, western blot of total and pS23/24

cTnI expression in whole heart lysates from young, old, shRNA-mBIN1 or shRNA-scrmb

transduced mice. Ponceau was used for normalization. **b**, histogram showing normalized total

cTnI levels relative to young (*N* = 3, average of 3 replicates). **c**, histogram showing cTnI pS23/24

normalized to total cTnI relative to young (*N* = 3, average of 2-3 replicates). **d**, western blot of

total and pS273 cMyBP-C expression in whole heart lysates from young, old, shRNA-mBIN1 or

shRNA-scrmb transduced mice. Ponceau was used for normalization. **e**, histogram showing

normalized total cMyBP-C levels relative to young (*N* = 3, average of 2-5 replicates). **f**, histogram

showing cMyBP-C pS273 normalized to total cMyBP-C relative to young (*N* = 3, average of 2-5

replicates).