Physiologically based pharmacokinetic (PBPK) modeling of the role of CYP2D6 polymorphism for metabolic phenotyping with dextromethorphan

Jan Grzegorzewski [1*], **Janosch Brandhorst** [1] **and Matthias König** [1]

[1] *Institute for Theoretical Biology, Institute of Biology, Humboldt University, Berlin, Germany*

Correspondence*:
Jan Grzegorzewski
grzegorj@hu-berlin.de

ABSTRACT

The cytochrome P450 2D6 (CYP2D6) is a key xenobiotic-metabolizing enzyme involved in
the clearance of many drugs. Genetic polymorphisms in CYP2D6 contribute to the large inter-
individual variability in drug metabolism and could affect metabolic phenotyping of CYP2D6 probe
substances such as dextromethorphan (DXM). To study this question, we (i) established an
extensive pharmacokinetics dataset for DXM; and (ii) developed and validated a physiologically
based pharmacokinetic (PBPK) model of DXM and its metabolites dextrorphan (DXO) and
dextrorphan O-glucuronide (DXO-Glu) based on the data. Drug-gene interactions (DGI) were
introduced by accounting for changes in CYP2D6 enzyme kinetics depending on activity score
(AS), which in combination with AS for individual polymorphisms allowed us to model CYP2D6
gene variants. Variability in CYP3A4 and CYP2D6 activity was modeled based on in vitro data from
human liver microsomes. Model predictions are in very good agreement with pharmacokinetics
data for CYP2D6 polymorphisms, CYP2D6 activity as described by the AS system, and CYP2D6
metabolic phenotypes (UM, EM, IM, PM). The model was applied to investigate the genotype-
phenotype association and the role of CYP2D6 polymorphisms for metabolic phenotyping using
the urinary cumulative metabolic ratio (UCMR), DXM/(DXO+DXO-Glu). The effect of parameters
on UCMR was studied via sensitivity analysis. Model predictions indicate very good robustness
against the intervention protocol (i.e. application form, dosing amount, dissolution rate, and
sampling time) and good robustness against physiological variation. The model is capable of
estimating the UCMR dispersion within and across populations depending on activity scores.
Moreover, the distribution of UCMR and the risk of genotype-phenotype mismatch could be
estimated for populations with known CYP2D6 genotype frequencies. The model can be applied
for individual prediction of UCMR and metabolic phenotype based on CYP2D6 genotype. Both,
model and database are freely available for reuse.

Keywords: Dextromethorphan, CYP2D6, Physiologically based pharmacokinetic model, PBPK, Pharmacokinetics, Pharmacogenomics
27

1 INTRODUCTION

The cytochrome P450 (CYPs) superfamily of enzymes has a central role in the clearance of many
substances and drugs, with the isoform 2D6 (CYP2D6) being one of the most important xenobiotic-
metabolizing enzymes. CYP2D6 is involved in the clearance of around 20% of the most prescribed
drugs (Saravanakumar et al., 2019) including antiarrhythmics having a small therapeutic range (e.g.
flecainide, procainamide, mexiletine), anticancer agents (e.g. tamoxifen), antidepressants (e.g. citalopram,
fluoxetine, duloxetine: venlafaxine), antipsychotics (e.g. aripiprazole, haloperidol, thioridazine), β-blockers
(metoprolol), analgesics (tramadol, oxycodone, codeine), and antitussives (dextromethorphan) (Kibaly
et al., 2021; Hurtado et al., 2020). CYP2D6-mediated drug response exhibits a particularly large inter-
individual variability which poses a challenge for personalized dosage of medication by underdosing on
the one hand and toxic side effects on the other. The activity of CYP2D6 is known to be majorly dependent
on genetic variants (Preskorn et al., 2013; Berm et al., 2013; Shah and Smith, 2015) with polymorphism of
CYP2D6 being related to the risk of adverse effects, non-response during treatment, and death by drug
intoxication (Gasche et al., 2004; Kawanishi et al., 2004; Rau et al., 2004; Zackrisson et al., 2010).

In the late 70s, a polymorphism in debrisoquine hydroxylation (Mahgoub et al., 1977) and sparteine
oxidation (Eichelbaum et al., 1979) was discovered and subsequently attributed to allelic variants of
the CYP2D6 gene. In the following years, CYP2D6 became one of the most studied drug-metabolizing
enzymes. Genetic variants were classified into distinct phenotypes and subjects carrying combinations
of these variants were categorized as poor metabolizer (gPM), intermediate metabolizer (gIM), extensive
metabolizer (gEM), and ultra rapid metabolizer (gUM) (Zanger et al., 2004; Gaedigk et al., 2017). This
classification is based on the relationship between genetic variants and CYP2D6-mediated drug response.
For these genetically predicted phenotypes, we use the 'g' nomenclature as they can be easily confused
with the actual *in vivo* metabolic phenotype, determined based on pharmacokinetic measurements after
the administration of CYP2D6 test drugs. Nowadays, the CYP2D6 activity score (AS) system, a more
refined metric, is often applied to characterize genotype-phenotype associations (Gaedigk et al., 2018a).
In the system, discrete values between 0 and 1 are assigned to gene variants. The final activity score is
calculated by the sum of the activity scores of both alleles. For instance, a person with diplotype *1/*3
(the variant *1 has an AS of 1 and the variant *3 has no activity with an AS of 0) has an overall AS
of 1. Higher activity scores than 2 and additional complexity arise from copy number variation (CNV),
chimeras, and hybrids with the pseudo gene CYP2D7. This can result in ambiguities and difficulties in the
assignment of the correct diplotype and activity score (Gaedigk et al., 2007; Nofziger and Paulmichl, 2018;
Gaedigk et al., 2019). Of note, AS specifics are still under heavy debate and regularly updated (Caudle
et al., 2020). A multitude of population studies have been conducted to identify and associate allele variants
with metabolic phenotypes within and across populations (Gaedigk et al., 2017). Over 130 CYP2D6 star
(*) allele haplotypes have been identified and subsequently cataloged by the Pharmacogene Variation
(PharmVar) Consortium into PharmGKB with their respective activity score contribution (Gaedigk et al.,
2018b; Whirl-Carrillo et al., 2021).

Various methods exist for the metabolic phenotyping based on test substances. The gold standard is
plasma concentration sampling of probe substances and their metabolites at various time points after the
administration. (Partial) clearance rates and the relative enzyme activities can be calculated from these
plasma time profiles. Simplified methods have been established for many probe substances which do not
require repeated sampling of blood, e.g., the (cumulative) metabolic ratios between the probe substance and
one or several of its metabolites at a single time point in blood, plasma, or urine are utilized as such proxy
measures. Large-scale population studies often tend to employ urinary ratios of metabolites. Alternatively,

sampling of saliva and breath are worth considering (De Kesel et al., 2016). Probe substances for metabolic
phenotyping of CYP2D6 are debrisoquine, dextromethorphan, metoprolol, or sparteine (Frank et al.,
2007; Fuhr et al., 2007). Bufuralol is less popular but well suited for *in vitro* investigations due to its
fluorescent properties (Zanger et al., 2004). Although debrisoquine and sparteine have excellent properties
for CYP2D6 phenotyping, they have been withdrawn from clinical use in most countries and are therefore
no longer readily available. Frequently in use for the phenotyping of CYP2D6 activity are metoprolol and
dextromethorphan.

Dextromethorphan (DXM) is an over-the-counter, antitussive, non-narcotic, synthetic analog of codeine
affecting the activity of numerous channels and receptors in the brain that trigger the cough reflex (Silva
and Dinis-Oliveira, 2020). It is generally well-tolerated, considered safe in therapeutic dosage, and easily
available (Fuhr et al., 2007). Besides therapeutic purposes, DXM is most commonly applied as a probe
substance for CY2D6 phenotyping, alone or with other probe substances in a cocktail. DXM can be
administered orally and intravenously, has low bioavailability ($\approx$50%) and a rapid first-pass effect in the
intestine and liver. Typically only about half of the dose is recovered in urine over at least 12 hours after
administration, primarily as glucuronides (Schadel et al., 1995; Capon et al., 1996; Tennezé et al., 1999;
Strauch et al., 2009). In the systemic circulation, $\approx$55-65% of DXM is non-specifically bound to plasma
proteins (Lutz and Isoherranen, 2012; Taylor et al., 2016).

The biotransformation of DXM is mostly confined to the liver, where DXM is O-demethylated by
CYP2D6 to the active metabolite dextrorphan (DXO). Subsequently to O-demethylation, most of the DXO
is rapidly transformed via UDP-glucuronosyltransferase (UGT) to dextrorphan O-glucuronide (DXO-Glu)
and excreted via the urine. In individuals without any functional variant of CYP2D6, the metabolization
of DXM to DXO is extremely slow but still present. Apparently, the O-demethylation is not exclusively
mediated by CYP2D6, and it has been demonstrated *in vitro* that O-demethylation of DXM can be
marginally mediated by CYP3A4, CYP3A5 and CYP2C9 (von Moltke et al., 1998; McGinnity et al., 2000;
Takashima et al., 2005; Yu and Haining, 2001). In line with this observation, inhibition of CYP2D6, e.g.,
barely affects poor metabolizer (Pope et al., 2004). The second pathway of DXM metabolization goes
via N-demethylation to 3-methoxymorphinan which is mainly catalyzed via CYP3A4. Subsequently, 3-
methoxymorphinan and DXO are biotransformed to 3-hydroxymorphinan which is then rapidly transformed
via glucuronidation to hydroxymorphian O-glucuronide and excreted in the urine. The urinary cumulative
metabolic ratio (UCMR) of DXM to its metabolites DXM/(DXO+DXO-Glu) is a widely applied measure
for the *in vivo* CYP2D6 phenotyping.

An crucial question for metabolic phenotyping and liver function testing is how CYP2D6 polymorphisms
affect the pharmacokinetics of DXM and metabolic phenotyping based on DXM, such as the UCMR. The
objective of this work was to answer this question by the means of physiologically based pharmacokinetic
(PBPK) modeling of DXM.

2 MATERIAL AND METHODS

2.1 Pharmacokinetics database of DXM

Pharmacokinetics data of DXM was systematically curated from literature for model development,
parameterization, and validation. Curation efforts were mainly focused on concentration-time profiles
of DXM, DXO, and DXO-Glu in plasma or serum and their amounts or ratios in urine. The data is
accompanied by metadata on the investigated subjects and groups (e.g. CYP2D6 genotype or activity
score) and the applied intervention (e.g. dose and application form of DXM). All data was curated using

an established curation pipeline (Grzegorzewski et al., 2022) and is available via the pharmacokinetics
database PK-DB (https://pk-db.com) (Grzegorzewski et al., 2021). As a first step, a PubMed search
for the pharmacokinetics of dextromethorphan in combination with genotyping and/or phenotyping
was performed with the search query https://pubmed.ncbi.nlm.nih.gov/?term=dextromethorphan+
AND+%28phenotype+OR+phenotyping%29+AND+genotype. The literature corpus was extended with
drug cocktail studies from PK-DB (Grzegorzewski et al., 2022), secondary literature from references,
and results from PKPDAI with the search query https://app.pkpdai.com/?term=dextromethorphan
Gonzalez Hernandez et al. (2021). Data was selected and curated based on eligibility criteria, see below.
During the curation process, the initial corpus was updated by additional publications from the references
and citations. A subset of the studies only reported pharmacokinetic parameters without timecourses. These
studies were curated but not further used in the following analyses.

To be eligible, studies had to report *in vivo* pharmacokinetics data for adult (age $\geq$ 18) humans after
administration of DXM or DXM hydrobromide. The application route of DXM was restricted to oral (PO)
or intravenous (IV). All application forms (e.g. tablet, capsule, solution) were accepted. No restrictions
were imposed on the dosing amount of DXM or coadministrations of other substances. Studies containing
coadminstrations that inhibit or induce the pharmacokinetics of DXM were identified during the modeling
process and excluded. The relevant outcome measures are concentration-time profiles in plasma, serum,
and urine amounts of DXM, DXM metabolites, or metabolic ratios of metabolites such as UCMR. Studies
containing pharmacokinetic parameters of DXM and its metabolites (e.g. clearance, half-life, AUC)
and (urinary cumulative) metabolic ratios of DXM and its metabolites were included. Data containing
timecourses and CYP2D6 genotype information were prioritized. Non-healthy subjects were excluded if
the disease is known to affect the pharmacokinetics of DXM or DXM metabolites. Study B from the PhD
thesis of Frank (2009) highly deviates from the remaining data and was therefore excluded. Further, Wyen
et al. (2008) was identified as a duplicate of Study E from the PhD thesis of Frank (2009) and excluded.
The final set of curated studies used in the presented analyses is provided in Tab. 1.

For the selection and evaluation of studies from the literature, the PRISMA-ScR guidelines were adopted
where applicable (Tricco et al., 2018). The initial corpus contained 404 studies. After screening, application
of eligibility criteria, and prioritization, a total of 47 studies were curated (see supplementary Fig. S1). Of
these studies, 36 contained data used in the present work (Tab. 1).

141　2.2　PBPK model of DXM

The PBPK model of DXM, DXO, and DXO-Glu (Fig. 1) was encoded in the Systems Biology Markup
Language (SBML) (Hucka et al., 2019; Keating et al., 2020). For development and visualization,
sbmlutils (König, 2021b) and cy3sbml (König et al., 2012; König and Rodriguez, 2019) were
used. The model utilizes ordinary differential equations (ODE) which were numerically solved by
sbmlsim (König, 2021a) based on the high-performance SBML simulator libroadrunner (Somogyi
et al., 2015; Welsh et al., 2022). It is available in SBML under CC-BY 4.0 license from
https://github.com/matthiaskoenig/dextromethorphan-model. Within this work, version 0.9.5 of the model
was used Grzegorzewski and König (2022).

The model is hierarchically organized with submodels coupled using hierarchical model
composition (Smith et al., 2015). The top layer represents the whole body with organs and tissues
connected via the blood flow. The lower layer describes substance-related processes within the tissues.
Tissues with minor influence on the pharmacokinetics of DXM, DXO, or DXO-Glu are lumped into
the 'rest' compartment. Intravenous and oral application of DXM appears in the venous and intestinal

compartments, respectively. A fraction of DXM is absorbed via the intestinal wall into the systemic
circulation. The remainder is excreted via the feces. The plasma concentration is evaluated at the median
cubital vein.

The distribution of DXM, DXO, and DXO-Glu between plasma and tissue compartments is based on
tissue-to-plasma partition coefficients (K_p) and the corresponding rates of tissue distribution (f_{tissue}).

The metabolism of DXM only includes processes relevant for the simulation of the reported
pharmacokinetics data (see Fig. 1B and C). Routes of minor contribution such as N-demethylation of DXM
in the liver were neglected. Metabolic reactions take place in the intestine and liver and are modeled using
irreversible Michaelis-Menten reaction kinetics of the form $v = V_{max} \cdot \frac{S}{S+Km}$, with V_{max} and K_m for
CYP3A4 and CYP2D6 sampled from distributions as described below. The conversion of DXM to DXO
can be either catalyzed via CYP2D6 (main route) or CYP3A4 (minor route) in the liver. Reactions with
other products than DXM, DXO, and DXO-Glu were modeled as annihilation, i.e. the products of the
reaction are not modeled explicitly. DXM, DXO, and DXO-Glu are eliminated into the urine via renal
excretion.

A subset of model parameters was fitted by minimizing the distance between model predictions and
subsets of the data in Fig. 4, Fig. 5, Fig. 6, Fig. 7, Fig. 8, and Fig. 9.

2.3 CYP3A4 and CYP2D6

CYP3A4 and CYP2D6 variability was modeled via correlated bivariate lognormal distributions fitted
to *in vitro* data for CYP2D6 (Yang et al., 2012; Storelli et al., 2019a) and CYP3A4 (Yang et al., 2012),
respectively. The data was log10 transformed and a Gaussian, parameterized by the mean (μ) and standard
deviation, was fitted by maximum likelihood estimation. The multivariate distribution was realized by a
Gaussian copula which in turn was parameterized by Kendall's tau correlation coefficient from the data
(see Fig. 2 for data and model).

In order to model the effect of the CYP2D6 AS on the activity, V_{max} was assumed to be proportional to the
AS, $V_{max} \propto AS$ and K_m was scaled by the activity score along the first principle component of $\log10(K_m)$
and $\log10(V_{max})$ (principal component regression). To model the effect of genetic polymorphisms of
CYP2D6, pharmacogenetic variants in the CYP2D6 gene were mapped to their AS and the total activity
calculated as the sum of the activity of the two alleles. The genotype-phenotype definitions (i.e. allele
variant to AS mapping) were used from PharmGKB (https://www.pharmgkb.org/page/cyp2d6RefMaterials,
accessed on 2022-01-10) (Whirl-Carrillo et al., 2021) (Tab. S1).

The stochastic model of CYP2D6 kinetics for a given population consists of a mixture model
comprised from the models for each AS weighted by their respective frequency $P(AS)$, i.e.,
$P(V_{max}, K_m) = \sum_{AS} P(AS)P(V_{max}, K_m|AS)$. To simulate a given AS, the respective K_m and V_{max}
values were used (see Fig. 2). The variability in pharmacokinetics was simulated by sampling
K_m and V_{max} from the CYP3A4 and CYP2D6 distributions. Distributions of CYP3A4 and
CYP2D6 parameters were assumed to be statistically independent. To simulate different populations,
the AS frequencies for the respective biogeographical population were used from PharmGKB
(https://www.pharmgkb.org/page/cyp2d6RefMaterials, accessed on 2022-01-10) (Whirl-Carrillo et al.,
2021) (Tab. S2).

2.4 CYP2D6 metabolic phenotype

The metabolic phenotypes ultrarapid metabolizer (UM), extensive metabolizer (EM), intermediate metabolizer (IM), and poor metabolizer (PM) were assigned based on the urinary cumulative metabolic ratio of DXM to total dextrorphan $\text{UCMR} = \frac{\text{DXM}}{\text{DXO}+\text{DXO}-\text{Glu}}$ with the following cutoffs: PM: $\text{UCMR} \geq 0.3$, IM: $0.03 \leq \text{UCMR} < 0.3$, EM: $0.0003 \leq \text{UCMR} < 0.03$, UM: $\text{UCMR} < 0.0003$. Some studies reported the extensive metabolizer as normal metabolizer (NM) with identical cutoffs to the EM. Such data was labeled as EM.

2.5 Sensitivity analysis

A local sensitivity analysis of the effect of model parameters on the UCMR was performed. Individual model parameters (p_i) were varied in both directions by 10% from the base model value ($p_{i,-\Delta} \xleftarrow{10\%} p_{i,0} \xrightarrow{10\%} p_{i,\Delta}$) and the change in the state variable describing the UCMR at 8 hours (q) was recorded. The local sensitivity ($S(q, p_i, AS)$) was calculated for a range of ASs (0, 0.25, 0.5, 1, 1.25, 1.5, 2.0, 3.0) by the following formula:

$$S(q, p_i, AS) = \frac{1}{2} \cdot \frac{q(p_{i,\Delta}, AS) - q(p_{i,-\Delta}, AS)}{p_{i,0}} \tag{1}$$

3 RESULTS

Within this work, a physiologically based pharmacokinetic (PBPK) model of DXM was developed and applied to study the role of the CYP2D6 polymorphism on the pharmacokinetics of DXM and metabolic phenotyping using DXM.

3.1 Pharmacokinetics database of DXM

For the development and evaluation of the model, a large pharmacokinetics dataset of DXM and its metabolites, consisting of 36 clinical studies, was established (Tab. 1). Most of the studies investigated either drug-gene interactions (DGI), drug-drug interactions (DDI), or the interplay of both (i.e. drug-drug-gene interactions). The large majority of studies applied DXM orally (n=35), whereas only a single publication studied DXM pharmacokinetics after intravenous application (n=1) (Duedahl et al., 2005). The application form (i.e. solution, syrup, capsule, table), the used DXM dose (2 mg to 3 mg/kg), and coadministrations (i.e. phenotyping cocktail, quinidine, cinacalcet hydrochloride, zuojin) vary between studies, as do sampling times and sampled tissues (i.e. urine, plasma, serum). Importantly, plenty of individual UCMR measurements with corresponding CYP2D6 genotype information are contained within this dataset (n=11 studies). To our knowledge, this is the first large freely available dataset of pharmacokinetics data for DXM with all data accessible from the pharmacokinetics database (PK-DB) (Grzegorzewski et al., 2021).

3.2 PBPK model of DXM

Within this work, a PBPK model was developed (Fig. 1) to study the role of CYP2D6 polymorphism on DXM pharmacokinetics and metabolic phenotyping with DXM. Important model parameters are provided in Tab. 2. The model is organized hierarchically, with the top layer representing the whole body (Fig. 1A) consisting of the liver, kidney, intestine, forearm, lung, and the rest compartment. Organs with minor relevance are not modeled explicitly and lumped into the rest compartment. Organs are coupled via the systemic circulation. DXM can be administered intravenously (IV) or orally (PO) with DXM appearing in

the venous blood or intestine, respectively. The intestinal model (Fig 1B) describes dissolution, absorption
and excretion of DXM. Only a fraction of DXM is absorbed, with the remainder excreted in the feces.
DXM enters the circulatory system by crossing the enterocytes of the intestinal wall. First pass metabolism
of DXM via CYP3A4 N-demethylation in the intestine reduces the amount of DXM appearing in the
systemic circulation. In the liver model (Fig 1C), DXM gets transformed via O-demethylation to DXO
and subsequently transformed to DXO-Glu. The reactions are modeled by Michaelis Menten kinetics and
characterized with K_m and V_{max} values. O-demethylation takes place via CYP3A4 and CYP2D6. The
K_m and V_{max} of CYP2D6 is modulated via the AS, details can be found in section 3.3. The kidney model
(Fig 1D) describes the urinary excretion of DXM, DXO, and DXO-Glu.

The model allows to predict concentrations and amounts of DXM, DXO, and DXO-Glu depending on
CYP2D6 polymorphism, CYP2D6 diplotype, and CYP2D6 AS with amounts and concentrations of DXM,
DXO, and DXO-Glu being evaluated in urine or the median cubital vein (plasma).

To our knowledge, this is the first freely accessible, reproducible, and reusable PBPK model of DXM
with the model available in SBML from https://github.com/matthiaskoenig/dextromethorphan-model.

3.3 CYP3A4 and CYP2D6 variability

Cytochrome P450 enzymes exhibit enormous inter-individual variability in enzyme activity. To
account for this variability a stochastic model of CYP2D6 and CYP3A4 activity based on bivariate
lognormal distributions of K_m and V_{max} was developed and fitted to experimental data from human liver
microsomes (Storelli et al., 2019a; Yang et al., 2012) (see Fig. 2).

For the CYP2D6 model, the V_{max} is assumed to be linearly related to the AS with $AS = 0$ having no
CYP2D6 activity. The dispersion of K_m and V_{max} are assumed to be constant for all activity scores. For
the mixture model, the frequencies of the individual activity scores $P(AS)$ are adopted from our curated
dataset (i.e. relative amount of subjects with reported activity scores and UCMRs). With increasing AS the
maximal reaction velocity V_{max} of DXM conversion via CYP2D6 increases as does the affinity for the
substrate DXM (K_m decreases). The models of CYP3A4 and CYP2D6 are capable of reproducing the data
from the literature, but limited information on CYP2D6 genetics within the data hinders the validation of
the AS-specific model.

As motivated in the introduction, even subjects carrying no functional variant of the CYP2D6 gene do
metabolize DXM to DXO, however extremely slow. This was implemented in the model via a secondary O-
demethylation via CYP3A4 with mean K_m for DXM adopted from Yu and Haining (2001). The dispersion
of K_m and V_{max} is assumed to be identical to the one measured by midazolam in Storelli et al. (2019a)
and Yang et al. (2012).

The resulting CYP3A4 and CYP2D6 enzyme model was coupled to the PBPK model and allowed to
account (i) for the variability in DXM pharmacokinetics due to the variability in CYPs parameters and (ii)
the effect of the AS on CYP2D6 activity and consequently DXM pharmacokinetics.

3.4 Effect of CYP2D6 activity score on DXM pharmacokinetics

Model performance was visually assessed for common pharmacokinetic measurements (i.e. DXM,
DXO, DXM/DXO in plasma, and DXM/(DXO+DXO-Glu) in urine) and for subjects with reported AS or
diplotype (Fig. 3). For each AS, a virtual population based on 2000 K_m and V_{max} samples was created
from the stochastic models of CYP3A4 and CYP2D6 model. For every AS, an oral application of 30mg
DXM was simulated and compared to the corresponding data. The model predicts large relative variance

within a AS group as well as across different AS groups. With increasing AS, and consequently CYP2D6
activity, plasma DXM decreases (Fig. 3A), plasma DXO increases (Fig. 3B) and the plasma DXM/DXO
decreases (Fig. 3C) in very good agreement with the data (Chen et al., 2017; Frank, 2009). The large
variability within a AS group is a consequence of the large variability of K_m and V_{max} in CYP2D6 activity
of a single AS (see Fig. 2). The large overlap between distributions of adjacent AS results in a large overlap
in the pharmacokinetics between neighboring AS.

The UCMR (Fig. 3D) is very stable over time with a good agreement with the data. With increasing AS,
the UCMR decreases and thereby shifts from PM via IM to the EM metabolic phenotype. The UCMR data
was pooled independently of the amount of applied DXM (in contrast to A-C only using data from 30mg
oral application) and compared to the simulation as the UCMR endpoint is very robust against the given
dose (see section 3.5).

Overall the model predictions of DXM pharmacokinetics depending on AS are in very good agreement
with the available data despite the limited availability of pharmacokinetics timecourses for the low AS 0,
0.25, and 0.5.

To further evaluate the model performance, simulations were compared to pharmacokinetics data for
DXM in plasma or serum (Fig. 4), DXO in plasma or serum (Fig. 5), and DXO+DXO-Glu in plasma or
serum (Fig. 6), DXM in urine (Fig. 7), DXO+DXO-Glu in urine (Fig. 8), and the UCMR (Fig. 9). With
expected variability in mind, the model is capable to reproduce all data from the pharmacokinetics dataset.
Minor shortcomings of the model are faster kinetics of DXO+DXO-Glu in plasma (Fig. 6).

3.5 Effect of parameters on metabolic phenotyping via UCMR

Analysis of the effect of parameter changes on UCMR is highly relevant as it can help to identify potential
confounding factors and bias in UCMR based phenotyping. Of special importance is the question if there
is a dependency on the genetic polymorphism (activity score) of these effects.

To answer this question, model parameters (i.e. liver volume, cardiac output, tissue-to-plasma partition
coefficient of DXM, and oral dose) were changed in reasonable ranges and the effect on UCMR at 8
hours after the application of 30 mg of DXM was investigated (Fig. 10A). Independent of the AS, UCMR
increased with increasing liver volume and decreased with increasing cardiac output. A change in the
tissue-to-plasma partition coefficient of DXM or the amount of oral DXM barely affected the UCMR.

CYP2D6 phenotyping by UCMR is very stable over time as demonstrated in the time course predictions
(see 3D and Fig. 9) and robust against changes in factors related to the intervention protocol (i.e. dosing
amount of DXM, dissolution rate) and to some extent against changes in physiological parameters (see
local sensitivity analysis of UCMR in Fig. 10B).

Liver volume, heart rate, cardiac output, kidney volume, and kidney elimination rate of DXM altered the
UCMR with a similar magnitude as the CYP2D6 reaction parameters. However, the biological variation
in these physiological parameters is orders of magnitude lower. The sensitivity analysis showed no effect
of UGT V_{max} and K_m on UCMR which is the reason why inter-individual variability of UGT activity
was not further investigated in this work. Local sensitivity of UCMR was almost identical at different AS
values for almost all parameters, i.e., the effect of physiological parameters is of similar relative magnitude
independent of AS. For AS=0, our model assumptions of minor DXM metabolism by CYP3A4 lead
to UCMR not being modulated by CYP2D6 but rather by CYP3A4 activity. Nonetheless, even across
studies with non-standardized intervention protocols the UCMR measurements seams to be a good but not
perfect endpoint to quantify and compare CYP2D6 enzyme activity. Importantly, our analysis indicates

that UCMR measurements can be pooled even across investigations with different intervention protocols
(as for instance performed in Fig. 3D). This still may lead to biases and errors, e.g. due to differences in the
quantification protocol.

3.6   Effect of CYP2D6 polymorphisms and activity score on UCMR

Next, we tested if the model is able to predict UCMR distributions for given genotypes and AS (Fig. 11).
Model predictions based on underlying genotype frequencies were compared with the experimental data.
UCMR distributions for individual AS groups are well described by the model. The AS impacts the
UCMR, with increasing AS resulting in an decrease in UCMR. However individual AS distributions
heavily overlap, as expected, due to the large overlap in CYP2D6 parameter distributions between different
AS. The predicted distributions tend to be slightly narrower than the actual data. Possible reasons are many
fold (e.g. omitted physiological variation, omitted variation in UGT activity, difficulties in correct genotype
assignment, unknown effect modifiers, and biases).

The AS system could be refined to better describe the data. The categorization of CYP2D6 genotypes
into discrete activity values (i. e. 0, 0.25, 0.5, 1) is an oversimplification, a continuous activity score
would probably perform better. The model and data indicate that gUM ($AS \geq 3$) is a very unreliable
predictor for ultra rapid metabolism and only gPMs ($AS = 0$) are almost perfectly distinguishable from
other metabolizers, see Fig. 11C, D, E, and F.

Another strength of the presented model is that it enables the prediction of the *in vivo* phenotype of
subjects based on *in vitro* data.

3.7   Population variability in UCMR

Finally, the model was also capable to predict UCMR distributions for different biogeographical
populations (Fig. 12) based on the underlying AS frequencies (Tab. S2). Based on the reported frequencies,
the UCMR distributions were simulated at 8 hours after the application of 30mg DXM for Oceanian,
Near Eastern, American, Latino, Central/South Asian, African American/Afro-Caribbean, Sub-Saharan
African, European, and East Asian populations (Fig. 12A). Data for Caucasian and East Asian populations
(Fig. 12B) was used for validation of the predictions (Fig. 12C). The data is in good agreement with
measurements of Caucasians and East Asians as reported by Abdelrahman et al. (1999); Frank (2009);
Gaedigk (2013); Köhler et al. (1997); Myrand et al. (2008).

4   DISCUSSION

During the last 20 years various modeling approaches and software solutions were utilized to
investigate various aspects of DXM pharmacokinetics, e.g. using GastroPlus (Bolger et al., 2019), P-
Pharm (Moghadamnia et al., 2003), SAS (Ito et al., 2010; Chiba et al., 2012), SimCYP (Dickinson et al.,
2007; Ke et al., 2013; Sager et al., 2014; Chen et al., 2016; Rougée et al., 2016; Adiwidjaja et al., 2018;
Machavaram et al., 2019; Storelli et al., 2019b), MATLAB (Kim et al., 2017), or PK-Sim (Rüdesheim
et al., 2022). However, most of the work is difficult/impossible to validate or to build up on due to a lack
of accessibility of models and software, and platform-dependency of the models. Here, we provide an
openly accessible, reproducible and platform-independent whole-body model of DXM metabolism, which
facilitates reusability, extensibility, and comparability.

Apart from that, modeling work which aims for high empirically evidence relies on trustworthy supporting
real world data. More and independent sources of data are highly beneficial for the scientific outcomes. For

that matter, guidelines like PRISMA for reporting transparency, completeness, and accuracy find very broad
endorsement in the field of systematic reviews and meta analysis. The present work faces somewhat similar
challenges for the evaluation and selection of data from literature. Therefore, PRISMA-ScR guidelines
were adopted where applicable. With this approach, bias within the used dataset could be mitigated or
at least identified. Importantly, we supplement our open and accessible model with a large, open, and
accessible database of pharmacokinetics data.

The presented PBPK model is able to predict the DXM metabolism of populations and individuals based
on their CYP2D6 genotype. It is probably the first model capable to predict individual UCMRs and the
expected distributions of UCMR. Moreover, it can reproduce a broad range of reported clinical data on
DXM and enables better intuition on how to interpret DXM related pharmacokinetics. E.g., an important
message is that CYP2D6 activity is not the only modulator of UCMR, as can be seen by the large variability
in activity score and overlap between activity scores. UCMR as a proxy of CYP2D6 metabolic phenotype
should therefore be interpreted carefully. The model shows that for extremely low CYP2D6 activity the
UCMR is not primarily governed by the CYP2D6 activity. This is consistent with the finding that CYP2D6
inhibition merely affects PMs (Pope et al., 2004).

The current version of the model is already very valuable, still there is plenty of room for improvement.
By providing the data and model in open and standardized formats we enable and encourage these
improvements by model extensions and updates.

Many of the physiological parameters in the model were fitted or estimated although they could be
measured in principle. E.g., relatively low DXM concentrations in plasma suggest substantial extra-
vascular binding of DXM. However, tissue-plasma partition coefficients (Kp) are difficult to assess and
only limited data is available. Steinberg et al. (1996) reported brain levels to be 68-fold higher and
cerebrospinal fluid levels 4-fold lower than serum levels, respectively. Others estimated Kp $\sim$ 1.65 from
n-octanol-water partition coefficients and again others suggested additional trapping mechanisms (i.e.
lysosomal trapping) (Bolger et al., 2019). In the model, the DXO-Glu kinetics is a bit to rapid (see Fig. 6),
probably due to the decision to model tissue distribution uniformly for all organs (i.e. identical K_p and
f_{tissue}). We decided for a more parsimonious model. Glucuronides, however, are generally much more
polar than their respective non-glucuronides which result in less plasma binding, higher urinary excretion,
lower lipid-solubility, and higher water-solubility. Transport into different tissues is affected differently by
polarity.

Most important for model improvements would be additional *in vitro* measurements on the association
between CYP2D6 genotype and phenotype which are very limited in literature (Storelli et al., 2019a; Ning
et al., 2019; Dalton et al., 2020). Furthermore, simultaneous *in vitro* and UCMR measurements do not exist
the literature. Both would be very important for the validation of the AS system and the development of
new models which e.g. take into account structural variation (Dalton et al., 2020). For instance, with the AS
system alone it is not possible to explain why CYP2D6 is inhibited differently for different genotypes Qiu
et al. (2016).

In conclusion, we developed and validated a PBPK model of DXM and applied it to study the effect of
the CYP2D6 polymorphism on metabolic phenotyping.

CONFLICT OF INTEREST STATEMENT

390 All authors declare that the research was conducted in the absence of any commercial or financial
391 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

JG and MK designed the study, developed the computational model, implemented and performed the analysis, and wrote the initial draft of the manuscript. JB provided support with data curation. All authors discussed the results. All authors contributed to and revised the manuscript critically.

FUNDING

JG, JB, and MK were supported by the Federal Ministry of Education and Research (BMBF, Germany) within the research network Systems Medicine of the Liver (LiSyM, grant number 031L0054). MK and JG were supported by the German Research Foundation (DFG) within the Research Unit Program FOR 5151 "QuaLiPerF (Quantifying Liver Perfusion-Function Relationship in Complex Resection - A Systems Medicine Approach)" by grant number 436883643 and MK by grant number 465194077 (Priority Programme SPP 2311, Subproject SimLivA). This work was supported by the BMBF-funded de.NBI Cloud within the German Network for Bioinformatics Infrastructure (de.NBI) (031A537B, 031A533A, 031A538A, 031A533B, 031A535A, 031A537C, 031A534A, 031A532B).

DATA AVAILABILITY STATEMENT

All clinical data of dextromethorphan pharmacokinetics that was used in this work can be found in PK-DB available from https://pk-db.com.

REFERENCES

Abdelrahman, S., Gotschall, R., Kauffman, R., Leeder, J., and Kearns, G. (1999). Investigation of terbinafine as a CYP2D6 inhibitor in vivo. *Clinical Pharmacology & Therapeutics* 65, 465–472. doi:10.1016/S0009-9236(99)70065-2

Abduljalil, K., Frank, D., Gaedigk, A., Klaassen, T., Tomalik-Scharte, D., Jetter, A., et al. (2010). Assessment of Activity Levels for CYP2D6*1, CYP2D6*2, and CYP2D6*41 Genes by Population Pharmacokinetics of Dextromethorphan. *Clinical Pharmacology & Therapeutics* 88, 643–651. doi:10.1038/clpt.2010.137

Adiwidjaja, J., Boddy, A. V., and McLachlan, A. J. (2018). A Strategy to Refine the Phenotyping Approach and Its Implementation to Predict Drug Clearance: A Physiologically Based Pharmacokinetic Simulation Study. *CPT: Pharmacometrics & Systems Pharmacology* 7, 798–808. doi:10.1002/psp4.12355

Armani, S., Ting, L., Sauter, N., Darstein, C., Tripathi, A. P., Wang, L., et al. (2017). Drug Interaction Potential of Osilodrostat (LCI699) Based on Its Effect on the Pharmacokinetics of Probe Drugs of Cytochrome P450 Enzymes in Healthy Adults. *Clinical drug investigation* 37, 465–472. doi:10.1007/s40261-017-0497-0

Barnhart, J. W. (1980). The urinary excretion of dextromethorphan and three metabolites in dogs and humans. *Toxicology and Applied Pharmacology* 55, 43–48. doi:10.1016/0041-008X(80)90218-5

Berm, E. J. J., Risselada, A. J., Mulder, H., Hak, E., and Wilffert, B. (2013). Phenoconversion of cytochrome P450 2D6: The need for identifying the intermediate metabolizer genotype. *The Journal of clinical psychiatry* 74, 1025. doi:10.4088/JCP.13lr08555

Bolger, M. B., Macwan, J. S., Sarfraz, M., Almukainzi, M., and Löbenberg, R. (2019). The Irrelevance of In Vitro Dissolution in Setting Product Specifications for Drugs Like Dextromethorphan That are Subject to Lysosomal Trapping. *Journal of pharmaceutical sciences* 108, 268–278. doi:10.1016/j.xphs.2018.09.036

Capon, D. A., Bochner, F., Kerry, N., Mikus, G., Danz, C., and Somogyi, A. A. (1996). The influence of
CYP2D6 polymorphism and quinidine on the disposition and antitussive effect of dextromethorphan in
humans. *Clinical Pharmacology & Therapeutics* 60, 295–307. doi:10.1016/S0009-9236(96)90056-9

Caudle, K. E., Sangkuhl, K., Whirl-Carrillo, M., Swen, J. J., Haidar, C. E., Klein, T. E., et al. (2020).
Standardizing CYP2D6 Genotype to Phenotype translation: Consensus Recommendations from the
Clinical Pharmacogenetics Implementation Consortium and Dutch Pharmacogenetics Working Group.
Clinical and Translational Science 13, 116–124. doi:10.1111/cts.12692

Chen, R., Rostami-Hodjegan, A., Wang, H., Berk, D., Shi, J., and Hu, P. (2016). Application of a
physiologically based pharmacokinetic model for the evaluation of single-point plasma phenotyping
method of CYP2D6. *European journal of pharmaceutical sciences : official journal of the European
Federation for Pharmaceutical Sciences* 92, 131–136. doi:10.1016/j.ejps.2016.07.001

Chen, R., Zheng, X., and Hu, P. (2017). CYP2D6 Phenotyping Using Urine, Plasma, and Saliva Metabolic
Ratios to Assess the Impact of CYP2D6$_*$10 on Interindividual Variation in a Chinese Population.
Frontiers in Pharmacology 8, 239. doi:10.3389/fphar.2017.00239

Chiba, K., Kato, M., Ito, T., Suwa, T., and Sugiyama, Y. (2012). Inter-individual variability of in
vivo CYP2D6 activity in different genotypes. *Drug metabolism and pharmacokinetics* 27, 405–413.
doi:10.2133/dmpk.dmpk-11-rg-078

Chládek, J., Zimová, G., Beránek, M., and Martínková, J. (2000). In-vivo indices of CYP2D6 activity:
Comparison of dextromethorphan metabolic ratios in 4-h urine and 3-h plasma. *European Journal of
Clinical Pharmacology* 56, 651–657. doi:10.1007/s002280000218

Dalton, R., Lee, S.-B., Claw, K. G., Prasad, B., Phillips, B. R., Shen, D. D., et al. (2020). Interrogation of
CYP2D6 Structural Variant Alleles Improves the Correlation Between CYP2D6 Genotype and CYP2D6-
Mediated Metabolic Activity. *Clinical and translational science* 13, 147–156. doi:10.1111/cts.1269
5

De Kesel, P. M. M., Lambert, W. E., and Stove, C. P. (2016). Alternative Sampling Strategies for
Cytochrome P450 Phenotyping. *Clinical Pharmacokinetics* 55, 169–184. doi:10.1007/s40262-015-030
6-y

de Simone, G., Devereux, R. B., Daniels, S. R., Mureddu, G., Roman, M. J., Kimball, T. R., et al. (1997).
Stroke volume and cardiac output in normotensive children and adults. Assessment of relations with
body size and impact of overweight. *Circulation* 95, 1837–1843. doi:10.1161/01.cir.95.7.1837

Demirbas, S., Reyderman, L., and Stavchansky, S. (1998). Bioavailability of dextromethorphan (as
dextrorphan) from sustained release formulations in the presence of guaifenesin in human volunteers.
Biopharmaceutics & Drug Disposition 19, 541–545. doi:10.1002/(SICI)1099-081X(1998110)19:8⟨541::
AID-BDD138⟩3.0.CO;2-8

Dickinson, G. L., Rezaee, S., Proctor, N. J., Lennard, M. S., Tucker, G. T., and Rostami-Hodjegan, A.
(2007). Incorporating In Vitro Information on Drug Metabolism Into Clinical Trial Simulations to Assess
the Effect of CYP2D6 Polymorphism on Pharmacokinetics and Pharmacodynamics: Dextromethorphan
as a Model Application. *The Journal of Clinical Pharmacology* 47, 175–186. doi:10.1177/0091270006
294279

Dorado, P., González, I., Naranjo, M. E. G., de Andrés, F., Peñas-Lledó, E. M., Calzadilla, L. R.,
et al. (2017). Lessons from Cuba for Global Precision Medicine: *CYP2D6* Genotype Is Not a Robust
Predictor of CYP2D6 Ultrarapid Metabolism. *OMICS: A Journal of Integrative Biology* 21, 17–26.
doi:10.1089/omi.2016.0166

Doroshyenko, O., Rokitta, D., Zadoyan, G., Klement, S., Schläfke, S., Dienel, A., et al. (2013). Drug
cocktail interaction study on the effect of the orally administered lavender oil preparation silexan on

cytochrome P450 enzymes in healthy volunteers. *Drug Metab. Dispos.* 41, 987–993. doi:10.1124/dmd.
112.050203

Duedahl, T. H., Dirks, J., Petersen, K. B., Romsing, J., Larsen, N.-E., and Dahl, J. B. (2005). Intravenous
dextromethorphan to human volunteers: Relationship between pharmacokinetics and anti-hyperalgesic
effect. *Pain* 113, 360–368. doi:10.1016/j.pain.2004.11.015

Dumond, J. B., Vourvahis, M., Rezk, N. L., Patterson, K. B., Tien, H.-C., White, N., et al. (2010). A
Phenotype–Genotype Approach to Predicting CYP450 and P-Glycoprotein Drug Interactions With the
Mixed Inhibitor/Inducer Tipranavir/Ritonavir. *Clinical Pharmacology & Therapeutics* 87, 735–742.
doi:10.1038/clpt.2009.253

Edwards, J. E., Eliot, L., Parkinson, A., Karan, S., and MacConell, L. (2017). Assessment
of Pharmacokinetic Interactions Between Obeticholic Acid and Caffeine, Midazolam, Warfarin,
Dextromethorphan, Omeprazole, Rosuvastatin, and Digoxin in Phase 1 Studies in Healthy Subjects.
Advances in Therapy 34, 2120–2138. doi:10.1007/s12325-017-0601-0

Eichelbaum, M., Spannbrucker, N., Steincke, B., and Dengler, H. J. (1979). Defective N-oxidation of
sparteine in man: A new pharmacogenetic defect. *European journal of clinical pharmacology* 16,
183–187. doi:10.1007/BF00562059

Eichhold, T. H., Greenfield, L. J., Hoke, S. H., and Wehmeyer, K. R. (1997). Determination
of dextromethorphan and dextrorphan in human plasma by liquid chromatography/tandem mass
spectrometry. *Journal of Mass Spectrometry* 32, 1205–1211. doi:10.1002/(SICI)1096-9888(199711)32:
11⟨1205::AID-JMS579⟩3.0.CO;2-C

Eichhold, T. H., McCauley-Myers, D. L., Khambe, D. A., Thompson, G. A., and Hoke, S. H. (2007).
Simultaneous determination of dextromethorphan, dextrorphan, and guaifenesin in human plasma using
semi-automated liquid/liquid extraction and gradient liquid chromatography tandem mass spectrometry.
Journal of Pharmaceutical and Biomedical Analysis 43, 586–600. doi:10.1016/j.jpba.2006.07.018

Frank, D. (2009). *Evaluation of pharmacokinetic metrics for phenotyping of the human CYP2D6 enzyme*
with dextromethorphan. Ph.D. thesis, Rheinische Friedrich-Wilhelms-Universität Bonn

Frank, D., Jaehde, U., and Fuhr, U. (2007). Evaluation of probe drugs and pharmacokinetic metrics for
CYP2D6 phenotyping. *European Journal of Clinical Pharmacology* 63, 321–333. doi:10.1007/s00228
-006-0250-8

Fuhr, U., Jetter, A., and Kirchheiner, J. (2007). Appropriate phenotyping procedures for drug metabolizing
enzymes and transporters in humans and their simultaneous use in the "cocktail" approach. *Clinical*
pharmacology and therapeutics 81, 270–283. doi:10.1038/sj.clpt.6100050

Gaedigk, A. (2013). Complexities of *CYP2D6* gene analysis and interpretation. *International Review of*
Psychiatry 25, 534–553. doi:10.3109/09540261.2013.825581

Gaedigk, A., Dinh, J., Jeong, H., Prasad, B., and Leeder, J. (2018a). Ten Years' Experience with the
CYP2D6 Activity Score: A Perspective on Future Investigations to Improve Clinical Predictions for
Precision Therapeutics. *Journal of Personalized Medicine* 8, 15. doi:10.3390/jpm8020015

Gaedigk, A., Eklund, J. D., Pearce, R. E., Leeder, J. S., Alander, S. W., Phillips, M. S., et al. (2007).
Identification and characterization of CYP2D6*56B, an allele associated with the poor metabolizer
phenotype. *Clinical pharmacology and therapeutics* 81, 817–820. doi:10.1038/sj.clpt.6100125

Gaedigk, A., Ingelman-Sundberg, M., Miller, N. A., Leeder, J. S., Whirl-Carrillo, M., Klein, T. E., et al.
(2018b). The pharmacogene variation (PharmVar) consortium: Incorporation of the human cytochrome
P450 (CYP) allele nomenclature database. *Clinical pharmacology and therapeutics* 103, 399–401.
doi:10.1002/cpt.910

Gaedigk, A., Sangkuhl, K., Whirl-Carrillo, M., Klein, T., and Leeder, J. S. (2017). Prediction of CYP2D6
phenotype from genotype across world populations. *Genetics in Medicine: Official Journal of the*
American College of Medical Genetics 19, 69–76. doi:10.1038/gim.2016.80

Gaedigk, A., Turner, A., Everts, R. E., Scott, S. A., Aggarwal, P., Broeckel, U., et al. (2019).
Characterization of Reference Materials for Genetic Testing of CYP2D6 Alleles. *The Journal of*
Molecular Diagnostics 21, 1034–1052. doi:10.1016/j.jmoldx.2019.06.007

Gasche, Y., Daali, Y., Fathi, M., Chiappe, A., Cottini, S., Dayer, P., et al. (2004). Codeine intoxication
associated with ultrarapid CYP2D6 metabolism. *The New England journal of medicine* 351, 2827–2831.
doi:10.1056/NEJMoa041888

Gonzalez Hernandez, F., Carter, S. J., Iso-Sipilä, J., Goldsmith, P., Almousa, A. A., Gastine, S., et al.
(2021). An automated approach to identify scientific publications reporting pharmacokinetic parameters.
Wellcome Open Research 6, 88. doi:10.12688/wellcomeopenres.16718.1

Grzegorzewski, J., Bartsch, F., Köller, A., and König, M. (2022). Pharmacokinetics of Caffeine: A
Systematic Analysis of Reported Data for Application in Metabolic Phenotyping and Liver Function
Testing. *Frontiers in Pharmacology* 12. doi:10.3389/fphar.2021.752826

Grzegorzewski, J., Brandhorst, J., Green, K., Eleftheriadou, D., Duport, Y., Barthorscht, F., et al. (2021).
PK-DB: Pharmacokinetics database for individualized and stratified computational modeling. *Nucleic*
Acids Research 49, D1358–D1364. doi:10.1093/nar/gkaa990

[Dataset] Grzegorzewski, J. and König, M. (2022). Physiologically based pharmacokinetic (PBPK) model
of dextromethorphan. https://doi.org/10.5281/zenodo.7025683. doi:10.5281/zenodo.7025683. Accessed:
2022-08-26

Herman, I. P. (2016). *Physics of the Human Body* (Springer)

Hou, Z.-Y., Pickle, L. W., Meyer, P. S., and Woosley, R. L. (1991). Salivary analysis for determination
of dextromethorphan metabolic phenotype. *Clinical Pharmacology and Therapeutics* 49, 410–419.
doi:10.1038/clpt.1991.48

Hu, L., Li, L., Yang, X., Liu, W., Yang, J., Jia, Y., et al. (2011). Floating matrix dosage form for
dextromethorphan hydrobromide based on gas forming technique: In vitro and in vivo evaluation in
healthy volunteers. *European Journal of Pharmaceutical Sciences* 42, 99–105. doi:10.1016/j.ejps.2010.
10.010

Hucka, M., Bergmann, F. T., Chaouiya, C., Dräger, A., Hoops, S., Keating, S. M., et al. (2019). The
systems biology markup language (SBML): Language specification for level 3 version 2 core release 2.
Journal of Integrative Bioinformatics doi:10.1515/jib-2019-0021

Hurtado, I., García-Sempere, A., Peiró, S., and Sanfélix-Gimeno, G. (2020). Increasing Trends in Opioid
Use From 2010 to 2018 in the Region of Valencia, Spain: A Real-World, Population-Based Study.
Frontiers in Pharmacology 11, 612556. doi:10.3389/fphar.2020.612556

ICRP (2002). Basic anatomical and physiological data for use in radiological protection: Reference values.
A report of age- and gender-related differences in the anatomical and physiological characteristics of
reference individuals. ICRP Publication 89. *Annals of the ICRP* 32, 5–265

Ito, T., Kato, M., Chiba, K., Okazaki, O., and Sugiyama, Y. (2010). Estimation of the interindividual
variability of cytochrome 2D6 activity from urinary metabolic ratios in the literature. *Drug Metabolism*
and Pharmacokinetics 25, 243–253. doi:10.2133/dmpk.25.243

Jones, D. R., Gorski, J. C., Haehner, B. D., O'Mara, E. M., and Hall, S. D. (1996). Determination
of cytochrome P450 3A4/5 activity in vivo with dextromethorphan N-demethylation. *Clinical*
Pharmacology and Therapeutics 60, 374–384. doi:10.1016/S0009-9236(96)90194-0

Jones, H. and Rowland-Yeo, K. (2013). Basic concepts in physiologically based pharmacokinetic modeling
in drug discovery and development. *CPT: pharmacometrics & systems pharmacology* 2, e63. doi:10.1
038/psp.2013.41

Kawanishi, C., Lundgren, S., Agren, H., and Bertilsson, L. (2004). Increased incidence of CYP2D6
gene duplication in patients with persistent mood disorders: Ultrarapid metabolism of antidepressants
as a cause of nonresponse. A pilot study. *European Journal of Clinical Pharmacology* 59, 803–807.
doi:10.1007/s00228-003-0701-4

Ke, A. B., Nallani, S. C., Zhao, P., Rostami-Hodjegan, A., Isoherranen, N., and Unadkat, J. D. (2013).
A Physiologically Based Pharmacokinetic Model to Predict Disposition of CYP2D6 and CYP1A2
Metabolized Drugs in Pregnant Women. *Drug Metabolism and Disposition* 41, 801–813. doi:10.1124/
dmd.112.050161

Keating, S. M., Waltemath, D., König, M., Zhang, F., Dräger, A., Chaouiya, C., et al. (2020). SBML Level
3: An extensible format for the exchange and reuse of biological models. *Molecular systems biology* 16,
e9110. doi:10.15252/msb.20199110

Kerry, N., Somogyi, A., Bochner, F., and Mikus, G. (1994). The role of CYP2D6 in primary and secondary
oxidative metabolism of dextromethorphan: In vitro studies using human liver microsomes. *British*
Journal of Clinical Pharmacology 38, 243–248. doi:10.1111/j.1365-2125.1994.tb04348.x

Kibaly, C., Alderete, J. A., Liu, S. H., Nasef, H. S., Law, P.-Y., Evans, C. J., et al. (2021). Oxycodone in the
Opioid Epidemic: High 'Liking', 'Wanting', and Abuse Liability. *Cellular and Molecular Neurobiology*
41, 899–926. doi:10.1007/s10571-020-01013-y

Kim, E.-Y., Shin, S.-G., and Shin, J.-G. (2017). Prediction and visualization of CYP2D6 genotype-
based phenotype using clustering algorithms. *Translational and Clinical Pharmacology* 25, 147–152.
doi:10.12793/tcp.2017.25.3.147

Köhler, D., Härtter, S., Fuchs, K., Sieghart, W., and Hiemke, C. (1997). CYP2D6 genotype and phenotyping
by determination of dextromethorphan and metabolites in serum of healthy controls and of patients under
psychotropic medication. *Pharmacogenetics* 7, 453–461. doi:10.1097/00008571-199712000-00003

[Dataset] König, M. (2021a). Matthiaskoenig/sbmlsim: 0.1.14 - SBML simulation made easy.
https://zenodo.org/record/5531088. doi:10.5281/zenodo.5531088. Accessed: 2022-08-23

[Dataset] König, M. (2021b). Sbmlutils: Python utilities for SBML. https://zenodo.org/record/6599299.
doi:10.5281/zenodo.5546603. Accessed: 2022-08-23

[Dataset] König, M. and Rodriguez, N. (2019). Matthiaskoenig/cy3sbml: Cy3sbml-v0.3.0 - SBML for
Cytoscape. https://zenodo.org/record/3451319. doi:10.5281/zenodo.3451319. Accessed: 2022-08-23

König, M., Dräger, A., and Holzhütter, H.-G. (2012). CySBML: A Cytoscape plugin for SBML.
Bioinformatics (Oxford, England) 28, 2402–2403. doi:10.1093/bioinformatics/bts432

Lenuzza, N., Duval, X., Nicolas, G., Thévenot, E., Job, S., Videau, O., et al. (2016). Safety and
pharmacokinetics of the CIME combination of drugs and their metabolites after a single oral dosing
in healthy volunteers. *European journal of drug metabolism and pharmacokinetics* 41, 125–138.
doi:10.1007/s13318-014-0239-0

López, M., Guerrero, J., Jung–Cook, H., and Alonso, M. E. (2005). CYP2D6 genotype and phenotype
determination in a Mexican Mestizo population. *European Journal of Clinical Pharmacology* 61,
749–754. doi:10.1007/s00228-005-0038-2

Lutz, J. D. and Isoherranen, N. (2012). Prediction of Relative In Vivo Metabolite Exposure from In Vitro
Data Using Two Model Drugs: Dextromethorphan and Omeprazole. *Drug Metabolism and Disposition*
40, 159–168. doi:10.1124/dmd.111.042200

Machavaram, K. K., Endo-Tsukude, C., Terao, K., Gill, K. L., Hatley, O. J., Gardner, I., et al. (2019).
Simulating the Impact of Elevated Levels of Interleukin-6 on the Pharmacokinetics of Various CYP450
Substrates in Patients with Neuromyelitis Optica or Neuromyelitis Optica Spectrum Disorders in
Different Ethnic Populations. *The AAPS journal* 21, 42. doi:10.1208/s12248-019-0309-y

Mahgoub, A., Idle, J. R., Dring, L. G., Lancaster, R., and Smith, R. L. (1977). Polymorphic hydroxylation
of Debrisoquine in man. *Lancet (London, England)* 2, 584–586. doi:10.1016/s0140-6736(77)91430-1

McGinnity, D. F., Parker, A. J., Soars, M., and Riley, R. J. (2000). Automated definition of the enzymology
of drug oxidation by the major human drug metabolizing cytochrome P450s. *Drug metabolism and*
disposition: the biological fate of chemicals 28, 1327–1334

Moghadamnia, A. A., Rostami-Hodjegan, A., Abdul-Manap, R., Wright, C. E., Morice, A. H., and Tucker,
G. T. (2003). Physiologically based modelling of inhibition of metabolism and assessment of the
relative potency of drug and metabolite: Dextromethorphan *vs* . dextrorphan using quinidine inhibition:
Dextromethorphan vs. dextrorphan using quinidine inhibition. *British Journal of Clinical Pharmacology*
56, 57–67. doi:10.1046/j.1365-2125.2003.01853.x

Montané Jaime, L. K., Lalla, A., Steimer, W., and Gaedigk, A. (2013). Characterization of the *CYP2D6*
gene locus and metabolic activity in Indo- and Afro-Trinidadians: Discovery of novel allelic variants.
Pharmacogenomics 14, 261–276. doi:10.2217/pgs.12.207

Myrand, S., Sekiguchi, K., Man, M., Lin, X., Tzeng, R.-Y., Teng, C.-H., et al. (2008).
Pharmacokinetics/Genotype Associations for Major Cytochrome P450 Enzymes in Native and First- and
Third-generation Japanese Populations: Comparison With Korean, Chinese, and Caucasian Populations.
Clinical Pharmacology & Therapeutics 84, 347–361. doi:10.1038/sj.clpt.6100482

Nagai, N., Kawakubo, T., Kaneko, F., Ishii, M., Shinohara, R., Saito, Y., et al. (1996). Pharmacokinetics
and polymorphic oxidation of dextromethorphan in a japanese population. *Biopharmaceutics & Drug*
Disposition 17, 421–433. doi:10.1002/(SICI)1099-081X(199607)17:5⟨421::AID-BDD421⟩3.0.CO;2-9

Nakashima, D., Takama, H., Ogasawara, Y., Kawakami, T., Nishitoba, T., Hoshi, S., et al. (2007). Effect of
Cinacalcet Hydrochloride, a New Calcimimetic Agent, on the Pharmacokinetics of Dextromethorphan:
In Vitro and Clinical Studies. *The Journal of Clinical Pharmacology* 47, 1311–1319. doi:10.1177/0091
270007304103

Ning, M., Duarte, J. D., Stevison, F., Isoherranen, N., Rubin, L. H., and Jeong, H. (2019). Determinants
of Cytochrome P450 2D6 mRNA Levels in Healthy Human Liver Tissue. *Clinical and Translational*
Science 12, 416–423. doi:10.1111/cts.12632

Nofziger, C. and Paulmichl, M. (2018). Accurately genotyping *CYP2D6* : Not for the faint of heart.
Pharmacogenomics 19, 999–1002. doi:10.2217/pgs-2018-0105

Nyunt, M. M., Becker, S., MacFarland, R. T., Chee, P., Scarborough, R., Everts, S., et al. (2008).
Pharmacokinetic Effect of AMD070, an Oral CXCR4 Antagonist, on CYP3A4 and CYP2D6 Substrates
Midazolam and Dextromethorphan in Healthy Volunteers. *JAIDS Journal of Acquired Immune Deficiency*
Syndromes 47, 559–565. doi:10.1097/QAI.0b013e3181627566

Oh, K.-S., Park, S.-J., Shinde, D. D., Shin, J.-G., and Kim, D.-H. (2012). High-sensitivity liquid
chromatography-tandem mass spectrometry for the simultaneous determination of five drugs and their
cytochrome P450-specific probe metabolites in human plasma. *Journal of chromatography. B, Analytical*
technologies in the biomedical and life sciences 895–896, 56–64. doi:10.1016/j.jchromb.2012.03.014

Pope, L. E., Khalil, M. H., Berg, J. E., Stiles, M., Yakatan, G. J., and Sellers, E. M. (2004).
Pharmacokinetics of Dextromethorphan After Single or Multiple Dosing in Combination With
Quinidine in Extensive and Poor Metabolizers. *The Journal of Clinical Pharmacology* 44, 1132–1142.
doi:10.1177/0091270004269521

Preskorn, S. H., Kane, C. P., Lobello, K., Nichols, A. I., Fayyad, R., Buckley, G., et al. (2013). Cytochrome
P450 2D6 phenoconversion is common in patients being treated for depression: Implications for
personalized medicine. *The Journal of clinical psychiatry* 74, 614–621. doi:10.4088/JCP.12m07807
Qiu, F., Liu, S., Miao, P., Zeng, J., Zhu, L., Zhao, T., et al. (2016). Effects of the Chinese herbal
formula "Zuojin Pill" on the pharmacokinetics of dextromethorphan in healthy Chinese volunteers with
CYP2D6*10 genotype. *European journal of clinical pharmacology* 72, 689–695. doi:10.1007/s00228-0
16-2048-7
Rau, T., Wohlleben, G., Wuttke, H., Thuerauf, N., Lunkenheimer, J., Lanczik, M., et al. (2004). CYP2D6
genotype: Impact on adverse effects and nonresponse during treatment with antidepressants-a pilot study.
Clinical pharmacology and therapeutics 75, 386–393. doi:10.1016/j.clpt.2003.12.015
[Dataset] RNAO (2022). Nursing best practice guideline by nurses' association
of ontario: Appendix g: Tools for assessing anxiety, depression, and stress.
https://bpgmobile.rnao.ca/sites/default/files/Appendix%20C%20Vein%20Anatomy.pdf. Accessed:
2022-08-23
Rougée, L. R. A., Mohutsky, M. A., Bedwell, D. W., Ruterbories, K. J., and Hall, S. D. (2016). The
Impact of the Hepatocyte-to-Plasma pH Gradient on the Prediction of Hepatic Clearance and Drug-Drug
Interactions for CYP2D6 Substrates. *Drug metabolism and disposition: the biological fate of chemicals*
44, 1819–1827. doi:10.1124/dmd.116.071761
Rüdesheim, S., Selzer, D., Fuhr, U., Schwab, M., and Lehr, T. (2022). Physiologically-based
pharmacokinetic modeling of dextromethorphan to investigate interindividual variability within CYP2D6
activity score groups. *CPT: Pharmacometrics & Systems Pharmacology* n/a. doi:10.1002/psp4.12776
Sager, J. E., Lutz, J. D., Foti, R. S., Davis, C., Kunze, K. L., and Isoherranen, N. (2014). Fluoxetine-
and norfluoxetine-mediated complex drug-drug interactions: In vitro to in vivo correlation of effects on
CYP2D6, CYP2C19, and CYP3A4. *Clinical pharmacology and therapeutics* 95, 653–662. doi:10.1038/
clpt.2014.50
Saravanakumar, A., Sadighi, A., Ryu, R., and Akhlaghi, F. (2019). Physicochemical Properties,
Biotransformation, and Transport Pathways of Established and Newly Approved Medications: A
Systematic Review of the Top 200 Most Prescribed Drugs vs. the FDA-Approved Drugs Between
2005 and 2016. *Clinical Pharmacokinetics* 58, 1281–1294. doi:10.1007/s40262-019-00750-8
Schadel, M., Wu, D., Otton, S. V., Kalow, W., and Sellers, E. M. (1995). Pharmacokinetics of
dextromethorphan and metabolites in humans: Influence of the CYP2D6 phenotype and quinidine
inhibition. *Journal of Clinical Psychopharmacology* 15, 263–269. doi:10.1097/00004714-199508000-0
0005
Schoedel, K. A., Pope, L. E., and Sellers, E. M. (2012). Randomized Open-Label Drug-Drug Interaction
Trial of Dextromethorphan/Quinidine and Paroxetine in Healthy Volunteers:. *Clinical Drug Investigation*
32, 157–169. doi:10.2165/11599870-000000000-00000
Shah, R. R. and Smith, R. L. (2015). Addressing phenoconversion: The Achilles' heel of personalized
medicine: Impact of phenoconversion. *British Journal of Clinical Pharmacology* 79, 222–240. doi:10.1
111/bcp.12441
Silva, A. R. and Dinis-Oliveira, R. J. (2020). Pharmacokinetics and pharmacodynamics of
dextromethorphan: Clinical and forensic aspects. *Drug Metabolism Reviews* 52, 258–282. doi:10
.1080/03602532.2020.1758712
Smith, L. P., Hucka, M., Hoops, S., Finney, A., Ginkel, M., Myers, C. J., et al. (2015). SBML Level 3
package: Hierarchical Model Composition, Version 1 Release 3. *J Integr Bioinform* 12, 268. doi:10.239
0/biecoll-jib-2015-268

Somogyi, E. T., Bouteiller, J.-M., Glazier, J. A., König, M., Medley, J. K., Swat, M. H., et al. (2015).
libRoadRunner: A high performance SBML simulation and analysis library. *Bioinformatics (Oxford,*
England) 31, 3315–3321. doi:10.1093/bioinformatics/btv363

Steinberg, G. K., Bell, T. E., and Yenari, M. A. (1996). Dose escalation safety and tolerance study of the
N-methyl-d-aspartate antagonist dextromethorphan in neurosurgery patients. *Journal of Neurosurgery*
84, 860–866. doi:10.3171/jns.1996.84.5.0860

Storelli, F., Desmeules, J., and Daali, Y. (2019a). Genotype-sensitive reversible and time-dependent
CYP2D6 inhibition in human liver microsomes. *Basic & clinical pharmacology & toxicology* 124,
170–180. doi:10.1111/bcpt.13124

Storelli, F., Desmeules, J., and Daali, Y. (2019b). Physiologically-Based Pharmacokinetic Modeling for
the Prediction of CYP2D6-Mediated Gene–Drug–Drug Interactions. *CPT: Pharmacometrics & Systems*
Pharmacology 8, 567–576. doi:10.1002/psp4.12411

Strauch, K., Lutz, U., Bittner, N., and Lutz, W. K. (2009). Dose–response relationship for the
pharmacokinetic interaction of grapefruit juice with dextromethorphan investigated by human urinary
metabolite profiles. *Food and Chemical Toxicology* 47, 1928–1935. doi:10.1016/j.fct.2009.05.004

Takashima, T., Murase, S., Iwasaki, K., and Shimada, K. (2005). Evaluation of Dextromethorphan
Metabolism Using Hepatocytes from CYP2D6 Poor and Extensive Metabolizers. *Drug Metabolism and*
Pharmacokinetics 20, 177–182. doi:10.2133/dmpk.20.177

Tamminga, W., Wemer, J., Oosterhuis, B., de Zeeuw, R., de Leij, L., and Jonkman, J. (2001). The
prevalence of CYP2D6 and CYP2C19 genotypes in a population of healthy Dutch volunteers. *European*
Journal of Clinical Pharmacology 57, 717–722. doi:10.1007/s002280100359

Taylor, C. P., Traynelis, S. F., Siffert, J., Pope, L. E., and Matsumoto, R. R. (2016). Pharmacology of
dextromethorphan: Relevance to dextromethorphan/quinidine (Nuedexta®) clinical use. *Pharmacology*
& Therapeutics 164, 170–182. doi:10.1016/j.pharmthera.2016.04.010

Tennezé, L., Verstuyft, C., Becquemont, L., Poirier, J. M., Wilkinson, G. R., and Funck-Brentano,
C. (1999). Assessment of CYP2D6 and CYP2C19 activity in vivo in humans: A cocktail study
with dextromethorphan and chloroguanide alone and in combination. *Clinical Pharmacology and*
Therapeutics 66, 582–588. doi:10.1053/cp.1999.v66.103401001

Tricco, A. C., Lillie, E., Zarin, W., O'Brien, K. K., Colquhoun, H., Levac, D., et al. (2018). PRISMA
Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Annals of Internal Medicine*
169, 467–473. doi:10/gfd8vk

Vander, J. S., A. (2001). Human physiology: The mechanisms of body function. *McGraw-Hill higher*
education

von Moltke, L. L., Greenblatt, D. J., Grassi, J. M., Granda, B. W., Venkatakrishnan, K., Schmider, J., et al.
(1998). Multiple human cytochromes contribute to biotransformation of dextromethorphan in-vitro:
Role of CYP2C9, CYP2C19, CYP2D6, and CYP3A. *The Journal of Pharmacy and Pharmacology* 50,
997–1004. doi:10.1111/j.2042-7158.1998.tb06914.x

[Dataset] Welsh, C., Xu, J., Smith, L., König, M., Choi, K., and Sauro, H. M. (2022). libRoadRunner 2.0:
A High-Performance SBML Simulation and Analysis Library. doi:10.48550/arXiv.2203.01175

Whirl-Carrillo, M., Huddart, R., Gong, L., Sangkuhl, K., Thorn, C. F., Whaley, R., et al. (2021). An
evidence-based framework for evaluating pharmacogenomics knowledge for personalized medicine.
Clinical pharmacology and therapeutics 110, 563–572. doi:10.1002/cpt.2350

Wyen, C., Fuhr, U., Frank, D., Aarnoutse, R. E., Klaassen, T., Lazar, A., et al. (2008). Effect of an
antiretroviral regimen containing ritonavir boosted lopinavir on intestinal and hepatic CYP3A, CYP2D6

and P-glycoprotein in HIV-infected patients. *Clinical pharmacology and therapeutics* 84, 75–82.
doi:10.1038/sj.clpt.6100452

Yamazaki, T., Desai, A., Goldwater, R., Han, D., Howieson, C., Akhtar, S., et al. (2017). Pharmacokinetic
Effects of Isavuconazole Coadministration With the Cytochrome P450 Enzyme Substrates Bupropion,
Repaglinide, Caffeine, Dextromethorphan, and Methadone in Healthy Subjects. *Clinical pharmacology*
in drug development 6, 54–65. doi:10.1002/cpdd.281

Yang, J., He, M. M., Niu, W., Wrighton, S. A., Li, L., Liu, Y., et al. (2012). Metabolic capabilities
of cytochrome P450 enzymes in Chinese liver microsomes compared with those in Caucasian liver
microsomes: Hepatic cytochromes P450 of Chinese and Caucasians. *British Journal of Clinical*
Pharmacology 73, 268–284. doi:10.1111/j.1365-2125.2011.04076.x

Yu, A. and Haining, R. L. (2001). Comparative contribution to dextromethorphan metabolism by
cytochrome P450 isoforms in vitro: Can dextromethorphan be used as a dual probe for both CTP2D6 and
CYP3A activities? *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 29, 1514–1520

Zackrisson, A. L., Lindblom, B., and Ahlner, J. (2010). High Frequency of Occurrence of CYP2D6
Gene Duplication/Multiduplication Indicating Ultrarapid Metabolism Among Suicide Cases. *Clinical*
Pharmacology & Therapeutics 88, 354–359. doi:10.1038/clpt.2009.216

Zanger, U. M., Raimundo, S., and Eichelbaum, M. (2004). Cytochrome P450 2D6: Overview and update on
pharmacology, genetics, biochemistry. *Naunyn-Schmiedeberg's archives of pharmacology* 369, 23–37.
doi:10.1007/s00210-003-0832-2

Zawertailo, L. A., Tyndale, R. F., Busto, U., and Sellers, E. M. (2010). Effect of metabolic blockade on the
psychoactive effects of dextromethorphan. *Human Psychopharmacology: Clinical and Experimental* 25,
71–79. doi:10.1002/hup.1086

Figure 1. PBPK model of dextromethorphan (DXM). A) whole body model consisting of liver, kidney, intestine, forearm, lung and the rest compartment. Organs with minor relevance are not modeled explicitly and lumped into the rest compartment. Organs are coupled via the systemic circulation with arrow width proportional to relative blood flow. DXM can be administered intravenously (IV) or orally (PO) with DXM appearing in the venous blood or intestine, respectively. **B)** intestine model consisting of dissolution, absorption and excretion of DXM. Only a fraction of DXM is absorbed with the remainder excreted in the feces. First pass metabolism of DXM via CYP3A4 in the intestine reduces the amount of DXM appearing in the circulation. **C)** liver model consisting of DXM → DXO conversion via CYP2D6 and CYP3A4 and subsequent glucuronidation to DXO-Glu. **D** kidney model for the urinary excretion of DXM, DXO and DXO-Glu. Created with BioRender.com.

Figure 2. Model of CYP3A4 and CYP2D6. A) CYP3A4 and CYP2D6 distributions. Conversion of DXM → DXO via CYP3A4 and CYP2D6 are modeled via Michaelis-Menten kinetics. Variability was included via two-dimensional lognormal distributions of Michaelis-Menten coefficient (K_m) and maximum rate of reaction (V_{max}). The distribution parameters were determine by fitting to *in vitro* data in human liver microsomes. Variability of CYP3A4 parameters was measured by midazolam (Yang et al., 2012), variability of CYP2D6 parameters via DXM (Yang et al., 2012; Storelli et al., 2019a). To transfer the CYP3A4 data from midazolam to DXM normalized values were used. The distribution of CYP2D6 was modeled as a mixture model of the underlying activity scores as depicted in B. The model CYP3A4 and CYP2D6 distributions were sampled with each point corresponding to a combination of V_{max} and K_m. CYP2D6 data was color-coded by the respective activity score. **B)** CYP2D6 activity score model. CYP2D6 activity was modeled via a mixture model of individual activity scores. With increasing activity score the V_{max} for the DXM → DXO conversion increases and the μ_{Km} for DXM decreases, i.e., reaction velocity and affinity for the substrate increase. The table provides AS, genetic phenotype (gPT), mean V_{max}, mean K_m, and AS frequency in curated UCMR data (P(AS)). In case of AS=0.0 CYP2D6 has no activity for the DXM → DXO conversion.

Figure 3. Time-dependency of DXM pharmacokinetics by activity score. A) DXM plasma concentration, **B)** DXO plasma concentration, **C)** DXM/DXO plasma ratio, **D)** UCMR (DXM/(DXO+DXO-Glu) in urine) Depicted is the subset of data in which 30 mg of DXM was applied orally. The upper rows in the panels depict the data in healthy adults from (Abdelrahman et al., 1999; Chen et al., 2017; Dorado et al., 2017; Frank, 2009; Gaedigk, 2013; Köhler et al., 1997; López et al., 2005; Montané Jaime et al., 2013; Myrand et al., 2008; Tamminga et al., 2001). Cocktail studies are included. Studies containing coadminstrations with established drug-drug interactions are excluded. The lower rows depict the respective simulation results. To visualize the large variability in the simulation box plots showing the quartiles along side the median and outliers for selected time points are used. Variables changed in the simulation are the CYP3A4 and CYP2D6 reaction parameters K_m and V_{max} according to the distributions in Fig. 2. For the different activity scores the respective CYP2D6 activity score model was used.

Figure 4. Dextromethorphan (DXM) concentration in plasma or serum. Studies were simulated according to the reported dosing protocol. In case of available activity score information the clinical data is color coded accordingly. Information on metabolizer phenotype (UM, EM, IM, PM) is provided where reported. Data from (Abduljalil et al., 2010; Armani et al., 2017; Capon et al., 1996; Doroshyenko et al., 2013; Duedahl et al., 2005; Dumond et al., 2010; Edwards et al., 2017; Eichhold et al., 1997, 2007; Frank, 2009; Köhler et al., 1997; Lenuzza et al., 2016; Nakashima et al., 2007; Nyunt et al., 2008; Oh et al., 2012; Qiu et al., 2016; Schadel et al., 1995; Tennezé et al., 1999; Yamazaki et al., 2017; Zawertailo et al., 2010).

Figure 5. Dextrorphan (DXO) concentration in plasma or serum. Studies were simulated according to the reported dosing protocol. In case of available activity score information the clinical data is color coded accordingly. Information on metabolizer phenotype (UM, EM, IM, PM) is provided where reported. Data from (Abduljalil et al., 2010; Eichhold et al., 1997, 2007; Frank, 2009; Lenuzza et al., 2016; Nakashima et al., 2007; Oh et al., 2012; Schadel et al., 1995; Zawertailo et al., 2010).

Figure 6. Total dextrorphan (DXO+DXO-Glu) concentration in plasma or serum. Studies were simulated according to the reported dosing protocol. In case of available activity score information the clinical data is color coded accordingly. Information on metabolizer phenotype (UM, EM, IM, PM) is provided where reported. Data from (Capon et al., 1996; Demirbas et al., 1998; Hu et al., 2011; Köhler et al., 1997; Tennezé et al., 1999; Yamazaki et al., 2017; Zawertailo et al., 2010).

Figure 7. Dextromethorphan (DXM) amount in urine. Studies were simulated according to the reported dosing protocol. In case of available activity score information the clinical data is color coded accordingly. Information on metabolizer phenotype (UM, EM, IM, PM) is provided where reported. Data from (Abduljalil et al., 2010; Barnhart, 1980; Frank, 2009; Jones et al., 1996; Nagai et al., 1996; Schadel et al., 1995).

Figure 8. Total dextrorphan (DXO+DXO-Glu) amount in urine. Studies were simulated according to the reported dosing protocol. In case of available activity score information the clinical data is color coded accordingly. Information on metabolizer phenotype (UM, EM, IM, PM) is provided where reported. Data from (Abduljalil et al., 2010; Barnhart, 1980; Frank, 2009; Jones et al., 1996).

Figure 9. Cumulative metabolic ratio between dextormethorphan and total dextrorphan (DXM/(DXO+DXO-GLU)) in urine (UCMR). Studies were simulated according to the reported dosing protocol. In case of available activity score information the clinical data is color coded accordingly. Information on metabolizer phenotype (UM, EM, IM, PM) is provided where reported. Data from (Abdelrahman et al., 1999; Chládek et al., 2000; Dorado et al., 2017; Frank, 2009; Gaedigk, 2013; Hou et al., 1991; Jones et al., 1996; Köhler et al., 1997; López et al., 2005; Montané Jaime et al., 2013; Myrand et al., 2008; Nagai et al., 1996; Tamminga et al., 2001). The metabolic phenotype definitions for UM, EM, IM, PM are depicted as gray areas.

Figure 10. Sensitivity analysis of UCMR by activity score. A) Dependency of UCMR (urinary cumulative ratio of DXM/(DXO-Glu) after 8 hours and 30 mg oral DXM) on selected physiological parameters and the DXM dose. Parameter scans were performed for all activity scores. Reference model parameters are depicted as dashed lines. **B)** Sensitivity analysis of model parameters. To systematically study the effect of parameter changes the local sensitivity of UCMR were calculated for all activity scores. Parameters were varied 10% in both direction around the reference parameter value and the relative change of UCMR was calculated (insensitive parameters with relative change of UCMR smaller than 1% were omitted). Positive sensitivities are depicted in red, negative sensitivities in blue. Parameters were sorted via agglomerative clustering. Representative parameters of the clusters (i.e. liver volume per bodyweight, cardiac output per bodyweight, DXM tissue/plasma partition coefficient, and oral DXM dose) are depicted in A. The local sensitivity for the activity scores in B corresponds to the normalized slope at the dashed lines in A.

Figure 11. CYP2D6 genotype-, activity score association of the UCMR. Simulation of urinary cumulative ratio of DXM/(DXO-Glu) (UCMR) based on activity score frequencies. UCMR data was measured at least 4 hours after the application of DXM (hydrobromide) in healthy adults. Cocktail studies were included in the analysis. Studies containing coadminstrations with established drug-drug interactions were excluded. The ranges for metabolic phenotypes (UM, EM, IM, PM) are depicted as gray shaded areas. For timecourse UCMRs, only the latest measurement after administration was included. Data from (Abdelrahman et al., 1999; Dorado et al., 2017; Frank, 2009; Gaedigk, 2013; Köhler et al., 1997; López et al., 2005; Montané Jaime et al., 2013; Myrand et al., 2008; Tamminga et al., 2001). **A)** Histogram of UCMR data stratified by CYP2D6 activity score. **B)** Corresponding simulation results (UCMR at 8hr) from the Monte Carlo simulation with random variables being the enzyme reaction parameter (i.e. K_m, V_{max}). See details in Fig. 2. **C)** Empirical CMFs stratified by the activity scores. **D)** Corresponding simulated CMFs stratified by CYP2D6 activity scores. **E)** Box plots of observed UCMRs stratified by CYP2D6 activity scores. **F)** Box plots of simulated UCMRs stratified by the activity scores. **G)** Box plots of observed UCMRs stratified by CYP2D6 diplotypes. **H)** Box plots of simulated UCMRs stratified by CYP2D6 diplotypes. For D, F, and H, 2000 samples were simulated for each activity score whereas in B and D a two-fold oversampling with the CYP2D6 activity score frequencies from the UCMR data was performed.

Figure 12. UCMR distributions for biogeographical populations A) Simulated UCMR distributions at 8 hours for various biogeographical populations based on reported CYP2D6 activity score frequencies as reported in PharmGKB (Whirl-Carrillo et al., 2021). Frequencies are provided in the Supplementary Tab. S2. **B)** Reported UCMRs depending on activity score for Caucasians and East Asians from (Abdelrahman et al., 1999; Frank, 2009; Gaedigk, 2013; Köhler et al., 1997; Myrand et al., 2008). Cocktail studies were included in the analysis. Studies containing coadminstrations with established drug-drug interactions were excluded. **C)** Simulated activity score frequency and metabolic phenotype frequency for the populations and comparison with data for Caucasian and East Asian populations (hatched bars).

Table 1   Clinical studies with pharmacokinetics used for model evaluation. NR: not reported, DXM: dextromethorphan.

Reference	PK-DB	PMID	DXM application	Dosing protocol	Description
Abdelrahman et al. (1999)	PKDB00573	10340911	DXM	Oral (syrup): 0.3 mg/kg	Investigation of terbinafine as a CYP2D6 inhibitor in vivo.
Abduljalil et al. (2010)	PKDB00574	20881950	DXM hydrobromide	Oral (capsule): 30 mg	Assessment of activity levels for CYP2D6*1, CYP2D6*2, and CYP2D6*41 genes by population pharmacokinetics of dextromethorphan.
Armani et al. (2017)	PKDB00428	10340911	DXM (in cocktail)	Oral (NR): 30 mg	The antitussive effect of dextromethorphan in relation to CYP2D6 activity.
Barnhart (1980)	PKDB00575	7423506	DXM hydrobromide	Oral (capsule): 30 mg	The urinary excretion of dextromethorphan and three metabolites in dogs and humans.
Capon et al. (1996)	PKDB00576	8841152	DXM hydrobromide	Oral (NR): 30 mg	The antitussive effect of dextromethorphan in relation to CYP2D6 activity.
Chen et al. (2017)	PKDB00577	28512430	DXM	Oral (tablet): 15 mg + water 300ml	CYP2D6 phenotyping using urine, plasma, and saliva metabolic ratios to assess the impact of CYP2D6*10 on inter-individual variation in a Chinese population.
Chládek et al. (2000)	PKDB00578	11214771	DXM hydrobromide	Oral (syrup): 30 mg	In-vivo indices of CYP2D6 activity: comparison of dextromethorphan metabolic ratios in 4-h urine and 3-h plasma.
Demirbas et al. (1998)	PKDB00579	9840216	DXM hydrobromide	Oral (sustained release tablet): 60 mg	Bioavailability of dextromethorphan (as dextrorphan) from sustained release formulations in the presence of guaifenesin in human volunteers.
Dorado et al. (2017)	PKDB00580	28271978	DXM	Oral (NR): 15 mg	Lessons from Cuba for global precision medicine: CYP2D6 genotype is not a robust predictor of CYP2D6 ultrarapid metabolism.
Doroshyenko et al. (2013)	PKDB00138	23401474	DXM (in cocktail)	Oral (capsule): 30 mg	Drug metabolism and disposition: the biological fate of chemicals.
Duedahl et al. (2005)	PKDB00597	15661445	DXM	Intravenous: 0.5 mg/kg	Intravenous dextromethorphan to human volunteers: relationship between pharmacokinetics and anti-hyperalgesic effect.
Dumond et al. (2010)	PKDB00499	20147896	DXM (in cocktail)	Oral (solution): 30 mg	A phenotype-genotype approach to predicting CYP450 and P-glycoprotein drug interactions with the mixed inhibitor/inducer tipranavir/ritonavir.
Edwards et al. (2017)	PKDB00496	28808886	DXM (in cocktail)	Oral (capsule): 30 mg	Assessment of pharmacokinetic interactions between obeticholic acid and caffeine, midazolam, warfarin, dextromethorphan, omeprazole, rosuvastatin, and digoxin in phase 1 studies in healthy subjects.
Eichhold et al. (1997)	PKDB00596	-	DXM hydrobromide	Oral (syrup): 30 mg	Determination of dextromethorphan and dextrorphan in human plasma by liquid chromatography/tandem mass spectrometry.
Eichhold et al. (2007)	PKDB00581	16930908	DXM hydrobromide	Oral (solution): 20 mg	Simultaneous determination of dextromethorphan, dextrorphan, and guaifenesin in human plasma using semi-automated liquid/liquid extraction and gradient liquid chromatography tandem mass spectrometry.
Frank (2009)	PKDB00582	-	DXM hydrobromide (in cocktail)	Oral (capsule): 30 mg	Evaluation of pharmacokinetic metrics for phenotyping of the human CYP2D6 enzyme with dextromethorphan.
Gaedigk (2013)	PKDB00583	24151800	DXM	Oral (syrup): 0.3 mg/kg	Complexities of CYP2D6 gene analysis and interpretation.
Hou et al. (1991)	PKDB00584	2015730	DXM hydrobromide	Oral (capsule): 50 mg	Salivary analysis for determination of dextromethorphan metabolic phenotype.
Hu et al. (2011)	PKDB00585	21050887	DXM hydrobromide	Oral (sustained release tablet): 30 mg	Floating matrix dosage form for dextromethorphan hydrobromide based on gas forming technique: in vitro and in vivo evaluation in healthy volunteers.
Jones et al. (1996)	PKDB00586	8873685	DXM hydrobromide	Oral (syrup): 30 mg	Determination of cytochrome P450 3A4/5 activity in vivo with dextromethorphan N-demethylation.
Köhler et al. (1997)	PKDB00587	9429230	DXM	Oral (syrup): 20 mg	CYP2D6 genotype and phenotyping by determination of dextromethorphan and metabolites in serum of healthy controls and of patients under psychotropic medication.
López et al. (2005)	PKDB00588	16249913	DXM hydrobromide	Oral (syrup): 30 mg	CYP2D6 genotype and phenotype determination in a Mexican Mestizo population.

(continued)

Reference	PK-DB	PMID	DXM application	Dosing protocol	Description
Lenuzza et al. (2016)	PKDB00598	25465228	DXM (in cocktail)	Oral (tablet): 18 mg	Safety and pharmacokinetics of the (CIME) Combination of Drugs and Their Metabolites after a single oral dosing in healthy volunteers.
Montané Jaime et al. (2013)	PKDB00589	23394389	DXM hydrobromide	Oral (NR): 30 mg + water	Characterization of the CYP2D6 gene locus and metabolic activity in Indo- and Afro-Trinidadians: discovery of novel allelic variants.
Myrand et al. (2008)	PKDB00497	18231117	DXM (in cocktail)	Oral (NR): 30 mg	Pharmacokinetics/genotype associations for major cytochrome P450 enzymes in native and first- and third-generation Japanese populations: comparison with Korean, Chinese, and Caucasian populations.
Nagai et al. (1996)	PKDB00590	8830977	DXM hydrobromide	Oral (tablet): 30 mg	Pharmacokinetics and polymorphic oxidation of dextromethorphan in a Japanese population.
Nakashima et al. (2007)	PKDB00599	17652181	DXM hydrobromide	Oral (tablet): 30 mg	Effect of cinacalcet hydrochloride, a new calcimimetic agent, on the pharmacokinetics of dextromethorphan: *in vitro* and clinical studies.
Nyunt et al. (2008)	PKDB00591	18362694	DXM	Oral (tablet): 30 mg	Pharmacokinetic effect of AMD070, an Oral CXCR4 antagonist, on CYP3A4 and CYP2D6 substrates midazolam and dextromethorphan in healthy volunteers.
Oh et al. (2012)	PKDB00054	22483397	DXM (in cocktail)	Oral (NR): 2 mg	High-sensitivity liquid chromatography-tandem mass spectrometry for the simultaneous determination of five drugs and their cytochrome P450-specific probe metabolites in human plasma.
Pope et al. (2004)	PKDB00592	15342614	DXM	Oral (capsule): 30 mg; 45 mg; 60mg	Pharmacokinetics of dextromethorphan after single or multiple dosing in combination with quinidine in extensive and poor metabolizers.
Qiu et al. (2016)	PKDB00600	27023460	DXM hydrobromide	Oral (tablet): 15 mg	Effects of the Chinese herbal formula "Zuojin Pill" on the pharmacokinetics of dextromethorphan in healthy Chinese volunteers with CYP2D6*10 genotype.
Schadel et al. (1995)	PKDB00593	7593709	DXM	Oral (capsule): 30 mg	Pharmacokinetics of dextromethorphan and metabolites in humans: influence of the CYP2D6 phenotype and quinidine inhibition.
Schoedel et al. (2012)	PKDB00594	22283559	DXM	Oral (capsule): twice daily for 8 days; 30 mg	Randomized open-label drug-drug interaction trial of dextromethorphan/quinidine and paroxetine in healthy volunteers.
Tamminga et al. (2001)	PKDB00498	11829201	DXM hydrobromide	Oral (tablet): 22 mg	The prevalence of CYP2D6 and CYP2C19 genotypes in a population of healthy Dutch volunteers.
Yamazaki et al. (2017)	PKDB00494	27273149	DXM (in cocktail)	Oral (NR): 30 mg	Pharmacokinetic Effects of isavuconazole coadministration with the cytochrome P450 enzyme substrates bupropion, repaglinide, caffeine, dextromethorphan, and methadone in healthy subjects.
Zawertailo et al. (2010)	PKDB00595	20041473	DXM	Oral (capsule): 3 mg/kg	Effect of metabolic blockade on the psychoactive effects of dextromethorphan.

Table 2. Model parameters in PBPK model of DXM. The complete information is available from the model repository. The prefixes GU__, LI__,KI__, correspond to the intestine/gut, liver, and kidneys, respectively. Values are either adopted from the references or fitted (**F**). During the robustness analysis of UCMR, various parameters were scanned (**S**) and a local sensitivity (**SA**) was performed, see section 3.5.

Parameter	Description	References	Value	Unit	F	S	SA
BW	Body weight	ICRP (2002) (male)	75	kg			✓
HEIGHT	Height	ICRP (2002) (male)	170	cm			✓
HR	Heart rate		70	1/min			✓
HRrest	Heart rate (resting)		70	1/min			✓
COBW	Cardiac output per bodyweight	ICRP (2002); de Simone et al. (1997)	1.548	ml/s/kg		✓	✓
HCT	Hematocrit	Vander (2001); Herman (2016) (upper range male)	0.51	-			
Kp_fo_dxm	Tissue/plasma partition coefficient DXM forearm		10	-	✓		✓
f_shunting_forearm	Shunting in forearm		0.2795	-	✓		
FVgu	Gut fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.0171	l/kg			✓
FVki	Kidney fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.0044	l/kg			✓
FVli	Liver fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.021	l/kg		✓	✓
FVlu	Lung fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.0076	l/kg			✓
FVsp	Spleen fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.0026	l/kg			✓
FVpa	Pancreas fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.01	l/kg			✓
FVfo	Fore arm fractional tissue volume		0.0048	l/kg	✓		✓
FVve	Venous fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.0514	l/kg			✓
FVar	Arterial fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.0257	l/kg			✓
FVpo	Portal fractional tissue volume	Jones and Rowland-Yeo (2013); ICRP (2002)	0.001	l/kg			✓
FQgu	Gut fractional tissue blood flow	Jones and Rowland-Yeo (2013)	0.146	-			✓
FQki	Kidney fractional tissue blood flow	Jones and Rowland-Yeo (2013)	0.19	-			✓
FQh	Hepatic (venous side) fractional tissue blood flow	Jones and Rowland-Yeo (2013)	0.215	-			
FQlu	Lung fractional tissue blood flow	Jones and Rowland-Yeo (2013)	1	-			✓
FQsp	Spleen fractional tissue blood flow	Jones and Rowland-Yeo (2013)	0.017	-			✓
FQfo	Fore arm fractional tissue blood flow	RNAO (2022)	0.0146	-			✓
FQpa	Pancreas fractional tissue blood flow	ICRP (2002)	0.017	-			✓
ftissue_dxm	Vmax tissue distribution DXM		1000	l/min	✓		✓
Kp_dxm	Tissue/plasma partition coefficient DXM		8.7346	-	✓	✓	✓
Ka_dis_dxm	DXM rate of dissolution & stomach passage		0.0217	1/hr	✓		✓
Mr_dxo	Molecular weight DXO	CHEBI:29133	257.3707	g/mole			
ftissue_dxo	Vmax tissue distribution DXO		100	l/min	✓		✓
Kp_dxo	Tissue/plasma partition coefficient DXO		4	-	✓		✓
Mr_dxo_glu	Molecular weight DXO_glu	CHEBI:32645	433.4948	g/mole			
ftissue_dxo_glu	Vmax tissue distribution DXO_glu		3	l/min	✓		✓
Kp_dxo_glu	Tissue/plasma partition coefficient DXO_glu		0.08	-	✓		✓
KI__DXMEX_k	DXM urinary excretion rate		0.017	1/min	✓		✓
KI__DXOEX_k	DXO urinary excretion rate		0.3	1/min	✓		✓
KI__DXOGLUEX_k	DXO glucuronide urinary excretion rate		10	1/min	✓		✓
LI__DXMCYP2D6_Vmax	DXM CYP2D6 Vmax		0.003	mmol/min/l	✓		✓
LI__DXMCYP2D6_Km	DXM CYP2D6 Km	Storelli et al. (2019a); Yang et al. (2012)	0.0079	mM			✓
LI__cyp2d6_ac	CYP2D6 activity score		0.0 - 3.0	-			✓
LI__lambda_1	Slope of Km by principal component regression of (Km, Vmax) in log space.	Storelli et al. (2019a); Yang et al. (2012)	-0.4	-	✓		
LI__DXMCYP3A4_Vmax	Vmax of DXO formation by CYP3A4		0.0004	mmol/min/l	✓		✓
LI__DXMCYP3A4_Km	Km of DXO formation by CYP3A4	Yu and Haining (2001)	0.157	mM			✓
LI__DXOUGT_Vmax	DXO UGT Vmax (glucuronidation)		0.8953	mmol/min/l	✓		
LI__DXOUGT_Km	DXO UGT Km (glucuronidation)	Lutz and Isoherranen (2012)	0.69	mM			
GU__F_dxm	Fraction absorbed DXM	Schadel et al. (1995)	0.55	-			✓
GU__Ka_abs_dxm	Ka_abs absorption DXM		3.4285	1/hr	✓		✓
GU__DXMCYP3A4_Vmax	DXM CYP3A4 Vmax		0.0002	mmol/min/l	✓		✓
GU__DXMCYP3A4_Km	DXM CYP3A4 Km	Kerry et al. (1994); Yu and Haining (2001)	0.7	mM			✓
PODOSE	DXM oral dose			mg		✓	✓