Individual differences in information processing during
sleep and wake predict sleep-based memory consolidation
of complex rules

Madison Richter[1,2]*, Zachariah R. Cross[1,3], Ina Bornkessel-Schlesewsky[1]

[1]Cognitive Neuroscience Laboratory – Australian Research Centre for Interactive and Virtual
Environments, University of South Australia, Adelaide, Australia.

[2]College of Nursing and Health Sciences, Flinders University, Adelaide, Australia.

[3]Department of Medical Social Sciences, Northwestern Feinberg School of Medicine,
Chicago, Illinois, United States of America.

*Corresponding author: madison_richter@outlook.com

ORCID: https://orcid.org/0000-0001-7411-368X

Manuscript details

Number of pages: 30

Number of figures: 4

Abstract word count: 264

Introduction word count: 1893

Discussion word count: 2034

Overall word count: 7621

18 Data available at:
19 https://osf.io/f5ksg/?view_only=738200b63e78433eb50e667ea772c08f

20 Code available at: https://tinyurl.com/turbidus

Funding and Acknowledgements: This work was supported by the Research Network for Undersea Decision Superiority 2020 Honours Scholarship awarded to MR and by an Australian Research Council Future Fellowship (FT160100437) awarded to IBS. ZC was also supported by Australian Commonwealth Government funding under the Research Training Program (RTP; number 212190) and Maurice de Rohan International Scholarship awarded to ZC. The authors would like to thank the research assistants Lena Zou, Erica Wilkinson, Nicole Vass, and Angela Osborn for help with data collection.

21 **Conflict of Interest**

22 The authors declare no competing financial interests.

23   **Abstract**

Memory is critical for many cognitive functions, from remembering facts, to learning

complex environmental rules. While memory encoding occurs during wake, memory

consolidation is associated with sleep-related neural activity. Further, research

suggests that individual differences in alpha frequency during wake (~ 7 – 13 Hz)

modulate memory processes, with higher individual alpha frequency (IAF)

associated with greater memory performance. However, the relationship between

wake-related EEG individual differences, such as IAF, and sleep-related neural

correlates of memory consolidation has been largely unexplored, particularly in a

complex rule-based memory context. Here, we aimed to investigate whether wake-

derived IAF and sleep neurophysiology interact to influence rule learning in a sample

of 35 healthy adults (16 males; mean age = 25.4, range: 18 – 40). Participants

learned rules of a modified miniature language prior to either 8hrs of sleep or wake,

after which they were tested on their knowledge of the rules in a grammaticality

judgement task. Results indicate that sleep neurophysiology and wake-derived IAF

do not interact but modulate memory for complex linguistic rules separately. Phase-

amplitude coupling between slow oscillations and spindles during non-rapid eye-

movement (NREM) sleep also promoted memory for rules that were analogous to

the canonical English word order. As an exploratory analysis, we found that rapid

eye-movement (REM) sleep theta power at posterior regions interacts with IAF to

predict rule learning and proportion of time in REM sleep predicts rule learning

differentially depending on grammatical rule type. Taken together, the current study

provides behavioural and electrophysiological evidence for a complex role of NREM

and REM sleep neurophysiology and wake-derived IAF in the consolidation of rule-

based information.

Keywords: memory, rule extraction, sleep, individual alpha frequency, phase-
49   amplitude coupling

1 INTRODUCTION

Rule learning involves forming overarching rules via the extraction of
regularities from stimuli that share similar features (Frank & Gibson, 2011). For
example, grammar within natural language requires rule extraction, as critical
features of a sentence allow for effective categorisation and extraction of meaning
(Pothos, 2007). Despite important theoretical and practical implications for
understanding the consolidation of rule-based information, the underlying
mechanisms that facilitate rule consolidation are poorly understood. Current
understanding of rule extraction implicates sleep as an important factor (Batterink et
al., 2014; Nieuwenhuis et al., 2013) as well as individual differences in baseline
memory performance (Diekelmann et al., 2010). However, individual differences in
neurophysiology (cf. Cross et al., 2020a; Klimesch, 1999) as related to rule
extraction have yet to be explored in detail.

1.1 Memory consolidation as repeated reactivation

The consolidation and generalisation of memory occurs via repeated
reactivation of new and existing memory traces (Yassa & Reagh, 2013).
Generalisation refers to the process by which the brain extracts and consolidates the
central features amongst multiple memory traces, often losing rich contextual details
(Stark et al., 2008; Stark & Stark, 2016) and is associated with activity within
hippocampus and neocortex (García-Lázaro et al., 2012; Yassa & Reagh, 2013).
During generalisation, only central features that overlap between the new memory
trace and an existing memory trace are strengthened and the specific, contextual
details (e.g., features that do not overlap with the original memory trace), compete
for representation in neocortex and are less likely to be consolidated due to mutual
inhibition (Yassa & Reagh, 2013). Therefore, over repeated reactivations of similar
regularities (e.g., grammar within a language), memory traces become highly
semanticised, containing important facts or rules that can be applied to future stimuli
(Yassa & Reagh, 2013). This process of consolidation can occur during both periods
of wake and sleep (Humiston et al., 2019; Klinzing et al., 2019; Tucker et al., 2020;
Wamsley, 2019). In particular, the systems consolidation perspective proposes that
memory consolidation and reactivation relies upon slow-wave sleep (SWS;
Diekelmann & Born, 2010; Lewis & Durrant, 2011).

1.2 The influence of sleep on memory and rule extraction: A systems consolidation perspective

It is now well established that sleep – particularly activity during slow-wave sleep (SWS) – benefits memory consolidation (Diekelmann et al., 2010; Diekelmann & Born, 2010; Lewis & Durrant, 2011; Nieuwenhuis et al., 2013; Stickgold & Walker, 2013; Tononi & Cirelli, 2006). According to the Active Systems Consolidation model (ASC; Diekelmann & Born, 2010), slow oscillations (SOs), thalamo-cortical spindles, and hippocampal ripples facilitate long-term memory consolidation, allowing for newly encoded memories to undergo qualitative changes (e.g., rule extraction) as they are gradually incorporated into existing neocortical memory networks (Klinzing et al., 2019; Lewis & Durrant, 2011). Experimental evidence (Helfrich et al., 2018, 2019; Staresina et al., 2015) demonstrates that phase-amplitude coupling (PAC; Canolty & Knight, 2010) between SOs and faster oscillatory activity (e.g., spindles and hippocampal ripples) facilitates the reactivation of neural assemblies that were active during initial encoding (Diekelmann & Born, 2010; Helfrich et al., 2019; Muehlroth et al., 2019). Additionally, both thalamo-cortical spindles and hippocampal ripples are implicated in the controlled transfer of information, by priming cortical storage areas and modulating selective activation of neural assemblies, respectively (Girardeau & Zugaro, 2011; Staresina et al., 2015). ASC proposes that the PAC between SO and faster oscillatory activity drives the controlled hippocampal-neocortical information transfer required for memory consolidation (Diekelmann & Born, 2010) and more specifically, rule extraction.

Studies exploring rule extraction suggest that sleep facilitates the extraction of overarching rules from regularities within stimuli (Batterink et al., 2014; Djonlagic et al., 2009; Durrant et al., 2011; Gómez et al., 2006; Lerner & Gluck, 2019; Lutz et al., 2017; Nieuwenhuis et al., 2013; Wilhelm et al., 2013), such as grammar within language. For example, a recent review by Lerner & Gluck, (2019) demonstrated that NREM sleep is most associated with the extraction of regularities, however this association is often dependent on the particular task used and whether the regularity contains temporal aspects. Experimental studies investigating the effect of sleep on linguistic rule learning found that greater SWS duration predicted subsequent sensitivity to a hidden rule, as well as being instrumental in stabilising memory traces related to a rule (Batterink et al., 2014; Nieuwenhuis et al., 2013). Additionally,

studies in non-linguistic contexts have reported similar results, in which SWS
duration was positively associated with the extraction of statistical rules and
subsequent performance on a task in which understanding of the rule was crucial
(Djonlagic et al., 2009; Durrant et al., 2011, 2013). However, only a few experiments
have explored how precise oscillatory events (e.g., SO-spindle coupling) that occur
during SWS may predict rule extraction. The limited work in this area has found that
stronger SO activity is positively associated with motor sequence learning in both
children and adults (Wilhelm et al., 2013), that SO-spindle coupling increases from
childhood to adolescence, and this increase predicts improvements in motor memory
(Hahn et al., 2020). However, while this work provides preliminary evidence for a role
of sleep-based oscillatory activity in motor memory, little is known regarding the
influence of SO-spindle coupling in other memory contexts, such as rule learning.
Due to SO-spindle coupling being linked with general memory consolidation, further
investigation of such activity in a rule-based context may increase our understanding
of the underlying mechanisms involved in rule extraction.

1.3 Beyond slow-wave sleep: A possible role for REM sleep in rule
generalisation

In addition to cortical activity within SWS, Rapid Eye Movement (REM) sleep
and its associated oscillatory activity has often been implicated in memory
consolidation (Ackermann & Rasch, 2014; Boyce et al., 2016; Grosmark et al.,
2012). Few studies have investigated the role of neurophysiology during REM (e.g.,
theta activity; ~4–7 Hz) within rule-based contexts, typically focussing on the
influence of REM sleep duration on processes such as gist extraction (Batterink et
al., 2014; Matorina & Poppenk, 2019), which refers to the extraction of the most
important elements or the general idea (Matorina & Poppenk, 2021). For example,
Matorina & Poppenk (2019) found that a higher proportion of time spent in REM
sleep led to decreased performance in a statistical learning task, suggesting that
REM sleep is disadvantageous to the extraction of rules. In contrast, Batterink et al.,
(2014) examined how NREM and REM sleep duration during a nap contributed to
rule generalisation and found that performance was positively associated with the
interaction between proportion of time spent in REM and NREM sleep, indicating a
role for REM sleep in rule extraction.

Other studies have indicated a beneficial effect of REM sleep on rule extraction, specifically when looking at acquisition of a rule for effective categorisation (Djonlagic et al., 2009) and within a task in which effective learning of sequential rules was critical (Maquet et al., 2000). However, research into the specific influence of neurophysiology that occurs during REM sleep on rule extraction is still limited, requiring further exploration. Interestingly, work into the facilitation of memory generalisation has implicated factors other than specific sleep neurophysiology, such as individual differences in memory performance (Diekelmann et al., 2010). However, individual differences in neural activity and their relation to sleep-associated memory consolidation are only just starting to be considered (Chatburn et al., 2021; Cross et al., 2020a; Fenn & Hambrick, 2012; Schabus et al., 2008).

1.4 Individual alpha frequency as a measure of individual differences in information processing

Research into inter-individual differences in cognitive processing have often focussed on alpha activity during wake (~8-12Hz) and its relation to information processing in a variety of domains, such as language and memory (Bornkessel et al., 2004; Bornkessel-Schlesewsky et al., 2022; Cross, Santamaria, et al., 2020; Furman et al., 2018; Klimesch, 2012; Samaha & Postle, 2015). The individual alpha frequency (IAF), measured during wake, refers to the individual dominant peak within the alpha frequency band, and has been linked to various aspects of cognition such as attention, processing speed and memory (Klimesch, 1997, 1999; Surwillo, 1961). IAF has been implicated in the controlled timing of information transmission (Klimesch, 2012) and it is hypothesised that those with a higher IAF have improved information transmission within cortex, resulting in better performance on tests of memory (Klimesch, 1999) and faster processing speed (Klimesch et al., 1996; Ociepka et al., 2022; Surwillo, 1961, 1963). Samaha and Postle (2015) suggest that IAF during wake may be related to temporal receptive windows of processing, determining the amount of information within each processing cycle. Here, higher IAF individuals are considered to have faster cycles with less information contained within, and lower IAF individuals have slower cycles that contain more information (Samaha & Postle, 2015). In relation to memory, research indicates that those with good memory performance tend to have an IAF of around 1 Hz higher than poor memory performers (Doppelmayr et al., 2005; Klimesch, 1997, 1999).

The proposed role of IAF in processing speed and information transfer
described above may have implications for processes dependent on information
processing and cortical communication. Therefore, as sleep-dependent memory
consolidation is proposed to rely heavily on information transmission within the brain,
mechanisms suggested to reflect this process (e.g., SO-spindle coupling) may
interact with IAF due to their shared association with information processing and
transfer. Whilst there is little experimental evidence for an explicit association
between sleep neurophysiology and wake-state IAF, Cross et al., (2020a)
investigated how these factors may interact to influence emotional memory
consolidation. Results revealed complex interactions between wake-derived IAF,
REM theta power and SO density modulating emotional memory outcomes,
indicating a relationship between individual processing capabilities during wake and
cortical activity during sleep, providing initial support for a link between IAF and sleep
neurophysiology. However, the interaction between these variables in other memory-
related contexts (e.g., rule extraction) is yet to be examined.

1.5   The current study

The present study is a reanalysis of Cross et al. (2021) and aimed to address
whether sleep-related (e.g., SO-spindle coupling) and intrinsic neural differences in
information processing (e.g., IAF) modulate rule extraction. Due to the significance of
individual differences in memory performance, exploring IAF as a measure of
individual differences will inform current models of sleep and memory as to how
underlying neurophysiology of an individual interacts with memory consolidation and
thus, rule extraction. Exploratory analyses further aimed to clarify the role of REM
sleep and its involvement in rule-based memory consolidation. Subjects were
randomly allocated to a sleep or wake condition and were required to learn
grammatical rules in a novel artificial language modelled on Mandarin Chinese
(Cross et al., 2020b). Participants then completed a baseline and delayed sentence
judgment task (after sleep or an equivalent period of wake) used to test knowledge
about underlying grammatical rules, performance on which served as a proxy for rule
extraction.

Based on previous literature suggesting the role of sleep in facilitating memory
performance, it was hypothesised that (H[1]) those in the sleep condition will perform

better on tasks assessing grammatical knowledge than those in the wake condition.
Additionally, due to previous literature indicating a beneficial effect of sleep and IAF
on memory performance, it was hypothesised that (H^2) differences in memory
performance relative to condition (sleep vs. wake) and grammatical rule type will be
modulated by both SO-spindle coupling and IAF and that this relationship may be (a)
additive or (b) interactive. Additionally, exploratory analyses were conducted to
assess whether REM sleep is associated with the extraction of complex rules.

2   METHODS

2.1   Participants

Thirty-five (16 males) adults ranging from 18 to 40 years of age (M = 25.4, SD =
7.10) were randomly allocated to either a sleep (*Mean age* = 23.60 [SD = 5.84], 9
males) or wake (*Mean age* = 27.11 [SD = 7.50], 8 males) condition (17 in sleep
condition). Eligible participants were healthy, right-handed, had no current or past
psychiatric conditions (including intellectual impairment) or history of substance
dependence, normal or corrected-to-normal vision and were not taking medication
that influenced neurophysiological measures. Participants were instructed to not
consume caffeine prior to the experiment. In addition, eligible participants were
monolingual, native English speakers with no previous exposure to Mandarin
Chinese and did not have any current conditions that affected sleep or circadian
rhythms and had not undertaken shift work in the past month to ensure no influence
of irregular or poor-quality sleep. Participants received an AUD$120 honorarium
upon completion of the study. Ethics approval was granted by the University of South
Australia's Human Research Ethics committee (I.D: 0000032556).

2.2   Materials and Measures

2.2.1 Electroencephalography (EEG)

The EEG was recorded with a 32-channel BrainCap (Brain Products GmbH,
Gilching, Germany) with Ag/AgCl electrodes mounted via the International 10-20
system. The ground electrode was located at AFz and the online reference was
located at FCz, with re-referencing to linked mastoids undertaken offline. The
electrooculogram (EOG) was recorded using electrodes located at the side of each
eye as well as above and below the left eye. All EEG and EOG channels were

amplified via a BrainAmp DC amplifier (Brain Products GmbH, Gilching, Germany),
band-pass filter of DC – 250Hz and a sampling rate of 1000 Hz. Impedances were
kept below 10kΩ. EEG signals were recorded throughout both the sleep durations
and experimental tasks.

2.2.2 Language Learning Task: Mini Pinyin

Mini Pinyin is a modified miniature language (MML) modelled on Mandarin
Chinese (Cross et al., 2020b). MML paradigms contain words that are combined into
sentences following grammatical rules adapted from the language they are based
upon (Mueller, 2006; Wang et al., 2012). Mini Pinyin contains 16 verbs, 25 noun
phrases, 2 coverbs, and 4 classifiers (for a full description of Mini Pinyin, see Cross
et al., 2020b). The nouns are divided into categories of human (10 nouns), animal
(10 nouns), and objects (5 nouns).

Mini Pinyin contains two coverbs that influence word order and the
interpretation of the noun phrases. For example, when bǎ is present in a sentence,
the first noun is the Actor (i.e., the participant instigating the event) and the second
noun is the Undergoer (i.e., the participant affected by the event). The second
coverb, bèi, reverses this pattern. When either of these coverbs are used, word order
becomes fixed and determines role assignment (see Table 1 for examples). Thus, bǎ
constructions instantiate Actor-first word orders, while bèi instantiates an Undergoer-
first word order. Additionally, verb positioning is an important element in Mini Pinyin:
sentences that contain coverbs must have the verb placed at the end of the
sentence, whereas sentences without coverbs (not examined in this analysis) require
the verb to be positioned between the noun phrases.

Based on the linguistic parameters described above, the process of rule
extraction can be explored by using the word order contingencies (e.g., the fixed
word order of the Actor-Undergoer relationship) of each of the coverbs. Each coverb
and the associated rules can be used to examine rule extraction across differing
levels of novelty in comparison to English (i.e., the native language of the
participants). While both coverbs require participants to learn a novel verb
positioning rule in comparison to English, Actor-first bǎ-constructions show some
similarity to canonical English word order in that the Actor precedes the Undergoer.
Undergoer-first bèi-constructions, by contrast, differ from English word order in two

respects: the position of the verb and the relative order of the nouns. Accurate rule extraction would allow the participant to correctly distinguish between the specific noun phrase order instantiated by Undergoer-first (bèi) and Actor-first (bǎ) constructions as well as the correct position of the verb. Table 2 summarises the degree of novelty of each construction in comparison to English.

Table 1. Examples of plausible and implausible sentence structures (adapted from Cross et al., 2020b).

Coverb	Sentence Examples
Bǎ Plausible (Actor-first)	zhi junma bǎ xi pingguo chile. (animal) horse bǎ (small object) apple eat. *"The horse eats the small apple."*
Bǎ Implausible (Actor-first)	xi pingguo bǎ zhi junma chile. (small object) apple bǎ (animal) horse eat. *"The small apple eats the horse."*
Bèi Plausible (Undergoer-first)	da shubao bèi ge yisheng dale. (large object) bag bèi (human) doctor hit. *"The doctor hits the large bag."*
Bèi Implausible (Undergoer-first)	ge yisheng bèi da shubao dale. (human) doctor bèi (large object) bag hit. *"The large bag hits the doctor."*

Table 2. Summary of similarities (+) and differences (-) to English word order for Actor-first (Bǎ) and Undergoer-first (Bèi) coverb constructions within Mini Pinyin.

Sentence Type	Noun Order	Verb Position
Bǎ **(Actor-first)**	**+**	**-**
Bèi **(Undergoer-first)**	**-**	**-**

Note: Other sentence types in Mini Pinyin (not examined in this analysis) required the verb to be positioned similarly to English. As such, participants were required to learn the verb-final positioning rule that was unique to the sentences with coverbs.

2.3 Procedure

Each participant received a picture-word pair booklet containing each of the 25 nouns approximately one week before testing to gain a basic vocabulary before the experimental tasks. Eligible participants visited the Cognitive Neuroscience Laboratory at the University of South Australia to undergo testing. Before continuing onto the main testing phase, participants were required to score above 90% on a vocabulary test designed to test their knowledge of the 25 nouns. All 36 participants included in this analysis scored above 90%.

The main testing session consisted of a sentence learning task, a baseline
sentence judgement task, and a delayed judgement task (approximately 12 hours
after the learning and baseline tasks). All sentence and picture stimuli were
presented via OpenSesame (Mathôt et al., 2012). The learning task used pictures to
depict events involving two entities and was designed for the participants to learn the
meaning of the verbs, coverbs, classifiers, and overall sentence structure. Each trial
began with a 1000ms fixation cross, followed by pictures depicting the event.
Subsequently, a sentence describing the event pictured was presented on a word-
by-word basis, with each word presented for 700ms. The inter-stimulus interval (ISI)
was 200ms. This trial structure was used for a total of 128 pseudorandomised
picture-sentence combinations.

Following the learning task, participants completed the baseline and delayed
judgement tasks. Both judgment tasks consisted of 288 pseudorandomised novel
sentences (without pictures) presented word-by-word, with each word being
presented for 600ms each with an ISI of 200ms. Participants were instructed prior to
the task to judge if the sentence presented was grammatically correct by pressing a
button (cued by a question mark presented on screen for 4000ms). During the
baseline judgment task, participants also received feedback as to whether their
response was correct. They did not receive feedback in the delayed judgement task.
Two stimuli lists were used to counterbalance the order of stimuli presentation
across participants. Half of the sentences were grammatical constructions, and the
other half were ungrammatical constructions. See Figure 1 for a schematic of both
the learning and baseline judgement tasks.

Figure 1. Schematic representation of trials within the (A) learning and (B) baseline judgement tasks (adapted from Cross et al., 2021).

Note: The sentence presented in the top row (A) demonstrates a trial within the learning phase, in which the event picture is presented followed by the sentence explaining said event. Word presentation was longer (700ms) in this task to allow more time for participants to encode the rules. The bottom row (B) depicts a trial from the baseline judgment task, in which the sentence is presented for judgement, followed by a prompt for a judgement. After this, feedback was presented as to whether the judgement was correct or incorrect. If the participant did not response within 4000ms, the trial ended, and the response was coded as incorrect. In both A and B translates specifically to *"Animal bear bǎ small apple to eat"* or *"The bear ate the apple"*.

Participants in the wake condition completed the learning and baseline judgment tasks at approximately 09:00hr before leaving the laboratory to undergo their normal daily routine. They were instructed not to nap or engage in any language learning activities that may interfere with the task. Participants returned the same day to complete the delayed judgment task at approximately 21:00hr.

Participants in the sleep condition completed the learning and baseline judgment tasks at approximately 21:00hr after EEG set up. Following this, participants were given an 8-hour sleep opportunity between 23:00hr – 07:00hr. For the sleep opportunity, EMG was recorded, EOG channels were removed from the vertical position (above and below left eye) and placed horizontally (either side of each eye), and polysomnography was continuously recorded. Upon waking, participants were given approximately 1 hour to overcome sleep inertia and 1 hour for EEG and poor impedances to be fixed before completing the delayed judgement task (at approximately 09:00hr). For a schematic of the experimental protocol refer to Figure 2.

341 **Figure 2.** Schematic representation of experimental protocol (adapted from Cross et al. 2021).

342
Note: This figure shows the stepwise procedure for the sleep condition (above) and the wake condition (below).
The x-axis shows the time of day in hours as the experiment progressed, starting with 8am on the left and ending
with 11am (the following day) on the right. For the wake condition (below), participants began at approximately
08:00hr (am), underwent setup and initial testing (at 09:00hr) followed by a period of wake and subsequent
testing (at 21:00hr). For the sleep condition (above), participants began at approximately 20:00hr (pm),
underwent setup and initial testing (at 21:00hr) followed by a period of sleep and subsequent testing (at 09:00hr).
The coloured legend indicates the task being completed at the approximate time and the dotted line at 0
represents 00.00hrs (midnight) where testing crossed into the following day.

2.4   Data Analysis

All statistical analyses were performed in *R* (R Core Team, 2017). Mixed-

effects regression models were fitted using maximum likelihood via the *lme4* (Bates

et al., 2015). Other packages included in mixed-model analysis and output reporting

included the *car* (Fox & Weisberg, 2018), *Rmisc* (Hope, 2013), *effects* (Fox &

Weisberg, 2018), *Hmisc* (Harrell Jr, 2019), *apaTables* (Stanley & Spence, 2018) and

performance (Lüdecke et al., 2021). Data was arranged using *tidyverse* (Wickham et

al., 2019), *reshape2* (Wickham, 2007), *plyr* (Wickham, 2011), *sjmisc* (Lüdecke,

2018a), and packages involved in plotting included *lattice* (Sarkar, 2008), *sjPlot*

(Lüdecke, 2018b), *ggpubr* (Kassambara, 2020), *RColorBrewer* (Neuwirth, 2014). All

effects were plotted using the *ggplot2* package (Wickham et al., 2021) and *ggeffects*

(Lüdecke et al., 2020). Rayleigh's test for non-uniformity was run using *CircStats*

(Lund & Agostinelli, 2012). Links to the raw data can be found in the supplementary

material (Appendix D.1) and analysis scripts can be found at the following Open

Science Framework repository: https://tinyurl.com/turbidus.

2.4.1 Behavioural analysis

Performance on both the baseline and delayed tasks were quantified using d-

prime (Lockhart & Murdock, 1970; Wixted, 2014). d-prime is a measure of

discriminability and is defined as the difference between the z-transformed scores of

hit rate (HR) and false alarm rate (FA; i.e., d-prime = z[HR] – z[FA]). To calculate d-

prime, each sentence type was isolated and scored according to the criteria

described in Table 3. For example, calculating d-prime for bǎ verb position

sentences consisted of isolating bǎ verb position sentences and scoring them. Here,

grammatical bǎ sentences that were correctly identified as grammatical were scored

as a hit, whilst ungrammatical bǎ verb position sentences that were incorrectly

identified as grammatical were scored as a false alarm. Following this, the number of

hits and false alarms per subject were converted to a proportional value before being

z-transformed and applying the d-prime calculation. Raw hit and false alarm scores

for each participant per sentence type can be found in the supplementary material.

For example, for bǎ verb position sentences, a formula of z(HR(bǎ grammatical) –

z(FA(bǎ ungrammatical)) was applied. d-prime calculations for the baseline testing

session were also calculated to be added into the behavioural linear mixed-effects

models for comparison to delayed scores. Baseline scores were also added into

subsequent sleep neurophysiology models to control for baseline performance when

predicting delayed memory outcomes. Positive d-prime scores indicate good

memory performance, whereas negative scores indicate poor memory performance.

d-prime scores of 0 reflect chance performance.

Table 3. Description of how responses were scored to calculate d-prime for the delayed testing phase
for measurement of memory performance.

Scoring	Description
Hit	A grammatical sentence is correctly identified as grammatical.
Miss	A grammatical sentence is incorrectly identified as ungrammatical.
Correct rejection	An ungrammatical sentence is correctly identified as ungrammatical.
False alarm	An ungrammatical sentence is incorrectly identified as grammatical.

390 2.4.2 Sleep EEG analysis

Sleep EEG was pre-processed according to the specifications detailed in
(Cross et al., 2021). Briefly, sleep EEG was scored according to standardised criteria
(AASM; Berry et al., 2012) by two sleep technicians using Compumedics Profusion 3
software (Melbourne, Australia). A high-pass filter of 0.3 Hz and a low-pass filter of
35 Hz were applied for viewing. For the analysis, the EEG was filtered from 0.1 – 40
Hz and re-referenced to linked mastoids. Artifact rejection was performed
automatically, in which a covariance-based artifact rejection method was used with
hypnograms (Barachant, 2015; Barthélemy et al., 2019). Additionally, the EEG was
down-sampled from 1000 Hz to 100 Hz to reduce computational processing time.

Phase-amplitude coupling (PAC) events were detected using published
algorithms (Dvorak & Fenton, 2014; Helfrich et al., 2019; Staresina et al., 2015) via
the YASA toolbox implemented in MNE Python (Vallat & Walker, 2021). An event-
locked method was utilized, in which the SOs were detected initially, and spindle
events that occur within the detected SOs were subsequently identified. The PAC
strength between SOs and spindles was calculated using the Mean Vector Length
(MVL) method (Canolty et al., 2006). MVL is calculated by combining the amplitude
of the frequency for amplitude (i.e. spindle) with the phase of the frequency for phase
(i.e. SO), in which the combined signals become complex vectors (Canolty et al.,
2006; Hülsemann et al., 2019; Munia & Aviyente, 2019). The average length of these
vectors can be delineated, whereby the vector length represents the magnitude (or
strength) of the coupling event (Hülsemann et al., 2019).

For the REM theta analysis, power spectral density (PSD) estimates were
calculated on consecutive 30s epochs with a 4-second sliding window. Intervals
were tapered by a single Hanning window before applying a fast Fourier
transformation that resulted in interval power spectra with a frequency resolution of
0.25 Hz. Power spectra were then averaged across all blocks (Welch's method)
~~normalised by the effective noise bandwidth~~ to obtain PSD estimates for relative
theta power (4 – 8 Hz) during REM epochs for each channel (for a detailed
description of this approach, see Vallat & Walker, 2021).

2.4.3 Individual alpha frequency calculation

Individual alpha frequency (IAF) estimates for each subject were calculated
from eyes-closed resting-state EEG recordings according to the method detailed by
Corcoran et al., (2018), via the *philistine* package (Alday, 2019). Resting-state
recordings were taken at the beginning of the learning task and after the judgment
task for 2 minutes (eyes open and eyes closed). Estimates were taken using the
peak alpha frequency (PAF) method, in which the most prominent peak in the alpha
bandwidth was delineated. The PAF for each participant was estimated from
occipital and parietal channels (P3/P4/P7/P8/Pz/O1/O2/Oz) as this is where the
alpha rhythm is most prominent during resting-state eyes closed periods (Babu
Henry Samuel et al., 2018; Capotosto et al., 2017). For a detailed description of the
IAF estimation procedure, please see Corcoran et al., (2018).

2.4.4 Statistical analysis

Data were analysed using linear mixed-effects models via the *lme4* package
(Bates et al., 2015). For all models, Participant ID was specified as a random effect
on the intercept (Brown, 2020) and significance of effects were examined using Type
II Wald tests from the *car* package (Fox & Weisberg, 2018).

Model predictors (fixed effects) included Condition (sleep, wake), Coverb (bǎ,
bèi), Violation Type (noun phrase order, verb position), ~~Session (baseline, delayed)~~,
PAC events (quantified as MVL), IAF, and Baseline d-prime scores. d-prime scores
from the delayed testing session were specified as the outcome variable. Categorical
variables were sum-to-zero contrast coded, meaning that factor-level estimates are
relative to the grand mean (Schad et al., 2020). The first model was run exploring the
effect of Condition (sleep vs. wake), Coverb, Violation Type, baseline d-prime

scores, and IAF on memory performance. A second model was run to investigate the

effect of PAC, Coverb, Violation Type, baseline d-prime scores and IAF on memory

performance to explore neurophysiological factors in detail.

Two additional exploratory models were run to investigate the role of REM

sleep and associated theta power on memory performance. Predictors of the initial

model included IAF, Coverb, Violation Type, Baseline d-prime scores, REM Theta

power and the topographical components saggitality (anterior, central, posterior) and

laterality (left, midline, right), to explore oscillatory influences during REM on delayed

d-prime scores. An additional model was run with predictors of IAF, Coverb, Violation

Type, Baseline d-prime Scores and Percentage of Time in REM Sleep, to investigate

how REM sleep influences delayed d-prime scores. REM theta power and

percentage of time spent in REM were converted to z-scores.

3   RESULTS

3.1   Descriptive Statistics

Overall, the average d-prime score across sentences for the wake group were

.66 (*SD = .97*) at baseline testing and .65 (*SD = 1.34*) at delayed testing. For the

sleep condition, the average d-prime scores across sentence types were .60 (*SD =*

.85) at the baseline testing session and .87 (*SD* = 1.40) at the delayed testing

session (refer to Table 4 for d-prime scores per sentence type). Raw values for hits

and false alarms per subject and sentence condition can be found in the

supplementary material (Appendix C). Sleep variables including total sleep time,

sleep onset latency, wake after sleep onset, and the total amount of time spent in

each sleep stage are reported in Table 4, while Figure 3 shows a comodulogram and

circular histogram to illustrate the estimated PAC metrics. Additionally, a correlogram

showing pairwise comparison between variables of interest can be found in the

supplementary material (Appendix B, Figure 3).

Table 4. Mean d-prime scores (and standard deviations) for baseline and delayed testing sessions in
the sleep and wake groups for each sentence type.

Violation Type	Sleep		Wake	
	Baseline	Delayed	Baseline	Delayed
Bǎ				
Noun Order	.26 (.57)	.40 (1.13)	.16 (.52)	.15 (.86)
Verb Position	.86 (.80)	1.37 (1.26)	1.40 (.91)	1.20 (1.43)
Bèi				
Noun Order	.29 (.68)	.49 (1.18)	-.01 (.68)	.07 (1.03)
Verb Position	.98 (1.08)	1.25 (1.82)	1.09 (.94)	1.21 (1.54)

Note: Standard deviation is in parentheses.

Figure 3. Examples of phase-amplitude coupling metrics for visualisation purposes.

Note: A. Comodulogram illustrating the phase-amplitude coupling (calculated via Mean Vector Length) between
the SO and the spindle from a single participant at channel Cz. Mean Vector Length is calculated by combining
the amplitude of the frequency for amplitude (i.e. spindle) with the phase of the frequency for phase (i.e. SO) into
a complex vector, in which the average can be delineated to reflect the magnitude of the coupling event. Here,
the warmer colours (i.e., yellow) depicts where the coupling is strongest. As shown, this is occurring at
approximately 0.5 Hz for the frequency for phase (the SO, x-axis) and approximately 15 Hz for the frequency for
amplitude (the spindle, y-axis). **B.** Circular histogram of the preferred phase of SO to maximal spindle amplitude
averaged across all participants during NREM sleep at channel Cz. Here, each instance of phase-amplitude
coupling is instantiated along a circular distribution, allowing the ability to see the most common angle of the SO
phase whenever the spindle amplitude is maximal (between -π/2 and 0, closer to 0).

Table 5. Descriptive statistics (means, standard deviations, and ranges) for sleep parameters measured during the 8-hour sleep opportunity between baseline and delayed testing.

Sleep Stage	Mean Minutes (SD)	% of TST (SD)
N1	38.06 (29.93)	10.05 (8.34)
N2	196.29 (47.01)	49.52 (10.52)
SWS	104.24 (42.93)	25.85 (9.75)
REM	61.29 (40.01)	14.58 (8.70)
Sleep Parameter	**Mean Minutes (SD)**	**Range**
TST	399.88 (68.06)	203.5 – 472.5
SOL	15.24 (12.42)	1 – 42.5
WASO	52.65 (56.47)	0 - 225
EEG Parameter	**Mean (SD)**	
Mean Vector Length (MVL)	.23 (.01)	.21 - .27
REM Theta Power	.20 (.04)	.02 - .33

Note: SD = standard deviation; TST = total sleep time; SOL = sleep onset latency; WASO = wake after sleep onset; N1 = stage 1; N2 = stage 2; SWS = slow wave sleep; REM = rapid eye movement sleep; MVL = PAC metric used to quantify the strength of coupling between SOs and spindles.

3.2   Behavioural Analysis

The results reported in the following summarise the effects most pertinent to the current behavioural hypotheses, drawing on mixed model output summaries (for β values) and Type II Wald Chi-square test for chi-square and p values. Full model and Wald test summaries can be found in the supplementary material (Appendix A.1).

Mixed model analyses revealed a significant interaction between Condition, IAF and Baseline d' predicting d-prime at delayed testing, as shown in Figure 4A (refer to Table 6 for inferential statistical information for all main effects and significant interactions). Overall, a positive relationship between baseline and delayed performance can be seen. However, the strength of this relationship differs with respect to IAF and Condition. In particular, the relationship between baseline and delayed performance is weaker for high IAF individuals in the wake condition, suggesting that wake appears to interfere with rule extraction for high-IAF individuals.

A significant interaction between IAF, Coverb and Baseline d' was also

revealed (statistical information can be found in Table 6 and the relationship is

visualised in Appendix B [Figure 1A] of the supplementary material). Figure 4A

illustrates that IAF differentially influenced memory performance at delayed testing

depending both on condition and baseline performance. Further, sleep influenced

memory performance, thus encouraging further examination of the effects of specific

sleep physiology (i.e., PAC strength between SOs and spindles) on memory

outcomes, which we report below.

Table 6. Statistical values for main effects and significant interaction effects within the behavioural
model examined via type II Wald chi-square tests.

Parameters	χ^2	df	β	p
Condition*	2.13	1	-1.96	.144
Coverb*	.25	1	-2.56	.618
Violation Type*	.37	1	-1.02	.541
Baseline d'	61.12	1	1.14	<.001
IAF	6.96	1	-.40	.008
Condition x IAF x Baseline d'	4.51	1	-.36	<.001
IAF x Coverb x Baseline d'	3.89	1	-.30	.043

515   **Note:** χ^2 = chi-square ratio, *df* = degrees of freedom, *β* = beta value, *p* = statistical significance, * = non-
516   significant.

3.3   Sleep Analysis

The results reported in the following summarise the effects most pertinent to

the current hypotheses related to SWS neurophysiology, drawing on mixed model

output summaries (for *β* values) and Type II Wald Chi-square tests for *p* values. Full

model and Wald test summaries can be found in the supplementary material

(Appendix A.2).

Mixed-effects modelling showed a significant interaction between PAC, Coverb

and Violation Type, suggesting that coupling strength modulated performance at

delayed testing, modulated by Coverb and Violation Type (refer to Table 7 for

statistical values). As shown in Figure 4B, PAC did not influence performance for

noun order violations of either coverb. However, for verb position violations, as

coupling strength increased, differences in performance can be seen depending on

coverb. Specifically, as coupling strength increased, performance for Undergoer-first
constructions decreased and performance for Actor-first constructions increased.

Significant three-way interactions were also observed for (1) IAF x Coverb x
Violation Type, (2) IAF x Coverb x Baseline d', and (3) IAF x Violation Type x
Baseline d', indicating that IAF influences post-sleep memory performance, as
modulated by grammatical rule type and pre-sleep performance (visualisations of
these interactions can be found in Appendix B [Figure 1B, 1C and 1D, respectively]
of the supplementary material. However, there was no signficiant interaction
between PAC and IAF.

Table 7. Statistical values for main effects and significant interaction effects within neurophysiological
model examined via type II Wald chi-square tests.

Parameters	χ^2	df	β	p
PAC*	.11	1	.83	.735
IAF*	1.76	1	-.53	.185
Coverb*	3.33	1	-4.2	.068
Violation Type	5.83	1	-2.50	.016
Baseline d'	37.30	1	-1.70	<.001
PAC x Coverb	9.99	1	-.30	.001
IAF x Coverb	13.16	1	.46	<.001
IAF x Violation Type	6.71	1	.25	.009
Coverb x Violation Type	7.60	1	2.00	.006
IAF x Baseline d'	10.60	1	.27	.001
PAC x Coverb x Violation Type	5.91	1	1.30	.015
IAF x Coverb x Violation Type	10.83	1	-.23	<.001
IAF x Coverb x Baseline d'	9.42	1	-.57	.002
IAF x Violation Type x Baseline d'	4.018	1	-.28	.045

540 **Note:** χ^2 = chi-square ratio, *df* = degrees of freedom, *β* = beta value, *p* = statistical significance, * = non-
541 significant.

3.4   Exploratory Analysis

The results reported in the following summarise the effects most pertinent to
exploratory investigation into REM sleep neurophysiology, drawing on mixed model
output summaries (for β values) and Type II Wald Chi-square tests for *p* values. As
literature suggests an effect of REM sleep on the consolidation of memory (Batterink
et al., 2014; Djonlagic et al., 2009; Maquet et al., 2000), two exploratory models were

run, with an initial model to investigate the effect of theta power during REM and a

subsequent model exploring the effect of percentage of time spent in REM on

memory. As the purpose of these models were to explore REM theta power and

percentage of time spent in REM within a rule-based context, only interactions

involving these predictors will be reported here. Full model and Wald test summaries

for the REM theta power and percentage of REM models can be found in the

supplementary material (Appendix A.3 & A.4, respectively).

555   The first mixed-model revealed a significant four-way REM Theta Power x IAF

x Baseline d' x Saggitality interaction, as shown in Figure 4C (refer to Table 8 for

statistical values). Here, the largest differences between IAF values can be seen for

higher pre-sleep performance scores and most prominently at posterior regions.

There is also a notable relationship between post-sleep performance and theta

power during REM for high IAF individuals, such that as theta power increases,

memory performance also increases. The opposite pattern of results can be seen for

low-IAF individuals. This pattern indicates both IAF and theta power modulate

memory performance at delayed testing, and thus, rule extraction. Additionally, there

was a significant REM Theta Power x IAF x Coverb x Saggitality interaction

(statistical values can be found within Table 8 and this interaction is visualised within

Appendix B [Figure 2A] of the supplementary material).

Table 8. Statistical values for main effects and significant interaction effects within the exploratory model (REM theta power) examined via type II Wald chi-square tests.

Parameters	χ^2	df	β	p
REM Theta*	.00	1	.54	.997
IAF*	2.00	1	-.67	.157
Saggitality*	.01	1	.02, .-09	.995
Laterality*	.10	1	-.10, .33	.950
Coverb	9.84	1	-2.40	.002
Violation Type	21.93	1	-1.80	<.001
Baseline d'	185.85	1	-4.30	<.001
REM Theta x Coverb	8.30	1	3.25	.004
IAF x Coverb	31.75	1	.26	<.001
IAF x Type	33.87	1	.16	<.001
Coverb x Violation Type	53.21	1	.32	<.001
IAF x Baseline d'	110.88	1	.55	<.001
Coverb x Baseline d'	28.30	1	3.00	<.001
Violation Type x Baseline d'	4.53	1	1.80	.033
IAF x Coverb x Violation Type	18.72	1	-.05	<.001
REM Theta x IAF x Baseline d'	9.36	1	.22	.002
IAF x Coverb x Baseline d'	22.10	1	-.36	<.001
IAF x Type x Baseline d'	7.55	1	-.17	.006
REM Theta x IAF x Coverb x Saggitality	7.73	1	.16, -.33	.021
REM Theta x IAF x Baseline d' x Saggitality	6.66	1	04, -.33	.036

Note: χ^2 = chi-square ratio, *df* = degrees of freedom, *β* = beta value, *p* = statistical significance, * = non-significant.

The second mixed model explored percentage of time spent in REM sleep revealed a significant REM x Coverb x Type interaction (χ^2 (1) = 7.41, β = -.69, p = .006), with the resolved interaction being shown in Figure 4D (refer to Table 9 for statistical values). Interestingly, percentage of time spent in REM also shows no visible effect on noun order violations. However, for verb positioning violations, there is a positive relationship between percentage of REM and post-sleep memory performance for Undergoer-first (bèi) constructions. In contrast, a negative relationship is seen for Actor-first (bǎ) constructions, indicating that processes during

REM sleep are benefitting one rule type, whilst being detrimental for another.

Additionally, there was a significant REM x Coverb x Baseline d' interaction

(statistical values can be found within Table 8 and this interaction is visualised within

Appendix B [Figure 2B] of the supplementary material). However, there was no

significant interaction between percentage of time spent in REM and IAF.

Table 9. Statistical values for main effects and significant interaction effects within the exploratory
model (percentage of time in REM) examined via type II Wald chi-square tests.

Parameters	χ^2	df	β	p
REM*	.02	1	-4.20	.888
IAF*	1.80	1	-.27	.180
Coverb*	2.85	1	-7.2	.091
Violation Type	4.43	1	-1.20	.035
Baseline d'	38.20	1	2.00	<.001
IAF x Coverb	17.75	1	.73	<.001
Coverb x Violation Type	8.08	1	3.01	.004
IAF x Baseline d'	5.45	1	-0.06	.020
IAF x Coverb x Violation Type	12.57	1	-.32	<.001
REM x Coverb x Violation Type	7.41	1	-.69	.006
REM x Coverb x Baseline d'	4.33	1	3.72	.037
IAF x Coverb x Baseline d'	10.66	1	-.86	.001

586   **Note:** χ^2 = chi-square ratio, *df* = degrees of freedom, *β* = beta value, *p* = statistical significance, * = non-
587   significant.

588   **Figure 4.** Plotted interactions from the behavioural model (A), sleep model (B), and exploratory
589   models examining REM Theta Power (C) and percentage of time in REM sleep (D).

590
Note: All interactions are predicting performance scores at delayed testing (on the y-axis) quantified using d-
prime (fitted model values). For all plots, the shaded areas of colour represent 83% confidence intervals and the
red dashed line intercepting the y-axis at 0 represents chance performance (d' = 0). **(A)** Condition x IAF x
Baseline d' interaction. Baseline performance values (d') are depicted on the x-axis, beginning with lower
performance values. Additionally, plot is faceted by IAF on the top axis, split into 2 facets, with the left facet
showing the relationship between baseline performance and delayed d-prime for lower IAF individuals and the
right facet showing this relationship for higher IAF individuals. IAF was dichotomised on this plot for visualisation
purposes only. **(B)** PAC x Coverb x Type interaction. PAC values are shown on the x-axis, from lower coupling
values (left) to higher coupling values (right), with higher coupling values indicating better coupling strength
between slow-oscillations (SOs) and spindles. Violation Type is shown on the top axis, split into 2 facets, with
noun order violations on the left and verb positioning violations on the right. **(C)** REM Theta Power x IAF x
Saggitality x Baseline d' interaction. REM Theta Power is shown on the x-axis, from lower power values (left) to
higher power values (right), with higher power values reflecting higher power spectral density values. Saggitality
is shown on the top axis, split into 3 facets, moving from anterior (left) to central (middle) and posterior (right),
depicting topographical positioning. Performance scores at baseline testing are shown along the right axis, from
lower values (top) to higher values (bottom), with higher values indicating better performance at baseline.
Baseline performance was trichotomized here for visualisation purposes only. **(D)** Percentage of REM x Coverb x
Violation Type interaction. Percentage of time spent in REM sleep is shown along the x-axis, from lower values
(left) to higher values (right), with higher values depicting higher percentage of time spent in REM. Violation Type
is shown on the top axis, split into 2 facets, with noun order violations on the left and verb positioning violations
on the right.

4   DISCUSSION

Slow oscillation-spindle coupling during SWS and REM sleep have been independently linked to memory consolidation (Diekelmann & Born, 2010; Klinzing et al., 2019; Lewis & Durrant, 2011). Here, we provide evidence that these factors modulate memory performance within a rule-based context in language processing. Behaviourally, results indicate a positive relationship between baseline performance and delayed memory performance in both sleep and wake conditions that is modulated by individual alpha frequency (IAF). Regarding sleep neurophysiology, results indicate an interaction between phase-amplitude coupling (PAC) and sentence type (violation type and coverb). Specifically, higher PAC was associated with increased acceptability judgement performance for Actor-first (bǎ) sentences and decreased performance for Undergoer-first (bèi) with respect to the verb position rule. Our exploratory analyses of post-sleep memory performance further revealed a significant interaction between REM theta power, IAF, saggitality (anterior, midline and posterior regions) and pre-sleep performance. For high pre-sleep performers with a high IAF, as REM theta power increased, so did post-sleep memory performance, with this effect being most prominent in posterior regions. Additionally, proportion of REM sleep interacted with coverb and violation type, such that greater time spent in REM sleep was associated with decreased post-sleep performance on Actor-first constructions and increased performance for Undergoer-first constructions with respect to the verb position rule. Overall, results indicate that sleep neurophysiology and IAF influence consolidation and subsequent performance at both behavioural and neural levels and that patterns of memory performance differ across distinct rule types.

4.1   The influence of IAF and sleep on memory performance at the behavioural level

Our results partially supported the hypothesis (H^1) that behavioural performance should be enhanced post-sleep relative to post-wake, however this relationship is modulated by IAF. Specifically, there was a generally positive relationship between baseline and delayed performance across conditions. However, for higher-IAF individuals in the wake condition, this relationship was weaker. As

such, this result divergences from previous work showing that high IAF is associated with better performance on an array of tasks, including those assessing memory (Doppelmayr et al., 2005; Klimesch, 1997, 1999). More recent results (e.g., Ociepka et al., 2022) indicate that the association between high IAF and greater performance on a variety of tasks is not as common as once believed. In fact, in another reanalysis of Cross et al., (2021) examining a different linguistic manipulation, Nalaye et al., (2022) found a negative relationship between IAF and performance on the Mini Pinyin grammaticality judgement task, with lower-IAF individuals performing better. In combination with our results, it appears that high IAF is not beneficial in every context.

The link between IAF and information processing (Ociepka et al., 2022; Samaha & Postle, 2015; Surwillo, 1963) may have influence the process of sleep-dependent memory consolidation which relies on hippocampal-to-neocortical information transfer (Yassa & Reagh, 2013). ~~Specifically, Samaha and Postle (2015) suggest that a higher IAF indicates faster cycles of information processing and a finer temporal resolution of incoming sensory information.~~ High IAF individuals in the wake condition may have been more susceptible to interference (Radvansky & Copeland, 2006) due to faster cycles of processing associated with higher IAF. Thus, lower IAF (slower cycles) may have been beneficial for those in the wake condition (Howard et al., 2017), with low IAF individuals preserving the initially encoded traces with less interference than high IAF individuals in the wake group. Our proposed role of IAF in information processing may also help to explain why baseline performance is critical when observing effects of IAF. It may be that the temporal receptive windows associated with IAF influence both the process of consolidation as well as initial encoding (Khader et al., 2010; Klimesch, 1997; Klimesch et al., 1996). However, this analysis did not specifically look at the process of encoding which limits the understanding of what may be occurring during this time (for an in-depth analysis of the encoding phase, see Cross et al., 2022). Despite this, analyses reported here provide further evidence for an effect of IAF on memory consolidation, consistent with previous literature (Doppelmayr et al., 2005; Klimesch, 1997, 1999; Richard Clark et al., 2004).

4.2 Neural mechanisms underlying sleep-based consolidation of rules

Both the behavioural and sleep analyses provide support for $H^2(a)$ which
hypothesised that both IAF and PAC would predict post-sleep memory performance
and that this relationship would be additive. Both models indicated that IAF
influenced delayed memory performance. Further, in the sleep model, results
showed an interaction between PAC, coverb, and violation type, such that as PAC
increased, post-sleep memory performance for Actor-first sentences increased with
respect to the verb position rule, while it decreased for Undergoer-first constructions.
Slow oscillation-spindle coupling is suggested to reflect hippocampal-neocortical
information transfer during NREM sleep (Diekelmann & Born, 2010; Klinzing et al.,
2019; Lewis & Durrant, 2011), and as such, increased PAC associated with post-
sleep performance suggests NREM sleep-based consolidation processes have
occurred for rule-based information. This result is broadly consistent with previous
literature linking NREM sleep neurophysiology to memory consolidation (Diekelmann
& Born, 2010; Lewis & Durrant, 2011). This result also builds upon previous research
linking SWS duration with rule extraction for both linguistic rules and non-linguistic
rules (e.g., statistical; Batterink et al., 2014; Djonlagic et al., 2009; Durrant et al.,
2013; Lerner & Gluck, 2019; Nieuwenhuis et al., 2013), linking specific PAC
mechanisms (e.g., SO-spindle coupling) to memory performance in a rule-based
context. In contrast, $H^2(b)$ was not supported, as PAC and IAF did not interact,
contrasting previous literature in which IAF was associated with neurophysiology
during SWS (Cross et al., 2020a).

Interestingly, PAC showed the greatest influence relative to the verb position
rule, suggesting the novelty of a rule may play a role. When exploring the verb
position rule, SO-spindle PAC demonstrated a differential effect depending on the
coverb. Specifically, increased PAC moderately improved memory performance for
Actor-first (bǎ) sentences. This may be due to native English speakers being more
familiar with sentences in which the Actor occurs before the Undergoer
(MacWhinney et al., 1984), which is also the case for bǎ sentences. As this rule type
is less novel than the others examined here, better regularity extraction may have
been influenced by such familiarity, leading to an effect of PAC for Actor-first
sentences. By contrast, PAC negatively affected performance for verb position in
Undergoer-first (bèi) sentences. This may have been due to the non-schema

conforming nature of these constructions, which differ from English both with respect

to noun order and verb position, resulting in a higher level of novelty. According to

the iOTA model of sleep-based memory consolidation, non-schema conforming

information is often downscaled to preferentially encode information analogous to

existing knowledge (Lewis et al., 2018; Lewis & Durrant, 2011). Additionally, theories

of memory consolidation, such as Complementary Learning Systems theory

(McClelland et al., 1995), suggest that information similar to that of previous

knowledge tends to be encoded and consolidated easier (Tse et al., 2011) and

quicker (Tse et al., 2007), due to associative schemas. As such, the associative

schema here is likely the grammatical elements of English, allowing the similar

information (Actor-first sentences) to be more easily consolidated. Thus, this may

explain how NREM mechanisms in our findings had the greatest effect on

information closest to existing knowledge (i.e., Actor-first constructions) and had the

opposite effect for information dissimilar to existing knowledge (i.e., Undergoer-first

constructions), consistent with the iOTA model (Lewis et al., 2018; Lewis & Durrant,

2011).

4.3   The role of REM sleep in memory consolidation and rule extraction

As an exploratory step, we investigated the effect of REM sleep

neurophysiology on rule extraction. Our results indicate differences in post-sleep

memory performance relative to REM theta activity in posterior regions, contrasting

with existing literature linking REM theta with central regions (Fogel et al., 2007) and

anterior regions (Simor et al., 2016). Posterior theta has been proposed as the

human correlate of the ponto-geniculo-occipital (PGO) wave (Frauscher et al., 2018),

which has been previously attributed to memory consolidation and neuronal plasticity

in experimental animal research (Datta, 2000; Datta et al., 2004). If so, the posterior

theta activity reported here may reflect changes in neuronal plasticity (Grosmark et

al., 2012; Hobson & Friston, 2012) and the consolidation of rule-based information.

Additionally, the relationship between REM theta and post-sleep memory outcomes

appears to differ depending on IAF. Our results revealed a positive relationship

between REM theta and post-sleep memory for higher-IAF and a negative

relationship for lower-IAF individuals. Cross et al., (2020a) also found a complex

relationship between IAF and theta power during REM sleep. Higher IAF may have

allowed for increased reactivations of memory during sleep or allowed the brain to
preferentially encode salient information (Cross et al., 2020a; Heib et al., 2015).

We also observed an interaction between REM sleep percentage, coverb and
violation type. Specifically, higher proportions of REM sleep positively affected
performance for Undergoer-first constructions (bèi) and led to decreased
performance for Actor-first constructions (bǎ), contrasting with the pattern seen
within NREM sleep mechanisms. This result partially converges with the results of
Matorina & Poppenk, (2019), in which higher proportions of REM sleep led to a
decrease in performance on a statistical learning task. Matorina & Poppenk, (2019)
proposed that REM sleep may in fact be working against processes of extracting
regularities, as it is more useful for a process called schema reconstruction, which
disbands and reintegrates current and existing information into abstract schemas
(Landmann et al., 2014). Further evidence for this comes from the iOTA model, in
which REM sleep is proposed as the ideal setting for forming novel and abstract
representations. Thus, NREM is likely most beneficial for associative memory
processes, while REM sleep may facilitate the construction of abstract schemas
(Lewis et al., 2018). The findings presented here provide support for this hypothesis,
as demonstrated by the favourable effect of NREM sleep on Actor-first constructions
(lower novelty) and REM sleep for Undergoer-first constructions (higher novelty).
Overall, despite tentative interpretation, our results provide preliminary evidence for
REM theta and IAF in rule-based contexts and demonstrate support for REM in non-
schema conforming memory consolidation, providing a foundation for future
research.

4.4 Limitations

As previously discussed, there are several limitations that should be addressed
in future work. Firstly, analysis of electrophysiological activity during the encoding
phase as they relate to delayed memory outcomes would provide additional
understanding of the mechanisms underlying rule-based learning and consolidation.
Additionally, whilst the focus here was on SO-spindle coupling, future research
would benefit from including a broader range of sleep physiology metrics such as
spindle-ripple coupling; albeit, this would need to be performed with invasive
monitoring techniques, such as intracranial EEG, as scalp-recorded EEG is not

capable of capturing activity in the ripple range (i.e., ~ 80 – 140 Hz). This would aid
in further understanding of the specific mechanisms driving rule-based consolidation
during sleep. Additionally, the differing times of the day at which participants were
tested may warrant further exploration into differences in circadian rhythms and any
potential influence on wake-EEG (Aeschbach et al., 1997, 1999) and IAF. The
relationship between IAF and circadian rhythms in particular is largely unexplored
and warrants further research. Despite these limitations, the current study provides a
novel exploration of the neurophysiological mechanisms underlying the consolidation
of rule-based information.

4.5   Conclusions

Here, we demonstrated that individual differences in intrinsic neural activity,
indexed by IAF, predict rule extraction and consolidation across sleep and wake.
Contrary to previous understanding of individual electrophysiological differences, we
found a beneficial influence of lower IAF on memory within a rule-based context,
proposing that higher-IAF and lower-IAF individuals draw on different information
processing strategies as opposed to simply showing differing performance
outcomes. We also demonstrated differential effects of SO-spindle coupling for
specific grammatical rules, favouring the consolidation of those analogous to existing
grammatical knowledge. We further provide support for the role of REM sleep in
consolidating non-schema conforming information, potentially through schema
reconstruction and the proposed role for REM for forming novel and abstract
representations. Finally, we provided preliminary evidence for a relationship between
REM sleep and IAF within rule extraction, a previously limited area of research.
Future research should focus on further delineating the relationship between REM
theta activity and individual processing capabilities in rule-based consolidation, which
would further our understanding of the complex relationship between individual
differences, sleep neurophysiology and memory consolidation.

5   REFERENCES

Ackermann, S., & Rasch, B. (2014). Differential effects of non-REM and REM sleep on memory consolidation? *Current Neurology and Neuroscience Reports*, *14*(2), Article 2.

Aeschbach, D., Matthews, J. R., Postolache, T. T., Jackson, M. A., Giesen, H. A., & Wehr, T. A. (1997). Dynamics of the human EEG during prolonged wakefulness: Evidence for frequency-specific circadian and homeostatic influences. *Neuroscience Letters*, *239*(2), 121–124. https://doi.org/10.1016/S0304-3940(97)00904-X

Aeschbach, D., Matthews, J. R., Postolache, T. T., Jackson, M. A., Giesen, H. A., & Wehr, T. A. (1999). Two circadian rhythms in the human electroencephalogram during wakefulness. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *277*(6), R1771–R1779. https://doi.org/10.1152/ajpregu.1999.277.6.R1771

Alday, P. M. (2019). *Philistine documentation*. https://philistine.readthedocs.io/_/downloads/en/latest/pdf/

Babu Henry Samuel, I., Wang, C., Hu, Z., & Ding, M. (2018). The frequency of alpha oscillations: Task-dependent modulation and its functional significance. *NeuroImage*, *183*, 897–906. https://doi.org/10.1016/j.neuroimage.2018.08.063

Bakker, I., Takashima, A., van Hell, J. G., Janzen, G., & McQueen, J. M. (2015). Changes in Theta and Beta Oscillations as Signatures of Novel Word Consolidation. *Journal of Cognitive Neuroscience*, *27*(7), Article 7. https://doi.org/10.1162/jocn_a_00801

Barachant, A. (2015). *PyRiemann: Biosignals classification with Riemannian Geometry—PyRiemann 0.2.6 documentation*. https://pyriemann.readthedocs.io/en/latest/index.html

Barthélemy, Q., Mayaud, L., Ojeda, D., & Congedo, M. (2019). The Riemannian Potato Field: A Tool for Online Signal Quality Index of EEG. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, *27*(2), Article 2. https://doi.org/10.1109/TNSRE.2019.2893113

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), Article 1. https://doi.org/doi:10.18637/jss.v067.i01

Batterink, L. J., Oudiette, D., Reber, P. J., & Paller, K. A. (2014). Sleep facilitates learning a new linguistic rule. *Neuropsychologia*, *65*, 169–179.

Berry, R. B., Budhiraja, R., Gottlieb, D. J., Gozal, D., Iber, C., Kapur, V. K., Marcus, C. L., Mehra, R., Parthasarathy, S., & Quan, S. F. (2012). Rules for scoring respiratory events in sleep: Update

of the 2007 AASM manual for the scoring of sleep and associated events. *Journal of Clinical Sleep Medicine*, *8*(05), Article 05.

Bornkessel, I. D., Fiebach, C. J., Friederici, A. D., & Schlesewsky, M. (2004). "Capacity" reconsidered: Interindividual differences in language comprehension and individual alpha frequency. *Experimental Psychology*, *51*(4), Article 4.

Bornkessel-Schlesewsky, I., Sharrad, I., Howlett, C. A., Alday, P. M., Corcoran, A. W., Bellan, V., Wilkinson, E., Kliegl, R., Lewis, R. L., Small, S. L., & Schlesewsky, M. (2022). Rapid adaptation of predictive models during language comprehension: Aperiodic EEG slope, individual alpha frequency and idea density modulate individual differences in real-time model updating. *Frontiers in Psychology*, *13*. https://www.frontiersin.org/articles/10.3389/fpsyg.2022.817516

Boyce, R., Glasgow, S. D., Williams, S., & Adamantidis, A. (2016). Causal evidence for the role of REM sleep theta rhythm in contextual memory consolidation. *Science (New York, N.Y.)*, *352*(6287), Article 6287. https://doi.org/10.1126/science.aad5252

Brown, V. A. (2020). *An approachable introduction to linear mixed effects modeling with implementation in R*. https://doi.org/10.31234/osf.io/9vghm

Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., Berger, M. S., Barbaro, N. M., & Knight, R. T. (2006). High Gamma Power Is Phase-Locked to Theta Oscillations in Human Neocortex. *Science*, *313*(5793), Article 5793. https://doi.org/10.1126/science.1128115

Canolty, R. T., & Knight, R. T. (2010). The functional role of cross-frequency coupling. *Trends in Cognitive Sciences*, *14*(11), 506–515.

Capotosto, P., Baldassarre, A., Sestieri, C., Spadone, S., Romani, G. L., & Corbetta, M. (2017). Task and Regions Specific Top-Down Modulation of Alpha Rhythms in Parietal Cortex. *Cerebral Cortex*, *27*(10), Article 10. https://doi.org/10.1093/cercor/bhw278

Chatburn, A., Lushington, K., & Kohler, M. J. (2021). Consolidation and generalisation across sleep depend on individual EEG factors and sleep spindle density. *Neurobiology of Learning and Memory*, *179*, 107384. https://doi.org/10.1016/j.nlm.2021.107384

Collette, F., Grandjean, J., Lorant, C., & Bastin, C. (2014). The Role of Memory Traces Quality in

Directed Forgetting: A Comparison of Young and Older Participants. *Psychologica Belgica*,

54(4), Article 4. https://doi.org/10.5334/pb.au

Corcoran, A. W., Alday, P. M., Schlesewsky, M., & Bornkessel-Schlesewsky, I. (2018). Toward a

reliable, automated method of individual alpha frequency (IAF) quantification.

Psychophysiology, *55*(7), Article 7. https://doi.org/10.1111/psyp.13064

Cross, Z. R., Corcoran, A. W., Schlesewsky, M., Kohler, M. J., & Bornkessel-Schlesewsky, I. (2022).

Oscillatory and Aperiodic Neural Activity Jointly Predict Language Learning. *Journal of*

Cognitive Neuroscience, 1–20. https://doi.org/10.1162/jocn_a_01878

Cross, Z. R., Helfrich, R. F., Corcoran, A. W., Kohler, M. J., Coussens, S., Zou-Williams, L.,

Schlesewsky, M., Gaskell, M. G., Knight, R. T., & Bornkessel-Schlesewsky, I. (2021). *Spindle-*

slow oscillation coupling during sleep predicts sequence-based language learning (p.

2020.02.13.948539). bioRxiv. https://doi.org/10.1101/2020.02.13.948539

Cross, Z. R., Santamaria, A., Corcoran, A. W., Chatburn, A., Alday, P. M., Coussens, S., & Kohler, M.

J. (2020). Individual alpha frequency modulates sleep-related emotional memory

consolidation. *Neuropsychologia*, *148*, 107660.

Cross, Z. R., Zou-Williams, L., Wilkinson, E., Schlesewsky, M., & Bornkessel-Schlesewsky, I. (2020).

Mini Pinyin: A modified miniature language for studying language learning and incremental

sentence processing [Preprint]. PsyArXiv. https://doi.org/10.31234/osf.io/b9erz

Datta, S. (2000). Avoidance Task Training Potentiates Phasic Pontine-Wave Density in the Rat: A

Mechanism for Sleep-Dependent Plasticity. *Journal of Neuroscience*, *20*(22), Article 22.

https://doi.org/10.1523/JNEUROSCI.20-22-08607.2000

Datta, S., Mavanji, V., Ulloor, J., & Patterson, E. H. (2004). Activation of Phasic Pontine-Wave

Generator Prevents Rapid Eye Movement Sleep Deprivation-Induced Learning Impairment in

the Rat: A Mechanism for Sleep-Dependent Plasticity. *Journal of Neuroscience*, *24*(6), Article

6. https://doi.org/10.1523/JNEUROSCI.4111-03.2004

Diekelmann, S., & Born, J. (2010). The memory function of sleep. *Nature Reviews Neuroscience*,

11(2), Article 2.

Diekelmann, S., Born, J., & Wagner, U. (2010). Sleep enhances false memories depending on

general memory performance. *Behavioural Brain Research*, *208*(2), Article 2.

https://doi.org/10.1016/j.bbr.2009.12.021

Djonlagic, I., Rosenfeld, A., Shohamy, D., Myers, C., Gluck, M., & Stickgold, R. (2009). Sleep

enhances category learning. *Learning & Memory*, *16*(12), Article 12.

Doppelmayr, M., Klimesch, W., Hödlmoser, K., Sauseng, P., & Gruber, W. (2005). Intelligence related

upper alpha desynchronization in a semantic memory task. *Brain Research Bulletin*, *66*(2),

Article 2.

Durrant, S. J., Cairney, S. A., & Lewis, P. A. (2013). Overnight consolidation aids the transfer of

statistical knowledge from the medial temporal lobe to the striatum. *Cerebral Cortex*, *23*(10),

Article 10.

Durrant, S. J., Taylor, C., Cairney, S., & Lewis, P. A. (2011). Sleep-dependent consolidation of

statistical learning. *Neuropsychologia*, *49*(5), Article 5.

Dvorak, D., & Fenton, A. A. (2014). Toward a proper estimation of phase–amplitude coupling in neural

oscillations. *Journal of Neuroscience Methods*, *225*, 42–56.

Fenn, K. M., & Hambrick, D. Z. (2012). Individual differences in working memory capacity predict

sleep-dependent memory consolidation. *Journal of Experimental Psychology. General*,

141(3), 404–410. https://doi.org/10.1037/a0025268

Fogel, S. M., Smith, C. T., & Cote, K. A. (2007). Dissociable learning-dependent changes in REM and

non-REM sleep in declarative and procedural memory systems. *Behavioural Brain Research*,

180(1), Article 1. https://doi.org/10.1016/j.bbr.2007.02.037

Fox, J., & Weisberg, S. (2018). *An R companion to applied regression*. Sage publications.

Frank, M. C., & Gibson, E. (2011). Overcoming Memory Limitations in Rule Learning. *Language

Learning and Development*, *7*(2), Article 2. https://doi.org/10.1080/15475441.2010.512522

Frauscher, B., Joshi, S., von Ellenrieder, N., Nguyen, D. K., Dubeau, F., & Gotman, J. (2018). Sharply

contoured theta waves are the human correlate of ponto-geniculo-occipital waves in the

primary visual cortex. *Clinical Neurophysiology*, *129*(8), Article 8.

https://doi.org/10.1016/j.clinph.2018.04.605

Furman, A. J., Meeker, T. J., Rietschel, J. C., Yoo, S., Muthulingam, J., Prokhorenko, M., Keaser, M.

L., Goodman, R. N., Mazaheri, A., & Seminowicz, D. A. (2018). Cerebral peak alpha

frequency predicts individual differences in pain sensitivity. *NeuroImage*, *167*, 203–210.

https://doi.org/10.1016/j.neuroimage.2017.11.042

García-Lázaro, H. G., Ramirez-Carmona, R., Lara-Romero, R., & Roldan-Valadez, E. (2012).

Neuroanatomy of episodic and semantic memory in humans: A brief review of neuroimaging

studies. *Neurology India*, *60*(6), Article 6. https://doi.org/10.4103/0028-3886.105196

Girardeau, G., & Zugaro, M. (2011). Hippocampal ripples and memory consolidation. *Current Opinion*

in Neurobiology, *21*(3), Article 3. https://doi.org/10.1016/j.conb.2011.02.005

Gómez, R. L., Bootzin, R. R., & Nadel, L. (2006). Naps promote abstraction in language-learning

infants. *Psychological Science*, *17*(8), Article 8. https://doi.org/10.1111/j.1467-

9280.2006.01764.x

Grosmark, A. D., Mizuseki, K., Pastalkova, E., Diba, K., & Buzsáki, G. (2012). REM Sleep

Reorganizes Hippocampal Excitability. *Neuron*, *75*(6), Article 6.

https://doi.org/10.1016/j.neuron.2012.08.015

Hahn, M. A., Heib, D., Schabus, M., Hoedlmoser, K., & Helfrich, R. F. (2020). Slow oscillation-spindle

coupling predicts enhanced memory formation from childhood to adolescence. *Elife*, *9*,

e53730.

Harrell Jr, F. E. (2019). Package 'hmisc.' *CRAN2018*, *2019*, 235–236.

Heib, D. P. J., Hoedlmoser, K., Anderer, P., Gruber, G., Zeitlhofer, J., & Schabus, M. (2015).

Oscillatory Theta Activity during Memory Formation and Its Impact on Overnight

Consolidation: A Missing Link? *Journal of Cognitive Neuroscience*, *27*(8), Article 8.

https://doi.org/10.1162/jocn_a_00804

Helfrich, R. F., Lendner, J. D., Mander, B. A., Guillen, H., Paff, M., Mnatsakanyan, L., Vadera, S.,

Walker, M. P., Lin, J. J., & Knight, R. T. (2019). Bidirectional prefrontal-hippocampal dynamics

organize information transfer during sleep in humans. *Nature Communications*, *10*(1), Article

1. https://doi.org/10.1038/s41467-019-11444-x

Helfrich, R. F., Mander, B. A., Jagust, W. J., Knight, R. T., & Walker, M. P. (2018). Old brains come

uncoupled in sleep: Slow wave-spindle synchrony, brain atrophy, and forgetting. *Neuron*,

97(1), Article 1.

Hobson, J. A., & Friston, K. J. (2012). Waking and dreaming consciousness: Neurobiological and

functional considerations. *Progress in Neurobiology*, *98*(1), Article 1.

https://doi.org/10.1016/j.pneurobio.2012.05.003

Hope, R. M. (2013). *Rmisc: Ryan miscellaneous. R package version 1.5*. CRAN.

Howard, C. J., Arnold, C. P. A., & Belmonte, M. K. (2017). Slower resting alpha frequency is

associated with superior localisation of moving targets. *Brain and Cognition*, *117*, 97–107.

https://doi.org/10.1016/j.bandc.2017.06.008

Hülsemann, M. J., Naumann, E., & Rasch, B. (2019). Quantification of Phase-Amplitude Coupling in

Neuronal Oscillations: Comparison of Phase-Locking Value, Mean Vector Length, Modulation

Index, and Generalized-Linear-Modeling-Cross-Frequency-Coupling. *Frontiers in*

Neuroscience, *13*. https://doi.org/10.3389/fnins.2019.00573

Humiston, G. B., Tucker, M. A., Summer, T., & Wamsley, E. J. (2019). Resting States and Memory

Consolidation: A Preregistered Replication and Meta-Analysis. *Scientific Reports*, *9*(1),

19345. https://doi.org/10.1038/s41598-019-56033-6

Kassambara, A. (2020). Package 'ggpubr.' *R Package Version 0.1*, *6*.

Khader, P. H., Jost, K., Ranganath, C., & Rösler, F. (2010). Theta and alpha oscillations during

working-memory maintenance predict successful long-term memory encoding. *Neuroscience*

Letters, *468*(3), Article 3. https://doi.org/10.1016/j.neulet.2009.11.028

Klimesch, W. (1997). EEG-alpha rhythms and memory processes. *International Journal of*

Psychophysiology, *26*(1–3), Article 1–3.

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: A

review and analysis. *Brain Research Reviews*, *29*(2–3), Article 2–3.

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored information.

Trends in Cognitive Sciences, *16*(12), Article 12.

Klimesch, W., Doppelmayr, M., Schimke, H., & Pachinger, T. (1996). Alpha frequency, reaction time,

and the speed of processing information. *Journal of Clinical Neurophysiology: Official*

Publication of the American Electroencephalographic Society, *13*(6), Article 6.

https://doi.org/10.1097/00004691-199611000-00006

Klinzing, J. G., Niethard, N., & Born, J. (2019). Mechanisms of systems memory consolidation during

sleep. *Nature Neuroscience*, *22*(10), Article 10. https://doi.org/10.1038/s41593-019-0467-3

Landmann, N., Kuhn, M., Piosczyk, H., Feige, B., Baglioni, C., Spiegelhalder, K., Frase, L., Riemann,

973   D., Sterr, A., & Nissen, C. (2014). The reorganisation of memory during sleep. *Sleep Medicine*

974   *Reviews*, *18*(6), Article 6. https://doi.org/10.1016/j.smrv.2014.03.005

975   Latchoumane, C.-F. V., Ngo, H.-V. V., Born, J., & Shin, H.-S. (2017). Thalamic Spindles Promote

976   Memory Formation during Sleep through Triple Phase-Locking of Cortical, Thalamic, and

977   Hippocampal Rhythms. *Neuron*, *95*(2), Article 2. https://doi.org/10.1016/j.neuron.2017.06.025

978   Lerner, I., & Gluck, M. A. (2019). Sleep and the extraction of hidden regularities: A systematic review

979   and the importance of temporal rules. *Sleep Medicine Reviews*, *47*, 39–50.

980   https://doi.org/10.1016/j.smrv.2019.05.004

981   Lewis, P. A., & Durrant, S. J. (2011). Overlapping memory replay during sleep builds cognitive

982   schemata. *Trends in Cognitive Sciences*, *15*(8), Article 8.

983   https://doi.org/10.1016/j.tics.2011.06.004

984   Lewis, P. A., Knoblich, G., & Poe, G. (2018). How Memory Replay in Sleep Boosts Creative Problem-

985   Solving. *Trends in Cognitive Sciences*, *22*(6), 491–503.

986   https://doi.org/10.1016/j.tics.2018.03.009

987   Lockhart, R. S., & Murdock, B. B. (1970). Memory and the theory of signal detection. *Psychological*

988   *Bulletin*, *74*(2), Article 2.

989   Lüdecke, D. (2018a). sjmisc: Data and variable transformation functions. *Journal of Open Source*

990   *Software*, *3*(26), 754.

991   Lüdecke, D. (2018b). sjPlot: Data visualization for statistics in social science. *R Package Version*,

992   *2*(1).

993   Lüdecke, D., Aust, F., Crawley, S., & Ben-Shachar, M. (2020). Package 'ggeffects.' *Create Tidy Data*

994   *Frames of Marginal Effects for "Ggplot" from Model Outputs*, *23*.

995   Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). performance: An R

996   package for assessment, comparison and testing of statistical models. *Journal of Open*

997   *Source Software*, *6*(60).

998   Lund, U., & Agostinelli, C. (2012). Circstats: Circular Statistics, from "Topics in Circular

999   Statistics"(2001). *R Package Version*, *2*(1).

Lutz, N. D., Diekelmann, S., Hinse-Stern, P., Born, J., & Rauss, K. (2017). Sleep Supports the Slow Abstraction of Gist from Visual Perceptual Memories. *Scientific Reports*, *7*(1), Article 1. https://doi.org/10.1038/srep42950

MacWhinney, B., Bates, E., & Kliegl, R. (1984). Cue validity and sentence interpretation in English, German, and Italian. *Journal of Verbal Learning and Verbal Behavior*, *23*(2), Article 2.

Maquet, P., Laureys, S., Peigneux, P., Fuchs, S., Petiau, C., Phillips, C., Aerts, J., Del Fiore, G., Degueldre, C., Meulemans, T., Luxen, A., Franck, G., Van Der Linden, M., Smith, C., & Cleeremans, A. (2000). Experience-dependent changes in cerebral activation during human REM sleep. *Nature Neuroscience*, *3*(8), Article 8. https://doi.org/10.1038/77744

Marshall, L., & Born, J. (2007). The contribution of sleep to hippocampus-dependent memory consolidation. *Trends in Cognitive Sciences*, *11*(10), Article 10. https://doi.org/10.1016/j.tics.2007.09.001

Mathôt, S., Schreij, D., & Theeuwes, J. (2012). OpenSesame: An open-source, graphical experiment builder for the social sciences. *Behavior Research Methods*, *44*(2), Article 2.

Matorina, N., & Poppenk, J. (2019). Sleep promotes relational overlapping memories for long-term generalization. *BioRxiv*.

Matorina, N., & Poppenk, J. (2021). Memory decay distinguishes subtypes of gist. *Neurobiology of Learning and Memory*, *185*, 107519. https://doi.org/10.1016/j.nlm.2021.107519

McClelland, J. L., McNaughton, B. L., & O'Reilly, R. C. (1995). Why there are complementary learning systems in the hippocampus and neocortex: Insights from the successes and failures of connectionist models of learning and memory. *Psychological Review*, *102*(3), Article 3. https://doi.org/10.1037/0033-295X.102.3.419

Muehlroth, B. E., Sander, M. C., Fandakova, Y., Grandy, T. H., Rasch, B., Shing, Y. L., & Werkle-Bergner, M. (2019). Precise Slow Oscillation–Spindle Coupling Promotes Memory Consolidation in Younger and Older Adults. *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-018-36557-z

Mueller, J. L. (2006). L2 in a nutshell: The investigation of second language processing in the miniature language model. *Language Learning*, *56*, 235–270.

Munia, T. T. K., & Aviyente, S. (2019). Time-Frequency Based Phase-Amplitude Coupling Measure

For Neuronal Oscillations. *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-

019-48870-2

Nalaye, H., Cross, Z. R., Schlesewsky, M., & Bornkessel-Schlesewsky, I. (2022). *Electrophysiological*

indices of individual differences in adult language learning (p. 2022.06.07.495229). bioRxiv.

https://doi.org/10.1101/2022.06.07.495229

Neuwirth, E. (2014). Package 'RColorBrewer.' *ColorBrewer Palettes*.

Nieuwenhuis, I. L., Folia, V., Forkstam, C., Jensen, O., & Petersson, K. M. (2013). Sleep promotes

the extraction of grammatical rules. *PloS One*, *8*(6), Article 6.

Ociepka, M., Kałamała, P., & Chuderski, A. (2022). High individual alpha frequency brains run fast,

but it does not make them smart. *Intelligence*, *92*, 101644.

https://doi.org/10.1016/j.intell.2022.101644

Pothos, E. M. (2007). Theories of artificial grammar learning. *Psychological Bulletin*, *133*(2), Article 2.

Prat, C. S., Yamasaki, B. L., Kluender, R. A., & Stocco, A. (2016). Resting-state qEEG predicts rate of

second language learning in adults. *Brain and Language*, *157–158*, 44–50.

https://doi.org/10.1016/j.bandl.2016.04.007

Radvansky, G. A., & Copeland, D. E. (2006). Memory retrieval and interference: Working memory

issues. *Journal of Memory and Language*, *55*(1), Article 1.

https://doi.org/10.1016/j.jml.2006.02.001

Richard Clark, C., Veltmeyer, M. D., Hamilton, R. J., Simms, E., Paul, R., Hermens, D., & Gordon, E.

(2004). Spontaneous alpha peak frequency predicts working memory performance across the

age span. *International Journal of Psychophysiology*, *53*(1), Article 1.

https://doi.org/10.1016/j.ijpsycho.2003.12.011

Samaha, J., & Postle, B. R. (2015). The Speed of Alpha-Band Oscillations Predicts the Temporal

Resolution of Visual Perception. *Current Biology*, *25*(22), Article 22.

https://doi.org/10.1016/j.cub.2015.10.007

Sarkar, D. (2008). *Lattice: Multivariate data visualization with R*. Springer Science & Business Media.

Schabus, M., Kerstin, H., Thomas, P., Anderer, P., Gruber, G., Parapatics, S., Sauter, C., Kloesch,

G., Klimesch, W., Saletu, B., & Zeitlhofer, J. (2008). Interindividual sleep spindle differences

and their relation to learning-related enhancements. *Brain Research*, *1191*, 127–135.

https://doi.org/10.1016/j.brainres.2007.10.106

Schad, D. J., Vasishth, S., Hohenstein, S., & Kliegl, R. (2020). How to capitalize on a priori contrasts

in linear (mixed) models: A tutorial. *Journal of Memory and Language*, *110*, 104038.

https://doi.org/10.1016/j.jml.2019.104038

Simor, P., Gombos, F., Szakadát, S., Sándor, P., & Bódizs, R. (2016). EEG spectral power in phasic

and tonic REM sleep: Different patterns in young adults and children. *Journal of Sleep

Research*, *25*(3), Article 3. https://doi.org/10.1111/jsr.12376

Stanley, D. J., & Spence, J. R. (2018). Reproducible Tables in Psychology Using the apaTables

Package. *Advances in Methods and Practices in Psychological Science*, *1*(3), 415–431.

https://doi.org/10.1177/2515245918773743

Staresina, B. P., Bergmann, T. O., Bonnefond, M., van der Meij, R., Jensen, O., Deuker, L., Elger, C.

E., Axmacher, N., & Fell, J. (2015). Hierarchical nesting of slow oscillations, spindles and

ripples in the human hippocampus during sleep. *Nature Neuroscience*, *18*(11), Article 11.

https://doi.org/10.1038/nn.4119

Stark, S., Gordon, B., & Stark, C. (2008). A case study of amnesia: Exploring a paradigm for new

semantic learning and generalization. *Brain Injury*, *22*(3), Article 3.

https://doi.org/10.1080/02699050801953081

Stark, S. M., & Stark, C. E. (2016). Introduction to Memory. In *Neurobiology of language* (pp. 841–

854). Elsevier.

Stickgold, R., & Walker, M. P. (2013). Sleep-dependent memory triage: Evolving generalization

through selective processing. *Nature Neuroscience*, *16*(2), Article 2.

Surwillo, W. W. (1961). Frequency of the 'alpha'rhythm, reaction time and age. *Nature*, *191*(4790),

Article 4790.

Surwillo, W. W. (1963). The relation of simple response time to brain-wave frequency and the effects

of age. *Electroencephalography and Clinical Neurophysiology*, *15*(1), Article 1.

Tononi, G., & Cirelli, C. (2006). Sleep function and synaptic homeostasis. *Sleep Medicine Reviews*,

10(1), Article 1.

Tse, D., Langston, R. F., Kakeyama, M., Bethus, I., Spooner, P. A., Wood, E. R., Witter, M. P., &

Morris, R. G. M. (2007). Schemas and Memory Consolidation. *Science*, *316*(5821), Article

5821. https://doi.org/10.1126/science.1135935

Tse, D., Takeuchi, T., Kakeyama, M., Kajii, Y., Okuno, H., Tohyama, C., Bito, H., & Morris, R. G. M.

(2011). Schema-Dependent Gene Activation and Memory Encoding in Neocortex. *Science*,

333(6044), Article 6044. https://doi.org/10.1126/science.1205274

Tucker, M. A., Humiston, G. B., Summer, T., & Wamsley, E. (2020). Comparing the Effects of Sleep

and Rest on Memory Consolidation. *Nature and Science of Sleep*, *12*, 79–91.

https://doi.org/10.2147/NSS.S223917

Vallat, R., & Walker, M. P. (2021). An open-source, high-performance tool for automated sleep

staging. *ELife*, *10*, e70092. https://doi.org/10.7554/eLife.70092

Wagner, U., & Born, J. (2008). Memory consolidation during sleep: Interactive effects of sleep stages

and HPA regulation. *Stress*, *11*(1), Article 1. https://doi.org/10.1080/10253890701408822

Wamsley, E. J. (2019). Memory Consolidation during Waking Rest. *Trends in Cognitive Sciences*,

23(3), Article 3. https://doi.org/10.1016/j.tics.2018.12.007

Wang, L., Schlesewsky, M., Philipp, M., & Bornkessel-Schlesewsky, I. (2012). The role of animacy in

online argument interpretation in Mandarin Chinese. In *Case, word order and prominence* (pp.

91–119). Springer.

Weiss, S., & Rappelsberger, P. (2000). Long-range EEG synchronization during word encoding

correlates with successful memory performance. *Cognitive Brain Research*, *9*(3), Article 3.

https://doi.org/10.1016/S0926-6410(00)00011-2

Wickham, H. (2007). Reshaping data with the reshape package. *Journal of Statistical Software*, *21*,

1–20.

Wickham, H. (2011). The Split-Apply-Combine Strategy for Data Analysis. *Journal of Statistical

Software*, *40*, 1–29. https://doi.org/10.18637/jss.v040.i01

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K.,

Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the

Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wickham, H., Chang, W., Henry, L., Pedersen, T. L., Takahashi, K., Wilke, C., Woo, K., Yutani, H.,

Dunnington, D., & RStudio. (2021). *ggplot2: Create Elegant Data Visualisations Using the*

Grammar of Graphics (3.3.5). https://CRAN.R-project.org/package=ggplot2

Wilhelm, I., Rose, M., Imhof, K. I., Rasch, B., Büchel, C., & Born, J. (2013). The sleeping child

outplays the adult's capacity to convert implicit into explicit knowledge. *Nature Neuroscience*,

16(4), Article 4. https://doi.org/10.1038/nn.3343

Wixted, J. T. (2014). Signal detection theory. *Wiley StatsRef: Statistics Reference Online*.

Yassa, M., & Reagh, Z. (2013). Competitive Trace Theory: A Role for the Hippocampus in Contextual

Interference during Retrieval. *Frontiers in Behavioral Neuroscience*, *7*, 107.

https://doi.org/10.3389/fnbeh.2013.00107

1124