Chimeric PRMT6 protein produced by an endogenous retrovirus promoter

regulates cell fate decision in mouse preimplantation embryos

Shinnosuke Honda, Maho Hatamura, Yuri Kunimoto, Shuntaro Ikeda, and Naojiro

Minami

Laboratory of Reproductive Biology, Graduate School of Agriculture, Kyoto University,

Kyoto 606-8502, Japan

Running Title: MERVL-PRMT6 regulates cell differentiation in mouse embryos.

Corresponding author: N Minami (e-mail: minami.naojirou.6r@kyoto-u.ac.jp)

Abstract

Murine endogenous retrovirus with leucine tRNA primer, also known as MERVL, is

expressed during zygotic genome activation in mammalian embryos. Here we show that

protein arginine N-methyltransferase 6 (*Prmt6*) forms a chimeric transcript with MT2B2,

one of the long terminal repeat sequences of MERVL, and is translated into an elongated

chimeric protein (PRMT6^{MT2B2}) whose function differs from that of the canonical

PRMT6 protein (PRMT6^{CAN}) in mouse preimplantation embryos. Overexpression of

PRMT6^{CAN} in fibroblast cells increased asymmetric dimethylation of the third arginine

residue of both histone H2A (H2AR3me2a) and histone H4 (H4R3me2a), while

overexpression of PRMT6^{MT2B2} increased only H2AR3me2a. In addition, overexpression

of PRMT6^{MT2B2} in one blastomere of mouse two-cell embryos promoted cell proliferation

and differentiation of the blastomere into epiblast cells at the blastocyst stage, while

overexpression of PRMT6^{CAN} repressed cell proliferation. This is the first report of the

translation of a chimeric protein (PRMT6^{MT2B2}) in mouse preimplantation embryos. Our

results suggest that analyzing chimeric transcripts with MERVL will provide insight into

the relationship between zygotic genome activation and subsequent intra- and extra-

cellular lineage determination.

Key words: Mouse, endogenous retrovirus, *Prmt6*, preimplantation embryo,

development

Introduction

Immediately after fertilization, mammalian embryos have low transcriptional activity and translate proteins from mRNA stored in oocytes. In mouse embryos, transcription subsequently begins with two waves of zygotic genome activation (ZGA), namely minor and major ZGA, the former during the S phase of the one-cell stage and the latter during the late two-cell stage (Minami et al., 2007; Schulz and Harrison, 2019). Before and after major ZGA, the epigenome of the fertilized egg changes dynamically (Aoki, 2022; Burton and Torres-Padilla, 2010; Xu et al., 2021). Epigenetic modifications, including DNA methylation and histone modifications, are known to be important for cell differentiation during development (Wu and Sun, 2006). For example, *Carm1*, encoding a protein arginine methyltransferase, has been reported to be involved in the differentiation of blastomeres into epiblast cells through the dimethylation of histone arginine residues in mouse preimplantation embryos (Torres-Padilla et al., 2007; White et al., 2016). Endogenous retroviruses (ERVs) are one of the components of transposable elements and occupy about 10% of human and mouse genomes (Ito et al., 2020; Mouse Genome Sequencing Consortium, 2002). Although ERVs are among the earliest transcribed genetic material in mouse preimplantation embryos (Kigami et al., 2003) and various studies have shown that ERVs are expressed in mammalian preimplantation

embryos (Evsikov et al., 2004; Poznanski and Calarco; Wang et al., 2001), there is little

direct evidence that ERVs regulate cell differentiation. The high abundance of ERVs in

the mammalian genome (Ito et al., 2020) and the fact that several ERV-derived genes,

such as *Syncytin* and *Peg10*, are essential for placentation (Dupressoir et al., 2009;

Dupressoir et al., 2012; Mi et al., 2000; Suzuki et al., 2007) suggest that ERVs were

involved in the evolution of the placenta in mammals. Murine or human endogenous

retrovirus with leucine tRNA primer (MERVL or HERVL, respectively) is a transposable

element expressed during major ZGA in mice and humans (Franke et al., 2017; Kigami

et al., 2003; Macfarlan et al., 2012; Wang et al., 2001). It has been reported that MERVL-

activated mouse embryonic stem cells injected into eight-cell embryos can also contribute

to extra-embryonic tissues (Yang et al., 2020), suggesting that some MERVL/HERVL

elements may regulate the totipotency of mammalian embryos. Recently, a MERVL-

derived non-coding RNA called LincGET was shown to control the differentiation of

embryonic and extra-embryonic tissues in mouse preimplantation embryos (Wang et al.,

2018), but the function of the vast majority of MERVL elements remains to be elucidated.

Several studies have shown that the long terminal repeat of MERVL generates chimeric

transcripts with host genes during early embryonic development (Macfarlan et al., 2012;

Peaston et al., 2004). Because MERVL has been reported to be one of the earliest

transcribed genomic sequences in preimplantation embryos (Kigami et al., 2003; Wang

et al., 2001), we hypothesized that MERVL chimeric transcripts contribute to the cell

lineage differentiation that occurs after major ZGA (Chazaud and Yamanaka, 2016).

Reanalysis of previously reported RNA-sequencing (RNA-seq) data revealed

that protein arginine N-methyltransferase 6 (*Prmt6*) expression is activated by the

MERVL promoter sequence in preimplantation embryos. Like *Carm1*, *Prmt6* encodes a

Type-I protein arginine methyltransferase (Di Lorenzo and Bedford, 2011) and has been

shown to regulate the epigenome by the asymmetric di-methylation of arginine residues

in histone proteins (Guccione et al., 2007; Hyllus et al., 2007; Kirmizis et al., 2007).

Furthermore, overexpression or suppression of *Prmt6* in embryonic stem cells results in

the loss of pluripotency and in the increased expression of differentiation marker genes

(Lee et al., 2012). In the course of analyzing transcripts involved in MERVL in mouse

preimplantation embryos, it was discovered that *Prmt6* has two transcripts of different

lengths. One is the canonical *Prmt6* transcript ($Prmt6^{CAN}$), and the other is $Prmt6^{MT2B2}$,

defined as the chimeric transcript of $Prmt6^{CAN}$ with MT2B2, the MERVL long terminal

repeat sequence. It was also found that the chimeric transcript produced an elongated

PRMT6 chimeric protein ($PRMT6^{MT2B2}$). In addition, the chimeric protein has different

histone modification specificity from that of the canonical PRMT6 protein ($PRMT6^{CAN}$),

and overexpression of the chimeric protein in one blastomere of two-cell embryos

promotes its contribution to the embryonic cell lineage. These findings shed light on a

new aspect of the differentiation mechanism during embryogenesis.

Results

Screening of candidate MERVL chimeric transcripts associated with cell lineage differentiation

Macfarlan et al. published a list of MERVL chimeric transcripts in mouse two-cell stage embryos (Macfarlan et al., 2012), and we compared these transcripts with histone modification–related genes (Fig. 1A). Although there was no statistically significant bias, eight genes (*Atxn7*, *Brms1l*, *Ctcf*, *Kdm4c*, *Pcgf5*, *Prmt6*, *Rnf20*, and *Smyd3*) overlapped, and half of them (*Ctcf*, *Kdm4c*, *Rnf20*, and *Smyd3*) had already been reported to be important in early embryonic development (Ooga et al., 2015; Suzuki et al., 2015; Wan et al., 2008; Wang et al., 2010). Analysis of a published RNA-seq dataset (Zhang et al., 2016) revealed that the expression levels of four of the eight genes (*Ctcf*, *Kdm4c*, *Pcgf5*, and *Prmt6*) were more than twice as high in late two-cell embryos compared with early two-cell embryos (Fig. 1B)

Expression dynamics of chimeric and canonical Prmt6 mRNA in mouse preimplantation embryos

Genomic analysis in mice revealed that MT2B2 is located 212 bp upstream of the first exon of *Prmt6*, and that the chimeric transcript is transcribed from around 380 bp

upstream of the canonical transcription start site. To determine the expression profiles of

$Prmt6^{MT2B2}$ and $Prmt6^{CAN}$, the amounts of total $Prmt6$ transcripts ($Prmt6^{MT2B2}$ and

$Prmt6^{CAN}$) and $Prmt6^{MT2B2}$ during preimplantation development were determined by

quantitative RT-PCR (Fig. 2A). As expected, the amounts of total $Prmt6$ transcripts and

$Prmt6^{MT2B2}$ transcripts in the late two-cell stage were increased at major ZGA and

gradually decreased by the blastocyst stage (Fig. 2B). At the blastocyst stage, $Prmt6^{MT2B2}$

transcripts had almost completely disappeared, but $Prmt6^{CAN}$ transcripts were still present,

albeit in small amounts (Fig. 2B). To compare the exact copy numbers of each transcript,

absolute quantification was performed using late two-cell embryos and blastocysts. In

late two-cell embryos, the amounts of total $Prmt6$ transcripts and chimeric transcripts

were almost the same (Fig. 2C). In blastocysts, on the other hand, the copy number of

$Prmt6^{MT2B2}$ transcripts was significantly lower than that of total $Prmt6$ transcripts (Fig.

2C). Furthermore, microinjecting zygotes with si$Prmt6^{MT2B2}$ (siRNA targeting the

sequence between the MT2B2 and $Prmt6^{CAN}$ transcription start sites) or si$Prmt6^{CAN}$-1 or

si$Prmt6^{CAN}$-2 (siRNAs targeting the $Prmt6$ exon sequence, see Fig. 2A) resulted in a

significant decrease in transcript levels of both total $Prmt6$ transcripts and $Prmt6^{MT2B2}$

transcripts at the late two-cell stage (Fig. 2D). Since trimethylation of the fourth lysine

residue of histone H3 (H3K4me3) is generally increased around transcription start sites,

we analyzed the previously reported chromatin immunoprecipitation sequencing (ChIP-

seq) data of H3K4me3 (Liu et al., 2016) in preimplantation embryos and found that the

weak H3K4me3 ChIP-seq peak appeared at the MT2B2 region from the 2-cell stage and

became stronger at the 4-cell stage (Fig. 2E). On the other hand, the H3K4me3 peak at

just before the first exon of $Prmt6^{CAN}$ was observed from the four-cell stage and get higher

at the eight-cell stage (Fig. 2E).

Identification of the translation initiation codon of the PRMT6^{MT2B2} chimeric protein

Since $Prmt6^{MT2B2}$ has an extended sequence on the 5' side of $Prmt6^{CAN}$ and two

putative translation initiation codons exist in this region, we sought to determine the site

where translation of the chimeric transcript began (Fig. 3A). Western blotting of the

PRMT6 protein using metaphase II (MII) oocytes and fertilized eggs at each

developmental stage revealed that ~55 kDa bands were detected in oocytes and embryos

during all developmental stages, and an additional ~42 kDa band was detected only in

embryos at the blastocyst stage (Fig. 3B). Since PRMT6^{CAN} has a molecular weight of

~42 kDa, mRNAs of $Prmt6^{CAN}$ and $Prmt6^{MT2B2}$ were generated by in vitro transcription

to identify the translation product of the ~55 kDa protein detected at each developmental

stage. The mRNAs transfected into NIH3T3 cells revealed that $Prmt6^{MT2B2}$ mRNA

encodes an elongated protein of ~55 kDa, while $Prmt6^{CAN}$ mRNA encodes a protein of

~42 kDa (Fig. 3C). Consistent with the amount of transcript, PRMT6^{MT2B2} with a

molecular weight of ~55 kDa was most abundant at the late two-cell stage and gradually

decreased until the blastocyst stage (Fig. 3B). There are two candidate translation

initiation codons (Met1 and Met2) for $Prmt6^{MT2B2}$, and Met3 is the translation initiation

codon for PRMT6^{CAN} (Fig. 3A). To determine which codon $Prmt6^{MT2B2}$ is translated from,

three types of mRNAs mutated in Met1, Met2, or Met3 (AUG to AAG) were transfected

into NIH3T3 cells. Among NIH3T3 cells transfected with the three mutant mRNAs, only

those with the Met1 mutant transcript failed to produce the ~55 kDa PRMT6^{MT2B2} (Fig.

3C). The same results were obtained using mouse four-cell embryos (Fig. 3C). Bands at

~55 kDa were detected in all four-cell embryos because these embryos express the

endogenous PRMT6^{MT2B2} (Fig. 3C). Knockdown of endogenous $Prmt6$ transcripts with

siRNAs (si$Prmt6^{CAN}$-1 or si$Prmt6^{CMT2B2}$) at the one-cell stage decreased the amount of

the ~55 kDa protein at the four-cell stage (Fig. 3D).

Functional analysis of the PRMT6^{MT2B2} chimeric protein

Three-dimensional structure prediction using AlphaFold (Jumper et al., 2021)

showed no structural change in the S-adenosylmethionine–dependent methyltransferase

domain of PRMT6^{MT2B2} compared to PRMT6^{CAN} (Fig. 4A). Therefore, the ~55 kDa

PRMT6^{MT2B2} was considered to have arginine methylation activity, and the function of

this protein and its involvement in cell differentiation in mammalian preimplantation

embryos were examined. Since PRMT6^{CAN} is known to be a type I protein arginine

methyltransferase (Di Lorenzo and Bedford, 2011), western blotting was performed using

various histone arginine asymmetric dimethylation antibodies to confirm the histone

arginine methylation status of NIH3T3 cells overexpressing PRMT6^{CAN} or PRMT6^{MT2B2}.

Overexpression of PRMT6^{CAN} significantly increased asymmetric dimethylation of the

third arginine residue of both histone H2A (H2AR3me2a) and histone H4 (H4R3me2a)

(Fig. 4B, 4C, 4D). By contrast, overexpression of PRMT6^{MT2B2} significantly increased

only H2AR3me2a, while the amount is lower than overexpression of PRMT6^{CAN} (Fig.

4B, 4C, 4D; $P<0.01$).

PRMT6^{MT2B2} regulates cell differentiation in mouse preimplantation embryos

Analysis of published single cell RNA-seq data (Goolam et al., 2016) showed that

Prmt6 is asymmetrically distributed among blastomeres in four- and eight-cell embryos

(Supplemental Fig. 1A). Next, we confirmed H2AR3me2a distribution among

blastomeres at each developmental stage. Immunostaining of H2AR3me2a showed an

asymmetrical distribution among blastomeres after the eight-cell stage and a greater

abundance in the trophectoderm (TE) than in the inner cell mass (ICM) at the blastocyst

stage (Supplemental Fig. 1B). $Prmt6^{CAN}$ or $Prmt6^{MT2B2}$ mRNA together with *H2B-EGFP*

mRNA as a lineage tracer was microinjected into one blastomere of two-cell embryos to

examine the effects on cell lineage specification in mouse preimplantation embryos (Fig.

5A). Overexpression of $PRMT6^{MT2B2}$ significantly increased the number of H2B-EGFP–

positive cells in blastocysts at 96 h post insemination (hpi) compared to overexpression

of H2B-EGFP alone, while $PRMT6^{CAN}$ overexpression reduced the number of H2B-

EGFP–positive cells in blastocysts at 96 hpi (Fig. 5B, 5C). In addition, overexpression of

$PRMT6^{MT2B2}$ significantly increased the percentage of NANOG-positive epiblast cells in

H2B-EGFP–positive cells in blastocysts than overexpression of H2B-EGFP alone (Fig.

5B, 5D), while this percentage was unaffected by $PRMT6^{CAN}$ overexpression (Fig. 5D).

The differentiation into trophectoderm (CDX2-positive cells) and primitive endoderm

(NANOG-negative and CDX2-negative cells) were not changed by $PRMT6^{CAN}$ or

$PRMT6^{MT2B2}$ overexpression (Supplementary Table 3). The si$Prmt6^{CAN}$-1 injection into

one blastomere of two-cell embryos resulted in the low number of H2B-EGFP–positive

cells in blastocyst, while neither si$Prmt6^{CAN}$-1 nor si$Prmt6^{MT2B2}$ injection influence cell

differentiation (Supplementary Table 4). When $Prmt6^{MT2B2}$ or $Prmt6^{CAN}$ mRNA was

injected in one-cell embryos, cell numbers were not changed comparing with EGFP

mRNA injected control embryos, while the percentage of CDX2-positive cells was

decreased in PRMT6^{MT2B2} overexpressed embryos (Supplementary Table 5).

Discussion

Among the histone modification–related genes expressed as chimeric transcripts with MERVL, the expression of four genes (*Ctcf*, *Kdm4c*, *Pcgf5*, and *Prmt6*) dramatically increased during major ZGA, suggesting that many genes expressed during this period may be mainly under control of the MERVL promoter. Indeed, the results of absolute quantitative RT-PCR (Fig. 2C) and siRNA knockdown experiments targeting the intermediate sequence between MT2B2 and *Prmt6* exon (Fig. 2D) showed that almost all *Prmt6* transcripts were *Prmt6^{MT2B2}* at the late two-cell stage. In contrast, most *Prmt6^{CAN}* transcripts were expressed at the blastocyst stage, indicating that the transcriptional machinery of *Prmt6* is dramatically altered during early embryogenesis. Analysis of H3K4me3 ChIP-seq data supported the dynamic change of promoter activity, but the H3K4me3 peak at the MT2B2 region at the two-cell stage is weaker than the four- and eight-cell stage. The temporal mismatch between H3K4me3 ChIP-seq data and our quantitative RT-PCR data could be due to the difference of mice strain (Bertozzi et al., 2020) or collection time of 2-cell embryos.

Overexpression experiments of *Prmt6^{MT2B2}* mRNA with a mutation in the translation initiation codon revealed that the *Prmt6^{MT2B2}* chimeric transcript is translated into the PRMT6^{MT2B2} chimeric protein from the 5'-extended uppermost ATG

codon (Fig. 3C, 3D). PRMT6^{MT2B2} is the first MERVL chimeric protein translated from

a chimeric transcript during mammalian development. Outside of early embryogenesis,

one case of a MERVL chimeric transcript translated into a longer protein was reported

in cancer cells (Lock et al., 2014). Since the extended 56-amino-acid sequence of

PRMT6^{MT2B2} showed no homology to retrovirus-derived proteins such as the GAG

protein (data not shown), further studies on the origin of the elongated sequence are

needed to understand the biological functions of PRMT6^{MT2B2} chimeric proteins.

Although many chimeric transcripts are expressed in cancer cells and embryonic stem

cells (Babaian and Mager, 2016; Macfarlan et al., 2012), their exact translation

initiation codons and biological functions remain unknown, and more comprehensive

proteomic analysis is needed to understand the precise function of chimeric transcripts,

including those expressed in preimplantation embryos.

Our results also revealed that the PRMT6^{MT2B2} chimeric protein has more

substrate-specific and/or lower histone arginine methylation activity than PRMT6^{CAN}

(Fig. 4B, 4C), suggesting that changes in epigenomic modifications may regulate

developmental progression and gene expression in early preimplantation embryos.

H2AR3me2a is asymmetrically distributed among blastomeres after the 8-cell stage and

is more abundant in trophectoderm at the blastocyst stage (Supplemental Fig. 1B). Since

PRMT6^{CAN} has a stronger ability to modify H2AR3me2a than PRMT6^{MT2B2} (Fig. 4C), it

is possible that the transcription shifts from *Prmt6^{MT2B2}* to *Prmt6^{CAN}* in some blastomeres

after the 8-cell stage, resulting in increased amounts of H2AR3me2a and enhanced

differentiation to trophectoderms. PRMT6 is known to be expressed in several cancer

tissues, including colorectal, breast, bladder, and lung adenocarcinomas (Avasarala et al.,

2020; Lim et al., 2018; Tang et al., 2020; Veland et al., 2017; Yoshimatsu et al., 2011).

Investigating how PRMT6^{MT2B2} is associated with cancer progression will provide insight

into the involvement of retroviruses in cancer development (Attermann et al., 2018;

Babaian and Mager, 2016; Kassiotis and Stoye, 2016; María et al., 2016) and a new

perspective on cancer research. The PRMT6^{CAN} protein contains a nuclear localization

signal (NLS) at the N-terminus (Fig. 3A) (Frankel et al., 2002; Herrmann et al., 2005).

Since it is known that the modification of surrounding amino acids and protein

conformation affect NLS function (Sorokin et al., 2007), it is possible that the extended

amino acid sequence of PRMT6^{MT2B2} leads to loss of NLS function and lower catalytic

activity for histone proteins. Since other arginine methyltransferases are known to

methylate proteins localized in the cytoplasm (Di Lorenzo and Bedford, 2011; Litt et al.,

2009), analysis of the subcellular localization of PRMT6^{MT2B2} could provide new insights

into its function.

In blastocysts, most *Prmt6* transcripts are presumably *Prmt6^{CAN}* based on

absolute quantification (Fig. 2C). Blastomeres overexpressing PRMT6^{MT2B2} were more

likely to consist of epiblast cells than non-injected blastomeres, while those

overexpressing PRMT6^{CAN} did not contribute to each cell lineage (Fig. 4G). In addition,

Prmt6^{MT2B2} mRNA microinjection to one-cell embryos showed lower CDX2-positive

cell rate at the blastocyst stage, which is stronger phenotype than *Prmt6^{CAN}* mRNA

injected embryos. These indicates that PRMT6^{MT2B2} and PRMT6^{CAN} have different

functions in cell differentiation during embryogenesis. It has been reported that

H3R2me2a mediated by PRMT6^{CAN} recruits Aurora B to chromosome arms (Kim et

al., 2020) and that overexpression of Aurora B facilitates cell contribution to the

placental cell lineage by regulating the nuclear dynamics of OCT4 in mouse

preimplantation embryos (Li et al., 2017; Plachta et al., 2011). Based on these results,

it is conceivable that an increased ratio of PRMT6^{CAN} to PRMT6^{MT2B2} promotes cell

differentiation during the blastocyst stage through localization of PRMT6^{CAN} in the

nucleus. In addition, several protein arginine methyltransferases have been reported to

act redundantly (Cheng et al., 2020; Wei et al., 2021). *Carm1* was found to promote

differentiation of blastomeres to epiblast cells via methylation of H3R26me2 (Goolam

et al., 2016; Torres-Padilla et al., 2007; White et al., 2016), and *Prmt1* enzymic inhibitor

promoted the differentiation of embryonic stem cells into primitive endoderm via Klf4

methylation (Zuo et al., 2022). These results support the possibility that PRMT6^{MT2B2},

but not PRMT6^{CAN}, may regulate the differentiation pathways in blastocysts.

Microinjection of PRMT6^{CAN} or si*Prmt6^{CAN}*-1 in a single blastomere in two-

cell stage embryos resulted in delayed cell proliferation (Fig. 4F), probably since

PRMT6^{CAN} controls the cell cycle, as reported in previous studies (Kim et al., 2020;

Schneider et al., 2021). On the other hand, overexpression of PRMT6^{MT2B2} promoted

cell proliferation (Fig. 4F), indicating that PRMT6^{CAN} and PRMT6^{MT2B2} affect cell

proliferation in different ways. Modzelewski et al. reported that the MERVL-*Cdk2ap1*

chimeric transcript is translated into a shorter protein (CDK2AP1$^{\Delta N}$) than the canonical

protein (CDK1AP1^{CAN}) in early mammalian embryos, and that CDK2AP1$^{\Delta N}$ promotes

cell proliferation while CDK2AP1^{CAN} inhibits the cell cycle (Modzelewski et al., 2021).

In mice, ZGA is known to occur earlier and to have a shorter implantation period than

in other mammals (Modzelewski et al., 2021). Adaptive evolution for rapid

development of mouse embryos may have occurred through the insertion of MT2B2

just before the Prmt6 locus, which allowed translation of the Prmt6MT2B2 chimeric

protein and may have promoted cell proliferation after ZGA. These findings and our

own indicate that chimeric transcripts expressed during major ZGA mediate different

functions than canonical transcripts.

In conclusion, we have shown that the chimeric transcript $Prmt6^{MT2B2}$ is

expressed in mouse preimplantation embryos and that it generates the chimeric protein

PRMT6^{MT2B2} during major ZGA. Furthermore, PRMT6^{MT2B2} expressed in early

preimplantation embryos is involved in determining cell differentiation during the

blastocyst stage. This study reveals that MERVL expressed at major ZGA regulates cell

differentiation during embryogenesis by regulating the expression of nearby genes and

the translation of their corresponding mRNA.

Materials and Methods

In vitro fertilization and embryo culture

Eight- to twelve-week-old ICR female mice (Japan SLC, Hamamatsu, Japan) were superovulated by injection of 7.5 IU of equine chorionic gonadotropin (eCG; ASKA Pharmaceutical, Tokyo, Japan) followed by 7.5 IU of human chorionic gonadotropin (hCG; ASKA Pharmaceutical) 48 h later. Cumulus oocyte complexes were harvested 14 h after hCG injection and placed in human tubal fluid (HTF) medium supplemented with 4 mg/ml bovine serum albumin (BSA, A3311; Sigma-Aldrich, St. Louis, MO) covered with paraffin oil (Nacalai Tesque, Kyoto, Japan). Spermatozoa were collected from the cauda epididymis of 13–20-week-old ICR male mice (Japan SLC) and cultured in HTF medium for 1 h. After capacitation, sperm were introduced into oocyte-containing droplets at a final concentration of 1×10^6 cells/ml. After 3-h incubation at 37°C in an atmosphere of 5% CO_2, fertilized oocytes were washed with K^+-modified simplex optimized medium (KSOM) supplemented with amino acids (Ho et al., 1995) and 1 mg/ml BSA, and then used for microinjection or cultured in the same medium under paraffin oil at 37°C in an atmosphere of 5% CO_2 to the following stages: one-cell (12 h post insemination (hpi)), early two-cell (24 hpi), late two-cell (36 hpi), four-cell (48 hpi), eight-cell (54 hpi), morula (72 hpi), and blastocyst (96 hpi). MII oocytes were collected

from cumulus oocyte complexes followed by treatment with 1% hyaluronidase to remove

cumulus cells.

Plasmid construction, in vitro transcription, and microinjection

Based on the transcription start site of *Prmt6^{CAN}* predicted from the NCBI

database (NM_178891.5) and that of *Prmt6^{MT2B2}* predicted from previously reported

RNA-seq data (Liu et al., 2016), the sequences of *Prmt6^{CAN}* and *Prmt6^{MT2B2}* were cloned

by PCR from the cDNA of late two-cell embryos. The amplified products were digested

with EcoRI and XbaI, and the fragments were incorporated into a pBluescript II SK (−)

vector plasmid (Agilent Technologies Japan, Hachioji, Japan). Start codon mutant

constructs were generated by mutation PCR using the PrimeSTAR Mutagenesis Basal Kit

(Takara Bio, Kusatsu, Japan) by replacing AUG with AAG at Met1, Met2, or Met3. The

AAG codon was chosen because this mutation has been reported to cause complete loss

of translational activity in eukaryotic cells compared to other single-nucleotide mutations

(Wei et al., 2013). Methylase-inactive KLA mutant constructs were generated using the

same method by replacing amino acids Val–Leu–Asp at positions 89–91 in PRMT6^{CAN}

with Lys–Leu–Ala to produce PRMT^{CAN}-KLA (Boulanger et al., 2005), or by replacing

these amino acids at positions 145–147 in PRMT6^{MT2B2} with Lys–Leu–Ala to produce

PRMT6^{MT2B2}-KLA. Each mRNA was synthesized with the mMESSAGE mMECHINE

T7 Ultra Kit (Thermo Fisher Scientific, Waltham, MA), purified with the RNeasy Mini

Kit (Qiagen, Hilden, Germany), and then dissolved in nuclease-free water. For

knockdown experiments, siRNAs (RNAi, Tokyo, Japan) dissolved in nuclease-free water

at a concentration of 100 μM were used for microinjection. For overexpression

experiments, *Prmt6^{CAN}* or *Prmt6^{MT2B2}* mRNA was diluted to 250 ng/μl and co-injected

with *H2B-EGFP* mRNA at a concentration of 100 ng/μl. Approximately 5–10 pl of

mRNA or siRNA was microinjected into the cytoplasm of one-cell embryos at 3–5 hpi or

into the cytoplasm of one blastomere of two-cell embryos at 24–26 hpi in HEPES-

buffered KSOM. Microinjection was performed under an inverted microscope (IX73;

Olympus, Tokyo, Japan) equipped with a piezo injector (PMAS-CT150; PRIME TECH,

Tokyo, Japan) and a micromanipulator (IM-11-2; Narishige, Tokyo, Japan). The primers

and siRNA sequences are listed in Supplementary Tables 1 and 2.

Cell culture and mRNA transfection

NIH/3T3 cells were cultured in Dulbecco's modified Eagle's medium

supplemented with 10% calf serum, 75 μg/ml penicillin, and 50 μg/ml streptomycin.

mRNAs transcribed in vitro were transfected into 50–60% confluent cells using

Lipofectamine MessengerMAX Transfection Reagent (Thermo Fisher Scientific).

Twenty-four hours after transfection, the cells were washed in phosphate-buffered saline

(PBS) once and then lysed in 1X Laemmli sample buffer.

RNA extraction and quantitative RT-PCR

RNA extraction and reverse transcription were performed using the SuperPrep

II Cell Lysis & RT Kit for qPCR (Toyobo, Osaka, Japan). Briefly, five embryos or

oocytes were washed three times in PBS containing 0.5% polyvinylpyrrolidone (PVP K-

30; Nacalai Tesque) (PVP-PBS) and collected in PCR tubes along with 0.5 µl of PVP-

PBS. After adding 3.5 µl of lysate, all lysates were subjected to reverse transcription.

Transcription levels were measured as previously described (Shikata et al., 2020) using

the $2^{-\Delta\Delta Ct}$ method normalized against the corresponding *H2afz* levels (Jeong et al., 2014;

Livak and Schmittgen, 2001). Absolute expression was determined using serial dilutions

of the chimeric *Prmt6* cloned plasmid for in vitro transcription or using the pTAC plasmid

in which an *H2afz* PCR fragment was cloned. The primers used for quantification are

listed in Supplementary Table 2.

Western blotting

Fifty embryos or oocytes, excluding blastocysts, were washed three times in

PVP-PBS and collected in 1X Laemmli sample buffer. Twenty-five blastocysts were

collected. After boiling at 95°C for 5 min, the samples were kept at −80°C until use.

NIH3T3 cells were directly lysed in 1X Laemmli sample buffer. To analyze histone

arginine methylation modifications in NIH3T3 cells, TagRFP, PRMT6^{CAN}-KLA, or

PRMT6^{MT2B2}-KLA were used as negative controls for overexpression. Total protein was

separated by SDS–polyacrylamide gel electrophoresis (SDS-PAGE) and transferred into

a 0.22-µm-pore polyvinylidene fluoride (PVDF) membrane (Cytiva, Tokyo, Japan). After

one wash in PBS containing 0.1% Tween (PBST), the membrane was blocked in PBST

containing 5% BSA for 1 h at room temperature, followed by incubation with a rabbit

anti-PRMT6 antibody (1:1500 dilution; 15395-1-AP; Proteintech, Rosemont, IL), a rabbit

anti-H2AR3me2a antibody (1:1000 dilution; ab21574; Abcam, Cambridge, UK), a rabbit

anti-H4R3me2a antibody (1:500 dilution; 39706; Active Motif, Carlsbad, CA), a rabbit

anti-H3R2me2a antibody (1:1000 dilution; ab175007; Abcam), a rabbit anti-H3R8me2a

antibody (1:500 dilution; 39652; Active Motif), a rabbit anti-H3R17me2a antibody

(1:2000 dilution; 39710; Active Motif), a rabbit anti-H3R26me2 antibody (1:8000

dilution; 07-215; Merck, Darmstadt, Germany), a mouse anti–β-actin antibody (1:5000

dilution; A5441; Merck), or a rabbit anti–whole-H3 antibody (1:1000 dilution; #9715;

Cell Signaling, Danvers, MA) in PBST containing 1% BSA overnight at 4°C. After three

washes in PBST, the membrane was incubated with a horseradish peroxidase–conjugated

anti-mouse secondary antibody (1:10000 dilution; NA931; Cytiva) or anti-rabbit

secondary antibody (1:10000 dilution; NA934; Cytiva) in PBST containing 1% BSA for

1 h at room temperature. After three washes in PBST, the membrane was developed using

ImmunoStar LD (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan) and

scanned with c-DiGit (LI-COR, Lincoln, NE). Densitometric quantification of the

immunoblot bands was performed using Image Studio software (LI-COR).

Immunofluorescence

Embryos were fixed in PBS containing 4% paraformaldehyde for 20 min at 28°C.

After three washes in PVP-PBS, embryos were permeabilized with 0.5% Triton X-100

(Sigma-Aldrich) in PBS for 40 min at 28°C. After washing three times in PVP-PBS,

embryos were blocked in PBS containing 1.5% BSA, 0.02% Tween 20, and 0.2% sodium

azide (blocking buffer) for 1 h at 28°C, and then incubated overnight at 4°C with a rabbit

anti-NANOG antibody (1:100 dilution; RCAB002P-F; ReproCELL, Kanagawa, Japan)

and a mouse anti-CDX2 antibody (1:100 dilution; MU392A-UC; Biogenex, Fremont,

CA). After three washes in blocking buffer, embryos were incubated in blocking buffer

containing secondary antibodies (Alexa Fluor 555 donkey anti-rabbit IgG, 1:500 dilution,

A31572, Thermo Fisher Scientific; Alexa Fluor 647 donkey anti-mouse IgG, 1:500

dilution, A31571, Thermo Fisher Scientific) for 1 h at 28°C. After washing three times in

blocking buffer, nuclei were stained in blocking buffer containing 10 µg/ml Hoechst

33342 (Sigma-Aldrich) for 10 min at 28°C. Stained embryos were mounted on slide glass

in blocking buffer containing 50% glycerol and signals were observed with a confocal

fluorescent microscope (LSM880; Zeiss, Oberkochen, Germany). H2B-EGFP protein

was detected based on its own fluorescence.

RNA-seq and ChIP-seq data processing

The bulk RNA-seq data (GSE71434) and the single cell RNA-seq data (E-

MTAB-3321) of embryos of each stage were downloaded and then mapped to the mm10

with STAR (v. 2.7.10a) (Dobin et al., 2013). The gene expression level was estimated

using RSEM (v. 1.3.3) (Li and Dewey, 2011) and normalized with the transcripts per

kilobase million method. The H3K4me3 ChIP-seq data of mouse preimplantation

embryos were downloaded from GSE73952 and mapped to the mm10 with bowtie2 (v.

2.4.5) (Langmead and Salzberg, 2012). In all cases, Bam files of biological replicates

were merged using samtools (v. 1.10) (Li et al., 2009), normalized by counts per million

using deeptools (v. 3.5.1) (Ramírez et al., 2016), and visualized using Integrative

Genomics Viewer software (v. 2.12.1) (Robinson et al., 2011).

Three-dimensional Structure Prediction of the PRMT6^{MT2B2} Protein

PRMT6^{CAN} and PRMT6^{MT2B2} protein structure prediction was performed using

Alphafold (Jumper et al., 2021) on the Google Colaboratory site

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.i

pynb). Obtained data were visualized using the Mol* site (Sehnal et al., 2021)

(https://molstar.org/).

Statistical analyses

Quantitative RT-PCR data were analyzed by Student's t-test for pairwise

comparisons or by Dunnett's test for multiple comparisons. Quantification of western

blotting was performed using the Tukey-Kramer test. Cell contribution assays were

analyzed with the Wilcoxon signed-rank test with holm's adjustment. All analyses were

performed using R (v. 4.2.1), and significance was accepted at P-values < 0.05.

Ethical approval for the use of animals

All animal experiments were approved by the Animal Research Committee of

Kyoto University (Permit Numbers: R3-17) and were performed in accordance with the

committee's guidelines.

Competing interest statement

The authors declare no competing interests.

Acknowledgments

Confocal fluorescent microscopy was performed at the iCeMS Analysis Center, Institute for Integrated Cell-Material Sciences (iCeMS), Kyoto University Institute for Advanced Study (KUIAS). This work was supported by a Grant-in-Aid for Scientific Research (no. 19H03136 to N.M.) and a Grant-in-Aid for JSPS Fellows (no. 19J23290 to S.H.) from the Japan Society for the Promotion of Science.

Author contributions: The experiments were designed by S.H. and N.M., and were performed by S.H., with contributions from Y.K. and M.H. The manuscript was written by S.H. and N.M., with contributions from S.I. All authors read and approved the final manuscript.

References

Aoki, F. (2022). Zygotic gene activation in mice: profile and regulation. *J. Reprod. Dev.*

68, 79–84.

Attermann, A. S., Bjerregaard, A.-M., Saini, S. K., Grønbæk, K. and Hadrup, S. R.

(2018). Human endogenous retroviruses and their implication for

immunotherapeutics of cancer. *Annals of Oncology* **29**, 2183–2191.

Avasarala, S., Wu, P.-Y., Khan, S. Q., Yanlin, S., Van Scoyk, M., Bao, J., Di Lorenzo,

A., David, O., Bedford, M. T., Gupta, V., et al. (2020). PRMT6 Promotes Lung

Tumor Progression via the Alternate Activation of Tumor-Associated

Macrophages. *Molecular Cancer Research* **18**, 166–178.

Babaian, A. and Mager, D. L. (2016). Endogenous retroviral promoter exaptation in

human cancer. *Mobile DNA* **7**, 24.

Bertozzi, T. M., Elmer, J. L., Macfarlan, T. S. and Ferguson-Smith, A. C. (2020).

KRAB zinc finger protein diversification drives mammalian interindividual

methylation variability. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 31290–31300.

Boulanger, M.-C., Liang, C., Russell, R. S., Lin, R., Bedford, M. T., Wainberg, M. A.

and Richard, S. (2005). Methylation of Tat by PRMT6 Regulates Human

Immunodeficiency Virus Type 1 Gene Expression. *J Virol* **79**, 124–131.

Burton, A. and Torres-Padilla, M.-E. (2010). Epigenetic reprogramming and

development: a unique heterochromatin organization in the preimplantation

mouse embryo. *Briefings in Functional Genomics* **9**, 444–454.

Chazaud, C. and Yamanaka, Y. (2016). Lineage specification in the mouse

preimplantation embryo. *Development* **143**, 1063–1074.

Cheng, D., Gao, G., Di Lorenzo, A., Jayne, S., Hottiger, M. O., Richard, S. and

Bedford, M. T. (2020). Genetic evidence for partial redundancy between the

arginine methyltransferases CARM1 and PRMT6. *Journal of Biological*

Chemistry **295**, 17060–17070.

Di Lorenzo, A. and Bedford, M. T. (2011). Histone arginine methylation. *FEBS Letters*

585, 2024–2031.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,

Chaisson, M. and Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq

aligner. *Bioinformatics* **29**, 15–21.

Dupressoir, A., Vernochet, C., Bawa, O., Harper, F., Pierron, G., Opolon, P. and

Heidmann, T. (2009). Syncytin-A knockout mice demonstrate the critical role in

placentation of a fusogenic, endogenous retrovirus-derived, envelope gene. *Proc.*

Natl. Acad. Sci. U.S.A. **106**, 12127–12132.

Dupressoir, A., Lavialle, C. and Heidmann, T. (2012). From ancestral infectious

retroviruses to bona fide cellular genes: Role of the captured syncytins in

placentation. *Placenta* **33**, 663–671.

Evsikov, A. V., de Vries, W. N., Peaston, A. E., Radford, E. E., Fancher, K. S., Chen,

F. H., Blake, J. A., Bult, C. J., Latham, K. E., Solter, D., et al. (2004). Systems

biology of the 2-cell mouse embryo. *Cytogenet Genome Res* **105**, 240–250.

Franke, V., Ganesh, S., Karlic, R., Malik, R., Pasulka, J., Horvat, F., Kuzman, M.,

Fulka, H., Cernohorska, M., Urbanova, J., et al. (2017). Long terminal repeats

power evolution of genes and gene expression programs in mammalian oocytes

and zygotes. *Genome Res.* **27**, 1384–1394.

Frankel, A., Yadav, N., Lee, J., Branscombe, T. L., Clarke, S. and Bedford, M. T.

(2002). The Novel Human Protein Arginine N-Methyltransferase PRMT6 Is a

Nuclear Enzyme Displaying Unique Substrate Specificity. *Journal of Biological*

Chemistry **277**, 3537–3543.

Goolam, M., Scialdone, A., Graham, S. J. L., Macaulay, I. C., Jedrusik, A.,

Hupalowska, A., Voet, T., Marioni, J. C. and Zernicka-Goetz, M. (2016).

Heterogeneity in Oct4 and Sox2 Targets Biases Cell Fate in 4-Cell Mouse

Embryos. *Cell* **165**, 61–74.

Guccione, E., Bassi, C., Casadio, F., Martinato, F., Cesaroni, M., Schuchlautz, H.,

Lüscher, B. and Amati, B. (2007). Methylation of histone H3R2 by PRMT6 and

H3K4 by an MLL complex are mutually exclusive. *Nature* **449**, 933–937.

Herrmann, F., Lee, J., Bedford, M. T. and Fackelmayer, F. O. (2005). Dynamics of

Human Protein Arginine Methyltransferase 1(PRMT1) in Vivo. *Journal of*

Biological Chemistry **280**, 38005–38010.

Ho, Y., Wigglesworth, K., Eppig, J. J. and Schultz, R. M. (1995). Preimplantation

development of mouse embryos in KSOM: Augmentation by amino acids and

analysis of gene expression. *Mol. Reprod. Dev.* **41**, 232–238.

Hyllus, D., Stein, C., Schnabel, K., Schiltz, E., Imhof, A., Dou, Y., Hsieh, J. and Bauer,

U.-M. (2007). PRMT6-mediated methylation of R2 in histone H3 antagonizes H3

K4 trimethylation. *Genes Dev.* **21**, 3369–3380.

Ito, J., Gifford, R. J. and Sato, K. (2020). Retroviruses drive the rapid evolution of

mammalian *APOBEC3* genes. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 610–618.

Jeong, J.-K., Kang, M.-H., Gurunathan, S., Cho, S.-G., Park, C., Seo, H. G. and Kim,

J.-H. (2014). Evaluation of reference genes in mouse preimplantation embryos

for gene expression studies using real-time quantitative RT-PCR (RT-qPCR).

BMC Res Notes **7**, 675.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly

accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589.

Kassiotis, G. and Stoye, J. P. (2016). Immune responses to endogenous retroelements:

taking the bad with the good. *Nat Rev Immunol* **16**, 207–219.

Kigami, D., Minami, N., Takayama, H. and Imai, H. (2003). MuERV-L Is One of the

Earliest Transcribed Genes in Mouse One-Cell Embryos1. *Biology of*

Reproduction **68**, 651–654.

Kim, S., Kim, N. H., Park, J. E., Hwang, J. W., Myung, N., Hwang, K.-T., Kim, Y.

A., Jang, C.-Y. and Kim, Y. K. (2020). PRMT6-mediated H3R2me2a guides

Aurora B to chromosome arms for proper chromosome segregation. *Nat Commun*

11, 612.

Kirmizis, A., Santos-Rosa, H., Penkett, C. J., Singer, M. A., Vermeulen, M., Mann,

M., Bähler, J., Green, R. D. and Kouzarides, T. (2007). Arginine methylation

at histone H3R2 controls deposition of H3K4 trimethylation. *Nature* **449**, 928–

932.

Langmead, B. and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2.

Nat Methods **9**, 357–359.

Lee, Y. H., Ma, H., Tan, T. Z., Ng, S. S., Soong, R., Mori, S., Fu, X.-Y., Zernicka-

Goetz, M. and Wu, Q. (2012). Protein Arginine Methyltransferase 6 Regulates

Embryonic Stem Cell Identity. *Stem Cells and Development* **21**, 2613–2622.

Li, B. and Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-

Seq data with or without a reference genome.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,

Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing

Subgroup (2009). The Sequence Alignment/Map format and SAMtools.

Bioinformatics **25**, 2078–2079.

Li, W., Wang, P., Zhang, B., Zhang, J., Ming, J., Xie, W. and Na, J. (2017). Differential

regulation of H3S10 phosphorylation, mitosis progression and cell fate by Aurora

Kinase B and C in mouse preimplantation embryos. *Protein & Cell* **8**, 662–674.

Lim, Y., Yu, S., Yun, J.-A., Do, I.-G., Cho, L., Kim, Y. H. and Kim, H. C. (2018). The

prognostic significance of protein arginine methyltransferase 6 expression in

colon cancer. *Oncotarget* **9**, 9010–9020.

Litt, M., Qiu, Y. and Huang, S. (2009). Histone arginine methylations: their roles in

chromatin dynamics and transcriptional regulation. *Bioscience Reports* **29**, 131–

141.

Liu, X., Wang, C., Liu, W., Li, J., Li, C., Kou, X., Chen, J., Zhao, Y., Gao, H., Wang, H., et al. (2016). Distinct features of H3K4me3 and H3K27me3 chromatin domains in pre-implantation embryos. *Nature* **537**, 558–562.

Livak, K. J. and Schmittgen, T. D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408.

Lock, F. E., Rebollo, R., Miceli-Royer, K., Gagnier, L., Kuah, S., Babaian, A., Sistiaga-Poveda, M., Lai, C. B., Nemirovsky, O., Serrano, I., et al. (2014). Distinct isoform of FABP7 revealed by screening for retroelement-activated genes in diffuse large B-cell lymphoma. *Proc. Natl. Acad. Sci. U.S.A.* **111**,.

Macfarlan, T. S., Gifford, W. D., Driscoll, S., Lettieri, K., Rowe, H. M., Bonanomi, D., Firth, A., Singer, O., Trono, D. and Pfaff, S. L. (2012). Embryonic stem cell potency fluctuates with endogenous retrovirus activity. *Nature* **487**, 57–63.

María, G.-C., Paola, I., Niki, K., Mariacarmela, S., Julià, B. and Rafael, R. (2016). Human endogenous retroviruses and cancer. *Cancer Biology & Medicine* **13**, 483.

Mi, S., Lee, X., Li, X., Veldman, G. M., Finnerty, H., Racie, L., LaVallie, E., Tang, X.-Y., Edouard, P., Howes, S., et al. (2000). Syncytin is a captive retroviral envelope protein involved in human placental morphogenesis. *Nature* **403**, 785–

789.

Minami, N., Suzuki, T. and Tsukamoto, S. (2007). Zygotic Gene Activation and Maternal Factors in Mammals. *J. Reprod. Dev.* **53**, 707–715.

Modzelewski, A. J., Shao, W., Chen, J., Lee, A., Qi, X., Noon, M., Tjokro, K., Sales, G., Biton, A., Anand, A., et al. (2021). A mouse-specific retrotransposon drives a conserved Cdk2ap1 isoform essential for development. *Cell* **184**, 5541-5558.e22.

Mouse Genome Sequencing Consortium (2002). Initial sequencing and comparative analysis of the mouse genome. *Nature* **420**, 520–562.

Ooga, M., Suzuki, M. G. and Aoki, F. (2015). Involvement of histone H2B monoubiquitination in the regulation of mouse preimplantation development. *Journal of Reproduction and Development* **61**, 179–184.

Peaston, A. E., Evsikov, A. V., Graber, J. H., de Vries, W. N., Holbrook, A. E., Solter, D. and Knowles, B. B. (2004). Retrotransposons Regulate Host Genes in Mouse Oocytes and Preimplantation Embryos. *Developmental Cell* **7**, 597–606.

Plachta, N., Bollenbach, T., Pease, S., Fraser, S. E. and Pantazis, P. (2011). Oct4 kinetics predict cell lineage patterning in the early mammalian embryo. *Nat Cell Biol* **13**, 117–123.

Poznanski, A. A. and Calarco, P. G. The Expression of lntracisternal A Particle Genes

in the Preimplantation Mouse Embryo.

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne,

S., Dündar, F. and Manke, T. (2016). deepTools2: a next generation web server

for deep-sequencing data analysis. *Nucleic Acids Res* **44**, W160–W165.

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz,

G. and Mesirov, J. P. (2011). Integrative genomics viewer. *Nat Biotechnol* **29**,

24–26.

Schneider, L., Herkt, S., Wang, L., Feld, C., Wesely, J., Kuvardina, O. N., Meyer, A.,

Oellerich, T., Häupl, B., Seifried, E., et al. (2021). PRMT6 activates cyclin D1

expression in conjunction with the transcription factor LEF1. *Oncogenesis* **10**, 42.

Schulz, K. N. and Harrison, M. M. (2019). Mechanisms regulating zygotic genome

activation. *Nat Rev Genet* **20**, 221–234.

Sehnal, D., Bittrich, S., Deshpande, M., Svobodová, R., Berka, K., Bazgier, V.,

Velankar, S., Burley, S. K., Koča, J. and Rose, A. S. (2021). Mol* Viewer:

modern web app for 3D visualization and analysis of large biomolecular structures.

Nucleic Acids Research **49**, W431–W437.

Shikata, D., Yamamoto, T., Honda, S., Ikeda, S. and Minami, N. (2020). H4K20

monomethylation inhibition causes loss of genomic integrity in mouse

624   preimplantation embryos. *J. Reprod. Dev.* **66**, 411–419.

625   **Sorokin, A. V., Kim, E. R. and Ovchinnikov, L. P.** (2007). Nucleocytoplasmic transport

626   of proteins. *Biochemistry Moscow* **72**, 1439–1457.

627   **Suzuki, S., Ono, R., Narita, T., Pask, A. J., Shaw, G., Wang, C., Kohda, T., Alsop, A.**

628   **E., Marshall Graves, J. A., Kohara, Y., et al.** (2007). Retrotransposon Silencing

629   by DNA Methylation Can Drive Mammalian Genomic Imprinting. *PLoS Genet* **3**,

630   e55.

631   **Suzuki, S., Nozawa, Y., Tsukamoto, S., Kaneko, T., Imai, H. and Minami, N.** (2015).

632   Histone methyltransferase Smyd3 regulates early embryonic lineage commitment

633   in mice. *REPRODUCTION* **150**, 21–30.

634   **Tang, J., Meng, Q., Shi, R. and Xu, Y.** (2020). PRMT6 serves an oncogenic role in lung

635   adenocarcinoma via regulating p18. *Mol Med Rep*.

636   **Torres-Padilla, M.-E., Parfitt, D.-E., Kouzarides, T. and Zernicka-Goetz, M.** (2007).

637   Histone arginine methylation regulates pluripotency in the early mouse embryo.

638   *Nature* **445**, 214–218.

639   **Veland, N., Hardikar, S., Zhong, Y., Gayatri, S., Dan, J., Strahl, B. D., Rothbart, S.**

640   **B., Bedford, M. T. and Chen, T.** (2017). The Arginine Methyltransferase PRMT6

641   Regulates DNA Methylation and Contributes to Global DNA Hypomethylation in

642   Cancer. *Cell Reports* **21**, 3390–3397.

643 **Wan, L.-B., Pan, H., Hannenhalli, S., Cheng, Y., Ma, J., Fedoriw, A., Lobanenkov,**

644   **V., Latham, K. E., Schultz, R. M. and Bartolomei, M. S.** (2008). Maternal

645   depletion of CTCF reveals multiple functions during oocyte and preimplantation

646   embryo development. *Development* **135**, 2729–2738.

647 **Wang, Q., Chung, Y. G., deVries, W. N., Struwe, M. and Latham, K. E.** (2001). Role

648   of Protein Synthesis in the Development of a Transcriptionally Permissive State

649   in One-Cell Stage Mouse Embryos1. *Biology of Reproduction* **65**, 748–754.

650 **Wang, J., Zhang, M., Zhang, Y., Kou, Z., Han, Z., Chen, D.-Y., Sun, Q.-Y. and Gao,**

651   **S.** (2010). The Histone Demethylase JMJD2C Is Stage-Specifically Expressed in

652   Preimplantation Mouse Embryos and Is Required for Embryonic Development1.

653   *Biology of Reproduction* **82**, 105–111.

654 **Wang, J., Wang, L., Feng, G., Wang, Y., Li, Y., Li, X., Liu, C., Jiao, G., Huang, C.,**

655   **Shi, J., et al.** (2018). Asymmetric Expression of LincGET Biases Cell Fate in

656   Two-Cell Mouse Embryos. *Cell* **175**, 1887-1901.e18.

657 **Wei, J., Zhang, Y., Ivanov, I. P. and Sachs, M. S.** (2013). The Stringency of Start Codon

658   Selection in the Filamentous Fungus Neurospora crassa. *Journal of Biological*

659   *Chemistry* **288**, 9549–9562.

Wei, H.-H., Fan, X.-J., Hu, Y., Tian, X.-X., Guo, M., Mao, M.-W., Fang, Z.-Y., Wu,

P., Gao, S.-X., Peng, C., et al. (2021). A systematic survey of PRMT interactomes

reveals the key roles of arginine methylation in the global control of RNA splicing

and translation. *Science Bulletin* **66**, 1342–1357.

White, M. D., Angiolini, J. F., Alvarez, Y. D., Kaur, G., Zhao, Z. W., Mocskos, E.,

Bruno, L., Bissiere, S., Levi, V. and Plachta, N. (2016). Long-Lived Binding of

Sox2 to DNA Predicts Cell Fate in the Four-Cell Mouse Embryo. *Cell* **165**, 75–

87.

Wu, H. and Sun, Y. E. (2006). Epigenetic Regulation of Stem Cell Differentiation.

Pediatr Res **59**, 21R-25R.

Xu, R., Li, C., Liu, X. and Gao, S. (2021). Insights into epigenetic patterns in

mammalian early embryos. *Protein Cell* **12**, 7–28.

Yang, F., Huang, X., Zang, R., Chen, J., Fidalgo, M., Sanchez-Priego, C., Yang, J.,

Caichen, A., Ma, F., Macfarlan, T., et al. (2020). DUX-miR-344-ZMYM2-

Mediated Activation of MERVL LTRs Induces a Totipotent 2C-like State. *Cell*

Stem Cell **26**, 234-250.e7.

Yoshimatsu, M., Toyokawa, G., Hayami, S., Unoki, M., Tsunoda, T., Field, H. I.,

Kelly, J. D., Neal, D. E., Maehara, Y., Ponder, B. A. J., et al. (2011).

Dysregulation of PRMT1 and PRMT6, Type I arginine methyltransferases, is

involved in various types of human cancers. *Int. J. Cancer* **128**, 562–573.

Zhang, B., Zheng, H., Huang, B., Li, W., Xiang, Y., Peng, X., Ming, J., Wu, X., Zhang,

Y., Xu, Q., et al. (2016). Allelic reprogramming of the histone modification

H3K4me3 in early mammalian development. *Nature* **537**, 553–557.

Zuo, Z., Yang, G., Wang, H., Liu, S., Zhang, Y., Cai, Y., Chen, F., Dai, H., Xiao, Y.,

Cheng, M., et al. (2022). Klf4 methylated by Prmt1 restrains the commitment of

primitive endoderm. *Nucleic Acids Research* **50**, 2005–2018.

Fig. 1. Verification of MERVL-mediated chimeric genes.

(A) Venn diagram showing overlap of genes expressed as chimeric transcripts with

MERVL- and histone modification–related genes. The list of MERVL chimeric genes

were obtained from published data (Macfarlan et al., 2012), and the list of histone

modification-related genes were obtained from MGI database

(http://www.informatics.jax.org) in Jan. 2019. (B) The expression patterns of eight

overlapping genes in MII oocytes and preimplantation embryos. Dots represent

replicates. Data were obtained from GSE71434. MII: MII oocytes, Zy: zygotes, E2C:

early two-cell embryos, 4C: four-cell embryos, 8C: eight-cell embryos, ICM: inner cell

mass cells.

Fig. 2. Gene structure and transcriptional profile of *Prmt6* in mice.

(A) Schematic diagram showing the positions of MT2B2 and *Prmt6* in the mouse

genome. MT2B2 is located 212 bp upstream from the transcription start site of *Prmt6*.

The names of primers and targeting loci (arrows) are indicated above the diagram, and

the names of siRNAs and their targeting sites (bars) are indicated underneath. (B) qRT-

PCR analysis of the *Prmt6* exon and *Prmt6*^{MT2B2} transcripts during mouse

preimplantation development. The amount of mRNA in one-cell embryos was defined

as 1, and the expression levels were normalized to *H2afz* as an internal control. Data

are expressed as mean ± s.e.m. (n=3). (C) Comparison of expression levels of the *Prmt6*

exon and *Prmt6^{MT2B2}* in late two-cell embryos and blastocysts based on absolute

quantification by qRT-PCR. Each graph shows the expression levels of each transcript

in late two-cell embryos and blastocysts. Copy numbers were normalized to *H2afz* copy

numbers. Data are expressed as mean ± s.e.m. (n=5). Student's t-test, ***P < 0.001;

n.s., not significant. (D) qRT-PCR analysis of *Prmt6* and *Prmt6^{MT2B2}* mRNA abundance

in embryos injected with si*Prmt6^{MT2B2}*, si*Prmt6*-1, si*Prmt6*-2, or siControl in the late

two-cell stage. Injection was performed 3–5 h after IVF. The amount of mRNA in

siControl-injected embryos was defined as 1, and the expression levels were

normalized to *H2afz* as an internal control. Data are expressed as mean ± s.e.m. (n=3).

Dunnett's test, ***P < 0.001. (E) Genome browser view of H3K4me3 ChIP-seq data

from each stage of mouse preimplantation embryos. Data for MII oocytes, two-cell

embryos, four-cell embryos, eight-cell embryos, morulae, inner cell mass (ICM) and

trophectoderm (TE) of blastocysts, embryonic stem cells (ESC), and trophoblast stem

cells (TSC) are shown. The *Prmt6^{CAN}* exon and MT2B2 locations are indicated at the

bottom. Data were obtained from GSE73952. A magnified view of the data for MII

oocytes and two-cell embryos are shown in the upper panel.

1C: one-cell embryo, E2C: early two-cell embryo, L2C: late two-cell embryo, 4C: four-

cell embryo, 8C: eight-cell embryo, Mo: morula, Bl: blastocyst

Fig. 3. Protein expression of Prmt6.

(A) Schematic diagram of the genome structure (upper) and genome and protein

sequences (lower) of *Prmt6* exon 1. Candidate translation initiation codons (Met1 and

Met2) and the canonical translation initiation codon (Met3) are indicated above the

schematic diagram and are shown in bold in the sequences. (B) Western blotting of

PRMT6 in mouse oocytes and preimplantation embryos. β-actin was used as a loading

control. Fifty oocytes or embryos were used in each lane except blastocysts. Twenty-

five blastocysts were used. The approximate molecular weight is on the right. The

representative result is shown from three independent experiments. (C) Western

blotting analysis for PRMT6 in NIH3T3 cells and four-cell embryos. NIH3T3 cells

were transfected with TagRFP mRNA, *Prmt6^{CAN}* mRNA, *Prmt6^{MT2B2}* mRNA,

Prmt6^{MT2B2} with Met1 mutant mRNA (Met1 AAG), *Prmt6^{MT2B2}* with Met2 mutant

mRNA (Met2 AAG), or *Prmt6^{MT2B2}* with Met3 mutant mRNA (Met3 AAG). Zygotes

were injected with the same mRNAs except *Prmt6^{MT2B2}* with Met3 mutant mRNA

(Met3 AAG) and cultured to the four-cell stage. Fifty embryos were used in each lane.

β-actin was used as a loading control. The approximate molecular weight is on the right.

The representative result is shown from three independent experiments. (D) Western

blotting analysis of PRMT6 protein in Prmt6-suppressed four-cell embryos.

si*Prmt6^{CAN}*-1, si*Prmt6^{MT2B2}*, or siControl was injected into zygote cytoplasm. β-actin

was used as a loading control. Fifty embryos were used for each lane. The approximate

molecular weight is on the right. The representative result is shown from three

independent experiments.

MII: metaphase II oocyte, 1C: one-cell embryo, E2C: early two-cell embryo, L2C: late

two-cell embryo, 4C: four-cell embryo, 8C: eight-cell embryo, Mo: morula, Bl:

blastocyst

Fig. 4. Substrate-specific histone arginine methylation activity of PRMT6^{MT2B2}.

(A) The 3D structures of PRMT6^{CAN} and PRMT6^{MT2B2} as predicted by AlphaFold2.

The blue structure represents PRMT6^{CAN} and the green structure represents

PRMT6^{MT2B2}. The five S-adenosylmethionine binding sites are shown in red. The right

panel shows a magnified view of the catalytic domain. (B) Western blotting analysis of

various histone arginine methylation modifications in NIH3T3 cells overexpressing

TagRFP, PRMT6^{CAN}, PRMT6^{CAN}-KLA, PRMT6^{MT2B2}, or PRMT6^{MT2B2}-KLA. β-actin

and whole H3 were used as loading controls. KLA-containing proteins are those in

which the enzyme activity is eliminated by replacing amino acids VLD with KLA. (C,

D) The amounts of H2AR3me2 (C) and H4R3me2 (D) based on the intensity

of (B). Overexpression of PRMT6^{CAN} increases H2AR3me2a and H4R3me2a intensity,

while overexpression of PRMT6^{MT2B2} increases only H2AR3me2a intensity. Data are

expressed as mean ± s.d. (n=3). Tukey-Kramer test, ***P < 0.001; **P < 0.01; *P <

0.05; n.s., not significant.

C	Blastocysts 3D	Total cells (n)	NANOG⁺ cells (%)	GFP⁺ cells (%)	NANOG⁺ cells that are GFP⁺ (%)	NANOG⁺ cells that are GFP⁻ (%)	GFP⁺ cells that are NANOG⁺ (%)
H2B-EGFP control	55	78.47 ± 13.69	(11.09 ± 5.34)	(48.63 ± 7.96)	(12.30 ± 7.61)	(10.39 ± 7.68)	(53.70 ± 23.92)
PRMT6^{CAN}-OE	49	75.00 ± 16.28	(12.09 ± 5.65)	(43.81 ± 8.56)*	(14.54 ± 11.14)	(10.26 ± 5.85)	(49.08 ± 23.97)
PRMT6^{MT2B2}-OE	58	75.83 ± 14.49	(13.44 ± 6.19)*	(51.77 ± 8.44)*	(15.65 ± 8.54)*	(10.87 ± 7.15)	(60.82 ± 21.92)

Fig. 5. PRMT6^{MT2B2} determines the cell fate of mouse preimplantation embryos.

(A) Schematic view of the cell fate contribution assay. *Prmt6^{CAN}* or *Prmt6^{MT2B2}* mRNA

together with *H2B-EGFP* mRNA was microinjected into one blastomere of two-cell

embryos. Embryos cultured until the blastocyst stage (96 hpi) were subjected to

immunostaining using antibodies against NANOG (red) and CDX2 (white). Cell nuclei

were stained with Hoechst (blue). The number of NANOG-positive cells was counted

separately for H2B-EGFP–positive and –negative cells. (B) Immunostaining of

NANOG and CDX2 in blastocysts overexpressing H2B-EGFP (control), PRMT6^{CAN}

with H2B-EGFP (PRMT6^{CAN}-OE), or PRMT6^{MT2B2} with H2B-EGFP (PRMT6^{MT2B2}-

OE). Photos are representative of immunostained blastocysts. Hoechst (blue), H2B-

EGFP (marker, green), NANOG (red), CDX2 (white), and 3D maximum projections of

merged images are shown. Scale bars, 20 μm. (C) Analysis of the distribution of

progeny of injected blastomere at the blastocyst stage based on (B). Data were

represented as mean ±s.d. Wilcoxon signed-rank test with holm's adjustment were used

for statistical analysis, *P < 0.05. Seven experimental replicates were performed. Key

to table headings: Blastocysts 3D is the number of blastocysts that were all analyzed,

Total cells (n) is the total number of cells in the blastocyst, NANOG$^+$ cells (%) is the

percentage of NANOG-positive cells out of the total number of cells in the blastocyst,

GFP$^+$ cells (%) is the percentage of GFP-positive cells out of the total number of cells

in the blastocyst, NANOG$^+$ cells that are GFP$^+$ (%) is the percentage of NANOG-

positive cells out of the total number of GFP-positive cells in the blastocyst, NANOG$^+$

cells that are GFP$^-$ (%) is the percentage of NANOG-positive cells out of the total

number of GFP-negative cells in the blastocyst, and GFP$^+$ cells that are NANOG$^+$ (%)

is the percentage of GFP-positive cells out of the total number of NANOG-positive

cells in the blastocyst.

Supplementary Information for

Chimeric protein of PRMT6 produced by endogenous retrovirus promoter regulates cell fate decision in mouse

preimplantation embryos

Shinnosuke Honda, Maho Hatamura, Yuri Kunimoto, Shuntaro Ikeda, and Naojiro Minami*

Laboratory of Reproductive Biology, Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

*Corresponding author

Email: minami.naojirou.6r@kyoto-u.ac.jp

Supplementary Fig. 1. Asymmetric distribution of *Prmt6* mRNA and H2AR3me2a.

(A) Average of the relative *Prmt6* mRNA levels in each blastomere at four- and eight-cell stage. Each blastomere at 4- or 8-cell stage is named B1 to B4 (4-cell, n=16) or B1 to B8 (8-cell, n=4) according to the amount of *Prmt6* expression. TPM (Transcripts Per Million) was normalized based on the blastomere with highest *Prmt6* expression in each embryo, and the mean averages are shown (± s.e.m.). Data were obtained from E-MTAB-3321. (B) Immunostaining for H2AR3me2a in embryos at each developmental stage. Hoechst (blue), H2AR3me2a (red), and CDX2 or OCT4 (green, only at the blastocyst stage) are shown. Scale bars, 50 μm. Arrowheads indicate the cell with high fluorescence and asterisks indicate the cell with low fluorescence in four- and eight-cell embryos. Dotted lines mark the inner cell mass in blastocysts.

Supplementary Table 1. Sequences of siRNAs used for microinjection

Names of siRNAs	Sense Strand (5'→3')	Antisense Strand (5'→3')
si*Prmt6* ^{MT2B2}	GCCAAGAUUGACUCUGGUUdTdT	AACCAGAGUCAAUCUUGGCdTdT
si*Prmt6* ^{CAN}-1	GCCGUCUGGUUUCAAGUGAdCdC	UCACUUGAAACCAGACGGCdGdG
si*Prmt6* ^{CAN}-2	CACCGAAGCCUACCGCUUAdGdG	UAAGCGGUAGGCUUCGGUGdCdC
siControl	GUACCGCACGUCAUUCGUAUC	UACGAAUGACGUGCGGUACGU

dT, 2'-deoxythymidine; dC, 2'-deoxycytidine; dG, 2'-deoxyguanosine

Supplementarry Table 2. Primer sequences

for quantitative PCR

Primer names		Sequences	Product sizes (bp)
Prmt6 exon	sense	AGTGCTACTCCGACGTCTCG	149
	anti-sense	GATGCTGAGAATGCCGGTGC	
Prmt6 ^{MT2B2}	sense	AGGGGTAAAAGGAAAGGGCAA	278
	anti-sense	GTCCCCCGACTCAAGCTTTC	
H2afz	sense	GGTAAAGCGTATCACCCCTCG	130
	anti-sense	CTTCCCGATCAGCGATTTGTG	

for plasmid construction

Primer names		Sequences	Product sizes (bp)
Prmt6 ^{MT2B2} cloning	sense	AAAGAATTCAGAGAACCCTGACAAGGGGTA	1408
	anti-sense	AAATCTAGATCAGTCCTCCATGGCAAAGTC	
Prmt6 ^{CAN} cloning	sense	AAAGAATTCATGTCGCTGAGCAAGAAAAGAA	1155
	anti-sense	AAATCTAGATCAGTCCTCCATGGCAAAGTC	
Met1 mutation	sense	CTTGTCAAGGCGTTAGAGGAAAATAA	—
	anti-sense	TAACGCCTTGACAAGTTGTTCTCTAA	
Met2 mutation	sense	AGCGGGAAGCAGGAAGCGGACTACGA	—
	anti-sense	TTCCTGCTTCCCGCTTCCTGAGGGCA	
Met3 mutation	sense	GCCAAGAAGTCGCTGAGCAAGAAAG	—
	anti-sense	CAGCGACTTCTTGGCCCCGGCAGCCG	
KLA mutation	sense	ACGAAGCTTGCGGTGGGCGCGGGCACCGGC	—
	anti-sense	CACCGCAAGCTTCGTCTTGCCTCGCAGCGC	

for in vitro transcription

Primer names		Sequences	Product sizes (bp)
Prmt6 ^{CAN}, *Prmt6* ^{MT2B2} IVT	sense	TAAGTTGGGTAACGCCAGGG	1351, 1604
	anti-sense	AGCGCGCAATTAACCCTCAC	
H2B-EGFP IVT	sense	TAAGTTGGGTAACGCCAGGG	1501
	anti-sense	GGAGCAGATACGAATGGCTACA	

Supplementary Table 3. Analysis of the distribution of progeny of injected blastomere at the blastocyst stage based on Fig. 5B.

	Blastocysts 3D	CDX2$^+$ cells (%)	CDX2$^+$ cells that are GFP$^+$ (%)	CDX2$^+$ cells that are GFP$^-$ (%)	NANOG$^-$ CDX2$^-$ cells (%)	NANOG$^-$ CDX2$^-$ cells that are GFP$^+$ (%)	NANOG$^-$ CDX2$^-$ cells that are GFP$^-$ (%)
H2B-EGFP control	55	(68.15 ± 9.23)	(68.26 ± 10.81)	(68.17 ± 13.09)	(23.83 ± 6.71)	(22.33 ± 9.69)	(23.12 ± 11.92)
PRMT6CAN-OE	49	(68.86 ± 7.30)	(68.43 ± 11.97)	(69.34 ± 8.13)	(20.08 ± 7.32)	(20.18 ± 8.89)	(21.25 ± 8.62)
PRMT6MT2B2-OE	58	(69.07 ± 7.43)	(68.29 ± 9.89)	(69.91 ± 11.57)	(18.38 ± 7.13)	(19.46 ± 9.04)	(20.87 ± 10.24)

Data were represented as mean ± s.d. Wilcoxon signed-rank test with holm's adjustment were used for statistical analysis. Seven experimental replicates were performed.

	Blastocysts 3D	CDX2$^+$ cells (%)	CDX2$^+$ cells that are GFP$^+$ (%)	CDX2$^+$ cells that are GFP$^-$ (%)	NANOG$^-$ CDX2$^-$ cells (%)	NANOG$^-$ CDX2$^-$ cells that are GFP$^+$ (%)	NANOG$^-$ CDX2$^-$ cells that are GFP$^-$ (%)

Supplementary Table 4. Analysis of the distribution of progeny of siRNA injected two-cell blastomere at the blastocyst stage.

	Blastocysts 3D	Total cells (n)	NANOG$^+$ cells (%)	CDX2$^+$ cells (%)	GFP$^+$ cells (%)	NANOG$^+$ cells that are GFP$^+$ (%)	NANOG$^+$ cells that are GFP$^-$ (%)	GFP$^+$ cells that are NANOG$^+$ (%)
siControl	37	77.62 ± 13.07	(10.41 ± 7.91)	(73.01 ± 7.94)	(49.20 ± 7.24)	(11.17 ± 13.49)	(9.79 ± 8.35)	(49.36 ± 31.70)
siPrmt6-1	29	70.79 ± 12.34	(11.74 ± 6.43)	(73.74 ± 7.99)	(44.19 ± 9.83)*	(14.40 ± 9.58)	(9.82 ± 7.47)	(54.58 ± 23.97)
siERVL-Prmt6	31	76.45 ± 12.63	(10.49 ± 4.93)	(70.13 ± 11.13)	(48.20 ± 7.26)	(11.23 ± 7.31)	(9.65 ± 6.19)	(49.52 ± 27.86)

Data were represented as mean ± s.d. Wilcoxon signed-rank test with holm's adjustment were used for statistical analysis, *P < 0.05. Three experimental replicates were performed.

Supplementary Table 5. Analysis of the cell fate distribution of *EGFP*, *Prmt6*[CAN] or *Prmt6*[MT2B2] mRNA injected one-cell embryos at the blastocyst stage.

	Blastcysts	Total cells (n)	NANOG$^+$ cells (%)	CDX2$^+$ cells (%)	NANOG$^-$ CDX2$^-$ cells (%)
EGFP control	39	77.49 ± 9.19	(15.50 ± 5.51)	(77.26 ± 6.07)	(9.62 ± 6.63)
Prmt6^{CAN}-OE	35	78.11 ± 10.15	(15.32 ± 6.98)	(73.85 ± 8.98)	(12.58 ± 7.54)
Prmt6^{MT2B2}-OE	34	78.21 ± 11.68	(14.34 ± 4.64)	(73.31 ± 8.16)*	(13.22 ± 8.59)*

Data were represented as mean ± s.d. Wilcoxon signed-rank test with holm's adjustment were used for statistical analysis, *P < 0.05. Three experimental replicates were performed.