The impact of maternal dietary folic acid or choline deficiencies on cerebral blood flow,
cardiac, aortic, and coronary function in young and middle-aged female mouse offspring
following ischemic stroke

Running Title: Impact of maternal diet on offspring blood flow

Kasey Pull BA[1], Robert Folk BS[1], Jeemin Kang MA[1,2], Shaley Jackson BSc[3], Brikena Gusek BS[1],
Mitra Esfandiarei PhD*[1,4,5], Nafisa M. Jadavji PhD*[1,2,3,6,7]

[1] Department of Biomedical Sciences, Midwestern University, Glendale, AZ, USA

[2] College of Osteopathic Medicine, Midwestern University, Glendale, AZ, USA

[3] College of Veterinary Medicine, Midwestern University, Glendale, AZ, USA

[4] Basic Medical Sciences, College of Medicine Phoenix, University of Arizona, Phoenix, AZ,
USA

[5] Anesthesiology, Pharmacology, & Therapeutics, Faculty of Medicine, University of British
Columbia, Vancouver, BC, Canada

[6] Department of Child Health, College of Medicine Phoenix, University of Arizona, Phoenix, AZ,
USA

[7] Department of Neuroscience, Carleton University, Ottawa, ON, Canada

* Correspondence:
18
Nafisa M. Jadavji, PhD
Department of Biomedical Sciences
Midwestern University
19555 N 59th Ave
Glendale, AZ 85308
Phone: 623-572-3665
Fax: 623-572-3647
nafisa.jadavji@mail.mcgill.ca
27
28
29 Mitra Esfandiarei, PhD
30 mesfan@midwestern.edu
31
32

Abstract

Background and Purpose

Adequate maternal dietary levels of one-carbon (1C) metabolites, such as folic acid and choline, play an important role in the closure of the neural tube *in utero*; however, the impact of deficiencies in 1C on offspring neurological function after birth remain undefined. Stroke is one of the leading causes of death and disability globally. The aim of our study was to determine the impact of maternal 1C nutritional deficiencies on cerebral and peripheral blood flow after ischemic stroke in adult female offspring.

Method

In this study, female mice were placed on either control (CD), folic acid (FADD), or choline (ChDD) deficient diets prior to pregnancy. Female offspring were weaned onto a CD for the duration of the study. Ischemic stroke was induced in offspring and after six weeks cerebral and peripheral blood flow velocity was measured using ultrasound imaging.

Results

Our data showed that 11.5-month-old female offspring from ChDD mothers had reduced blood flow in the posterior cerebral artery compared to controls. In peripheral blood flow velocity measurements, we report an aging effect.

Conclusions

These results emphasize the importance of maternal 1C diet in early life neuro-programming on long-term vasculature health.

Keywords

58 One-carbon metabolism, Ischemic stroke, Photothrombosis, Maternal diet, Posterior cerebral

59 artery, Blood flow

Introduction

Maternal nutrition during pregnancy and lactation is recognized as a critical factor determining the health of offspring [1–5]. In addition, early life nutritional cofactors are also critical for fetal development, as well as in determining offspring disease outcome later in life[6,7]. The Developmental Origins of Health and Disease (DOHaD) theory suggests that prospective chronic diseases are programmed *in utero* [8–11]. While already compensating for fetal nutrient accumulation and increased maternal metabolic demands [12], altered or insufficient maternal nutrition can impact both early development and future offspring health. In offspring, maternal dietary deficiencies have been associated with ventricular septal defects [13] and impaired glucose tolerance [14], as well as modified neural tube closure [15,16] and neurocognitive development [17–21]. Beyond this evidence of suboptimal structural development in offspring, maternal nutritional deficiencies have been linked to programming of offspring metabolic [22] and epigenetic [23–28] adaptations.

Epidemiological studies have demonstrated the effect of maternal diet on lifelong cardiovascular and neurological function [29–31]. Most of this population-level work reveals an effect of poor maternal health on birthweight and incidence of disease and cardiovascular risk factors in adulthood, such as hypertension and hyperlipidemia [4,10]. Such relationships have been shown in numerous global populations and are apparent from birth through early childhood [32]. Among nutritional cofactors, folates and choline are important players in healthy fetal neurodevelopment due to their involvement in the closure of the neural tube and are components of one-carbon metabolism [33,34]. Folates and its chemically synthesized form folic acid are important for fetal neurodevelopment [35], as folate requirements during pregnancy are increased by 5- to 10-fold compared to non-pregnant women [36]. Maternal folates and choline levels during pregnancy have also been shown to be important in the development of the cerebellum and hippocampus [37], as well as affecting postnatal myelination trajectories [38], short-term memory [39],

hyperactivity/attention [40], neurocognitive development [41], and risk of autism spectrum disorder

(ASD) [42].

Recent work in rodent models has improved mechanistic understanding of how maternal levels of

folate and choline impact neurodevelopmental processes [43]. Akin to human epidemiological

studies [29–31], murine maternal folate deficiencies have been implicated in adverse reproductive

performance, implantation, and fetal growth [44]. During pregnancy and lactation, maternal dietary

folic acid availability has been shown to influence progenitor cell mitosis, and apoptosis in the

fetal mouse forebrain [45] and hippocampus [39]. Investigations of maternal perinatal folate

deficiencies have revealed reduced hippocampal proliferation, impaired vesicular transport and

synaptic plasticity, as well as poor neurite outgrowth [46], modified cellular neocortex composition,

and diminished complexity and arborization of projection neurons [47] in offspring. In addition to

these structural observations, both genetic and epigenetic modifications have been observed. For

example maternal folate deficiencies has been shown to reduce expressions levels of brain

derived neurotropic factor (BDNF) and H3K9me2 (an epigenetic modification to the DNA

packaging protein Histone H3) in the fetal hippocampus, while folic acid deficiency for two

generations, being reported to significantly enhance *de novo* mutations accumulation during

meiosis [48].

Choline, another one-carbon metabolite implicated in a number of diverse biological processes [49],

has yielded variable results in animal models of neurodevelopment. Effects of maternal choline

deficiencies, such as defective layering of the cortex, reduced cortical size and brain weight [50],

and modified hippocampal electrophysiology [51] and neurogenesis [52], have been observed in

offspring. Beyond these physiological and histological findings, choline and folate deficiencies

have been shown to elicit similar adverse effects, such as impaired homocysteine re-methylation,

oxidative stress, and endothelial dysfunction in murine cerebrovasculature [53,54]; effectively

demonstrating the link between maternal one-carbon metabolites and typical fetal

neurodevelopment.

The link between maternal nutrition and fetal development is abundantly clear, but the long-term

(postnatal) effects of maternal nutritional deficiencies on adult offspring are not clearly

understood. In this study, we aim to improve our understanding of one-carbon metabolite dietary

requirements during pregnancy and its impact on early life programming of adult offspring

neurovascular diseases, such as ischemic stroke. Stroke is among the leading causes of death

globally and its prevalence as a major health concern is predicted to increase, as the global

population ages. Additionally, the demographics of populations change, for example there is a

rise in the number of people affected by hypertension, diabetes, and obesity, all increasing risk

factors for ischemic stroke[55–57]. Furthermore, in stroke survivors there is an increase in disability,

which leads to a reduction in the quality of life for a patient [58–60]. One of the many reasons these

problems exist is that the majority of preclinical studies are targeted only towards male subjects

[61]. Over 90% of preclinical studies use strictly male mice whereas all clinical studies use equal

part male and female participants [61,62(p201)]. This makes clinical pharmaceutical findings favor

better outcomes in males [63,64]. To address the gap in the literature, we investigated the impact of

perinatal maternal nutritional deficiencies in folic acid or choline on cerebral blood flow velocity

after ischemic stroke in young adult and middle-age female offspring. Since cerebral vasculature

blood flow is also directly affected by changes in cardiac and peripheral vascular function and

structure [65–67], we have further investigated the biophysical properties of the heart and aorta in our

experimental groups.

Methods

Experimental Design

All methods were performed in accordance with the relevant guidelines and regulations. All

animal experimentation was performed following approval by the Midwestern University

Institutional Animal Care and Use Committee (Protocol 2983) in accordance with animal welfare

and ARRIVE guidelines.

Experimental conditions are summarized in Figure 2. Briefly, for the maternal cohort (n = 30)

female and (n = 30) male C57BL/6J mice (RRID: IMSR_JAX:000664, Jackson Laboratories)

were purchased from Jackson Laboratories. The mice were acclimatized for one week to

controlled housing conditions (22 ± 1°C, 12h-light/12h-dark cycle) with *ad libitum* access to food

and water. At two months of age (Day 0), females were randomized to control (CD, TD.190790),

and commercially (Envigo) prepared folic acid (FADD, TD.01546) or choline (ChDD,

TD.06119) deficient diets and maintained on these diets for four weeks prior to mating, and later

throughout pregnancy and lactation (Figure 1). Levels of folic acid, choline bitartrate, and

macronutrients in experimental diets are listed in in Table 1 [39,53,54,68].

At three weeks of age newly born pubs were weaned, and all female offspring were maintained

on the CD *ad libitum* for the rest of the experimental timeline. Offspring were randomized to one

of two cohorts undergoing photothrombotic (PT) stroke at either 2- or 10-months of age, followed

by ultrasound measurements at 1.5 months post-stroke.

Photothrombosis

When female offspring reached 2 or 10 months of age, ischemia was induced using the

photothrombosis model. They were anesthetized with isoflurane (1.5%) in a 70:30 nitrous

oxide:oxygen mixture. Core body temperature was monitored with a rectal thermometer (catalog

number, 521591, Harvard Apparatus) and maintained at 37 ± 0.2 ºC using a heating blanket. 10

mg/kg of the photosensitive Rose Bengal dye (catalog number, 198250, Sigma-Aldrich) was

injected intraperitoneally 5 minutes prior to irradiation. A 532 nm green laser (MGM20 (20-

25mW, Beta Electronics) was placed 3 cm above the animal and directed to the sensorimotor

cortex (mediolateral + 0.24mm) for 15 minutes [53,69–71]. At the completion of the study brain

tissue was collected, sectioned and stained to confirm ischemic damage [53,69–74].

Ultrasound imaging

Approximately 1.5 months after ischemic stroke, *in vivo* measurements of posterior cerebral

artery (PCA) and aortic function were performed using Vevo® 2100 ultrasound imaging system

(FUJIFILM, VisualSonics, Toronto, Canada). Measurements were completed in random order by

investigators blinded to treatment groups. The high-frequency, high-resolution ultrasound system

used in this study was equipped with a 40 MHz transducer (MS550S) with a focal length of 7.0

mm, frame rate of 557 fps (single zone, 5.08 mm width, B-mode), and a maximum two-

dimensional field of view of 14.1×15.0 mm with a spatial resolution of 90 µm lateral by 40 µm

axial.

As described in our previous published reports[75–77], Mice were anesthetized in an induction

chamber with 3% isoflurane and 1 L/min flow of 100% oxygen for 1–2 mins, then placed supine

on a heated platform and maintained with 1.5–2% isoflurane. Heart rate, electrocardiogram

(ECG), and respiratory rate were measured by the four ECG electrodes embedded in the platform.

Using a heat lamp and heated platform, body temperature was maintained at 36–38°C and

monitored by a rectal probe throughout. The left ventricular (LV) structural and functional

parameters, including stroke volume, ejection fraction, fractional shortening, and cardiac output,

were calculated from the LV parasternal short-axis M-mode view and recorded at the level of two

papillary muscles. An M-mode cursor was positioned perpendicular to the anterior and posterior

walls in the middle of the LV for measuring wall thickness. Interventricular septal wall (IVS)

thickness during diastole (IVSd) and systole (IVSs) were also obtained from LV parasternal long-

axis M-mode view.

Aortic diameters at the annulus, sinuses of Valsalva, and sinotubular junction were measured

from the B-mode aortic arch view. Ascending and descending aortic, and posterior cerebral artery

(PCA) peak velocities were measured from the pulse wave (PW) Doppler-mode. Aortic pulse

wave velocity (PWV) was obtained from the B-mode and Doppler-mode aortic arch view,

calculated as PWV (mm/s) = aortic arch distance (d2-d1)/transit time (T1-T2). The PW Doppler

mode sample volume was placed in the ascending aorta to verify the time from the onset of the

QRS complex to the onset of the ascending aortic Doppler waveform (T1). Using the same image

plane, the time from the onset of the QRS complex to the onset of the descending aortic Doppler

waveform (T2) was also measured, and the average values for T1 and T2 over 10 cardiac cycles

were calculated. Furthermore, the aortic arch distance was measured between the two sample

volume positions along the central axis of aortic arch on the B-mode image.

Transcranial Doppler sonography is a non-invasive, non-ionizing, inexpensive, portable, and safe

technique that uses a pulsed Doppler transducer for assessment of intracerebral blood flow in the

clinical practice [78,79] and has become an important translational tool to evaluate the intracerebral

blood flow in animal models. The PCA peak blood flow was measured using the 24MHz

(MS250) transducer. The trans occipital window was used to visualize the posterior cerebral

arteries and pulsed wave (PW), Doppler-mode was used to measure the PCA peak blood flow

velocity.

Statistics

Ultrasound data was analyzed by two individuals that were blinded to experimental treatment

groups using Vevo Lab ultrasound analysis software (FUJIFILM, VisualSonics, Toronto,

Canada). Using GraphPad Prism 9.0, two-way ANOVA analysis was performed to assess

maternal diet and age effects. Significant main effects of two-way ANOVAs were followed up

with Tukey's HSD post-hoc test to adjust for multiple comparisons. All data are presented as

mean $\pm$ standard error of the mean (SEM). Statistical tests were performed using a significance

level $p < 0.05$.

Results

Ischemic damage in brain tissue

To confirm that mice had a ischemic stroke, brain tissue was collected from all animals and

damage was confirmed by staining sectioned tissue and quantifying damage volume using

microscopy (data not shown) [80].

Cerebral blood flow

After ischemic stroke was induced, approximately 1.5 months, cerebral blood flow was measured

in posterior cerebral artery using ultrasound. Representation of posterior cerebral artery is shown

in Figure 1A. Maternal diet impacted blood flow velocity within the posterior cerebral artery

(Figure 1B; $F_{(2,28)} = 5.47$, $p = 0.009$). There was significant decrease cerebral blood flow in 11.5-

month-old offspring from ChDD mothers ($p < 0.05$). There was no effect of offspring age ($F_{(1,28)}$

$= 0.82$, $p = 0.37$.) and no interaction between both factors ($F_{(2,28)} = 1.25$, $p = 0.30$).

Measurements of cardiac function and structure

Cardiac function and structure were evaluated at 1.5 months timepoint after ischemic stroke using

ultrasound. Both maternal diet ($F_{(2,28)} = 4.16$, $p = 0.03$) and offspring age ($F_{(1,28)} = 13.55$, $p =$

0.001) impacted average heart rate (Table 2). Aged (11.5-month-old) offspring from CD ($p <$

0.01) FADD ($p < 0.01$) and ChDD ($p < 0.01$) mothers had higher heart rates compared to 3.5-

month-old controls (Table 3). While exclusively offspring age affected ejection fraction ($F_{(1,29)} =$

7.79, $p = 0.01$) and cardiac output ($F_{(1,28)} = 35.66$, $p < 0.0001$), there were no pairwise differences

(Table 3). There were interactions between maternal diet and offspring age for stroke volume

($F_{(2,29)} = 4.05$, $p = 0.03$), there was a difference between 3.5- and 11.5-month old FADD

offspring (Table 3). There were no differences between groups for fractional shortening and

internal diameter in systolic and diastolic measurements.

Aortic pulse wave velocity

Aortic pulse wave velocity is a measure of aortic wall stiffness and was evaluated 1.5 months

after ischemic stroke using ultrasound. Offspring age impacted pulse wave velocity (Figure 3A, F

$(_{1,25}) = 33.70$, $p < 0.0001$), there were differences between 3.5-and 11.5-month-old CD ($p < 0.01$)

and ChDD ($p < 0.01$) offspring. There was no impact of maternal diet ($F_{(2,25)} = 0.21$, $p = 0.81$) or

interaction between maternal diet and offspring age ($F_{(2,25)} = 2.05$, $p = 0.15$).

Measurements of coronary artery function

Coronary artery function was measured using ultrasound 1.5 months after ischemic stroke using

ultrasound. Significant interaction between maternal diet and age for coronary artery velocity

ratio (Figure 3B, $F_{(2,28)} = 4.28$, $p = 0.02$). There was no impact of material diet ($F_{(2,28)} = 0.10$, $p =$

0.38) or age ($F_{(1,28)} = 0.004$, $p = 0.95$).

Discussion

The Developmental Origins of Health and Disease (DOHaD) theory suggests that prospective

chronic diseases are programmed *in utero*- giving rise to programming of offspring

cardiovascular, metabolic, and neuroendocrine dysfunction [8–11]. Despite impressive evidence of

the importance of the maternal environment for fetal growth and development *in utero*, there is

not much information available on the impact on offspring as they age, and particularly under

neurovascular complications, such as ischemic stroke. Nutrition is a modifiable risk factor for

ischemic tissue [81] and can also impact outcome after ischemic damage[53,69–73,80,82]. Using an

experimental model of ischemic stroke, our study aimed to determine the impact of perinatal

maternal dietary deficiencies in folic acid and choline on measures of cerebral, cardiac, aorta, and

coronary in offspring, following ischemic injury. Our results demonstrate that maternal diet

impact a cerebral blood in offspring after an ischemic stroke in female offspring. The effects of

both maternal diet and offspring age, as well as interactions between these variables were

observed for numerous cardiac, aortic, and coronary artery function after ischemic stroke in

female offspring.

The neurovascular unit (NVU) is comprised of a number of unique neuronal, glial, and

endothelial cell types, and recent findings indicate unique cross-talk between neurons and the

cerebral vasculature [83–86], emphasizing the complex, pivotal role the NVU plays during

development and in the progression of neurovascular pathologies like ischemic stroke and

cerebral aneurysm [87–92]. Further, the NVU is responsible for the maintenance of a highly selective

blood–brain barrier (BBB) and cerebral homeostasis, as well as the control of cerebral blood flow

(CBF) [93]. The impact of maternal diet on the NVU, modulating integrity of cerebral blood vessels

and closure of the neural tube, has been established. Our study adds to these investigations by

assessing the response of blood flow within the posterior cerebral artery (PCA) in both young and

aged offspring. The contra-lesional PCA was selected as an index of cerebral blood flow due to

its spatial and functional independence from the sensorimotor cortex targeted during

photothrombotic stroke [94], and evident correlation to measures of the Middle Cerebral Artery

(MCA) [95].

In line with studies detailing the impact of maternal choline on neurovascular development [16], our

results suggest that maternal choline levels during pregnancy and lactation impair cerebral blood

flow in young mice following ischemic stroke. In rodent models, the importance of choline for

optimal neurodevelopment is well-established [96,97]. Recent work has examined the role of choline

in neurovascular interactions as well, modulating levels of anti-angiogenic factors during

gestation [98], fetal hippocampal angiogenesis [37], and promoting the proliferation of rat endothelial

cells following hypoxic injury in cerebral vessels [99]. In this way, choline has been shown to

influence neurovascular health across the lifespan and may be implicated in both neurovascular

structure and functional response to injury.

The cardiovascular system has also demonstrated effects of choline deficiency, including heart

defects [100,101], while higher intake of choline was associated with reduced risk of adult

cardiovascular disease [102] and amelioration of impaired vagal activity and inflammation in

hypertensive rodents [103]. Our study revealed a diet effect of both maternal choline and folic acid,

where deficiencies in either nutrient significantly increased offspring heart rate, regardless of

offspring age. While heart rate is a well-known risk factor for cardiovascular disease, our results

align well with recent findings associating low heart rate with better functional and cognitive

outcomes following ischemic stroke [104] and high heart rate with impaired endothelial function and

increased ischemic lesion size following stroke [105], as well as death due to vascular diseases [104].

Overall, maternal diet has an established developmental influence on basic measures of

cardiovascular and neurovascular health, and may impact offspring programming of the NVU,

thereby influencing stroke outcome via endothelial homeostasis via endothelial NO synthase

(eNOS) [106,107].

We did not observe the effect of maternal diet on cerebral blood flow in 11.5-month-old

offspring. This could be due to the well-investigated aging-associated changes in the structural

and functional integrity of the vasculature [65,108–111]. Therefore, we propose that the difference in

effect between young (3.5-month-old) and old (11.5-month-old) offspring is a result of the aging

of the control mice. In addition, the presumed damage or endothelial dysfunction induced by the

maternal choline and folic acid deficient diets is long-lasting and may contribute to premature

aging, generating a mathematically significant difference when compared to young, healthy

controls, but only a minor difference when compared to senescent offspring displaying similar

levels of vascular dysfunction. This result is supported by the interaction effect of diet and

offspring age and requires further investigation. In addition, unique mechanisms drive vascular

senescence in males and females [112], so our results may be obscured by our study of exclusively

female mice. Another interesting result from our study is the significance difference of the

coronary artery velocity (S/D) ratio between 3.5 and 11.5-month-old and maternal diet offspring.

This result may indicate cardiovascular impairment related to a maternal diet deficient in choline

and control diets. However, because the incidence of coronary artery disease (CAD), valvular

disease, rhythm disorders, and heart failure increases with age[113], it appears that folic acid may

play a role in programming resistance to this age-related dysfunction.

Outside of heart rate, which is discussed above, age effects were observed for offspring ejection

fraction, cardiac output, and pulse wave velocity. Ejection fraction, an index of the left ventricular

output, has recently been used as a measure of cardiac mortality risk[114], with lower percentages

indicating cardiac dysfunction. In our study, older mice displayed a significantly reduced ejection

fraction, indicating cardiovascular impairment. In a similar manner, our cardiac output data

indicate the expected increased cardiac dysfunction as a product of aging[115]. Aortic pulse wave

velocity (PWV) was also found to be significantly increased on the older cohort, in line with

clinical findings[116]. Controlling cardiovascular risk factors, such as hypertension can decrease risk

for ischemic stroke [117].

The present study has demonstrated novel data in the field of maternal dietary deficiencies and

blood flow after ischemic stroke in offspring. Our data has demonstrated that there are changes in

vasculature of female offspring after ischemic stroke, because of maternal diet, this is the first

study of its kind. Furthermore, our data points to the need for rodent models spanning a variety of

ages for research in age-related diseases such as stroke and vascular dysfunction. We recognize

that the exclusion of male subjects in this study may limit our ability to draw conclusions with

respect to the impact of sex hormone in observed phenomenon. In future studies, we plan to

include male animals, and design experiments that would also allow us to investigate the role of

paternal dietary effects. Additionally, we plan to further age animals to ~20mo after ischemic

stroke as well as investigate the role of over supplementation on blood flow after stroke. A

detailed analysis of angiogenesis after ischemic stroke might also be prudent.

In conclusion, maternal dietary deficiencies in folic acid and choline have unique roles in cerebral

and peripheral blood flow after ischemic stroke in female offspring. Maternal nutrition during

pregnancy and lactation has effects, even after infancy and childhood. Our work demonstrated an

age effect in animal models encourages further comprehensive longitudinal time-point studies

that includes older age animals.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

Funding

Grants awarded to Mitra Esfandiarei; R15HL145646 (NIH) and Nafisa M. Jadavji;
20AIREA35050015 (American Heart Association).

Acknowledgments

Figure 2A was created with BioRender.com

References

1. Gernand AD, Schulze KJ, Stewart CP, West KP, Christian P. Micronutrient deficiencies in pregnancy worldwide: health effects and prevention. *Nat Rev Endocrinol* 2016;12:274–289.

2. Badon SE, Miller RS, Qiu C, Sorensen TK, Williams MA, Enquobahrie DA. Maternal healthy lifestyle during early pregnancy and offspring birthweight: differences by offspring sex. *J Matern Fetal Neonatal Med* 2018;31:1111–1117.

3. Lu M-S, Chen Q-Z, He J-R, Wei X-L, Lu J-H, Li S-H, et al. Maternal Dietary Patterns and Fetal Growth: A Large Prospective Cohort Study in China. *Nutrients* 2016;8:E257.

4. Ae-Ngibise KA, Wylie BJ, Boamah-Kaali E, Jack DW, Oppong FB, Chillrud SN, et al. Prenatal maternal stress and birth outcomes in rural Ghana: sex-specific associations. *BMC Pregnancy Childbirth* 2019;19:391.

5. Borge TC, Aase H, Brantsæter AL, Biele G. The importance of maternal diet quality during pregnancy on cognitive and behavioural outcomes in children: a systematic review and meta-analysis. *BMJ Open* 2017;7:e016777.

6. Warner MJ, Ozanne SE. Mechanisms involved in the developmental programming of adulthood disease. *Biochem J* 2010;427:333–347.

7. Steegers-Theunissen RP, Obermann-Borst SA, Kremer D, Lindemans J, Siebel C, Steegers EA, et al. Periconceptional Maternal Folic Acid Use of 400 µg per Day Is Related to Increased Methylation of the IGF2 Gene in the Very Young Child. *PLOS ONE* 2009;4:e7845.

8. Barker DJP, Osmond C. INFANT MORTALITY, CHILDHOOD NUTRITION, AND ISCHAEMIC HEART DISEASE IN ENGLAND AND WALES. *The Lancet* 1986;327:1077–1081.

9. Barker DJP, Osmond C, Winter PD, Margetts B, Simmonds SJ. WEIGHT IN INFANCY AND DEATH FROM ISCHAEMIC HEART DISEASE. *The Lancet* 1989;334:577–580.

10. Barker DJP, Godfrey KM, Gluckman PD, Harding JE, Owens JA, Robinson JS. Fetal nutrition and cardiovascular disease in adult life. *The Lancet* 1993;341:938–941.

11. Barker D. Intrauterine programming of coronary heart disease and stroke. *Acta Paediatrica* 1997;86:178–182.

12. Christian P, Stewart CP. Maternal micronutrient deficiency, fetal development, and the risk of chronic disease. *J Nutr* 2010;140:437–445.

13. Song X, Wei J, Shu J, Liu Y, Sun M, Zhu P, et al. Association of polymorphisms of FOLR1 gene and FOLR2 gene and maternal folic acid supplementation with risk of ventricular septal defect: a case-control study. *Eur J Clin Nutr* https://doi.org/10.1038/s41430-022-01110-9.

14. Ravelli A, van der Meulen J, Michels R, Osmond C, Barker D, Hales C, et al. Glucose tolerance in adults after prenatal exposure to famine. *The Lancet* 1998;351:173–177.

15.   Pitkin RM. Folate and neural tube defects. *Am J Clin Nutr* 2007;85:285S-288S.

16.   Shaw GM, Carmichael SL, Yang W, Selvin S, Schaffer DM. Periconceptional dietary
intake of choline and betaine and neural tube defects in offspring. *Am J Epidemiol*
2004;160:102–109.

17.   Nyaradi A, Li J, Hickling S, Foster J, Oddy WH. The role of nutrition in children's
neurocognitive development, from pregnancy through childhood. *Front Hum Neurosci*
2013;7:97.

18.   Wachs TD. Nutritional deficits and behavioural development. *International Journal of*
Behavioral Development 2000;24:435–441.

19.   Georgieff MK. Nutrition and the developing brain: nutrient priorities and measurement. *Am*
J Clin Nutr 2007;85:614S-620S.

20.   Georgieff MK, Ramel SE, Cusick SE. Nutritional Influences on Brain Development. *Acta*
Paediatr 2018;107:1310–1321.

21.   Caffrey A, McNulty H, Rollins M, Prasad G, Gaur P, Talcott JB, et al. Effects of maternal
folic acid supplementation during the second and third trimesters of pregnancy on
neurocognitive development in the child: an 11-year follow-up from a randomised
controlled trial. *BMC Med* 2021;19:73.

22.   Huang R-C, Mori TA, Beilin LJ. Early life programming of cardiometabolic disease in the
Western Australian pregnancy cohort (Raine) study. *Clin Exp Pharmacol Physiol*
2012;39:973–978.

23.   Heijmans BT, Tobi EW, Stein AD, Putter H, Blauw GJ, Susser ES, et al. Persistent
epigenetic differences associated with prenatal exposure to famine in humans. *Proceedings*
of the National Academy of Sciences 2008;105:17046–17049.

24.   Dominguez-Salas P, Cox SE, Prentice AM, Hennig BJ, Moore SE. Maternal nutritional
status, C1 metabolism and offspring DNA methylation: a review of current evidence in
human subjects. *Proc Nutr Soc* 2012;71:154–165.

25.   James P, Sajjadi S, Tomar AS, Saffari A, Fall CHD, Prentice AM, et al. Candidate genes
linking maternal nutrient exposure to offspring health via DNA methylation: a review of
existing evidence in humans with specific focus on one-carbon metabolism. *Int J*
Epidemiol 2018;47:1910–1937.

26.   Bokor S, Vass RA, Funke S, Ertl T, Molnár D. Epigenetic Effect of Maternal Methyl-
Group Donor Intake on Offspring's Health and Disease. *Life (Basel)* 2022;12:609.

27.   Peter CJ, Fischer LK, Kundakovic M, Garg P, Jakovcevski M, Dincer A, et al. DNA
METHYLATION SIGNATURES OF EARLY CHILDHOOD MALNUTRITION
ASSOCIATED WITH IMPAIRMENTS IN ATTENTION AND COGNITION. *Biol*
Psychiatry 2016;80:765–774.

28.   Guéant J-L, Namour F, Guéant-Rodriguez R-M, Daval J-L. Folate and fetal programming:
a play in epigenomics? *Trends in Endocrinology & Metabolism* 2013;24:279–289.

29.   Arima Y, Fukuoka H. Developmental origins of health and disease theory in cardiology. *J*
Cardiol 2020;76:14–17.

30.   Forsdahl A. Living conditions in childhood and subsequent development of risk factors for
426   arteriosclerotic heart disease. The cardiovascular survey in Finnmark 1974-75. *J Epidemiol*
427   *Community Health* 1978;32:34–37.

428   31.   Thornburg KL, O'Tierney PF, Louey S. Review: The placenta is a programming agent for
429   cardiovascular disease. *Placenta* 2010;31 Suppl:S54-59.

430   32.   Law CM, Shiell AW. Is blood pressure inversely related to birth weight? The strength of
431   evidence from a systematic review of the literature. *J Hypertens* 1996;14:935–941.

432   33.   McKee SE, Reyes TM. Effect of supplementation with methyl-donor nutrients on
433   neurodevelopment and cognition: considerations for future research. *Nutr Rev*
434   2018;76:497–511.

435   34.   Fernandez-Osornio LF, Gomez-Diaz RA, Mondragon-Gonzalez R, Gonzalez-Carranza E,
436   Diaz-Flores M, Sharma T, et al. Micronutrients of the one-carbon metabolism cycle are
437   altered in mothers and neonates by gestational diabetes and are associated with weight,
438   height and head circumference at birth. *The Journal of Nutritional Biochemistry*
439   2022;105:108996.

440   35.   Naninck EFG, Stijger PC, Brouwer-Brolsma EM. The Importance of Maternal Folate
441   Status for Brain Development and Function of Offspring. *Adv Nutr* 2019;10:502–519.

442   36.   Stamm RA, Houghton LA. Nutrient intake values for folate during pregnancy and lactation
443   vary widely around the world. *Nutrients* 2013;5:3920–3947.

444   37.   Mehedint MG, Craciunescu CN, Zeisel SH. Maternal dietary choline deficiency alters
445   angiogenesis in fetal mouse hippocampus. *Proc Natl Acad Sci U S A* 2010;107:12834–
446   12839.

447   38.   Deoni S, Dean D, Joelson S, O'Regan J, Schneider N. Early nutrition influences
448   developmental myelination and cognition in infants and young children. *Neuroimage*
449   2018;178:649–659.

450   39.   Jadavji NM, Deng L, Malysheva O, Caudill MA, Rozen R. MTHFR deficiency or reduced
451   intake of folate or choline in pregnant mice results in impaired short-term memory and
452   increased apoptosis in the hippocampus of wild-type offspring. *Neuroscience* 2015;300:1–
453   9.

454   40.   Schlotz W, Jones A, Phillips DIW, Gale CR, Robinson SM, Godfrey KM. Lower maternal
455   folate status in early pregnancy is associated with childhood hyperactivity and peer
456   problems in offspring. *J Child Psychol Psychiatry* 2010;51:594–602.

457   41.   Breimer LH, Nilsson TK. Has folate a role in the developing nervous system after birth and
458   not just during embryogenesis and gestation? *Scand J Clin Lab Invest* 2012;72:185–191.

459   42.   Li M, Francis E, Hinkle SN, Ajjarapu AS, Zhang C. Preconception and Prenatal Nutrition
460   and Neurodevelopmental Disorders: A Systematic Review and Meta-Analysis. *Nutrients*
461   2019;11:E1628.

462   43.   Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutr Rev*
463   2014;72:267–284.

464   44.   Antony AC. In utero physiology: role of folic acid in nutrient delivery and fetal
465   development. *Am J Clin Nutr* 2007;85:598S-603S.

466   45.   Craciunescu CN, Brown EC, Mar M-H, Albright CD, Nadeau MR, Zeisel SH. Folic Acid
467   Deficiency During Late Gestation Decreases Progenitor Cell Proliferation and Increases
468   Apoptosis in Fetal Mouse Brain. *The Journal of Nutrition* 2004;134:162–166.

469   46.   Akchiche N, Bossenmeyer-Pourié C, Kerek R, Martin N, Pourié G, Koziel V, et al.
470   Homocysteinylation of neuronal proteins contributes to folate deficiency-associated
471   alterations of differentiation, vesicular transport, and plasticity in hippocampal neuronal
472   cells. *The FASEB Journal* 2012;26:3980–3992.

473   47.   Harlan De Crescenzo A, Panoutsopoulos AA, Tat L, Schaaf Z, Racherla S, Henderson L, et
474   al. Deficient or Excess Folic Acid Supply During Pregnancy Alter Cortical
475   Neurodevelopment in Mouse Offspring. *Cerebral Cortex* 2021;31:635–649.

476   48.   Zhao Y, Chen D, Tang J, Zheng Y, Qi J, Wang H. Parental folate deficiency induces birth
477   defects in mice accompanied with increased de novo mutations. *Cell Discov* 2022;8:1–13.

478   49.   FREEDMAN R, ROSS RG. Prenatal choline and the development of schizophrenia.
479   *Shanghai Arch Psychiatry* 2015;27:90–102.

480   50.   Wang Y, Surzenko N, Friday WB, Zeisel SH. Maternal dietary intake of choline in mice
481   regulates development of the cerebral cortex in the offspring. *FASEB J* 2016;30:1566–
482   1578.

483   51.   Jones JP, H. Meck W, Williams CL, Wilson WA, Swartzwelder HS. Choline availability to
484   the developing rat fetus alters adult hippocampal long-term potentiation. *Developmental
485   Brain Research* 1999;118:159–167.

486   52.   Glenn MJ, Gibson EM, Kirby ED, Mellott TJ, Blusztajn JK, Williams CL. Prenatal choline
487   availability modulates hippocampal neurogenesis and neurogenic responses to enriching
488   experiences in adult female rats. *Eur J Neurosci* 2007;25:2473–2482.

489   53.   Jadavji NM, Emmerson JT, MacFarlane AJ, Willmore WG, Smith PD. B-vitamin and
490   choline supplementation increases neuroplasticity and recovery after stroke. *Neurobiol Dis*
491   2017;103:89–100.

492   54.   Jadavji NM, Farr TD, Lips J, Khalil AA, Boehm-Sturm P, Foddis M, et al. Elevated levels
493   of plasma homocysteine, deficiencies in dietary folic acid and uracil-DNA glycosylase
494   impair learning in a mouse model of vascular cognitive impairment. *Behav Brain Res*
495   2015;283:215–226.

496   55.   Ovbiagele B, Goldstein LB, Higashida RT, Howard VJ, Johnston SC, Khavjou OA, et al.
497   Forecasting the Future of Stroke in the United States: A Policy Statement From the
498   American Heart Association and American Stroke Association. *Stroke* 2013;44:2361–2375.

499   56.   Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart
500   Disease and Stroke Statistics—2016 Update.

501   57.   Cabral NL, Freire AT, Conforto AB, dos Santos N, Reis FI, Nagel V, et al. Increase of
502   Stroke Incidence in Young Adults in a Middle-Income Country. *Stroke* 2017;48:2925–
503   2930.

58.   Kissela BM, Khoury JC, Alwell K, Moomaw CJ, Woo D, Adeoye O, et al. Age at stroke:
Temporal trends in stroke incidence in a large, biracial population. *Neurology*
2012;79:1781–1787.

59.   Dhamoon MS, Moon YP, Paik MC, Boden-Albala B, Rundek T, Sacco RL, et al. Quality
of life declines after first ischemic stroke: The Northern Manhattan Study. *Neurology*
2010;75:328–334.

60.   Bako AT, Pan A, Potter T, Tannous J, Johnson C, Baig E, et al. Contemporary Trends in
the Nationwide Incidence of Primary Intracerebral Hemorrhage. *Stroke* 2022;53:e70–e74.

61.   Persky RW, Turtzo LC, McCullough LD. Stroke in Women: Disparities and Outcomes.
Curr Cardiol Rep 2010;12:6–13.

62.   the Women Initiative for Stroke in Europe (WISE) group, Cordonnier C, Sprigg N, Sandset
EC, Pavlovic A, Sunnerhagen KS, et al. Stroke in women — from evidence to inequalities.
Nat Rev Neurol 2017;13:521–532.

63.   Macrae I. Preclinical stroke research - advantages and disadvantages of the most common
rodent models of focal ischaemia: Animal models of focal cerebral ischaemia. *British*
Journal of Pharmacology 2011;164:1062–1078.

64.   Singh D, Wasan H, Reeta KH. Preclinical Stroke Research and Translational Failure: A
Bird's Eye View on Preventable Variables. *Cell Mol Neurobiol*
https://doi.org/10.1007/s10571-021-01083-6.

65.   Yabluchanskiy A, Nyul-Toth A, Csiszar A, Gulej R, Saunders D, Towner R, et al. Age-
related alterations in the cerebrovasculature affect neurovascular coupling and BOLD fMRI
responses: Insights from animal models of aging. *Psychophysiology* 2021;58:e13718.

66.   Ivey FM, Hafer-Macko CE, Ryan AS, Macko RF. Impaired Leg Vasodilatory Function
After Stroke. *Stroke* 2010;41:2913–2917.

67.   Wang SJ, Lee C-H, Kang HG, Kim KW, Kim M, Jeong H-J, et al. Peripheral vasoreactivity
in acute ischemic stroke with hemiplegia. *Scientific Reports* 2021;11.

68.   Dam K, Füchtemeier M, Farr TD, Boehm-Sturm P, Foddis M, Dirnagl U, et al. Increased
homocysteine levels impair reference memory and reduce cortical levels of acetylcholine in
a mouse model of vascular cognitive impairment. *Behav Brain Res* 2017;321:201–208.

69.   Jadavji NM, Emmerson JT, Shanmugalingam U, MacFarlane AJ, Willmore WG, Smith
PD. A genetic deficiency in folic acid metabolism impairs recovery after ischemic stroke.
Exp Neurol 2018;309:14–22.

70.   Jadavji NM, Mosnier H, Kelly E, Lawrence K, Cruickshank S, Stacey S, et al. One-carbon
metabolism supplementation improves outcome after stroke in aged male MTHFR-
deficient mice. *Neurobiol Dis* 2019;132:104613.

71.   Abato JE, Moftah M, Cron GO, Smith PD, Jadavji NM. Methylenetetrahydrofolate
reductase deficiency alters cellular response after ischemic stroke in male mice. *Nutritional*
Neuroscience https://doi.org/10.1080/1028415X.2020.1769412.

72. Poole J, Jasbi P, Pascual AS, North S, Kwatra N, Weissig V, et al. Reduced stroke outcome
in old-aged female mice maintained on a dietary vitamin B12 deficiency.
https://doi.org/10.1101/2022.04.04.487028.

73. Hurley L, Jauhal J, Ille S, Pull K, Malysheva OV, Jadavji NM. Maternal Dietary
Deficiencies in Folic Acid and Choline Result in Larger Damage Volume, Reduced Neuro-
Degeneration and -Inflammation and Changes in Choline Metabolites after Ischemic Stroke
in Middle-Aged Offspring. *Nutrients* 2023;15:1556.

74. Mbs GBY, Wasek B, Bottiglieri T, Malysheva O, Caudill MA, Jadavji NM. Dietary
vitamin B12 deficiency impairs motor function and changes neuronal survival and choline
metabolism after ischemic stroke in middle-aged male and female mice. *Nutr Neurosci*
https://doi.org/10.1080/1028415X.2023.2188639.

75. Esfandiarei M, Hoxha B, Talley NA, Anderson MR, Alkhouli MF, Squire MA, et al.
Beneficial effects of resveratrol and exercise training on cardiac and aortic function and
structure in the 3xTg mouse model of Alzheimer's disease. *Drug Des Devel Ther*
2019;13:1197–1211.

76. Kuechenmeister B, Gusek B, Jones TB, Esfandiarei M. Evaluation of the effects of Marfan
pathogenesis and losartan treatment on coronary and cerebral arteries in the well-
established Marfan Syndrome mouse model. *The FASEB Journal* 2021;35.

77. Cui JZ, Lee L, Sheng X, Chu F, Gibson CP, Aydinian T, et al. In vivo characterization of
doxycycline-mediated protection of aortic function and structure in a mouse model of
Marfan syndrome-associated aortic aneurysm. *Scientific Reports* 2019;9:2071–2071.

78. Purkayastha S, Sorond F. Transcranial Doppler Ultrasound: Technique and Application.
Semin Neurol 2012;32:411–420.

79. Aaslid R, Markwalder TM, Nornes H. Noninvasive transcranial Doppler ultrasound
recording of flow velocity in basal cerebral arteries. *J Neurosurg* 1982;57:769–774.

80. Clementson M, Hurley L, Coonrod S, Bennett C, Marella P, Pascual AS, et al. Maternal
dietary deficiencies in folic acid or choline worsen stroke outcomes in adult male and
female mouse offspring. *Neural Regen Res* 2023;18:2443–2448.

81. Hankey GJ. Nutrition and the risk of stroke. *The Lancet Neurology* 2012;11:66–81.

82. Ahmed S, Bogiatzi C, Hackam DG, Rutledge AC, Sposato LA, Khaw A, et al. Vitamin B
12 deficiency and hyperhomocysteinaemia in outpatients with stroke or transient ischaemic
attack: A cohort study at an academic medical centre. *BMJ Open* 2019;9:1–8.

83. Whiteus C, Freitas C, Grutzendler J. Perturbed neural activity disrupts cerebral
angiogenesis during a postnatal critical period. *Nature* 2014;505:407–411.

84. Peguera B, Segarra M, Acker-Palmer A. Neurovascular crosstalk coordinates the central
nervous system development. *Curr Opin Neurobiol* 2021;69:202–213.

85. Paredes I, Himmels P, Ruiz de Almodóvar C. Neurovascular Communication during CNS
Development. *Dev Cell* 2018;45:10–32.

86. Tata M, Ruhrberg C. Cross-talk between blood vessels and neural progenitors in the
developing brain. *Neuronal Signal* 2018;2:NS20170139.

87. Wang L, Xiong X, Zhang L, Shen J. Neurovascular Unit: A critical role in ischemic stroke.
CNS Neuroscience & Therapeutics 2021;27:7–16.

88. Lecrux C, Hamel E. The neurovascular unit in brain function and disease. *Acta Physiol*
(Oxf) 2011;203:47–59.

89. Guan J, Pavlovic D, Dalkie N, Waldvogel HJ, O'Carroll SJ, Green CR, et al. Vascular
Degeneration in Parkinson's Disease. *Brain Pathol* 2012;23:154–164.

90. Murphy MJ, Grace GM, Tartaglia MC, Orange JB, Chen X, Rowe A, et al. Widespread
cerebral haemodynamics disturbances occur early in amyotrophic lateral sclerosis.
Amyotroph Lateral Scler 2012;13:202–209.

91. Kisler K, Nelson AR, Montagne A, Zlokovic BV. Cerebral blood flow regulation and
neurovascular dysfunction in Alzheimer's disease. *Nat Rev Neurosci* 2017;18:419–434.

92. Toledo JB, Arnold SE, Raible K, Brettschneider J, Xie SX, Grossman M, et al.
Contribution of cerebrovascular disease in autopsy confirmed neurodegenerative disease
cases in the National Alzheimer's Coordinating Centre. *Brain* 2013;136:2697–2706.

93. Muoio V, Persson PB, Sendeski MM. The neurovascular unit – concept review. *Acta*
Physiologica 2014;210:790–798.

94. Xiong B, Li A, Lou Y, Chen S, Long B, Peng J, et al. Precise Cerebral Vascular Atlas in
Stereotaxic Coordinates of Whole Mouse Brain. *Frontiers in Neuroanatomy* 2017;11.

95. Haubrich C, Wendt A, Diehl R r., Klötzsch C. Dynamic Autoregulation Testing in the
Posterior Cerebral Artery. *Stroke* 2004;35:848–852.

96. Zeisel SH. Nutritional importance of choline for brain development. *J Am Coll Nutr*
2004;23:621S-626S.

97. Radziejewska A, Chmurzynska A. Folate and choline absorption and uptake: Their role in
fetal development. *Biochimie* 2019;158:10–19.

98. Jiang X, Bar HY, Yan J, Jones S, Brannon PM, West AA, et al. A higher maternal choline
intake among third-trimester pregnant women lowers placental and circulating
concentrations of the antiangiogenic factor fms-like tyrosine kinase-1 (sFLT1). *The FASEB*
Journal 2013;27:1245–1253.

99. Zhang L-C, Jin X, Huang Z, Yan Z-N, Li P-B, Duan R-F, et al. Protective effects of
choline against hypoxia-induced injuries of vessels and endothelial cells. *Exp Ther Med*
2017;13:2316–2324.

100. Chan J, Deng L, Mikael LG, Yan J, Pickell L, Wu Q, et al. Low dietary choline and low
dietary riboflavin during pregnancy influence reproductive outcomes and heart
development in mice. *The American Journal of Clinical Nutrition* 2010;91:1035–1043.

101. Sotres-Alvarez D, Siega-Riz AM, Herring AH, Carmichael SL, Feldkamp ML, Hobbs CA,
et al. Maternal Dietary Patterns are Associated With Risk of Neural Tube and Congenital
Heart Defects. *American Journal of Epidemiology* 2013;177:1279–1288.

102. Golzarand M, Mirmiran P, Azizi F. Association between dietary choline and betaine intake
and 10.6-year cardiovascular disease in adults. *Nutrition Journal* 2022;21:1.

103. Liu L, Lu Y, Bi X, Xu M, Yu X, Xue R, et al. Choline ameliorates cardiovascular damage
by improving vagal activity and inhibiting the inflammatory response in spontaneously
hypertensive rats. *Sci Rep* 2017;7:42553.

104. Bohm M, Cotton D, Foster L, Custodis F, Laufs U, Sacco R, et al. Impact of resting heart
rate on mortality, disability and cognitive decline in patients after ischaemic stroke.
European Heart Journal 2012;33:2804–2812.

105. Custodis F, Gertz K, Balkaya M, Prinz V, Mathar I, Stamm C, et al. Heart Rate Contributes
to the Vascular Effects of Chronic Mental Stress. *Stroke* 2011;42:1742–1749.

106. Laufs U, Gertz K, Dirnagl U, Böhm M, Nickenig G, Endres M. Rosuvastatin, a new HMG-
CoA reductase inhibitor, upregulates endothelial nitric oxide synthase and protects from
ischemic stroke in mice. *Brain Research* 2002;942:23–30.

107. Bolduc V, Drouin A, Gillis M-A, Duquette N, Thorin-Trescases N, Frayne-Robillard I, et
al. Heart rate-associated mechanical stress impairs carotid but not cerebral artery
compliance in dyslipidemic atherosclerotic mice. *American Journal of Physiology-Heart
and Circulatory Physiology* 2011;301:H2081–H2092.

108. Hakim MA, Chum PP, Buchholz JN, Behringer EJ. Aging Alters Cerebrovascular
Endothelial GPCR and K+ Channel Function: Divergent Role of Biological Sex. *J
Gerontol A Biol Sci Med Sci* 2020;75:2064–2073.

109. Mandalà M, Pedatella AL, Morales Palomares S, Cipolla MJ, Osol G. Maturation is
associated with changes in rat cerebral artery structure, biomechanical properties and tone.
Acta Physiol (Oxf) 2012;205:363–371.

110. Caporarello N, Meridew JA, Aravamudhan A, Jones DL, Austin SA, Pham TX, et al.
Vascular dysfunction in aged mice contributes to persistent lung fibrosis. *Aging Cell*
2020;19:e13196.

111. Donato AJ, Machin DR, Lesniewski LA. Mechanisms of Dysfunction in the Aging
Vasculature and Role in Age-Related Disease. *Circulation Research* 2018;123:825–848.

112. Barros PR, Costa TJ, Akamine EH, Tostes RC. Vascular Aging in Rodent Models:
Contrasting Mechanisms Driving the Female and Male Vascular Senescence. *Front Aging*
2021;2:727604.

113. Global, regional, and national incidence, prevalence, and years lived with disability for 354
diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for
the Global Burden of Disease Study 2017. *Lancet* 2018;392:1789–1858.

114. Kosaraju A, Goyal A, Grigorova Y, Makaryus AN. Left Ventricular Ejection Fraction.
StatPearls. Treasure Island (FL): StatPearls Publishing, 2022.

115. Becklake MR, Frank H, Dagenais GR, Ostiguy GL, Guzman CA. Influence of age and sex
on exercise cardiac output. *Journal of Applied Physiology* 1965;20:938–947.

116. McEniery CM, Yasmin, Hall IR, Qasem A, Wilkinson IB, Cockcroft JR. Normal Vascular
Aging: Differential Effects on Wave Reflection and Aortic Pulse Wave Velocity: The
Anglo-Cardiff Collaborative Trial (ACCT). *Journal of the American College of Cardiology*
2005;46:1753–1760.

117. Arboix A. Cardiovascular risk factors for acute stroke: Risk profiles in the different
subtypes of ischemic stroke. *World J Clin Cases* 2015;3:418–429.

Tables

Table 1. List of micro- and macro-nutrient contents in Envigo control (TD.01369) folic acid (TD.01546), choline (TD.06119) deficient diets. The Envigo folic acid deficient (TD.95247) was identical to the control diet (TD.04194) except it contained 0.2 mg/kg folic acid. The Envigo choline deficient (TD.06119) was identical to the control diet (TD.01369) except it contained 0.3 mg/kg choline bitartrate.

Category	Concentration	Category	Concentration	Category	Concentration	Category	Concentration
Amino acids	**(g/kg)**	**Lipids**	**(g/kg)**	**Fat-soluble Vitamins**	**(IU/kg)**	**Nutrient**	**(mg/kg)**
Lysine	14	Total fat	60.0	Vitamin A	4000	Choline Bitartrate	1150 (300)
Methionine	4.7	Saturated fat	9.0	Vitamin D	1000	**Fat-soluble Vitamins**	**(IU/kg)**
Cystine	3.6	Monounsaturated fat	13.8	Vitamin E	75	Vitamin A	4000
Arginine	6.6	Polyunsaturated fat	36.6	Vitamin K	0.8	Vitamin D	1000
Phenylalanine	8.8	4:0 Butyric acid	0.0	**Minerals**	**(mg/kg)**	Vitamin E	75
Tyrosine	9.2	6:0 Caproic Acid	0.0	Calcium	5100	Vitamin K	0.8
Histidine	5.1	8:0 Caprylic Acid	0.0	Phosphorus	3000	**Water-soluble Vitamins**	**(mg/kg)**
Isoleucine	10.1	10:0 Capric acid	0.0	Potassium	3600	Biotin	0.20
Leucine	16	12:0 Lauric acid	0.0	Sodium	1000	Choline	1147.5
Threonine	7.6	14:0 Myristic Acid	0.0	Chlorine	1600	Folic acid	2.0 (0.3)
Tryptophan	2.1	16:0 Palmitic acid	6.6	Magnesium	515	Niacin	30.0
Valine	11.7	16:1 Palmitoleic acid	0.0	Copper	6.0	Pantothenate	14.7
Aspartic Acid	12.1	18:0 Stearic acid	2.4	Iron	36.5	Riboflavin	6.0
Glutamic acid	36.6	18:1 Oleic acid	13.8	Zinc	35.6	Thiamin	4.9
Alanine	5.2	18:2 Linoleic acid	31.8	Manganese	10.5	Vitamin B_6	5.8

Glycine	3.2	18:3 Linolenic acid	4.8	Iodine	0.21	Vitamin B$_{12}$	0.03
Proline	18.1	Cholesterol	8.8 mg/kg	Selenium	0.15	Vitamin C	0.0
Serine	10			Molybdenum	0.15		
				Chromium	1.00		

Table 2. Descriptive statistics (Mean ± SEM) and *p*-values results from two-way ANOVA analysis of cerebral and peripheral blood flow in 3.5 and 11.5-month-old female mouse offspring. Maternal diets included a control diet (CD), a folic acid deficient diet (FADD), and a choline deficient diet (ChDD). Mean ± SEM of 5 to 7 mice per group. * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001 significant diet, age or interaction (diet and age).

		Offspring Age		*p-values*		
Measurement	**Maternal Diet**	**3.5 months**	**11.5 months**	**Diet**	**Age**	**Diet x Age**
Average Heart Rate (BPM)	CD	494.3±13.23	524.78 ± 8.82			
	FADD	504.5 ±12.68	531.42 ± 4.02	0.0261*	0.0010**	0.1374
	ChDD	513.0 ± 9.55	531.30 ± 6.02			
Average Stroke Volume (μL)	CD	24.04 ± 2.54	23.55 ± 4.55			
	FADD	28.67 ± 2.17	37.47 ± 5.56	0.2744	0.2319	0.0282*
	ChDD	25.26 ± 2.06	38.24 ± 2.08			
Average Ejection Fraction (%)	CD	86.58 ± 4.24	84.88 ± 3.55			
	FADD	84.68 ± 3.09	73.10 ± 4.61	0.1638	0.009*	0.2929
	ChDD	86.46 ± 2.30	74.05 ± 2.49			
Average Fractional Shortening (%)	CD	57.94 ± 6.28	53.65 ± 4.18			
	FADD	53.89 ± 3.67	53.92 ± 3.88	0.3924	0.1811	0.4998
	ChDD	54.89 ± 3.13	43.70 ± 4.04			
Average Cardiac Output (mL/min	CD	11.77 ± 1.04	20.32 ± 0.78			
	FADD	14.43 ± 1.08	19.91 ± 2.94	0.7197	<0.0001***	0.5400
	ChDD	12.94 ± 1.03	20.28 ± 0.96			
Left Internal Diameter in Systolic (IVSs) (mm)	CD	0.90 ± 0.08	0.90 ± 0.10			
	FADD	1.13 ± 0.10	0.91 ± 0.05	0.2428	0.4138	0.3333
	ChDD	0.95 ± 0.09	0.99 ± 0.12			
Left Internal Diameter in Diastolic (IVSd)(mm)	CD	1.37 ± 0.10	1.56 ± 0.12			
	FADD	1.71 ± 0.04	1.44 ± 0.18	0.08098	0.5384	0.4547
	ChDD	1.61 ± 0.15	1.53 ± 0.12			

Table 3. Tukey's HSD post hoc pairwise analysis of cardiac function in 3.5 and 11.5-month-old
female mouse offspring. Maternal diets included a control diet (CD), a folic acid deficient diet
(FADD), and a choline deficient diet (ChDD). Presented are the p-values of 5 to 7 mice per
group. * $p < 0.05$, ** $p < 0.01$, post-hoc pairwise comparison.

	Comparison				
Measurement	Maternal Diet	Offspring Age	Maternal Diet	Offspring Age	*p-value*
Average Heart Rate (BPM)	CD	3.5	CD	11.5	0.0061**
	CD	3.5	FADD	11.5	0.0027**
	CD	3.5	ChDD	11.5	0.0028**
	FADD	3.5	CD	11.5	0.9755
	FADD	3.5	FADD	11.5	0.8215
	FADD	3.5	ChDD	11.5	0.8263
	ChDD	3.5	CD	11.5	0.9290
	ChDD	3.5	FADD	11.5	0.7478
	ChDD	3.5	ChDD	11.5	0.7525
Average Stroke Volume (µL)	CD	3.5	CD	11.5	0.9985
	CD	3.5	FADD	11.5	0.2006
	CD	3.5	ChDD	11.5	0.9997
	FADD	3.5	CD	11.5	>0.9999
	FADD	3.5	FADD	11.5	0.0426*
	FADD	3.5	ChDD	11.5	0.9331
	ChDD	3.5	CD	11.5	0.8983
	ChDD	3.5	FADD	11.5	0.6806
	ChDD	3.5	ChDD	11.5	0.9978
	CD	3.5	CD	11.5	0.9985
Average Ejection Fraction (%)	CD	3.5	CD	11.5	0.9993
	CD	3.5	FADD	11.5	0.1809
	CD	3.5	ChDD	11.5	0.1476
	FADD	3.5	CD	11.5	>0.9999
	FADD	3.5	FADD	11.5	0.2741
	FADD	3.5	ChDD	11.5	0.2275
	ChDD	3.5	CD	11.5	0.9997
	ChDD	3.5	FADD	11.5	0.2825
	ChDD	3.5	ChDD	11.5	0.2493
Average Cardiac Output (mL/min)	CD	3.5	CD	11.5	0.0020*
	CD	3.5	FADD	11.5	0.0057*
	CD	3.5	ChDD	11.5	0.0036*
	FADD	3.5	CD	11.5	0.0355*
	FADD	3.5	FADD	11.5	0.0825
	FADD	3.5	ChDD	11.5	0.0543

ChDD	3.5	CD	11.5	0.0245*
ChDD	3.5	FADD	11.5	0.0499*
ChDD	3.5	ChDD	11.5	0.0343*

Figure Legends

Figure 1. Experimental timeline. Beginning at 2-months-of age female mice were fed either

control (CD), folic acid (FADD) or choline (ChDD) deficient diets. The female mice were

maintained on these diets throughout the pregnancy and lactation until the offspring were weaned.

Once the offspring were weaned, they were fed the CD. Separate cohorts of female offspring at 2

or 10 months of age had ischemic stroke induced via the photothrombosis (PT) model. At 3.5

(CD, n = 6; FADD, n = 7; ChDD, n = 6) and 11.5 (CD, n = 6; FADD, n = 6; ChDD, n = 6)

months of age all female mouse offspring underwent ultrasound imaging (U).

Figure 2. The impact of maternal diet on posterior artery blood flow velocity. (A) Visual

representation of mouse cerebral vasculature, posterior cerebral artery (PCA) and location of

ischemic stroke. **(B)** Blood flow velocity in the PCA after ischemic stroke in 3.5- and 11.5-

month-old female offspring from control (CD), folic acid (FADD) and choline (ChDD) deficient

diet mothers. Scatter plot with mean ± SEM of 5 to 7 mice per group. * $p < 0.05$, Tukey's

pairwise comparison.

Figure 3. The impact of maternal diet on aorta and coronary artery function. (A) Aortic

pulse wave velocity as an index of aortic wall stiffness and **(B)** ratio of systolic and diastolic

coronary flow velocity after ischemic stroke in 3.5- and 11.5- month-old female offspring from

control (CD), folic acid (FADD) and choline (ChDD) deficient diet mothers. Scatter plot with

mean ± SEM of 5-7 mice per group. ** $p < 0.05$, Tukey's pairwise comparison.