Basins of attraction of microbiome structure and soil ecosystem functions

Hiroaki Fujita[1], Shigenobu Yoshida[2], Kenta Suzuki[3,4] and Hirokazu Toju[1,5,6†]

[1]Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2133, Japan

[2]Institute for Plant Protection, National Agriculture and Food Research Organization, Tsukuba, Ibaraki 305-8666, Japan

[3]Integrated Bioresource Information Division, BioResource Research Center, RIKEN, Tsukuba, Ibaraki 305-0074, Japan

[4]Institute of Multidisciplinary Sciences, Yokohama National University. Yokohama, Kanagawa 240-8501, Japan.

[5]Laboratory of Ecosystems and Coevolution, Graduate School of Biostudies, Kyoto University, Kyoto 606-8501, Japan

[6]Center for Living Systems Information Science (CeLiSIS), Graduate School of Biostudies, Kyoto University, Kyoto 606-8501, Japan

For correspondence:

toju.hirokazu.4c@kyoto-u.ac.jp

Competing interest:

HT is the founder and director of Sunlit Seedlings Ltd. The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

This work was financially supported by JST PRESTO (JPMJPR16Q6), JST FOREST

(JPMJFR2048), Human Frontier Science Program (RGP0029/2019), JSPS Grant-in-Aid for

Scientific Research (20K20586), NEDO Moonshot Research and Development Program

(JPNP18016), and JST CREST (JPMJCR23N5) to H.T., JSPS Grant-in-Aid for Scientific

Research (20K06820 and 20H03010) to K.S., and JSPS Fellowship to H.F.

Abstract

Theory predicts that biological communities can have multiple basins of attraction in terms of their species/taxonomic compositions. The presence of such basins of community structure has been examined in classic empirical studies on forest–savanna transitions and those on eutrophication in freshwater lakes. Nonetheless, it remains a major challenge to extend the investigations of multistability to species-rich microbial communities. By targeting soil microbiomes, we infer the stability landscapes of community structure based on the concepts of statistical physics. Our analysis on the compiled dataset involving 11 archaeal, 332 bacterial, and 240 fungal families detected from > 1,500 agroecosystem soil samples suggested that both prokaryotic and fungal community compositions could be classified into several basins of attraction. We also found that the basins differed greatly in their associations with crop disease prevalence in agroecosystems. A further analysis highlighted microbial taxa potentially playing key roles in transitions between basins with different ecosystem-scale functions. The statistical framework commonly applicable to diverse microbial and non-microbial communities will reorganize our understanding of relationship among community structure, stability, and functions.

Introduction

The idea that the state space of biological community structure can comprise multiple basins of attraction have inspired both empirical and theoretical ecologists since the late 1960s (Beisner et al., 2003; May, 1977; Scheffer et al., 1993). The concept of multistability has been examined in aquatic and terrestrial ecosystems (Scheffer and Carpenter, 2003; Schröder et al., 2005; Suding and Hobbs, 2009). Community structure in shallow lakes, for example, is known to show two discrete states depending on nutrient (phosphorus) concentrations as represented by the bistability of charophyte densities (Ibelings et al., 2007; Scheffer et al., 1993; Smith and Schindler, 2009). Likewise, worldwide inventories of tree cover have shown that forest and savanna vegetation types possibly represent different basins of attraction (Hirota et al., 2011; Staver et al., 2011b, 2011a). Those studies have further shown that ecosystem functions (e.g., fishery, agricultural, and forestry production) can differ greatly between such basins of biological community structure (Gunderson, 2000; Scheffer et al., 2001, 1993; Scheffer and Carpenter, 2003). Consequently, understanding how structure, stability, and biological functions are organized in real communities and ecosystems has been one of the major goals in ecology.

While classic studies targeting freshwater and terrestrial biomes have explored basins of attraction based on simple characterization of community states (e.g., tree cover percentages), recent technical advances in microbial community (microbiome) research have come to provide opportunities for deepening our knowledge of biological community stability (Amor et al., 2020; Costea et al., 2017; Faust et al., 2015; Shaw et al., 2019; Toju et al., 2018; Zaneveld et al., 2017). Based on amplicon and shotgun sequencing technologies, large datasets of microbial species/taxonomic compositions have been made available, providing a basis for exploring reproducible states in microbiome community structure (Amor et al., 2020; H Fujita et al., 2023; Hayashi et al., 2024). Such high-throughput DNA sequencing studies in medicine, for example, have shown that human individuals can be classified into three or four semi-discrete clusters in terms of their intestinal microbiome compositions (Arumugam et al., 2011; Wu et al., 2011) [see also (Jeffery et al., 2012; Knights et al., 2014)]. Intriguingly, these alternative gut microbiomes ("enterotypes") differ in their associations with human disease such as type II diabetes and Crohn's disease (Costea et al., 2017). In addition to those studies on animal-associated microbiomes (Arumugam et al., 2011; Moeller et al., 2012; Yajima et al., 2023), studies on plant-associated microbiomes have started to reorganize our recognition of how multistability of phyllosphere/rhizosphere microbiome structure is associated with ecosystem-scale processes and

functions (Toju et al., 2018, 2016). Because hundreds or thousands of replicate community

samples are available in such microbiome studies, it is now possible to discuss potential

relationship between community structure and ecosystem functions based on statistical signs of

the presence of multiple basins (and background attractors).

In theoretical ecology, stability of community states (taxonomic or species compositions) is

often discussed in the framework of stability landscapes (Beisner et al., 2003; Hastings et al.,

2018; Lewontin, 1969; Scheffer and Carpenter, 2003; Suzuki et al., 2021). On the landscape

representing stability/instability of community structure, basins of attraction are split by "tipping

points" representing unstable equilibria (Beisner et al., 2003; Scheffer et al., 2001; Scheffer and

Carpenter, 2003; Suzuki et al., 2021) (Figure 1). As these basins differ in the biological functions

of constituent communities, stable and highly functional community states can be explored within

the stability landscapes. With the application of a recently proposed mathematical approach

developed based on statistical physics (Becker and Karplus, 1997; Watanabe et al., 2014), it is

now possible to infer "energy landscapes", which represent structure of stability landscapes, from

empirical datasets of ecological communities (Dakos and Kéfi, 2022; Sánchez-Pinillos et al.,

2024; Suzuki et al., 2021). The statistical framework allows us to explore the probabilities of

community compositions within the "assembly graphs" (Coyte et al., 2021; Serván and Allesina,

2021) representing paths of possible community assembly (H Fujita et al., 2023; Suzuki et al.,

2021) (Figure 1). Although hundreds or thousands of community compositional data points are

required to apply the statistical approach (H Fujita et al., 2023; Suzuki et al., 2021), such energy

landscape analyses will allow us to define key features of stable and highly functional

microbiome states out of numerous possible combinations of microbial species or taxa. Despite

the potential for systematically profiling the relationship among community structure, stability,

and functions based on massive community datasets, the energy landscape analysis has been

applied only to a few microbial community datasets (H Fujita et al., 2023; Suzuki et al., 2021).

We here apply the emerging statistical framework to soil microbiomes, which often show

highest levels of structural diversity in nature. We compile a cropland soil microbiome dataset

consisting of > 1,500 sampling positions across the Japan Archipelago (Fujita et al., 2024). With

the massive dataset, we infer the compositional stability of prokaryotic and fungal communities

based on maximum entropy models of the energy landscape analysis (Suzuki et al., 2021). We

then examine whether the basins of attraction of soil microbiomes can differ in ecosystem-scale

functions by focusing on potential relationship between soil microbial compositions and the

prevalence of crop plant disease. We also explore key microbial taxa whose abundance critically

divide the basins representing favorable and unfavorable ecosystem functions. The results of the

energy landscape analysis are further used to infer tipping points splitting the inferred basins.

Overall, this study illustrates how we can integrate the information of community structure,

stability, and functions based on a statistical platform commonly applicable to diverse microbial

and non-microbial communities.

Methods

Dataset compilation

We compiled a publicly available dataset of cropland soil microbiomes (DDBJ accession:

DRA015491; Figure 2) with its metadata of the samples (Fujita et al., 2024). In the previous

study reporting the data (Fujita et al., 2024), 2,903 bulk soil samples collected from the field of

19 crop plant species (apple, broccoli, cabbage, celery, Chinese cabbage, eggplant, ginger,

komatsuna, lettuce, onion, potato, radish, rice, satsuma mandarin, soybean, spinach, strawberry,

sweet corn, tomato) across the Japan Archipelago from January 23, 2006 to July 28, 2014

(latitudes of the sampling positions: 26.1–42.8 °N) were subjected to the amplicon sequencing

analysis of the prokaryotic 16S rRNA region and the fungal internal transcribed spacer 1 (ITS1)

region (Fujita et al., 2024). The information of dry soil pH, electrical conductivity,

carbon/nitrogen (C/N) ratio, and available phosphorous concentration was available for 2,830,

2,610, 2,346, and 2,249 samples, respectively. Likewise, the information of crop plant disease

[the percentage of diseased plants or disease severity index (Chiang et al., 2017)] was available

for 1,471 samples (Fujita et al., 2024). The plant pathogens surveyed were *Colletotrichum*

gloeosporioides on the strawberry, *Fusarium oxysporum* on the celery, the lettuce, the strawberry,

and the tomato, *Phytophthora sojae* on the soybean, *Plasmodiophora brassicae* on Cruciferae

plants, *Pyrenochaeta lycopersici* on the tomato, *Pythium myriotylum* on the ginger, *Ralstonia*

solanacearum on the eggplant and the tomato, and *Verticillium* spp. on Chinese cabbage (Fujita

et al., 2024). After a series of quality filtering, prokaryotic and fungal community data were

available for 2,318 and 2,186 samples, respectively. In total, 579 archaeal amplicon sequence

variants (ASVs) representing 11 families, 26,640 bacterial ASVs representing 332 families, and

6,306 fungal ASVs representing 240 families were detected (Fujita et al., 2024) (Figures 2;

Figure 2–figure supplement 1).

Community structure along environmental gradients

We first inspected how prokaryotic and fungal community structure varied along environmental

gradients. For each data matrix representing the family-level compositions of prokaryotes or

fungi, a principal coordinate analysis (PCoA) was performed based on Bray-Curtis β-diversity.

The PCoA1 and PCoA2 scores were then plotted, respectively, along the axes of soil

environmental factors. Specifically, the axes of the environmental factors were defined based on

a principal component analysis (PCA) of soil pH, electrical conductivity, C/N ratio, and available

phosphorous concentration. In total, 1,771 and 1,664 samples for which the information of both

community structure and all the four environmental variables was available were included in the

analyses of prokaryotes and fungi, respectively. For each plot representing relationship between

environmental conditions and community structure, the density of data points was visualized with

the ggplot2 3.3.6 package (Wickham, 2011) of R v.4.1.2 (R Core Team, 2020).

Energy landscape analysis

We examined the stability landscape of soil microbiome structure based on the framework of an

energy landscape analysis (H Fujita et al., 2023; Suzuki et al., 2021; Watanabe et al., 2014)

(tutorials of energy landscape analyses are available at https://github.com/kecosz/rELA). In the

framework, the term "energy" is defined by the following equations based on statistical physics

(Suzuki et al., 2021; Watanabe et al., 2014). Within the "assembly graphs" representing paths of

community dynamics (Coyte et al., 2021; Serván and Allesina, 2021), probabilities of observing

specific community compositions can be explored as detailed previously (Suzuki et al., 2021). In

brief, probabilities of community states $\mathrm{p}\big(\vec{\sigma}^{(k)}\big)$ are given by

$$P\big(\vec{\sigma}^{(k)}|\varepsilon\big) = e^{-E(\vec{\sigma}^{(k)},\varepsilon)}/Z \qquad [\text{eq. 1}]$$

$$Z = \sum_{i=1}^{2^S} e^{-E(\vec{\sigma}^{(i)},\varepsilon)} \qquad [\text{eq. 2}],$$

where $\vec{\sigma}^{(k)} = (\sigma_1^{(k)}, \sigma_2^{(k)}, \dots, \sigma_S^{(k)})$ is a community state vector of k-th community state and S is

the total number of taxa (e.g., ASVs, species, genera, or families) examined. $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_M)$

is an array of continuous values representing environmental factors (e.g., soil pH and electrical

conductivity) and M is the total number of environmental parameters. $\sigma_i^{(k)}$ is a binary variable

that represents presence (1) or absence (0) of taxon *i*: i.e., there are a total of 2^S community

states. As the exploration of the 2^S community states were computationally intensive, we coded

community states at the family-level taxonomic compositions. Specifically, for each sample,

families whose relative abundance exceeds a certain threshold value (threshold for binarization)

were coded as 1, while the remaining minor families were coded as 0. Subsequently, families

whose occurrence ratios (i.e., the proportions of samples in which target families were coded as

1) were less than a certain threshold (occurrence threshold) were excluded from the dataset.

Likewise, families that appeared in almost all samples (1 – occurrence threshold) were excluded.

Note that without such thinning of input data, the dimensions of community states are too high to

be explored even using supercomputers. Therefore, exclusion of the taxa that contribute little to

the classification of community states (i.e., taxa appearing only in a small fraction of samples or

those appearing in most samples) is inevitable in the energy landscape analysis. Through

intensive preliminary computational runs with various combinations of binarization and

occurrence thresholds, we found that the number of taxa (*S*) should be kept less than 65 as

detailed in the next subsection.

When input community matrix is defined, the energy of the community state $\vec{\sigma}^{(k)}$ is given

by the extended pairwise maximum entropy model:

$$E(\vec{\sigma}^{(k)}, \varepsilon) = -\sum_{i=1}^{S} h_i \vec{\sigma}_i^{(k)} - \sum_{j=1}^{S} \sum_{i=1}^{M} g_{ij} \varepsilon_i^{(k)} \sigma_j^{(k)} - \sum_{i=1}^{S} \sum_{j=1}^{S} J_{ij} \vec{\sigma}_i^{(k)} \vec{\sigma}_j^{(k)} / 2 \qquad \text{[eq. 3]},$$

where h_i represents the net effect of implicit abiotic factors, by which *i*-th taxon is more likely

to present ($h_i > 0$) or not ($h_i < 0$), g_{ij} represents the effect of the *i*-th observed environmental

factor, and J_{ij} represents a co-occurrence pattern of *i*-th and *j*-th taxa. Since the logarithm of the

probability of a community state is inversely proportional to $E(\vec{\sigma}^{(k)})$, a community state having

lower *E* is more frequently observed. To consider dynamics on an assembly graph defined as a

network whose 2^S nodes represent possible community states and the edges represents transition

path between them (two community states are adjacent only if they have the opposite

presence/absence status for just one species), we assigned energy to nodes with the above

equation, and so imposed the directionality in state transitions. Then, by using the steepest

descent algorithm (Suzuki et al., 2021), we identified nodes having the lowest energy compared

to all its neighbors within the weighted network, and determined their basins of attraction

(Lewontin, 1969; Suzuki et al., 2021). These community states whose energy was lower than that

of all adjacent community states represent estimated point equilibria (attractors), around which

community states are expected to show transient fluctuations due to demographic stochasticity as

considered in the statistical framework (H Fujita et al., 2023; Suzuki et al., 2021) (Figure 1). Soil

pH, electrical conductivity, C/N ratio, and available phosphorous concentration were included as

environmental variables in the model after normalization within the ranges from 0 to 1.

Energy landscape structure

The energy landscapes of community structure were inferred, respectively, for three types of

datasets, namely, the prokaryotic community matrix, the fungal matrix, and the matrix including

both prokaryotes and fungi. As mentioned above, various combinations of binarization and

occurrence thresholds were examined to check the reproducibility of the results. In addition to the

energy landscape analysis based on the above-mentioned family-level delineation of community

states, analyses based on community-state delineation at the order-level were performed. In the

main body and supplementary figures of this study, we show the results at the following settings:

prokaryotes (family), binarization = 0.020, occurrence = 0.10; prokaryotes (order), binarization =

0.020, occurrence = 0.10; fungi (family), binarization = 0.001, occurrence = 0.05; fungi (family),

binarization = 0.001, occurrence = 0.10; prokaryotes + fungi (family), binarization = 0.030,

occurrence = 0.10; prokaryotes + fungi (order), binarization = 0.030, occurrence = 0.10. Note that

these thresholds were selected to make the state space (2^S) neither too simplified (e.g., $S < 30$)

nor too complex ($S < 65$).

For each setting, the parameters of the extended pairwise maximum entropy model [eq. 3]

were adjusted to the empirical data. More precisely, the maximum likelihood estimates of h_i, g_{ij},

and J_{ij} was obtained by a stochastic approximation method as detailed elsewhere (Suzuki et al.,

2021). The parameters were regularized by a logistic prior with location 0 and scale 2.0 (for

environmental responses) or 0.5 (for pairwise relationships) (Harris, 2016). Hyperparameters for

the algorithm, criterion value for judging the convergence of parameters qth = 10^{-5}, were set

according to a series of preliminary analyses. Based on the inferred maximum entropy model, we

determined basins of attraction (Lewontin, 1969) within the energy landscape based on a steepest

descent procedure (Suzuki et al., 2021). The structure of the energy landscape was visualized by

showing the energy of each soil sample on the two-dimensional surface of the community state

space defined with the abovementioned PCoA scores. The default setting of environmental

variables (the mean value for each of soil pH, electrical conductivity, C/N ratio, and available

phosphorous concentration) was used in the energy calculation. Spline smoothing of the energy

landscape was performed with optimized penalty scores using the mgcv v.1.8-40 package (Wood,

2022) of R. For each analysis of the prokaryote, fungi, and prokaryote + fungi datasets, 1,771,

1,664, and 1,474 samples for which the information of both community structure and all the four

environmental variables was available were subjected to the analysis, respectively.

Ecosystem functions and key taxa

For the inferred basins of microbial community compositions, associations with crop disease

prevalence were examined. We first constructed the list of soil samples whose community

structure was located within each basin of attraction. We then evaluated the ecosystem-scale

properties of the basins in light of the metadata of crop disease symptoms (Fujita et al., 2024).

Specifically, for each basin, we calculated the proportion of constituent soil samples with the

minimal level of crop disease symptoms (the percentage of diseased plants < 20 or disease

severity index < 20; (Fujita et al., 2024)). The bottoms of basins representing different levels of

crop disease prevalence were then compared in terms of taxonomic compositions in order to

explore microbial taxa that were keys to distinguish potentially disease-suppressive and disease-

promotive soil ecosystems.

Disconnectivity graphs

For the reconstructed energy landscape, we inferred "disconnectivity graphs" (Suzuki et al.,

2021) representing how basins of attraction were split by tipping points (Figure 1A). Within a

disconnectivity graph, community states whose energy is much lower than the energy of

connected tipping points are expected to be resistant to perturbations (demographic stochasticity).

In contrast, community states with small energy gaps to tipping points may be shifted from

current basins to adjacent basins with minimal perturbations.

Results

Community structure along environmental gradients

On each plot showing community compositions (PCoA1 or PCoA2 scores) along the soil

environmental gradient (Figure 3), multiple clusters of data points were observed for both

prokaryotes and fungi (Figure 3–figure supplements 1-2). In other words, community states are

expected to be classified into some clusters even under equivalent edaphic conditions.

Energy landscape structure

The energy landscape of the family-level prokaryotic data included several major basins differing

remarkably in associations with the prevalence of crop plant disease (Figure 4). Specifically,

59.6% of soil samples located within a basin (basin ID = 0IK1G2) were associated with the

minimal plant-disease level, while the proportion was only 10.7% for another basin (LQWZ02)

(Figure 4C-D). The presence of basins differing greatly in their associations with plant-disease

levels was inferred as well at the order-level analysis of the prokaryotic data (Figure 4–figure

supplement 1). Such variation in crop disease prevalence among inferred basins was observed

also for the family-level analysis of fungal community structure (Figure 5). Specifically, while

57.9% of samples belonging to the basin 7QH9moTf8Xa, but none of the samples belonging to

the basin 68C0849W020, were associated with the minimal plant-disease level (Figure 5D).

Meanwhile, such difference in associations with disease prevalence was moderate in an analysis

in which a smaller number of fungal families were examined to define community states (Figure

5–figure supplement 1). The presence of multiple basins, which differed in associations with

crop-disease prevalence, was suggested even when prokaryotic and fungal community data were

simultaneously analyzed (Figure 4–figure supplements 2-3).

Ecosystem functions and key taxa

In an analysis of the prokaryotic community structure, 19 families were keys to distinguish basins

differing in associations with crop-disease prevalence (Figure 4D). The presence of

Pyrinomonadaceae and Vicinamibacteraceae, for example, was unique to the basin with the

highest proportion of samples showing the minimal plant-disease level (Figure 4D). Likewise, in

an analysis of the fungal community structure, the basin associated closely with the minimal

plant-disease prevalence (7QH9moTf8Xa) was defined by the presence of several families such

as Basidiobolaceae, Cordycipitaceae, and Gelatinodiscaceae (Figure 5D). The exploration of

microbial taxa keys to distinguish basins with different ecosystem-level functions can be

performed at other taxonomic levels (e.g., order-level; Figures 4–figure supplements 1 and 3).

Disconnectivity graphs

Within the energy landscape of the family-level analysis of prokaryotes (Figure 4), both the

basins associated with the least-diseased (OIK1G2) and most-diseased (N21H04) crop status

were the deepest among the inferred basins (i.e., showing the largest energy gaps from the bottom

to tipping points; Figure 6A-B). In the family-level analysis of fungi, the basin associated with

the least-diseased status (7QH9moTf8Xa) was the deepest, while the other basin representing the

most-diseased status (68C0849W020) was the shallowest (Figure 6C).

Discussion

We have estimated the stability landscape structure of complex microbiomes based on a

statistical framework commonly applicable to diverse types of biological communities. The

energy landscape analysis allows systematic analyses of taxon-rich community datasets by

incorporating the information of multiple environmental factors (Dakos and Kéfi, 2022; Sánchez-

Pinillos et al., 2024; Suzuki et al., 2021). While classic studies on community multistability have

discussed ecological processes spanning a few intuitively distinguishable community states

[high/low tree cover in forest-savanna transitions (Hirota et al., 2011; Staver et al., 2011a, 2011b)

or macrophyte-/phytoplankton-dominated state in shallow lakes (Ibelings et al., 2007; Scheffer

and Carpenter, 2003)], it is now made possible to define basins of attraction based on high-

dimensional community datasets involving hundreds of species/taxa (Arumugam et al., 2011;

Costea et al., 2017; H Fujita et al., 2023; Guim Aguadé-Gorgorió et al., 2023; Hayashi et al.,

2024). Application of the general statistical platform will enhance our understanding of how

stability landscape properties differ among diverse microbial and non-microbial systems.

Despite numerous potential compositions (2^S community states; S is the number of

considered species/taxa), the prokaryotic and fungal community states were grouped into small

numbers of basins within energy landscapes (Figures 4-5). This result suggests that soil

microbiome structure remain within certain regions even after demographic perturbations. In

other words, once trapped in a basin of attraction, large shifts in community structure would not occur without perturbations whose strength exceed certain thresholds (Beisner et al., 2003; Lewontin, 1969; May, 1977; Scheffer et al., 1993). Importantly, the threshold strength of perturbations is estimated as the energy gap between bottoms of basins and tipping points (Suzuki et al., 2021) (Figure 6A). Furthermore, potential paths of community structural transitions can be quantitatively inferred as illustrated in disconnectivity graphs (Suzuki et al., 2021) (Figures 6B-C). Such statistical framework of quantitative science will entail novel opportunities for testing theories on biological community processes in the era of massive datasets.

Among potential processes or mechanisms underlying the multistability of community structure, historical contingency is of particular interest (Fukami, 2015). In the local assembly of microbial communities, early colonizers or residents can prevent the settlement of followers by constructing physical barriers (e.g., biofilms and mycelia) (Baümler and Sperandio, 2016; Fukami, 2015; Leopold et al., 2017; Verbruggen et al., 2013; Werner and Kiers, 2015) or emitting antibiotics (Mendes et al., 2013; Raaijmakers et al., 2002). In addition to those antagonistic effects on late colonizers, webs of mutualistic or commensalistic interactions within the microbiomes of early colonizers (Elias and Banin, 2012; Hiroaki Fujita et al., 2023; Zelezniak et al., 2015) would influence community dynamics. Due to such "priority effects" (Fukami, 2015), bacterial and fungal community compositions may persist within limited ranges of community states without substantial perturbations. Given that abilities to form physical or chemical barriers can differ greatly among microbial species/taxa (Mendes et al., 2013; Raaijmakers et al., 2002; Werner and Kiers, 2015), such variation in constituent species' priority effects may underly the observed variation in the depth of basins (Figure 6B-C).

The inference of stability landscape structure provided an opportunity for evaluating relationship between community stability and ecosystem-scale functions. The basins of attraction of prokaryotic/fungal community structure differed considerably in associations with crop disease prevalence (Figure 5), suggesting the presence of "stable and favorable" and "stable but unfavorable" states of microbiomes (Mendes et al., 2011; Schlatter et al., 2017; Yuan et al., 2020) in terms of agricultural productivity. This finding adds an important dimension of discussion on the use of microbes in agriculture. Beyond investigations on single species/strains of microbes, microbiome studies have explored sets of microbes that collectively maximize biological functions (Jansson and Hofmockel, 2018; Toju et al., 2018; Trivedi et al., 2020; Vorholt et al., 2017). In particular, experimental studies on "synthetic" communities have reorganized our

knowledge of microbiome functions (Jansson and Hofmockel, 2018; Trivedi et al., 2020; Vorholt et al., 2017). Nonetheless, such microbial functions cannot be realized in real agroecosystems if the synthesized or designed microbiome compositions are vulnerable to biotic and abiotic environmental changes in the wild (Mazzola and Freilich, 2017). Thus, in addition to functional properties, compositional stability is the key to manage microbiomes in agroecosystems (Faust and Raes, 2012; Toju et al., 2020; Vorholt et al., 2017).

In our analysis across the Japan Archipelago, prokaryotic and fungal taxa keys to distinguish least-diseased and severely-diseased states of soil microbiomes were highlighted (Figures 4-5). Among them, Basidiobolaceae and Cordycipitaceae are of particular interest because they include many species potentially utilized as biological control agents for suppressing pest insects (Meyling and Eilenberg, 2007; Möckel et al., 2022). Gelatinodiscaceae is another fungal taxon playing potentially important roles as symbionts of plants (Johnston et al., 2019). These results illuminate the hypothesis that plant disease could be suppressed under the coexistence of multiple prokaryotic and fungal taxa with favorable ecosystem functions (Toju et al., 2018; Toju and Tanaka, 2019). Thus, statistical analyses of stability landscapes allow the exploration of key species or taxa (Paine, 1966; Power et al., 1996), whose management could result in transitions from unfavorable ecosystem states to favorable ones (Gunderson, 2000; Scheffer et al., 2001, 1993; Scheffer and Carpenter, 2003). Given that most prokaryotic and fungal families highlighted in our analysis have cosmopolitan distributions, a next crucial step is to test whether the basins defined across the Japan Archipelago can be used to categorize disease-suppressive and disease-susceptible microbiomes in other regions on the globe.

Although the energy landscape analysis enhances our understanding of community stability and functions, its results should be interpreted with caution. First, given that classic empirical studies examined community multistability with system-specific simple criteria [e.g., high/low tree cover (Hirota et al., 2011; Staver et al., 2011a, 2011b)], special care should be taken when we extend the approach to species-rich (high-dimensional) community datasets (Guim Aguadé-Gorgorió et al., 2023). In other words, unambiguous and broadly applicable criteria based on statistical evaluation are the prerequisite for comparative analyses of community multistability. Although we applied a straightforward statistical definition of basins of attraction (Suzuki et al., 2021) (Figure 1) in light of classic theoretical studies (Beisner et al., 2003; Lewontin, 1969; May, 1977; Scheffer et al., 1993), continuous methodological improvements should be explored towards further comprehensive analyses. Second, our analysis on hyper-diverse soil microbiomes

incurred substantial computational costs, forcing us to limit the energy landscape analysis to

family-level input data. Further improvements of codes are necessary for inferring stability

landscapes at genus-, species-, or strain-level analyses. Third, it should be acknowledged that

detailed discussion on ecological processes require time-series datasets (Davidson et al., 2023;

Scheffer et al., 2012, 2009). Because our present data lacked the information of temporal changes

in community structure, we are unable to discuss the frequency and pace of community structural

transitions between basins of attraction. Monitoring of microbiome compositions (Faust et al.,

2015; Hayashi et al., 2024; Yajima et al., 2023) is necessary for filling the gap between

theoretical and empirical studies (Long et al., 2024).

The energy landscape framework of multistability analysis is readily applicable to a wide

range of microbiome datasets. Application to human microbiome data is of particular interest in

terms of the confirmation of the existence of multiple basins of attraction (Jeffery et al., 2012). In

addition, insights into the key microbial species/taxa that would play key roles in the transitions

from disease-associated microbiome states to healthy ones will open new directions of

microbiome therapy. Furthermore, time-series analyses of community dynamics on stability

landscapes will allow us to forecast and prevent transitions into unfavorable community states

[e.g., dysbiosis (Carding et al., 2015; H Fujita et al., 2023; Long et al., 2024)]. Along with such

extensions of observational research, experimental studies controlling key species/taxa or

environmental parameters (Schröder et al., 2005) will promote both basic and applied sciences of

ecosystem functions.

Acknowledgements. We thank the SuperComputer System, Institute for Chemical Research,

Kyoto University for the use of super computers.

Author contributions

H.F., S.Y. and H.T. designed the work. H.F. performed molecular experiments. H.F. and H.T.

analyzed the data. H.T. wrote the paper with H.F., S.Y., and K.S.

Author ORCIDs

Hiroaki Fujita: https://orcid.org/0009-0003-5422-9775

Shigenobu Yoshida: https://orcid.org/0000-0001-5105-0807

Kenta Suzuki: https://orcid.org/0000-0002-0639-2303

Hirokazu Toju: https://orcid.org/0000-0002-3362-3285

Data accessibility

423   The accession number of the DDBJ Sequence Read Archive: DRA015491. The microbial

424   community matrices are provided with codes at our GitHub repository (https://github.com/hiro-

425   toju/Soil_EnergyLandscape_NARO3000) [to be released after the acceptance of the manuscript].

References

Amor DR, Ratzke C, Gore J. 2020. Transient invaders can induce shifts between alternative
stable states of microbial communities. *Sci Adv* **6**:eaay8676. doi:10.1126/sciadv.aay8676

Arumugam M, Raes J, Pelletier E, Le Paslier D, Yamada T, Mende DR, Fernandes GR, Tap J,
Bruls T, Batto JM, Bertalan M, Borruel N, Casellas F, Fernandez L, Gautier L, Hansen T,
Hattori M, Hayashi T, Kleerebezem M, Kurokawa K, Leclerc M, Levenez F, Manichanh C,
Nielsen HB, Nielsen T, Pons N, Poulain J, Qin J, Sicheritz-Ponten T, Tims S, Torrents D,
Ugarte E, Zoetendal EG, Wang J, Guarner F, Pedersen O, De Vos WM, Brunak S, Doré J,
Weissenbach J, Ehrlich SD, Bork P. 2011. Enterotypes of the human gut microbiome.
Nature **473**:174–180. doi:10.1038/nature09944

Baümler AJ, Sperandio V. 2016. Interactions between the microbiota and pathogenic bacteria in
the gut. *Nature*. doi:10.1038/nature18849

Becker OM, Karplus M. 1997. The topology of multidimensional potential energy surfaces:
Theory and application to peptide structure and kinetics. *Journal of Chemical Physics*
106:1495. doi:10.1063/1.473299

Beisner BE, Haydon DT, Cuddington K. 2003. Alternative stable states in ecology. *Front Ecol*
Environ **1**:376–382. doi:10.1890/1540-9295(2003)001[0376:ASSIE]2.0.CO;2

Carding S, Verbeke K, Vipond DT, Corfe BM, Owen LJ. 2015. Dysbiosis of the gut microbiota in
disease. *Microb Ecol Health Dis* **26**:26191. doi:10.3402/mehd.v26.26191

Chiang KS, Liu HI, Bock CH. 2017. A discussion on disease severity index values. Part I: warning
on inherent errors and suggestions to maximise accuracy. *Annals of Applied Biology*
171:139–154. doi:10.1111/aab.12362

Costea PI, Hildebrand F, Manimozhiyan A, Bäckhed F, Blaser MJ, Bushman FD, De Vos WM,
Ehrlich SD, Fraser CM, Hattori M, Huttenhower C, Jeffery IB, Knights D, Lewis JD, Ley
RE, Ochman H, O'Toole PW, Quince C, Relman DA, Shanahan F, Sunagawa S, Wang J,
Weinstock GM, Wu GD, Zeller G, Zhao L, Raes J, Knight R, Bork P. 2017. Enterotypes in
the landscape of gut microbial community composition. *Nat Microbiol* **3**:8–16.
doi:10.1038/s41564-017-0072-8

Coyte KZ, Rao C, Rakoff-Nahoum S, Foster KR. 2021. Ecological rules for the assembly of
microbiome communities. *PLoS Biol* **19**:e3001116. doi:10.1371/JOURNAL.PBIO.3001116

Dakos V, Kéfi S. 2022. Ecological resilience: What to measure and how. *Environmental Research*
Letters **17**:043003. doi:10.1088/1748-9326/ac5767

Davidson TA, Sayer CD, Jeppesen E, Søndergaard M, Lauridsen TL, Johansson LS, Baker A,
Graeber D. 2023. Bimodality and alternative equilibria do not help explain long-term
patterns in shallow lake chlorophyll-a. *Nat Commun* **14**:398. doi:10.1038/s41467-023-
36043-9

Elias S, Banin E. 2012. Multi-species biofilms: Living with friendly neighbors. *FEMS Microbiol*
Rev **36**:990–1004. doi:10.1111/j.1574-6976.2012.00325.x

Faust K, Lahti L, Gonze D, de Vos WM, Raes J. 2015. Metagenomics meets time series analysis:
Unraveling microbial community dynamics. *Curr Opin Microbiol* **25**:56–66.
doi:10.1016/j.mib.2015.04.004

Faust K, Raes J. 2012. Microbial interactions: From networks to models. *Nat Rev Microbiol*
10:538–550. doi:10.1038/nrmicro2832

Fujita H, Ushio M, Suzuki K, Abe M, Yamamichi M, Iwayama K, Canarini A, Hayashi I,
Fukushima K, Fukuda S, Kiers E, Toju H. 2023. Alternative stable states, nonlinear behavior,
and predictability of microbiome dynamics. *Microbiome* **11**:63. doi:10.1186/s40168-023-
01474-5

Fujita Hiroaki, Ushio M, Suzuki K, Abe MS, Yamamichi M, Okazaki Y, Canarini A, Hayashi I,
Fukushima K, Fukuda S, Kiers ET, Toju H. 2023. Facilitative interaction networks in
experimental microbial community dynamics. *Front Microbiol* **14**:1153952.
doi:10.3389/fmicb.2023.1153952

Fujita H, Yoshida S, Suzuki K, Toju H. 2024. Soil prokaryotic and fungal biome structures
associated with crop disease status across the Japan Archipelago. *mSphere* in press.

Fukami T. 2015. Historical contingency in community assembly: integrating niches, species
pools, and priority effects. *Annu Rev Ecol Evol Syst* **46**:1–23. doi:10.1146/annurev-ecolsys-
110411-160340

Guim Aguadé-Gorgorió, Jean-François Arnoldi, Matthieu Barbier, Sonia Kéfi. 2023. A taxonomy
of multiple stable states in complex ecological communities. *bioRxiv*
https://doi.org/10.1101/2023.08.30.555051.

Gunderson LH. 2000. Ecological resilience - In theory and application. *Annu Rev Ecol Syst*
31:425–439. doi:10.1146/annurev.ecolsys.31.1.425

Harris DJ. 2016. Inferring species interactions from co-occurrence data with Markov networks.
Ecology **97**:3308–3314. doi:10.1002/ecy.1605

Hastings A, Abbott KC, Cuddington K, Francis T, Gellner G, Lai YC, Morozov A, Petrovskii S,
Scranton K, Zeeman M Lou. 2018. Transient phenomena in ecology. *Science (1979)*
361:eaat6412. doi:10.1126/science.aat6412

Hayashi I, Fujita H, Toju H. 2024. Deterministic and stochastic processes generating alternative
states of microbiomes. *ISME Communications* **4**:ycae007. doi:10.1093/ismeco/ycae007

Hirota M, Holmgren M, Van Nes EH, Scheffer M. 2011. Global resilience of tropical forest and
savanna to critical transitions. *Science (1979)* **334**:232–235. doi:10.1126/science.1210657

Ibelings BW, Portielje R, Lammens EHRR, Noordhuis R, Van Den Berg MS, Joosse W, Meijer
ML. 2007. Resilience of alternative stable states during the recovery of shallow lakes from
eutrophication: Lake Veluwe as a case study. *Ecosystems* **10**:4–16. doi:10.1007/s10021-006-
9009-4

Jansson JK, Hofmockel KS. 2018. The soil microbiome — from metagenomics to metaphenomics.
Curr Opin Microbiol **43**:162–168. doi:10.1016/j.mib.2018.01.013

Jeffery IB, Claesson MJ, O'Toole PW, Shanahan F. 2012. Categorization of the gut microbiota:
Enterotypes or gradients? *Nat Rev Microbiol* **10**:591–592. doi:10.1038/nrmicro2859

Johnston PR, Quijada L, Smith CA, Baral HO, Hosoya T, Baschien C, Pärtel K, Zhuang WY,
Haelewaters D, Park D, Carl S, López-Giráldez F, Wang Z, Townsend JP. 2019. A
multigene phylogeny toward a new phylogenetic classification of Leotiomycetes. *IMA*
Fungus **10**:1–22. doi:10.1186/s43008-019-0002-x

Knights D, Ward TL, McKinlay CE, Miller H, Gonzalez A, McDonald D, Knight R. 2014.
Rethinking enterotypes. *Cell Host Microbe* **16**:433–437. doi:10.1016/j.chom.2014.09.013

Leopold DR, Wilkie JP, Dickie IA, Allen RB, Buchanan PK, Fukami T. 2017. Priority effects are
interactively regulated by top-down and bottom-up forces: evidence from wood decomposer
communities. *Ecol Lett* **20**:1054–1063. doi:10.1111/ele.12803

Lewontin RC. 1969. The meaning of stability. *Brookhaven Symp Biol* **22**:13–23.
doi:10.2307/j.ctv14161w4.26

Long C, Deng J, Nguyen J, Liu Y-Y, Alm EJ, Solé R, Saavedra S. 2024. Structured community
transitions explain the switching capacity of microbial systems. *Proceedings of the National*
Academy of Sciences **121**:e2312521121. doi:10.1073/pnas.2312521121

May RM. 1977. Thresholds and breakpoints in ecosystems with a multiplicity of stable states.
Nature **269**:471–477. doi:10.1038/269471a0

Mazzola M, Freilich S. 2017. Prospects for biological soilborne disease control: Application of
indigenous versus synthetic microbiomes. *Phytopathology* **107**:256–263.
doi:10.1094/PHYTO-09-16-0330-RVW

Mendes R, Garbeva P, Raaijmakers JM. 2013. The rhizosphere microbiome: Significance of plant
beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiol Rev*
37:634–663. doi:10.1111/1574-6976.12028

Mendes R, Kruijt M, de Bruijn I, Dekkers E, van der Voort M, Schneider JHM, Piceno YM,
DeSantis TZ, Andersen GL, Bakker P a HM, Raaijmakers JM. 2011. Deciphering the
rhizosphere microbiome for disease-suppressive bacteria. *Science* **332**:1097–100.
doi:10.1126/science.1203980

Meyling N V., Eilenberg J. 2007. Ecology of the entomopathogenic fungi Beauveria bassiana and
Metarhizium anisopliae in temperate agroecosystems: Potential for conservation biological
control. *Biological Control* **43**:145–155. doi:10.1016/j.biocontrol.2007.07.007

Möckel L, Meusemann K, Misof B, Schwartze VU, De Fine Licht HH, Voigt K, Stielow B, de
Hoog S, Beutel RG, Buellesbach J. 2022. Phylogenetic Revision and Patterns of Host
Specificity in the Fungal Subphylum Entomophthoromycotina. *Microorganisms* **10**:256.
doi:10.3390/microorganisms10020256

Moeller AH, Degnan PH, Pusey AE, Wilson ML, Hahn BH, Ochman H. 2012. Chimpanzees and
humans harbour compositionally similar gut enterotypes. *Nat Commun* **3**:1179.
doi:10.1038/ncomms2159

Paine RT. 1966. Food Web Complexity and Species Diversity. *Am Nat* **100**:65–75.
doi:10.1086/282400

Power ME, Tilman D, Estes JA, Menge BA, Bond WJ, Mills LS, Daily G, Castilla JC, Lubchenco
J, Paine RT. 1996. Challenges in the quest for keystones. *Bioscience* **46**:609–620.
doi:10.2307/1312990

R Core Team. 2020. R: A language and environment for statistical computing. *R: A language and*
environment for statistical computing R Foundation for Statistical Computing, Vienna,
Austria URL https://wwwR-project.org/.

Raaijmakers JM, Vlami M, de Souza JT. 2002. Antibiotic production by bacterial biocontrol
agents. *Antonie Van Leeuwenhoek* **81**:537–547. doi:10.1023/A:1020501420831

Sánchez-Pinillos M, Dakos V, Kéfi S. 2024. Ecological dynamic regimes: A key concept for
assessing ecological resilience. *Biol Conserv* **289**:110409. doi:10.1016/j.biocon.2023.110409

Scheffer M, Bascompte J, Brock WA, Brovkin V, Carpenter SR, Dakos V, Held H, Van Nes EH,
Rietkerk M, Sugihara G. 2009. Early-warning signals for critical transitions. *Nature* **461**:53–
59. doi:10.1038/nature08227

Scheffer M, Carpenter S, Foley J a, Folke C, Walker B. 2001. Catastrophic shifts in ecosystems.
Nature **413**:591–596. doi:10.1038/35098000

Scheffer M, Carpenter SR. 2003. Catastrophic regime shifts in ecosystems: Linking theory to
observation. *Trends Ecol Evol* **18**:648–656. doi:10.1016/j.tree.2003.09.002

Scheffer M, Carpenter SR, Lenton TM, Bascompte J, Brock W, Dakos V, Van De Koppel J, Van
De Leemput IA, Levin SA, Van Nes EH, Pascual M, Vandermeer J. 2012. Anticipating
critical transitions. *Science (1979)* **338**:344–348. doi:10.1126/science.1225244

Scheffer M, Hosper SH, Meijer ML, Moss B, Jeppesen E. 1993. Alternative equilibria in shallow
lakes. *Trends Ecol Evol* **8**:275–279. doi:10.1016/0169-5347(93)90254-M

Schlatter D, Kinkel L, Thomashow L, Weller D, Paulitz T. 2017. Disease suppressive soils: New
insights from the soil microbiome. *Phytopathology* **107**:1284–1297. doi:10.1094/PHYTO-
03-17-0111-RVW

Schröder A, Persson L, De Roos AM. 2005. Direct experimental evidence for alternative stable
states: A review. *Oikos* **110**:3–19. doi:10.1111/j.0030-1299.2005.13962.x

Serván CA, Allesina S. 2021. Tractable models of ecological assembly. *Ecol Lett* **24**:1029–1037.
doi:10.1111/ele.13702

Shaw LP, Bassam H, Barnes CP, Walker AS, Klein N, Balloux F. 2019. Modelling microbiome
recovery after antibiotics using a stability landscape framework. *ISME Journal* **13**:1845–
1856. doi:10.1038/s41396-019-0392-1

Smith VH, Schindler DW. 2009. Eutrophication science: where do we go from here? *Trends Ecol*
Evol **24**:201–207. doi:10.1016/j.tree.2008.11.009

Staver AC, Archibald S, Levin S. 2011a. Tree cover in sub-Saharan Africa: Rainfall and fire
constrain forest and savanna as alternative stable states. *Ecology* **92**:1063–1072.
doi:10.1890/10-1684.1

Staver AC, Archibald S, Levin SA. 2011b. The global extent and determinants of savanna and
forest as alternative biome states. *Science (1979)* **334**:230–232.
doi:10.1126/science.1210465

Suding KN, Hobbs RJ. 2009. Threshold models in restoration and conservation: a developing
framework. *Trends Ecol Evol* **24**:271–279. doi:10.1016/j.tree.2008.11.012

Suzuki K, Nakaoka S, Fukuda S, Masuya H. 2021. Energy landscape analysis elucidates the
multistability of ecological communities across environmental gradients. *Ecol Monogr* **91**:1–
21. doi:10.1002/ecm.1469

Toju H, Abe MS, Ishii C, Hori Y, Fujita H, Fukuda S. 2020. Scoring species for synthetic
community design: network analyses of functional core microbiomes. *Front Microbiol*
11:1361–1361. doi:10.3389/fmicb.2020.01361

Toju H, Peay KG, Yamamichi M, Narisawa K, Hiruma K, Naito K, Fukuda S, Ushio M, Nakaoka
S, Onoda Y, Yoshida K, Schlaeppi K, Bai Y, Sugiura R, Ichihashi Y, Minamisawa K, Kiers
ET. 2018. Core microbiomes for sustainable agroecosystems. *Nat Plants* **4**:247–257.
doi:10.1038/s41477-018-0139-4

Toju H, Tanaka Y. 2019. Consortia of anti-nematode fungi and bacteria in the rhizosphere of
soybean plants attacked by root-knot nematodes. *R Soc Open Sci* **6**:181693.
doi:10.1098/rsos.181693

Toju H, Yamamoto S, Tanabe AS, Hayakawa T, Ishii HS. 2016. Network modules and hubs in
plant-root fungal biomes. *J R Soc Interface* **13**:20151097–20151097.
doi:10.1098/rsif.2015.1097

Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. 2020. Plant–microbiome interactions: from
community assembly to plant health. *Nat Rev Microbiol* **18**:607–621. doi:10.1038/s41579-
020-0412-1

Verbruggen E, van der Heijden MGA, Rillig MC, Kiers ET. 2013. Mycorrhizal fungal
establishment in agricultural soils: Factors determining inoculation success. *New Phytologist*
197:1104–1109. doi:10.1111/j.1469-8137.2012.04348.x

Vorholt JA, Vogel C, Carlström CI, Müller DB. 2017. Establishing causality: opportunities of
synthetic communities for plant microbiome research. *Cell Host Microbe* **22**:142–155.
doi:10.1016/j.chom.2017.07.004

Watanabe T, Masuda N, Megumi F, Kanai R, Rees G. 2014. Energy landscape and dynamics of
brain activity during human bistable perception. *Nat Commun* **5**:4765.
doi:10.1038/ncomms5765

Werner GDA, Kiers ET. 2015. Order of arrival structures arbuscular mycorrhizal colonization of
plants. *New Phytologist* **205**:1515–1524. doi:10.1111/nph.13092

Wickham H. 2011. ggplot2. *Wiley Interdiscip Rev Comput Stat* **3**:180–185. doi:10.1002/wics.147

Wood S. 2022. mgcv: Mixed GAM computation vehicle with automatic smoothness estimation
available at https://cran.r-project.org/web/packages/mgcv/index.html.

Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, Bewtra M, Knights D,
Walters WA, Knight R, Sinha R, Gilroy E, Gupta K, Baldassano R, Nessel L, Li H, Bushman
FD, Lewis JD. 2011. Linking long-term dietary patterns with gut microbial enterotypes.
Science (1979) **334**:105–108. doi:10.1126/science.1208344

Yajima D, Fujita H, Hayashi I, Shima G, Suzuki K, Toju H. 2023. Core species and interactions
prominent in fish-associated microbiome dynamics. *Microbiome* **11**:53.
doi:10.1186/s40168-023-01498-x

Yuan J, Wen T, Zhang H, Zhao M, Penton CR, Thomashow LS, Shen Q. 2020. Predicting disease
occurrence with high accuracy based on soil macroecological patterns of Fusarium wilt.
ISME Journal **14**:2936–2950. doi:10.1038/s41396-020-0720-5

Zaneveld JR, McMinds R, Thurber RV. 2017. Stress and stability: Applying the Anna Karenina
principle to animal microbiomes. *Nat Microbiol* **2**:17121. doi:10.1038/nmicrobiol.2017.121

Zelezniak A, Andrejev S, Ponomarova O, Mende DR, Bork P, Patil KR. 2015. Metabolic
dependencies drive species co-occurrence in diverse microbial communities. *Proc Natl Acad*
Sci U S A **112**:6449–6454. doi:10.1073/pnas.1421834112

Figure 1. Schema of multistability of ecological communities. (**A**) Basins of attraction and tipping points. The structure of "stability landscapes" showing relationship between community states (species or taxonomic compositions) and their stability is inferred based on the energy landscape analysis. The "energy" of each community state is calculated with maximum entropy models as detailed in Methods. Lower energy represents a more stable community state on a stability landscape. Transient fluctuations around the bottoms of basins (i.e., point attractors) are assumed as probabilistic phenomena in the statistical approach. (**B**) Assembly graph. To explore numerous possible states of real ecological communities, input data are binarized in the energy landscape analysis. Potential transitions between community states are then considered within "assembly graphs", in which paths between different species/taxonomic compositions are treated as network links. Thus, by the assembly-graph approach, the energy landscape analysis provides a general framework for inferring the structure of stability landscapes in empirical studies of complex microbiome datasets.

Figure 2. Community structure of the source data. The family-level compositions of prokaryotes (**A**) and fungi (**B**) are shown based on the source dataset (Fujita et al., 2024). The soil samples from which DNA sequence data were unavailable for either prokaryotic 16S rRNA or fungal ITS regions are indicated as blanks.

Figure supplement 1. Community structure of the source data (order- and genus-level compositions).

Figure 3. Community structure along environmental gradients. The scores representing prokaryotic/fungal community compositions (community PCoA1 scores) are shown along each PCA axis of soil environmental conditions. Regarding the environmental PCA axes, factor loadings of environmental variables examined (pH, electrical conductivity, C/N ratio, and available phosphorous concentration) are shown separately for prokaryotic ($N = 1{,}771$) and fungal ($N = 1{,}664$) datasets.

Figure supplement 1. Prokaryotic community structure along environmental gradients (detailed analyses).

Figure supplement 2. Fungal community structure along environmental gradients (detailed analyses).

Figure 4. Energy landscape of prokaryotic communities. (**A**) Inferred energy landscape of family-level prokaryotic community structure (threshold for binarization = 0.020; occurrence threshold = 0.10; *S* = 35). The surface of energy levels was reconstructed across the PCoA space of fungal community structure (community PCoA1 and PCoA2 scores in Figures 2–figure supplement 1) based on spline smoothing. Community states with lower energy are inferred to be more stable. (**B**) Landscape of crop disease prevalence. Across the PCoA space of prokaryotic compositions, the proportion of samples with disease severity index < 20 is shown based on spline smoothing. (**C**) Community data points on the energy landscape. The axis of "energy of community state" is more expanded than that in panel **A** in order to cover the range of samples. Data points (samples) indicated by the same color belong to the same basins of attraction, which are represented by the IDs of the community states whose energy is lower than that of any adjacent community states (i.e., bottoms of basins). (**D**) Key taxa whose abundance represent basins. In the upper panel, the mean proportion of soil samples with the minimum level of plant (crop) disease symptoms (the percentage of diseased plants < 20 or disease severity index < 20) is shown for each basin. The lower panel indicates the key taxa whose abundance characterizes difference among the bottoms of the basins.

Figure supplement 1. Energy landscape of prokaryotic communities (order-level compositions;

threshold for binarization = 0.020; occurrence threshold = 0.10; S = 32).

Figure supplement 2. Energy landscape of communities including both prokaryotes and fungi

(family-level compositions; threshold for binarization = 0.030; occurrence threshold = 0.10; S =

31).

Figure supplement 3. Energy landscape of communities including both prokaryotes and fungi

(order-level compositions; threshold for binarization = 0.030; occurrence threshold = 0.10; S =

32).

Figure 5. Energy landscape of fungal communities. (**A**) Inferred energy landscape of family-level

fungal community structure (threshold for binarization = 0.001; occurrence threshold = 0.05; S =

62). The surface of energy levels was reconstructed across the PCoA space of fungal community

structure (community PCoA1 and PCoA2 scores in in Figures 2–figure supplement 2) based on

spline smoothing. Community states with lower energy are inferred to be more stable. (**B**)

Landscape of crop disease prevalence. Across the PCoA space of prokaryotic compositions, the

proportion of samples with disease severity index < 20 is shown based on spline smoothing. (**C**)

Community data points on the energy landscape. The axis of "energy of community state" is

more expanded than that in panel **A** in order to cover the range of samples. Data points (samples)

indicated by the same color belong to the same basins of attraction, which are represented by the

IDs of the community states whose energy is lower than that of any adjacent community states

(i.e., bottoms of basins). (**D**) Key taxa whose abundance represent basins. In the upper panel, the

mean proportion of soil samples with the minimum level of plant (crop) disease symptoms (the

percentage of diseased plants < 20 or disease severity index < 20) is shown for each basin. The

lower panel indicates the key taxa whose abundance characterizes difference among the bottoms

of the basins.

713   **Figure supplement 1.** Energy landscape of fungal communities (family-level compositions;
714   threshold for binarization = 0.001; occurrence threshold = 0.10; $S = 42$).

Figure 6. Disconnectivity graphs of the energy landscapes. (**A**) Schema of a disconnectivity graph. The energy of the "tipping points" splitting basins of attraction are presented across the axis of 2^S possible community states, where S denotes the number of the species or taxa examined. The energy of the bottom of each basin is shown. (**B**) Tipping points and basins on the energy landscape of prokaryotes. The major basins of attraction with ≥ 10 samples with plant-disease information are highlighted with the colors defined in Figure 4. (**C**) Tipping points and basins on the energy landscape of fungi. The major basins of attraction with ≥ 10 samples with plant-disease information are highlighted with the colors defined in Figure 5.