Stable states and keystone species in soil microbiomes of agroecosystems

Hiroaki Fujita[1], Shigenobu Yoshida[2], Kenta Suzuki[3] and Hirokazu Toju[1†]

[1]Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2133, Japan

[2]Institute for Plant Protection, National Agriculture and Food Research Organization, Tsukuba, Ibaraki 305-8666, Japan

[3]Integrated Bioresource Information Division, BioResource Research Center, RIKEN, Tsukuba, Ibaraki 305-0074, Japan

[†]**Correspondence:** Hirokazu Toju (toju.hirokazu.4c@kyoto-u.ac.jp).

This article includes 6 Figures, 7 Extended Data Figures, 2 Supplementary Tables and 1 Source Data.

Abstract

Archaea, bacteria, and fungi in the soil are increasingly recognized as major determinants of

agricultural productivity and sustainability. Although our knowledge of soil biodiversity has

been greatly enhanced for the last decade, defining key characteristics of stable microbiome

structure and their ecosystem functions remains a major challenge. We here show that "stable

states" of soil microbiomes can be defined with their disease-suppressive effects on crop

plants. We integrated agricultural field metadata with microbial community analyses by

targeting > 2,000 agricultural soil samples collected across the Japanese Archipelago,

profiling microbiomes constituted, in total, by 579 archaeal, 26,640 bacterial, and 6,306

fungal species/strains. A DNA-metabarcoding-based evaluation of balance between

prokaryote and fungal abundance indicated that fungus-dominated soil is more resistant to

crop plant disease. A statistical physics approach for reconstructing the "energy landscape" of

community structure indicated that disease-suppressive and disease-susceptible soils could be

distinguished as alternative stable states of fungal communities. A network theoretical

analysis further indicated that the entire microbiomes were partitioned into several

substructures differing in potential impacts on crop disease level. We then highlighted

potential keystone prokaryotes and fungi governing ecological processes within/across such

substructures. These results suggest that agroecosystem performance and sustainability can be

managed by identifying key structure and species within microbiomes.

The ongoing global-scale degradation of agroecosystems is threatening food production[1–3].

Maximizing the functions of microbial communities is a prerequisite for building bases of

sustainable agriculture[4–9]. Archaea, bacteria, and fungi not only promote soil nutrient cycles

but also protect crop plants from pests/pathogens and abiotic stresses[10–14]. Therefore,

diagnosing soil microbiome states is of central interest when we try to design and manage

resource-efficient and disease-tolerant agroecosystems.

Despite the promise that informatics of microbiomes renovates agricultural practices,

defining favorable and unfavorable states of microbial communities[4,15,16] is notoriously

difficult due to tremendous biodiversity in soil ecosystems. Microbiome compositions in the

wild are consequences of multiple ecological processes[17–22]. Filtering by environmental

conditions, for example, is a primary mechanism determining establishment of microbial

populations[21,23,24]. Furthermore, competitive and facilitative interactions between microbial

species are essential factors governing local community structure[25–27]. Continually influenced

by these mechanisms and factors, local microbial communities differ in their stability and

functions[16,28–30]. Consequently, mapping "alternative stable states[17,31–33]" within immensely

complex state space of microbiome structure is an essential step for managing agroecosystem

functions in light of possible community-level consequences with benign/adverse effects.

We here perform systematic profiling of cropland microbiome structure across the

Japanese Archipelago in order to define general characteristics of disease-suppressive and

disease-susceptible soil based on theoretical ecology. We integrated DNA metabarcoding

results of prokaryote and fungal communities with environmental meta-datasets representing

> 2,000 cropland soil samples of 19 plant species. With the bird's-eye view of possible

microbiome structure, we explore backbone characteristics associated with crop disease

conditions. Specifically, we reconstruct the "energy landscape" of microbial community

structure, drastically reducing dimensions for diagnosing disease-suppressive and disease-

susceptible states of agroecosystems. With a network theoretical approach, we further reveal

how soil microbiomes are partitioned into substructures potentially differing in agricultural

performance. Within the reconstructed microbe-to-microbe network, we identify candidates

of keystone species that potentially trigger switches between benign and adverse

agroecosystem states. Overall, this study provides frameworks and criteria for diagnosing

agroecosystem functions based on microbiome structure.

Results

Diversity of microbiome structure

We compiled the field metadata of 2,903 soil samples collected in research projects of

National Agricultural and Food Research Organization (NARO). The rhizosphere/bulk soils

of farmlands were sampled from subtropical to cool-temperate regions (26.1–42.8 ºN) across

the Japanese Archipelago, targeting 19 crop plant species (Fig. 1a) from 2006 to 2014. Most

of the farmlands were managed with conventional agricultural practices (characterized by

intensive tillage and chemical fertilizer/pesticide application), while some were

experimentally controlled as organic agricultural fields (Source Data S1). The metadata

included the information of chemical [e.g., pH, electrical conductivity, carbon/nitrogen (C/N)

ratio, and available phosphorous concentration], physical (e.g., soil taxonomy), and biological

(e.g., crop disease level) properties, providing an invaluable platform for profiling and

categorizing ecosystem states of cropland soil.

To integrate the metadata with the information of microbial community structure, we

performed DNA metabarcoding analyses of both prokaryotes (archaea and bacteria) and

fungi. After a series of quality filtering, prokaryote and fungal community data were

successfully obtained from 2,534 and 2,384 samples, respectively. In total, 579 archaeal

amplicon sequence variants (ASVs) representing 14 genera (11 families), 26,640 bacterial

ASVs representing 875 genera (332 families), and 6,306 fungal archaeal ASVs representing

573 genera (240 families) were detected (Fig. 1b; Extended Data Fig. 1).

The prokaryote communities did not have dominant taxa at the genus and family levels,

although the archaeal family Nitrososphaeraceae had the highest abundance at some research

sites (Fig. 1b). In contrast, the fungal communities were dominated by fungi in the families

Mortierellaceae, Chetomiaceae, and Nectriaceae (Fig. 1b). A reference database profiling of

fungal functional groups suggested that the fungal communities were dominated by plant

pathogenic and soil saprotrophic fungi (Fig. 1c) as characterized by the dominance of

Fusarium and *Mortierella* at the genus level (Extended Data Fig. 1). Meanwhile,

mycoparasitic fungi had exceptionally high proportions at some research sites, as represented

by the dominance of *Trichoderma* (Hypocreaceae) at those sites (Fig. 1b; Extended Data Fig.

1).

The prokaryote and fungal communities differed in basic biodiversity patterns. In terms

of α-diversity at the family level, prokaryote communities showed higher Shannon's entropy

than fungal communities in all but one of the research sites (Fig. 1d). In contrast, variability

of community structure among samples, i.e., Bray-Curtis β-diversity, was higher in fungi than

in prokaryotes in most study sites (Fig. 1e). Likewise, fungi showed higher levels of

community structural difference between research sites than prokaryotes (Welch's test; $t = $ -

6.6, df = 168.9, $P < 0.0001$; Fig. 1f). These results indicate that diverse taxonomic groups of

archaea and bacteria are present without dominance by particular taxa, while fungal

community structure is spatially dynamic both within and across agroecosystems.

Compiling the metadata of edaphic factors, we found that variation in the community

structure of prokaryotes and fungi was significantly explained by crop plant identity and soil

taxonomy as well as by soil chemical properties such as pH, electrical conductivity, C/N ratio,

and available phosphorous concentration (Fig. 2a-b; Extended Data Fig. 2-3; Supplementary

Information). Meanwhile, both prokaryote and fungal community structure was significantly

associated with the severity of crop disease (Fig. 2c-d). Thus, the dataset as a whole offers an

opportunity for understanding mechanisms spanning physical, chemical, and biological

processes in agroecosystems.

Crop disease and microbiome structure

We then focused on factors determining disease-suppressive and disease-susceptible

properties of agroecosystem soil. In addition to chemical environmental factors (e.g., soil pH

and electrical conductivity), whose associations with agricultural productivity have been

routinely examined[34,35], we explored biological factors critical to crop-plant disease level. The

α-diversity of prokaryotes and/or fungi, for example, is an index that can be estimated with

DNA-based profiling of soil biological communities. In further reconstructing biological

factors in a high-throughput manner, we focused on relative abundance of prokaryotes and

fungi. As the classic chromatography-based method for quantifying phospholipid fatty acids

specific to bacteria and fungi[36] could require impractically long time for processing > 2,000

samples, we developed an alternative approach for estimating the ratio of prokaryote

abundance to fungal abundance (hereafter, prokaryote/fungus ratio) based on DNA

metabarcoding. Specifically, the relative concentrations of prokaryote 16S rRNA gene

sequences to fungal ITS gene sequences were calculated based on a quantitative amplicon

sequencing analysis (Fig. 3a; see Methods for details). We then constructed statistical models

comparing effects on crop-plant disease level among various types of chemical/biological

factors.

A series of analysis indicated prokaryote/fungus ratio had strong impacts on crop-plant

disease level. While soil chemistry (soil pH) had significant effects on the proportion of

diseased crop plants, the prokaryote/fungus ratio showed even stronger associations with crop

plant disease level (Model 1 in Fig. 3b). Specifically, crop plant disease was more likely to

occur when prokaryotes were abundant relative to fungi (Fig. 3c). The effects of

prokaryote/fungus ratio on plant disease level were conspicuous even when they were

compared with effects of prokaryote and fungal α-diversity (Model 2 in Fig. 3b). The

threshold values over which more than 50 % of plants showed disease levels 2–5 were

estimated to be around 0.7 in a log-ratio unit (Fig. 3c), although the values potentially vary

depending on various factors such as crop plants, geography, and climatic conditions.

Alternative stables states

We next reconstructed alternative stable states[17,31,33] of microbiomes in order to define

community structure of disease-suppressive and disease-susceptible soil. In an energy

landscape analysis of statistical physics[32], we modeled how microbial community

compositions (community states) depend on soil chemical factors and interactions between

microbial ASVs. The modeling approach allowed us to calculate the "energy" of each

community state. We then estimated the shape of an energy landscape, on which lower energy

represents more stable community state. Stable states were then defined as the community

compositions whose energy is lower than all possible adjacent community compositions[32].

The results of energy landscape reconstruction fundamentally differed between

prokaryotes and fungi. Even if community states were defined in a binary format at the family

or order level to reduce the complexity of the input data (i.e., 2^S possible community states,

where S denotes the number of families; see Methods), more than 10^3 stable states were

inferred to exist for the prokaryote dataset (Fig. 4). This result suggests that ecological

processes of soil prokaryote communities are described as shifts between many equally

possible compositions and that categorization of ecosystem functions based on prokaryote

community structure is challenging. In contrast, the energy landscape of fungal community

structure was represented by much smaller number of alternative stable states (Fig. 4). While

the estimated number of stables states varied depending on threshold values in the

binarization of community structural data, fungal communities consistently showed lower

number of stable states than prokaryote communities (Supplementary Information; Extended

Data Fig. 4).

We then filtered the (alternative) stable states of fungal communities based on the

"depth of basins" and the number of samples belonging to the stable states[32], obtaining 10

major stable states representing the energy landscape (Fig. 5a; see also Extended Data Fig. 5).

By integrating the information of energy landscape (Fig. 5b) with that of crop-plant disease

level (Fig. 5c), we ranked the stable states in terms of possible disease-suppressive functions

(Fig. 5d). We then found that all soil samples belonging to the top-three stable states had

associations with the lowest crop-disease level, while 92 % of samples belonging to the

lowest-rank stable state showed disease levels 2–5.

The 10 stable states were distinguished in light of balance among 22 fungal families

(Fig. 5c). Hypocreaceae, which embraced well-known mycoparasitic taxa (e.g.,

Trichoderma[37]), appeared at high proportions in stable states with potential disease-

suppressive functions. In contrast, Aspergillaceae, which included both plant pathogens and

saprotrophs[38], were relatively abundant in stable states with high disease potential. In addition
to these basic patterns, several families were keys to define disease-suppressive or disease-
susceptible soils. For example, high abundance of Didymellaceae, Lasiosphaeriaceae,
Mrakiaceae, Orbilliaceae, and Piptocephalidaceae characterized the two stable states with the
highest performance (Fig. 3d). Among the families, Orbiliaceae and Piptocephalidaceae were
known as nematophagous and mycoparasitic taxa, respectively[39,40], presumably working as
suppressors of plant-parasitic nematodes and fungi in the soil.

Microbiome networks and keystones

To understand potential mechanisms and factors driving shifts between disease-suppressive
and disease-susceptible microbiomes, we reconstructed the network architecture of microbe-
to-microbe associations. As the number of possible interactions inflates with that of examined
microbes (i.e., X^2 pairs, where X denotes the number of microbial species/ASVs), we targeted
microbial ASVs that appeared in ≥ 10 % of samples in the network statistical analysis. The
reconstructed network of coexistence was subdivided into several "modules", in which
archaeal, bacterial, and fungal ASVs sharing environmental preferences and/or those in
positive interactions were lined with each other (Fig. 6a). We found that the network modules
differed considerably in their associations with crop-plant disease level (Fig. 6a-b; Extended
Data Fig. 6). Modules 3 and 7, for example, were constituted by microbes with potentially
negative impacts on crop plants, while Modules 1, 2, 4, and 5 were characterized by microbes
positively associated with plant disease level (Fig. 6b).

Based on specificity to crop disease status (Extended Data Fig. 6) and topological
positions within the network (Extended Data Fig. 7), candidates of keystone microbes, which
impose profound impacts on ecosystem-level processes, were highlighted. We found that
some ASVs allied to a clade of ammonium-oxidizing archaea[41] (*Nitrocosmicus* and other
Nitrososphaeraceae ASVs) were associated with low disease level of plants and they were
placed at hub positions as indicated by the betweenness centrality index (Fig. 6c). Among
microbes associated with severely damaged plants, an *Chaetomium* fungus had the highest
network centrality (Fig. 6c). When the microbial ASVs were ranked in terms of potential roles
interlinking multiple modules, potentially ammonium-oxidizing archaea in the family
Nitrososphaeraceae[41] and nitrite-oxidizing bacteria in the genus *Nitrospira*[42] exhibited were
highlighted (Fig. 5d). Meanwhile, the above-mentioned *Chaetomium* fungus served as a hub
within Module 7 (Fig. 6d), which consisted of microbes associated with crop disease (Fig.

6b). These microbes occupying key topological positions are prioritized targets of research

because control/manipulation of them can impose cascade effects throughout the network,

triggering shifts between alternative stable states of microbiomes[26].

Discussion

By integrating large datasets of agricultural field properties and DNA metabarcoding, we

found that seemingly diverse microbiome structure could be categorized into several stable

states varying in ecosystem-level performance. Our analysis based on statistical physics

suggested that prokaryote communities had numerous local compositional optima,

presumably due to their rapid and continuous responses to changes in environmental

conditions. In contrast, ecological processes of fungal communities could be interpreted as

shifts between alternative stable states (Fig. 5). Basically, fungi can form complex hyphal

structures and their individual lifespan is much longer than most archaea and bacteria. Given

these biological properties, structure and functions of fungal communities are more likely to

be influenced by legacy effects of past ecological processes[43,44] than those of prokaryotes.

Consequently, agroecosystem functions can be managed by keeping fungal community

structure at targeted stable states with positive impacts on crop production. From the

perspective of pest/pathogen control, increasing the agroecosystem-level resistance to

devastating nematodes and fungi is a pressing issue[45,46]. Our finding that specific taxa of

nematophagous and mycoparasitic fungi are keys to distinguish disease-suppressive and

disease-susceptible soils will help us reduce the use of chemical pesticides/fungicides by

maximizing of ecosystem stability and functions.

The intensive profiling of microbiomes further allowed us to pinpoint potential

keystone species/taxa, which could have disproportionate impacts on ecosystem-level

processes through networks of microbe-to-microbe interactions. The fact that archaea and

bacteria involved in ammonium/nitrite oxidization processes[41,42] were designated as candidate

keystones (Fig. 6) is of particular interest. Control of these archaea/bacteria, for example, can

alter the efficiency or pace of soil nitrogen cycles, resulting in increased/decreased N_2O

emission from crop lands[47,48]. Thus, these potential keystone microbes deserve future

prioritized research, in which we will explore the ground design of sustainable

agroecosystems with efficient nutrient cycles, reduced greenhouse-gas emission, and

enhanced defense against pests/pathogens.

While we herein profiled alternative stable states of microbiomes and potential keystone microbes by focusing mainly on conventional agricultural fields, soil microbiomes with higher functions and stability may be explored in the framework of conservation agriculture[49] or regenerative agriculture[50]. Minimum tillage[51], for example, can decrease damage to fungal hyphae, increasing plants' nutrient uptake through mycorrhizal fungi and suppressing pests/pathogens by the functions of diverse entomophagous, nematophagous, and mycoparasitic fungi. In this respect, the fact that crop disease decreased towards the range of dominance by fungi over prokaryotes (Fig. 3) is of particular interest. Given that the prokaryote/fungus ratio varied considerably among soil samples with similar community compositions (Fig. 3a), the byproduct index resulting from high-throughput DNA metabarcoding is expected to provide a diagnostic criterion of agroecosystem health in lines with stable-state analyses of microbiome structure.

Knowledge of ecosystem-level patterns and processes is indispensable for maximizing biodiversity functions for sustainable food production. Bird's-eye views of stable microbiome compositions and keystone species will provide platforms for integrating ecology, genomics, and microbiology towards the realization of "nature-positive" world[52].

References

1. Springmann, M. *et al.* Options for keeping the food system within environmental limits. *Nature* **562**, 519–525 (2018).

2. Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R. & Polasky, S. Agricultural sustainability and intensive production practices. *Nature* **418**, 671–677 (2002).

3. Tilman, D., Balzer, C., Hill, J. & Befort, B. L. Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences* **108**, 20260–20264 (2011).

4. Toju, H. *et al.* Core microbiomes for sustainable agroecosystems. *Nat Plants* **4**, 247–257 (2018).

5. Busby, P. E. *et al.* Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol* **15**, e2001793 (2017).

6. Berendsen, R. L., Pieterse, C. M. J. & Bakker, P. A. H. M. The rhizosphere microbiome and plant health. *Trends Plant Sci* **17**, 478–486 (2012).

7. Bommarco, R., Kleijn, D. & Potts, S. G. Ecological intensification: Harnessing ecosystem services for food security. *Trends Ecol Evol* **28**, 230–238 (2013).

8. de Vries, F. T. & Wallenstein, M. D. Below-ground connections underlying above-
ground food production: a framework for optimising ecological connections in the
rhizosphere. *Journal of Ecology* **105**, 913–920 (2017).

9. Chaparro, J. M., Sheflin, A. M., Manter, D. K. & Vivanco, J. M. Manipulating the soil
microbiome to increase soil health and plant fertility. *Biol Fertil Soils* **48**, 489–499
(2012).

10. Peay, K. G., Kennedy, P. G. & Talbot, J. M. Dimensions of biodiversity in the Earth
mycobiome. *Nat Rev Microbiol* **14**, 434–447 (2016).

11. Baker, B. J. *et al.* Diversity, ecology and evolution of Archaea. *Nat Microbiol* **5**, 887–
900 (2020).

12. Hayat, R., Ali, S., Amara, U., Khalid, R. & Ahmed, I. Soil beneficial bacteria and their
role in plant growth promotion: A review. *Ann Microbiol* **60**, 579–598 (2010).

13. Newsham, K. K. A meta-analysis of plant responses to dark septate root endophytes.
New Phytol **190**, 783–93 (2011).

14. van der Heijden, M. G. A. *et al.* Mycorrhizal fungal diversity determines plant
biodiversity, ecosystem variability and productivity. *Nature* **396**, 69–72 (1998).

15. Schlatter, D., Kinkel, L., Thomashow, L., Weller, D. & Paulitz, T. Disease suppressive
soils: New insights from the soil microbiome. *Phytopathology* **107**, 1284–1297 (2017).

16. Jansson, J. K. & Hofmockel, K. S. The soil microbiome — from metagenomics to
metaphenomics. *Curr Opin Microbiol* **43**, 162–168 (2018).

17. Fukami, T. Historical contingency in community assembly: integrating niches, species
pools, and priority effects. *Annu Rev Ecol Evol Syst* **46**, 1–23 (2015).

18. Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T. & Singh, B. K. Plant–microbiome
interactions: from community assembly to plant health. *Nat Rev Microbiol* **18**, 607–621
(2020).

19. Jiao, S., Yang, Y., Xu, Y., Zhang, J. & Lu, Y. Balance between community assembly
processes mediates species coexistence in agricultural soil microbiomes across eastern
China. *ISME Journal* **14**, 202–216 (2020).

20. Faust, K., Lahti, L., Gonze, D., de Vos, W. M. & Raes, J. Metagenomics meets time
series analysis: Unraveling microbial community dynamics. *Curr Opin Microbiol* **25**,
56–66 (2015).

21. Tedersoo, L. *et al.* Global diversity and geography of soil fungi. *Science (1979)* **346**,
1256688 (2014).

22. Toju, H., Tanabe, A. S. & Sato, H. Network hubs in root-associated fungal
metacommunities. *Microbiome* **6**, 116 (2018).

23. Talbot, J. M. *et al.* Endemism and functional convergence across the North American
soil mycobiome. *Proceedings of the National Academy of Sciences* **111**, 6341–6346
(2014).

24. Bahram, M. *et al.* Structure and function of the global topsoil microbiome. *Nature* **560**,
233–237 (2018).

25. Faust, K. & Raes, J. Microbial interactions: From networks to models. *Nat Rev*
Microbiol **10**, 538–550 (2012).

26. Toju, H., Yamamoto, S., Tanabe, A. S. A. S., Hayakawa, T. & Ishii, H. S. H. S.
Network modules and hubs in plant-root fungal biomes. *J R Soc Interface* **13**,
20151097–20151097 (2016).

27. van der Heijden, M. G. A. & Hartmann, M. Networking in the plant microbiome. *PLoS*
Biol **14**, 1–9 (2016).

28. Hartman, K. *et al.* Deciphering composition and function of the root microbiome of a
legume plant. *Microbiome* **5**, 1–13 (2017).

29. Vorholt, J. A., Vogel, C., Carlström, C. I. & Müller, D. B. Establishing causality:
opportunities of dynthetic vommunities for plant microbiome tesearch. *Cell Host*
Microbe **22**, 142–155 (2017).

30. Mendes, L. W. *et al.* Soil-borne microbiome: linking diversity to function. *Microb Ecol*
70, 255–265 (2015).

31. Beisner, B. E., Haydon, D. T. & Cuddington, K. Alternative stable states in ecology.
Front Ecol Environ **1**, 376–382 (2003).

32. Suzuki, K., Nakaoka, S., Fukuda, S. & Masuya, H. Energy landscape analysis
elucidates the multistability of ecological communities across environmental gradients.
Ecol Monogr **91**, 1–21 (2021).

33. Scheffer, M., Carpenter, S., Foley, J. a, Folke, C. & Walker, B. Catastrophic shifts in
ecosystems. *Nature* **413**, 591–596 (2001).

34. Goulding, K. W. T. Soil acidification and the importance of liming agricultural soils
with particular reference to the United Kingdom. *Soil Use Manag* **32**, 390–399 (2016).

35. Corwin, D. L. & Lesch, S. M. Apparent soil electrical conductivity measurements in
agriculture. *Comput Electron Agric* **46**, 11–43 (2005).

36. Frostegård, A. & Bååth, E. The use of phospholipid fatty acid analysis to estimate
bacterial and fungal biomass in soil. *Biol Fertil Soils* **22**, 59–65 (1996).

37. Poveda, J. Trichoderma as biocontrol agent against pests: New uses for a mycoparasite.
Biological Control vol. 159 Preprint at
https://doi.org/10.1016/j.biocontrol.2021.104634 (2021).

38. Tekaia, F. & Latgé, J. P. Aspergillus fumigatus: Saprophyte or pathogen? *Current*
Opinion in Microbiology vol. 8 Preprint at https://doi.org/10.1016/j.mib.2005.06.017
(2005).

39. Yang, Y., Yang, E., An, Z. & Liu, X. Evolution of nematode-trapping cells of
predatory fungi of the Orbiliaceae based on evidence from rRNA-encoding DNA and
multiprotein sequences. *Proc Natl Acad Sci U S A* **104**, 8379–8384 (2007).

40. Lazarus, K. L., Benny, G. L., Ho, H. M. & Smith, M. E. Phylogenetic systematics of
syncephalis (zoopagales, zoopagomycotina), a genus of ubiquitous mycoparasites.
Mycologia **109**, 333–349 (2017).

41. Zhalnina, K. v. *et al.* Genome sequence of Candidatus nitrososphaera evergladensis
from group I.1b enriched from everglades soil reveals novel genomic features of the
ammonia-oxidizing archaea. *PLoS One* **9**, e101648 (2014).

42. Lücker, S. *et al.* A Nitrospira metagenome illuminates the physiology and evolution of
globally important nitrite-oxidizing bacteria. *Proc Natl Acad Sci U S A* **107**, 13479–84
(2010).

43. Kennedy, P. G., Peay, K. G. & Bruns, T. D. Root tip competition among
ectomycorrhizal fungi: are priority effects a rule or an exception? *Ecology* **90**, 2098–
107 (2009).

44. Werner, G. D. A. & Kiers, E. T. Order of arrival structures arbuscular mycorrhizal
colonization of plants. *New Phytologist* **205**, 1515–1524 (2015).

45. Callaway, E. Devastating wheat fungus appears in Asia for first time. *Nature* vol. 532
421–422 Preprint at https://doi.org/10.1038/532421a (2016).

46. Onkendi, E. M., Kariuki, G. M., Marais, M. & Moleleki, L. N. The threat of root-knot
nematodes (Meloidogyne spp.) in Africa: A review. *Plant Pathology* vol. 63 727–737
Preprint at https://doi.org/10.1111/ppa.12202 (2014).

47. Tian, H. *et al.* A comprehensive quantification of global nitrous oxide sources and
sinks. *Nature* **586**, 248–256 (2020).

48. Lugato, E., Leip, A. & Jones, A. Mitigation potential of soil carbon management
overestimated by neglecting N2O emissions. *Nat Clim Chang* **8**, 219–223 (2018).

49. Kassam, A., Friedrich, T. & Derpsch, R. Global spread of conservation agriculture.
International Journal of Environmental Studies **76**, 29–51 (2019).

50. Schreefel, L., Schulte, R. P. O., de Boer, I. J. M., Schrijver, A. P. & van Zanten, H. H.
E. Regenerative agriculture – the soil is the base. *Glob Food Sec* **26**,
https://doi.org/10.1016/j.gfs.2020.100404 (2020).

51.   Li, Y., Zhang, Q., Cai, Y., Yang, Q. & Chang, S. X. Minimum tillage and residue
retention increase soil microbial population size and diversity: Implications for
conservation tillage. *Science of the Total Environment* **716**, 137164 (2020).

52.   UNDP, SCBD & UNEP-WCMC. *Creating a nature-positive future: the contribution of*
protected areas and other effective area-based conservation measures. (2021).

53.   Caporaso, J. G. *et al.* Global patterns of 16S rRNA diversity at a depth of millions of
sequences per sample. *Proc Natl Acad Sci U S A* **108**, 4516–4522 (2011).

54.   Apprill, A., Mcnally, S., Parsons, R. & Weber, L. Minor revision to V4 region SSU
rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton.
Aquatic Microbial Ecology **75**, 129–137 (2015).

55.   Lundberg, D. S., Yourstone, S., Mieczkowski, P., Jones, C. D. & Dangl, J. L. Practical
innovations for high-throughput amplicon sequencing. *Nat Methods* **10**, 999–1002
(2013).

56.   Toju, H., Kurokawa, H. & Kenta, T. Factors influencing leaf- and root-associated
communities of bacteria and fungi across 33 plant orders in a grassland. *Front*
Microbiol **10**, 1–14 (2019).

57.   Stevens, J. L., Jackson, R. L. & Olson, J. B. Slowing PCR ramp speed reduces chimera
formation from environmental samples. *J Microbiol Methods* **93**, 203–205 (2013).

58.   Ushio, M. Interaction capacity as a potential driver of community diversity.
Proceedings of the Royal Society B: Biological Sciences **289**, 20212690 (2022).

59.   Ushio, M. *et al.* Quantitative monitoring of multispecies fish environmental DNA
using high-throughput sequencing. *Metabarcoding Metagenom* **2**, 1–15 (2018).

60.   Fujita, H. *et al.* Alternative stable states, nonlinear behavior, and predictability of
microbiome dynamics. *bioRxiv*.

61.   Toju, H., Tanabe, A. S. A., Yamamoto, S. & Sato, H. High-coverage ITS primers for
the DNA-based identification of ascomycetes and basidiomycetes in environmental
samples. *PLoS One* **7**, e40863 (2012).

62.   Hamady, M., Walker, J. J., Harris, J. K., Gold, N. J. & Knight, R. Error-correcting
barcoded primers for pyrosequencing hundreds of samples in multiplex. *Nat Methods*
5, 235–237 (2008).

63.   Tanabe, A. S. & Toju, H. Two new computational methods for universal DNA
barcoding: a benchmark using barcode sequences of bacteria, archaea, animals, fungi,
and land plants. *PLoS One* **8**, e76910 (2013).

64.   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina
amplicon data. *Nat Methods* **13**, 581–583 (2016).

65.   R Core Team. R: A language and environment for statistical computing. *R: A language*
419   *and environment for statistical computing. R Foundation for Statistical Computing,*
420   *Vienna, Austria. URL https://www.R-project.org/* (2020).

421   66.   Kurtz, Z. D. *et al.* Sparse and compositionally robust inference of microbial ecological
422   networks. *PLoS Comput Biol* **11**, e1004226 (2015).

423   67.   Põlme, S. *et al.* FungalTraits: a user-friendly traits database of fungi and fungus-like
424   stramenopiles. *Fungal Divers* **105**, 1–16 (2020).

425   68.   Watanabe, T., Masuda, N., Megumi, F., Kanai, R. & Rees, G. Energy landscape and
426   dynamics of brain activity during human bistable perception. *Nat Commun* **5**, 4765
427   (2014).

428   69.   Wood, S. mgcv: Mixed GAM computation vehicle with automatic smoothness
429   estimation available at https://cran.r-project.org/web/packages/mgcv/index.html.
430   (2022).

431   70.   Newman, M. & Girvan, M. Finding and evaluating community structure in networks.
432   *Phys Rev E* **69**, 026113 (2004).

433   71.   Csardi, G. & Nepusz, T. The igraph software package for complex network research.
434   *InterJournal Complex Systems* **1695**, 1–9 (2006).

435   72.   Freeman, L. C. A set of measures of centrality based on betweenness. *Sociometry* **40**,
436   35–41 (1977).

437   73.   Olesen, J. M., Bascompte, J., Dupont, Y. L. & Jordano, P. The modularity of
438   pollination networks. *Proc Natl Acad Sci U S A* **104**, 19891–19869 (2007).

439   74.   Bonacich, P. Factoring and weighting approaches to status scores and clique
440   identification. *J Math Sociol* **2**, 113–120 (1972).

Methods

Soil samples and metadata

Over research projects of National Agricultural and Food Research Organization (NARO), which were carried out through five national research programs funded by Ministry of Agriculture, Forestry and Fisheries, 2,903 rhizosphere/bulk soil samples were collected from conventional agricultural fields across the Japanese Archipelago from January 23, 2006, to July 28, 2014 (Source Data S1). When the latitude and longitude of the sampling positions were round to one decimal place, 42 research sites were identified. Across the metadata of the 2,903 samples, the information of 19 crop plants, 34 former crop plants (including "no crop"), 13 soil taxonomic groups (e.g., "Andosol"), 60 experimental/research purposes (e.g., "soil comparison between organic and conventional management") was described. Likewise, the metadata included the information of dry soil pH, electrical conductivity, carbon/nitrogen (C/N) ratio, and available phosphorous concentration from, 2,830, 2,610, 2,346, and 2,249 samples, respectively. In addition, the information of the severity of crop plant disease was available for 1,472 samples. The values of the proportion of diseased plants or disease severity index was normalized within the ranges from 0 to 100, and they were then categorized into five levels (level 1, 0-20; level 2, 20-40; level3; 40-60; level 4, 60-80; level 5, 80-100). The plant pathogens examined in the disease-level evaluation were *Colletotrichum gloeosporioides* on the strawberry, *Fusarium oxysporum* on the celery, the lettuce, the strawberry, and the tomato, *Phytopthora sojyae* on the soybean, *Plasmodiophora brassicae* on Cruciferae plants, *Pyrenochaeta lycopersici* on the tomato, *Pythium myriotylum* on the ginger, *Ralstonia solanacearum* on the eggplant and the tomato, and *Verticillium* spp. on Chinese cabbage. For continuous variables within the metadata, emergent outliers (mean + 5 SD) were converted into "NA" in the data matrix used in the following statistical analyses as potential measurement/recording errors. Unrealistic electrical conductivity records (> 20) were converted into "NA" as well.

For the collection of each sample, ca. 500 g of soil was mixed and sieved with 2-mm mesh in the field. The samples were stored at -20 ºC until DNA extraction. In laboratory conditions, 0.4 g of soil (fresh weight) was subjected to DNA extraction with FastDNA SPIN Kit for Soil (Q-BioGene).

DNA amplification and sequencing

Profiling of soil microbial biodiversity was performed by targeting archaea, bacteria, and

fungi. For the amplification of the 16S rRNA V4 region of archaea and bacteria (prokaryotes),

the set of the forward primer 515f[53] and the reverse primer 806rB[54] were used. The primers

were fused with 3–6-mer Ns for improved Illumina sequencing quality[55] and Illumina

sequencing primers as detailed in a previous study[56]. PCR was performed using KOD ONE

PCR Master Mix (TOYOBO, Osaka) with the temperature profile of 35 cycles at 98 ºC for 10

seconds (denaturation), 55 ºC for 5 seconds (annealing of primers), and 68 ºC for 30 seconds

(extension), and a final extension at 68 ºC for 2 minutes. The ramp rate through the thermal

cycles was set to 1 ºC/sec to prevent the generation of chimeric sequences[57]. In the PCR, we

added five artificial DNA sequence variants with different concentrations (i.e., standard DNA

gradients; 1.0×10^{-4}, 5.0×10^{-5}, 2.0×10^{-5}, 1.0×10^{-5}, and 5.0×10^{-6} nM) to the PCR master mix

solution in order to calibrate 16S rRNA gene copy numbers in the template DNA samples as

detailed elseshere[58–60]. Likewise, the ITS1 region of fungi was amplified using the set of the

forward primer ITS1F_KYO1 and the reverse primer ITS2_KYO2[61]. PCR was performed

using the Illumina-sequencing fusion primer design mentioned above with the temperature

profile of 35 cycles at 98 ºC for 10 seconds, 53 ºC for 5 s seconds, and 68 ºC for 5 seconds,

and a final extension at 68 ºC for 2 minutes (ramp rate = 1 ºC/sec). Newly designed artificial

sequence variants (1.0×10^{-5}, 7.0×10^{-6}, 5.0×10^{-6}, 2.0×10^{-6}, and 1.0×10^{-6} nM; Supplementary

Information) were added to the PCR master mix as standard DNA gradients for the

calibration of the ITS sequence concentrations in the template DNA samples.

The PCR products of the prokaryiote 16S rRNA and fungal ITS1 regions were

respectively subjected to the additional PCR step for linking Illumina sequencing adaptors

and 8-mer sample identifier indexes[62] with the amplicons as detailed elsewhere[56]. The

temperature profile in the PCR was 8 cycles at 98 ºC for 10 seconds, 55 ºC for 5 seconds, and

68 ºC for 5 seconds, and a final extension at 68 ºC for 2 minutes. The PCR products were then

pooled for each of the 16S rRNA and fungal ITS1 regions after a purification/equalization

process with the AMPureXP Kit (Beckman Coulter, Inc., Brea). Primer dimers, which were

shorter than 200 bp, were removed from the pooled library by supplemental purification with

AMpureXP: the ratio of AMPureXP reagent to the pooled library was set to 0.8 (v/v) in this

process. The sequencing libraries of the two regions were processed in an Illumina MiSeq

sequencer (10% PhiX spike-in). Because the quality of forward sequences is generally higher

than that of reverse sequences in Illumina sequencing, we optimized the MiSeq run setting in

order to use only forward sequences. Specifically, the run length was set 271 forward (R1)

and 31 reverse (R4) cycles to enhance forward sequencing data: the reverse sequences were

used only for discriminating between prokaryote 16S and fungal ITS1 sequences in the

following bioinformatic pipeline.

Bioinformatics

In total, 23,573,405 sequencing reads were obtained in the Illumina sequencing (16S rRNA,

11,647,166 sequencing reads; ITS, 11,926,239 sequencing reads). The raw sequencing data

were converted into FASTQ files using the program bcl2fastq 1.8.4 distributed by Illumina.

For each of the 16S rRNA and fungal ITS1 regions, the output FASTQ files were

demultiplexed using Claident v0.9.2022.01.26[63]. The sequencing data were deposited to DNA

Data Bank of Japan (DDBJ) (Bioproject accession no.: PSUB018361). The removal of low-

quality sequences and ASV inferences were done using DADA2[64] v1.17.5 of R v3.6.3[65]. The

mean number of filtered sequencing reads obtained per sample was 3,949 and 4,075 for the

prokaryote and fungal, datasets, respectively. Taxonomic annotation of prokaryotes was

conducted based on the SILVA 138 SSU (Quast et al., 2013) the assignTaxonomy function of

DADA2. For the taxonomic annotation of fungi, the QCauto method[63] was applied to the

NCBI nucleotide sequence database bundled with Claident v0.9.2022.01.26. ASVs that were

not assigned to the domain Archaea/Bacteria and the kingdom Fungi were removed from the

16S rRNA and ITS1 datasets, respectively. For each target organismal group (prokaryotes and

fungi), we then obtained a sample × ASV matrix, in which a cell entry depicted the number of

sequencing reads of an ASV in a sample. The samples with less than 1,000 reads were

discarded from the matrices. The matrix was rarefied to 1,000 reads per sample for the

downstream statistical pipeline except for the network analysis, which had an original

standardization protocol of input data[66]. In total, prokaryote and fungal community data were

obtained for 2,534 and 2,384 samples, respectively. The number of ASVs included in the

rarefied matrices were 27,219 and 6,306 for prokaryotes and fungi, respectively. For fungal

ASVs, putative functional groups (e.g., "plant pathogen") were inferred using the program

FungalTraits[67]. The estimation of DNA concentrations of the prokaryote 16S rRNA and

fungal ITS regions was performed, respectively, based on the calibration with the standard

DNA gradients (artificial DNA variants introduced to the PCR master mix solutions) as

detailed elsewhere[60].

Diversity of microbiome structure

For each sample, Shannon's α-diversity estimates of prokaryotes and fungi were respectively

calculated at the family level. Likewise, mean Bray-Curtis dissimilarity (β-diversity) in

family-level compositions within each of the 42 research sites was calculated as a measure of

within-site variability of prokaryote/fungal community structure. In addition, the degree of

among-site community differentiation was evaluated for prokaryotes and fungi. Specifically,

after calculating mean family-level compositions at each research site, Bray-Curtis β-diversity

was calculated in all pairs of research sites. Those α-diversity, within-site β-diversity, and

among-site β-diversity estimates were respectively compared between prokaryotes and fungi.

For each of the prokaryote and fungal datasets, permutational analyses of variance

(PERMANOVA) were performed to examine associations between family-level community

compositions and variables in the metadata. Prior to the PEMANOVA, Bray-Curtis β-

diversity was calculated based on the rarefied sample × ASV matrices. Two types of

PEMANOVA models were constructed (10,000 iterations). Specifically, one is constituted by

categorical explanatory variables (crop plant, former crop plant, soil taxonomy, research site,

and sampling date), while the other included continuous explanatory variables (soil pH,

electrical conductivity, C/N ratio, and available phosphorous concentration, latitude, and

longitude).

To reduce the dimensions of the community composition data, a principal coordinate

analysis (PCoA) was performed based on the Bray-Curtis β-diversity calculated with the

rarefied sample × ASV matrix of prokaryotes or fungi. For each PCoA axis (axes 1 to 5) in

each of the prokaryote and fungal analyses, Pearson's correlation with each chemical

environmental factor (soil pH, electrical conductivity, C/N ratio, and available phosphorous

concentration) was calculated.

Crop disease and microbiome structure

We then evaluated how community structure of prokaryotes and fungi are associated with

crop disease. For each of the prokaryote and fungal data sets, a generalized linear mixed

model (GLMM) of crop-disease level was constructed by included the PCoA axes 1 to 5 as

fixed effects and experimental/research purposes in the metadata as random effects. A logit-

link function and binomial error was assumed in the GLMM after converting the response

variable into a binary format [disease level 1 (= 1) vs. disease levels 2–5 (= 0)].

We also examined how balance between prokaryote and fungal abundance could be

associated with crop disease levels. Based on the estimated concentrations of prokaryote 16S

rRNA and fungal ITS sequences in template DNA solutions, we calculated the ratio of

prokaryote DNA concentrations to fungal DNA concentrations in respective samples

(prokaryote/fungus ratio) as follows:

$log \dfrac{\text{prokaryote 16S rRNA gene concentration(DNA copies/μL)}}{\text{fungal ITS gene concentration (DNA copies/μL)}}$.

Although potential variation in DNA extraction skills of researchers might affect absolute

DNA concentrations in the template DNA solutions, balance between prokaryote and fungal

DNA in each template DNA sample could be used as a reliable measure.

Potential effects of the prokaryote/fungus ratio on crop-disease level were examined in

a GLMM with a logit-link function and binomial errors. In the GLMM, soil pH and electrical

conductivity were included as fixed effects in addition to the prokaryote/fungus ratio. In the

GLMM, C/N ratio and available phosphorous concentration were not included due to the

small number of samples with both crop-disease information and the chemical variables. The

experimental/research purpose information in the metadata was included as random effects in

the GLMM. To examine the effects of other aspects of biological factors, we constructed

another GLMM including the ASV-level Shannon diversity of prokaryotes, that of fungi, and

the prokaryote/fungus ratio. All the explanatory variables were z-standardized (zero-mean and

unit-variance) before the GLMM analysis. To visualize how the proportion of least-diseased

(disease-level 1) samples varied along the axis of the prokaryote/fungus ratio, we performed a

supplementary analysis with generalized linear models (GLMs). Specifically, for the five

experimental/research purpose categories from which 10 or more disease-level 1 (least-

diseased) samples and 10 or more disease-level 2–5 (severely diseased) samples were

obtained, a GLM of disease level was constructed by including the prokaryote/fungus ratio as

an explanatory variable.

Alternative stables states

599   We estimated the alternative stable states of microbial community structure based on the

600   framework of an energy landscape analysis[32,68] (tutorials of energy landscape analyses are

601   available at https://community.wolfram.com/groups/-/m/t/2358581). Probabilities of

602   community states $p(\vec{\sigma}^{(k)})$ are given by

$$P\left(\vec{\sigma}^{(k)}|\varepsilon\right) = e^{-H\left(\vec{\sigma}^{(k)},\varepsilon\right)}/Z$$

$$Z = \sum_{i=1}^{2^S} e^{-H\left(\vec{\sigma}^{(i)},\varepsilon\right)},$$

where $\vec{\sigma}^{(k)} = (\sigma_1^{(k)}, \sigma_2^{(k)}, \dots, \sigma_S^{(k)})$ is a community state vector of k-th community state and S

is the total number of taxa (e.g., ASVs, species, genera, or families) examined. $\varepsilon =$

$(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_M)$ is an array of continuous values representing environmental factors (e.g., soil

pH and electrical conductivity) and M is the total number of environmental parameters. $\sigma_i^{(k)}$

is a binary variable that represents presence (1) or absence (0) of taxon i: i.e., there are a total

of 2^S community states. As the exploration of the 2^S community states were

computationally intensive, we coded community states at the family level taxonomic

compositions. Specifically, families that represented 10 % or more of the total sequencing

reads of a target sample were coded as 1, while those representing less than 10 % of total

reads were coded as 0. Then, the energy of the community state $\vec{\sigma}^{(k)}$ is given by the

extended pairwise maximum entropy model:

$\quad H(\vec{\sigma}^{(k)}, \varepsilon) = -\sum_{i=1}^S h_i \vec{\sigma}_i^{(k)} - \sum_{j=1}^S \sum_{i=1}^M g_{ij} \varepsilon_i^{(k)} \sigma_j^{(k)} - \sum_{i=1}^S \sum_{j=1}^S J_{ij} \vec{\sigma}_i^{(k)} \vec{\sigma}_j^{(k)}/2,$

where h_i represents the net effect of implicit abiotic factors, by which i-th taxon is more

likely to present ($h_i > 0$) or not ($h_i < 0$), g_{ij} represents the effect of the i-th observed

environmental factor, and J_{ij} represents a co-occurrence pattern of i-th and j-th taxa. Since

the logarithm of the probability of a community state is inversely proportional to $H(\vec{\sigma}^{(k)})$, a

community state having lower H is more frequently observed. To consider dynamics on an

assembly graph defined as a network whose 2^S nodes represent possible community states

and the edges represents transition path between them (two community states are adjacent

only if they have the opposite presence/absence status for just one species), we assigned

energy to nodes with the above equation, and so imposed the directionality in state transitions.

Then, by using the steepest descent algorithm[32] we identified the stable state communities as

the energy minima of the weighted network (nodes having the lowest energy compared to all

its neighbors), and determined their basins of attraction based on a steepest descent procedure

starting from each node. As environmental variables, soil pH, electrical conductivity, C/N

ratio, and available phosphorous concentration were included after normalization within the

ranges from 0 to 1.

The number of stable states within energy landscapes of community structure was

inferred, respectively, for three types of datasets, namely, the prokaryote community matrix,

the fungal matrix, and the matrix including both prokaryotes and fungi. In addition to the

energy landscape analyses based on the abovementioned family-level delineation of

community states, analyses based on community-state delineation at the order-level were

performed. Likewise, additional analyses based on another threshold value (5 %) in the

binarization of community states were conducted. Consequently, we performed energy

landscape analyses in each of the combinations of three types of data (prokaryotes, fungi, and

prokaryotes + fungi), two taxonomic levels (family or order), and two threshold levels (5 %

and 10 %). For each combination (setting), the parameters of the extended pairwise maximum

entropy model were adjusted to fit the model to the empirical data. The number of stable

states was then estimated by using the gradient descent algorithm for 20,000 randomly

generated community states within the 2^X possible community states. In the heuristic search,

sets of values of the environmental variables (soil pH, electrical conductivity, C/N ratio, and

available phosphorous concentration) were drawn from those of sampling points.

We then reconstructed energy landscapes for the settings with which community

structure could be characterized by relatively small numbers of stable states (i.e., the family-

level fungal data matrix defined at the 10-% threshold and the order-level fungal data matrix

defined at the 10-% threshold; Fig. 4; Extended Data Fig. 4). Minor stable states whose

shallowest "energy barrier" (i.e., deviation of energy from adjacent "tipping points") was less

than 1/10 of that of the deepest alternative stables states were incorporated into the basins of

deeper stable states (i.e., pruning of alternative stable states[32]). The "energy" estimates were

then plotted against the PCoA axes mentioned above. Spline smoothing of the landscape was

performed with optimized penalty scores using the mgcv v.1.8-40 package[69] of R.

We then showed the energy of each sample on the two-dimensional surface of the state

space inferred with the above-mentioned PCoA: the horizontal axes indicated PCoA 1 and 2

scores and the vertical axis indicated the energy of target samples' community states. On the

energy landscape, samples belonging to the same alternative stable states were indicated by

colors. For the alternative stable states that represented 10 or more samples, the proportions of

disease-level-1 samples were calculated.

Microbiome network modules

To infer potential interactions between microbial ASVs, the algorithm of sparse inverse

covariance estimation for ecological association inference (SPIEC-EASI) was applied based

on the Meinshausen-Bühlmann (MB) method as implemented in the SpiecEasi package[66] of

R. The network inference based on co-occurrence patterns allowed us to detect pairs of

microbial ASVs that potentially interact with each other in facilitative ways and/or those that

might share ecological niches (e.g., preference for edaphic factors). Because estimation of co-

occurrence patterns was not feasible for rare nodes, the prokaryote and fungal ASVs that

appeared in more than 10 % of the sequenced samples were included in the input matrix of

the network analysis. Network modules, within which closely associated ASVs were

interlinked with each other, were identified with the algorithm based on edge betweenness[70]

using the igraph package[71] or R.

We then characterized the inferred network modules based on their potential impacts on

crop-plant disease level. For each microbial ASV constituting the modules, we evaluated

specificity of occurrences in samples differing in crop disease levels based on a

randomization analysis. For the calculation, the rarefied matrices of prokaryotes and fungi

were combined. Within the combined sample × ASV matrix, samples were categorized into

the five disease levels. Mean read counts across samples displaying each of the five disease

levels were then calculated for each ASV. Meanwhile, mean read counts for respective

disease levels were calculated as well for randomized matrices, in which disease labels of the

samples were shuffled (10,000 permutations). For i-th ASV, standardized specificity to

disease level j (s_{ij}) was obtained as follows:

$$s_{ij} = \frac{O_{ij} - Mean(R_{ij})}{SD(R_{ij})},$$

where O_{ij} and R_{ij} is the mean read counts of i-th ASV across disease-level-j samples in the

observed and randomized matrices, respectively, and $Mean(R_{ij})$ and $SD(R_{ij})$ indicate mean

and standard deviation across the randomized matrices. An s_{ij} value higher than 4 or that

lower than -4 indicates statistically strong specificity of the i-th ASV to the disease level j

(Extended Data Fig. 6). For each module in the inferred co-occurrence network, we then

calculated mean standardized specificity to disease level 1 across constituent ASVs.

Network hubs

To explore potential keystone microbes within the network, we scored respective ASVs on

the basis of their topological positions. Among the indices used for evaluating ASVs,

betweenness centrality[72], which measures the extent to which a given nodes (ASV) is located

within the shortest paths connecting pairs of other nodes in a network, is commonly used to

find hubs mediating flow of effects in a network. The network centrality scores were

normalized as implemented in the igraph packages of R[71]. In addition, by focusing on the

above-mentioned network modules, we ranked ASVs based on their within-module degree

and among-module connectivity[73]. The former index is obtained as the number of nodes

linked with a target node within a target network module, suggesting the topological

importance of a node within the module it belongs to. The latter index represents the extent to

which a node interlinks nodes belonging to different network modules. Within-module degree

was z-standardized (i.e., zero-mean and unit-variance) within each module, while among-

module connectivity was defined to vary between 0 to 1. In addition to those indices for

evaluating topological roles within a network, eigenvector centrality[74] was calculated for

respective nodes.

Data availability

711 The 16S rRNA and ITS sequencing data are available from the DNA Data Bank of Japan

712 (DDBJ: Bioproject: PSUB018361). The microbial community data are deposited at our

713 GitHub repository (https://github.com/hiro-toju/Soil_Microbiome_NARO3000).

Code availability

All the R scripts used to analyze the data are available at the GitHub repository

(https://github.com/hiro-toju/Soil_Microbiome_NARO3000).

Acknowledgements. We thank the SuperComputer System, Institute for Chemical Research,

Kyoto University for the use of super computers. This work was financially supported by JST

PRESTO (JPMJPR16Q6), JST FOREST (JPMJFR2048), Human Frontier Science Program

(RGP0029/2019), JSPS Grant-in-Aid for Scientific Research (20K20586), NEDO Moonshot

Research and Development Program (JPNP18016), and JST FOREST (JPMJFR2048) to

H.T., JSPS Grant-in-Aid for Scientific Research (20K06820 and 20H03010) to K.S., and

JSPS Fellowship to H.F..

Author Contributions. H.F., S.Y. and H.T. designed the work. H.F. performed molecular

experiments. H.F. and H.T. analyzed the data. H.T. wrote the paper with H.F., S.Y., and K.S..

Competing Interests. The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at xxxx [URL to be supplied by the

publisher].

Correspondence and requests for materials should be addressed to

toju.hirokazu.4c@kyoto-u.ac.jp.

Fig. 1 | Diversity of agroecosystem microbiomes. **a**, Map of research sites across the

Japanese Archipelago. The 2,903 soil samples were grouped into 42 research sits were when

their latitude and longitude profiles were round to one decimal place. **b**, Taxonomic

compositions of prokaryotes (archaea and bacteria; top) and fungi (bottom) at the family

level. See Extended Data Fig. 1 for results at the genus, order, and class levels. **c**,

Compositions of functional groups of fungi. **d**, Comparison of α-diversity between

prokaryotes and fungi. Mean Shannon diversity at the family level was shown for each

research site. Bars represent standard deviations. **e**, Mean community dissimilarity (β-

diversity) between pairs of samples within each research site. **f**, Community dissimilarity (β-

diversity) in mean community compositions between pairs of research sites.

Fig. 2 | Dimensions of soil microbiome structure. **a**, Principal co-ordinate analysis (PCoA)
of prokaryote and fungal community structure. The identify of crop plants is shown by colors.
See Extended Data Figs. 2-3 for relationship between community structure and environmental
factors. **b**, Correlation between PCoA scores and soil environmental factors. **c**, Microbiome
structure and crop disease levels. **d**, Relationship between prokaryote/fungal community
structure on the disease level of crop plants. A GLMM of crop plants' disease level (disease
level 1 vs. disease levels 2-5) with a logit-link function and binomial errors was constructed
by setting principal coordinates of prokaryote/fungal community structure as explanatory
variables (fixed effects). The identity of research projects was included as random effects in
the GLMM.

Fig. 3 | Balance between prokaryote and fungal abundance. **a**, Prokaryote/fungus ratio.

The relative abundance of prokaryotes and fungi was evaluated based on estimated DNA

concentrations ($log \frac{\text{prokaryote 16S rRNA gene concentration(DNA copies/μL)}}{\text{fungal ITS gene concentration (DNA copies/μL)}}$). The prokaryote/fungus

ratio is shown for each sample on the PCoA surfaces of prokaryote and fungus community

compositions. **b**, Effects of prokaryote/fungus ratio on the disease level of crop plants. In each

GLMM with a logit-link function and binomial errors, crop plants disease level [least-

diseased (disease level 1) = 1, severely-diseased (disease level 2-5) = 0] was included as

response variable. In addition to bacterium/fungal abundance ratio, soil environmental

variables (pH and electrical conductivity; Model 1) and α-diversity estimates of prokaryotes

and fungi (Model 2) were included as explanatory variables (fixed effects). The identity of

research projects was included as random effects in the GLMM. A negative estimate indicates

that the proportion of least-diseased plants decreased with increasing values of a target

explanatory variable. **c**, Representative relationship between prokaryote/fungus ratio on crop

disease. For each of the five research projects with enough number of disese-level-1 and

disease-level-2-5 samples, a GLM was constructed. For the two research projects with

significant effects of prokaryote/fungus ratio on crop disease level, the prokaryote/fungus

ratios at which 50 % of samples display disease-level-1 symptoms are shown.

Fig. 4 | **Number of the stable states of community structure**. The parameters of the extended pairwise maximum entropy model were adjusted to fit the model to the empirical data (see Methods). The number of stable states in the energy landscape was then estimated by using the gradient descent algorithm for 20,000 randomly generated community states within the 2^S possible community states, where S denotes the number of families or orders. In the heuristic search, sets of values of the environmental variables (soil pH, electrical conductivity, C/N ratio, and available phosphorous concentration) were drawn from those of sampling points. The boxplots show variation in the number of stable states across different sets of environmental variables. A box represents the first quartile, median and the third quartile, while bars and circles indicate 1.5 × inter-quartile range and outliers, respectively. The analysis was performed, respectively, for the dataset representing both prokaryotes and fungi, the prokaryote dataset, and the fungus dataset. The threshold value in the binarization of the community data was set at 10 %: see Extended Data Figure 4 for results based on another threshold value (5 %).

Fig. 5 | Alternative stable states of microbiomes. a, Energy landscape of fungal community

structure. The surface of energy levels was reconstructed on the PCoA dimensions (Fig. 2) of

fungal community structure based on spline smoothing. Community states with lower energy

are inferred to be more stable. Data points (samples) indicated by the same color belong to the

basins of the same alternative stable states, whose energy is lower than any adjacent

community structure. See panel **c** for details about the alternative stable states. **b**, Surface of

crop disease conditions. The area of community structure with less diseased states and lower

energy (i.e., community stability; panel **a**) are expected to represent stably-manageable

microbiomes with potential disease-suppressive functions. **c**, Definitions of the stable states.

To reduce the complexity of input data to a computationally feasible range, alternative stable

states were explored based on family-level community compositions (threshold for

binarization = 10 %). The 10 alternative stable states representing ≥ 10 samples were

compared in terms of their associations with crop disease conditions.

Fig. 6 | Network modules and hubs within the microbiomes. **a**, Co-occurrence networks of archaea, bacteria, and fungi. Specificity of occurrences to disease-level-1 (the lowest disease level) samples is shown for each ASV within the network. The specificity is shown as node size separately for negative (left) and positive (right) associations with least-diseased states of crop plants. Colors indicate network modules, in which microbial ASVs in commensalistic/mutualistic interactions and/or those sharing environmental preferences are densely linked with each other. See Extended Data Figure 7 for taxonomy (archaea, bacteria, or fungi) of respective nodes. **b**, Characteristics of network modules. Mean specificity to disease-level-1 samples is shown for each network module. **c**, Betweenness centrality of ASVs. For each ASV, betweenness centrality within the co-occurrence network is shown along the horizontal axis. Specificity to disease-level-1 samples is shown along the vertical axis. **d**, Topological roles of ASVs within and across network modules. For each ASV, within-module degree represents the number of the ASVs linked with the target ASV within a module (z-standardized). Among-module connectivity represents the extent to which an ASV interlinks ASVs belonging to different network modules.

829 **Extended Data Figures**

832   **Extended Data Fig. 1 | Microbial taxonomic compositions at the genus, order, and class**

833   **levels**. **a**, Genus-level taxonomic compositions of prokaryotes and fungi. See Figure 1a for the

834   map of the research sites. **b**, Order-level taxonomic compositions. **c**, Class-level taxonomic

835   compositions.

Extended Data Fig. 2 | Community structure and metadata properties of the samples. **a**, Prokaryote/fungal community structure and research-site profiles. Research sites are indicated by colors on the PCoA surface of prokaryote/fungal community structure. **b**, Soil taxonomy profile. **c**, Former crop plant.

Extended Data Fig. 3 | Community structure and soil chemical properties. **a**,

Prokaryote/fungal community structure and soil pH of the samples. **b**, Electrical conductivity.

c, Available phosphorous concentration. **d**, Carbon to nitrogen ratio.

Extended Data Fig. 4 | Number of the stable states of community structure (threshold

for binarization = 5 %). The parameters of the extended pairwise maximum entropy model

were adjusted to fit the model to the empirical data (see Methods). The number of stable

states in the energy landscape was then estimated by using the gradient descent algorithm for

20,000 randomly generated community states within the 2^S possible community states, where

S denotes the number of families or orders. In the heuristic search, sets of values of the

environmental variables (soil pH, electrical conductivity, C/N ratio, and available

phosphorous concentration) were drawn from those of sampling points. The boxplots show

variation in the number of stable states across different sets of environmental variables. A box

represents the first quartile, median and the third quartile, while bars and circles indicate 1.5 ×

inter-quartile range and outliers, respectively. The analysis was performed, respectively, for

the dataset representing both prokaryotes and fungi, the prokaryote dataset, and the fungus

dataset. The threshold value in the binarization of the community data was set at 10 %: see

Extended Data Figure 4 for results based on another threshold value (5 %).

Extended Data Fig. 5 | Alternative stable states defined for order-level fungal data
(threshold for binarization = 5 %). a, Energy landscape of fungal community structure. The
surface of energy levels was reconstructed on the PCoA dimensions (Fig. 2) of fungal
community structure based on spline smoothing. Community states with lower energy are
inferred to be more stable. Data points (samples) indicated by the same color belong to the
basins of the same alternative stable states, whose energy is lower than any adjacent
community structure. See panel c for details about the alternative stable states. b, Definitions
of the stable states. To reduce the complexity of input data to a computationally feasible

873   range, alternative stable states were explored based on order-level community compositions

874   (threshold for binarization = 5 %). The 7 alternative stable states representing $\geq$ 10 samples

875   were compared in terms of their associations with crop disease conditions.

a

b

Extended Data Fig. 6 | Each ASV's specificity to disease levels. **a**, Standardized specificity
to disease level (s_{ij}). Based on a randomization approach, each ASV's specificity to samples
differing in crop disease levels was evaluated. A higher value of the specificity index
indicates that a microbial ASV displayed higher abundance in samples of the target disease
level than that expected by chance. **b**, Specificity scores and statistical significance. An s_{ij}
value higher than 4 or that lower than -4 indicates statistically strong specificity of an ASV
to a target disease level.
885

887   **Extended Data Fig. 7 | Network centrality scores**. **a**, Distribution of archaea, bacteria, and

888   fungi within the co-occurrence network. **b**, Each ASV's betweenness centrality score within

889   the network. The node size represents relative betweenness centrality scores. **c**, Each ASV's

890   eigenvector centrality score within the network. The node size represents relative eigenvector

891   centrality scores.