Alternative stable states in soil microbiomes of agroecosystems

Hiroaki Fujita[a], Shigenobu Yoshida[b], Kenta Suzuki[c] and Hirokazu Toju[a*]

[a]Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2133, Japan

[b]Institute for Plant Protection, National Agriculture and Food Research Organization, Tsukuba, Ibaraki 305-8666, Japan

[c]Integrated Bioresource Information Division, BioResource Research Center, RIKEN, Tsukuba, Ibaraki 305-0074, Japan

*Hirokazu Toju

Email: toju.hirokazu.4c@kyoto-u.ac.jp

Author Contributions: H.T. conceived the project; H.F., S.Y. and H.T. designed the work; H.F. performed molecular experiments; H.F. and H.T. analyzed the data; H.T. wrote the paper with H.F., S.Y., and K.S..

Competing Interest Statement: The authors declare that they have no competing interests.

Classification: Biological Sciences: Environmental Sciences and Ecology

Keywords: biodiversity; community stability; ecosystem functions; multistability; species coexistence

This PDF file includes:

      Main Text
      Figures 1 to 5

Abstract

Archaea, bacteria, and fungi in the soil are increasingly recognized as major determinants of agricultural
productivity and sustainability. Although our knowledge of soil biodiversity has been greatly enhanced for
the last decade, defining key characteristics of stable and highly functional microbiomes remains a major
challenge. We here show that "stable states" of soil microbiomes can be defined with their disease-
suppressive effects on crop plants. We integrated agricultural field metadata with microbial community
analyses by targeting > 2,000 soil samples collected across a latitudinal gradient from cool-temperate to
subtropical regions, profiling microbiomes constituted, in total, by 11 archaeal, 332 bacterial, and 240 fungal
families. A DNA-metabarcoding-based evaluation of balance between prokaryote and fungal abundance
indicated that fungus-dominated soil is more resistant to crop plant disease. We then examined whether
disease-suppressive soil can be defined based on prokaryote and/or fungal taxonomic compositions. A
statistical-physics-based approach for reconstructing "energy landscapes" of community structure indicated
that disease-suppressive and disease-susceptible soils could be distinguished as alternative stable states of
fungal communities. On the other hand, bacterial community structure could not be classified into a small
number of alternative stable states, making it difficult to build simplified criteria for diagnosing disease-
suppressive functions of agroecosystems with prokaryote taxonomic compositions alone. As the 22 fungal
families keys to define the alternative stable states have cosmopolitan distributions, their roles and
compositional balance are potentially keys to maintain agroecosystem functions and stability in diverse
types of cropland soils.

Significance Statement

49   Managing "healthy" soil microbial communities is a prerequisite of sustainable crop production on the Earth.
50   Based on intensive profiling of soil archaea, bacteria, and fungi in cool-temperate to subtropical regions, we
51   found general community-level characteristics common to disease-resistant agroecosystems. Coexistence of
52   multiple fungal lineages that can suppress devastating pests and pathogens is the key to stabilize and
53   maximize ecosystem functions in agriculture.

Main Text

Introduction

The ongoing global-scale degradation of agroecosystems is threatening food production (1, 2). Maximizing
the functions of microbial communities is a prerequisite for building bases of sustainable agriculture (3–6).
Archaea, bacteria, and fungi not only promote soil nutrient cycles but also protect crop plants from
pests/pathogens and abiotic stresses (7–11). Therefore, diagnosing soil microbiome states is of central
interest when we try to design and manage resource-efficient and disease-tolerant agroecosystems.

A number of studies have explored specific microbes and specific combinations of microbes promoting
crop plants' growth (3, 12–14). However, such microbial functions can be lost due to unexpected changes in
community structure in soil microbiomes (15). In general, microbiome compositions in the wild are
consequences of multiple ecological processes (13, 16–19). Selection by environmental conditions, for
example, is a primary mechanism determining establishment of microbial populations (18, 20, 21).

Furthermore, competitive and facilitative interactions between microbial species are essential factors
governing local community structure (22–24). Continually influenced by these mechanisms and factors, local
microbial communities differ in their functions and stability (14, 25–27). Thus, understanding of not only
functions but also community-level stability is essential for maximizing the benefits provided by
microbiomes (12, 22, 28). Stability in soil or plant-associated microbiomes have started to be examined in
studies based on laboratory experiments (29, 30). Nonetheless, we still have limited knowledge of soil
microbiome stability in wild ecosystems.

In theoretical ecology, stability of community states (taxonomic or species compositions) is often
discussed in the framework of stability landscapes (31–33). On the landscape representing
stability/instability of community structure (*SI Appendix*; Fig. S1), stable community states are split by
unstable community states (31–34). If those "alternative stable states" (31, 34, 35) [see also (36, 37)] differ
in their community-level biological functions, stable and highly functional community states can be explored
within the stability landscapes. Based on a mathematical approach developed in statistical physics (38, 39), it
is now possible to reconstruct "energy landscapes", which represent structure of stability landscapes, from
empirical datasets of ecological communities (32). Although hundreds or thousands of community
compositional data points are required to apply the statistical approach (32), such energy landscape analyses
will allow us to define key features of stable and highly functional microbiome states out of numerous
possible combinations of microbes.

We here perform systematic profiling of cropland microbiome structure across the Japanese
Archipelago in order to define general characteristics of disease-suppressive and disease-susceptible soils
(40–43) based on theoretical ecology. We integrate DNA metabarcoding results of prokaryote and fungal
communities with environmental meta-datasets including > 2,000 soil samples collected from the croplands
of 19 plant species. With the bird's-eye view of possible microbiome structure, we map alternative stable
states within the reconstructed energy landscape of microbiome structure. We then test the hypothesis that
disease-suppressive and disease-susceptible states of soil microbiomes are defined with compositions of
cosmopolitan microbial taxa. Overall, this study shows how we can reduce the complexity of microbiome
data to build simplified diagnostic criteria for exploring optimal soil microbiomes with high controllability
and agricultural functionality.

97   Results

98   Diversity of microbiome structure

We compiled the field metadata of 2,903 soil samples collected in the research projects of National
Agricultural and Food Research Organization (NARO). The bulk soil of farmlands was sampled from
subtropical to cool-temperate regions (26.1–42.8 °N) across the Japanese Archipelago, targeting 19 crop
plant species (Fig. 1a) from 2006 to 2014. Most of the croplands were managed with conventional
agricultural practices (characterized by intensive tillage and chemical fertilizer/pesticide application), while
some were experimentally controlled as organic agricultural fields. The metadata included the information
of chemical [e.g., pH, electrical conductivity, carbon/nitrogen (C/N) ratio, and available phosphorous
concentration], physical (e.g., soil taxonomy), and biological (e.g., crop disease level) properties, providing an
invaluable platform for profiling and categorizing ecosystem states of cropland soil.

To integrate the metadata with the information of microbial community structure, we performed DNA
metabarcoding analyses of both prokaryotes (archaea and bacteria) and fungi. After a series of quality
filtering, prokaryote and fungal community data were obtained from 2,534 and 2,384 samples, respectively.
In total, 579 archaeal amplicon sequence variants (ASVs) representing 14 genera (11 families), 26,640
bacterial ASVs representing 875 genera (332 families), and 6,306 fungal ASVs representing 573 genera (240
families) were detected (Fig. 1B; *SI Appendix*: Fig. S2).

The prokaryote communities did not have dominant taxa at the genus and family levels, although the
archaeal family Nitrososphaeraceae had the highest abundance at some research sites (Fig. 1B). In contrast,
the fungal communities were dominated by fungi in the families Mortierellaceae, Chetomiaceae, and
Nectriaceae (Fig. 1B). A reference database profiling of fungal functional groups suggested that the fungal
communities were dominated by plant pathogenic and soil saprotrophic fungi (Fig. 1C) as characterized by
the dominance of *Fusarium* and *Mortierella* at the genus level (*SI Appendix*: Fig. S2). Meanwhile,
mycoparasitic fungi had exceptionally high proportions at some research sites, as represented by the
dominance of *Trichoderma* (Hypocreaceae) at those sites (Fig. 1B; *SI Appendix*: Fig. S2).

The prokaryote and fungal communities differed in basic biodiversity patterns. In terms of α-diversity
at the family level, prokaryote communities showed higher Shannon's entropy than fungal communities in
all but one of the research sites (Fig. 1D). In contrast, variability of community structure among samples, i.e.,
Bray-Curtis β-diversity, was higher in fungi than in prokaryotes in most study sites (Fig. 1E). Likewise, fungi
showed higher levels of community structural difference between research sites than prokaryotes (Welch's
test; t = -6.6, df = 168.9, P < 0.0001; Fig. 1F). These results indicate that diverse taxonomic groups of archaea
and bacteria are present without dominance by particular taxa, while fungal community structure is spatially
dynamic both within and across agroecosystems.

Compiling the metadata of edaphic factors, we found that variation in the community structure of
prokaryotes and fungi was significantly explained by crop plant identity and soil taxonomy as well as by soil
chemical properties such as pH, electrical conductivity, C/N ratio, and available phosphorous concentration
(Fig. 2A-B; *SI Appendix*: Figs. S3-4; Table S1). Meanwhile, both prokaryote and fungal community structure
was significantly associated with the severity of crop disease (Fig. 2C-D). Thus, the dataset as a whole offers
an opportunity for understanding mechanisms spanning physical, chemical, and biological processes in
agroecosystems.

138   Crop disease and microbiome structure

We then focused on factors determining disease-suppressive and disease-susceptible properties of
agroecosystem soil. In addition to chemical environmental factors (e.g., soil pH and electrical conductivity),
whose associations with agricultural productivity have been routinely examined (44, 45), we explored
biological factors critical to crop-plant disease level. The α-diversity of prokaryotes and/or fungi, for
example, is an index that can be estimated with DNA-based profiling of soil biological communities. In
further reconstructing biological factors in a high-throughput manner, we focused on the ratio of prokaryote
abundance to fungal abundance. As the classic chromatography-based method for quantifying phospholipid
fatty acids specific to bacteria or fungi (46) could require impractically long time for processing > 2,000
samples, we developed an alternative approach for estimating the ratio of prokaryote abundance to fungal
abundance (hereafter, prokaryote/fungus ratio) based on DNA metabarcoding. Specifically, the relative
concentrations of prokaryote 16S rRNA gene sequences to fungal ITS gene sequences were calculated based

on a quantitative amplicon sequencing analysis (Fig. 3A; see Materials and Methods for details). We then
constructed statistical models comparing effects on crop-plant disease level among various types of
chemical/biological factors.

A series of analysis indicated prokaryote/fungus ratio had strong impacts on crop-plant disease level.
While soil chemistry (soil pH) had marginally significant effects on the proportion of diseased crop plants, the
prokaryote/fungus ratio showed much stronger associations with crop plant disease level (Model 1 in Fig.
3B). Specifically, crop plant disease was more likely to occur when prokaryotes were abundant relative to
fungi (Fig. 3C). The effects of prokaryote/fungus ratio on plant disease level were conspicuous even when
they were compared with effects of prokaryote and fungal α-diversity (Model 2 in Fig. 3B). The threshold
values over which more than 50 % of plants showed disease levels 2–5 were estimated to be around 0.7 in a
log-ratio unit (Fig. 3C), although the values potentially vary depending on various factors such as crop plants,
geography, and climatic conditions.

Alternative stables states

We next reconstructed alternative stable states (31, 34, 37) of microbiomes in order to define community
structure of disease-suppressive and disease-susceptible soils. In an energy landscape analysis of statistical
physics (32), we modeled how microbial community compositions (community states) depend on soil
chemical factors and interactions between microbes. The modeling approach allowed us to calculate the
"energy" of each community state. We then estimated the shape of an energy landscape, on which lower
energy represents more stable community states (*SI Appendix*; Fig. S1). Stable states were then defined as
the community compositions whose energy is lower than all possible adjacent community compositions (32).

The results of energy landscape reconstruction fundamentally differed between prokaryotes and
fungi. Even if community states were defined in a binary format at the family or order level to reduce the
complexity of the input data (i.e., 2^S possible community states, where S denotes the number of families; see
Materials and Methods), more than 10^3 stable states were inferred to exist for the prokaryote dataset (Fig.
4). This result suggests that ecological processes of soil prokaryote communities are described as shifts
between many equally possible compositions and that categorization of ecosystem functions based on
prokaryote community structure is challenging. In contrast, the energy landscape of fungal community
structure was represented by much smaller number of stable states (Fig. 4). While the estimated number of
stable states varied depending on threshold values in the binarization of community structural data, fungal
communities consistently showed lower number of stable states than prokaryote communities (*SI Appendix*:
Fig. S5).

We then filtered the alternative stable states of fungal communities based on the "depth of basins" (*SI*
Appendix: Fig. S1) and the number of samples belonging to the basins of the stable states (32), obtaining 10
major stable states representing the energy landscape (Fig. 5A; see also *SI Appendix*: Fig. S6). By integrating
the information of the energy landscape (Fig. 5A) with that of crop-plant disease level (Fig. 5B), we ranked
the stable states in terms of possible disease-suppressive functions (Fig. 5C). We then found that all soil
samples belonging to the top-three stable states had associations with the lowest crop-disease level, while
92 % of samples belonging to the lowest-rank stable state showed higher disease levels (Fig. 5C).

The 10 stable states were distinguished in light of balance among 22 fungal families with
cosmopolitan distributions (Fig. 5C). Hypocreaceae, which embraced well-known mycoparasitic taxa [e.g.,

Trichoderma (47)], appeared at high proportions in stable states with potential disease-suppressive
functions. In contrast, Aspergillaceae, which included both plant pathogens and saprotrophs (48), were
relatively abundant in stable states with high disease potential. In addition to these basic patterns, several
families were keys to define disease-suppressive or disease-susceptible soils. For example, high abundance
of Didymellaceae, Lasiosphaeriaceae, Mrakiaceae, Orbilliaceae, Piptocephalidaceae, and
Plectosphaerellaceae characterized the two stable states with the highest performance (Fig. 3D). Among the
families, Orbiliaceae, Piptocephalidaceae, and Plectosphaerellaceae have been known to include
nematophagous and mycoparasitic taxa (49–51), presumably working as suppressors of plant-parasitic
nematodes and fungi in the soil.

200
Discussion

By integrating large datasets of agricultural field properties and DNA metabarcoding, we found that
seemingly diverse microbiome structure could be categorized into several stable states varying in
ecosystem-level properties. The analysis of energy landscapes suggested that prokaryote communities had
numerous local compositional equilibria or quasi-equilibria (Fig. 4), presumably due to their rapid and
continuous responses to changes in environmental conditions. In contrast, ecological processes of fungal
communities could be interpreted as shifts between a small number of alternative stable states (Fig. 5).
Basically, fungi can form complex hyphal structures and their individual lifespan is much longer than most
archaea and bacteria. Because of these biological properties, structure and functions of fungal communities
are more likely to be influenced by legacy effects of past ecological processes (7, 37, 52, 53) than those of
prokaryotes. Consequently, fungal taxonomic compositions can give essential insights into current status of
agroecosystem functions, while prokaryote community structure is too flexible to be used for building
simplified diagnostic criteria for soil microbiome functions.

These insights into alternative stable states add an important dimension of discussion on the use of
microbiomes in agriculture. Beyond investigations on single species/strains of microbes, microbiome studies
have explored sets of microbes that collectively maximize biological functions (3, 12–14). In particular,
experimental studies on "synthetic" communities have reorganized our understanding of microbiome
functions (12–14). Nonetheless, such microbial functions cannot be realized in real agroecosystems if the
synthesized or designed microbiome compositions are vulnerable to biotic and abiotic environmental
changes in the wild (15). Thus, in addition to functional properties, compositional stability is the key to
manage microbiomes in agroecosystems (12, 22, 28). The finding that the number of alternative stable
states is much fewer in fungi than in prokaryotes (Fig. 4) highlights the possibility that agroecosystem
functions are managed by keeping fungal community structure at targeted stable states with positive
impacts on crop production. The fact that prokaryote communities could not be classified into a small
number of alternative stable states does not indicate lack of their roles in crop production. Bacteria and
archaea play fundamental roles in carbon- and nitrogen-cycles of soil ecosystems (54, 55). The point is that
information of fungal taxonomic compositions is indispensable for adaptively managing microbiome
structure in agricultural practices.

In our analysis across the Japanese Archipelago, 22 out of the 240 fungal families detected were keys
to distinguish disease-suppressive and disease-susceptible soil microbiomes (Fig. 5C). Among them,
Hypocreaceae, Orbiliaceae, Piptocephalidaceae, and Plectosphaerellaceae are of particular interest because
they include many species utilized as biological control agents for suppressing plant pathogenic fungi and/or
plant parasitic nematodes (47, 49–51). Fungal pathogens and cyst/root-knot nematodes have continued to
have devastating impacts on global crop production (43, 56, 57). Because those fungi and nematodes cannot

be eradicated with chemical fungicide/pesticides, understanding of the biological processes in which their
outbreaks are prevented in nature is an essential step for implementing sustainable agriculture. Our results
showed that plant disease could be suppressed under coexistence of multiple nematophagous and
mycoparasitic fungal taxa (Fig. 5). Therefore, shifting fungal community structure towards the stable states
represented by the nematophagous/mycoparasitic families will provide more comprehensive solutions than
choosing single biological control agents with best disease-suppressive functions.

Although we focused on macroscopic structure of microbial communities by performing energy
landscape analysis at the family and order levels, analyses at lower taxonomic levels will enhance our
understanding of microbiome stability. Positive or adverse impacts on plants often differ substantially
among microbial species or strains (58). Thus, analyses at the level of ASVs or operational taxonomic units
(OTUs) are necessary if we try to find "keystone" microbial species (59, 60) at each local ecosystem.
Meanwhile, such knowledge of specific species or strains are not utilized in other climatic or geographic
regions in which the focal species/strains are absent (19). Thus, for the purpose of building broadly
applicable criteria of stable and functional microbiomes, energy landscape analyses of taxonomic
compositions, rather than those of species- or strain-level compositions, are advocated. Given that all the 22
fungal families highlighted in our analysis have cosmopolitan distributions, a next crucial step is to test
whether the alternative stable states defined across the Japanese Archipelago can be used to categorize
disease-suppressive and disease-susceptible microbiomes in other regions on the globe.

Although we herein profiled alternative stable states of microbiomes by focusing mainly on
conventional agricultural fields, soil microbiomes with higher functions and stability are potentially explored
in the framework of conservation agriculture (61) or regenerative agriculture (62). Minimum tillage (63), for
example, can decrease physical damage to fungal hyphae (64), thereby increasing plants' nutrient uptake
through mycorrhizal hyphal networks and suppressing pests/pathogens by the functions of entomophagous,
nematophagous, and mycoparasitic fungi. In this respect, the fact that crop disease frequency decreased
towards the range of dominance by fungi over prokaryotes (Fig. 3) is of particular interest. We also found
that the prokaryote/fungus ratio varied considerably among soil samples with similar prokaryote/fungal
community compositions (Fig. 3A). Thus, analyses of balance between prokaryote and fungal abundance, in
line with energy landscape analyses of community states, will help us monitor and control fungal biomes for
further exploration of sustainable crop production.

We herein explored stable microbial community structure of disease-suppressive soil based on an
intensive profiling of farmland soil across the Japanese Archipelago. Although fungi are often dismissed in
studies on soil microbiomes, analyses of fungal biomes give essential insights into agroecosystem states (7,
43). For further understanding relationship among structure, stability, and functions of soil microbiomes, it is
necessary to incorporate additional measures of microbiome performance (e.g., crop yield and nitrogen
utilization efficiency) into community ecological analyses. Moreover, developing ways to shift microbiome
structure from unfavorable to favorable stable states is a crucial challenge for controlling agroecosystem
functions with minimum anthropogenic intervention. Bird's-eye views of stable microbiome compositions
will provide platforms for integrating ecology, genomics, and microbiology towards the realization of
"nature-positive" world (65).

274
275
276 **Materials and Methods**
277

Soil samples and metadata

Over research projects of National Agricultural and Food Research Organization (NARO), which were carried
out through five national research programs funded by Ministry of Agriculture, Forestry and Fisheries, 2,903
rhizosphere/bulk soil samples were collected from conventional agricultural fields across the Japanese
Archipelago from January 23, 2006, to July 28, 2014 (*SI Appendix*: Dataset S1). When the latitude and
longitude of the sampling positions were round to one decimal place, 42 research sites were distinguished.
Across the metadata of the 2,903 samples, the information of 19 crop plants, 34 former crop plants
(including "no crop"), 13 soil taxonomic groups (e.g., "Andosol"), 60 experimental/research purposes (e.g.,
"soil comparison between organic and conventional management") was described. Likewise, the metadata
included the information of dry soil pH, electrical conductivity, carbon/nitrogen (C/N) ratio, and available
phosphorous concentration from, 2,830, 2,610, 2,346, and 2,249 samples, respectively. In addition, the
information of the severity of crop plant disease was available for 1,472 samples (tomato, 637 samples;
Chinese cabbage, 336 samples; eggplant, 202 samples; celery, 97 samples; Broccoli, 96 samples, etc.). The
values of the proportion of diseased plants or disease severity index was normalized within the ranges from
0 to 100, and they were then categorized into five levels (level 1, 0-20; level 2, 20-40; level3; 40-60; level 4,
60-80; level 5, 80-100). The plant pathogens examined in the disease-level evaluation were *Colletotrichum*
gloeosporioides on the strawberry, *Fusarium oxysporum* on the celery, the lettuce, the strawberry, and the
tomato, *Phytophthora sojae* on the soybean, *Plasmodiophora brassicae* on Cruciferae plants, *Pyrenochaeta*
lycopersici on the tomato, *Pythium myriotylum* on the ginger, *Ralstonia solanacearum* on the eggplant and
the tomato, and *Verticillium* spp. on Chinese cabbage. For continuous variables within the metadata,
emergent outliers (mean + 5 SD) were converted into "NA" in the data matrix used in the following statistical
analyses as potential measurement/recording errors. Unrealistic electrical conductivity records (> 20) were
converted into "NA" as well.

At each sampling position, five soil sub-samples collected from the upper layer (0-10 cm in depth) at
five points (ca. 100 g each) were mixed. The mixed soil sample (ca. 500 g) was then sieved with 2-mm mesh
in the field. The samples were stored at -20 °C until DNA extraction. In laboratory conditions, 0.4 g of soil
(fresh weight) was subjected to DNA extraction with FastDNA SPIN Kit for Soil (Q-BioGene).

DNA amplification and sequencing

Profiling of soil microbial biodiversity was performed by targeting archaea, bacteria, and fungi. For the
amplification of the 16S rRNA V4 region of archaea and bacteria (prokaryotes), the set of the forward primer
515f (66) and the reverse primer 806rB (67) were used. The primers were fused with 3–6-mer Ns for
improved Illumina sequencing quality (68) and Illumina sequencing primers as detailed in a previous study
(69). PCR was performed using KOD ONE PCR Master Mix (TOYOBO, Osaka) with the temperature profile of
35 cycles at 98 °C for 10 seconds (denaturation), 55 °C for 5 seconds (annealing of primers), and 68 °C for 30
seconds (extension), and a final extension at 68 °C for 2 minutes. The ramp rate through the thermal cycles
was set to 1 °C/sec to prevent the generation of chimeric sequences (70). In the PCR, we added five artificial
DNA sequence variants with different concentrations (i.e., standard DNA gradients; 1.0×10^{-4}, 5.0×10^{-5}, 2.0
$\times 10^{-5}$, 1.0×10^{-5}, and 5.0×10^{-6} nM; *SI Appendix*: Table S2) to the PCR master mix solution in order to calibrate
16S rRNA gene copy numbers in the template DNA samples as detailed elsewhere (30, 71, 72). Likewise, the
internal transcribed spacer 1 (ITS1) region of fungi was amplified using the set of the forward primer
ITS1F_KYO1 and the reverse primer ITS2_KYO2 (73). PCR was performed using the Illumina-sequencing

fusion primer design mentioned above with the temperature profile of 35 cycles at 98 °C for 10 seconds, 53
°C for 5 s seconds, and 68 °C for 5 seconds, and a final extension at 68 °C for 2 minutes (ramp rate = 1
°C/sec). Newly designed artificial sequence variants (1.0×10^{-5}, 7.0×10^{-6}, 5.0×10^{-6}, 2.0×10^{-6}, and 1.0×10^{-6}
nM; *SI Appendix*: Table S2) were added to the PCR master mix as standard DNA gradients for the calibration
of the ITS sequence concentrations in the template DNA samples.

The PCR products of the prokaryote 16S rRNA and fungal ITS1 regions were respectively subjected to
the additional PCR step for linking Illumina sequencing adaptors and 8-mer sample identifier indexes (74)
with the amplicons as detailed elsewhere (69). The temperature profile in the PCR was 8 cycles at 98 °C for
10 seconds, 55 °C for 5 seconds, and 68 °C for 5 seconds, and a final extension at 68 °C for 2 minutes. The
PCR products were then pooled for each of the 16S rRNA and fungal ITS1 regions after a
purification/equalization process with the AMPureXP Kit (Beckman Coulter, Inc., Brea). Primer dimers, which
were shorter than 200 bp, were removed from the pooled library by supplemental purification with
AMPureXP: the ratio of AMPureXP reagent to the pooled library was set to 0.8 (v/v) in this process. The
sequencing libraries of the two regions were processed in an Illumina MiSeq sequencer (10% PhiX spike-in).
Because the quality of forward sequences is generally higher than that of reverse sequences in Illumina
sequencing, we optimized the MiSeq run setting in order to use only forward sequences. Specifically, the run
length was set 271 forward (R1) and 31 reverse (R4) cycles to enhance forward sequencing data: the reverse
sequences were used only for discriminating between prokaryote 16S and fungal ITS1 sequences in the
following bioinformatic pipeline.

Bioinformatics

In total, 23,573,405 sequencing reads were obtained in the Illumina sequencing (16S rRNA, 11,647,166
sequencing reads; ITS, 11,926,239 sequencing reads). The raw sequencing data were converted into FASTQ
files using the program bcl2fastq 1.8.4 distributed by Illumina. For each of the 16S rRNA and fungal ITS1
regions, the output FASTQ files were demultiplexed using Claident v0.9.2022.01.26 (75). The sequencing
data were deposited to DNA Data Bank of Japan (DDBJ) (Bioproject accession no.: PSUB018361). The
removal of low-quality sequences and ASV inferences were done using DADA2 (76) v1.17.5 of R v3.6.3 (77).
The mean number of filtered sequencing reads obtained per sample was 3,949 and 4,075 for the prokaryote
and fungal datasets, respectively. Taxonomic annotation of prokaryotes was conducted based on the SILVA
138 SSU (Quast et al., 2013) the assignTaxonomy function of DADA2. For the taxonomic annotation of fungi,
the QCauto method (75) was applied to the NCBI nucleotide sequence database bundled with Claident
v0.9.2022.01.26. ASVs that were not assigned to the domain Archaea/Bacteria and the kingdom Fungi were
removed from the 16S rRNA and ITS1 datasets, respectively. For each target organismal group (prokaryotes
and fungi), we then obtained a sample × ASV matrix, in which a cell entry depicted the number of
sequencing reads of an ASV in a sample. The samples with less than 1,000 reads were discarded from the
matrices. The matrix was rarefied to 1,000 reads per sample for the downstream statistical pipeline. The
number of reads was insufficient for comprehensively profiling rare microbial species, which are often
targets of soil microbiome studies. However, because data matrices including numerous rare ASVs or OTUs
could not be subjected to the computationally intensive ecological analyses detailed below even if we used
supercomputers, we focused on major components of soil prokaryote and fungal biomes. In other words,
our purpose here was to extract major components of agroecosystem soil microbiomes across the Japanese
Archipelago, thereby developing simple diagnostic criteria for distinguishing disease-suppressive and
disease-susceptible soils.

In total, prokaryote and fungal community data were obtained for 2,534 and 2,384 samples,
respectively. The number of ASVs included in the rarefied matrices were 27,219 and 6,306 for prokaryotes
and fungi, respectively. For fungal ASVs, putative functional groups (e.g., "plant pathogen") were inferred
using the program FungalTraits (78). The estimation of DNA concentrations of the prokaryote 16S rRNA and
fungal ITS regions was performed, respectively, based on the calibration with the standard DNA gradients
(artificial DNA variants introduced to the PCR master mix solutions) as detailed elsewhere (30).

Diversity of microbiome structure

For each sample, Shannon's α-diversity estimates of prokaryotes and fungi were respectively calculated.
Because biological entities of ASVs or operational taxonomic units (OTUs) differ considerably between
prokaryotes and fungi, we compared α-diversity between the two organismal groups at the family level.
Likewise, mean Bray-Curtis dissimilarity (β-diversity) in family-level compositions within each of the 42
research sites was calculated as a measure of within-site variability of prokaryote/fungal community
structure. In addition, the degree of among-site community differentiation was evaluated for prokaryotes
and fungi. Specifically, after calculating mean family-level compositions at each research site, Bray-Curtis β-
diversity was calculated in all pairs of research sites. The α-diversity, within-site β-diversity, and among-site
β-diversity estimates were respectively compared between prokaryotes and fungi.

For each of the prokaryote and fungal datasets, permutational analyses of variance (PERMANOVA)
were performed to examine associations between family-level community compositions and variables in the
metadata. Two types of PERMANOVA models were constructed based on Bray-Curtis β-diversity (10,000
iterations). Specifically, one is constituted by categorical explanatory variables (crop plant, former crop plant,
soil taxonomy, research site, and sampling month), while the other included continuous explanatory
variables (soil pH, electrical conductivity, C/N ratio, and available phosphorous concentration, latitude, and
longitude).

To reduce the dimensions of the community composition data, a principal coordinate analysis (PCoA)
was performed based on the Bray-Curtis β-diversity calculated with the rarefied sample × ASV matrix of
prokaryotes or fungi. For each PCoA axis (axes 1 to 5) in each of the prokaryote and fungal analyses,
Pearson's correlation with each chemical environmental factor (soil pH, electrical conductivity, C/N ratio,
and available phosphorous concentration) was calculated.

Crop disease and microbiome structure

We then evaluated how community structure of prokaryotes and fungi were associated with crop disease.
For each of the prokaryote and fungal datasets, a generalized linear mixed model (GLMM) of crop-disease
level was constructed by including the PCoA axes 1 to 5 as fixed effects. Sampling month and
experimental/research purposes in the metadata were set as random effects. A logit-link function and
binomial error was assumed in the GLMM after converting the response variable into a binary format
[disease level 1 (= 1) vs. disease levels 2–5 (= 0)].

We also examined how balance between prokaryote and fungal abundance could be associated with
crop disease levels. Based on the estimated concentrations of prokaryote 16S rRNA and fungal ITS sequences

in template DNA solutions, we calculated the ratio of prokaryote DNA concentrations to fungal DNA
concentrations in respective samples (prokaryote/fungus ratio) as follows:

404
$$\log\left(\frac{\text{prokaryote 16S rRNA gene concentration(DNA copies/}\mu\text{L)}}{\text{fungal ITS gene concentration (DNA copies/}\mu\text{L)}}\right).$$

Although potential variation in DNA extraction skills of researchers might affect absolute DNA
concentrations in the template DNA solutions, balance between prokaryote and fungal DNA in each
template DNA sample could be used as a reliable measure. The DNA-metabarcoding-based approach of
estimating prokaryote/fungus ratio has advantage over quantitative-PCR-based approaches. Specifically, the
former approach allows us to eliminate effects of nonspecific PCR amplification based on DNA sequencing
data, while the latter is affected by "contamination" of nontarget amplicons (e.g., plastid DNA in 16S rRNA
sequencing and plant DNA in ITS sequencing).

Potential effects of the prokaryote/fungus ratio on crop-disease level were examined in a GLMM with
a logit-link function and binomial errors. In the GLMM, soil pH and electrical conductivity were included as
fixed effects in addition to the prokaryote/fungus ratio. In the GLMM, C/N ratio and available phosphorous
concentration were not included due to the small number of samples with both crop-disease information
and the chemical variables. Sampling month and experimental/research purpose information in the
metadata were included as random effects in the GLMM. To examine the effects of other aspects of
biological factors, we constructed another GLMM including the ASV-level Shannon diversity of prokaryotes,
that of fungi, and the prokaryote/fungus ratio. All the explanatory variables were *z*-standardized (zero-mean
and unit-variance) before the GLMM analysis. To visualize how the proportion of least-diseased (disease-
level 1) samples varied along the axis of the prokaryote/fungus ratio, we performed a supplementary
analysis with generalized linear models (GLMs). Specifically, for the five experimental/research purpose
categories from which 10 or more disease-level 1 (least-diseased) samples and 10 or more disease-level 2–5
(severely diseased) samples were obtained, a GLM of disease level was constructed by including the
prokaryote/fungus ratio as an explanatory variable.

Alternative stables states

428 We estimated the alternative stable states of microbial community structure based on the framework of an
429 energy landscape analysis(32, 38) (tutorials of energy landscape analyses are available at
430 https://community.wolfram.com/groups/-/m/t/2358581). Probabilities of community states $\mathrm{p}\left(\vec{\sigma}^{(k)}\right)$ are
431 given by

432
$$P\left(\vec{\sigma}^{(k)}\middle|\pmb{\varepsilon}\right) = e^{-H\left(\vec{\sigma}^{(k)},\varepsilon\right)}/Z$$

433
$$Z = \sum_{i=1}^{2^S} e^{-H\left(\vec{\sigma}^{(i)},\varepsilon\right)},$$

where $\vec{\pmb{\sigma}}^{(k)} = \left(\sigma_1^{(k)}, \sigma_2^{(k)}, \dots, \sigma_S^{(k)}\right)$ is a community state vector of *k*-th community state and *S* is the total
number of taxa (e.g., ASVs, species, genera, or families) examined. $\pmb{\varepsilon} = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_M)$ is an array of
continuous values representing environmental factors (e.g., soil pH and electrical conductivity) and *M* is the
total number of environmental parameters. $\sigma_i^{(k)}$ is a binary variable that represents presence (1) or absence
(0) of taxon *i*: i.e., there are a total of 2^S community states. As the exploration of the 2^S community states

were computationally intensive, we coded community states at the family-level taxonomic compositions.
Specifically, families that represented 10 % or more of the total sequencing reads of a target sample were
coded as 1, while those representing less than 10 % of total reads were coded as 0. Then, the energy of the
community state $\vec{\sigma}^{(k)}$ is given by the extended pairwise maximum entropy model:

$$H(\vec{\sigma}^{(k)}, \varepsilon) = -\sum_{i=1}^{S} h_i \vec{\sigma}_i^{(k)} - \sum_{j=1}^{S} \sum_{i=1}^{M} g_{ij} \varepsilon_i^{(k)} \sigma_j^{(k)} - \sum_{i=1}^{S} \sum_{j=1}^{S} J_{ij} \vec{\sigma}_i^{(k)} \vec{\sigma}_j^{(k)} / 2,$$

where h_i represents the net effect of implicit abiotic factors, by which i-th taxon is more likely to present (h_i
> 0) or not ($h_i < 0$), g_{ij} represents the effect of the i-th observed environmental factor, and J_{ij} represents a
co-occurrence pattern of i-th and j-th taxa. Since the logarithm of the probability of a community state is
inversely proportional to $H(\vec{\sigma}^{(k)})$, a community state having lower H is more frequently observed. To
consider dynamics on an assembly graph defined as a network whose 2^S nodes represent possible
community states and the edges represents transition path between them (two community states are
adjacent only if they have the opposite presence/absence status for just one species), we assigned energy to
nodes with the above equation, and so imposed the directionality in state transitions. Then, by using the
steepest descent algorithm (32) we identified the stable state communities as the energy minima of the
weighted network (nodes having the lowest energy compared to all its neighbors), and determined their
basins of attraction based on a steepest descent procedure starting from each node. As environmental
variables, soil pH, electrical conductivity, C/N ratio, and available phosphorous concentration were included
after normalization within the ranges from 0 to 1.

The number of stable states within energy landscapes of community structure was inferred,
respectively, for three types of datasets, namely, the prokaryote community matrix, the fungal matrix, and
the matrix including both prokaryotes and fungi. In addition to the energy landscape analyses based on the
abovementioned family-level delineation of community states, analyses based on community-state
delineation at the order-level were performed. Likewise, additional analyses based on another threshold
value (5 %) in the binarization of community states were conducted. Consequently, we performed energy
landscape analyses in each of the combinations of three types of data (prokaryotes, fungi, and prokaryotes +
fungi), two taxonomic levels (family or order), and two threshold levels (5 % and 10 %). For each
combination (setting), the parameters of the extended pairwise maximum entropy model were adjusted to
fit the model to the empirical data. The number of stable states was then estimated by using the gradient
descent algorithm for 20,000 randomly generated community states within the 2^S possible community
states. In the heuristic search, sets of values of the environmental variables (soil pH, electrical conductivity,
C/N ratio, and available phosphorous concentration) were drawn from those of sampling points.

We then reconstructed energy landscapes for the settings with which community structure could be
characterized by relatively small numbers of stable states (i.e., the family-level fungal data matrix defined at
the 10-% threshold and the order-level fungal data matrix defined at the 10-% threshold; Fig. 4; *SI Appendix*:
Fig. S5). Minor stable states whose shallowest "energy barrier" (i.e., deviation of energy from adjacent
"tipping points") was less than 1/10 of that of the deepest alternative stables states were incorporated into
the basins of deeper stable states [i.e., pruning of alternative stable states (32)]. The "energy" estimates
were then plotted against the PCoA axes mentioned above. Spline smoothing of the landscape was
performed with optimized penalty scores using the mgcv v.1.8-40 package (79) of R.

We then showed the energy of each sample on the two-dimensional surface of the state space
inferred with the above-mentioned PCoA: the horizontal axes indicated PCoA 1 and 2 scores and the vertical
axis indicated the energy of target samples' community states defined at the ASV level. On the energy

landscape, samples belonging to the same alternative stable states were indicated by colors. For the
alternative stable states that represented 10 or more samples, the proportions of disease-level-1 samples
were calculated.

Acknowledgments

We thank the SuperComputer System, Institute for Chemical Research, Kyoto University for the use of super
computers. This work was financially supported by JST PRESTO (JPMJPR16Q6), Human Frontier Science
Program (RGP0029/2019), JSPS Grant-in-Aid for Scientific Research (20K20586), NEDO Moonshot Research
and Development Program (JPNP18016), and JST FOREST (JPMJFR2048) to H.T., JSPS Grant-in-Aid for
Scientific Research (20K06820 and 20H03010) to K.S., and JSPS Fellowship to H.F..

498
499 **References**
500
501

1.   M. Springmann, *et al.*, Options for keeping the food system within environmental limits. *Nature* **562**,
519–525 (2018).

2.   D. Tilman, C. Balzer, J. Hill, B. L. Befort, Global food demand and the sustainable intensification of
agriculture. *Proceedings of the National Academy of Sciences* **108**, 20260–20264 (2011).

3.   H. Toju, *et al.*, Core microbiomes for sustainable agroecosystems. *Nat Plants* **4**, 247–257 (2018).

4.   P. E. Busby, *et al.*, Research priorities for harnessing plant microbiomes in sustainable agriculture.
PLoS Biol **15**, e2001793 (2017).

5.   R. L. Berendsen, C. M. J. Pieterse, P. A. H. M. Bakker, The rhizosphere microbiome and plant health.
Trends Plant Sci **17**, 478–486 (2012).

6.   R. Bommarco, D. Kleijn, S. G. Potts, Ecological intensification: Harnessing ecosystem services for food
security. *Trends Ecol Evol* **28**, 230–238 (2013).

7.   K. G. Peay, P. G. Kennedy, J. M. Talbot, Dimensions of biodiversity in the Earth mycobiome. *Nat Rev
Microbiol* **14**, 434–447 (2016).

8.   B. J. Baker, *et al.*, Diversity, ecology and evolution of Archaea. *Nat Microbiol* **5**, 887–900 (2020).

9.   R. Hayat, S. Ali, U. Amara, R. Khalid, I. Ahmed, Soil beneficial bacteria and their role in plant growth
promotion: A review. *Ann Microbiol* **60**, 579–598 (2010).

10. K. K. Newsham, A meta-analysis of plant responses to dark septate root endophytes. *New Phytol* **190**,
783–93 (2011).

11. M. G. A. van der Heijden, *et al.*, Mycorrhizal fungal diversity determines plant biodiversity, ecosystem
variability and productivity. *Nature* **396**, 69–72 (1998).

12. J. A. Vorholt, C. Vogel, C. I. Carlström, D. B. Müller, Establishing Causality: Opportunities of Synthetic
Communities for Plant Microbiome Research. *Cell Host Microbe* **22** (2017).

13. P. Trivedi, J. E. Leach, S. G. Tringe, T. Sa, B. K. Singh, Plant–microbiome interactions: from community
assembly to plant health. *Nat Rev Microbiol* **18**, 607–621 (2020).

14. J. K. Jansson, K. S. Hofmockel, The soil microbiome — from metagenomics to metaphenomics. *Curr*
Opin Microbiol **43**, 162–168 (2018).

15. M. Mazzola, S. Freilich, Prospects for biological soilborne disease control: Application of indigenous
versus synthetic microbiomes. *Phytopathology* **107**, 256–263 (2017).

16. S. Jiao, Y. Yang, Y. Xu, J. Zhang, Y. Lu, Balance between community assembly processes mediates
species coexistence in agricultural soil microbiomes across eastern China. *ISME Journal* **14**, 202–216
(2020).

17. K. Faust, L. Lahti, D. Gonze, W. M. de Vos, J. Raes, Metagenomics meets time series analysis:
Unraveling microbial community dynamics. *Curr Opin Microbiol* **25**, 56–66 (2015).

18. L. Tedersoo, *et al.*, Global diversity and geography of soil fungi. *Science (1979)* **346**, 1256688 (2014).

19. H. Toju, A. S. Tanabe, H. Sato, Network hubs in root-associated fungal metacommunities. *Microbiome*
6, 116 (2018).

20. J. M. Talbot, *et al.*, Endemism and functional convergence across the North American soil
mycobiome. *Proceedings of the National Academy of Sciences* **111**, 6341–6346 (2014).

21. M. Bahram, *et al.*, Structure and function of the global topsoil microbiome. *Nature* **560**, 233–237
(2018).

22. K. Faust, J. Raes, Microbial interactions: From networks to models. *Nat Rev Microbiol* **10**, 538–550
(2012).

23. H. Toju, S. Yamamoto, A. S. A. S. Tanabe, T. Hayakawa, H. S. H. S. Ishii, Network modules and hubs in
plant-root fungal biomes. *J R Soc Interface* **13**, 20151097–20151097 (2016).

24. M. G. A. van der Heijden, M. Hartmann, Networking in the plant microbiome. *PLoS Biol* **14**, 1–9
(2016).

25. K. Hartman, *et al.*, Deciphering composition and function of the root microbiome of a legume plant.
Microbiome **5**, 1–13 (2017).

26. J. A. Vorholt, C. Vogel, C. I. Carlström, D. B. Müller, Establishing causality: opportunities of dynthetic
vommunities for plant microbiome tesearch. *Cell Host Microbe* **22**, 142–155 (2017).

27.   L. W. Mendes, *et al.*, Soil-borne microbiome: linking diversity to function. *Microb Ecol* **70**, 255–265
553   (2015).

554   28.   H. Toju, *et al.*, Scoring species for synthetic community design: network analyses of functional core
555   microbiomes. *Front Microbiol* **11**, 1361–1361 (2020).

556   29.   J. E. Goldford, *et al.*, Emergent simplicity in microbial community assembly. *Science (1979)* **361**, 469–
557   474 (2018).

558   30.   H. Fujita, *et al.*, Alternative stable states, nonlinear behavior, and predictability of microbiome
559   dynamics. *bioRxiv*, https://doi.org/10.1101/2022.08.23.505041 (2022).

560   31.   B. E. Beisner, D. T. Haydon, K. Cuddington, Alternative stable states in ecology. *Front Ecol Environ* **1**,
561   376–382 (2003).

562   32.   K. Suzuki, S. Nakaoka, S. Fukuda, H. Masuya, Energy landscape analysis elucidates the multistability of
563   ecological communities across environmental gradients. *Ecol Monogr* **91**, 1–21 (2021).

564   33.   M. Scheffer, S. R. Carpenter, Catastrophic regime shifts in ecosystems: Linking theory to observation.
565   *Trends Ecol Evol* **18**, 648–656 (2003).

566   34.   M. Scheffer, S. Carpenter, J. a Foley, C. Folke, B. Walker, Catastrophic shifts in ecosystems. *Nature*
567   **413**, 591–596 (2001).

568   35.   P. J. Mumby, A. Hastings, H. J. Edwards, Thresholds and the resilience of Caribbean coral reefs. *Nature*
569   **450**, 98–101 (2007).

570   36.   A. Hastings, *et al.*, Transient phenomena in ecology. *Science (1979)* **361**, eaat6412 (2018).

571   37.   T. Fukami, Historical contingency in community assembly: integrating niches, species pools, and
572   priority effects. *Annu Rev Ecol Evol Syst* **46**, 1–23 (2015).

573   38.   T. Watanabe, N. Masuda, F. Megumi, R. Kanai, G. Rees, Energy landscape and dynamics of brain
574   activity during human bistable perception. *Nat Commun* **5**, 4765 (2014).

575   39.   O. M. Becker, M. Karplus, The topology of multidimensional potential energy surfaces: Theory and
576   application to peptide structure and kinetics. *Journal of Chemical Physics* **106**, 1495 (1997).

577   40.   D. Schlatter, L. Kinkel, L. Thomashow, D. Weller, T. Paulitz, Disease suppressive soils: New insights
578   from the soil microbiome. *Phytopathology* **107**, 1284–1297 (2017).

579   41.   L. L. Kinkel, M. G. Bakker, D. C. Schlatter, A coevolutionary framework for managing disease-
580   suppressive soils. *Annu Rev Phytopathol* **49**, 47–67 (2011).

581   42.   R. Mendes, *et al.*, Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science*
582   **332**, 1097–100 (2011).

583   43.   J. Yuan, *et al.*, Predicting disease occurrence with high accuracy based on soil macroecological
584   patterns of Fusarium wilt. *ISME Journal* **14**, 2936–2950 (2020).

585   44.   K. W. T. Goulding, Soil acidification and the importance of liming agricultural soils with particular
586   reference to the United Kingdom. *Soil Use Manag* **32**, 390–399 (2016).

587   45.   D. L. Corwin, S. M. Lesch, Apparent soil electrical conductivity measurements in agriculture. *Comput
588   Electron Agric* **46**, 11–43 (2005).

589   46.   A. Frostegård, E. Bååth, The use of phospholipid fatty acid analysis to estimate bacterial and fungal
590   biomass in soil. *Biol Fertil Soils* **22**, 59–65 (1996).

591   47.   J. Poveda, *Trichoderma* as biocontrol agent against pests: New uses for a mycoparasite. *Biological
592   Control* **159**, 104634 (2021).

593   48.   F. Tekaia, J. P. Latgé, *Aspergillus fumigatus*: Saprophyte or pathogen? *Curr Opin Microbiol* **8**, 385–392
594   (2005).

595   49.   Y. Yang, E. Yang, Z. An, X. Liu, Evolution of nematode-trapping cells of predatory fungi of the
596   Orbiliaceae based on evidence from rRNA-encoding DNA and multiprotein sequences. *Proc Natl Acad
597   Sci U S A* **104**, 8379–8384 (2007).

598   50.   K. L. Lazarus, G. L. Benny, H. M. Ho, M. E. Smith, Phylogenetic systematics of syncephalis (zoopagales,
599   zoopagomycotina), a genus of ubiquitous mycoparasites. *Mycologia* **109**, 333–349 (2017).

600   51.   J. M. Bourne, B. R. Kerry, F. A. A. M. de Leij, The importance of the host plant on the interaction
601   between root-knot nematodes (*Meloidogyne* spp.) and the Nematophagous fungus, *Verticillium
602   chlamydosporium* Goddard. *Biocontrol Sci Technol* **6**, 539–548 (1996).

603   52.   P. G. Kennedy, K. G. Peay, T. D. Bruns, Root tip competition among ectomycorrhizal fungi: are priority
604   effects a rule or an exception? *Ecology* **90**, 2098–107 (2009).

605   53.   G. D. A. Werner, E. T. Kiers, Order of arrival structures arbuscular mycorrhizal colonization of plants.
606   *New Phytologist* **205**, 1515–1524 (2015).

607   54.   M. M. M. Kuypers, H. K. Marchant, B. Kartal, The microbial nitrogen-cycling network. *Nat Rev
608   Microbiol* **16**, 263–276 (2018).

609   55.   J. K. Jansson, K. S. Hofmockel, Soil microbiomes and climate change. *Nat Rev Microbiol* **18**, 35–46
610   (2020).

611   56.   E. M. Onkendi, G. M. Kariuki, M. Marais, L. N. Moleleki, The threat of root-knot nematodes
612   (Meloidogyne spp.) in Africa: A review. *Plant Pathol* **63**, 727–737 (2014).

613   57.   M. Dita, M. Barquero, D. Heck, E. S. G. Mizubuti, C. P. Staver, Fusarium wilt of banana: Current
614   knowledge on epidemiology and research needs toward sustainable disease management. *Front
615   Plant Sci* **9**, 1468 (2018).

616   58.   A. Hartmann, M. Schmid, D. van Tuinen, G. Berg, Plant-driven selection of microbes. *Plant Soil* **321**,
617   235–257 (2008).

618   59.   S. Banerjee, K. Schlaeppi, M. G. A. van der Heijden, Keystone taxa as drivers of microbiome structure
619   and functioning. *Nat Rev Microbiol* **16**, 567–576 (2018).

60.   H. Toju, *et al.*, Species-rich networks and eco-evolutionary synthesis at the metacommunity level. *Nat*
Ecol Evol **1**, 0024 (2017).

61.   A. Kassam, T. Friedrich, R. Derpsch, Global spread of conservation agriculture. *International Journal of*
Environmental Studies **76**, 29–51 (2019).

62.   L. Schreefel, R. P. O. Schulte, I. J. M. de Boer, A. P. Schrijver, H. H. E. van Zanten, Regenerative
agriculture – the soil is the base. *Glob Food Sec* **26**, 100404 (2020).

63.   Y. Li, Q. Zhang, Y. Cai, Q. Yang, S. X. Chang, Minimum tillage and residue retention increase soil
microbial population size and diversity: Implications for conservation tillage. *Science of the Total*
Environment **716**, 137164 (2020).

64.   E. Verbruggen, E. Toby Kiers, Evolutionary ecology of mycorrhizal functional diversity in agricultural
systems. *Evol Appl* **3**, 547–560 (2010).

65.   UNDP, SCBD, UNEP-WCMC, "Creating a nature-positive future: the contribution of protected areas
and other effective area-based conservation measures" (2021).

66.   J. G. Caporaso, *et al.*, Global patterns of 16S rRNA diversity at a depth of millions of sequences per
sample. *Proc Natl Acad Sci U S A* **108**, 4516–4522 (2011).

67.   A. Apprill, S. Mcnally, R. Parsons, L. Weber, Minor revision to V4 region SSU rRNA 806R gene primer
greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology* **75**, 129–137 (2015).

68.   D. S. Lundberg, S. Yourstone, P. Mieczkowski, C. D. Jones, J. L. Dangl, Practical innovations for high-
throughput amplicon sequencing. *Nat Methods* **10**, 999–1002 (2013).

69.   H. Toju, H. Kurokawa, T. Kenta, Factors influencing leaf- and root-associated communities of bacteria
and fungi across 33 plant orders in a grassland. *Front Microbiol* **10**, 1–14 (2019).

70.   J. L. Stevens, R. L. Jackson, J. B. Olson, Slowing PCR ramp speed reduces chimera formation from
environmental samples. *J Microbiol Methods* **93**, 203–205 (2013).

71.   M. Ushio, Interaction capacity as a potential driver of community diversity. *Proceedings of the Royal*
Society B: Biological Sciences **289**, 20212690 (2022).

72.   M. Ushio, *et al.*, Quantitative monitoring of multispecies fish environmental DNA using high-
throughput sequencing. *Metabarcoding Metagenom* **2**, 1–15 (2018).

73.   H. Toju, A. S. A. Tanabe, S. Yamamoto, H. Sato, High-coverage ITS primers for the DNA-based
identification of ascomycetes and basidiomycetes in environmental samples. *PLoS One* **7**, e40863
(2012).

74.   M. Hamady, J. J. Walker, J. K. Harris, N. J. Gold, R. Knight, Error-correcting barcoded primers for
pyrosequencing hundreds of samples in multiplex. *Nat Methods* **5**, 235–237 (2008).

75.   A. S. Tanabe, H. Toju, Two new computational methods for universal DNA barcoding: a benchmark
using barcode sequences of bacteria, archaea, animals, fungi, and land plants. *PLoS One* **8**, e76910
(2013).

76. B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data. *Nat*
Methods **13**, 581–583 (2016).

77. R Core Team, R: A language and environment for statistical computing. *R: A language and*
environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL
https://www.R-project.org/ (2020).

78. S. Põlme, *et al.*, FungalTraits: a user-friendly traits database of fungi and fungus-like stramenopiles.
Fungal Divers **105**, 1–16 (2020).

79. S. Wood, mgcv: Mixed GAM computation vehicle with automatic smoothness estimation available at
https://cran.r-project.org/web/packages/mgcv/index.html (2022).

664
665

666
667

Figure 1. Diversity of agroecosystem microbiomes. (*A*) Map of research sites across the Japanese
Archipelago. The 2,903 soil samples were grouped into 42 research sits were when their latitude and
longitude profiles were round to one decimal place. (*B*) Taxonomic compositions of prokaryotes (archaea
and bacteria; top) and fungi (bottom) at the family level. See *SI Appendix*: Fig. S2 for results at the genus,
order, and class levels. (*C*) Compositions of functional groups of fungi. (*D*) Comparison of α-diversity
between prokaryotes and fungi. Mean Shannon diversity at the family level was shown for each research
site. Bars represent standard deviations. (*E*) Mean community dissimilarity (β-diversity) between pairs of
samples within each research site. (*F*) Community dissimilarity (β-diversity) in mean community
compositions between pairs of research sites.

678
679
Figure 2. Dimensions of soil microbiome structure. (*A*) Prokaryote and fungal community structure. Principal
co-ordinate analyses (PCoA) were performed based on ASV-level compositional matrices respectively for the
prokaryote and fungal communities. The identify of crop plants is shown by colors. See *SI Appendix*: Figs S3-4
for relationship between community structure and environmental factors. (*B*) Correlation between PCoA
scores and soil environmental factors. (*C*) Microbiome structure and crop disease levels. (*D*) Relationship
between prokaryote/fungal community structure on the disease level of crop plants. A GLMM of crop plants'
disease level (disease level 1 vs. disease levels 2-5) with a logit-link function and binomial errors was
constructed by setting principal coordinates of prokaryote/fungal community structure as explanatory
variables (fixed effects). The identity of research projects and sampling month were included as random
effects in the GLMM.
690
691

692
693
Figure 3. Balance between prokaryote and fungal abundance. (*A*) Prokaryote/fungus ratio. The ratio of
prokaryote abundance to fungal abundance was evaluated based on estimated DNA concentrations
[$\log \left(\frac{\text{prokaryote 16S rRNA gene concentration(DNA copies/μL)}}{\text{fungal ITS gene concentration (DNA copies/μL)}} \right)$]. The prokaryote/fungus ratio is shown for each
sample on the PCoA surfaces of prokaryote and fungus community compositions. (*B*) Effects of
prokaryote/fungus ratio on the disease level of crop plants. In each GLMM with a logit-link function and
binomial errors, crop plants disease level [least-diseased (disease level 1) = 1, severely-diseased (disease
level 2-5) = 0] was included as response variable. In addition to bacterium/fungal abundance ratio, soil
environmental variables (pH and electrical conductivity; Model 1) and α-diversity estimates of prokaryotes
and fungi (Model 2) were included as explanatory variables (fixed effects). The identity of research projects
and sampling month were included as random effects in the GLMM. A negative estimate indicates that the
proportion of least-diseased plants decreased with increasing values of a target explanatory variable. (*C*)
Representative relationship between prokaryote/fungus ratio on crop disease. For each of the five research
projects with enough number of disese-level-1 and disease-level-2-5 samples, a GLM was constructed. For
the two research projects with significant effects of prokaryote/fungus ratio on crop disease level, the
prokaryote/fungus ratios at which 50 % of samples display disease-level-1 symptoms are shown.
709
710

711
712
Figure 4. Number of the stable states of community structure. The parameters of the extended pairwise
maximum entropy model were adjusted to fit the model to the empirical data (see Methods). The number of
stable states in the energy landscape was then estimated by using the gradient descent algorithm for 20,000
randomly generated community states within the 2^S possible community states, where S denotes the
number of families or orders. In the heuristic search, sets of values of the environmental variables (soil pH,
electrical conductivity, C/N ratio, and available phosphorous concentration) were drawn from those of
sampling points. The boxplots show variation in the number of stable states across different sets of
environmental variables. A box represents the first quartile, median and the third quartile, while bars
indicate the largest/smallest values within 1.5 × inter-quartile ranges from the box: outliers were indicated
as circles. The analysis was performed, respectively, for the dataset representing both prokaryotes and fungi,
the prokaryote dataset, and the fungus dataset. The threshold value in the binarization of the community
data was set at 10 %: see *SI Appendix*: Fig. S5 for results based on another threshold value (5 %).

726
727
Figure 5. Alternative stable states of microbiomes. (*A*) Energy landscape of fungal community structure. The
surface of energy levels was reconstructed on the PCoA dimensions (Fig. 2) of fungal community structure
based on spline smoothing. Community states with lower energy are inferred to be more stable. Data points
(samples) indicated by the same color belong to the basins of the same alternative stable states (defined at
the family-level taxonomic compositions), whose energy is lower than any adjacent community structure.
See panel *C* for details about the alternative stable states. (*B*) Surface of crop disease conditions. The area of
community structure with less diseased states and lower energy (i.e., community stability; panel *A*) are
expected to represent stably-manageable microbiomes with potential disease-suppressive functions. (*C*)
Definitions of the stable states. To reduce the complexity of input data to a computationally feasible range,
alternative stable states were explored based on family-level community compositions (threshold for
binarization = 10 %). The 10 alternative stable states representing ≥ 10 samples were compared in terms of
their associations with crop disease conditions.

740
741