1   **Enterotoxigenic *Escherichia coli* heat-labile toxin drives enteropathic changes in small**
2   **intestinal epithelia**
3

4   Alaullah Sheikh[1], Brunda Tumala[1], Tim J. Vickers[1], John C. Martin[2], Bruce A. Rosa[1,2], Subrata Sabui[5,6], Supratim Basu[1], Rita D. Simoes[1],
5   Makedonka Mitreva[1,2], Chad Storer[2,3], Erik Tyksen[2,3], Richard D. Head[2,3], Wandy Beatty[4], Hamid M. Said[5,6], and James M. Fleckenstein[1,7].
6
7   Division of Infectious Diseases[1], Department of Medicine, Washington University School of Medicine, St. Louis, MO 63110, USA.
8
9   The McDonnell Genome Institute[2], Washington University School of Medicine, St. Louis, MO 63110, USA.
10
11   Department of Genetics[3], Washington University School of Medicine, St. Louis, MO 63110, USA.
12
13   Department of Molecular Microbiology[4], Washington University School of Medicine, Saint Louis, Missouri, USA 63110.
14
15   Departments of Medicine and Physiology/Biophysics, School of Medicine, University of California-Irvine, Irvine, CA[5]; 92697.
16
17   Department of Research, VA Medical Center, Long Beach, CA[6]; 90822
18
19   Infectious Diseases, Medicine Service, Veterans Affairs Saint Louis Health Care System, Saint Louis Missouri, USA 63106[7].
20

corresponding author
James M. Fleckenstein
Division of Infectious Diseases
Department of Medicine
Washington University School of Medicine
Campus Box 8051
660 South Euclid Avenue
Saint Louis Missouri, USA 63110
p 314-362-9218
jfleckenstein@wustl.edu

31
32
33

abstract

Enterotoxigenic *E. coli* (ETEC), produce heat-labile (LT) and/or heat-stable (ST) enterotoxins, and

are a common cause of diarrhea in children of resource-poor regions. ETEC have also been linked

repeatedly to poorly understood sequelae including enteropathy, malnutrition, and growth

impairment. While the cellular actions of ETEC enterotoxins leading to diarrhea are well-established,

their potential contribution to subsequent pathology is unclear. LT stimulates cellular cAMP

production to activate protein kinase A (PKA) which phosphorylates cellular ion channels that drive

export of salt and water into the intestinal lumen resulting in diarrhea. However, as PKA exhibits

broad kinase activity and its activated catalytic subunits modulate transcription of many genes, we

interrogated the transcriptional profiles of LT-treated small intestinal epithelia. These studies

demonstrated toxin-induced changes in hundreds of genes including those required for biogenesis

and function of the brush border, the major site absorption of nutrients, and suppression of a key

transcription factor, HNF4, critical to differentiation of intestinal epithelia. Accordingly, LT treatment

of intestinal epithelial cells significantly disrupted the absorptive microvillus architecture and altered

transport of essential nutrients. In addition, challenge of neonatal mice with LT-producing ETEC

recapitulated the architectural derangement of the brush border while maternal vaccination with LT

prevented brush border disruption in ETEC-challenged neonatal mice. These findings highlight

impacts of ETEC enterotoxins beyond acute diarrheal illness and may inform approaches to mitigate

and prevent major sequelae including malnutrition that impact millions of young children.

Introduction

Infectious diarrhea remains a leading cause of death and morbidity among young children in low-

middle income countries where access to clean water and sanitation remain in short supply[1].

Enterotoxigenic *E. coli* (ETEC), initially discovered as a cause of severe, cholera-like illness[2], are one

of the most common pathogens associated with moderate-severe diarrhea among children under

the age of five years[3,4], and are perennially the most common cause of diarrhea in travelers[5] to

endemic regions where these organisms are thought to account for hundreds of millions of cases of

diarrheal illness each year[6].

Importantly, ETEC infections have been linked to non-diarrheal sequelae including

"environmental enteric dysfunction (EED)," a condition characterized by impaired nutrient

absorption, impaired growth[7,8] and malnutrition,[9,10] adding significantly to the morbidity as well as

deaths from diarrhea and other infections[11]. The risk of stunting multiplies with each episode of

diarrheal illness in children under the age of two years[12], a period during which children residing in

impoverished areas commonly sustain multiple ETEC infections[8]. However, the molecular

pathogenesis underlying the intestinal changes associated with EED, and the contribution of

individual pathogens including ETEC remain poorly understood.

Similarly, toxin-producing *E. coli* have also been repeatedly identified in patients with tropical

sprue[13-15], a condition classically described in adults residing for extended periods of time in areas

where ETEC diarrheal disease is common. Like EED, tropical sprue is associated with changes to the

small intestinal villous architecture, including ultrastructural alteration of the epithelial brush border

formed by the microvilli[16,17], nutrient malabsorption, and wasting.

The basic molecular mechanisms underpinning acute watery diarrhea caused by ETEC are well-

established[18]. ETEC produce heat-labile (LT) and/or heat-stable (ST) enterotoxins that activate

production of cAMP and cGMP second messengers respectively, leading to activation of cellular

kinases that in turn modulate the activity of sodium and chloride channels in apical membrane of

intestinal epithelial cells to promote net efflux of salt and water into the intestinal lumen resulting in

watery diarrhea.

LT and cholera toxin (CT) share ~85 % amino acid identity, and both toxins exert their major

effects on the cell through the ADP-ribosylation of the alpha subunit of Gs (Gsα), a stimulatory

intracellular guanine nucleotide-binding protein. Inhibition of Gsα GTPase activity leads to

constitutive activation of adenylate cyclase and increased production of intracellular cAMP[19].

In its central role as a second messenger, cAMP governs a diverse array of cellular processes[20]

and modulates transcription of multiple genes through a number of cAMP-responsive transcriptional

activators and repressors[21] . cAMP activates protein kinase A (PKA), a heterotetramer, by liberating

its two regulatory subunits from the catalytic subunits which are then free to phosphorylate a wide

variety of cytoplasmic and nuclear protein substrates[22]. PKA largely regulates transcription by

phosphorylation of transcription factors including the cyclic AMP response element binding protein

(CREB) and the cAMP-response element modulator (CREM) which bind cAMP response elements

(CRE) in the promoter regions of target genes[21-23].

Notably, cholera toxin (CT), LT, and dibutyryl-cyclic AMP all induce hypersecretion and impact the

architecture of gastrointestinal epithelia in rodent small intestine [24], while small intestinal biopsies of

patients with acute cholera exhibit marked changes in the ultrastructure of the intestinal brush

border, the major absorptive surface in the small intestine[25,26], including shortening and disruption

of the microvilli. Consistent with these observations, studies of young children less than two years of

age in Bangladesh have specifically associated LT-producing ETEC with undernutrition[27], suggesting

that heat-labile toxin may exert effects on intestinal mucosa that extend beyond acute diarrheal

illness.

Here we demonstrate that in addition to the canonical effects of LT on cellular export of salt and

water into the intestinal lumen, this toxin impacts multiple genes involved in formation of microvilli,

resulting in marked alteration of the architecture of the intestinal brush border, the major site of

nutrient absorption in the small intestine. These effects are compounded by alteration of solute

transporters within the brush border epithelia, potentially disrupting the absorption of multiple

essential nutrients.

Results

heat-labile toxin markedly alters the transcriptomes of intestinal epithelial cells.

Commensurate with the importance of cAMP as a second messenger, we found that compared to

either untreated controls or cells treated with a catalytically inactive (E211K) mutant of LT, wild type

LT holotoxin substantially modulated transcription of many genes in intestinal epithelial cells. In

RNAseq studies of polarized Caco-2 intestinal epithelial cells we found that 3,832 genes were

significantly ($p \leq 10^{-5}$) upregulated and 3687 downregulated in response to LT, while binding of the

inactive toxin failed to induce significant changes in the transcriptome (figure 1a). However, Caco-2

cells are derived from distant metastases of a colon cancer tumor in which transcriptomes would

likely be altered relative to untransformed intestinal epithelia[28,29], and cAMP signaling is known to be

aberrant in some transformed cells[23]. Therefore, to examine a more physiologically relevant target,

we next examined the impact of LT on differentiated small intestinal enteroids. Here, we found that

far fewer genes were differentially expressed ($\leq 10^{-5}$) with 746 significantly upregulated, and 561

downregulated in response to intoxication with LT (figure 1b). Notably, however we found

substantial statistically significant overlap in genes significantly modulated in Caco-2 cells and

enteroids (supplemental table 3, supplemental dataset 1) with the transcription of hundreds of

genes significantly up- or down-regulated in both groups. Gene ontology enrichment analysis as

well as ontology-independent investigation of genes (CompBio, supplemental figure 1) modulated

by the toxin highlighted multiple cellular components associated with both the development and

function of the absorptive surface of the small intestine (supplemental dataset 2).

heat-labile toxin impairs development of small intestinal microvilli.

Small intestinal enterocytes are each covered with hundreds of microvilli, complex structures

comprised of a central core of actin filaments within protrusions of the plasma membrane.

Collectively, the luminal surface of the intestine formed by these microvilli, known as the brush

border, represents the major absorptive surface of the gastrointestinal tract. Three major classes of

proteins are required for the biogenesis of microvilli[30] (figure 2a). These include (1) proteins such as

villin, epsin, plastin, and EPS8 that bundle parallel clusters of actin filaments; BAIP2L1 (IRTKS)

responsible for recruiting the EPS bundling protein to the tips of microvilli[31]; (2) ezrin, myo1a[32],

myo6 that link the actin cytoskeleton with the plasma membrane; and (3) protocadherin molecules

CDHR2 and CDHR5 engaged in extracellular heterotypic complexes between the tips of the

microvilli[30] that are stabilized by a tripartite complex of MYO7B, ANKS4B[33], and USH1C[33].

Interrogation of transcriptional profiles indicated that the transcription of each of these classes of

genes was significantly altered following exposure to wild type heat-labile toxin (figure 2b). Similarly,

RT-PCR confirmed decreased expression of multiple genes involved in microvilli biogenesis,

including VIL1 encoding villin (figure 3a), and we were able to demonstrate that production of villin

was depressed in polarized small intestinal enteroids (figure 3b). In addition, TEM images of

polarized small intestinal enteroids exposed to heat-labile toxin demonstrated significantly

shortened and disorganized microvillus structures on the apical surface of enterocytes (figure 3c).

heat-labile toxin modulates the transcription of multiple brush border nutrient transport

genes.

The human solute carrier (SLC) gene superfamily is comprised of more than 50 gene families

thought to encode more than 300 functional transporters[34]. Many of the SLC proteins are enriched in

the small intestinal brush border where they transport critical nutrients including amino acids,

oligopeptides, sugars, and vitamins. We found that transcription of many SLC genes was altered in

Caco-2 cells as well as small intestinal enteroids (figure 4a). These included transporters for Zinc,

known both to be deficient in children with enteropathy[35], and a micronutrient critical for intestinal

homeostasis. Likewise transcription of SLC19A3 encoding the principal SLC responsible for uptake

of the water-soluble B vitamin thiamine (vitamin B1)[36] by differentiated intestinal epithelial cells lining

the surface of intestinal villi of the proximal small intestine[37,38](figure 4b) was repressed. Moreover,

we found that LT treatment of human small intestinal organoids also interfered with transcription of

the cis regulatory element specificity protein 1 (SP-1) previously shown to govern the transcription of

SLC19A3[39-41]. Finally, we found that thiamine transport was significantly depressed following

exposure to LT (figure 4c) providing additional evidence that ETEC can impair transport of critical

nutrients.

LT-producing ETEC disrupt the absorptive architecture of the small intestine

To further study the potential impact of ETEC toxins on the intestinal architecture, we performed

challenge studies in infant mice. Compared to sham-challenged (PBS) controls, or mice challenged

with a toxin deficient strain of ETEC, we again noted down-regulation of genes involved in F-actin

bundling, membrane cross-linking, and intermicrovillus adhesion complex formation, all required for

intestinal microvilli (figure 5a) biogenesis. Likewise, on examination of small intestinal villi we found

that production of villin in enterocyte brush borders was substantially decreased relative to sham

challenged controls (figure 5b-d). In mice challenged with a wild type ETEC isolate that makes ST

and LT, but not a LT/ST-toxin-negative mutant (jf4763, supplemental table 1), we observed

significant alteration in architecture of the intestinal brush border with significant shortening and

disorganization of the microvilli (figure 5e,f; supplemental figure 2a-d) reminiscent of the earlier

ultrastructural studies of patients with tropical sprue [17] and *V. cholerae* infections[25]. Similarly, we

found that in mice challenged with a strain containing an isogenic mutation in *eltA* (jf571,

supplemental table 1) encoding the LT A subunit, which still makes heat-stable toxins, the microvillus

architecture was preserved (figure 5g, h), suggesting that LT is the principal toxin underlying the

enteropathic changes to the enterocyte surfaces. However, despite the dramatic toxin-dependent

changes to the absorptive surface of the intestine, the growth kinetics of suckling mice challenged

with either wild type bacteria or toxin deletion mutants were surprisingly similar (supplemental figure

3a,b) and paralleled those of sham challenged controls, suggesting that other elements including

caloric restriction, and alterations in gut microbiota may be required to replicate growth faltering

observed in children[42].

maternal vaccination with LT prevents brush border disruption

To address whether enteropathic changes to the small intestine can be prevented by vaccination,

and to further define the role of LT, we vaccinated mouse dams with heat-labile toxin and examined

the brush border ultrastructure in suckling mice. Vaccinated dams but not sham vaccinated controls

expressed significant levels of IgA and IgG in breast milk (figure 6a), consistent with increased levels

of antibodies in the stomachs of infant mice (figure 6b). Notably, we found that maternal vaccination

with LT completely abrogated changes to the microvilli (figure 6c,d) further substantiating the

importance of LT in driving changes to the epithelial architecture.

LT modulates key transcription factors that govern enterocyte development

Despite the marked alteration in transcription mediated by LT the majority of genes critical for

brush border development lacked conserved CRE sites [23]. Therefore, we performed transcription

factor target enrichment analysis[43,44] to identify potential transcription factors responsible for

differential regulation of genes significantly ($p < 0.05$) upregulated or downregulated by LT in both

Caco-2 and enteroid RNA-seq datasets (supplemental table 4, supplemental dataset 1). The

upregulated genes were most significantly ($p = 4.2 \times 10^{-3}$) linked to transcription factor targets of AP-

1 encoded by the *c-jun* gene, previously shown to be regulated by cAMP[45], and to be involved in

intestinal epithelial repair[46]. Notably however, downregulated genes were most significantly

enriched in targets for the HNF4α transcription factor or its intestine specific paralog HNF4γ (P =

1.9×10^{-4}) that were recently shown to regulate multiple genes required for brush border

development[47,48]. Importantly, PKA has also been shown to phosphorylate HNF4 α at a consensus

recognition site within the DNA binding domain, shared with HNF4 γ, inhibiting transcription[49].

Of note, the transcription of both paralogs was found to be significantly depressed following

exposure of intestinal epithelia to LT (figure 7A,B), as were levels of HNF4γ in nuclear fractions from

toxin-treated cells (figure 7C), suggesting that activation of cAMP can interfere with transcription

mediated by HNF4. To further examine the impact of LT on HNF4-mediated transcription we

introduced a transcriptional reporter plasmid containing 6 tandem copies of the HNF4

transcriptional response element (5'- CAAAGGTCA-3') linked to a human codon optimized *Gaussia*

princeps luciferase into Caco-2 cells. These assays demonstrated that HNF4-mediated transcription

was dramatically reduced in cells treated with LT (figure 7D). Similarly, we found that relative to

nuclei of intestinal epithelia cells in ileal segments from sham-challenged control mice, those from

ETEC-challenged mice exhibited significantly less HNFγ (figure 7E) further supporting a role for

ETEC in modulating the production of this important transcription factor. Chen, *et al* described a

"feed-forward regulatory module" essential to enterocyte differentiation in which HNF4 and SMAD4

transcription factors reciprocally activate each other's transcription. As would be predicted from this

model, we found that transcription of SMAD4 was also impaired by LT (supplemental figure 4A),

leading us to propose a model in which LT mediates phosphorylation of HNF4 by PKA, interrupting

this critical transcription module (Supplemental figure 4B-D).

Collectively, the current studies demonstrate that in addition to their known canonical effects on

ion transport that culminate in watery diarrhea, ETEC toxins can drive appreciable derangement of

enterocyte architecture and function by interfering with key pathways in intestinal epithelia that

govern the formation of mature enterocytes capable of effective nutrient absorption. These findings

have important implications for our understanding and prevention of enteropathic conditions linked

to ETEC.

Discussion

Understanding the molecular events that lead to sequelae of undernutrition and growth faltering

following ETEC infections may be key to the effective design of prevention strategies including

vaccines[50]. Although the molecular mechanisms involved in the fluid and ion fluxes into the intestinal

lumen leading to diarrhea are firmly established, only recently has evidence emerged to suggest

that ETEC toxins may incite previously unappreciated changes in small intestinal epithelia[51,52]. The

present studies, initiated to identify additional effects of LT, were prompted by an appreciation that

cyclic nucleotides, particularly cAMP, govern a multitude of cellular pathways potentially resulting in

collateral impacts that extend beyond the acute episodes of diarrhea. Consistent with this model, we

found that exposure of intestinal cells to heat-labile toxin altered the transcription of hundreds of

genes. A central theme highlighted in the analysis of these transcriptional alterations is that LT

affects major classes of genes involved in biogenesis of microvilli and function of the intestinal brush

border, the major site of nutrient uptake in the small intestine, potentially offering a direct molecular

link to sequelae of malnutrition and impaired growth in children.

The potential clinical relevance of the observations reported here is highlighted by remarkably

similar ultrastructural alteration of intestinal epithelial cells seen in small intestinal biopsies of

patients with acute cholera[25] and tropical sprue[17]. In both entities, the brush border was noted to be

abnormal, with shortened, irregular microvilli. Importantly, however despite the structural and

functional similarity between LT and CT, clinical cholera, unlike ETEC infections, has not been linked

to enteropathy or attendant sequelae. Whether this relates to the repetitive nature of ETEC

infections compared to the durable protective immunity that follows a single *V. cholerae* infection[53]

is not presently clear. Similarly, while tropical sprue remains a leading cause of malabsorption in

regions where infectious diarrhea is prevalent[54-56], the most debilitating forms of this illness have not

typically followed isolated cases of traveler's diarrhea, but occur in resident populations or

expatriates[57] repeatedly assailed by diarrhea while residing in endemic regions[58].

The negative impacts of LT on brush border architecture with commensurate reduction in surface

area available for nutrient absorption, are compounded by the alteration of multiple SLC genes that

encode transporters critical for uptake of essential vitamins and other molecules. Importantly, small

intestinal biopsies obtained from Zambian children with enteropathy and refractory stunting

exhibited similar changes in SLC gene expression profiles[59]. The decreased transcription of

molecules required for intestinal zinc uptake, in cells treated with LT is particularly intriguing given

the known aberrations in zinc absorption in children with enteropathy[35,60,61], the possible

contribution of zinc deficiency to enteropathic changes[62], and the salutary effects of zinc in treatment

of children with diarrhea[63,64].

Further study will be needed to precisely delineate the role of ETEC LT and ST enterotoxins, alone

and in combination, in driving enteropathic changes to the intestine and sequelae. ST-producing

ETEC were most strongly associated with moderate to severe diarrhea in GEMS[3], and follow-on

studies of children enrolled in these studies have also linked infections with ETEC encoding heat-

stable toxin to growth faltering[7]. However, our earlier analysis of a global collection of more than

1100 ETEC isolates, including those collected in GEMS, demonstrated that slightly more than half of

all ST-encoding strains also encoded LT, and that roughly one third of the isolates overall encoded

LT alone, ST-LT, or ST only[65]. While LT-producing ETEC have been specifically linked to malnutrition

among children in Bangladesh[27], and our *in vitro* and animal studies point to the potential

importance of LT, additional effort will be needed to correlate toxin-induced morphologic and

functional perturbation of the intestinal brush border with outcomes in children. Importantly, the

long-term morbidity associated with ETEC infections does not appear to correlate with the severity

of diarrhea as both mild illness[4], and perhaps asymptomatic colonization may lead to growth

faltering.

Further refinement of animal models that can faithfully recapitulate features of enteropathy are

also needed. Despite marked derangement of the brush border following infection, single infections

of neonatal mice did not lead to sustained growth defects suggesting it may be difficult to

completely capture the full effects of ETEC in a small animal model. Indeed, conventional mice lack

genes that could be required to reproduce the full effects of ETEC. For instance, each of the

carcinoembryonic antigen cell adhesion molecules (CEACAMs) that are substantially upregulated on

human small intestinal epithelia in response to LT, and which we have recently shown to play a

critical role in ETEC interactions with human small intestine[51], are absent in mice.

While the precise mechanism underlying LT-mediated modulation of genes required for

microvillus biogenesis and absorptive function of the brush border is presently unclear, stimulation

of adenylate cyclase invokes many cAMP-responsive nuclear factors that may serve either to activate

or repress transcription. Genes implicated in development of microvilli are mostly devoid of

consensus palindromic (TGACGTCA) or "half" (CGTCA) cAMP response element (CRE) sites within

their promoter regions for direct modulation by CREB[66], which typically is involved as a

transcriptional activator, and both CREB and CREM can yield several alternatively spliced variants

that may act as either activators or repressors[67]. cAMP second messaging also engages multiple

signaling pathways converging at CREB[68], and PKA can phosphorylate and modulate the activity of

multiple transcription factors to act either as transcriptional activators or repressors, including SP1[69].

Notably, putative binding sites for HNF4, a cAMP-modulated transcription factor[49] known to regulate

genes needed for formation of microvilli[47,48], were significantly enriched in the promotors of genes

downregulated by LT. Both HNF4α and HNF4γ possess canonical PKA recognition sites within their

DNA binding motifs, and PKA phosphorylation of these sites interrupts transcription[70]. HNF4

activates the transcription of SMAD4, and in turn SMAD4 activates transcription of HNF4[71]. Both

transcription factors then engage genes needed for effective differentiation of stem cells to mature

enterocytes[71]. Modulation of the activity of this transcription factor by LT would therefore be

predicted to have a marked impact on pathways critical to intestinal epithelial homeostasis.

We should also note that increases in cellular cAMP can impact multiple cellular pathways

independent of PKA. Included among these are pathways governed by a more recently discovered

family of cellular cAMP-binding molecules, exchange proteins activated by cAMP (EPACs). EPACs

appear to play critical roles as guanine exchange factors that regulate GTPase proteins[72] and are in

involved in complex signaling networks implicated in cell growth, differentiation, and

morphogenesis[73,74].

cAMP can also exert potent epigenetic influences on transcription. CREB-binding protein (CBP)

possesses intrinsic histone acetyl-transferase (HAT) activity[75], and can therefore modulate chromatin

remodeling, enhancing access of transcription factors. In addition, cAMP messaging through PKA

leads to phosphorylation-dependent activation of the histone demethylase enzyme PHF2 to

promote the transcription of multiple genes that can impact transition from stem cells to epithelial

cells[76].

Altogether it seems likely that multiple pathways governed by increases in cellular cAMP may

underlie the morphologic and functional disruption of the brush border epithelial observed in our

studies. Nevertheless, the data presented here provide compelling evidence that the heat-labile

toxin of ETEC ultimately impacts multiple genes required for the biogenesis and function of the

brush border, the major site of nutrient absorption in the human small intestine. The findings may

have significant implications for our understanding of sequelae linked to ETEC including

environmental enteropathy in young children, and tropical sprue in adults. The increased

acknowledgement of long-term morbidity linked to ETEC and an improved understanding of the

role of ETEC enterotoxins as drivers of this morbidity may also strengthen the case for vaccines[50]

specifically engineered to prevent both the acute illness and sequelae.

Methods

Culture and differentiation of Enteroids

Biopsy samples from adult patients undergoing routine endoscopy were obtained at Washington

University School of Medicine after patient consent and approval from the Institutional review board.

Purified crypt cells from the ileum were re-suspended in Matrigel (BD Bisosciences, San Jose,

California, USA) and 15 μL of re-suspended matrix gel was added to each well in 24 well plates.

Plates were incubated at 37° C and 5% CO2 with 50% L-WRN conditioned media (CM) and 50%

primary culture medium (Advanced DMEM/F12, Invitrogen) supplemented with 20% FBS, 2 mM L-

glutamine, 100 units/mL penicillin, 0.1 mg/mL streptomycin, 10 μM Y-27632 (ROCK inhibitor, Tocris

Bioscience, R&D systems, Minneapolis, MN, USA), and 10 μM SB431541 (TGFBR1 inhibitor, Tocris

Bioscience, R&D systems).

To induce differentiation and polarization of enteroids, cells were washed once to remove

Matrigel, followed by trypsinization and centrifugation at 1100 x g for 5 minutes. Cells were then re-

suspended in 1:1 CM and primary medium with Y-27632 and SB431541 as described above. Cells

were then plated on semiporous filters (Transwells®, 6.5 mm insert, 24 well plate, 0.4 μm polyester

membrane, Corning Incorporated, Kennebunk, ME, USA) that had been previously coated with

Collagen IV (Millipore Sigma). Transwell® inserts were rinsed with DMEM/F12 with Hepes, 10% FBS,

L-glutamine, Penicillin, and streptomycin. Cells were allowed to grow to confluency in 50%

conditioned media (CM) and then changed to differentiation medium (5% CM in primary medium +

ROCK inhibitor) for 48 hours. Differentiated cells were used for toxin treatment.

propagation of Caco-2 cells

Caco-2 intestinal epithelial cells were obtained from ATCC (ATCC HTB-37) and cultured at 37 °C, 5%

CO_2, in Eagle's MEM supplemented with 20% of fetal bovine serum (FBS). To generate polarized

monolayers, ~1 x10^5 cells were seeded onto polystyrene membrane filters (0.4µM, 6.5mm diameter

insert, Transwell, Corning) and grew for at least a week prior to toxin treatment. Media was replaced

with fresh media every two days.

Toxin treatment

Polarized differentiated cells were treated with heat-labile toxin (LT) in differentiation medium

(100 ng/mL) in a volume of 700 µL at the basolateral side of the Transwell insert and 100 µL volume

on the apical aspect of the monolayer. Treated cells were incubated at 37° C for 18 hours and then

fixed for transmission electron microscopy or fluorescence microscopy, or lysed for RNA extraction

(GE Healthcare, Buckinghamshire, UK).

RNA extraction and cDNA synthesis

RNA was extracted using Illustra RNAspin Mini RNA extraction kit (GE Healthcare,

Buckinghamshire, UK). Three biological replicates per treatment (untreated and LT treated) were

submitted to Genome Technology Access Center (GTAC) at Washington University in St. Louis

School of Medicine for RNA-seq library preparation. An Agilent Bioanalyzer was used to determine

the integrity of RNA samples. cDNA was generated using Superscript™ Vilo™ cDNA synthesis kit

(Invitrogen by Thermo Fisher) after normalizing RNA concentrations.

RNA-seq Library Preparation

Library preparation was performed with 10 ng of total RNA, integrity was determined using an

Agilent bioanalyzer. ds-cDNA was prepared using the SMARTer Ultra Low RNA kit for Illumina

Sequencing (Clontech) per manufacturer's protocol. cDNA was fragmented using a Covaris E220

sonicator using peak incident power 18, duty factor 20%, cycles/burst 50, time 120 seconds. cDNA

was blunt ended, had an A base added to the 3'ends, and then had Illumina sequencing adapters

ligated to the ends. Ligated fragments were then amplified for 12-15 cycles using primers

incorporating unique index tags. Fragments were sequenced on an Illumina HiSeq-2500 using

single reads extending 50 bases.

RNA-seq data processing and analysis

RNA-seq reads were aligned to the Ensembl top-level human genome assembly with STAR

version 2.7.3a. Gene counts were derived from the number of uniquely aligned unambiguous

reads by Subread:featureCount version 1.54.1. Read counts were used as input for DESeq2

differential gene expression analysis (version 1.24.0)[77] with default settings, and a minimum *P*-value

significance threshold of 10^{-5} (after False Discovery Rate [FDR[78]] correction for the number of tests).

Fragments per kilobase per million reads mapped (FPKM) values for relative gene expression were

calculated from DESeq2-normalized read counts, and Z-scores were calculated per gene using the

average and standard deviations of FPKM values across samples. Log_2 fold changes were identified

from DESeq2 differential expression output. Pathway enrichment analysis for KEGG[79] and Gene

Ontology (GO)[80] pathways among gene sets of interest was performed using the over-

representation analysis tool provided on the WebGestalt[43] web server (version 2019). Heatmap and

bar graph visualization was performed with Microsoft Excel.

Transcript counts were produced by Sailfish version 0.6.3. Sequencing performance was

assessed for total number of aligned reads, total number of uniquely aligned reads, genes and

transcripts detected, ribosomal fraction known junction saturation and read distribution over known

gene models with RSeQC version 2.3.

All gene-level and transcript counts were then imported into the R/Bioconductor package EdgeR

and TMM normalization size factors were calculated to adjust for samples for differences in library

size. Ribosomal features as well as any feature not expressed in at least the smallest condition size

minus one sample were excluded from further analysis and TMM size factors were recalculated to

created effective TMM size factors. The TMM size factors and the matrix of counts were then

imported into R/Bioconductor package Limma and weighted likelihoods based on the observed

mean-variance relationship of every gene/transcript and sample were then calculated for all samples

with the voomWithQualityWeights function. Performance of the samples was assessed with a

spearman correlation matrix and multi-dimensional scaling plots. Gene/transcript performance was

assessed with plots of residual standard deviation of every gene to their average log-count with a

robustly fitted trend line of the residuals. Generalized linear models were then created to test for

gene/transcript level differential expression. Differentially expressed genes and transcripts were

then filtered for FDR adjusted p-values less than or equal to 0.05.

The biological interpretation of the large set of features found in the Limma results were then

elucidated for global transcriptomic changes in known Gene Ontology (GO) and KEGG terms with

the R/Bioconductor packages GAGE and Pathview. Briefly, GAGE measures for perturbations in GO

or KEGG terms based on changes in observed $\log_2$ fold-changes for the genes within that term

versus the background $\log_2$ fold-changes observed across features not contained in the respective

term as reported by Limma. For GO terms with an adjusted statistical significance of FDR <= 0.05,

heatmaps were automatically generated for each respective term to show how genes co-vary or co-

express across the term in relation to a given biological process or molecular function. In the case of

KEGG curated signaling and metabolism pathways, Pathview was used to generate annotated

pathway maps of any perturbed pathway with an unadjusted statistical significance of p-value <=

0.05. Genes significantly modulated in both Caco-2 cells and enteroids were subjected to further

ontology-free analysis via CompBio v1.4 (PercayAI, Inc., www.percayai.com/compbio)[81] to identify

unifying biological themes in sets of genes differentially expressed between pairwise comparisons of

different groups.

Quantitative real time PCR

Quantitative real time PCR was performed using a QuantStudio 3 real-time detection system

(Applied Biosystems). Fast SYBR Green master mix (Applied Biosystems / Thermo Fisher) was used

for qPCR analysis. Disassociation curve analysis was performed to assess the specificity of

amplification for each sample, and PCR product size was verified by agarose gel electrophoresis.

Percent expression was normalized by GAPDH and analyzed using the comparative threshold cycle

(Ct) method. Amplification of SLC19A3 transcripts from small intestinal enteroids was performed as

recently described[82] with relative gene expression normalized to β actin. HNF4a, HNFg, and SMAD4

gene expression was determined by TaqMan (ThermoFisher) using validated primer and probe sets.

Primers and TaqMan probes used in this study are listed in Supplemental Table 2.

HNF4 transcriptional reporter assays

To assess the impact of LT on HNF4-mediated transcription, pTR104 (GeneCopoeia) carrying 6

copies of the HNF4 transcriptional response element upstream of a secreted *Gaussia* luciferase

(Gluc) reporter gene was transfected (FuGENE, Promega) into confluent Caco-2 cells. Media (EMEM,

ATCC) were exchanged 24 hours after transfection, and after an additional 24 hours baseline

samples of media were removed from each well and stored at -80°C. Wells were then treated with

heat labile toxin (100 ng/ml) overnight (16 hours). At the end of treatment cell culture supernatants

media were transferred to black walled microplates (Greiner 655086) and processed as directed

(Secrete-Pair Luminescence, GeneCopoeia, Inc. Rockville, MD), followed by luminescence detection

(Synergy H1, BioTek). Data are expressed as Relative Light Units (RLU) before and after treatment

with LT.

Thiamin Uptake

Initial rates (30 min; at 37 °C) of carrier-mediated thiamin uptake were examined in LT-treated (100

ng/ml; overnight) and untreated control differentiated small intestinal enteroid monolayers

incubated in Krebs–Ringer buffer (pH 7.4) containing [^{3}H]-thiamin (15 nM). Enteroid monolayers

were then washed with ice-cold Krebs–Ringer buffer followed by lysis with NaOH and neutralization

with 10 N HCl. The radioactive content was counted using a liquid scintillation counter as described

previously[82]. Uptake of thiamin by its respective and distinct carrier-mediated mechanism was

determined by subtracting uptake of [^{3}H]-thiamin in the presence of a high pharmacological

concentration (1 mM) of unlabeled thiamin from uptake in their absence; all uptake data points were

calculated relative to total protein content (in milligrams) of the different preparations and presented

as percentage relative to simultaneously performed controls.

confocal microscopy

Cell monolayers were fixed with 2% paraformaldehyde for 30 min at 37 °C and then washed 3× with

PBS prior to blocking with 1% BSA in PBS for 1 h at room temperature. Villin expression was

detected using 1:100 dilution of anti-villin antibodies raised in mice (Santa Cruz Biotechnology) for 1

hour at room temperature followed by additional 1 hour incubation with 1:200 dilution of

fluorescence-tagged goat anti-mouse IgG Alexa Fluor 488 (Thermofisher) (Supplemental Table 5).

Cell membranes were stained with 1:2000 dilution of CellMask orange (Invitrogen) and nuclei were

counterstained with DAPI (1:1000) and mounted on glass slides using Prolong Gold antifade reagent

(Invitrogen). Images were captured and analyzed on a Nikon C2 confocal microscope equipped with

NIS-Elements AR 5.11.01 software (Nikon).

electron microscopy

For ultrastructural analyses, *in vitro* grown differentiated polarized monolayers of human ileal

enteroid samples as well as mouse intestinal biopsy samples were fixed in 2%

paraformaldehyde/2.5% glutaraldehyde (Ted Pella, Inc., Redding, CA) in 100 mM sodium cacodylate

buffer, pH 7.2 for 2 hours at room temperature and then overnight at 4°C. Samples were washed in

sodium cacodylate buffer and postfixed in 2% osmium tetroxide (Ted Pella, Inc) 1 hour at room

temperature. Samples were then rinsed in dH$_2$0, dehydrated in a graded series of ethanol, and

embedded in Eponate 12 resin (Ted Pella, Inc.). Sections of 95 nm were cut with a Leica Ultracut

UCT ultramicrotome (Leica Microsystems Inc., Bannockburn, IL), stained with uranyl acetate and lead

citrate, and viewed on a JEOL 1200 EX transmission electron microscope (JEOL USA Inc., Peabody,

MA) equipped with an AMT 8 megapixel digital camera and AMT Image Capture Engine V602

software (Advanced Microscopy Techniques, Woburn, MA). Images were analyzed using ImageJ

software for microvilli length and structures.

identification of potential transcription factor target sites

The CREB targets database[66] (http://natural.salk.edu/CREB/) was used to identify potential

categorical CRE full (TGACGTCA) or half (TGACG/CGTCA) sites within the promotor regions of

genes modulated by heat-labile toxin. Following previously described protocol[23], 5 kilobase

upstream sequences from each human gene were retrieved from Ensembl[83] and were searched for

the CREB binding sites TGACGTCA and TGACGTAA (and their reverse complements), based on

sequences retrieved from the TRANSFAC[84] database (matrix ID: V$CREB_01). CREB sequences

matches were identified in the upstream sequences of 1,302 genes. WebGestalt[85] was used for

overall functional enrichment of potential transcription factor binding sequences upstream from

genes that were significantly ($p < 0.05$) up- and down-regulated in both Caco-2 cells and enteroids.

suckling mouse challenge studies

Studies in mice were performed under protocol 20-0438 approved by the Washington University

in Saint Louis Institutional Animal Care and Use Committee. In neonatal mouse challenge studies, 3-

day-old CD-1 mice were inoculated with wild type ETEC (H10407), isogenic toxin deletion mutants,

or sterile PBS (sham negative controls). Inocula were prepared from frozen glycerol stocks

maintained at -80°C grown overnight (~16 hours) in 2 ml Luria Bertani (LB) broth in a shaking

incubator (at 37°C, 225 rpm). Cultures were then diluted 1:100 in 20 ml of fresh LB, grown for

additional 2 hours, and bacteria harvested by centrifugation (6000 rpm, 5 minutes at 4°C). Pellets

were then washed once in ice-cold sterile PBS, then resuspended to a density of ~$1*10^5$ Colony

forming units (CFU) per 20 µl inoculum. CFU in each inoculum were determined by plating serial

dilutions LB-agar plates. Mice were then inoculated with ETEC directly into the stomach through the

abdominal wall using a 31g insulin needle. Following infection, mice were marked with tattoo ink to

define groups and returned to the cage. Seven days post infection mice were sacrificed, and the ileal

tissue collected fixed for microscopy, or preserved for RNA extraction.

Vaccination and immunologic assessment

Adult female mouse dams were vaccinated intranasally three times at 2-week intervals with 10 µg LT

holotoxin in PBS, with the last dose administered approximately 2 weeks prior to anticipated

delivery. Control mice were mock vaccinated with PBS. Fecal pellets collected from vaccinated dams

were used to assess mucosal responses to LT. To collect breast milk the dam was separated from the

litter ~2hours prior to milking. Oxytoxin ([Sigma O3251](#)), 0.002 IU/g, was injected IP to stimulate milk

production, followed by collection via pipette tip.   Anti-LT immune responses in maternal fecal

extracts and breast milk, and stomach contents and sera of neonates were assessed by GM1

ganglioside[86] kinetic ELISA[87] for anti-LT IgA and IgG, using plates coated with GM1 (Sigma G7641),

as previously described[88].

⁶⁶

Figures
figure 1

515
figure 1. Heat-labile toxin modulates expression of multiple genes in intestinal epithelia.
Model at left depicts the *E. coli* heat-labile toxin[89] based on PDB structure <u>1LTS</u> with the A1 subunit
in blue the A2 region of in yellow and pentameric B subunit in green. The E211K mutation of mLT is
in the active site of the A1 subunit. **A.** Scatterplot of RNAseq data right depicts differential
expression profiles of Caco-2 cells following exposure to heat-labile toxin relative to untreated cells
or cells treated with the biologically inactive mLT. **B.** RNAseq data from polarized small intestinal
ileal enteroids treated with LT compared to control untreated cells.
523

524
525
figure 2

figure 2. heat labile toxin modulates multiple genes involved in microvillus assembly.
a. diagram at top (adapted from reference[30]) depicts molecules involved in key elements of
microvillus development. **b.** heat maps of RNAseq data obtained following treatment of Caco-2
intestinal cells (left) with mLT, or LT; and ileal enteroids (right) treated with LT relative to control
untreated cells. RNAseq fold change data comparisons were made with DESeq2[90]. *p≤0.05, **
p≤0.01, *** p≤0.001, **** p≤10^{-4}, *****p≤10^{-5}.
535

536   **figure 3**
537

538
figure 3. heat labile toxin impairs effective formation of small intestinal microvilli.
a. quantitative RT-PCR data for selected microvillus genes comparing untreated (-) and LT-treated
(+) ileal epithelial cells. **b.** Villin production is suppressed in small intestinal enteroids following
treatment (t+72h) with LT . Shown are representative confocal images obtained showing membrane
(CellMask, blue), villin (gold) nuclei (white), and merged image. Graph at right shows villin mean
fluorescence intensity data normalized to the nuclear signal following overnight (~18 hours), and
treatment for 72 hours. ****p<0.0001, **p=0.008, Mann-Whitney 2 tailed testing. **c.** TEM images of
small intestinal microvilli following treatment LT (right) compared to control untreated cells (left).
Graph at right shows length of microvilli when enteroids are treated before (pre) and after (post)
differentiation on polarized ileal cells ****<0.0001 by ANOVA (Kruskal-Wallis, nonparametric
testing).
550

figure 4

figure 4. heat-labile toxin alters the transcription of multiple brush border SLC genes.

a. heat-map indicating key SLC genes modulated by heat labile toxin (LT) compared to enzymatically

inactive E112K LT mutant (mLT), or untreated (ø) Caco-2 cells (left) and human small intestinal (ileal)

enteroids (Hu235D, right). **b.** qRT-PCR confirming LT-mediated modulation of genes in ileal

(Hu235D) enteroids encoding the major thiamine transporter SLC19A3 and the SP1 cis regulatory

element. Data reflect two independent experiments with 2 replicates each. **c.** uptake of [^{3}H]-thiamine

by Hu235 D cells is impaired following LT treatment. Data presented are from two independent

experiments with n=3 replicates each. (* <0.05, **<0.01 Mann-Whitney nonparametric comparisons).

562
563
564

565
566   **figure 5.**

567
figure 5. ETEC disrupts *in vivo* formation of small intestinal microvilli
Timeline at top depicts challenge with ETEC or control nontoxigenic isolate, or sham (PBS)
challenge. **a.** Quantitative PCR results for genes involved in brush border development in small
intestinal samples obtained from infant mice (n=9/group) 7 days after challenge with toxigenic ETEC
(jf876), nontoxigenic ETEC (jf4763, LT⁻/ST⁻) PBS controls. Comparisons between data represent
ANOVA, Kruskal-Wallis testing where ***p≤0.001, **p≤0.01, and *p≤0.05. **b.** Immunofluorescence
images of small intestinal sections showing villin expression (blue), and nuclei (white). **c.** data
represent mean villin fluorescence intensity normalized to the intensity of the nuclear signal (DAPI)

(n=4mice/group) **d.** mean villin fluorescence intensity normalized per enterocyte. **e.** representative
transmission electron microscopy (TEM) images of the small intestinal brush border from mice
challenged with toxin-negative (Δ, left) and toxigenic ETEC (wt, right). **f.** microvillus length
****p≤0.0001 by Mann Whitney (two-tailed) nonparametric comparisons. Data represent geometric
mean length from n=5 mice per group in three independent experiments. g. TEM images from mice
challenged with (clockwise from left) jf570 (*eltA*::Km^R), sham PBS controls, or mice challenged with
wild type ETEC. h. length of microvilli (dashed horizontal lines represent geometric means).
****p<0.0001 by Kruskal-Wallis.
584

figure 6

figure 6. Maternal vaccination with LT mitigates microvillus disruption in neonatal mice.
Timeline depicts vaccination and challenge (top): maternal intranasal (i.n.) vaccinations with 10 μg
LT/immunization (yellow arrows on days 0, 14, 28) and neonatal challenge at 3 days of age (green
arrow) followed by sacrifice and tissue collection at 7 days post infection. **A**. Kinetic ELISA data of
breast milk anti-LT (IgA, and IgG in immunized dams and un-immunized controls). *<0.05 by Mann-
Whitney two-tailed nonparametric testing. **B.** Anti-LT antibodies in the gastric contents of neonatal
mice at day 53. **C.** Representative transmission electron microscopy images of brush border
microvilli from unvaccinated mice challenged with wild type ETEC (left), vaccinated mice challenged
with wild type ETEC, and vaccinated un-challenged controls. **D.** Microvillus lengths (based on image
analysis of n=5 mice group) ****<0.0001 Kruskal-Wallis comparisons.
597
598

599   **figure 7**

600
601
Enterotoxigenic E. coli heat-labile toxin impairs production of HNF4 nuclear receptors.
A. heatmap demonstrating the impact of LT on transcription of paralogous transcription factors
HNF4α and HNF4γ in Caco-2cells (left) and ileal enteroids, (Hu-235D, right) ø = untreated, mLT =
mutant LT. **B.** qRT-PCR (TaqMan) data confirming decreased transcription of HNF4 transcription
factors following treatment of enteroids with LT. **C**. HNF4γ is decreased in nuclear fractions obtained
from small intestinal enteroids following treatment with LT. Shown in the HNF4γ immunoblot are
samples from 4 independent experiments with graph below representing quantitation of signal
intensity normalized to the lamin B1 nuclear protein (p=0.0017, paired t-test, one-tailed). **D.** HNF4
transcription *Gaussia* luciferase reporter assay showing decrease in signal following treatment of
TR104-transfected Caco-2 cells with LT. 2 experimental replicates (n=15 samples total) *** p<0.001
Wilcoxon matched pairs, one-tailed). **E.** Confocal microscopy of representative Ileal sections from
sham-challenged (PBS) left, and ETEC-infected mice (right). Immunofluorescence intensity of HNF4γ
signal in sections from n=5 control mice, and n=6 ETEC-challenged mice. Membranes (yellow) were
strained with CellMask orange (ThermoFisher C10045), nuclei (blue) stained with DAPI, and HNFγ
immunostaining represented in white. Each symbol represents a microscopic region of interest. Bars
represent geographic means (p<0.0001, Mann-Whitney two-tailed comparisons).
618

supplemental figure 1. image from representative CombBio (v1.4) analysis of differential gene expression

in LT-treated enteroids compared to untreated controls, highlighting themes (spheres) related to

biogenesis and function of small intestinal microvilli. Concepts linked within each theme are listed on the

corresponding sphere.

630   supplemental figure 2
631

632
supplemental figure 2. ETEC perturbs the ordered array structure of microvilli. Shown at top are
representative TEM cross sections of infected and control mice (n=5) in each group. **A.** density of
microvilli/μm^2 (n=83 areas in infected and control). p<0.0001. **B.** distance between centers of adjacent
microvilli (n=1001 control, n=2278 infected), p<0.0001 Mann-Whitney. **C.** cross sectional area of
microvilli (n=550 control, n=1559 infected) p<0.0001. **D.** g angle distributions of microvilli from control
and ETEC-infected mice. (analysis by Mann-Whitney, 2-tailed, nonparametric comparisons).
639

643
644   **supplemental figure 3. growth kinetics following challenge of suckling mice with ETEC**
645   **a.** challenge of mice with wild type toxigenic ETEC H10407 (wt), *ΔeltA* mutant (jf571), vs sham
646   challenge (PBS) (n=24/group). data points reflect mean ± sd. **b.** challenge of mice with wild type
647   toxigenic ETEC H10407 (wt) (n=21); *ΔeltA, ΔestH, ΔestP* nontoxigenic mutant (jf4763), vs sham challenge
648   (PBS) (n=20/group).
649

650   supplemental figure 4

651
Supplemental figure 4. LT interrupts HNF4-SMAD4 mediated transcription. **A.** treatment of human-
derived small intestinal enteroids (Hu235D) with LT leads to decreased SMAD4 transcription. Shown are
TaqMan probe results from two independent experiments of polarized epithelial monolayers treated with
LT. n=3 biological replicates x 3 technical replicates/ experiment =18 data points. ****p<0.0001 by Mann
Whitney, two-tailed nonparametric comparisons to untreated cells. **B.** depiction of the HNF4-SMAD4
reciprocal transcription factor activation in which HNF4 activates transcription of SMAD4 and *vice versa* as
(Chen, *et* al[71]). **C.** PKA-mediated phosphorylation of HNF4 paralogs interrupts a DNA-binding module
(Viollet, *et al*[70].). **D.** Summary of proposed mechanism by which LT leads to depletion of transcription
factors essential for enterocyte differentiation. LT-mediated activation of PKA leads to phosphorylation of
HNF4 rendering these transcription factors incapable of binding to *cis* transcriptional response elements
(TRE) in promotors of target genes including SMAD4. Decreased SMAD4 transcription in turn depletes
available HNF4.
664

665
666   **supplemental tables**
667   supplemental table 1

strains and plasmids used in these studies		
bacterial strains		
strain designation	description	source
H10407	wild type ETEC strain, *eltAB, estP, estH*	91
jf571	*eltA*::Km^R derivative of H10407	92
jf876	*lacZYA*::Km^R derivative of H10407	92
jf4763	*eltAB*::Sp^R, estP::Km^R, estH::Cm^R toxin-negative derivative of H10407.	this study
plasmids		
designation	description	source
TR104	6 copies of HNF4 transcriptional response element upstream of Gaussia luciferase reporter (Gaussia-Dura Luc). Km^R, Pm^R	GeneCopoeia
SEAP-PA01	secreted alkaline phosphatase; CMV promotor; Amp^R	GeneCopoeia
Sp^R spectinomycin resistant; Km^R kanamycin resistant; Cm^R chloramphenicol resistant; Pm^R puromycin resistant, Amp^R ampicillin resistant		

668
669

671   supplemental table 2 primers and probes for gene expression

RT-PCR primers			
gene	species	primer sequence (5'-3')	
		forward	reverse
SLC19A3	Human	TTCCTGGATTTACCCCACTG	TATGTCCAAACGGGGAAGA
VIL1	Human	CTGAGCGCCCAAGTCAAAG	AGCAGTCACCATCGAAGAAGC
MYO1A	Human	ACAGGCGAGAGTGGATCAG	GCACTGGGTTAGACTGTAGCAG
MYO7B	Human	TGGACGCCTCAGACGTGAT	CTGGTGACAAATTCCCCTCGG
SP-1	Human	CCATACCCCTTAACCCCG	GAATTTTCACTAATGTTTCCCACC
GAPDH	Human	CATTTCCTGGTATGACAACGA	CTTCCTCTTGTGCTCTTGCT
b-actin	Human	CATCCTGCGTCTGGACCT	TAATGTCACGCACGATTTCC
Ezr	Mouse	CAATCAACGTCCGGGTGAC	GCCAATCGTCTTTACCACCTGA
Vil1	Mouse	TCAAAGGCTCTCTCAACATCAC	AGCAGTCACCATCGAAGAAGC
Espn	Mouse	GGAAATACAGAGGGCGAAAGA	AGAAGCCGAGAGACTCCTATTA
Cdhr2	Mouse	GGCCTCGATTCTACAGCCAAT	GCTCAGAGCCATTCTCAGTCAC
Myo7B	Mouse	AGGGGACCTGAATGAGGCT	GCGGCTGTAGTAGATTTGTACCT
Ebp50	Mouse	AGGGTCCAAATGGCTACGG	CTCCACCAATCGGTCTCCAG
GAPDH	Mouse	AGGTCGGTGTGAACGGATTTG	TGTAGACCATGTAGTTGAGGTCA
TaqMan probes			
gene	species	chromosome	location
HNF4α	Human	20	44355801 - 44432845
HNF4γ	Human	8	75407567 - 75566834
SMAD4	Human	18	51030213 - 51085042

675   supplemental table 3

differential gene expression in gastrointestinal cells following treatment with LT			
response to LT	dataset	genes (n) p<0.05*	genes (n) p<10^{-5}*
higher	Caco-2	5,631	3,832
	ileal enteroids	2138	746
	both[†]	877	268
lower	Caco-2	5,263	3,687
	ileal enteroids	1,839	561
	both[†]	1,013	311
*differentially expressed genes identified by DESeq2[90] [†] overlap in up- and downregulated genes was statistically higher than expected by random selection (p<10^{-10}, binomial distribution testing)			

676
677

678   supplemental table 4.

supplemental table 4. Transcription factor target sequence enrichment among genes differentially regulated in both RNA-seq datasets

		transcription factor target			
transcription factor	target designation	target motif	Total pathway size	Number of significant genes	FDR-adjusted P value
*877 genes upregulated by LT in both RNA-seq datasets**					
AP1 / JUN	TGANTCA_V$AP1_C	TGANTCA	1108	91	4.2E-03
LEF1	CTTTGT_V$LEF1_Q2	CTTTGT	1939	142	4.2E-03
USF	V$USF_C	NCACGTGN	278	32	5.2E-03
*1,013 genes downregulated by LT in both RNA-seq datasets**					
HNF4αγ	V$HNF4_01	NNNRGGNCAAAGKTCANNN	269	37	1.9E-04
HNF4αγ	V$HNF4_01_B	NRGGNCAAAGGTCAN	249	32	2.6E-03
USF	V$USF_02	NNRNCACGTGNYNN	270	32	8.8E-03
SREBP1	TCANNTGAY_V$SREBP1_01	TCANNTGAY	469	47	9.9E-03
NR2F2/COUP	V$COUP_01	TGAMCTTTGMMCYT	260	30	0.015
IRF1	STTTCRNTTT_V$IRF_Q6	STTTCRNTTT	185	23	0.025
USF	V$USF_01	NNRYCACGTGRYNN	254	28	0.038
HNF1	RGTTAMWNATT_V$HNF1_01	RGTTAMWNATT	72	12	0.044

679   *differentially expressed genes identified by DESeq2[90] p<0.05
680

682   supplemental table 5.

antibodies used in these studies		
Specificity	description	source/reference
villin	Villin antibody (BDID2C3), mouse monoclonal IgG1	Santa Cruz sc-66022
mouse IgG	Goat anti-Mouse (H&L) IgG, highly cross-adsorbed antibody conjugated to Alexa Fluor 488	ThermoFisher A-11029
mouse IgA	Pre-absorbed, affinity purified goat anti-mouse IgA conjugated to HRP	Santa Cruz sc-3793
mouse IgG	Affinity purified horse anti-mouse IgG conjugated to HRP	Cell Signaling 7076
HNF4γ	rabbit polyclonal IgG against residues 327-408 of human HNF4γ	ThermoFisher 25801-1-AP
HNF4α	monoclonal IgG2a clone K9218) vs Baculovirus-expressed recombinant human HNF4α (aa 3-49)	ThermoFisher MA1-199

685   supplemental table 6

links to original images	
item	DOI
figure 4b	10.6084/m9.figshare.19251848
figure 4c	10.6084/m9.figshare.19251845
figure 5b	10.6084/m9.figshare.19287353
figure 5e	10.6084/m9.figshare.19251911
figure 5g	10.6084/m9.figshare.19251926
figure 6c	10.6084/m9.figshare.19251938
Figure 7c	10.6084/m9.figshare.19742413

695
696
697
698

699
700

references

Collaborators, G. B. D. D. D. Quantifying risks and interventions that have affected the burden of diarrhoea among children younger than 5 years: an analysis of the Global Burden of Disease Study 2017. *Lancet Infect Dis* **20**, 37-59, doi:10.1016/S1473-3099(19)30401-3 (2020).

Sack, R. B. *et al.* Enterotoxigenic Escherichia coli isolated from patients with severe cholera-like disease. *J Infect Dis* **123**, 378-385 (1971).

Kotloff, K. L. *et al.* Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet* **382**, 209-222, doi:10.1016/S0140-6736(13)60844-2 (2013).

Kotloff, K. L. *et al.* The incidence, aetiology, and adverse clinical consequences of less severe diarrhoeal episodes among infants and children residing in low-income and middle-income countries: a 12-month case-control study as a follow-on to the Global Enteric Multicenter Study (GEMS). *Lancet Glob Health* **7**, e568-e584, doi:10.1016/S2214-109X(19)30076-2 (2019).

Steffen, R., Hill, D. R. & DuPont, H. L. Traveler's diarrhea: a clinical review. *JAMA* **313**, 71-80, doi:10.1001/jama.2014.17006 (2015).

Khalil, I. A. *et al.* Morbidity and mortality due to shigella and enterotoxigenic Escherichia coli diarrhoea: the Global Burden of Disease Study 1990-2016. *Lancet Infect Dis* **18**, 1229-1240, doi:10.1016/S1473-3099(18)30475-4 (2018).

Nasrin, D. *et al.* Pathogens associated with linear growth faltering in children with diarrhea and impact of antibiotic treatment: The Global Enteric Multicenter Study. *J Infect Dis*, doi:10.1093/infdis/jiab434 (2021).

Qadri, F. *et al.* Disease burden due to enterotoxigenic Escherichia coli in the first 2 years of life in an urban community in Bangladesh. *Infect Immun* **75**, 3961-3968, doi:10.1128/IAI.00459-07 (2007).

Black, R. E., Brown, K. H. & Becker, S. Effects of diarrhea associated with specific enteropathogens on the growth of children in rural Bangladesh. *Pediatrics* **73**, 799-805 (1984).

Anderson, J. D. t. *et al.* Burden of enterotoxigenic Escherichia coli and shigella non-fatal diarrhoeal infections in 79 low-income and lower middle-income countries: a modelling analysis. *Lancet Glob Health* **7**, e321-e330, doi:10.1016/S2214-109X(18)30483-2 (2019).

Caulfield, L. E., de Onis, M., Blossner, M. & Black, R. E. Undernutrition as an underlying cause of child deaths associated with diarrhea, pneumonia, malaria, and measles. *Am J Clin Nutr* **80**, 193-198, doi:10.1093/ajcn/80.1.193 (2004).

Checkley, W. *et al.* Multi-country analysis of the effects of diarrhoea on childhood stunting. *Int J Epidemiol* **37**, 816-830, doi:10.1093/ije/dyn099 (2008).

Klipstein, F. A., Short, H. B., Engert, R. F., Jean, L. & Weaver, G. A. Contamination of the small intestine by enterotoxigenic coliform bacteria among the rural population of Haiti. *Gastroenterology* **70**, 1035-1041 (1976).

Klipstein, F. A., Holdeman, L. V., Corcino, J. J. & Moore, W. E. Enterotoxigenic intestinal bacteria in tropical sprue. *Ann Intern Med* **79**, 632-641 (1973).

Klipstein, F. A., Engert, R. F. & Short, H. B. Enterotoxigenicity of colonising coliform bacteria in tropical sprue and blind-loop syndrome. *Lancet* **2**, 342-344 (1978).

Brunser, O., Eidelman, S. & Klipstein, F. A. Intestinal morphology of rural Haitians. A comparison between overt tropical sprue and asymptomatic subjects. *Gastroenterology* **58**, 655-668 (1970).

Mathan, M., Mathan, V. I. & Baker, S. J. An electron-microscopic study of jejunal mucosal morphology in control subjects and in patients with tropical sprue in southern India. *Gastroenterology* **68**, 17-32 (1975).

Fleckenstein, J. M. *et al.* Molecular mechanisms of enterotoxigenic Escherichia coli infection. *Microbes Infect* **12**, 89-98, doi:S1286-4579(09)00235-4 [pii]
10.1016/j.micinf.2009.10.002 (2010).

Tsai, S. C., Noda, M., Adamik, R., Moss, J. & Vaughan, M. Enhancement of choleragen ADP-ribosyltransferase activities by guanyl nucleotides and a 19-kDa membrane protein. *Proc Natl Acad Sci U S A* **84**, 5139-5142. (1987).

Lodish H, B. A., Zipursky SL, Matsudaira P, Baltimore D, Darnell J. in *Molecular Cell Biology*   (W. H. Freeman, 2000).

Sassone-Corsi, P. Transcription factors responsive to cAMP. *Annu Rev Cell Dev Biol* **11**, 355-377, doi:10.1146/annurev.cb.11.110195.002035 (1995).

Sassone-Corsi, P. The cyclic AMP pathway. *Cold Spring Harb Perspect Biol* **4**, doi:10.1101/cshperspect.a011148 (2012).

Zambon, A. C. *et al.* Gene expression patterns define key transcriptional events in cell-cycle regulation by cAMP and protein kinase A. *Proc Natl Acad Sci U S A* **102**, 8561-8566, doi:10.1073/pnas.0503363102 (2005).

Hansson, H. A., Lange, S., Lonnroth, I., Rozell, B. & Tinberg, H. Cytoskeletal disarrangement in rat intestinal epithelium after in vivo exposure to secretagogues. *Acta Pathol Microbiol Immunol Scand [B]* **92**, 93-99 (1984).

Mathan, M. M., Chandy, G. & Mathan, V. I. Ultrastructural changes in the upper small intestinal mucosa in patients with cholera. *Gastroenterology* **109**, 422-430, doi:10.1016/0016-5085(95)90329-1 (1995).

Asakura, H., Tsuchiya, M., Watanabe, Y., Enomoto, Y. & Morita, A. Electron microscopic study on the jejunal mucosa in human cholera. *Gut* **15**, 531-544, doi:10.1136/gut.15.7.531 (1974).

27   Platts-Mills, J. A. *et al.* Association between enteropathogens and malnutrition in children aged 6-23 mo in Bangladesh: a case-
control study. *Am J Clin Nutr* **105**, 1132-1138, doi:10.3945/ajcn.116.138800 (2017).
28   Ilantzis, C., DeMarte, L., Screaton, R. A. & Stanners, C. P. Deregulated expression of the human tumor marker CEA and CEA
family member CEACAM6 disrupts tissue architecture and blocks colonocyte differentiation. *Neoplasia* **4**, 151-163,
doi:10.1038/sj.neo.7900201 (2002).
29   Tadjali, M., Seidelin, J. B., Olsen, J. & Troelsen, J. T. Transcriptome changes during intestinal cell differentiation. *Biochim Biophys*
Acta **1589**, 160-167, doi:10.1016/s0167-4889(02)00170-2 (2002).
30   Crawley, S. W., Mooseker, M. S. & Tyska, M. J. Shaping the intestinal brush border. *J Cell Biol* **207**, 441-451,
doi:10.1083/jcb.201407015 (2014).
31   Postema, M. M., Grega-Larson, N. E., Neininger, A. C. & Tyska, M. J. IRTKS (BAIAP2L1) Elongates Epithelial Microvilli Using EPS8-
Dependent and Independent Mechanisms. *Curr Biol* **28**, 2876-2888 e2874, doi:10.1016/j.cub.2018.07.022 (2018).
32   Tyska, M. J. *et al.* Myosin-1a is critical for normal brush border structure and composition. *Mol Biol Cell* **16**, 2443-2457,
doi:10.1091/mbc.e04-12-1116 (2005).
33   Crawley, S. W., Weck, M. L., Grega-Larson, N. E., Shifrin, D. A., Jr. & Tyska, M. J. ANKS4B Is Essential for Intermicrovillar Adhesion
Complex Formation. *Dev Cell* **36**, 190-200, doi:10.1016/j.devcel.2015.12.022 (2016).
34   He, L., Vasiliou, K. & Nebert, D. W. Analysis and update of the human solute carrier (SLC) gene superfamily. *Hum Genomics* **3**,
195-206, doi:10.1186/1479-7364-3-2-195 (2009).
35   Manary, M. J. *et al.* Perturbed zinc homeostasis in rural 3-5-y-old Malawian children is associated with abnormalities in intestinal
permeability attributed to tropical enteropathy. *Pediatr Res* **67**, 671-675, doi:10.1203/PDR.0b013e3181da44dc (2010).
36   Reidling, J. C., Lambrecht, N., Kassir, M. & Said, H. M. Impaired intestinal vitamin B1 (thiamin) uptake in thiamin transporter-2-
deficient mice. *Gastroenterology* **138**, 1802-1809, doi:10.1053/j.gastro.2009.10.042 (2010).
37   Subramanian, V. S., Marchant, J. S. & Said, H. M. Targeting and trafficking of the human thiamine transporter-2 in epithelial cells.
J Biol Chem **281**, 5233-5245, doi:10.1074/jbc.M512765200 (2006).
38   Said, H. M., Balamurugan, K., Subramanian, V. S. & Marchant, J. S. Expression and functional contribution of hTHTR-2 in thiamin
absorption in human intestine. *American journal of physiology. Gastrointestinal and liver physiology* **286**, G491-498,
doi:10.1152/ajpgi.00361.2003 (2004).
39   Nabokina, S. M., Subramanian, V. S., Valle, J. E. & Said, H. M. Adaptive regulation of human intestinal thiamine uptake by
extracellular substrate level: a role for THTR-2 transcriptional regulation. *American journal of physiology. Gastrointestinal and*
liver physiology **305**, G593-599, doi:10.1152/ajpgi.00237.2013 (2013).
40   Nabokina, S. M. & Said, H. M. Characterization of the 5'-regulatory region of the human thiamin transporter SLC19A3: in vitro
and in vivo studies. *Am J Physiol Gastrointest Liver Physiol* **287**, G822-829, doi:10.1152/ajpgi.00234.2004 (2004).
41   Nabokina, S. M., Reidling, J. C. & Said, H. M. Differentiation-dependent up-regulation of intestinal thiamin uptake: cellular and
molecular mechanisms. *J Biol Chem* **280**, 32676-32682, doi:10.1074/jbc.M505243200 (2005).
42   Blanton, L. V., Barratt, M. J., Charbonneau, M. R., Ahmed, T. & Gordon, J. I. Childhood undernutrition, the gut microbiota, and
microbiota-directed therapeutics. *Science* **352**, 1533, doi:10.1126/science.aad9359 (2016).
43   Wang, J., Vasaikar, S., Shi, Z., Greer, M. & Zhang, B. WebGestalt 2017: a more comprehensive, powerful, flexible and interactive
gene set enrichment analysis toolkit. *Nucleic Acids Res.* **45**, W130-W137, doi:10.1093/nar/gkx356 (2017).
44   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-425,
doi:10.1016/j.cels.2015.12.004 (2015).
45   Nakamura, T., Datta, R., Sherman, M. L. & Kufe, D. Regulation of c-jun gene expression by cAMP in HL-60 myeloid leukemia cells.
J Biol Chem **265**, 22011-22015 (1990).
46   Wang, P. Y. *et al.* c-Jun enhances intestinal epithelial restitution after wounding by increasing phospholipase C-gamma1
transcription. *Am J Physiol Cell Physiol* **312**, C367-C375, doi:10.1152/ajpcell.00330.2016 (2017).
47   San Roman, A. K., Aronson, B. E., Krasinski, S. D., Shivdasani, R. A. & Verzi, M. P. Transcription factors GATA4 and HNF4A control
distinct aspects of intestinal homeostasis in conjunction with transcription factor CDX2. *J Biol Chem* **290**, 1850-1860,
doi:10.1074/jbc.M114.620211 (2015).
48   Chen, L. *et al.* The nuclear receptor HNF4 drives a brush border gene program conserved across murine intestine, kidney, and
embryonic yolk sac. *Nature communications* **12**, 2886, doi:10.1038/s41467-021-22761-5 (2021).
49   Dankel, S. N., Hoang, T., Flageng, M. H., Sagen, J. V. & Mellgren, G. cAMP-mediated regulation of HNF-4alpha depends on the
level of coactivator PGC-1alpha. *Biochim Biophys Acta* **1803**, 1013-1019, doi:10.1016/j.bbamcr.2010.05.008 (2010).
50   Khalil, I. *et al.* Enterotoxigenic Escherichia coli (ETEC) vaccines: Priority activities to enable product development, licensure, and
global access. *Vaccine* **39**, 4266-4277, doi:10.1016/j.vaccine.2021.04.018 (2021).
51   Sheikh, A. *et al.* CEACAMs serve as toxin-stimulated receptors for enterotoxigenic Escherichia coli. *Proc Natl Acad Sci U S A* **117**,
29055-29062, doi:10.1073/pnas.2012480117 (2020).
52   Sheikh, A. *et al.* Enterotoxigenic Escherichia coli degrades the host MUC2 mucin barrier to facilitate critical pathogen-enterocyte
interactions in human small intestine. *Infect Immun*, IAI0057221, doi:10.1128/IAI.00572-21 (2021).
53   Harris, J. B. Cholera: Immunity and Prospects in Vaccine Development. *J Infect Dis* **218**, S141-S146, doi:10.1093/infdis/jiy414
(2018).

54   Pipaliya, N. *et al*. Spectrum of chronic small bowel diarrhea with malabsorption in Indian subcontinent: is the trend really
changing? *Intest Res* **14**, 75-82, doi:10.5217/ir.2016.14.1.75 (2016).
55   Ghoshal, U. C., Srivastava, D., Verma, A. & Ghoshal, U. Tropical sprue in 2014: the new face of an old disease. *Curr Gastroenterol*
Rep **16**, 391, doi:10.1007/s11894-014-0391-3 (2014).
56   Dutta, A. K., Balekuduru, A. & Chacko, A. Spectrum of malabsorption in India--tropical sprue is still the leader. *J Assoc Physicians*
India **59**, 420-422 (2011).
57   Klipstein, F. A. & Falaiye, J. M. Tropical sprue in expatriates from the tropics living in the continental United States. *Medicine*
(Baltimore) **48**, 475-491 (1969).
58   Klipstein, F. A. Tropical sprue in travelers and expatriates living abroad. *Gastroenterology* **80**, 590-600 (1981).
59   Kelly, P. *et al*. Gene expression profiles compared in environmental and malnutrition enteropathy in Zambian children and
adults. *EBioMedicine* **70**, 103509, doi:10.1016/j.ebiom.2021.103509 (2021).
60   Zyambo, K. *et al*. Reduced Fractional Absorption of Zinc in Children With Environmental Enteropathy in Zambia. *J Pediatr*
Gastroenterol Nutr **74**, 277-283, doi:10.1097/MPG.0000000000003292 (2022).
61   Long, J. M. *et al*. Zinc Absorption from Micronutrient Powders Is Low in Bangladeshi Toddlers at Risk of Environmental Enteric
Dysfunction and May Increase Dietary Zinc Requirements. *J Nutr* **149**, 98-105, doi:10.1093/jn/nxy245 (2019).
62   Lindenmayer, G. W., Stoltzfus, R. J. & Prendergast, A. J. Interactions between zinc deficiency and environmental enteropathy in
developing countries. *Adv Nutr* **5**, 1-6, doi:10.3945/an.113.004838 (2014).
63   Roy, S. K. *et al*. Impact of zinc supplementation on intestinal permeability in Bangladeshi children with acute diarrhoea and
persistent diarrhoea syndrome. *J Pediatr Gastroenterol Nutr* **15**, 289-296, doi:10.1097/00005176-199210000-00010 (1992).
64   Dhingra, U. *et al*. Lower-Dose Zinc for Childhood Diarrhea - A Randomized, Multicenter Trial. *N Engl J Med* **383**, 1231-1241,
doi:10.1056/NEJMoa1915905 (2020).
65   Kuhlmann, F. M. *et al*. Conservation and global distribution of non-canonical antigens in Enterotoxigenic Escherichia coli. *PLoS*
Negl Trop Dis **13**, e0007825, doi:10.1371/journal.pntd.0007825 (2019).
66   Zhang, X. *et al*. Genome-wide analysis of cAMP-response element binding protein occupancy, phosphorylation, and target gene
activation in human tissues. *Proc Natl Acad Sci U S A* **102**, 4459-4464, doi:10.1073/pnas.0501076102 (2005).
67   Mayr, B. & Montminy, M. Transcriptional regulation by the phosphorylation-dependent factor CREB. *Nat Rev Mol Cell Biol* **2**, 599-
609 (2001).
68   Delghandi, M. P., Johannessen, M. & Moens, U. The cAMP signalling pathway activates CREB through PKA, p38 and MSK1 in NIH
3T3 cells. *Cell Signal* **17**, 1343-1351, doi:10.1016/j.cellsig.2005.02.003 (2005).
69   Rohlff, C., Ahmad, S., Borellini, F., Lei, J. & Glazer, R. I. Modulation of transcription factor Sp1 by cAMP-dependent protein
kinase. *J Biol Chem* **272**, 21137-21141, doi:10.1074/jbc.272.34.21137 (1997).
70   Viollet, B., Kahn, A. & Raymondjean, M. Protein kinase A-dependent phosphorylation modulates DNA-binding activity of
hepatocyte nuclear factor 4. *Molecular and cellular biology* **17**, 4208-4219, doi:10.1128/MCB.17.8.4208 (1997).
71   Chen, L. *et al*. A reinforcing HNF4-SMAD4 feed-forward module stabilizes enterocyte identity. *Nat Genet* **51**, 777-785,
doi:10.1038/s41588-019-0384-0 (2019).
72   de Rooij, J. *et al*. Epac is a Rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP. *Nature* **396**, 474-477,
doi:10.1038/24884 (1998).
73   Bos, J. L., de Rooij, J. & Reedquist, K. A. Rap1 signalling: adhering to new models. *Nat Rev Mol Cell Biol* **2**, 369-377,
doi:10.1038/35073073 (2001).
74   Frische, E. W. & Zwartkruis, F. J. Rap1, a mercenary among the Ras-like GTPases. *Dev Biol* **340**, 1-9,
doi:10.1016/j.ydbio.2009.12.043 (2010).
75   Ogryzko, V. V., Schiltz, R. L., Russanova, V., Howard, B. H. & Nakatani, Y. The transcriptional coactivators p300 and CBP are
histone acetyltransferases. *Cell* **87**, 953-959, doi:10.1016/s0092-8674(00)82001-2 (1996).
76   Pattabiraman, D. R. *et al*. Activation of PKA leads to mesenchymal-to-epithelial transition and loss of tumor-initiating ability.
Science **351**, aad3680, doi:10.1126/science.aad3680 (2016).
77   Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol* **11**, R106, doi:10.1186/gb-2010-
11-10-r106 (2010).
78   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing.
Journal of the Royal Statistical Society. Series B (Methodological) **57**, 289-300, doi:10.2307/2346101 (1995).
79   Kanehisa, M., Furumichi, M., Sato, Y., Ishiguro-Watanabe, M. & Tanabe, M. KEGG: integrating viruses and cellular organisms.
Nucleic Acids Res **49**, D545-D551, doi:10.1093/nar/gkaa970 (2021).
80   Gene Ontology, C. The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res* **49**, D325-D334,
doi:10.1093/nar/gkaa1113 (2021).
81   Delannoy-Bruno, O. *et al*. Evaluating microbiome-directed fibre snacks in gnotobiotic mice and humans. *Nature* **595**, 91-95,
doi:10.1038/s41586-021-03671-4 (2021).
82   Sabui, S. *et al*. Hypoxia inhibits colonic uptake of the microbiota-generated forms of vitamin B1 via HIF-1alpha-mediated
transcriptional regulation of their transporters. *J Biol Chem* **298**, 101562, doi:10.1016/j.jbc.2022.101562 (2022).
83   Howe, K. L. *et al*. Ensembl 2021. *Nucleic Acids Res* **49**, D884-D891, doi:10.1093/nar/gkaa942 (2021).

84   Matys, V. *et al*. TRANSFAC and its module TRANSCompel: transcriptional gene regulation in eukaryotes. *Nucleic Acids Res* **34**,
D108-110, doi:10.1093/nar/gkj143 (2006).
85   Wang, J., Vasaikar, S., Shi, Z., Greer, M. & Zhang, B. WebGestalt 2017: a more comprehensive, powerful, flexible and interactive
gene set enrichment analysis toolkit. *Nucleic Acids Res* **45**, W130-W137, doi:10.1093/nar/gkx356 (2017).
86   Sack, D. A., Huda, S., Neogi, P. K., Daniel, R. R. & Spira, W. M. Microtiter ganglioside enzyme-linked immunosorbent assay for
vibrio and Escherichia coli heat-labile enterotoxins and antitoxin. *J Clin Microbiol* **11**, 35-40 (1980).
87   Tsang, V. C., Wilson, B. C. & Maddison, S. E. Kinetic studies of a quantitative single-tube enzyme-linked immunosorbent assay.
Clinical chemistry **26**, 1255-1260 (1980).
88   Roy, K., Hamilton, D. J., Munson, G. P. & Fleckenstein, J. M. Outer Membrane Vesicles Induce Immune Responses to Virulence
Proteins and Protect against Colonization by Enterotoxigenic Escherichia coli. *Clin Vaccine Immunol* **18**, 1803-1808,
doi:10.1128/CVI.05217-11 (2011).
89   Sixma, T. K. *et al*. Refined structure of Escherichia coli heat-labile enterotoxin, a close relative of cholera toxin. *J Mol Biol* **230**,
890-918, doi:10.1006/jmbi.1993.1209 (1993).
90   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome
Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).
91   Evans, D. J., Jr. & Evans, D. G. Three characteristics associated with enterotoxigenic Escherichia coli isolated from man. *Infect
Immun* **8**, 322-328 (1973).
92   Dorsey, F. C., Fischer, J. F. & Fleckenstein, J. M. Directed delivery of heat-labile enterotoxin by enterotoxigenic Escherichia coli.
Cell Microbiol **8**, 1516-1527, doi:10.1111/j.1462-5822.2006.00736.x (2006).
892

acknowledgements

893
JMF was supported by funding from the National Institute of Allergy and Infectious Diseases (NIAID)

of the National Institutes of Health (NIH) R01 AI126887, R01 AI089894, U01 AI095473; and the

Department of Veterans Affairs (5I01BX001469-05). Research conducted by AS was also supported

by National Institute of Allergy and Infectious Diseases of the National Institutes of Health under

Award Number T32AI007172. We thank the Genome Technology Access Center in the Department

of Genetics at Washington University School of Medicine for help with genomic analysis. The Center

is partially supported by NCI Cancer Center Support Grant #P30 CA91842 to the Siteman Cancer

Center and by ICTS/CTSA Grant# UL1TR002345 from the National Center for Research Resources

(NCRR), a component of the National Institutes of Health (NIH), and NIH Roadmap for Medical

Research. HS was supported by funding from the Department of Veterans Affairs (I01BX001142) and

the National Institutes of Health (DK5606 and AA018071). The content is solely the responsibility of

the authors and does not necessarily represent the official views of the National Institutes of Health,

or the Department of Veterans Affairs.

interests declaration

R.D.H. and C.S. may receive royalty income based on the CompBio technology developed by R.D.H. and licensed by Washington University
to PercayAI.
910