Distinct Nrf2 Signaling Thresholds Mediate Lung Tumor Initiation and Progression

Janine M. DeBlasi[1,2], Aimee Falzone[1], Samantha Caldwell[1], Yun Pyo Kang[1], Nicolas Prieto-Farigua[1], Justin R. Prigge[3], Edward E. Schmidt[3], Iok In Christine Chio[4,5], Florian A. Karreth[6], Gina M. DeNicola[1*]

[1]Department of Cancer Physiology, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, 33612, USA.
[2]Cancer Biology PhD Program, University of South Florida, Tampa, FL 33612, USA.
[3]Microbiology & Cell Biology Department, Montana State University, Bozeman, MT 59717, USA.
[4]Institute for Cancer Genetics, Department of Genetics and Development, Columbia University Irving Medical Center, New York, NY 10032, USA.
[5]Herbert Irving Comprehensive Cancer Center, Columbia University Irving Medical Center, New York, NY 10032, USA.
[6]Department of Molecular Oncology, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, 33612, USA.

*Corresponding author. E-mail: gina.denicola@moffitt.org

Keywords: Nuclear factor erythroid 2-related factor 2 (**NRF2**), Kelch-like ECH-associated protein 1 (**KEAP1**), Non-small cell lung cancer (**NSCLC**), Genetically engineered mouse model (**GEMM**)

Abstract

Mutations in the KEAP1-NRF2 pathway occur in up to a third of non-small cell lung cancer (NSCLC) cases and often confer resistance to therapy and poor outcomes. Here, we developed murine alleles of the KEAP1 and NRF2 mutations found in human NSCLC and comprehensively interrogated their impact on tumor initiation and progression. Chronic Nrf2 stabilization by Keap1 or Nrf2 mutation was not sufficient to induce tumorigenesis, even in the absence of tumor suppressors p53 or Lkb1. When combined with Kras$^{G12D/+}$, constitutive Nrf2 activation promoted lung tumor initiation and early progression of hyperplasia to low-grade tumors but impaired their progression to advanced-grade tumors, which was reversed by Nrf2 deletion. Finally, NRF2 overexpression in KEAP1 mutant NSCLC cell lines was detrimental to cell proliferation, viability, and anchorage-independent colony formation. Collectively, our results

establish the context-dependence and activity threshold for NRF2 during the lung tumorigenic process.

Main

NRF2 (nuclear factor-erythroid 2 p45-related factor 2) is a stress-responsive transcription factor that regulates the detoxification of reactive oxygen species (ROS), maintains cellular homeostasis, and regulates many facets of metabolism[1,2]. NRF2 is negatively regulated by KEAP1 (Kelch-like ECH-associated protein 1), a substrate adaptor protein for the cullin 3 (CUL3)-based E3 ubiquitin ligase that facilitates NRF2 ubiquitination and proteasomal degradation in the absence of oxidative or xenobiotic stress[3]. NRF2 promotes the detoxification of carcinogens to limit deleterious mutations that initiate cancer[4-7] and NRF2 activators are being explored as chemopreventative agents[8-10]. However, NRF2 is frequently stabilized in many cancers, particularly non-small cell lung cancer (NSCLC), where mutations in the KEAP1-NRF2 pathway are found in up to 30% of cases[11,12]. NRF2 stabilization is associated with poor prognosis[13-18], resistance to chemo- and radiotherapy[19,20], cancer cell survival[20], proliferation[21], metabolic reprogramming[22-24], and metastasis[25]. It remains unclear, however, whether chronic NRF2 stabilization transforms normal cells. Thus, it is important to understand the contexts and mechanisms by which NRF2 can prevent and promote cancer phenotypes.

Preclinical genetically engineered mouse models (GEMMs) have advanced our understanding of the role of NRF2 in lung tumorigenesis[21,25-35]. NRF2 activation in NSCLC has been modeled in GEMMs by inactivating Keap1 via conditional knockout[26-29,31,33] or CRISPR-mediated deletion[25,32,34,35], in contrast to the KEAP1 mutations found in human lung cancer. These studies have shown that Nrf2 promotes lung tumor initiation[27,28], tumor size[32,36], progression[31,32], and metastasis[25]. However, other studies have failed to see an effect of Nrf2 on lung tumor initiation[36,37] or size[34,35,37], and we reported that Nrf2 activation significantly decreases tumor size[30]. Study conditions, time

points and phenotypes assayed varied across these studies. Therefore, the role NRF2 activation plays at various stages of tumor initiation and progression remains unclear.

In the current study, we generated lung cancer GEMMs expressing Keap1[R554Q] and Nrf2[D29H] mutations to comprehensively investigate how Nrf2 activation affects each stage of the tumorigenic process. We found that constitutive Nrf2 stabilization induced by these mutations was insufficient for lung tumor development, even in the context of tumor suppressor loss. In contrast, these mutations promoted lung tumor initiation in the Kras[G12D/+] model of early lung adenocarcinoma, consistent with previous studies[21,27,28]. Using the Kras[G12D/+]; p53[fl/fl] adenocarcinoma model, we found that homozygous Keap1 mutation unexpectedly blocked tumor progression. Moreover, Nrf2 expression and activity was downregulated in advanced tumors, and Nrf2 deletion could rescue the Keap1 mutation-mediated progression impairment. Overall, our data suggest that NRF2 has distinct, threshold-dependent effects during lung tumor initiation and progression.

Results
Keap1[R554Q] and *Nfe2l2[D29H]* alleles activate the Nrf2 transcriptional program
To study the role of NRF2 activation in lung cancer, we developed alleles harboring either the Keap1[R554Q] or the Nrf2[D29H] mutation found in human NSCLC and compared the effects of these alleles to their wild-type (WT) counterparts **(Fig. 1a-d)**. Both Keap1[R554Q] and Nrf2[D29H] mutations prevent Keap1-mediated ubiquitination of Nrf2, allowing for constitutive expression of Nrf2 and transcription of Nrf2 target genes[38,39]. To generate the conditionally active *(CA)-Keap1[R554Q]* allele, we inserted a wild-type *Keap1* cDNA containing exons 3-5 flanked by loxP sites upstream of the R554Q mutation in endogenous exon 4 of the *Keap1* gene **(Fig. 1a)**[30]. For the Lox-STOP-Lox *(LSL)-Nfe2l2[D29H]* allele, we inserted a loxP-flanked transcriptional and translational STOP (LSL) cassette upstream of the D29H mutation in exon 2 of the endogenous *Nfe2l2* gene **(Fig. 1c).** For both alleles, Cre-mediated excision of loxP-flanked cassettes allows for physiological expression of Keap1[R554Q] or Nrf2[D29H], recapitulating NRF2 activation in human NSCLC. To validate the functionality of these alleles, we first generated mouse embryonic fibroblasts (MEFs), which allowed the switching from a Nrf2 deficient state

(Nrf2$^{LSL/LSL}$) to a Nrf2 stabilized state (Nrf2$^{D29H/D29H}$), or from a basal Nrf2 state (Keap1$^{+/+}$) to a stabilized Nrf2 state (Keap1$^{R554Q/R554Q}$) in an isogenic system. Using these MEFs, we performed both transcriptomic and metabolomic profiling **(Fig. 1e-h)**. RNA-sequencing indicated that both Keap1$^{R554Q/R554Q}$ and Nrf2$^{D29H/D29H}$ MEFs demonstrated increased transcription of canonical Nrf2 target genes, including Nqo1, Srxn1, Txnrd1, and Gclc **(Fig. 1e,f)**. Non-targeted metabolomics revealed a robust increase in sulfur-containing metabolites, including glutathione, N-acetyl-L-cysteine, hypotaurine, and L-cystathionine in the Nrf2$^{D29H/D29H}$ MEFs, consistent with our previous findings in the Keap1$^{R554Q/R554Q}$ MEFs **(Fig. 1g,h)**[30]. Targeting the murine *Keap1* locus to generate a *Keap1^{flox}* allele resulted in the generation of a hypomorphic allele prior to Cre-mediated recombination, leading to decreased Keap1 levels and increased Nrf2 transcriptional activity throughout the whole animal[40]. Importantly, we found no differences in expression of Keap1, Nrf2, or Nrf2 target proteins Nqo1 and Gclc between CA-Keap1^{R554Q} and WT Keap1 MEFs, indicating that the *CA-Keap1^{R554Q}* allele is not hypomorphic (**Extended Data Fig. 1**). Collectively, these results indicate that the mutant *Keap1^{R554Q}* and *Nfe2l2^{D29H}* alleles activate the Nrf2 transcriptional program and similarly rewire cellular metabolism.

Keap1 or Nrf2 mutation is not sufficient to initiate lung tumorigenesis

Given the importance of NRF2 in cytoprotection and redox homeostasis, there has been a longstanding interest in activating NRF2 pharmacologically for chemoprevention[8-10]. The long-term safety of this approach, and whether the chronic activation of NRF2 can transform healthy cells *in vivo*, remains unknown. Moreover, the whole body deletion of Keap1 in mice results in postnatal lethality because of constitutive Nrf2 activation[41]. To test whether constitutive Nrf2 activation is sufficient to initiate lung tumor formation, we induced the recombination of *Keap1* and *Nfe2l2* alleles in the lungs of Keap1$^{R554Q/+}$, Keap1$^{R554Q/R554Q}$, or Nrf2$^{D29H/+}$ mice using adenoviral-Cre **(Fig. 2)**. Importantly, Keap1$^{R554Q/R554Q}$ is most reflective of human disease, as biallelic inactivation of *KEAP1* is common in lung cancer[13]. In contrast, *NFE2L2* mutations in lung cancer patient tumors are frequently heterozygous[42]. First, we analyzed the overall survival between the different groups. The median survival across genotypes ranged from approximately

650-750 days, with no significant survival differences observed between wild-type and Keap1/Nrf2 mutant groups **(Fig. 2a)**. While mice did develop tumors, they comprised age-associated tumors like lymphoma. Upon examination of mouse lung histology for the presence of lung tumors, lung tumor-free survival was also not different between the groups **(Fig. 2b)**. Finally, histological analysis of lung tissues revealed that both alveolar and bronchiolar cells appeared phenotypically normal across the genotypes **(Fig. 2c)**. These results indicate that constitutive Nrf2 activation is not sufficient to induce lung tumor formation.

Keap1 or Nrf2 mutation is not sufficient to initiate lung tumorigenesis in combination with tumor suppressor loss

To determine whether tumor suppressor loss was required for mutant Keap1 or Nrf2 to initiate lung tumor formation, we crossed Keap1 and Nrf2 mutant mice with p53^{flox} and Lkb1^{flox} alleles to concomitantly activate Nrf2 and delete these tumor suppressors in the lung **(Fig. 3)**. We first examined the consequence of Nrf2$^{D29H/+}$, Keap1$^{R554Q/+}$, or Keap1$^{R554Q/R554Q}$ in combination with p53 deletion. Mice were aged to 500 days, at which time all mice were euthanized and examined for evidence of lung tumor formation. While a small number of these mice did succumb to disease prior to 500 days, they developed age-associated tumors including lymphoma, and we did not observe any differences in overall or lung tumor-free survival between genotypes **(Fig. 3a,b)**. However, when examining the lung tissue histology, we observed dysplasia in Keap1$^{R554Q/R554Q}$ bronchioles following loss of p53 **(Fig. 3c)**. This observation is consistent with previous work showing that tracheospheres derived from *Keap1$^{-/-}$;Trp53$^{-/-}$* cells had an aberrant morphology[29]. We next examined the consequence of Nrf2$^{D29H/+}$, Keap1$^{R554Q/+}$ or Keap1$^{R554Q/R554Q}$ in combination with Lkb1 deletion. Similar to what was observed with p53, we also did not find any differences in overall or lung tumor-free survival between cohorts **(Fig. 3d,e)**. Moreover, the bronchiolar and alveolar morphology was normal across genotypes, in contrast to what was observed upon p53 loss **(Fig. 3f)**. Our findings indicate that Keap1/Nrf2 mutation is not sufficient to initiate lung tumor formation in combination with tumor suppressor loss.

Nrf2 activation cooperates with mutant Kras to promote lung tumor initiation and early progression

We and others have reported that Nrf2 activation is important for Kras mutant lung tumorigenesis[21,27,28]. To understand how Keap1 or Nrf2 mutation affect lung tumor initiation and early progression, we crossed Keap1/Nrf2 mutant mice with the Kras$^{G12D/+}$ model of early lung adenocarcinoma[43]. While all mice succumbed to lung tumors with a median survival of around 200 days, we observed no difference in survival between cohorts **(Fig. 4a,b)**. We next validated that these mutations were activating toward Nrf2 in tumors by performing immunohistochemical staining for Nrf2 and the Nrf2 target Nqo1. We observed that Keap1$^{R554Q/R554Q}$ expression resulted in the greatest degree of Nrf2 activation, followed by Nrf2$^{D29H/+}$, and then Keap1$^{R554Q/+}$ compared to Keap1/Nrf2$^{+/+}$ expression **(Fig. 4c-e)**. To examine the influence of Nrf2 activation on tumor initiation, we quantified tumor number across the genotypes and found that Keap1$^{R554Q/R554Q}$ and Nrf2$^{D29H/+}$ significantly increased tumor number in the Kras$^{G12D/+}$ model **(Fig. 4f,g)**, consistent with prior reports using Keap1 deletion models[27,28]. We then examined the influence of Nrf2 on tumor progression by analyzing tumor grade. The distribution of atypical adenomatous and bronchiolar hyperplasia (AAH and BH, respectively) and tumors from grades 1 (adenoma) to 5 (adenocarcinoma) was determined. We observed an increased proportion of grade 1 tumors in Keap1$^{R554Q/R554Q}$ and Nrf2$^{D29H/+}$ mice compared to Keap1$^{R554Q/+}$ and Keap1/Nrf2$^{+/+}$ mice **(Fig. 4h)**. Surprisingly, we found that there was a decrease in grade 3 tumor burden across all Keap1 and Nrf2 mutant models **(Fig. 4i)**, although these tumors were rare. Our findings indicate that Kras$^{G12D/+}$ mutation cooperates with Keap1/Nrf2 mutation to promote formation of lung tumors and early progression to low-grade tumors.

Nrf2 activation impairs lung adenocarcinoma progression

The decrease in grade 3 tumor burden in the Keap1/Nrf2 mutant models suggested that Nrf2 activation may impair progression to higher grade tumors. We next used the Kras$^{G12D/+}$; p53$^{fl/fl}$ (KP) model, which develops advanced-grade lung adenocarcinomas[44]. We previously reported that the expression of Keap1$^{R554Q/R554Q}$ dramatically decreased overall tumor size in this model[30], but other groups have found differing effects of Nrf2

activation. While some studies reported that Keap1 inactivation promoted adenocarcinoma progression[31,32], others reported that Keap1 deletion did not affect tumor size[34,35], but the conditions used and phenotypes assayed varied across these studies. Thus, we decided to perform comprehensive phenotyping on KP tumors following Nrf2 activation. Similar to what we observed in the Kras$^{G12D/+}$ model, we found that overall survival of the KP model was not affected by Keap1 or Nrf2 mutation **(Fig. 5a,b)**. Moreover, we found that Keap1/Nrf2 mutation affected Nrf2 activation in a similar manner to the Kras$^{G12D/+}$ model, with Keap1$^{R554Q/R554Q}$ being the most activating, followed by Nrf2$^{D29H/+}$, and then Keap1$^{R554Q/+}$ compared to Keap1/Nrf2$^{+/+}$ mice **(Fig. 5c-e)**. Next, we examined tumor progression by tumor grading and found a significant decrease in the proportion of grade 3 and 4 tumors in the Keap1$^{R554Q/R554Q}$ cohort **(Fig. 5f,g)**. We also observed an increase in the proportion of grade 1 tumors with Keap1$^{R554Q/R554Q}$, suggesting that there may be a threshold for Nrf2 activation to promote early progression, but impair late progression **(Fig. 5g)**. Moreover, we found similar trends by analyzing tumor burden by grade, as seen with an increase in grade 1 tumor burden and a decrease in grade 3 tumor burden with Keap1$^{R554Q/R554Q}$ **(Fig. 5h)**. Although not statistically significant, the Nrf2$^{D29H/+}$ cohort also had a modest decrease in grade 3 and 4 tumors, further supporting a threshold for Nrf2 to impair tumor progression **(Fig. 5f-h)**. We next investigated whether tumors that progressed to the adenocarcinoma stage altered Nrf2 expression and/or activity. To this end, we analyzed Nrf2 and Nqo1 levels across all tumor grades and hyperplasia (AAH, BH). We found that Nrf2 and Nqo1 levels were highly elevated in Keap1/Nrf2 mutant grade 1 tumors, with Nqo1 demonstrating increased nuclear localization in homozygous Keap1 mutant tumors compared to Nrf2 mutant tumors **(Fig. 6a)**. However, as tumors progressed to higher grades, Nrf2 and Nqo1 expression were reduced in the Keap1/Nrf2 mutant models **(Fig. 6a-c)**. These results suggest that Nrf2 activation beyond a certain threshold impairs advanced-grade tumor progression, requiring selection for a more tolerable level of Nrf2 expression and activity in high-grade tumors.

NRF2 overexpression impairs NSCLC cell proliferation, viability, and anchorage-independent colony formation

KEAP1 has other substrates[45-50], raising the question of whether NRF2 plays a causal role in tumor suppression. It was previously reported that KEAP1 mutant lung cancer cell lines are "NRF2 addicted" and dependent on NRF2 for proliferation[51]. Supportingly, analysis of DepMap data[52] revealed that NSCLC cell lines with high NRF2 activity[22], which were enriched for KEAP1 mutations, exhibited NRF2 dependence **(Fig. 7a)**. To directly test the hypothesis that excessive NRF2 activation above a specific threshold is detrimental to lung tumor cell growth, we used lentiviral transduction to overexpress NRF2 in five KEAP1 mutant lung cancer cell lines (H1944, H322, A549, HCC15, H460). We confirmed overexpression of NRF2 by western blot analysis of NRF2 and target genes GCLC, xCT, and GSR **(Fig. 7b)**. Lentiviral transduction increased the expression of NRF2 in all cell lines, and also increased the expression of NRF2 targets, demonstrating that NRF2 binding sites were not saturated by the level of NRF2 in these cell lines **(Fig. 7b)**. Next, we determined the influence of NRF2 overexpression on cellular proliferation and death over the course of four days using live cell imaging **(Fig. 7c-g)**. We found that NRF2 overexpression decreased cell proliferation in all cell lines, and increased cell death in 4 out of 5 cell lines **(Fig. 7f,g)**. Finally, we observed impaired anchorage-independent growth in soft agar in all cell lines **(Fig. 7h,i)**. These results indicate that there is an optimal threshold of NRF2 activity, and that excess NRF2 activation can impair lung cancer phenotypes.

Single copy Nrf2 deletion rescues homozygous Keap1^{R554Q}-mediated tumor progression impairment

To directly examine whether reducing Nrf2 levels could alleviate the block in adenocarcinoma progression in the Keap1$^{R554Q/R554Q}$ model, we crossed a Nrf2^{flox} allele into both our Kras$^{G12D/+}$; p53$^{fl/fl}$ and Kras$^{G12D/+}$; p53$^{fl/fl}$; Keap1$^{R554Q/R554Q}$ models. Because we previously found that complete Nrf2 deficiency impairs tumor initiation[21], we examined the consequence of single copy Nrf2 deletion on tumor phenotypes. Again, we found no difference in overall survival between groups **(Fig. 8a)**. However, histological examination of the lungs revealed a striking difference in tumor number and size, with Nrf2 heterozygous deletion having a minimal effect on the Kras$^{G12D/+}$; p53$^{fl/fl}$ model while dramatically increasing tumor burden in the Kras$^{G12D/+}$; p53$^{fl/fl}$;

Keap1$^{R554Q/R554Q}$ model **(Fig. 8b).** Next, we examined tumor progression in these models. In agreement with our previous experiment **(Fig. 5g)**, we found that the Keap1$^{R554Q/R554Q}$ cohort had a significant reduction in grade 3 and grade 4 tumors upon Nrf2^{WT} expression **(Fig. 8c)**. However, upon single copy Nrf2 deletion (Nrf2$^{flox/+}$), Keap1$^{R554Q/R554Q}$ failed to suppress tumor progression **(Fig. 8c)**. Similar findings were observed when analyzing tumor burden by grade, with a decrease in grade 3 and grade 4 tumor burden induced by Keap1$^{R554Q/R554Q}$ that was alleviated by single copy deletion of Nrf2 **(Fig. 8d)**. These results demonstrate that there is a threshold by which Nrf2 activation can promote or impair tumor initiation or progression.

Discussion

Using genetically engineered mouse models of Keap1/Nrf2 mutation, we find that Keap1 or Nrf2 mutations alone are insufficient to cause lung tumor formation. Even in combination with tumor suppressor loss, we did not observe lung tumor formation after 500 days, suggesting that chronic NRF2 activation would be a safe strategy for chemoprevention. These findings corroborate previous studies showing that Keap1 deletion does not induce lung tumor development[26], even in the absence of tumor suppressors p53 or Lkb1[27], out to 12-15 months. In combination with Kras$^{G12D/+}$ mutation, we found that Nrf2 activation promoted tumor initiation, regardless of Nrf2 dosage. Our results are consistent with previous work indicating that Nrf2 activation via Keap1 deletion promotes Kras$^{G12D/+}$ tumor initiation[27,28]. Surprisingly, we find that Nrf2 activation impairs tumor progression, which is correlated with Nrf2 dosage. In our previous study with Keap1$^{R554Q/R554Q}$ mice, we reported smaller tumors in the KP model[30], which we now find is due to impaired tumor progression mediated by Nrf2 hyperactivation. Because most prior studies from other groups did not specifically analyze tumor grade[25,27,34,35], they may not have captured this effect of Nrf2 on tumor progression. Alternatively, there may be biological differences between Keap1 deletion and mutation.

Our findings that tumors downregulate Nrf2 to select for a level permissive for tumor progression and that ectopic NRF2 expression antagonizes the proliferation and viability

of human NSCLC cells are supported by our finding that single copy deletion of Nrf2 can rescue the homozygous Keap1^{R554Q}-mediated block in tumor progression. This result demonstrates a direct role for Nrf2 hyperactivation but we cannot exclude the possibility that alternative KEAP1 substrates, such as PGAM5[45], PALB2[46], MCM3[47] or EMSY[48], contribute to the block in tumor progression. In agreement with our findings, a Keap1-binding defective Nrf2^{E79Q} mouse model of small-cell lung cancer (SCLC) with p53/ p16 inactivation[53] also displayed Nrf2 downregulation in aggressive SCLC tumors. The exact mechanism(s) by which high NRF2 activity impairs tumor cell proliferation and tumor progression remains to be determined. The dosage-dependent effects of Nrf2 are reminiscent of what has been observed for oncogenes like Ras and Myc, where low levels promote transformation and proliferation, and high levels promote senescence or death[54,55].

Given that KEAP1 and NRF2 mutations are found with a high frequency in human NSCLC and are associated with poor outcomes, these findings raise the question of under which contexts NRF2 activation provides an advantage. There are multiple features of patient tumors not captured by our mouse models that remain to be examined. Our models did not develop metastases with sufficient frequency, precluding an examination of the influence of Nrf2 activation on metastasis as reported previously[25]. Moreover, NRF2 protects against carcinogens[4-7] and mice are not exposed to smoking and other environmental toxins under which NRF2 activation may promote survival. Finally, we have not tested the response of these models to therapy. Overall, our work demonstrates the context-dependence of the role of NRF2 during the complex stages of tumorigenesis and warrants further investigation into how NRF2 elicits distinct responses.

Methods

Mice
Mice were housed and bred in accordance with the ethical regulations and approval of the IACUC (protocols #: IS00003893R and IS00007922R). Generation of the Keap1

targeting vector was previously described[30]. Briefly, the *CA-Keap1^R554Q* allele (Keap1^tm1Gmdn, MGI: 7327097) was made by inserting a wild-type *Keap1* cDNA containing exons 3-5 flanked by loxP sites upstream of the R554Q mutation in exon 4 of the *Keap1* gene. *Keap1* was targeted in C10 murine ES cells and cells were selected with blasticidin. To make the *LSL-Nfe2l2^D29H* allele (Nfe2l2^tm1Gmdn, MGI: 7327101), a STOP cassette flanked by loxP sites was inserted into intron 1 and codon 29 in endogenous exon 2 was mutated from an aspartic acid to a histidine. The endogenous *Nfe2l2* locus was targeted in C10 murine ES cells and puromycin was used to select positive cells. For both alleles, positive clones were screened by copy number real-time PCR and injected into blastocysts. Genotyping primers were as follows: for the *Keap1^R554Q* allele: Mutant forward: 5'-ATGGCCACACTTTTCTGGAC-3'; wild-type (WT) forward: 5'-GGGGGTAGAGGGAGGAGAAT-3'; Common reverse: 5'-GCCACCCTATTCACAGACCA-3'. The WT PCR product was 326 bp and the mutant PCR product 584 bp. For the *Nfe2l2^D29H* allele: WT forward: 5'-GAGGCAGGTAGTTCTCTGAGTTTG-3'; Common reverse: 5'-GCAAATGCACTGAGACACTCAT-3'; Mutant forward: 5'-CTAGCCACCATGGCTTGAGT-3'. The WT PCR product was 189 bp and the mutant PCR product 282 bp. All mice were maintained on a mixed C57BL/6 genetic background. In addition to Keap1^R554Q and Nrf2^D29H mice, p53^flox (RRID:IMSR_JAX:008462); Lkb1^flox (RRID:IMSR_JAX:014143); Nrf2^flox (RRID:IMSR_JAX:025433); and LSL-Kras^G12D/+ (RRID:IMSR_JAX:008179) mice were used. For mouse lung tumor studies, intranasal installation of 2.5 x 10^7 PFU adenoviral-Cre (University of Iowa) was used to induce lung tumors as previously described[43]. Adenoviral infections were performed under isofluorane anesthesia.

Murine embryonic fibroblast generation and culture

MEFs were isolated from E13.5-14.5-day old embryos and maintained in pyruvate-free DMEM (Corning) containing 10% FBS, 100 units/mL penicillin and 100 μg/mL streptomycin (Gibco) in a humidified incubator with 5% CO_2 and 95% air at 37°C. MEFs were used within four passages and infected with control empty adenovirus or adenoviral-Cre (University of Iowa) at an approximate multiplicity of infection of 500.

RNA-sequencing preparation and analysis

Samples were prepared using the RNeasy plus mini kit (Qiagen, 74134). RNA quality was checked with the QIAxcel RNA QC kit (Qiagen, 929104). Additional RNA QC, sequencing, mapping to the mouse genome, and analysis were performed by Novogene. Differentially expressed genes (DESeq2) with $p < 0.05$ were included in the volcano plot.

LC-HRMS conditions for MEF non-targeted metabolomics

Sample preparation and conditions were previously described[30]. Briefly, cells were extracted in 80% methanol and analyzed by liquid chromatography-high resolutions mass spectrometry (LC-HRMS) using a Vanquish UPLC systems coupled to a Q Exactive HF mass spectrometer. A Luna 3 μm HILIC 200 Å, LC Column 100 × 2 mm (Phenomenex, Torrance, CA, Part N: 00D-4449-B0) was used for chromatographic separation. Both positive and negative mode were used for the MS1 scan. Data analysis was done using El Maven v0.3.1[56]. The identification of metabolites was based on comparison of retention time and m/z value of sample peaks with an internal library (MSMLS Library, Sigma Aldrich).

Immunohistochemistry (IHC)

Mouse lung tissue was fixed with 10% formalin overnight, transferred to 70% ethanol and paraffin embedded to be sectioned. Unstained tissue sections were de-paraffinized in xylene followed by rehydration in a graded alcohol series. Antigen retrieval was performed by boiling in 10mM citrate buffer (pH 6). Antibodies used for IHC include affinity-purified NRF2 (1:150)[57] and NQO1 (Sigma Aldrich, RRID:AB_1079501, 1:500). Following overnight incubation at 4°C in primary antibody, the ImmPRESS HRP goat anti-rabbit kit (Vector Laboratories, RRID:AB_2631198) was used as directed by manufacturer's instructions. DAB peroxidase (HRP) substrate (Vector Laboratories, SK-4105) was used to develop immunohistochemical staining, followed by counterstaining with hematoxylin (Vector Laboratories, H-3404). Slides were scanned with the Aperio imager and the DAB optical density of at least five representative regions/ mouse was

analyzed with QuPath software[58]. Representative images were captured using the Axio Lab.A1 microscope (Carl Zeiss Microimaging Inc.).

Tumor grading analysis and histology

Lung tumor grading was performed manually as previously described[44]. Tumor grading distribution percentages were calculated by dividing the number of tumors in a specific grade by the total number of tumors per mouse. Tumor burden by grade was calculated by dividing the area of the lung covered by a specific tumor grade by the total lung area.

NSCLC cell lines and culture

Human lung cancer cell lines used include H1944 (RRID:CVCL_1508), H322 (RRID:CVCL_1556), A549 (RRID:CVCL_0023), HCC15 (RRID:CVCL_2057), and H460 (RRID:CVCL_0459) and were previously described[22]. Cells were cultured in RPMI 1640 (Gibco) containing 5% FBS without antibiotics in a humidified incubator with 5% CO_2 and 95% air at 37°C. Cells were screened regularly and confirmed to be free of mycoplasma with the MycoAlert kit (Lonza).

Lentivirus generation and infection of NSCLC cells

Lentiviruses were made by transfecting Lenti-X 293T cells (Takara 632180) overnight with polyethylenimine (PEI), lentiviral plasmid (pLX317-NRF2[59] or the control pLX317 empty vector[30]), and packaging plasmids pCMV-dR8.2 dvpr (RRID:Addgene_8455) and pCMV-VSV-G (RRID:Addgene_8454) in DMEM containing 10% FBS. To generate NRF2-overexpressing cells, NSCLC cells were transduced for 24 hrs with lentiviruses in medium containing polybrene (8 µg/mL). After transduction, infected cells were selected with 0.5 µg/mL (H1944, H322, H460) or 1 µg/mL (A549, HCC15) puromycin for 72 hrs. Immediately following selection, cells were seeded in respective puromycin concentrations for the indicated assays.

Cell proliferation and cell death assays

NSCLC cells were monitored with the CELLCYTE X[TM] live cell imaging instrument (Cytena) over the course of 96 hours. Prior to imaging, SYTOX Green nucleic acid stain

(Thermo Fisher Scientific, S7020) was added to medium at a final concentration of 20 nM. Images were acquired from each well at 8-hour intervals and analyzed using CellCyte Studio (CELLINK). Cell confluency was represented as the % of the image covered by cells. The number of dead cells was normalized to cell confluency [number of Sytox Green positive cells/ mm^2/ cell confluency]. The area under the curve (AUC) values were calculated by summing the proliferation or normalized dead cell number at each time point.

Western blotting

Cells were lysed in ice-cold RIPA buffer with protease inhibitors (Fisher Scientific, PIA32955) followed by sonication in a water bath sonicator (Diagenode). Protein was quantified using the DC protein assay (Bio-Rad). Lysates were prepared with 6X SDS sample buffer containing 12% (v/v) β-ME (VWR) and separated on Bolt^{TM} or NuPAGE 4-12% Bis-Tris gels (Invitrogen). SDS-PAGE separation was followed by transfer to 0.45 μm nitrocellulose membranes (GE Healthcare). The membranes were blocked in 5% non-fat milk in Tris-buffered saline with 0.1% Tween 20 (TBST). For immunoblotting, the following antibodies diluted in 5% milk in TBST were used: KEAP1 (Millipore Sigma, RRID:AB_2921362, 1:2000), NRF2 (Cell Signaling Technologies, D1Z9C, RRID:AB_2715528, 1:1000), NQO1 (Sigma Aldrich, RRID:AB_1079501, 1:1000), GCLC (Sana Cruz Biotechnology, H-5, RRID:AB_2736837, 1:1000), xCT (abcam, RRID:AB_778944, 1:1000), GSR (Santa Cruz Biotechnology, RRID:AB_2295121, 1:1000), β-actin (Invitrogen AM4302, RRID:AB_2536382, 1:100,000). HRP secondary antibodies used include goat anti-rabbit IgG (Jackson ImmunoResearch Labs, RRID: AB_23135627), goat anti-mouse IgG (Jackson ImmunoResearch Labs, RRID:AB_10015289), and goat anti-rat IgG (Jackson ImmunoResearch Labs, RRID:AB_2338128). Membranes were developed with Clarity ECL substrate (Bio-Rad) or a luminol-based homemade ECL substrate.

Soft agar colony formation assays

6-well plates were coated with a 0.8% agar prepared in RPMI. NSCLC cells were then seeded in 0.4% agar in RPMI. After the cell/ agar mixture solidified, RPMI medium

containing 10% FBS, Pen/Strep and puromycin was added to each well and replenished every few days. Colonies were allowed to form for 10–16 days, and wells were stained with 0.01% crystal violet in a 4% paraformaldehyde in PBS solution. Plates were scanned on a flatbed scanner and ImageJ was used to quantify colonies.

DepMap Analysis

NFE2L2 dependency scores were downloaded from the DepMap database v. 22Q2[52]. Values were plotted from CRISPR (DepMap 22Q2 Public+Score, Chronos) for non-small cell lung cancer cell lines that we previously evaluated for high or low NRF2 activity[22].

Statistical analysis

Graphpad Prism9 software was used for statistical analyses and P values < 0.05 were considered significant, with symbols as follows: *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. All data is represented as mean +/- standard deviation unless otherwise stated. For all experiments, similar variances between groups were observed.

Acknowledgements

This work was supported by grants from the NIH/NCI (R37-CA230042), the American Lung Association (LCDA-498544) and the American Cancer Society's Institutional Research Grant to G.M.D. This work was also supported by the Analytical Microscopy, the Tissue, and Proteomics/Metabolomics Core Facilities at the Moffitt Cancer Center, an NCI designated Comprehensive Cancer Center (P30-CA076292). We would like to thank the DeNicola lab members and Dr. Ana Gomes for the helpful discussions. We thank Cheyenne Schneider for proofreading.

Author Contributions

G.M.D. conceived the project and G.M.D. and J.M.D. designed the experiments. G.M.D., I.I.C.C. and F.A.K. designed and generated the Nrf2 and Keap1 mutant mice. G.M.D. performed the RNA sequencing. G.M.D. and Y.P.K. performed and analyzed the

metabolomics. Maintenance of animal colonies, generation of experimental animals, and collections of tissues were performed by A.F., with assistance from S.C. and N.F.P. J.M.D. performed cell line experiments, immunohistochemistry, and tumor histology analyses. J.R.P. and E.E.S. generated the affinity-purified NRF2 antibody used for immunohistochemistry. J.M.D. and G.M.D. wrote the manuscript, and all authors reviewed it. G.M.D. acquired funding and supervised the study.

Competing interests

The authors declare no competing financial or personal interests.

Additional information

Correspondence and requests for materials should be addressed to G.M.D.

Figure 1. Development of mutant *Keap1* and *Nfe2l2* alleles found in human lung cancer. (A) The conditionally active (CA)-*Keap1^{R554Q}* allele was generated by inserting a loxP-flanked, wild type (WT) *Keap1* cDNA containing exons 3-5 into intron 2 and introducing the R554Q mutation into endogenous exon 4 of the *Keap1* gene. Prior to intranasal installation of adenoviral-Cre recombinase Keap1 is wild type. Once the floxed cargo is excised, mutant Keap1^{R554Q} is expressed. PA = poly A signal. **(B)** WT *Keap1* allele. WT Keap1 is expressed before or after adenoviral-Cre recombinase. **(C)** The Lox-STOP-Lox *(LSL)-Nfe2l2^{D29H}* allele was created by inserting a STOP cassette flanked by loxP sites into intron 1 and introducing the D29H mutation in endogenous exon 2 of the *Nfe2l2* gene. Following Cre-mediated excision of the STOP cassette, mutant Nrf2^{D29H} is expressed. **(D)** WT *Nfe2l2* allele. WT Nrf2 is expressed before or after adenoviral-Cre recombinase. **(E)** Volcano plot of RNA-sequencing data from murine embryonic fibroblasts (MEFs) expressing Keap1^{R554Q/R554Q} compared to Keap1^{+/+}. N=3, representative of two individual MEF lines. **(F)** Volcano plot of RNA-sequencing data from MEFs expressing Nrf2^{D29H/D29H} compared to Nrf2^{LSL/LSL}, which lack Nrf2 expression. N=3, representative of two individual MEF lines. **(G)** Volcano plot of LC-HRMS metabolomics profiling comparing Keap1^{R554Q/R554Q} MEFs to Keap1^{+/+} MEFs from our previous study[30]. N=3. **(H)** LC-HRMS metabolomics profiling comparing Nrf2^{D29H/D29H} MEFs to Nrf2^{LSL/LSL} MEFs. N=3, representative of two individual MEF lines.

Figure 2. Mutation of Keap1 or Nrf2 is not sufficient to initiate lung tumorigenesis.
(A) Overall survival of Keap1/Nrf2 mutant mice. Keap1/Nrf2$^{+/+}$ (n=17), Keap1$^{R554Q/+}$ (n=8); Keap1$^{R554Q/R554Q}$ (n=11); Nrf2$^{D29H/+}$ (n=15). Ns= not significant (Log-rank (Mantel-Cox) test). **(B)** Lung tumor-free survival of Keap1/Nrf2 mutant mice. Keap1/Nrf2$^{+/+}$ (n=11), Keap1$^{R554Q/+}$ (n=4); Keap1$^{R554Q/R554Q}$ (n=8); Nrf2$^{D29H/+}$ (n=11). **(C)** Representative H&E of mouse lung depicting normal bronchiolar and alveolar cells (scale bars = 100µM (top panel), 20 µM (bottom panel)).

Figure 3. Mutation of Keap1 or Nrf2 is not sufficient to initiate lung tumorigenesis with p53 or Lkb1 loss. (A) Overall survival of p53$^{fl/fl}$ mice expressing wild-type or mutant Keap1/Nrf2. Keap1/Nrf2$^{+/+}$ (n=15); Keap1$^{R554Q/+}$ (n=7); Keap1$^{R554Q/R554Q}$ (n=18); Nrf2$^{D29H/+}$ (n=18). Ns= not significant (Log-rank (Mantel-Cox) test). **(B)** Lung tumor-free survival of p53$^{fl/fl}$ mice expressing wild-type or mutant Keap1/Nrf2. Keap1/Nrf2$^{+/+}$ (n=11); Keap1$^{R554Q/+}$ (n=6); Keap1$^{R554Q/R554Q}$ (n=16); Nrf2$^{D29H/+}$ (n=10). **(C)** Representative H&E of mouse lung depicting bronchiolar and alveolar cells of the p53$^{fl/fl}$ models (scale bars = 100μM (top panel), 20μM (middle panel), 10μM (bottom panel)). **(D)** Overall survival of Lkb1$^{fl/fl}$ mice expressing wild-type or mutant Keap1/Nrf2. Keap1/Nrf2$^{+/+}$ (n=11); Keap1$^{R554Q/+}$ (n=7); Keap1$^{R554Q/R554Q}$ (n=11); Nrf2$^{D29H/+}$ (n=5). Ns= not significant (Log-rank (Mantel-Cox) test). **(E)** Lung tumor-free survival of Lkb1$^{fl/fl}$ mice expressing wild-type or mutant Keap1/Nrf2. Keap1/Nrf2$^{+/+}$ (n=11); Keap1$^{R554Q/+}$ (n=6); Keap1$^{R554Q/R554Q}$ (n=9); Nrf2$^{D29H/+}$ (n=4). **(F)** Representative H&E of mouse lung depicting bronchiolar and alveolar cells of the Lkb1$^{fl/fl}$ models (scale bars = 100μM (top panel), 20μM (middle panel), 10μM (bottom panel)). For **A, B, D, E,** mice were infected intranasally with adenoviral-Cre, followed by collection at 500 days to analyze lung tissue histology.

Figure 4. Keap1/Nrf2 mutation cooperates with Kras$^{G12D/+}$ to promote lung tumor initiation and early progression. (A) Overall survival of Keap1 mutant mice with Kras$^{G12D/+}$ mutation. Keap1$^{+/+}$ (n=16); Keap1$^{R554Q/+}$ (n=18); Keap1$^{R554Q/R554Q}$ (n=18). **(B)** Overall survival of Nrf2 mutant mice with Kras$^{G12D/+}$ mutation. Nrf2$^{+/+}$ (n=14); Nrf2$^{D29H/+}$ (n=12). Ns = not significant (Log-rank (Mantel-Cox) test). **(C)** Representative immunohistochemical (IHC) staining of Nrf2 in Keap1/Nrf2 mutant tumors with Kras$^{G12D/+}$ mutation (scale bars = 20 μM). **(D, E)** Quantification of DAB optical density (OD) in AAH and BH for Nrf2 (nuclear) **(D)** and Nrf2 target Nqo1 (whole cell) **(E)** IHC staining. N= 3 mice and >9,500 tumor cells per genotype. *$p<0.05$ (one-way ANOVA). **(F)** Representative whole lung H&E-stained section (scale bars = 2000 μM). **(G)** Tumor number per mouse in Keap1/Nrf2 mutant models normalized to lung area. *$p<0.05$ (one-way ANOVA). **(H)** Distribution of tumor grades across Keap1/Nrf2 mutant models. *$p<0.05$ (unpaired t test with Holm-Sidak's multiple comparisons test). AAH = atypical adenomatous hyperplasia. BH = bronchiolar hyperplasia. **(I)** Fraction of lung tumor burden by grade (lung tumor area/ total lung area per grade). *$p<0.05$ (unpaired t test with Holm-Sidak's multiple comparisons test). For both **(H)** and **(I)** n=10 mice and ≥2,000 tumors per genotype.

Figure 5. Homozygous Keap1^{R554Q} impairs adenocarcinoma progression in the Kras$^{G12D/+}$; p53$^{fl/fl}$ model. (A) Overall survival of Kras$^{G12D/+}$; p53$^{fl/fl}$ mice with Keap1 mutation. Keap1$^{+/+}$ (n=20); Keap1$^{R554Q/+}$ (n=11); Keap1$^{R554Q/R554Q}$ (n=25). **(B)** Overall survival of Kras$^{G12D/+}$; p53$^{fl/fl}$ mice with Nrf2 mutation. Nrf2$^{+/+}$ (n=25); Nrf2$^{D29H/+}$ (n=29). Ns = not significant (Log-rank (Mantel-Cox) test). **(C)** Representative immunohistochemical (IHC) staining of Nrf2 in Kras$^{G12D/+}$; p53$^{fl/fl}$ mice with Keap1/Nrf2 mutation (scale bars = 20 µM). **(D, E)** Quantification of DAB optical density (OD) in AAH and BH for Nrf2 (nuclear) **(D)** and Nrf2 target Nqo1 (whole cell) **(E)** IHC staining. N= 3 mice and >7,800 tumor cells per genotype. *$p<0.05$ (one-way ANOVA). **(F)** Representative whole lung H&E stained section (scale bars = 2000 µM). **(G)** Distribution of tumor grades across Keap1/Nrf2 mutant models. *$p<0.05$ (unpaired t test with Holm-Sidak's multiple comparisons test). **(H)** Fraction of lung tumor burden by grade (lung tumor area per grade/ total lung area). *$p<0.05$ (unpaired t test with Holm-Sidak's multiple comparisons test). For both **(G)** and **(H)** n$\geq$9 mice and $\geq$1,900 tumors per genotype. Only one grade 5 tumor was found in the Keap1$^{R554Q/R554Q}$ cohort, and therefore was excluded from these analyses.

Figure 6. Nrf2 expression and activity is reduced in higher-grade tumors. (A) Representative Nrf2 and Nqo1 IHC staining in grade 1 and 4 tumors from Kras[G12D/+]; p53[fl/fl] mice with Keap1 or Nrf2 mutation (scale bars = 20 µM). **(B, C)** Heatmaps depicting the median DAB optical density from IHC staining for Nrf2 (nuclear) **(B)** and the Nrf2 target Nqo1 (whole cell) **(C)**. N = 3 mice per genotype, $\geq$20,000 tumor cells per mouse. Only one grade 5 tumor was found in the Keap1[R554Q/R554Q] cohort, and therefore was excluded from these analyses.

Figure 7. NRF2 overexpression impairs lung cancer cell proliferation, viability, and soft agar colony formation. (A) Dependency scores obtained from DepMap[52] and represented as *NFEL2* 22Q2 Public+Score, Chronos for NSCLC cell lines previously determined to have high or low NRF2 activity[22]. NRF2 mutant line symbols are represented by yellow, KEAP1 mutant lines by dark red or dark blue, and KEAP1 deleted lines by green. ***p<0.0001 (unpaired t-test). **(B)** Western blot analysis of NRF2, β-actin, and NRF2 target GCLC, xCT, and GSR expression in *KEAP1* mutant lung cancer cell lines transduced with PLX317-empty vector (EV) or PLX317-NRF2 (NRF2). **(C)** Representative images of HCC15 cells transduced with EV or NRF2 demonstrating cell confluency (enhanced contour) and cell death (Sytox Green) (scale bars = 100 μM). **(D, E)** Analysis of EV and NRF2 HCC15 cell proliferation and death over 96 hours. Proliferation is represented as % confluency at each time point, and cell death as the number of Sytox Green positive cells per area normalized to % confluency. N=3 technical replicates per cell line, and two independent experiments. **(F, G)** Area under the curve (AUC) analysis of cell proliferation **(F)** and Sytox Green-positive cell death **(G)** in KEAP1 mutant lung cancer cells lines +/- NRF2, normalized to empty vector control. *p<0.05 (one-way ANOVA). For **(C-G)** NSCLC cells were seeded in triplicate in 96-well plates at a density of 2,500 cells/ well. **(H)** Representative images of H460 and HCC15 soft agar colony formation +/- NRF2. **(I)** Quantification of soft agar colony number of *KEAP1* mutant lung cancer cell lines. *p<0.05 (one-way ANOVA). N=3 technical replicates per cell line, two independent experiments. For **(H, I),** 5,000 cells per well were seeded in 6-well plates in triplicate.

A

Kras$^{G12D/+}$; p53$^{fl/fl}$ Survival

Overall Survival vs Days (post-Cre)

- Keap1$^{+/+}$; Nrf2$^{+/+}$
- Keap1$^{+/+}$; Nrf2$^{fl/+}$
- Keap1$^{R554Q/R554Q}$; Nrf2$^{+/+}$
- Keap1$^{R554Q/R554Q}$; Nrf2$^{fl/+}$

ns

B

Kras$^{G12D/+}$; p53$^{fl/fl}$

| Keap1$^{+/+}$; Nrf2$^{+/+}$ | Keap1$^{+/+}$; Nrf2$^{fl/+}$ | Keap1$^{R554Q/R554Q}$; Nrf2$^{+/+}$ | Keap1$^{R554Q/R554Q}$; Nrf2$^{fl/+}$ |

C

Kras$^{G12D/+}$; p53$^{fl/fl}$ Tumor Grade Distribution

% Tumors vs AAH, BH, Grade 1, Grade 2, Grade 3, Grade 4, Grade 5

- Keap1$^{+/+}$; Nrf2$^{+/+}$
- Keap1$^{+/+}$; Nrf2$^{fl/+}$
- Keap1$^{R554Q/R554Q}$; Nrf2$^{+/+}$
- Keap1$^{R554Q/R554Q}$; Nrf2$^{fl/+}$

D

Kras$^{G12D/+}$; p53$^{fl/fl}$ Tumor Burden

Tumor burden (% of whole lung) vs AAH, BH, Grade 1, Grade 2, Grade 3, Grade 4, Grade 5

- Keap1$^{+/+}$; Nrf2$^{+/+}$
- Keap1$^{+/+}$; Nrf2$^{fl/+}$
- Keap1$^{R554Q/R554Q}$; Nrf2$^{+/+}$
- Keap1$^{R554Q/R554Q}$; Nrf2$^{fl/+}$

Figure 8. Single copy Nrf2 deletion rescues homozygous Keap1^{R554Q}-mediated adenocarcinoma progression impairment in the Kras$^{G12D/+}$; p53$^{fl/fl}$ model. (A) Overall survival of Kras$^{G12D/+}$; p53$^{fl/fl}$ mice with Keap1 mutation and/ or single copy Nrf2 deletion. Keap1$^{+/+}$; Nrf2$^{+/+}$ (n=11), Keap1$^{+/+}$; Nrf2$^{fl/+}$ (n=7), Keap1$^{R554Q/R554Q}$; Nrf2$^{+/+}$ (n=6), Keap1$^{R554Q/R554Q}$; Nrf2$^{fl/+}$ (n=5). Ns = not significant (Log-rank (Mantel-Cox) test). **(B)** Representative whole lung H&E-stained sections (scale bars = 2000 μM). **(C)** Distribution of tumor grades across Keap1 mutant/ Nrf2-deleted models. *p<0.05 (unpaired t test with Holm-Sidak's multiple comparisons test). **(D)** Fraction of lung tumor burden by grade (lung tumor area per grade/ total lung area). *p<0.05 (unpaired t test with Holm-Sidak's multiple comparisons test). For both **(C)** and **(D)** n≥4 mice and ≥700 tumors per genotype were analyzed. No grade 5 tumors were found in the Keap1$^{R554Q/R554Q}$; Nrf2$^{+/+}$ cohort.

References

DeBlasi, J. M. & DeNicola, G. M. Dissecting the Crosstalk between NRF2 Signaling and Metabolic Processes in Cancer. *Cancers (Basel)* **12**, doi:10.3390/cancers12103023 (2020).

He, F., Ru, X. & Wen, T. NRF2, a Transcription Factor for Stress Response and Beyond. *Int J Mol Sci* **21**, doi:10.3390/ijms21134777 (2020).

Itoh, K. *et al.* Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes Dev* **13**, 76-86, doi:10.1101/gad.13.1.76 (1999).

Ramos-Gomez, M., Dolan, P. M., Itoh, K., Yamamoto, M. & Kensler, T. W. Interactive effects of nrf2 genotype and oltipraz on benzo[a]pyrene–DNA adducts and tumor yield in mice. *Carcinogenesis* **24**, 461-467, doi:10.1093/carcin/24.3.461 (2003).

Ramos-Gomez, M. *et al.* Sensitivity to carcinogenesis is increased and chemoprotective efficacy of enzyme inducers is lost in <i>nrf2</i> transcription factor-deficient mice. *Proceedings of the National Academy of Sciences* **98**, 3410-3415, doi:doi:10.1073/pnas.051618798 (2001).

Aoki, Y. *et al.* Accelerated DNA adduct formation in the lung of the Nrf2 knockout mouse exposed to diesel exhaust. *Toxicol Appl Pharmacol* **173**, 154-160, doi:10.1006/taap.2001.9176 (2001).

Iida, K. *et al.* Nrf2 Is Essential for the Chemopreventive Efficacy of Oltipraz against Urinary Bladder Carcinogenesis. *Cancer Research* **64**, 6424-6431, doi:10.1158/0008-5472.Can-04-1906 (2004).

Kensler, T. W. & Wakabayashi, N. Nrf2: friend or foe for chemoprevention? *Carcinogenesis* **31**, 90-99, doi:10.1093/carcin/bgp231 (2010).

Schmidlin, C. J., Shakya, A., Dodson, M., Chapman, E. & Zhang, D. D. The intricacies of NRF2 regulation in cancer. *Semin Cancer Biol* **76**, 110-119, doi:10.1016/j.semcancer.2021.05.016 (2021).

Telkoparan-Akillilar, P., Suzen, S. & Saso, L. Pharmacological Applications of Nrf2 Inhibitors as Potential Antineoplastic Drugs. *Int J Mol Sci* **20**, doi:10.3390/ijms20082025 (2019).

Network, C. G. A. R. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* **511**, 543-550, doi:10.1038/nature13385 (2014).

Network, C. G. A. R. Comprehensive genomic characterization of squamous cell lung cancers. *Nature* **489**, 519-525, doi:10.1038/nature11404 (2012).

Singh, A. *et al.* Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. *PLoS Med* **3**, e420, doi:10.1371/journal.pmed.0030420 (2006).

Shibata, T. *et al.* Cancer related mutations in NRF2 impair its recognition by Keap1-Cul3 E3 ligase and promote malignancy. *Proc Natl Acad Sci U S A* **105**, 13568-13573, doi:10.1073/pnas.0806268105 (2008).

Wang, R. *et al.* Hypermethylation of the Keap1 gene in human lung cancer cell lines and lung cancer tissues. *Biochem Biophys Res Commun* **373**, 151-154, doi:10.1016/j.bbrc.2008.06.004 (2008).

Kim, Y. R. *et al.* Oncogenic NRF2 mutations in squamous cell carcinomas of oesophagus and skin. *J Pathol* **220**, 446-451, doi:10.1002/path.2653 (2010).

Zhang, P. *et al.* Loss of Kelch-like ECH-associated protein 1 function in prostate cancer cells causes chemoresistance and radioresistance and promotes tumor growth. *Mol Cancer Ther* **9**, 336-346, doi:10.1158/1535-7163.MCT-09-0589 (2010).

Solis, L. M. *et al.* Nrf2 and Keap1 abnormalities in non-small cell lung carcinoma and association with clinicopathologic features. *Clin Cancer Res* **16**, 3743-3753, doi:10.1158/1078-0432.CCR-09-3352 (2010).

Sporn, M. B. & Liby, K. T. NRF2 and cancer: the good, the bad and the importance of context. *Nat Rev Cancer* **12**, 564-571, doi:10.1038/nrc3278 (2012).

Singh, A., Bodas, M., Wakabayashi, N., Bunz, F. & Biswal, S. Gain of Nrf2 function in non-small-cell lung cancer cells confers radioresistance. *Antioxidants and Redox Signaling*, doi:10.1089/ars.2010.3219 (2010).

Denicola, G. M. *et al.* Oncogene-induced Nrf2 transcription promotes ROS detoxification and tumorigenesis. *Nature* **475**, 106-110, doi:10.1038/nature10189 (2011).

DeNicola, G. M. *et al.* NRF2 regulates serine biosynthesis in non-small cell lung cancer. *Nature Genetics* **47**, 1475-1481, doi:10.1038/ng.3421 (2015).

Mitsuishi, Y. *et al.* Nrf2 redirects glucose and glutamine into anabolic pathways in metabolic reprogramming. *Cancer Cell* **22**, 66-79, doi:10.1016/j.ccr.2012.05.016 (2012).

Sayin, V. I. *et al.* Activation of the NRF2 antioxidant program generates an imbalance in central carbon metabolism in cancer. *Elife* **6**, doi:10.7554/eLife.28083 (2017).

Lignitto, L. *et al.* Nrf2 Activation Promotes Lung Cancer Metastasis by Inhibiting the Degradation of Bach1. *Cell* **178**, 316-329.e318, doi:10.1016/j.cell.2019.06.003 (2019).

Best, S. A. *et al.* Synergy between the KEAP1/NRF2 and PI3K Pathways Drives Non-Small-Cell Lung Cancer with an Altered Immune Microenvironment. *Cell Metabolism* **27**, 935-943.e934, doi:10.1016/j.cmet.2018.02.006 (2018).

Best, S. A. *et al.* Distinct initiating events underpin the immune and metabolic heterogeneity of KRAS-mutant lung adenocarcinoma. *Nature Communications* **10**, 1-14, doi:10.1038/s41467-019-12164-y (2019).

Hayashi, M. *et al.* Microenvironmental Activation of Nrf2 Restricts the Progression of Nrf2-Activated Malignant Tumors. *Cancer Res* **80**, 3331-3344, doi:10.1158/0008-5472.Can-19-2888 (2020).

Jeong, Y. *et al.* Role of KEAP1/NRF2 and TP53 mutations in lung squamous cell carcinoma development and radiation resistance. *Cancer Discovery* **7**, 86-101, doi:10.1158/2159-8290.CD-16-0127 (2017).

Kang, Y. P. *et al.* Cysteine dioxygenase 1 is a metabolic liability for non-small cell lung cancer. *eLife*, doi:10.7554/eLife.45572 (2019).

Romero, R. *et al.* Keap1 mutation renders lung adenocarcinomas dependent on Slc33a1. *Nature Cancer* **1**, 589-602, doi:10.1038/s43018-020-0071-1 (2020).

Romero, R. *et al.* Keap1 loss promotes Kras-driven lung cancer and results in dependence on glutaminolysis. *Nature Medicine*, doi:10.1038/nm.4407 (2017).

Singh, A. *et al.* NRF2 Activation Promotes Aggressive Lung Cancer and Associates with Poor Clinical Outcomes. *Clinical Cancer Research* **27**, 877, doi:10.1158/1078-0432.CCR-20-1985 (2021).

Foggetti, G. *et al.* Genetic Determinants of EGFR-Driven Lung Cancer Growth and Therapeutic Response In Vivo. *Cancer Discovery* **11**, 1736-1753, doi:10.1158/2159-8290.Cd-20-1385 (2021).

Rogers, Z. N. *et al.* Mapping the in vivo fitness landscape of lung adenocarcinoma tumor suppression in mice. *Nature Genetics*, doi:10.1038/s41588-018-0083-2 (2018).

Cai, H. *et al.* A Functional Taxonomy of Tumor Suppression in Oncogenic KRAS-Driven Lung Cancer. *Cancer Discov* **11**, 1754-1773, doi:10.1158/2159-8290.Cd-20-1325 (2021).

Rogers, Z. N. *et al.* A quantitative and multiplexed approach to uncover the fitness landscape of tumor suppression in vivo. *Nature Methods* **14**, 737-742, doi:10.1038/nmeth.4297 (2017).

Hast, B. E. *et al.* Cancer-derived mutations in KEAP1 impair NRF2 degradation but not ubiquitination. *Cancer Research*, doi:10.1158/0008-5472.CAN-13-1655 (2014).

Tong, K. I. *et al.* Different Electrostatic Potentials Define ETGE and DLG Motifs as Hinge and Latch in Oxidative Stress Response. *Molecular and Cellular Biology*, doi:10.1128/mcb.00753-07 (2007).

Taguchi, K. *et al.* Genetic Analysis of Cytoprotective Functions Supported by Graded Expression of Keap1. *Molecular and Cellular Biology*, doi:10.1128/mcb.01591-09 (2010).

Wakabayashi, N. *et al.* Keap1-null mutation leads to postnatal lethality due to constitutive Nrf2 activation. *Nature Genetics*, doi:10.1038/ng1248 (2003).

Huppke, P. *et al.* Activating de novo mutations in NFE2L2 encoding NRF2 cause a multisystem disorder. *Nature Communications* **8**, 818, doi:10.1038/s41467-017-00932-7 (2017).

Jackson, E. L. *et al.* Analysis of lung tumor initiation and progression using conditional expression of oncogenic K-ras. *Genes and Development*, doi:10.1101/gad.943001 (2001).

Jackson, E. L. *et al.* The differential effects of mutant p53 alleles on advanced murine lung cancer. *Cancer Research* **65**, 10280-10288, doi:10.1158/0008-5472.CAN-05-2193 (2005).

Lo, S. C. & Hannink, M. PGAM5 tethers a ternary complex containing Keap1 and Nrf2 to mitochondria. *Exp Cell Res* **314**, 1789-1803, doi:10.1016/j.yexcr.2008.02.014 (2008).

Ma, J. *et al.* PALB2 interacts with KEAP1 to promote NRF2 nuclear accumulation and function. *Mol Cell Biol* **32**, 1506-1517, doi:10.1128/MCB.06271-11 (2012).

Mulvaney, K. M. *et al.* Identification and characterization of MCM3 as a kelch-like ECH-associated protein 1 (KEAP1) substrate. *Journal of Biological Chemistry* **291**, 23719-23733, doi:10.1074/jbc.M116.729418 (2016).

Marzio, A. *et al.* EMSY inhibits homologous recombination repair and the interferon response, promoting lung cancer immune evasion. *Cell* **185**, 169-183.e119, doi:10.1016/j.cell.2021.12.005 (2022).

Zhang, Y. *et al.* Emerging Substrate Proteins of Kelch-like ECH Associated Protein 1 (Keap1) and Potential Challenges for the Development of Small-Molecule Inhibitors of the Keap1-Nuclear Factor Erythroid 2-Related Factor 2 (Nrf2) Protein–Protein Interaction. *Journal of Medicinal Chemistry* **1**, doi:10.1021/acs.jmedchem.9b01865 (2020).

McCutcheon, D. C., Lee, G., Carlos, A., Montgomery, J. E. & Moellering, R. E. Photoproximity Profiling of Protein-Protein Interactions in Cells. *J Am Chem Soc* **142**, 146-153, doi:10.1021/jacs.9b06528 (2020).

Kitamura, H. & Motohashi, H. NRF2 addiction in cancer cells. *Cancer Sci* **109**, 900-911, doi:10.1111/cas.13537 (2018).

Institute, B. DepMap Public 22Q2 (2022).

Hamad, S. H. *et al.* TP53, CDKN2A/P16, and NFE2L2/NRF2 regulate the incidence of pure- and combined-small cell lung cancer in mice. *Oncogene* **41**, 3423-3432, doi:10.1038/s41388-022-02348-0 (2022).

Tuveson, D. A. *et al.* Endogenous oncogenic K-ras(G12D) stimulates proliferation and widespread neoplastic and developmental defects. *Cancer Cell* **5**, 375-387, doi:10.1016/s1535-6108(04)00085-6 (2004).

Murphy, D. J. *et al.* Distinct thresholds govern Myc's biological output in vivo. *Cancer Cell* **14**, 447-457, doi:10.1016/j.ccr.2008.10.018 (2008).

Clasquin, M. F., Melamud, E. & Rabinowitz, J. D. LC-MS data processing with MAVEN: a metabolomic analysis and visualization engine. *Curr Protoc Bioinformatics* **Chapter 14**, Unit14.11, doi:10.1002/0471250953.bi1411s37 (2012).

McLoughlin, M. R. *et al.* TrxR1, Gsr, and oxidative stress determine hepatocellular carcinoma malignancy. *Proc Natl Acad Sci U S A* **116**, 11408-11417, doi:10.1073/pnas.1903244116 (2019).

Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci Rep* **7**, 16878, doi:10.1038/s41598-017-17204-5 (2017).

Berger, A. H. *et al.* High-throughput Phenotyping of Lung Cancer Somatic Mutations. *Cancer Cell* **30**, 214-228, doi:10.1016/j.ccell.2016.06.022 (2016).