Leptospiral lipopolysaccharide dampens inflammation through upregulation of autophagy adaptor p62 and NRF2 signaling in macrophages

Delphine Bonhomme [1], Ignacio Santecchia [1], Pedro Escoll [2], Frédérique Vernel-Pauillac [1], Ivo G. Boneca [1] & Catherine Werts [1*]

[1] Institut Pasteur, Université Paris Cité, CNRS UMR6047, INSERM U1306, Unité de Biologie et Génétique de la Paroi Bactérienne, Paris, France

[2] Institut Pasteur, Université Paris Cité, CNRS UMR6047, Unité Biologie des Bactéries Intracellulaires, Paris, France

* corresponding author cwerts@pasteur.fr

ABSTRACT

Leptospira interrogans are pathogenic bacteria responsible for leptospirosis, a worldwide neglected zoonosis affecting yearly 1 million people. All vertebrates can be infected by leptospires. Some species like humans are susceptible to the disease whereas rodents such as mice are resistant and can even become asymptomatic renal carriers. Leptospires have been shown to avoid the activation of some innate immune receptors of the Toll-like receptors (TLRs) and NOD-like receptors families through their atypical microbe associated molecular patterns, and to escape some phagocyte functions. Autophagy is an essential nutrient-recycling cellular mechanism that is also crucial to clear invasive pathogens and is one of the many antimicrobial mechanisms that may be subverted upon bacterial infection. Here we showed that leptospires reduced the quantity of autophagosomes in murine and human macrophages. Moreover, leptospires infection induced a time-dependent specific accumulation of the autophagy adaptor p62 in puncta-like structures. Interestingly, we showed that these effects were not associated with an active leptospiral mechanism but were mediated by the recognition of the atypical leptospiral lipopolysaccharide (LPS) through TLR4 and TLR2. Moreover, we demonstrate in murine macrophages that leptospires induced, *via* p62 accumulation, the nuclear translocation of transcription factor NRF2, consequently triggering the retro-amplification of the p62/NRF2 axis. Finally, inhibition of p62/NRF2 signaling led to increased cytokines transcription and secretion. Overall, this suggests that the p62/NRF2 axis induced by leptospires contributes to dampening inflammation. These findings highlight a novel escape mechanism of leptospires, potentially participating in the stealthiness of the infection.

AUTHOR SUMMARY

Leptospira interrogans are pathogenic bacteria responsible for leptospirosis, a worldwide neglected zoonosis. Some species like humans may suffer from severe disease whereas rodents such as mice are asymptomatic renal carriers. Leptospires have been shown to escape different innate immune receptors of the host immune defenses, and largely escape phagocytosis. Autophagy is a host cell mechanism essential for nutrient-recycling but also crucial to fight and clear intracellular pathogens. Here we show by Western blot and automated microscopy, that infection with leptospires in human and murine macrophages did not trigger autophagy, although it triggered the specific accumulation of the autophagic adaptor p62. Interestingly, we show in murine macrophages that leptospiral infection also activated the nuclear translocation of the transcription factor NRF2. These effects were recapitulated with inactivated bacteria or using the leptospiral lipopolysaccharide (LPS). Finally, we show using RNA interference that inhibiting p62/NRF2 signaling led to increased inflammation. Overall, this suggests that the p62/NRF2 axis induced by the leptospiral LPS contributes to limiting the inflammation in macrophages. These findings highlight a novel mechanism of *Leptospira*, potentially contributing to their stealthiness.

INTRODUCTION

Leptospira interrogans are spirochete bacteria and the causative agent of leptospirosis, a worldwide neglected zoonosis that is currently reemerging because of climate change and causes around 60 000 deaths *per* year worldwide [1] Although all vertebrates can be infected by leptospires, they do not all present the same symptoms and susceptibility to the disease [2]. Humans as sensitive hosts may suffer from acute leptospirosis ranging from flu-like symptoms to potential multi-organ failure in 5-10% cases [1]. On the other hand, rodents such as mice and rats are resistant to acute illness and get chronically colonized upon infection [3]. Leptospires reach the proximal tubules in the kidneys and establish a stable colonization that leads to their excretion in the urine throughout the life of the animal, contributing to the spread of the zoonosis [3, 4].

The host immune defense relies on humoral and cellular components, such as the complement system, immunoglobulins, and phagocytes that patrol to fight microbes. Activation of immune cells such as macrophages are mediated by the sensing of microbial associated molecular patterns (MAMPs) by pattern recognition receptors (PRRs). Leptospires possess numerous lipoproteins, conferring them the ability to potently activate Toll-Like receptor 2 (TLR2) [5, 6]. However, they are stealth pathogens and escape recognition by several other PRRs through unique mechanisms. Our group showed that the leptospiral peptidoglycan escapes NOD receptors recognition by binding to LipL21, one of the numerous leptospiral lipoproteins [7]. We also showed that the leptospiral endoflagella escapes recognition by TLR5 [8]. Furthermore, previous studies on the leptospiral lipopolysaccharide (LPS), a virulence factor [9], revealed that the lipid A (lipidic anchor) possesses structural peculiarities [10] and that it does not activate the human TLR4 whereas it activates the murine TLR4. However, we recently showed that the recognition of the leptospiral LPS in murine macrophages is only partial, because the LPS escapes TLR4 endocytosis and subsequent activation of the TLR4-TRIF pathway [11]. We further demonstrated that the leptospiral LPS activates TLR2 [5], presumably through copurified lipoproteins such as LipL32 (the major leptospiral lipoprotein). On the other hand, our group also showed that leptospires activate the NOD-Like Receptors Pyrin 3 (NLRP) inflammasome and trigger IL1β production [12].

In addition to escaping recognition by several PRRs, leptospires also escape phagocytes' functions. Indeed, although leptospires are mostly extracellular bacteria, we reported that leptospires can be found within macrophages without intracellular replication, and that they evade killing by the classical microbicidal mechanisms such as xenophagy [13]. Autophagy is an essential mechanism allowing nutrient recycling upon starvation. Recently, xenophagy arose as a specific form of autophagy promoting the degradation of intracellular pathogens and dampening inflammation [14, 15]. Xenophagy has been shown essential to clear intracellular bacteria, such as *Mycobacterium tuberculosis*, *Shigella flexneri* and *Salmonella typhimurium* [16-18].

Autophagy induction is a tightly regulated process: upon stress, class III phosphoinositide

3-kinase (PI3K) complex (VSP34, Beclin-1, Atg14) promotes the nucleation of a double membrane

called autophagophore [19]. In addition, ULK1/2 phosphorylates Atg13 to form the initiation complex

[19]. The expansion of the phagophore is then possible *via* the Atg protein conjugation system [20].

Finally, the LC3 conjugation system is required to promote complete autophagosome formation. After

translation, pro-LC3 proteins get cleaved (22 amino acids in C-ter) by protease Atg4 producing LC3-I.

In turn, LC3-I gets conjugated by Atg7 (E1-like ligase) [21] and Atg3 that promote the addition of a

phosphatidylethanolamine (PE) group [22]. The resulting LC3-II localizes on the membrane of the

phagophore promoting its maturation in an autophagosome [23] and its fusion with lysosomes, hence

leading to cargo degradation by lysosomal enzymes such as cathepsins. Although LC3-II of the inner

leaflet will be degraded, LC3-II present in the outer leaflet will be recycled by Atg4, *via* de-lipidation

and return to the LC3-I form [21]. Consequently, LC3-II turnover has been proposed as a good way to

monitor autophagy [24].

Upon infection, several intracellular and extracellular immune signaling pathways can lead to

autophagy induction. Toll-Like Receptors (TLRs) signaling facilitates autophagy through interaction

of MyD88 complexes with Beclin-1 of the nucleation complex [15]. Furthermore, LPS recognition by

TLR4 has been shown to be a potent activator of autophagy, *via* the endosomal TRIF pathway [25].

Other studies also demonstrated that NOD1 and NOD2 induce autophagy [26]. Given that leptospires

escape both NODs and TLR4-TRIF pathways [7, 11], our study aimed to decipher the interplay

between *L. interrogans* and the autophagy machinery of mouse macrophages. Therefore, we assessed

autophagy by monitoring LC3-II, and autophagy adaptors, namely p62. Then, considering the

importance of bacterial LPS in autophagy modulation [25], we investigated the potential role of the

leptospiral LPS in autophagy modulation.

RESULTS

L. interrogans **reduces the quantity of autophagosomes in macrophages**

First, to assess the potential induction of autophagy by leptospires, we infected C57BL/6 mice with

L. interrogans serovar Manilae (one of the most pathogenic serotypes) strain MFLum1 [4]. We

injected intraperitoneally 5×10^7 leptospires/mouse and assessed LC3 levels in peritoneal cells by

Western blot 4 h post-infection. Our results showed that leptospires did not trigger an increase in

LC3-II levels, unlike injection of positive controls for autophagy induction: a mix of muropeptides,

agonists of NOD1 and NOD2, or rapamycin (**Fig 1A**). Then, we infected bone marrow derived

macrophages (BMDMs) *in vitro* with the same strain and monitored LC3-II 24 h post-infection. Here

again, in contrast to autophagy induction triggered by rapamycin, we observed no increase in LC3-II

after *Leptospira* infection (**Fig 1B**). The same observation was made upon stimulation with heat-killed

(HK) leptospires (**Fig 1B**). Furthermore, to address autophagy induction in cells from other vertebrate

hosts, we infected human THP1 stably transfected with CD14 (hereafter THP1-CD14) with

L. interrogans and monitored LC3-II by WB. Our results showed no accumulation of LC3-II after 4 h

infection (**Fig S1**). Overall, these data showed that leptospires did not trigger autophagy in

macrophages.

Second, to address potential modulation of the autophagy flux upon infection, we infected BMDMs

with *L. interrogans* and then treated the cells prior to collection with Bafilomycin A1 (BafA1). BafA1

inhibits the fusion with lysosomes and arrests autophagy before degradation of cargo, including

LC3-II, allowing the accumulation of autophagosomes and consequent analysis of the overall flux.

Our results showed that there was an expected accumulation of LC3-II in BafA1 treated cells, but no

major difference with or without infection, suggesting that leptospires do not alter the autophagy flux

(**Fig 1C**). To ensure that infection with leptospires did not modulate autophagy at earlier time points,

we then performed kinetics analyses of LC3-II. Interestingly, we observed a slight yet robust time-

dependent reduction of LC3-II levels 4 h and 24 h post-infection of BMDMs with *L. interrogans*, as

evidenced by Western blot and their quantifications (**Fig 1D, lower panel**). Many bacteria interfere

actively with autophagy functions via effectors or RNA interference [17, 27-32]. We therefore

analyzed BMDMs after stimulation with HK leptospires and we observed similar LC3-II decrease

over time (**Fig 1D, right panel**), excluding an active mechanism for LC3-II downregulation. To

decipher the origin of LC3-II decrease upon infection, we infected BMDMs and analyzed LC3 mRNA

levels by RT-qPCR 24 h post-infection. Our results showed that LC3 was not transcriptionally

regulated upon infection with leptospires (**Fig 1E**), suggesting a post-transcriptional regulatory

mechanism.

Last, to confirm this paradoxical LC3-II decrease despite no flux alteration upon infection, we used

RAW-mLC3-DiFluo reporter cell line to specifically monitor autophagosomes and autolysosomes

formation. These macrophages are stably transfected with LC3 tagged with tandem GFP-RFP. Both

GFP and RFP fluoresce under basal conditions and when LC3 is embedded in the autophagosome

membrane. However, when autophagosomes become autolysosomes, acidification of the environment

quenches GFP fluorescence, and only RFP fluorescence remains. Using High-Content (HC) automated

microscopy, we analyzed these cells 4 h post-infection with *L. interrogans*. Our results showed that

both autophagosomes (RFP+GFP) and autolysosomes (RFP) were visible in the non-infected cells,

and in the starved cells (EBSS) (**Fig 1F, white arrows**). Interestingly, and consistent with the Western

blots results, we observed a decrease of the overall LC3-RFP fluorescence intensity upon infection as

quantified at 4 h post-infection (**Fig 1G**). Unexpecttedly, we found by quantification of the number of

autophagosomes and autolysosomes that leptospires triggered a decrease in autophagosome number,

without altering the maturation to autolysosomes (**Fig 1G**). Here again, these results confirmed that

leptospires did not block the autophagy flux while reducing the LC3-II autophagosomes structures.

***L. interrogans* induces specific accumulation of autophagy adaptor p62 in puncta-like structures**

The adaptor proteins, such as Sequestosome 1 (SQSTM1, also known as p62), Nuclear Dot Protein 52

(NDP52) and optineurin (OPTN) [17, 33-36] are important players of xenophagy that allow selective

bridging between the bacterial cargo and the autophagosome. The levels of these adaptors in the cells

are regulated by autophagy: they are constitutively degraded upon lysosome / autophagosome fusion.

Considering the LC3-II decrease induced by *L. interrogans*, we then addressed the consequences of

the infection on the autophagy adaptors. We infected BMDMs with *L. interrogans* serovar Manilae

strain MFLum1 and analyzed by Western blots the levels of the different adaptors. Strickingly, we

observed a time-dependent accumulation of p62 upon infection (**Fig 2A**). We confirmed this

phenotype upon infection of BMDMs with two other strains from pathogenic serotypes of

L. interrogans: serovar Copenhageni strain Fiocruz L1-130 and serovar Icterohaemorrhagiae strain

Verdun, and with the saprophytic *Leptospira biflexa* serovar Patoc strain Patoc I (**Fig 2B**). Such

accumulation of the adaptor p62 could have been non-specifically associated with the reduction of

autophagosomes. However, while performing infection of RAW264.7 cells, we observed no difference

in the levels of other adaptors OPTN and NDP52 upon infection (**Fig S2A**). We concluded that

L. interrogans triggered specific accumulation of p62 in murine BMDMs.

To investigate the specific mechanism of p62 accumulation triggered by leptospires, we analyzed the

transcriptional regulation of p62 upon infection by RT-qPCR. In BMDMs infected with

L. interrogans, we observed a significant upregulation of p62 mRNA 24 h post-infection (**Fig 2C**).

This corroborated the idea that p62 accumulation is not only mediated by autophagy downregulation.

Furthermore, we also analyzed infected BMDMs by fluorescence microscopy and we observed that,

4 h post-infection, p62 accumulated in puncta-like structures (**Fig 2D**). The kinetics of these p62

puncta formation were further characterized using automated HC confocal microscopy. Upon infection

of RAW264.7 cells by *L. interrogans*, we quantified the number of p62 puncta *per* cell, as well as the

percentage of cells with p62 puncta. Results showed time-dependent increase of both p62 puncta *per*

cell and percentage of positive cells and showed around 80% of cells positive for p62 with an average

of 6 puncta *per* cell 24 h post-infection (**Fig 2E**). Similar analyses were performed after infection with

L. biflexa, confirming a higher p62 puncta formation with the saprophytic strain, compared to the

pathogenic Manilae (**Fig S2B**). Such findings were confirmed by single cell analysis of RAW264.7

cells infected with *L. interrogans*, that showed an average of 5-10 puncta *per* cell with some

macrophages containing up to 60 p62 puncta (**Fig 2F**). Of note, BafA1 that prevents the acidification

of lysosomes and consequently blocks autophagy flux also led to p62 accumulation, but to a much

lower extent than infection with leptospires (**Fig 2E and 2F**).

Finally, considering the host species-specificity of different escape mechanisms triggered by

leptospires, we also addressed in human cells the effect of infection on p62. Therefore, we infected

THP1-CD14 cells with *L. interrogans* serovar Manilae strain MFLum1. We also observed a specific

p62 accumulation and no NDP52 accumulation after infection with leptospires (**Fig S2C**), suggesting

a conserved mechanism in human and murine macrophages.

Autophagy modulation is mediated by TLR2/TLR4

Considering that leptospires did not reduce LC3-II levels through an active mechanism, we then

addressed the role of PRRs in autophagy modulation. Since leptospires are potent agonists of TLR2

and signal through TLR4-MyD88 in murine cells [11], we analyzed autophagy modulation in

macrophages lacking both TLR2 and TLR4. To do so, we infected wild-type (WT) and TLR2/TLR4

double knock-out (TLR2/4 dko) BMDMs with *L. interrogans*. After 24 h of infection, we assessed by

Western blot the levels of LC3-II and p62. Our results showed that, in contrast to WT cells, there was

no reduction in LC3-II and no accumulation of p62 in TLR2/4 dko BMDMs (**Fig 3A**), suggesting that

autophagy modulation by leptospires is mediated by these two receptors. To further confirm such

results, we also infected WT and MyD88/TRIF dko macrophages, that lack both adaptors for TLRs

signaling. Here again, we observed no accumulation of p62 in MyD88/TRIF BMDMs compared with

the WT BMDMs (**Fig 3B**).

Finally, we investigated the role of TLR2/TLR4 in the specific accumulation of p62. We first analyzed

infected BMDMs by RT-qPCR and observed that the increase in p62 mRNA levels observed in WT

BMDMs was abolished in TLR2/4 dko BMDMs (**Fig 3C**). Furthermore, we performed single cell

analysis of p62 puncta formation in WT, TLR2/4 dko and MyD88/TRIF dko BMDMs upon 24 h

infection by three serotypes of *L. interrogans* (serovar Manilae strain L495, serovar Copenhageni

strain Fiocruz L1-130 and serovar Icterohaemorrhagiae strain Verdun). Our results showed that the

number of p62 puncta *per* cell was greatly reduced in TLR2/TLR4 dko cells and almost completely

abolished in MyD88/TRIF dko cells (**Fig 3D**), which may suggest the involvement of other TLRs than

TLR2 and TLR4s in the leptospiral p62 upregulation.

Given that leptospires modulate autophagy in a TLR2/TLR4 dependent manner, we finally addressed

the potential role of inflammasome in autophagy dampening. Indeed, leptospires have been shown to

activate NLRP3 in a TLR2/TLR4-dependent manner [12], and NLRPs inflammasomes have been

shown to modulate autophagy [37-40]. We therefore stimulated BMDMs with both live and heat-

killed leptospires in presence or absence of glibenclamide, a NLRP3 inhibitor. We first confirmed the

efficiency of the treatment by measuring IL1β production after 24 h, as a readout for inflammasome

activation. Our result showed that IL1β was produced upon infection by leptospires, but reduced in

presence of glibenclamide, confirming that glibenclamide inhibited NLRP3 (**Fig S3A**). We then

analyzed by Western blot the levels of LC3-II, and we observed a reduction of LC3-II upon infection

with either live or heat-killed leptospires in both control and glibenclamide treated cells (**Fig S3B**).

This suggests that the modulation of autophagy by leptospires, although TLR2/TLR4 dependent, is not

mediated by activation of NLRP3 inflammasome.

p62 accumulation is mediated by the leptospiral LPS

With the predominant role of TLR2 and TLR4 in autophagy modulation, and transcriptional regulation

of p62, we then hypothesized that these effects could be induced by the leptospiral LPS. To address

the role of the leptospiral LPS in the p62 accumulation phenotype, we first infected BMDMs with both

heat-killed leptospires (recapitulating MAMPs) and purified leptospiral LPS. Western blot analyses

showed that p62 got accumulated in a time-dependent manner in response to both heat-killed

leptospires and also 24 h post-stimulation with their LPS (**Fig 4A**). We further characterized by HC

microscopy the kinetics of p62 puncta formation in BMDMs after incubation with live, heat-killed

L. interrogans serovar Manilae strain L495 and purified LPS. Results indicated that LPS stimulation

induced the exact same p62 puncta formation than the one observed with bacteria (**Fig 4B**). Next, we

also analyzed BMDMs at the single cell level 24 h post-infection. Here again, HK leptospires or the

leptospiral LPS were sufficient to recapitulate p62 puncta formation in cells (**Fig 4C**). Overall, our

results showed that p62 accumulation was mediated by the leptospiral LPS.

Leptospiral LPS also triggers translocation of NRF2 transcription factor

When autophagy adaptors accumulate in cell (in the absence of functional autophagy or because of

specific upregulation), stress pathways are induced. For instance, p62 accumulation has been shown to

induce translocation of stress-responsive nuclear factor erythroid 2-related factor 2 (NRF2) [41-43].

Subsequently, NRF2 triggers antioxidant and anti-apoptotic programs [41, 42], and promotes p62

upregulation, creating a loop that counteracts stress in autophagy-deficient conditions [41]. We

therefore infected BMDMs with *L. interrogans* serovar Manilae strain L495 and analyzed NRF2 by

immunofluorescence 4 h post-infection. Results showed that NRF2 staining was barely visible in the

non-infected conditions, whereas NRF2 was visible in the nucleus of cells upon infection (**Fig 5A**).

This suggests that leptospires triggered NRF2 translocation in murine macrophages. Interestingly, we

also observed that not all macrophages seem activated to the same extent (**Fig 5A**). Then, we decided

to perform single cell analysis of NRF2 fluorescence intensity in the nucleus of cells infected with

three different serovars of *L. interrogans*. Similarly, upon infection, the levels of NRF2 fluorescence

in the nucleus increased (**Fig 5B**). Next, we calculated the ratio of NRF2 intensities

[nucleus/cytoplasm] and estimated that cells positive for NRF2 had a ratio > 1.4. We then plotted the

percentage of cells with NRF2 ratio superior to 1.4 and observed that up to 60% of cells were positive

after 4 h of infection (**Fig 5C**). Finally, we addressed the role of the leptospiral LPS and performed

similar analyses on BMDMs stimulated for 4 h with either live, heat-killed leptospires or their purified

LPS. All conditions triggered similar NRF2 translocation (**Fig 5D and 5E**), suggesting that the
leptospiral LPS is responsible for NRF2 activation.

p62 and NRF2 constitute a retro-activation loop upon infection with *L. interrogans*

Consequently to p62 accumulation and NRF2 translocation upon infection with *L. interrogans*, we
investigated the potential regulation mechanisms between these two phenotypes. Thus, we performed
small-interfering RNA (siRNA) transfection in BMDMs to knock-down specifically p62 or NRF2. We
then infected the transfected BMDMs and analyzed NRF2 and p62 4 h post-infection by HC
microscopy. First, we analyzed the NRF2 ratio [nucleus/cytoplasm] during p62 knocking-down. Our
results showed that NRF2 ratio was lower in p62 siRNA condition (**Fig 6A**). Absence of p62 was
confirmed and quantified by confocal microscopy (**Fig 6B**). Conversely, we then analyzed p62 puncta
formation upon NRF2 siRNA treatment. Our results showed that p62 puncta formation was altered in
NRF2 siRNA conditions (**Fig 6C**). Similalrly, NRF2 knocked-down was confirmed and quantified
(**Fig 6D**). Overall, these results showed that p62 contributed to NRF2 translocation, and that, in turn,
NRF2 regulated p62 puncta formation, suggesting that p62 and NRF2 are in a retro-activation loop
upon infection by *L. interrogans*.

To further characterize the involvement of NRF2 in p62 retro-activation, we analyzed BMDMs
transfected with NRF2 siRNA and infected with *L. interrogans* by Western blot and RT-qPCR 24 h
post-infection, allowing us to measure both the protein and mRNA levels of p62. Interestingly, we
observed that NRF2 silencing prevented p62 protein accumulation (**Fig 6E**) but also reduced mRNA
upregulation (**Fig 6F**), suggesting that NRF2 contributes transcriptionally to p62 retro-activation.

NRF2 translocation prevents inflammation upon infection

NRF2 is a stress response transcription factor that promotes antioxidant and antiapoptotic programs
upon activation [41, 42]. Therefore, we first monitored the transcription of targets genes involved in
fighting oxidative stress in macrophages in response to infection. We infected BMDMs with
L. interrogans and analyzed by RT-qPCR the levels of gluthatione S-reductase and catalase mRNA at
4 h and 24 h post-infection. Surprisingly, we observed no upregulation of these transcripts (**Fig S4A**
and S4B), suggesting that no antioxidant program is activated upon infection with *L. interrogans*.

Interestingly, NRF2 was shown to also have repressor functions and to dampen inflammation by
inhibiting the transcription of cytokines [44, 45], hence conferring resistance to inflammatory disease
such as sepsis [44]. We therefore investigated the role of NRF2 translocation in inflammation upon
infection by leptospires. We transfected RAW624.7 cells with siRNA targeted against NRF2 and then
infected them with *L. interrogans*. First, we observed a decrease in macrophage viability upon
infection of NRF2-silenced cells (**Fig 7A, left panel**), as expected considering that NRF2 is known to

prevent cell death. Interestingly, this enhanced loss of viability was not associated with lactate

dehydrogenase (LDH) release from the cytosol, illustrating that no membrane damage or cell lysis was

occurring (**Fig 7A, right panel**). We then analyzed the mRNA levels of several cytokines and

enzymes, namely iNOS (nitric oxide inducible synthase), IL6, TNF and IL10. Upon infection, all

mRNA levels were higher in NRF2-silenced cells than in control cells (**Fig 7B**), suggesting that NRF2

does play a role in repressing the expression of inflammatory targets. Finally, to address whether this

transcriptional regulation was strong enough to alter cytokines levels, we analyzed nitric oxide (NO),

IL6, TNF and IL10 production in cell supernatant. Consistent with the mRNA analyses, we observed

higher NO, IL6 and TNF upon infection of NRF2-silenced RAW264.7 (**Fig 7C**). Interestingly, IL10

levels were not modulated in NRF2-silenced cells. All these findings were even more striking when

normalizing the cytokines production with the cell viability (**Fig 7C, grey area**). Overall, NRF2

played a repressor role to dampen NO, TNF and IL6 production in mouse macrophages upon infection

with *L. interrogans*.

DISCUSSION

Our results showed that *L. interrogans* does not induce autophagy in macrophages either

in vivo in mice or *in vitro* in murine and human cells. These findings are consistent with previous

studies from our laboratory showing that leptospires escape NOD1/NOD2 [7] and TLR4-TRIF [11]

activation, since both pathways have been described as autophagy inducers [25, 26]. Furthermore,

infection with leptospires unexpectedly triggered a slight reduction of the number of autophagosomes,

without alteration of the autophagy flux. To this date, we do not know what the functional

consequences of such downregulation are, warranting further investigations. Interestingly, using

heat-killed leptospires we observed similar results, excluding a potential active mechanism. We

conclude that the leptospires behave differently from other pathogenic bacteria such as

Legionella pneumophila, Shigella flexneri or *Listeria monocytogenes* for which autophagy modulation

is mediated by effector proteins, namely RavZ, IcsB and ActA respectively [31, 32]. An

effector-independent mechanism is even more consistent with the fact that leptospires do not possess

traditional secretion systems (SS) such as type-III SS or type-IV SS. Indeed, only the presence of an

incomplete type-II SS has been identified in *Leptospira* species [46]. Consistently, a study revealed

that 59% of the pathogen-specific leptospiral genes have unknown functions, strongly suggesting that

the leptospiral virulence may rely on unique and novel mechanisms [3]. Our results further showed

that the modulation of autophagy was induced by leptospiral LPS signaling through TLR2/TLR4. The

unique leptospiral LPS, the best characterized virulence factor [9], plays an important role in

leptospiral host-pathogen interactions. Indeed, we previously showed that leptospiral LPS avoids

human TLR4 and mouse TLR4-TRIF activation. Recently, through the atypical structures of its

lipid A and O antigen [11, 47, 48], LPS has also been shown to prevent cell death by pyroptosis.

Therefore, it was not surprising to attribute the autophagy modulation phenotype to leptospiral LPS.

Interestingly, the LPS of *L. biflexa* was shown to be a more potent TLR4 agonist [11], potentially

explaining the stronger p62 accumulation observed after infection with the saprophytic strain.

TLRs signaling has classically been associated to autophagy induction, mainly through TLR4-

TRIF [25] but also *via* TLR2, TLR3, TLR4, TLR5 and TLR7 signaling with MyD88 [15]. However,

consistent with our results, some studies also reported that TLRs signaling can also block autophagy.

For instance, LPS signaling downregulates autophagy in non-inflammatory dendritic cells (DCs) [49]

and in microglial cells [50], evidencing the complexity of the autophagy/TLR interplay. In DCs, TLR4

has been associated with mTOR activation, hence reducing the autophagy flux [49]. Whether this

mechanism could also be used by leptospires in macrophages remains to be answered. In addition,

leptospires have been shown to induce NLRP3 in a TLR2/TLR4-dependent manner [12]. Although the

interplay between NLRP3 and autophagy is unclear, several studies reported that NLRP3 can dampen

autophagy [37-40]. Therefore, we hypothesized that *Leptospira*-induced NLRP3 could modulate

autophagy. However, our results suggest that the NLRP3 does not contribute to the autophagy

modulation.

A consequence of autophagy downregulation is the accumulation of a central player of the

autophagy pathway, the p62/SQSTM-1 adaptor [43]. This protein has been shown to accumulate *via*

TLRs signaling [51], most likely through NFκB-mediated transcriptional regulation. Our results

showed that TLR2/TLR4 control the accumulation of p62 upon infection by leptospires. Nevertheless,

we cannot differentiate a direct mechanism (NFκB) from an indirect one (autophagy dysregulation).

Our data showing that p62 is transcriptionally regulated led us to favor the first hypothesis. Moreover,

the levels of p62 upon infection are higher than after BafA1 treatment, suggesting that the

accumulation of p62 is mostly independent of the slight autophagy downregulation induced by

leptospires.

The TLRs-induced accumulation of p62 has been shown to trigger the translocation of the

transcription factor NRF2 [51]. Under physiological conditions, NRF2 is classically targeted for

degradation by the proteasome *via* interaction with its inhibitor Keap1 [43, 52]. The accumulation and

direct binding of p62 to Keap1, leads to its sequestration and subsequent release from NRF2, which

then can translocate in the nucleus [41-43, 52]. Our results confirmed that p62 induces translocation of

NRF2 upon infection with leptospires. Other microbes such as Epstein-Barr virus (EBV) or parasite

Leshmania major have been shown to activate the p62/NRF2 axis [53, 54], hence showing an

important role of NRF2 in response to pathogens. However, to date, the role of NRF2 translocation in

response to pathogens remains unclear.

We then illustrated that the retro-activation loop between p62 and NRF2, described in the

literature [41], is activated upon *L. interrogans* infection. Our results showed that p62 accumulation

triggered translocation of NRF2 and that in turn, NRF2 contributed to the transcriptional regulation of
p62. Interestingly, this loop has few negative regulators, and it is hypothesized that its main regulator
would be autophagy, through degradation of p62 [41]. However, considering the lack of autophagy
induction upon infection with leptospires, it is not surprising that the loop remains active and that p62
puncta did not disappear even at 24 h post-infection. This suggests that leptospires are potent
activators of the p62/NRF2 axis in macrophages.

NRF2 is involved in numerous pathways including the induction of antioxidant programs [41].
We were therefore surprised to observe no modulation in the mRNA levels of antioxidant targets, such
as catalase or gluthatione S-reductase, either at 4 h or 24 h post-infection. Of note, pathogenic
leptospires are equipped to fight against oxidative stress with inducible catalase, peroxidase, and
peroxiredoxin [55]. We therefore hypothesize that leptospires would not benefit from induction of
antioxidant program in the host cells. Interestingly, active repression of these NRF2-dependent
antioxidant targets was previously shown upon infection with live *Leshmania major* parasite [54, 56]
Whether leptospires could also actively prevent upregulation of NRF2 antioxidant targets remains to
be addressed.

Alternatively, NRF2 has been shown to have transcription inhibition properties. Indeed, in
BMDMs, NRF2 was described as an inhibitor of transcription for pro-inflammatory targets such as
IL6, IL1β and IL12 [45, 57]. Our results showed that this was also the case upon infection with
L. interrogans. Accordingly, we did not observe dampening of the anti-inflammatory IL10 upon
infection. NRF2 inhibits polymerase III recruitment and hence prevents transcription of cytokines in
response to stimulation [45]. Consequently, NRF2 was shown to downregulate neutrophils activation
and migration [58]. Interestingly, neutrophils are barely observed in tissues of mice and humans
infected with leptospires, although they are found in numbers in blood [59, 60]. Whether NRF2
translocation in specific cell type could have a role in this phenotype remains to be studied. NRF2 was
also associated to inflammation reduction *via* indirect mechanism relying on STING and type I IFN
downregulation [61], which remains to be studied in the case of leptospiral infection.

Finally, we showed that accumulation of p62 observed in murine macrophages was conserved
in human cells. Although we could not address NRF2 translocation in human cells due to lack of
specific antibodies and tools, we may speculate that inflammation dampening could therefore be
conserved in human macrophages. Although the functional consequences of this mechanism remain to
be addressed, we believe it could play a key role in the discretion of leptospires in hosts, lack of
immune cell recruitment and potentially contribute to the establishment of chronic infections.

MATERIALS AND METHODS

Leptospira interrogans **cultures.** *Leptospira interrogans* used in this work (serovar Manilae strain L495 and bioluminescent derivative MFLum1 [4], serovar Copenhageni strain Fiocruz L1-130, serovar Icterohaemorragiae strain Verdun) and *Leptospira biflexa* (serovar Patoc strain Patoc I) were grown in Ellinghausen-McCullough-Johnson-Harris (EMJH) medium at 28°C without agitation and diluted weekly, or twice a week for *L. biflexa*, to obtain exponential phase growth culture at the time of experiments. For infection, leptospires cultures were centrifuged (3250 g, 25 min), resuspended in endotoxin-free PBS (Lonza), enumerated using a Petroff-Hauser chamber and bacterial concentration was adjusted accordingly. To inactivate bacteria (thereafter called "heat-killed") leptospires were heated at 56 °C for 30 min with mild agitation. For fluorescent labelling, 10 mL of exponential culture was centrifuged (3250 g, 25 min) and the pellet was resuspended in the same volume of PBS (Lonza) with addition of 10 µM CFSE (Sigma). The bacterial suspension was incubated at room temperature for 30 min protected from light and washed 1 time with 20 mL of PBS before counting and infection.

Mice experiments. Adults male and female mice of the following genotypes in a C57/BL6 background were used: WT (Janvier Labs, Le Genest-Saint-Isle, France); TLR2$^{-/-}$TLR4$^{-/-}$ and MyD88$^{-/-}$/TRIF^{lps2} (both from Institut Pasteur). For peritoneal cavity analyses, mice were injected *via* intraperitoneal route with 5×10^7 leptospires/mouse, 30 µg/mouse of rapamycin (Sigma) or 30 µg/mouse of NOD1 agonist (FK156) [62] and NOD2 agonist (MDP, Invivogen) in PBS. Next, 4 h post-infection mice were euthanized, and peritoneal content was recovered as previously described [63]. Briefly, skin was peeled and clamped, followed by the injection of 2 mL of RPMI media. Next an incision was made on the pre-clamped skin to recover the peritoneal content. Peritoneal cells were then used for sample preparation and Western blot analyses, as described below.

Ethic statement. All protocols were undertaken in compliance with EU Directive 2010/63 EU and the French regulation on the protection of laboratory animals issued on February 1st, 2013. They are part of projects approved by the Institut Pasteur ethics committee for animal experimentation (Comité d'Ethique en Expérimentation Animale CETEA registered under #89) under # HA0036 for the bone marrow cells, and #2014-0049 for the leptospiral infections, which was authorized by the French Ministry of Research, the French Competent Authority under #8562.

Cell culture. RAW264.7 murine macrophage-like cell line was cultured in complete RPMI containing glutamine (Lonza) and supplemented with 10 % v/v heat-inactivated fetal calf serum (HI-FCS, Gibco), 1 mM non-essential amino acids (MEM, Gibco) and 1 mM sodium pyruvate (NaPy, Gibco). RAW-mLC3-DiFluo (Invivogen) autophagy reporter cells stably transfected with LC3-GFP-RFD were cultivated in the same conditions, with the addition of 100 µg/mL zeocin (Invivogen) for efficient selective pressure. All the cells were seeded in 96-, 24-, 12-well plates (TPP) the day before

infection at a concentration of $0,3 \times 10^6$ cells/mL in 200 µL, 1 mL or 2 mL respectively, in
antibiotic-free medium.

Human THP-1 monocyte-like cell line, stably transfected with CD14 (THP1-CD14), was culture in
complete RPMI, and the cells were differentiated for 48 hours with 20 ng/mL of phorbol 12-myristate
13-acetate (PMA, Invivogen) at 1×10^6 cells/mL. The cells were then washed 3 times with PBS and
allowed 24 hours of resting before infection.

BMDMs were obtained from different genotypes of C57BL/6 background mice. After euthanasia by
cervical dislocation, the head of each bone was removed, and 2 mL of medium were passed through
the bones using a 22 G needle to flush out the bone marrow. Bone marrow cells were centrifuged
(300 g, 7 min) and treated with Red Blood Cell Lysis Buffer (Sigma-Aldrich) for 10 minutes, followed
by another centrifugation. The cells were then enumerated, and 5×10^6 cells were seeded in 100 cm^2
cell culture dishes in 12 mL of complete RPMI supplemented with 10% v/v L929 cell supernatant and
1x PenStrep (Gibco). Cells were incubated at 37 °C, 5 % CO_2. At day 3, 3 mL of the same medium
was added. At day 7, the cells were collected by scraping in Cell Dissociation buffer (Gibco),
centrifuged, enumerated, and seeded for infection at 1×10^6 cells/mL.

In addition to the infections with *L. interrogans*, other treatments were performed. Starvation was
induced in minimum EBSS media (Gibco) after 2 washes to remove complete cell culture media.
Autophagy was induced with 500 mM rapamycin (Sigma). For blockage of autophagy flux, 100 nM
bafilomycin A1 (BafA1, Sigma-Aldrich) was added to the infected cells 4 h before cell collection and
analyses. The inflammasome inhibitor glibenclamide (Thermo Fisher) was added at 25 µM, 30 min
before infection. Finally, stimulations by leptospiral LPS were performed by incubating the cells with
1 µg/mL of LPS of *L. interrogans* (serovar Manilae strain L495, serovar Copenhageni strain Fiocruz
L1-130 or serovar Icterohaemorragiae strain Verdun) extracted from the phenol phase of the hot
water/phenol extraction protocol, as we recently reviewed[64].

Immunofluorescence and High Content (HC) microscopy. For classical IF analyses, cells were
seeded in 24-well plate containing 1 cover glass per well (18 mm diameter, # 1.5 thickness, Electron
Microscopy Science). After infection, cells were fixed in 4 % v/v *para*-formaldehyde for 10 min,
followed by three washes with PBS. Cells were blocked 1 h in PBS with 2.5% w/v bovine serum
albumin (BSA) and 2.5 µg/mL anti-CD16-CD32 (FcBlock, Thermo Fisher). Cells were stained after
blocking with primary antibodies (**Table 1**) overnight at 4 °C in PBS 1% w/v BSA, 0.05% w/v
saponin. Cells were then washed three times with PBS and were incubated with fluorescent secondary
antibodies (**Table 1**) and 1µg/mL DAPI in PBS 1% w/v BSA, 0.05% w/v saponin for 1 hour at room
temperature. For NRF2 nuclear staining, permeabilization was initially performed by 10 min
incubation in PBS 1% w/v BSA, 0.5% v/v Triton X-100 prior to blocking, and saponin was not used.
In all cases, cells were washed three times with PBS and image acquisition was performed on Leica

SP5 confocal microscope (63x - 1.4 NA, oil immersion). Default settings were used and both UV and
argon lasers were used at power between 5-40 %.

For HC microscopy analyses, cells were seeded in dark, transparent bottom 96-well plates
(Greiner, µClear) at $0.1x10^6$ cells/mL. Staining protocols are identical to the ones described above,
except for RAW-mLC3-DiFluo cells that were fixed to 10 min in cold ethanol/acetone (1:1) buffer.
All cells were then kept in PBS for imaging by Opera Phenix HCS (Perkin Elmer) using confocal
settings with a 63X water-immersion objective (NA 1.15). Light source was kept at 20 to 50% power
and exposure time was set for each experiment to obtain intensities around 1000-5000. Automated
acquisition was launched to acquire a minimum of 500-1000 cells/well in technical triplicates for each
experiment. Automated image analysis workflow and puncta quantification was performed using
Columbus image data storage and analysis system (PerkinElmer), with the following steps: *(i)* import
data; *(ii)* find nuclei with DAPI signal; *(iii)* find cytoplasm with the background of the target protein
signal (NRF2 or p62); *(iv)* find spots using method C if relevant (for p62); *(v)* formulate classical
outputs: number of p62 puncta, NRF2 intensity in nucleus/cytoplasm; *(vi)* formulate calculated
outputs: %p62 positive cells, NRF2 ratio, %NRF2 positive cells*; (vii)* save script and run batch
analysis and *(viii)* export data.

Table 1. Antibodies list for immunofluorescence and Western blot analyses

Application*	Target	Clonality**	Conjugated	Host	Dilution	Reference	Supplier
IF	**p62**	mAb	Purified	Rabbit	1:500	ab109012	Abcam
	Rabbit IgG	pAb	AlexaFluor 488	Goat	1:1.000	A-11034	ThermoFisher
IF	**Nrf2**	mAb	Purified	Rat	1:250	#14596	CST
	Rat IgG	pAb	AlexaFluor 647	Goat	1:1.000	A-21247	ThermoFisher
WB	**LC3**	pAb	Purified	Rabbit	1:1.000	L7543	Sigma
	p62	mAb	Purified	Mouse	1:1.000	ab109012	Abcam
	OPTN	pAb	Purified	Rabbit	1:1.000	ab23666	Abcam
	NDP52	pAb	Purified	Rabbit	1:1.000	GTX115378	GeneTex
	Rabbit IgG	pAb	HRP	Goat	1:10.000	7074S	CST

* IF: immunofluorescence, WB: Western blot, ** mAb: monoclonal antibody, pAb: polyclonal antibody

SDS-PAGE and Western blot. For proteins collection and Western blot analyses, cell supernatants
were removed, and 1 mL of Cell Dissociation buffer (Gibco) was added and incubated for 5 minutes,
followed by scrapping, collection, and centrifugation (400 g, 10 min, 4 °C). The pellets were
resuspended in 50 µL of ice-cold RIPA lysis buffer supplemented with 1x complete Mini, EDTA-free
proteases inhibition cocktail (Roche). Cells were lysed on ice for 20 min followed by a centrifugation
step (12000 g, 40 min, 4 °C) to remove the non-soluble material. Proteins were dosed using the
Bradford assay and the samples were denatured in Laemmli buffer (99 °C, 10 min). SDS-PAGE was

done using 4-15% gradient acrylamide Stain-free gels (BioRad), loaded with 5-10 µg of protein extract *per* lane and gels were run at 100 V in Tris-Glycine-SDS buffer (BioRad). Proteins were visualized using stain free reagent in ChemiDoc (BioRad) with a 5 min exposition time and transferred onto a 0.22 µm PVDF membrane (BioRad) using Mixed MW fast transfer of 1 miniGel (BioRad). The membrane was blocked with 5% BSA in TBS with 0,05% v/v Tween 20 (TBS-T) for 60 minutes at room temperature with mild agitation. The membrane was probed with either anti-LC3 or anti-p62, anti-NDP52, anti-OPTN (**Table.1**) in 5% BSA in TBS-T overnight at 4 °C. The membranes were then washed in TBS-T 3 times for 15 minutes followed by incubation with secondary anti-rabbit IgG HRP-linked (**Table 1**) in 5% BSA in TBS-T for 60 minutes protected from light at room temperature, followed by 3 washes in TBS-T for 15 minutes. Blots were revealed using the Clarity reagent (BioRad) with automatic exposure time. Western blot quantifications were performed using Image Lab software (BioRad), and LC3-II levels were quantified after normalization by the strain free (total amounts of proteins/lane).

Small interfering RNA transfection. For siRNA transfection, cells were seeded at confluence of $0,1x10^6$ cells/mL in 24-well plate (TPP) the day before transfection. In each well, transfection was performed using 4,5 µL of Lipofectamine siRNA Max reagent (ThermoFisher) in 650 µL of OptiMEM media (Gibco) and according to the manufacturer's instructions. The final concentration of commercial predesigned siRNA targeting *p62* and *nrf2* (FlexiTube, Qiagen) was 80 nM and the corresponding scramble siRNA (all start negative, Qiagen) was performed as control. The siRNA preparations in lipofectamine were incubated at room temperature for 25 min before delicate transfection of the cells. After 12 hours, 350 µL of complete RPMI was added on the cells. The infection was performed 24 hours post-transfection in complete RPMI and the reduction in p62 and NRF2 levels was checked by HCS microscopy.

ELISA and Griess reaction. Fresh cell culture supernatants were used 24 h post-infection for dosage of nitric oxide (NO) by the Griess reaction. On the other hand, cytokine dosage was performed on frozen supernatants by ELISA using DuoSet kits (R&D Systems), according to the manufacturer's instructions.

Viability and LDH release assays. LDH release was measured using the CyQuant LDH Cytotoxicity Assay kit (fresh supernatant) (Invitrogen), according to the manufacturers' instructions. On the other hand, viability was measured using MTT assay: after infection, cells were incubated for 2 h in 1 mg/mL MTT solution (Sigma) in complete RPMI. MTT crystals were then dissolved in HCl 1 M, isopropanol (1:24) and absorbance was read at 595 nm.

RT-qPCR. RNAs were extracted from frozen cells using RNeasy Mini Kit (Qiagen) according to the manufacturer's instructions. cDNAs were then obtained by retro-transcription (RT) using SuperScript II RT (Invitrogen). The equivalent of 20ng of cDNA were used for RT-qPCR with

518   specific primers and probes (**Table 2**). qPCR was performed using Taqman Universal MasterMix

519   (Applied Biosystems) on a StepOne Plus Real Time PCR System (Applied Biosystems) with the

520   standard protocol. Fold changes were calculated with the $2^{-\Delta\Delta Ct}$ method, using hypoxanthine guanine

521   phosphoribosyl transferase (HPRT) as internal control.

523 **Table 2.** Primers and probe list for RT-qPCR analyses

Target	Forward primer sequence	Reverse primer sequence	Probe sequence
p62	5'-ATGTGGAACATGGAGGGAAGAG-3'	5'-TTCTGTGCCTGTGCTGGAACT-3'	5'-CCGCCTGACACCCACTACCCCA-3'
lc3	5'-GCCCCACCCCTGAAAGG-3'	5'-TGCAGAGGAAATGACCACAGAT-3'	5'-TGGCTCCCCTTGTCCTGACTCGG-3'
iNOS	5'-CATGACTGCTAATGTCAGAG-3'	5'-TCCTAGTCCATCCGGATAAA-3'	5'-GAAATCTGAGTTTGGCCTGAGG-3'
tnf	5'-CATCTTCTCAAAATTCGAGTGACAA-3'	5'-TGGGAGTAGACAAGGTACAACCC-3'	5'-CACGTCGTAGCAAACCACCAAGTGGA-3'
il6	5'-CTCCAGCTTATCTGTTAGGA-3'	5'-CTTCAACCAAGAGGTAAAAGA-3'	5'-AAATTGGGGTAGGAAGGACTATTTTATG-3'
il10	5'-GGCGCTGTCATCGATTTCTC-3'	5'-GACACCTTGGTCTTGGAGCTTATTAA-3'	5'-AAAATAAGAGCAAGGCAGTGGAGCAGGTG-3'
hprt	5'-CTGGTGAAAAGGACCTCTCGA-3'	5'-TGCCCTTGACTATAATGAGTACTTCA-3'	5'-TGTTGGATACAGGCCAGACTTTGTTGGAT-3'
cat	Pre-designed: Mm00437992_m1 (ThermoFisher)		
gsr	Pre-designed: Mm01197925_m1 (ThermoFisher)		
hprt	Pre-designed: Mm03024075_m1 (ThermoFisher)		

525 **Statistical analyses.** All statistical analyses were performed using Student's *t*-test with
526 corresponding *p* values: * for $p < 0.05$; ** for $p < 0.01$ and *** for $p < 0.001$.

ACKNOWLEDGMENTS

528 We acknowledge Nathalie Aulner and Anne Danckaert (Institut Pasteur) as well as the Photonic
529 BioImaging Unit of Technology and Service, a member of the France-BioImaging infrastructure, for
530 support in conducting this study.

FUNDING

This work was funded by Institut Pasteur grant PTR2017-66 to CW and by Agence Nationale de la
Recherche (ANR) grant ANR-10-LABX-62-IBEID to IGB. DB was funded by Université Paris Cité
(formerly Université Paris Diderot) through Doctoral school FIRE (ED FIRE474). IS has been part of
the Pasteur-Paris University (PPU) International PhD program. This program has received funding
from the Institut Carnot Pasteur Microbes & Santé, and the European Union's Horizon 2020 research
and innovation program under the Marie Sklodowska-Curie grant agreement no. 665807. IS has
additionally benefited from a scholarship "Fin de thèse de science" number FDT201805005258
granted by "Fondation pour la recherche médicale (FRM)". France-BioImaging infrastructure network
is supported by the French National Research Agency (ANR-10–INSB–04, Investments for the
future). The use of the Opera Phenix system in the PBI core facility has been made possible by the
kind financial support of the Institut Pasteur (Paris) and the Région Ile-de-France (program
DIM1Health). The authors declare no conflict of interest.

AUTHORS CONTRIBUTIONS

Conception, administration, and supervision of the project: CW.
Investigation: DB, IS, and CW.
Methodology: IS, DB and PE.
Data analysis: DB, IS, PE and CW.
Visualization: DB.
Validation: IS, DB and CW.
Resources: FVP.
Funding acquisition: CW and IGB.
DB and IS wrote the original draft, under CW's supervision.
All authors contributed to the review of the manuscript and approved it for submission.

REFERENCES

1. Costa F, Hagan JE, Calcagno J, Kane M, Torgerson P, Martinez-Silveira MS, et al. Global Morbidity and Mortality of Leptospirosis: A Systematic Review. PLoS Negl Trop Dis. 2015;9(9):e0003898. Epub 20150917. doi: 10.1371/journal.pntd.0003898. PubMed PMID: 26379143; PubMed Central PMCID: PMCPMC4574773.

2. Adler B. History of leptospirosis and leptospira. Curr Top Microbiol Immunol. 2015;387:1-9. doi: 10.1007/978-3-662-45059-8_1. PubMed PMID: 25388129.

3. Ko AI, Goarant C, Picardeau M. Leptospira: the dawn of the molecular genetics era for an emerging zoonotic pathogen. Nat Rev Microbiol. 2009;7(10):736-47. doi: 10.1038/nrmicro2208. PubMed PMID: 19756012; PubMed Central PMCID: PMCPMC3384523.

4. Ratet G, Veyrier FJ, Fanton d'Andon M, Kammerscheit X, Nicola MA, Picardeau M, et al. Live imaging of bioluminescent leptospira interrogans in mice reveals renal colonization as a stealth escape from the blood defenses and antibiotics. PLoS Negl Trop Dis. 2014;8(12):e3359. Epub 20141204. doi: 10.1371/journal.pntd.0003359. PubMed PMID: 25474719; PubMed Central PMCID: PMCPMC4256284.

5. Werts C, Tapping RI, Mathison JC, Chuang TH, Kravchenko V, Saint Girons I, et al. Leptospiral lipopolysaccharide activates cells through a TLR2-dependent mechanism. Nat Immunol. 2001;2(4):346-52. doi: 10.1038/86354. PubMed PMID: 11276206.

6. Hsu SH, Hung CC, Chang MY, Ko YC, Yang HY, Hsu HH, et al. Active Components of Leptospira Outer Membrane Protein LipL32 to Toll-Like Receptor 2. Sci Rep. 2017;7(1):8363. Epub 20170821. doi: 10.1038/s41598-017-08743-y. PubMed PMID: 28827637; PubMed Central PMCID: PMCPMC5566480.

7. Ratet G, Santecchia I, Fanton d'Andon M, Vernel-Pauillac F, Wheeler R, Lenormand P, et al. LipL21 lipoprotein binding to peptidoglycan enables Leptospira interrogans to escape NOD1 and NOD2 recognition. PLoS Pathog. 2017;13(12):e1006725. Epub 20171206. doi: 10.1371/journal.ppat.1006725. PubMed PMID: 29211798; PubMed Central PMCID: PMCPMC5764436.

8. Holzapfel M, Bonhomme D, Cagliero J, Vernel-Pauillac F, Fanton d'Andon M, Bortolussi S, et al. Escape of TLR5 Recognition by Leptospira spp.: A Rationale for Atypical Endoflagella. Front Immunol. 2020;11:2007. Epub 20200811. doi: 10.3389/fimmu.2020.02007. PubMed PMID: 32849665; PubMed Central PMCID: PMCPMC7431986.

9. Murray GL, Srikram A, Henry R, Hartskeerl RA, Sermswan RW, Adler B. Mutations affecting Leptospira interrogans lipopolysaccharide attenuate virulence. Mol Microbiol. 2010;78(3):701-9. Epub 20100914. doi: 10.1111/j.1365-2958.2010.07360.x. PubMed PMID: 20807198.

10. Que-Gewirth NL, Ribeiro AA, Kalb SR, Cotter RJ, Bulach DM, Adler B, et al. A methylated phosphate group and four amide-linked acyl chains in leptospira interrogans lipid A. The membrane anchor of an unusual lipopolysaccharide that activates TLR2. J Biol Chem. 2004;279(24):25420-9. Epub 20040324. doi: 10.1074/jbc.M400598200. PubMed PMID: 15044492; PubMed Central PMCID: PMCPMC2556802.

11. Bonhomme D, Santecchia I, Vernel-Pauillac F, Caroff M, Germon P, Murray G, et al. Leptospiral LPS
escapes mouse TLR4 internalization and TRIFassociated antimicrobial responses through O antigen and
associated   lipoproteins.   PLoS   Pathog.   2020;16(8):e1008639.   Epub   20200813.   doi:
10.1371/journal.ppat.1008639. PubMed PMID: 32790743; PubMed Central PMCID: PMCPMC7447051.

12. Lacroix-Lamande S, d'Andon MF, Michel E, Ratet G, Philpott DJ, Girardin SE, et al. Downregulation of
the Na/K-ATPase pump by leptospiral glycolipoprotein activates the NLRP3 inflammasome. J Immunol.
2012;188(6):2805-14. Epub 20120208. doi: 10.4049/jimmunol.1101987. PubMed PMID: 22323544.

13. Santecchia I, Bonhomme D, Papadopoulos S, Escoll P, Giraud-Gatineau A, Moya-Nilges M, et al. Alive
Pathogenic and Saprophytic Leptospires Enter and Exit Human and Mouse Macrophages With No
Intracellular   Replication.   Front   Cell   Infect   Microbiol   2022;12:936931.   doi: doi:
10.3389/fcimb.2022.936931.

14. Sorbara MT, Ellison LK, Ramjeet M, Travassos LH, Jones NL, Girardin SE, et al. The protein ATG16L1
suppresses inflammatory cytokines induced by the intracellular sensors Nod1 and Nod2 in an autophagy-
independent manner. Immunity. 2013;39(5):858-73. doi: 10.1016/j.immuni.2013.10.013. PubMed PMID:
24238340.

15. Wang Z, Li C. Xenophagy in innate immunity: A battle between host and pathogen. Dev Comp Immunol.
2020;109:103693. Epub 20200331. doi: 10.1016/j.dci.2020.103693. PubMed PMID: 32243873.

16. Travassos LH, Carneiro LA, Girardin S, Philpott DJ. Nod proteins link bacterial sensing and autophagy.
Autophagy. 2010;6(3):409-11. Epub 20100426. doi: 10.4161/auto.6.3.11305. PubMed PMID: 20200479.

17. Birmingham CL, Smith AC, Bakowski MA, Yoshimori T, Brumell JH. Autophagy controls Salmonella
infection in response to damage to the Salmonella-containing vacuole. J Biol Chem. 2006;281(16):11374-
83. Epub 20060222. doi: 10.1074/jbc.M509157200. PubMed PMID: 16495224.

18. Gutierrez MG, Master SS, Singh SB, Taylor GA, Colombo MI, Deretic V. Autophagy is a defense
mechanism inhibiting BCG and Mycobacterium tuberculosis survival in infected macrophages. Cell.
2004;119(6):753-66. doi: 10.1016/j.cell.2004.11.038. PubMed PMID: 15607973.

19. He C, Klionsky DJ. Regulation mechanisms and signaling pathways of autophagy. Annu Rev Genet.
2009;43:67-93. doi: 10.1146/annurev-genet-102808-114910. PubMed PMID: 19653858; PubMed Central
PMCID: PMCPMC2831538.

20. Mizushima N, Kuma A, Kobayashi Y, Yamamoto A, Matsubae M, Takao T, et al. Mouse Apg16L, a novel
WD-repeat protein, targets to the autophagic isolation membrane with the Apg12-Apg5 conjugate. J Cell
Sci. 2003;116(Pt 9):1679-88. doi: 10.1242/jcs.00381. PubMed PMID: 12665549.

21. Tanida I, Ueno T, Kominami E. LC3 conjugation system in mammalian autophagy. Int J Biochem Cell
Biol. 2004;36(12):2503-18. doi: 10.1016/j.biocel.2004.05.009. PubMed PMID: 15325588; PubMed Central
PMCID: PMCPMC7129593.

22. Hanada T, Noda NN, Satomi Y, Ichimura Y, Fujioka Y, Takao T, et al. The Atg12-Atg5 conjugate has a
novel E3-like activity for protein lipidation in autophagy. J Biol Chem. 2007;282(52):37298-302. Epub
20071106. doi: 10.1074/jbc.C700195200. PubMed PMID: 17986448.

23. Kabeya Y, Mizushima N, Ueno T, Yamamoto A, Kirisako T, Noda T, et al. LC3, a mammalian homologue
of yeast Apg8p, is localized in autophagosome membranes after processing. EMBO J. 2000;19(21):5720-8.
doi: 10.1093/emboj/19.21.5720. PubMed PMID: 11060023; PubMed Central PMCID: PMCPMC305793.

24. Klionsky DJ, Abdel-Aziz AK, Abdelfatah S, Abdellatif M, Abdoli A, Abel S, et al. Guidelines for the use
and interpretation of assays for monitoring autophagy (4th edition)(1). Autophagy. 2021;17(1):1-382. Epub
20210208. doi: 10.1080/15548627.2020.1797280. PubMed PMID: 33634751; PubMed Central PMCID:
PMCPMC7996087.

25. Xu Y, Jagannath C, Liu XD, Sharafkhaneh A, Kolodziejska KE, Eissa NT. Toll-like receptor 4 is a sensor
for autophagy associated with innate immunity. Immunity. 2007;27(1):135-44. Epub 20070719. doi:
10.1016/j.immuni.2007.05.022. PubMed PMID: 17658277; PubMed Central PMCID: PMCPMC2680670.

26. Travassos LH, Carneiro LA, Ramjeet M, Hussey S, Kim YG, Magalhaes JG, et al. Nod1 and Nod2 direct
autophagy by recruiting ATG16L1 to the plasma membrane at the site of bacterial entry. Nat Immunol.
2010;11(1):55-62. Epub 20091108. doi: 10.1038/ni.1823. PubMed PMID: 19898471.

27. Birmingham CL, Canadien V, Gouin E, Troy EB, Yoshimori T, Cossart P, et al. Listeria monocytogenes
evades killing by autophagy during colonization of host cells. Autophagy. 2007;3(5):442-51. Epub
20070518. doi: 10.4161/auto.4450. PubMed PMID: 17568179.

28. Case ED, Chong A, Wehrly TD, Hansen B, Child R, Hwang S, et al. The Francisella O-antigen mediates
survival in the macrophage cytosol via autophagy avoidance. Cell Microbiol. 2014;16(6):862-77. Epub
20131216. doi: 10.1111/cmi.12246. PubMed PMID: 24286610; PubMed Central PMCID:
PMCPMC4028363.

29. Kreibich S, Emmenlauer M, Fredlund J, Ramo P, Munz C, Dehio C, et al. Autophagy Proteins Promote
Repair of Endosomal Membranes Damaged by the Salmonella Type Three Secretion System 1. Cell Host
Microbe. 2015;18(5):527-37. doi: 10.1016/j.chom.2015.10.015. PubMed PMID: 26567507.

30. Chen Z, Wang T, Liu Z, Zhang G, Wang J, Feng S, et al. Inhibition of Autophagy by MiR-30A Induced by
Mycobacteria tuberculosis as a Possible Mechanism of Immune Escape in Human Macrophages. Jpn J
Infect Dis. 2015;68(5):420-4. Epub 20150410. doi: 10.7883/yoken.JJID.2014.466. PubMed PMID:
25866116.

31. Choy A, Dancourt J, Mugo B, O'Connor TJ, Isberg RR, Melia TJ, et al. The Legionella effector RavZ
inhibits host autophagy through irreversible Atg8 deconjugation. Science. 2012;338(6110):1072-6. Epub
20121025. doi: 10.1126/science.1227026. PubMed PMID: 23112293; PubMed Central PMCID:
PMCPMC3682818.

32. Ogawa M, Yoshimori T, Suzuki T, Sagara H, Mizushima N, Sasakawa C. Escape of intracellular Shigella
from autophagy. Science. 2005;307(5710):727-31. Epub 20041202. doi: 10.1126/science.1106036.
PubMed PMID: 15576571.

33. Manzanillo PS, Ayres JS, Watson RO, Collins AC, Souza G, Rae CS, et al. The ubiquitin ligase parkin
mediates resistance to intracellular pathogens. Nature. 2013;501(7468):512-6. Epub 20130904. doi:
10.1038/nature12566. PubMed PMID: 24005326; PubMed Central PMCID: PMCPMC3886920.

34. Mostowy S, Sancho-Shimizu V, Hamon MA, Simeone R, Brosch R, Johansen T, et al. p62 and NDP52
proteins target intracytosolic Shigella and Listeria to different autophagy pathways. J Biol Chem.
2011;286(30):26987-95. Epub 20110606. doi: 10.1074/jbc.M111.223610. PubMed PMID: 21646350;
PubMed Central PMCID: PMCPMC3143657.

35. Watson RO, Bell SL, MacDuff DA, Kimmey JM, Diner EJ, Olivas J, et al. The Cytosolic Sensor cGAS
Detects Mycobacterium tuberculosis DNA to Induce Type I Interferons and Activate Autophagy. Cell Host
Microbe. 2015;17(6):811-9. Epub 20150602. doi: 10.1016/j.chom.2015.05.004. PubMed PMID: 26048136;
PubMed Central PMCID: PMCPMC4466081.

36. Zheng YT, Shahnazari S, Brech A, Lamark T, Johansen T, Brumell JH. The adaptor protein p62/SQSTM1
targets invading bacteria to the autophagy pathway. J Immunol. 2009;183(9):5909-16. Epub 20091007. doi:
10.4049/jimmunol.0900441. PubMed PMID: 19812211.

37. Wlodarska M, Thaiss CA, Nowarski R, Henao-Mejia J, Zhang JP, Brown EM, et al. NLRP6 inflammasome
orchestrates the colonic host-microbial interface by regulating goblet cell mucus secretion. Cell.
2014;156(5):1045-59. doi: 10.1016/j.cell.2014.01.026. PubMed PMID: 24581500; PubMed Central
PMCID: PMCPMC4017640.

38. Jounai N, Kobiyama K, Shiina M, Ogata K, Ishii KJ, Takeshita F. NLRP4 negatively regulates autophagic
processes through an association with beclin1. J Immunol. 2011;186(3):1646-55. Epub 20110105. doi:
10.4049/jimmunol.1001654. PubMed PMID: 21209283.

39. Deng Q, Wang Y, Zhang Y, Li M, Li D, Huang X, et al. Pseudomonas aeruginosa Triggers Macrophage
Autophagy To Escape Intracellular Killing by Activation of the NLRP3 Inflammasome. Infect Immun.
2016;84(1):56-66. Epub 20151014. doi: 10.1128/IAI.00945-15. PubMed PMID: 26467446; PubMed
Central PMCID: PMCPMC4694000.

40. Lai M, Yao H, Shah SZA, Wu W, Wang D, Zhao Y, et al. The NLRP3-Caspase 1 Inflammasome
Negatively Regulates Autophagy via TLR4-TRIF in Prion Peptide-Infected Microglia. Front Aging
Neurosci. 2018;10:116. Epub 20180418. doi: 10.3389/fnagi.2018.00116. PubMed PMID: 29720937;
PubMed Central PMCID: PMCPMC5915529.

41. Jain A, Lamark T, Sjottem E, Larsen KB, Awuh JA, Overvatn A, et al. p62/SQSTM1 is a target gene for
transcription factor NRF2 and creates a positive feedback loop by inducing antioxidant response element-

driven gene transcription. J Biol Chem. 2010;285(29):22576-91. Epub 20100507. doi:
10.1074/jbc.M110.118976. PubMed PMID: 20452972; PubMed Central PMCID: PMCPMC2903417.

42. Komatsu M, Kurokawa H, Waguri S, Taguchi K, Kobayashi A, Ichimura Y, et al. The selective autophagy
substrate p62 activates the stress responsive transcription factor Nrf2 through inactivation of Keap1. Nat
Cell Biol. 2010;12(3):213-23. Epub 20100221. doi: 10.1038/ncb2021. PubMed PMID: 20173742.

43. Lau A, Wang XJ, Zhao F, Villeneuve NF, Wu T, Jiang T, et al. A noncanonical mechanism of Nrf2
activation by autophagy deficiency: direct interaction between Keap1 and p62. Mol Cell Biol.
2010;30(13):3275-85. Epub 20100426. doi: 10.1128/MCB.00248-10. PubMed PMID: 20421418; PubMed
Central PMCID: PMCPMC2897585.

44. Kong X, Thimmulappa R, Craciun F, Harvey C, Singh A, Kombairaju P, et al. Enhancing Nrf2 pathway by
disruption of Keap1 in myeloid leukocytes protects against sepsis. Am J Respir Crit Care Med.
2011;184(8):928-38. Epub 20110728. doi: 10.1164/rccm.201102-0271OC. PubMed PMID: 21799073;
PubMed Central PMCID: PMCPMC3208662.

45. Kobayashi EH, Suzuki T, Funayama R, Nagashima T, Hayashi M, Sekine H, et al. Nrf2 suppresses
macrophage inflammatory response by blocking proinflammatory cytokine transcription. Nat Commun.
2016;7:11624. Epub 20160523. doi: 10.1038/ncomms11624. PubMed PMID: 27211851; PubMed Central
PMCID: PMCPMC4879264.

46. Abby SS, Cury J, Guglielmini J, Neron B, Touchon M, Rocha EP. Identification of protein secretion
systems in bacterial genomes. Sci Rep. 2016;6:23080. Epub 20160316. doi: 10.1038/srep23080. PubMed
PMID: 26979785; PubMed Central PMCID: PMCPMC4793230.

47. Nahori MA, Fournie-Amazouz E, Que-Gewirth NS, Balloy V, Chignard M, Raetz CR, et al. Differential
TLR recognition of leptospiral lipid A and lipopolysaccharide in murine and human cells. J Immunol.
2005;175(9):6022-31. doi: 10.4049/jimmunol.175.9.6022. PubMed PMID: 16237097.

48. Bonhomme D, Hernandez-Trejo V, Papadopoulos S, Pigache R, Fanton d'Andon M, Outlia A, et al.
Leptospira interrogans prevents macrophage cell death and pyroptotic IL1ß releasethrough atypical
lipopolysaccharide. BioRXiv. doi: 10.1101/2022.07.25.501344.

49. Terawaki S, Camosseto V, Prete F, Wenger T, Papadopoulos A, Rondeau C, et al. RUN and FYVE
domain-containing protein 4 enhances autophagy and lysosome tethering in response to Interleukin-4. J
Cell Biol. 2015;210(7):1133-52. doi: 10.1083/jcb.201501059. PubMed PMID: 26416964; PubMed Central
PMCID: PMCPMC4586740.

50. Ye X, Zhu M, Che X, Wang H, Liang XJ, Wu C, et al. Lipopolysaccharide induces neuroinflammation in
microglia by activating the MTOR pathway and downregulating Vps34 to inhibit autophagosome
formation. J Neuroinflammation. 2020;17(1):18. Epub 20200111. doi: 10.1186/s12974-019-1644-8.
PubMed PMID: 31926553; PubMed Central PMCID: PMCPMC6954631.

51. Yin S, Cao W. Toll-Like Receptor Signaling Induces Nrf2 Pathway Activation through p62-Triggered
Keap1 Degradation. Mol Cell Biol. 2015;35(15):2673-83. Epub 20150526. doi: 10.1128/MCB.00105-15.
PubMed PMID: 26012548; PubMed Central PMCID: PMCPMC4524114.

52. Jiang T, Harder B, Rojo de la Vega M, Wong PK, Chapman E, Zhang DD. p62 links autophagy and Nrf2
signaling. Free Radic Biol Med. 2015;88(Pt B):199-204. Epub 20150624. doi:
10.1016/j.freeradbiomed.2015.06.014. PubMed PMID: 26117325; PubMed Central PMCID:
PMCPMC4628872.

53. Gilardini Montani MS, Santarelli R, Granato M, Gonnella R, Torrisi MR, Faggioni A, et al. EBV reduces
autophagy, intracellular ROS and mitochondria to impair monocyte survival and differentiation.
Autophagy. 2019;15(4):652-67. Epub 20181023. doi: 10.1080/15548627.2018.1536530. PubMed PMID:
30324853; PubMed Central PMCID: PMCPMC6526866.

54. Bichiou H, Rabhi S, Ben Hamda C, Bouabid C, Belghith M, Piquemal D, et al. Leishmania Parasites
Differently Regulate Antioxidant Genes in Macrophages Derived From Resistant and Susceptible Mice.
Front Cell Infect Microbiol. 2021;11:748738. Epub 20211015. doi: 10.3389/fcimb.2021.748738. PubMed
PMID: 34722338; PubMed Central PMCID: PMCPMC8554229.

55. Zavala-Alvarado C, Sismeiro O, Legendre R, Varet H, Bussotti G, Bayram J, et al. The transcriptional
response of pathogenic Leptospira to peroxide reveals new defenses against infection-related oxidative
stress. PLoS Pathog. 2020;16(10):e1008904. Epub 20201006. doi: 10.1371/journal.ppat.1008904. PubMed
PMID: 33021995; PubMed Central PMCID: PMCPMC7567364.

56. Vivarini AC, Calegari-Silva TC, Saliba AM, Boaventura VS, Franca-Costa J, Khouri R, et al. Systems
Approach Reveals Nuclear Factor Erythroid 2-Related Factor 2/Protein Kinase R Crosstalk in Human
Cutaneous Leishmaniasis. Front Immunol. 2017;8:1127. Epub 20170915. doi: 10.3389/fimmu.2017.01127.
PubMed PMID: 28959260; PubMed Central PMCID: PMCPMC5605755.

57. Macoch M, Morzadec C, Genard R, Pallardy M, Kerdine-Romer S, Fardel O, et al. Nrf2-dependent
repression of interleukin-12 expression in human dendritic cells exposed to inorganic arsenic. Free Radic
Biol Med. 2015;88(Pt B):381-90. Epub 20150211. doi: 10.1016/j.freeradbiomed.2015.02.003. PubMed
PMID: 25680285.

58. Helou DG, Braham S, De Chaisemartin L, Granger V, Damien MH, Pallardy M, et al. Nrf2 downregulates
zymosan-induced neutrophil activation and modulates migration. PLoS One. 2019;14(8):e0216465. Epub
20190816. doi: 10.1371/journal.pone.0216465. PubMed PMID: 31419224; PubMed Central PMCID:
PMCPMC6697320.

59. Fanton d'Andon M, Quellard N, Fernandez B, Ratet G, Lacroix-Lamande S, Vandewalle A, et al.
Leptospira Interrogans induces fibrosis in the mouse kidney through Inos-dependent, TLR- and NLR-
independent signaling pathways. PLoS Negl Trop Dis. 2014;8(1):e2664. Epub 20140130. doi:
10.1371/journal.pntd.0002664. PubMed PMID: 24498450; PubMed Central PMCID: PMCPMC3907306.

60. Raffray L, Giry C, Vandroux D, Kuli B, Randrianjohany A, Pequin AM, et al. Major Neutrophilia
Observed in Acute Phase of Human Leptospirosis Is Not Associated with Increased Expression of
Granulocyte Cell Activation Markers. PLoS One. 2016;11(11):e0165716. Epub 20161101. doi:
10.1371/journal.pone.0165716. PubMed PMID: 27802348; PubMed Central PMCID: PMCPMC5089758.

61. Olagnier D, Brandtoft AM, Gunderstofte C, Villadsen NL, Krapp C, Thielke AL, et al. Nrf2 negatively
regulates STING indicating a link between antiviral sensing and metabolic reprogramming. Nat Commun.
2018;9(1):3506. Epub 20180829. doi: 10.1038/s41467-018-05861-7. PubMed PMID: 30158636; PubMed
Central PMCID: PMCPMC6115435.

62. Werts C, le Bourhis L, Liu J, Magalhaes JG, Carneiro LA, Fritz JH, et al. Nod1 and Nod2 induce
CCL5/RANTES through the NF-kappaB pathway. Eur J Immunol. 2007;37(9):2499-508. doi:
10.1002/eji.200737069. PubMed PMID: 17705131.

63. Santecchia I, Vernel-Pauillac F, Rasid O, Quintin J, Gomes-Solecki M, Boneca IG, et al. Innate immune
memory through TLR2 and NOD2 contributes to the control of Leptospira interrogans infection. PLoS
Pathog. 2019;15(5):e1007811. Epub 20190520. doi: 10.1371/journal.ppat.1007811. PubMed PMID:
31107928; PubMed Central PMCID: PMCPMC6544334.

64. Bonhomme D, Werts C. Purification of LPS from Leptospira. Methods Mol Biol. 2020;2134:53-65. doi:
10.1007/978-1-0716-0459-5_6. PubMed PMID: 32632859.

 FIGURES LEGENDS

Figure 1. *L. interrogans* **reduces the quantity of autophagosomes in macrophages**

A) WB of LC3 in C57/BL6 mouse peritoneal cells 4 h post-intraperitoneal injection of PBS, 5 x 10^7
L. interrogans serovar Manilae strain MFLum1 (*L.i*), 30 µg of muropeptides (murP) (MDP+FK156) or
30 µg of rapamycin (Rap). **B)** WB of LC3 in BMDMs 24 h post-infection with either live or
heat-killed (HK) *L. interrogans* serovar Manilae strain MFLum1 at MOI 100, or stimulation with
500 nM rapamycin. **C)** WB of LC3 in BMDMs 24 h post-infection with *L. interrogans* serovar
Manilae strain MFLum1 at MOI 100, with or without bafilomycin A1 (BafA1) addition 4 h before cell
collection. **D)** WB analyses and quantifications of LC3 in BMDMs 1 h, 4 h and 24 h post-infection
with either live or HK *L. interrogans* serovar Manilae strain MFLum1 at MOI 100. Quantifications
represent the ratio of LC3-II in stimulated cells over non-stimulated cells, and after stain free
normalization. Each dot corresponds to the quantification of a blot from an independent experiment (*n*
from 3 to 6). **E)** RT-qPCR analyses of LC3 mRNA levels in BMDMs 24 h post-infection with MOI
100 of *L. interrogans* serovar Manilae strain L495. Bars correspond to mean +/- SD of technical
replicates (*n*=4). **F-G)** High Content (HC) confocal microscopy images and quantifications of
RAW-mLC3-DiFluo cells 4 h post-infection with *L. interrogans* serovar Manilae strain MFLum1 at
MOI 100, with or without bafilomycin A1 (BafA1) addition 2 h before cell fixation. Bars correspond
to mean +/- SD of technical replicates (*n*=4). **A-E)** Data presented are representative of at least 3
independent experiments.

Figure 2. *L. interrogans* **induces specific accumulation of p62 adaptor in puncta-like structures**

A) WB of p62 in BMDMs 1 h, 4 h and 24 h post-infection with *L. interrogans* serovar Manilae strain
MFLum1. **B)** WB of p62 in BMDMs 24 h post-infection with *L. interrogans* serovar Manilae strain
MFLum1, serovar Copenhageni strain Fiocruz L1-130, serovar Icterohaemorrhagiae strain Verdun,
and *L. biflexa* serovar Patoc strain Patoc I. **C)** RT-qPCR analyses of p62 mRNA levels in BMDMs
24 h post-infection with *L. interrogans* serovar Manilae strain L495. Bars correspond to mean +/- SD
of technical replicates (*n*=4). **D)** Epifluorescence analyses of p62 in BMDMs 4 h post-infection with
L. interrogans serovar Manilae strain MFLum1. **E)** High Content (HC) confocal microscopy analyses
of the number of p62 puncta *per* cell and the percentage of p62 positive cells 1 h, 4 h and 24 h
post-infection of RAW264.7 cells with *L. interrogans* serovar Manilae strain L495. **F)** HC single cell
microscopy analysis of the number of p62 puncta *per* cell 24 h post-infection of RAW264.7 cells with
L. interrogans serovar Manilae strain L495. **A-F)** Data presented are representative of at least 3
independent experiments.

Figure 3. Autophagy modulation is mediated by TLR2/TLR4

A) WB of LC3 and p62 in WT and TLR2/4 dko BMDMs 24 h post-infection with *L. interrogans*
serovar Manilae strain MFLum1 at MOI 100, or upon starvation induction by incubation in EBSS
medium. **B)** WB of p62 in WT and MyD88/TRIF dko BMDMs 24 h post-infection with *L. interrogans*
serovar Manilae strain MFLum1. **C)** RT-qPCR analyses of p62 mRNA levels in WT and TLR2/4 dko
BMDMs 24 h post-infection with *L. interrogans* serovar Manilae strain L495 at MOI 100. Bars
correspond to mean of independent experiments (*n*=3). **D)** High Content (HC) single cell microscopy
analysis of the number of p62 puncta *per* cell 24 h post-infection of WT, TLR2/TLR4 dko and

MyD88/TRIF dko BMDMs with *L. interrogans* serovar Manilae strain L495, serovar Copenhageni
strain Fiocruz L1-130 and serovar Icterohaemorrhagiae strain Verdun. **A-D)** Data presented are
representative of at least 2 independent experiments.

Figure 4. p62 accumulation is mediated by the leptospiral LPS

A) WB of p62 in BMDMs 1 h, 4 h and 24 h post-stimulation with heat-killed (56°C, 30 min)
L. interrogans serovar Manilae strain MFLum1 at MOI 100 or 24 h post-stimulation with 1 µg/mL of
the corresponding purified leptospiral LPS. **B)** High Content (HC) confocal microscopy analyses of
the number of p62 puncta *per* cell and the percentage of p62 positive cells 1 h, 4 h and 24 h post-
stimulation of RAW264.7 cells with MOI 100 of either live or heat-killed (56°C, 30 min)
L. interrogans serovar Manilae strain L495 or stimulation with 1 µg/mL of the corresponding purified
leptospiral LPS. **C)** HC single cell microscopy analysis of the number of p62 puncta *per* cell 24 h
post-infection of RAW264.7 cells with MOI 100 of either live or heat-killed (56°C, 30 min)
L. interrogans serovar Manilae strain L495 or stimulation with 1 µg/mL of the corresponding purified
leptospiral LPS. **A-C)** Data presented are representative of at least 3 independent experiments.

Figure 5. Leptospiral LPS also triggers translocation of NRF2 transcription factor

A) Confocal microscopy analyses of NRF2 in BMDMs 4 h post-infection with *L. interrogans* serovar
Manilae strain L495 at MOI 100. **B)** High Content (HC) single cell microscopy analysis of the levels
of NRF2 intensity in the nucleus and 4 h post-infection of RAW264.7 cells with *L. interrogans*
serovar Manilae strain L495, serovar Copenhageni strain Fiocruz L1-130 or serovar
Icterohaemorrhagiae strain Verdun, at MOI 100. Fluorescence is expressed in arbitrary units (AU),
dashed lines correspond to median. **C)** HC analysis of the number of cells with NRF2 intensity ratio
[nucleus/cytoplasm] > 1.4 4 h post-infection of RAW264.7 cells with *L. interrogans* serovar Manilae
strain L495, serovar Copenhageni strain Fiocruz L1-130 or serovar Icterohaemorrhagiae strain
Verdun, at MOI 100. **D)** HC single cell microscopy analysis of the levels of NRF2 intensity in the
nucleus 4 h post-infection of RAW264.7 cells with MOI 100 of either live or heat-killed (56°C,
30 min) *L. interrogans* serovar Manilae strain L495 or stimulation with 1 µg/mL of the corresponding
purified leptospiral LPS. Fluorescence is expressed in arbitrary units (AU), dashed lines correspond to
median. **E)** HC analysis of the number of cells with NRF2 intensity ratio [nucleus/cytoplasm] > 1.4
4 h post-infection of RAW264.7 cells with MOI 100 of either live or heat-killed (56°C, 30 min)
L. interrogans serovar Manilae strain L495 or stimulation with 1 µg/mL of the corresponding purified
leptospiral LPS. **A-E)** Data presented are representative of at least 2 independent experiments.

Figure 6. p62 and NRF2 constitute a retro-activation loop upon infection with *L. interrogans*

A) High Content (HC) single cell microscopy analysis of NRF2 intensity ratio [nucleus/cytoplasm] in
BMDMs transfected with scramble (scrbl) or p62 siRNA and infected for 4 h with *L. interrogans*
serovar Manilae strain L495 at MOI 100. **B)** Corresponding controls are microscopy analyses and
quantification of p62 intensity in the same transfection and infection conditions. Lines correspond to
mean +/- SD of technical replicates (*n*=3). **C)** HC single cell microscopy analysis of p62 puncta
formation in BMDMs transfected with scrbl or NRF2 siRNA and infected for 4 h with *L. interrogans*
serovar Manilae strain L495 at MOI 100. **D)** Corresponding controls are microscopy analyses and
quantification of NRF2 intensity in the same transfection and infection conditions. Lines correspond to
mean +/- SD of technical replicates (*n*=3). **E)** WB of p62 in BMDMs transfected with scrbl or NRF2

siRNA and infected for 24 h with *L. interrogans* serovar Manilae strain L495 at MOI 10 or 100.
F) RT-qPCR analyses of p62 mRNA levels in RAW264.7 cells transfected with scrbl or NRF2 siRNA
and infected for 24 h with *L. interrogans* serovar Manilae strain L495 at MOI 100. Bars correspond to
mean of independent experiments (*n*=2). **A-E)** Data presented are representative of at least 3
independent experiments.

Figure 7. NRF2 translocation prevents inflammation upon infection

A) Cell viability (measured by MTT assay) and **B)** LDH release (measured by CyQuant assay) in
RAW264.7 cells transfected with scramble (scrbl) or NRF2 siRNA and infected for 24 h with
L. interrogans serovar Manilae strain L495 at MOI 100. Bars correspond to mean +/- SD of technical
replicates (*n*=4). **B)** RT-qPCR analyses of iNOS, IL6, TNF and IL10 mRNA levels in RAW264.7 cells
transfected with scrbl or NRF2 siRNA and infected for 10 h with *L. interrogans* serovar Manilae strain
L495 at MOI 100. Bars correspond to mean of independent experiments (*n*=3). **C)** NO dosage by
Griess reaction and, IL6, TNF and IL10 by ELISA in RAW264.7 cells transfected with scrbl or NRF2
siRNA and infected for 24 h with *L. interrogans* serovar Manilae strain L495 at MOI 100. Bars
correspond to mean +/- SD of technical replicates (*n*=4). **A-C)** Data presented are representative of at
least 3 independent experiments.

A

NI   *L. i*

1h   4h   24h   1h   4h   24h

p62

Stain free

(BMDMs)

B

   L. interrogans   *L. biflexa*

NI   L495   Fiocruz   Verdun   Patoc

p62

Stain free

(BMDMs)

C

p62 mRNA levels
(BMDMs, 24h pi)

Fold change

NI   *L.i*

D

Confocal microscopy
(BMDMs, 4h pi)

NI   *L. i*

DAPI - *L.interrogans* - p62

E

Kinetics of p62 puncta formation
(RAW264.7)

Number of puncta / cell

Time (hours)

○ NS
● NI
○ *L. i*
● BafA1

% cells with p62 puncta

Time (hours)

○ NS
● NI
○ *L. i*
● BafA1

F

Single cell analysis of p62 puncta
(RAW264.7, 24h pi)

Number of puncta / cell

NS   NI   *L. i*   BafA1

A

WT | TLR2/4 dko
NI EBSS *L.i* | NI EBSS *L.i*

LC3-I
LC3-II

Stain
free

(BMDMs)

WT | TLR2/4 dko
NI EBSS *L.i* | NI EBSS *L.i*

p62

Actin

(BMDMs)

B

WT | MyD88/Trif dko
NI *L.i* | NI *L.i*

p62

Stain
free

(BMDMs)

C

p62 mRNA levels
(BMDMs, 24h pi)

□ WT
■ TLR2/4 dko

Fold change

NS   *L.i*

**

D

Single cell analysis of p62 puncta upon infection
(BMDMs, 24h pi)

Number of puncta / cell

● WT
● TLR2/TLR4 dko
● MyD88/Trif dko

NI   L495 Fiocruz Verdun | NI   L495 Fiocruz Verdun | NI   L495 Fiocruz Verdun
      L.i | *L.i* | *L.i*

*** ** *** *** *** ***

A

NI   HK *L.i*   NI   *L.i* LPS

1h   4h   24h   1h   4h   24h   24h   24h

LC3-I
LC3-II

Stain
free

(BMDMs)

B

Kinetics of p62 puncta formation
(RAW264.7)

NS
NI
Live
HK ⎤
LPS ⎦ *L.i*
BafA1

C

Single cell analysis of p62 puncta
(RAW264.7, 24h pi)

NI   Live   HK   LPS   BafA1

L. interrogans

A

NI   *L. i*

Confocal microscopy
(BMDMs, 4h pi)

DAPI - *L.interrogans* - NRF2

B   *L.i* serovars

**Single cell analysis
of NRF2 intensity
in the nucleus**
(RAW264.7, 4h pi)

***

AU

NS   NI   L495   Fiocruz   Verdun

L. i

C   **Percentage of cells with NRF2
nucleus / cytoplasm ratio > 1.4**
(RAW264.7, 4h pi)

**

*

% cells

NS   NI   L495   Fiocruz   Verdun

L. i

D   HK and LPS

**Single cell analysis
of NRF2 intensity
in the nucleus**
(RAW264.7, 4h pi)

***

AU

NS   NI   Live   HK   LPS

L. i

E   **Percentage of cells with NRF2
nucleus / cytoplasm ratio > 1.4**
(RAW264.7, 4h pi)

**

*

% cells

NS   NI   Live   HK   LPS

L. i