Targeted IS-element sequencing uncovers transposition dynamics during selective

pressure in enterococci

Joshua M. Kirsch[1], Shannon Ely[1], Madison E. Stellfox[2], Karthik Hullahalli[3], Phat Luong[1], Kelli L.

Palmer[3], Daria Van Tyne[2], Breck A. Duerkop[1*]

[1]Department of Immunology and Microbiology, University of Colorado School of Medicine,

Aurora, CO 80045, USA

[2]Department of Medicine, Division of Infectious Diseases, University of Pittsburgh, Pittsburgh,

PA 15214, USA

[3]Department of Biological Sciences, University of Texas at Dallas, Richardson, TX 75080, USA

*Correspondence: Breck A. Duerkop breck.duerkop@cuanschutz.edu

Abstract

Insertion sequences (IS) are simple transposons implicated in the genome evolution of diverse

pathogenic bacterial species. Enterococci have emerged as important human intestinal

pathogens with newly adapted virulence potential and antibiotic resistance. These genetic

features arose in tandem with large-scale genome evolution mediated by mobile elements.

Pathoadaptation in enterococci is thought to be mediated in part by the IS element IS256

through gene inactivation and recombination events. However, the regulation of IS256 and the

mechanisms controlling its activation are not well understood. Here, we adapt an IS256-specfic

deep sequencing method to describe how chronic lytic phage infection drives a widespread

expansion of IS256 in *E. faecalis* and how antibiotic exposure is associated with IS256

expansion in both *E. faecalis* and *E. faecium* during a clinical human infection. We show through

comparative genomics that IS256 is primarily found in hospital-adapted enterococcal isolates.

IS256 transposase gene expression analyses reveal that IS256 mobility is regulated at the

transcriptional and translational levels in *E. faecalis*, indicating tight control of IS256 activation in

the absence of selective pressure. Our findings reveal that stressors such as phages and

antibiotic exposure drives rapid genome-scale transposition in the enterococci. IS256 expansion

can therefore explain how selective pressures mediate evolution of the enterococcal genome,

ultimately leading to the emergence of dominant nosocomial lineages that threaten human

health.

Introduction

Enterococci, including the human commensals *Enterococcus faecalis* and *Enterococcus faecium*, are opportunistic pathogens of public health concern due to their acquisition of antibiotic resistance and virulence traits[1]. Enterococcal infections are a leading cause of infective endocarditis[2, 3] and 30% of nosocomial infections are vancomycin resistant[4, 5]. Antibiotic resistance in enterococci is often encoded on mobile genetic elements, including plasmids and composite transposons[6]. Plasmids, especially those containing Inc18, Rep_3, and RepA-N replicon-containing plasmids, frequently encode resistance genes to diverse antibiotics, including chloramphenicol, erythromycin, and gentamicin[7, 8]. Composite transposons are typically organized with a cargo gene (such as an antibiotic resistance gene) flanked by insertion sequence (IS) elements. Examples in the enterococci include Tn4001 or Tn4031, which carries gentamicin resistance genes[9] and Tn1546 and Tn1547, which carry the vancomycin resistance *vanA* and *vanB* operons, respectively[10, 11].

        To combat clinically relevant enterococcal strains, new therapeutics are urgently needed. Recently, the concept of using bacterial viruses (bacteriophages or phages) to treat multi-drug resistant (MDR) bacteria is being revisited[12]. Phage therapy case studies show that these viruses can be efficacious against refractory MDR bacteria in human patients[13, 14, 15, 16, 17]. However, bacterial resistance to phage infection develops quickly[18, 19, 20, 21] and the lessons learned from the past century of antibiotic use suggest that bacteria will gain resistance to phage infection after widespread phage treatment[22]. Acquired phage resistance, although beneficial to bacteria in the face of phage pressure, can come with fitness costs that dampen antibiotic resistance and virulence[12, 23, 24, 25]. Therefore, it is imperative to fully understand the mechanisms that promote phage-resistance in bacteria, and the phenotypic outcomes of phage resistance. With this knowledge it will be possible to use phage resistance as a tool that can be leveraged against bacteria to enhance current antibacterial therapeutics.

Enterococci have remarkably diverse genomes containing numerous mobile elements

that contribute to their adaptation and evolution[26, 27, 28]. These include plasmids, prophages, and

transposable insertion sequence (IS) elements. Pathogenic enterococci frequently contain

multiple IS elements, and IS elements appear to be a feature of newly-adapted nosocomial

strains[29]. In the widely studied nosocomial type-strain *E. faecalis* V583, there are 38 IS

elements consisting of 11 different types[26]. IS256 is the most abundant IS element in the *E.*

faecalis V583 genome, with six chromosomal copies and four plasmid copies. IS256 is also

common in other gram-positive bacteria, including staphylococci, where it has been widely

studied[30, 31, 32]. In the staphylococci, IS256 mediates clinically-relevant phenotypes, such as

small colony variation[33], biofilm formation[31], and antibiotic resistance[34]. The transcription factor

σ-B negatively regulates IS256 transposition in the staphylococci through the production of a 3'

antisense RNA[31, 35]. IS256 is a crucial component for enterococcal genome adaptation. In *E.*

faecalis V583, the virulence factor cytolysin is attenuated by IS256 and related IS905

insertions[36]. Additionally, copies of IS256 in the *E. faecalis* V583 pheromone-responsive

plasmids pTEF1 and pTEF2 recombine with chromosomal IS256 copies to mobilize broad

regions of the *E. faecalis* genome[37]. A related IS element, IS16, has been shown to be highly

abundant in hospital-adapted *E. faecium* isolates, suggesting that IS16 may aid in the success

of *E. faecium* as a nosocomial pathogen[38]. Considering IS elements are likely involved in the

pathoadaptation of the enterococci, little is known about how these elements are regulated and

what events lead to their activation.

In this work, we investigated both the regulation and spread of IS256 transposition in *E.*

faecalis and *E. faecium*. We found that IS256 is present in multiple, genetically disparate

isolates of both species, and is enriched in hospital adapted lineages. At steady state, IS256

produces a wealth of low abundance insertions throughout the chromosome and is regulated at

both the transcriptional and translational levels to tightly control activation. We discovered that

phage infection and antibiotic usage in different biological settings were associated with a

dramatic expansion of IS256 insertion events. In the case of phage infection, cell populations

experiencing IS256 expansion maintained phage genomes as independently replicating

episomes and a subpopulation of these cells chronically shed phage particles throughout

growth. Phage shedding provided a competitive advantage during co-culture with phage-

susceptible enterococci, but this advantage is lost when co-cultured with phage-resistant

enterococci. This suggests that phage genome carriage is a strong selective pressure that

drives IS256 mobilization and can be used to occupy an environmental niche when competing

with phage-susceptible bacterial strains. Lastly, we investigated how IS256 diversifies

enterococcal genomes in a chronically infected patient, and found that IS256 insertion

abundances in vivo coincided with specific antibiotic usage. Together, this work sheds light on

how IS elements are regulated and expanded in enterococcal genomes during both phage

predation and clinical infection, and provides evidence for phage carriage as an important

selective pressure that promotes IS256 mobility. Furthermore, this work suggests that

therapeutic use of both phages and antibiotics could cause rapid and widespread enterococcal

genome evolution for which the physiological consequences are unknown.

Results

IS256 is common within enterococcal lineages associated with hospital adaptation and

pathogenesis

IS256 is an important factor driving the pathoadaptation of *E. faecalis*[36], yet it is unknown

how widely IS256 is distributed among *E. faecalis* strains. To answer this, we searched all

available *E. faecalis* genomes from NCBI RefSeq (2065 genomes) for IS256 transposases with

100% amino acid similarity to the IS256 transposase copies found in *E. faecalis* V583. A total of

232 genomes contained one or more IS256 sequences (Fig. 1A). The sporadic distribution of

IS256 within the *E. faecalis* phylogenetic tree suggests that IS256 is a recent addition to many

E. faecalis genomes and has arisen within multiple discrete lineages of this species. To further

characterize the distribution of IS256 across *E. faecalis* strains, we performed in silico multilocus

sequence typing (MLST) on all *E. faecalis* genomes and compared the proportions of genomes

with and without IS256 in each sequence type (ST) (Fig. 1B). IS256 occurs within a variety of

ST clades and is most abundant in in ST6, ST103, ST778, and ST388 genomes, all of which

are associated with hospital-adapted, opportunistic pathogenic lineages (Table S1A)[39, 40, 41, 42].

We compared the abundances of different virulence factors between genomes with and without

IS256 (Fig. 1C & Table S1C). IS256-containing genomes are enriched for multiple virulence

factors compared to genomes lacking IS256, demonstrating that IS256 preferentially occurs in

hospital-acquired and virulent *E. faecalis* strains. We next expanded this analysis to include all

E. faecium genomes from RefSeq (2306 genomes) and similar to *E. faecalis*, *E. faecium* IS256

is primarily found in the nosocomial lineages ST17, ST664, ST736, and ST18 (Fig. 1D-E and

Table S1B)[43, 44]. Although *E. faecium* IS256-containing genomes have a significant enrichment

in some virulence factors, this was less than *E. faecalis* (Fig. 1F and Table S1D). Finally, IS256

is more frequently found in *E. faecium* genomes compared to *E. faecalis* genomes, confirming

prior research (Fig. 1G)[45]. Together these data show that IS256 is widely distributed within the

enterococci and is preferentially found in nosocomial and virulent isolates.

IS-Seq identifies widespread movement of IS256 in E. faecalis that is transcriptionally

and translationally controlled

148   Previously, IS256 sequences were identified as hot spots in *E. faecalis* V583 that

facilitated the integration of the endogenous plasmids pTEF1 and pTEF2, leading to the

mobilization of the pathogenicity island and other chromosomal regions[37]. Instances of IS256

insertions leading to the inactivation of *E. faecalis* genes involved in diverse phenotypes have

been described[19, 46, 47, 48]. However, we lack a complete understanding of IS256 mobility across

the *E. faecalis* genome during selective pressures that would potentially drive IS256 activation

and genome diversification. To assess genome-wide IS256 insertions in *E. faecalis* we adapted

a next-generation sequencing (NGS) enrichment technique for use with IS256. This technique,

IS-Seq, includes IS256 amplicon enrichment during NGS library construction followed by read

mapping of IS256-chromosomal junctions to identify specific IS256 insertion locations (Fig.

S1)[49]. Similar techniques have been used to identify IS element insertions in *Acinetobacter*[50]

and *Mycobacterium*[51]. IS-Seq of wild-type (WT) *E. faecalis* V583 under steady state conditions

identified the known IS256 locations in the chromosome and on the three pTEF plasmids (Fig.

2A & S2). One of the IS256 copies on pTEF1 lacks a canonical left inverted repeat (IR),

preventing binning of the IS-Seq reads originating from this locus. Numerous low abundance

IS256 insertions are observed throughout the *E. faecalis* genome, indicating promiscuous

movement of the element within subpopulations of cells (Fig. 2A). The three replicate cultures

tested in Fig. 2A show that each *E. faecalis* population analyzed has a unique repertoire of

IS256 insertions. We hypothesize that these low abundance steady-state insertion events

provide a mechanism for rapid genome diversification following exposure to selective pressures.

To understand how IS256 is regulated in *E. faecalis* under steady state conditions, we

investigated possible mechanisms of repression of the IS element. First, we assessed the

transcription of the IS256 transposase (Tnp) gene. During growth in rich media, IS256 gene

transcription modestly increases during logarithmic growth, and doubles in comparison to the

expression of the reference gene *clpX* (Fig. 2B). Considering there are 10 copies of IS256 in *E.*

faecalis V583, each IS256 copy achieves 1/5th the transcriptional activity of a single *clpX* gene.

Thus, the relatively weak IS256 promoter may limit over-activation of the element. Next, we

investigated whether IS256 is translationally regulated. Other IS elements have been reported

to be controlled by expression of antisense small RNAs (asRNA)[52]. These asRNAs bind to

sense transcripts at or before the start codon and ribosome binding site to prevent translation of

the sense transcript. We reanalyzed a publicly available stranded RNA-Seq dataset of *E.*

faecalis V583[53] by aligning reads to an IS256 element with delineation of the sense and

antisense alignments (Fig. 2C). The translational start site (TSS) of IS256 has not been

experimentally confirmed in any bacterial species and is predicted to occur in the 5' terminus of

the element[54]. We demonstrate that IS256 has two TSSs indicated by high antisense read

mapping coverage, demonstrating that antisense inhibition is present at both TSSs. If translation

initiates at the site-2 TSS this would produce a truncated IS256 Tnp. It is unclear whether this

truncated IS256 Tnp would function similar to the full-length IS256 Tnp. Upon further

investigation of these TSSs and their asRNAs, we found that the site-1 asRNA is likely under

the control of a canonical σ-70 promoter. To identify if this asRNA controls IS256 Tnp

translation, we constructed an IS256 Tnp-GFP translational reporter. This construct consists of

an IS256 Tnp coding sequence fused in frame at its C-terminus to a green fluorescence protein

(GFP) gene. GFP fluorescence serves as a proxy for translation of the IS256 Tnp. In addition,

we built a version of the IS256 Tnp-GFP fusion lacking the predicted -10 element of the site-1

asRNA. We transformed these constructs into *E. faecalis* OG1RF, a strain that lacks native

IS256 elements and IS256-derived asRNAs in its genome, and measured GFP fluorescence in

these cells (Fig. 2D). In cells lacking the site-1 asRNA, there is significantly higher GFP

fluorescence compared to cells with an intact site-1 asRNA promoter, demonstrating that these

asRNAs repress IS256 Tnp translation. These results show that in *E. faecalis* IS256 is regulated

by multiple mechanisms at both the transcriptional and translational levels.

IS256 selection during phage predation

Our previous work suggested that phage predation increases IS256 transposition in *E.*

faecalis V583 and is used as a mechanism of phage resistance[18]. To determine if phage

selective pressure results in the expansion of IS256 mobility in *E. faecalis*, we challenged *E.*

faecalis V583 with the phage phi19[18]. Following phage exposure, we isolated single colonies

that were resistant to phi19. Whole genome sequencing (WGS) of these phage-resistant

isolates predicted an expansion of new IS256 insertions in phi19 resistant isolates (referred to

as 19RS strains) (Fig. 3A). Among the new IS256 insertions in 19RS strains, we found that an

insertion was predicted in the 3' end of *epaX*, a glycosyltransferase gene involved in the

biosynthesis of the enterococcal polysaccharide antigen that is required for infection by phi19[18].

To verify this insertion and its orientation, we performed PCR using genomic DNA (gDNA) from

four 19RS strains and wild type (WT) *E. faecalis* V583 (Fig. 3B). We found that all 19RS strains

harbored an IS256 insertion in *epaX* in both the positive and negative strand orientations. This

IS256 insertion likely prevents functional EpaX from being expressed blocking phage infection.

Recently, Lossouarn et al. identified IS256 insertions inactivating *epaX*, confirming that an

IS256 insertion in *epaX* is sufficient to prevent phage infection[48]. These results demonstrate that

19RS strains contain novel IS256 insertions not found in WT *E. faecalis* V583. These results

also suggest that 19RS strains, even though they were isolated as individual colonies, are

heterogeneous populations with varied abundances of IS256 insertions.

To determine the global effect of IS256 transposition in the 19RS strains, we performed

Southern blotting using a probe against IS256 (Fig. 3C). We found differential IS256 banding

patterns in the 19RS strains that are not found in WT *E. faecalis* V583, demonstrating that

phage infection modulates IS256 transposition. To test if IS256 provides an advantage to *E.*

faecalis against phage infection, we transformed a multi-copy plasmid containing either an

IS256 gene or an inactive IS256 gene lacking the coding sequences for the two catalytic

aspartic acid residues of the Tnp's DDE motif into *E. faecalis* OG1RF, which lacks native IS256

and infected these cells with phi19 (Fig. 3D). We found that cells containing a functional IS256

gene were more resistant to phage infection compared to cells containing the catalytically dead

Tnp or the empty vector, indicating that IS256 carriage increases mutation frequency. To

understand if IS256 activation was initiated by phi19 infection, we quantified the number of

IS256 circular intermediates. IS256 elements circularize during transposition, which can be used

to measure active transposition[30, 31]. During the course of phi19 infection, IS256 circular

intermediates significantly increased relative to a mock-infected cells (Fig. 3E).

Southern blotting and WGS analysis indicate that phage predation increases IS256

expansion across the *E. faecalis* genome. However, these methods are low resolution. To

quantify the exact locations and abundances of IS256 insertions following phage pressure, we

performed IS-Seq on 19RS strains and compared these to WT *E. faecalis* V583 (Fig. 4A). IS-

Seq revealed a genome-wide expansion of IS256 insertions at new locations in 19RS strains

(Fig. 4A). Pairwise comparisons determined that 19RS strains are significantly enriched in the

number of novel IS256 insertions compared WT *E. faecalis* V583 (Figure 4B & Table S2). New

IS256 insertions in the pTEF plasmids were also found in the 19RS strains (Fig. S3A-C).

Additionally, we measured IS256 circular intermediates in our IS-Seq dataset by aligning IS256

reads (which originate from the left terminus of IS256) to the right terminus of an IS256 element.

We found that 19RS strains have a significantly greater number of IS256 circular intermediates

compared to WT *E. faecalis* V583 (Fig. 4C).

We focused on two genomic regions that experienced a high level of IS256 insertions in

the 19RS strains. The first was the *epa* locus. In addition to numerous novel insertions in *epaX*,

we found insertions within other *epa* genes including *epaR*, *epaS*, *epaAA* and *epaAB*. (Fig.

S4A). IS256 insertions in these *epa* genes would alter the teichoic acid decoration of the core

rhammnose-containing Epa, likely altering the surface chemistry of *E. faecalis* and rendering

them resistant to phage infection[18]. We also identified multiple insertions in the *vex/vnc* operon

(Fig. S4B), which is purported to have both virulence and antibiotic resistance functions[55, 56].

E. faecalis is a native member of the intestinal microbiota and can outgrow and cause

disease during intestinal dysbiosis. Phages have been proposed as a treatment option for

difficult to control enterococcal intestinal blooms[57]. To determine if phage predation supported

IS256 expansion in the intestine, we colonized mice with *E. faecalis* V583 and orally challenged

these animals with phi19 (Fig. S5). We performed IS-Seq on fecal gDNA from these animals

and found that phi19 exposure in the intestine also increased new IS256 insertions, albeit to a

lesser extent than in vitro (Fig. 4D-F, Fig. S3D-F, & Table S3). We found that similar to cells

exposed to phi19 in vitro, IS256 insertions within the same location in *epaX* and the *vex/vnc*

operons are observed in the mouse intestine following phage administration (Fig. S4C-D). This

suggests that phage selection plays a role in IS256 expansion with the context of the native

intestinal habitat.

Enterococci with expanded IS256 genomes chronically shed phi19 phages.

Phages can lysogenize their host bacterium[58]. During lysogeny the phage genome is

maintained in a dormant state as an integrated element within the bacterial chromosome or as a

cytoplasmic episome. Episomal phage genomes can also be in an intermediate state between

lytic replication and lysogenic conversion referred to as a pseudolysogen or carrier state. Such

phages can enter the lytic cycle spontaneously or following cell stress, and reports indicate that

lysogenic phages can be continuously shed from cells with little to no reduction in viability of the

overall bacterial population[59, 60]. During the analysis of bacterial colonies originating from *E.*

faecalis 19RS strains, we discovered that the phi19 genome is maintained in these phage-

resistant bacteria, and is not integrated in the bacterial chromosome. This carriage of the phi19

genome may explain its ability to influence IS256 transposition. We found that 19RS strains

continually release infectious phage particles with little impact on the viability of the bacterial

population (Fig. 5A). These particles were confirmed to be phi19 using PCR (Fig. 5B). We

discovered that ~10% of 19RS colonies produced infectious phages (Fig. 5C). This suggests

that a minority of each population exerts a selective pressure to maintain phage resistance by

producing infectious phage progeny. Loss of phage resistance in this population would lead to

phage infection and population collapse. Colonies that produced zones of clearing were isolated

and serially passaged for three passages. All cells that initially produced zones of clearing

continued to produce zones for all three passages (Fig. S6), demonstrating that phi19 is stably

carried in this subpopulation of cells. Finally, we reanalyzed reads from our WGS experiments

and found that we could recover the phi19 genome with high coverage from 19RS strains (Fig.

5D). Wild type *E. faecalis* V583 did not have any detectable phi19 genomic DNA. Collectively,

these results show that 19RS strains release phages from a subpopulation of cells, which may

impose stress on *E. faecalis* influencing IS256 expansion.

Considering that 19RS strains have IS256-rich genomes and release infectious phages,

we hypothesized that phage carriage may increase the competitiveness of these 19RS strains

when co-cultured with phage sensitive strains. We performed competition assays where *E.*

faecalis V583 WT or 19RS strains were competed against *E. faecalis* OG1RF, which is sensitive

to phi19 infection, or an OG1RF Δ*epaOX* mutant strain (homologous to *E. faecalis* V583 *epaX*)

[61] that is resistant to phi19 infection, without the addition of exogenous phages to the cultures

(Fig. 6). 19RS strains that were shedding phi19 outcompeted WT *E. faecalis* OG1RF. However,

they were unable to outcompete *E. faecalis* OG1RF Δ*epaOX*. These results demonstrate that

phi19 carriage provides a selective advantage to *E. faecalis* cells when competing with related

bacteria that are phage susceptible but if competitors are phage resistant, phi19 carriage comes

with a significant fitness cost.

IS256 expansion occurs in enterococcal genomes during human infection.

IS elements contribute to genome diversification and adaptive plasticity that is important

for bacterial evolution[62]. Environmental pressure is a strong driver of IS element mobility[63, 64, 65],

and the work described here expands this knowledge to phage predation. Antibiotics are a

strong selective pressure that can promote bacterial evolution, and IS256 insertions have been

tied to pathoadaptation in enterococci during human infections[36]. To assess if environmental

cues such as antibiotics may be driving IS element mobility, we investigated whether

enterococcal populations within a single individual infected with *E. faecium* experience IS256

expansion during therapeutic intervention. Stools samples were collected at various timepoints

over 109 days, during which the individual was treated with regimens containing daptomycin,

vancomycin, and the vancomycin analogue oritavancin for recurrent *E. faecium* bacteremia. IS-

Seq was performed on these samples across the time series described above (Fig. 7A). Nearly

all of the insertions found using IS-Seq occurred in contigs which were likely generated from *E.*

faecium, rather than *E. faecalis* (Fig. S7). We found that these *E. faecium* populations had a

major expansion of IS256 insertions following oritavancin therapy compared to daptomycin

and/or vancomycin therapy. Comparing differentially abundant insertions between vancomycin-

containing and oritavancin treatment revealed multiple enriched insertions in the coding

sequences of a putative Rib transcriptional regulator, and the *vex/vncRS* operon also identified

as a IS256 insertion hotspot in *E. faecalis* 19RS strains (Fig. 7B-C). These two operons are

likely involved in antibiotic resistance and virulence in *E. faecium* and IS256 insertions likely

render these genes nonfunctional. These insertions increase in sequencing depth during the

course of treatment and peak during oritavancin administration. The *vex/vncRS* operon in *E.*

faecalis V583 contains an IS element belonging to the ISL3 family[26], indicating that the genome

evolution of *E. faecium* during blood stream infection may follow a similar trajectory to *E.*

faecalis V583[26]. We also found that IS256 regularly interrupts an aminoglycoside resistance

operon (Fig. S8). Oscillating insertions both in coding and non-coding regions in this operon

suggest that this region may be an IS256 insertion hotspot. Lastly, we found that oritavancin-

treated samples had an increase in IS256 circle formation indicating that in addition to novel

insertion events, active IS256 transposition is occurring during blood stream infection (Fig. 7D).

Overall, these findings suggest that IS256 transposition is frequent in the intestine during *E.*

faecium infection, and this transposition likely impacts the antibiotic resistance and virulence

profiles of the infecting bacterial population.

Discussion

IS elements directly shape bacterial physiological responses involved in antibiotic

resistance and virulence, and are key drivers of bacterial genome evolution[66, 67, 68]. Considering

the importance of IS element biology to bacterial fitness, the selective pressures that guide IS

element regulation and mobility remain poorly defined. Using the Gram-positive opportunistic

pathogen *E. faecalis,* we extend our knowledge of IS elements by detailing the population

biology, regulation, and expansion dynamics of IS256. IS256 is a widely distributed IS element

in Gram-positive pathogens. Through the use of traditional bacterial genetics, comparative

genomics, and IS-Seq we have revealed that the dynamics of IS256 insertion events are

strongly dictated by phage predation and the mammalian host environment. Together these

results suggest that IS256 mobility creates a genetically flexible genome that enables

enterococci to rapidly adapt to diverse environmental conditions.

IS256 is common in *E. faecalis* genomes and is found more frequently in hospital

adapted lineages (Fig. 1). Consistent with this idea, IS256 element abundance is a defining

feature of hospital adapted *E. faecium* isolates[43, 44]. Movement of IS elements between cells

requires other mobile elements, such as phages and plasmids, as IS elements lack any

transforming or transducing ability[27]. The pattern of intermittent IS256 abundance in the

enterococci is likely due to periodic exposure to IS256-containing mobile elements. In other

words, genetically-related strains may interact with IS256-containing mobile elements at

different rates, resulting in the dissemination of IS256 to only a subset of the population. The

enrichment of IS256 among hospital adapted strains may reflect the need for increased

genomic plasticity in these strains as they adapt to an altered host environment[44, 69]. Pathogens

can utilize IS elements to form pseudogenes and to condense genome size through IS-

mediated recombination, which can inhibit immune detection through loss of virulence factors

and reduces the genome's metabolic burden[68, 69].

Regulation of IS256 has not been investigated in the enterococci and is key to

understanding this element's biology. Here we show that IS256 is basally active in *E. faecalis*

V583, yet it can be regulated at both the transcriptional and translational level (Fig. 2B-D). This

is important because overactive IS256 could be catastrophic to *E. faecalis* viability and layers of

regulation likely temper the elements activity until appropriate conditions require IS256

activation. Additionally, we show that IS256 has two different TSSs as part of this regulation and

that both sites are regulated by asRNAs. The protein product occurring from the site-2 TSS is

predicted to be truncated and it is unknown if this IS256 variant would be active. Other IS

elements encode two or more protein coding frames and these different protein products can

contribute to IS element regulation through inhibition[67, 70]. Using IS-Seq as a readout of basal

IS256 mobility, we show that IS256 insertions function as a mechanism of genome-wide

mutation (Fig. 2A). This is the highest resolution IS-Seq map created, to our knowledge, and

demonstrates that this technique can identify low-abundance insertions with high confidence[50,

51]. Baseline IS256 movement likely adds mutational diversity to an enterococcal population

which allows for adaption to changing environmental conditions through selection.

Previous work from our group determined that *E. faecalis* infection by the phage phi19

was associated with an expansion of IS256 insertions[18] (Fig. 3A). This was found using WGS

that was based on less than 10 reads per insertion. We extended this observation by performing

IS-Seq to accurately identify these novel insertion sites both in vitro and in vivo. We found that

E. faecalis challenged with phi19 (19RS strains) had a major increase in new IS256 insertions

(Fig. 4A-B). 19RS strains were chronically infected by phi19 and cells within the population

carried the phi19 genome and this chronic infection may provide a continuous selective

pressure that aids in IS256 mobility. This correlated with 19RS strains harboring elevated IS256

circularization intermediates (Fig. 4C). These circles are formed during IS256 copy-paste

transposition and have been studied extensively[30].

We also observed that in a mouse model of *E. faecalis* V583 intestinal colonization,

exposure to phi19 resulted in a similar expansion of IS256 (Fig. 4D-F). IS256 expansion was

less robust in the mouse intestine compared to the levels of expansion seen in vitro. This may

be for several reasons, including a lower frequency of insertion events per location in vivo and a

lower number of samples sequenced. An additional explanation is that Epa is crucial for

bacterial colonization and *epa* mutations may lead to these mutants being outcompeted in the

intestine[18, 71].

Although we found evidence for phage-mediated IS256 activation in *E. faecalis* 19RS

strains, we also determined that selection is important for the location of IS256 expansion (Fig.

4G-H). Cells that were not exposed to phi19 had lower IS256 read density in *epaX*. *EpaX* is a

glycosyltransferase involved in teichoic acid decoration of the core rhamnose containing Epa

exopolysaccharide backbone. Mutations in *epaX* result in poor phage adsorption[18]. We found no

other insertions in genes known to be involved in phage infection indicating that *epaX* and other

epa genes are likely hotspots for IS256 integration following phage pressure. In light of these

discoveries, we hypothesize that initial phage infection may select for *epa*-IS256 insertions and

that prolonged phage exposure though chronic infection leads to increased and non-specific

IS256 transposition as a stress response. We suspect that phage carriage and release will

continually pressure intermittently resistant cells and force fully resistant cells to maintain high

levels of resistance.

IS elements are involved in pseudogene formation during pathoadaption. Examples

include IS256 insertions in the cytolysin operon of *E. faecalis* V583[36] and a dependence on IS

insertions in *E. coli* to evolve resistance to macrophage killing[72]. Additionally, how enterococci

evolve during human infection is not well understood and may involve a variety of genetic

changes in cell wall modifying[73] and cellular respiration[74] genes. While IS elements are directly

involved in enterococcal pathoadaptation [36], an in-depth analysis of IS-mediated mutations

during human infection had never been performed. Previously, such dynamics were studied by

laboriously sequencing individual isolates. While this approach can successfully identify new IS

element insertion sites, it overlooks IS element mobility within the entirety of the bacterial

population. Here, we show that IS-Seq can be used to study bacterial pathoadaptation at the

population level and can reveal multiple pathogen-associated insertions within such a

population (Fig. 7). Following oritavancin therapy an increase in IS256 insertions occurred in the

vex/*vncRS* operon, which has been implicated in both virulence[56] and vancomycin resistance[55].

Additionally, there is an increase in insertions in a putative transcriptional regulator that may

regulate a Rib/Alpha adhesin[75, 76]. Loss of these potential virulence factors may lead to

modulation of virulence and avoidance of immune activation or detection. Additionally, we found

highly abundant IS256 insertions in an aminoglycoside acetyltransferase. Vancomycin is

reported to act synergistically with gentamicin against enterococci[77], likely due to increased

gentamicin transport through the vancomycin-permeabilized cell wall[78]. In recent years,

enterococci with aminoglycoside resistance have emerged[79] with acquisition of resistance

genes carried in an IS256 composite transposon[80]. Our results suggest that genetic modulation

of aminoglycoside resistance fluctuates in the host during infection, with different highly

abundant insertions prominent at different timepoints. This data suggests that aminoglycoside

resistance is repeatedly modulated and the intestinal enterococcal population regularly samples

different versions of this aminoglycoside resistance operon. Future studies to identify if

glycopeptide treatment potentiates aminoglycoside sensitivity through IS256 insertions is

warranted. We additionally show that oritavancin treatment is associated with IS256 expansion

in this patient. Oritavancin is an analog of vancomycin, but is also reported to disrupt cell

membrane integrity and possess additional mechanisms for inhibiting cell wall synthesis[81]. This

may explain the difference in IS256 expandability, as this antibiotic may impose stronger

pressure resulting in a broader range of mutations. Likewise, oritavancin treatment was

associated with an increase in IS256 circles. In *Staphylococcus aureus*, vancomycin treatment

increases IS256 transposition[35, 82], reflecting a similar glycopeptide-induced IS256 activation.

In summary, we show how the widespread and clinically-relevant IS element IS256

creates genome diversification during phage predation and human infection using sensitive

NGS IS-Seq. This work provides a high-resolution picture of IS256 movement during both

steady-state and stress inducing conditions. Understanding the outcomes of these IS256

insertion events will be critical to the study of the evolution of enterococcal pathogenesis in

response to both phage and antibiotic therapies. We also show that IS256 is regulated at both

the transcriptional and translational level in *E. faecalis* and have discovered that IS256 mobility

is biased for highly mobilizable insertion events in nosocomial and virulent enterococcal

isolates, further demonstrating that IS elements are domesticated and specialized in their choice

of bacterial host.

Materials and Methods

Bacteria and bacteriophages.

E. faecalis strains were grown in BHI broth with aeration at 37°C. A list of bacterial and

bacteriophage strains can be found here (Table S4). *Escherichia coli* cultures were grown in

Lennox L broth at 37°C with aeration. 15 µg/mL or 5 µg/mL chloramphenicol was used for

selection of transformed *E. coli* and *E. faecalis*, respectively. The generation of phage-resistant

19RS strains was recently described[18].

Enterococcal phylogeny and Identification of IS256 in enterococcal genomes

All available *E. faecalis* and *E. faecium* genomes were downloaded from NCBI RefSeq

on 3/21/2022. Phylogenetic tree was constructed with GToTree v1.6.31 using default

parameters[83], pruned using treeshrink v1.3.9[84], and visualized using the iTOL viewer[85]. IS256

transposase genes were identified using GToTree with the IS256 pfam "PF00872" and 100%

amino acid matches to the IS256 Tnp from *E. faecalis* V583 were found using seqkit v0.16.1[86].

MLST and virulence factor abundance

Enterococcal genomes obtained from RefSeq were assigned an MLST type using mlst

v2.19.0[87] with the *E. faecalis* and *E. faecium* MLST scheme from pubmlst[88]. Virulence factors

were identified by calling all predicted ORFs in all downloaded RefSeq genomes using prodigal

v2.6.3[89] and running diamond blastp v0.9.21[90] against this database with known *E. faecalis* and

E. faecium virulence factors from the Virulence Factor Database[91]. All further analyses and

statistics were performed in R.

DNA extraction and IS-Seq

Bacterial genomic DNA was extracted from overnight cultures grown in 5 mL BHI broth

using the Zymo BIOmics DNA Miniprep Kit (Zymo, Irvine, CA). For library preparation, 200 ng of

genomic DNA was used as input in the Illumina DNA Prep Kit following the manufacturer's

instructions (Illumina, San Diego, CA). Following library construction, IS256 enrichment was

performed as follows. First, 10 ng of library DNA was amplified with the p7 primer and the IS-

Seq Step1 primer (Table S5) for 13 cycles using Q5 2X Master Mix (NEB, Ipswich, MA). The

products of this reaction were diluted 1:100 and 10 µL was added to a PCR reaction with the p7

primer and the IS-Seq Step2 primer for 9 cycles using Q5 2X Master Mix. The products of this

second reaction were sequenced on an Illumina NovaSEQ 6000 by the University of Colorado

Anschutz Medical Center Genomics Core with paired-end 150 cycle chemistry. 23.3 ± 3.8 M

reads, 17.9 ± 2.5 M reads, and 16.3 ± 2.6 M reads were obtained from sequencing the human,

mouse, and *in vitro* IS-Seq libraries, respectively. IS-seq reads have been deposited in the

European Nucleotide Archive under project accession number PRJEB55280.

IS-Seq analysis

Paired end reads obtained from IS-Seq were partitioned into read1 and read2 files, and

reads from read1 files were used in the downstream analysis. First, cutadapt v1.18 was used to

bin reads with a full IS256 5' terminus. the IS256 5' terminus was then trimmed from these

reads[92] suing cutadapt with the following flags: '-O 6 -g

"^CGTAAAAGGACTGTTATATGGCCTTTTTACTTTTACACAATTATACGGACTTTATC" '.

These trimmed reads were aligned to the *E. faecalis* V583 chromosome using Bowtie2

v2.3.5.1[93] and the location of each insertion was determined using samtools v1.13[94] and

bedtools genomecov v2.30.0[95]. All downstream analysis was performed in R. Insertion site

abundances were normalized using DESeq2 estimateSizeFactors[96] and significantly enriched

unique insertion sites were found using DESeq2. Both significantly enriched and insignificant

insertions with at least 100 reads were reported in tables and figures.

E. faecalis phi19 infection in the mouse intestine

All animal protocols were approved by the Institutional Animal Care and Use Committee

of the University of Colorado School of Medicine (protocol number 00253). Conventional 6-

week-old C57BL/6 mice were divided into two groups (Control: 2 female/2 male, Experimental: 2

female/2male). All mice were treated with an antibiotic cocktail (streptomycin [1 mg/ml],

gentamicin [1 mg/ml], erythromycin [200 µg/ml]) by first dosing via oral gavage with 100 µl of

the cocktail and replacing their drinking water with the same antibiotic cocktail for 1 week. On

day 7, water bottles were replaced with antibiotic free-drinking water. After 24 hours mice were

colonized with *E. faecalis* V583 suspended in phosphate buffered saline (PBS) by oral gavage

(10^9 CFU in 100 µl). After 24 hours mice were orally gavaged daily as follows: 100 µl of 1M

sodium bicarbonate (all mice) and 100 µl PBS per control mouse or 10^9 PFU phi19 in 100 µl per

experimental mouse. Feces were collected daily and homogenized in 1ml PBS. 10µl of the fecal

slurry was serially diluted and plated on BHI agar containing 100 µg/ml gentamicin or 100 µg/ml

gentamicin and 10^8 pfu/ml phi19. From the experimental samples, 5 µl of slurry was mixed with

45µl chloroform, spun at 16,363 RCF for 1 min and the supernatant was enumerated for phage

particles by agar overlay plaque assay as described previously[19].

RNA extraction and qPCR

RNA was isolated from logarithmically growing (OD=0.3) *E. faecalis* using a modified

protocol for the RNeasy Mini-Prep Kit (Qiagen, Hilden, Germany). First, cell pellets were

collected by centrifugation of 5 mL of cell culture at 8,228 RCF for 5 minutes. Cell pellets were

resuspended in 1 mL RNAlater and centrifuged again at 8,228 RCF for 10 minutes. The cell

pellets were stored at -80 C until sample processing. To isolate RNA, the pellets were thawed

and resuspended in 100 µL TE buffer with 15 mg/mL lysozyme and incubated at room

temperature for 30 minutes. Following this, 700 µL Buffer RLT with 0.01% beta-mercaptoethanol

was added and the sample was transferred to a Lysing Matrix B tube (MPBio, Irvine, CA). The

tubes were bead-beaten in a Mini-Beadbeater-16 (Biospec Products, Bartlesville, OK) on the

fastest setting at 30-second intervals with 1 minute rests on ice between cycles. Following bead

beating, the supernatant was removed after centrifugation at 16,363 RCF for 30 seconds and

processed following the manufacturer's instructions. RNA was eluted from the column in

RNase/DNase free H_2O and any residual DNA contamination was removed with a 1 hour off-

column DNase treatment (Qiagen). The RNA was purified following the RNeasy Mini-Prep Kit

following manufacturer's instructions. cDNA was synthesized using Qscript Master Mix

(Quantabio, Beverley, MA) and 1 µg RNA. For qPCR of gDNA targets, 1 ng of gDNA template

was used per reaction. qPCR was run with PowerUp SYBR Green Master Mix (Thermo Fisher,

Waltham, MA) on an Applied Biosystems QuantStudio 7 Flex qPCR system. qPCR primers are

listed in Table S5.

DNA manipulation and cloning

535   All cloning primers and DNA constructs used in this study are listed in Table S5. All

536   plasmid constructs used the pLZ12 shuttle vector backbone[97]. For cloning, all inserts were

537   amplified using 2X Q5 DNA polymerase Master Mix and assembled using 2X Gibson Assembly

538   Master Mix (NEB).

540   *Phage carriage identification and serial passaging*

541   To enumerate extracellular viral particles, we collected supernatant from cultures after

centrifugation for 1 minute at 10,000 RCF, treated with 1/10 volume chloroform, centrifuged for

1 min at 21,000 RCF and extracted the aqueous phase. The resulting supernatant was serially

diluted in SM+ buffer (50 mM Tris-HCl, 100 mM NaCl, 8 mM $MgSO_4$, 5 mM $CaCl_2$ [pH 7.4]),

and 10 µL was mixed with 130 µL of a 1:10 dilution of *E. faecalis* V583 cells grown overnight.

This mixture was combined with 5 mL of 0.35% Todd Hewitt agar (THA) supplemented with

10mM $MgSO_4$, and poured onto THA+10mM $MgSO_4$ plates.

To identify 19RS *E. faecalis* cells actively shedding phages, glycerol stocks of phage-

resistant cultures were streaked on BHI agar and incubated at 37°C. After overnight growth, 182

µL of a 1:10 dilution in SM+ of an overnight wild type *E. faecalis* V583 culture was mixed with 7

mL of 0.35% Todd Hewitt agar (THA) and 10mM $MgSO_4$ and solidified on THA+10mM $MgSO_4$

plates for 30 minutes. Individual colonies were picked from the overnight plate patched onto the

plate containing *E. faecalis* V583 embedded in 0.35% THA, with care not to break the surface of

the agar, and incubated at 37°C overnight.

To serially passage phage-positive colonies, colonies producing zones of clearing were

patched onto both THA+10mM $MgSO_4$ plates and THA+10mM $MgSO_4$ with an *E. faecalis* V583

top agar layer as described above. The next day, the colony on the THA+10mM $MgSO_4$ plate

was patched onto both types of plates and this was continued for two subsequent passages.

Competition assays

E. faecalis cultures were grown overnight in BHI broth and diluted to an OD_{600} of 1 in PBS. 30

µL of each strain was added to 3 mL BHI and incubated at 37°C for 24 hours. The cultures were

plated on both BHI gentamicin 100 µg/mL to select for *E. faecalis* V583 strains and BHI

rifampicin 25 µg/mL, fusidic acid 50 µg/mL to select for *E. faecalis* OG1RF or E. faecalis

OG1RF Δ*epaOX*.

Enterococcal isolation from stool samples and gDNA extraction

Enterococcal stool populations were obtained over a span of 109 days from a patient being

treated for recurrent *E. faecium* bacteremia. Stool samples were diluted and plated onto bile

esculin azide – Enterococcosel agar (Becton Dickinson, Franklin Lakes, NJ), and 100-1,000

colonies were collected and pooled into BHI containing 16.7% glycerol and stored at -80°C. One

mL aliquots of the frozen population stocks were thawed and washed with BHI. Populations

were allowed to briefly expand at 37°C at 170 rpm for 5.5 hours. The cultures were pelleted and

genomic DNA was extracted into DNase free water using the DNeasy Blood and Tissue Kit

(Qiagen). gDNA concentrations were determined with the Qubit 1X dsDNA High-Sensitivity Kit

and fluorometer (Invitrogen, Thermo Fisher).

Statistics and Visualization

Graphs were made in Graphpad Prism or in R using dplyr and ggplot2. T-tests and one-way

ANOVAs were calculated in Graphpad Prism and insertion site differential abundance and

proportional tests (prop.test) were calculated in R.

Acknowledgements

This work was supported by National Institutes of Health grants R01AI141479 (B.A.D.),

R01AI165519 (D.V.T.), R01AI116610 (K.L.P.), T32AI052066 (J.M.K), F31AI169976 (J.M.K.),

and T32AI138954 (M.E.S.).

Figure Legends

Figure 1. Distribution of IS256 in *E. faecalis* and *E. faecium*. A) Phylogenetic tree of *E.*

faecalis genomes. Genomes with IS256 are labeled in red and genomes without IS256 are

labeled in black. **B)** Presence or absence of IS256 in the genomes of hospital adapted *E.*

faecalis MLST types. **C)** Comparison of *E. faecalis* virulence factor abundances in the genomes

of hospital adapted *E. faecalis* MLST types. **D)** Phylogenetic tree of *E. faecium genomes.*

Genomes with IS256 are labeled in red and genomes without IS256 are labeled in black. **E)**

Presence or absence of IS256 in the genomes of hospital adapted *E. faecium* MLST types. **F)**

Comparison of *E. faecium* virulence factor abundances in the genomes of hospital adapted *E.*

faecium MLST types. **G)** Percent of *E. faecalis* and *E. faecium* genomes from RefSeq that

contain IS256. (* $p<0.05$, ** $p<10^{-5}$, *** $p<10^{-25}$, determined with chi-squared test)

Figure 2. Regulation of IS256 Tnp activity during steady state conditions. A) IS256

sequencing (IS-Seq) from three individual *E. faecalis* V583 cultures. Peaks indicated in black

represent the six endogenous IS256 insertions identified in the sequenced genome of V583. **B)**

qPCR of IS256 *tnp* gene expression relative to the housekeeping protease gene *clpX*. **C)**

Stranded RNA-Seq of IS256 tnp gene expression in *E. faecalis* V583. **D)** IS256 Tnp protein

expression indicated by measuring GFP fluorescence of an IS256-GFP and IS256-GFP

ΔasRNA translational fusions. (* $p<0.05$, determined with multiple t-tests)

Figure 3. IS256 expansion confers phage resistance and is elevated in *E. faecalis* isolates

resistant to phi19. A) Enumeration of predicted IS256 insertions determined from whole

genome sequencing of *E. faecalis* phage resistant 19RS strains. **B)** PCR showing that IS256

insertions occur in both the positive and negative strand orientations in the *epaX* gene 19RS

strains. **C)** Southern blot analysis of global IS256 insertions in wild type (WT) *E. faecalis* V583

and 19RS strains. Red arrows indicate novel IS256 banding patterns. **D)** Emergence of phage

resistant *E. faecalis* OG1RF with IS256 or a catalytically dead (ΔDDE) IS256. EV indicates *E.*

faecalis OG1RF carrying the empty vector. **E)** qPCR quantification of IS256 circular

intermediates normalized to total cell density following phi19 infection or mock infection (MI). (*

$p<0.05$, ** $p<0.01$, determined with multiple t-tests).

Figure 4. IS-Seq to identify IS256 insertion sites in *E. faecalis* phage resistant isolates in

vitro and in vivo. A) IS-Seq of *E. faecalis* V583, and 19RS strains. **B)** Quantification of

significantly enriched insertions between pairwise comparisons of each group in A. In each

square, the top value is the number of insertions overabundant in the column group and the

bottom value is number of insertions overabundant in the row group. **C)** IS256 circle formation

determined with IS-Seq from samples shown in A. **D)** IS-Seq of E. faecalis V583 and phage

resistant colonies isolated from the feces of mice colonized with (Mouse 5 & 6) and without

(Mouse 1 & 2) oral phi19 inoculation. **E)** Quantification of significantly enriched insertions

between pairwise comparisons of each group in D. In each square, the top value is the number

of insertions overabundant in the column group and the bottom value is number of insertions

overabundant in the row group. **F)** IS256 circle formation determined with IS-Seq from samples

shown in D.

Figure 5. *E. faecalis* 19RS strains chronically shed phi19 virus and harbor the phage

genome intracellularly. A) Quantification of shed infectious phi19 phage particles. **B)** PCR

confirmation of phi19 recovered from culture fluid of *E. faecalis* 19RS strains. **C)** Proportion of

cells actively replicating and releasing infectious phi19 phage particles. **D)** WGS read mapping

alignments to the phi19 genome from *E. faecalis* V583 and 19RS strains.

Figure 6. phi19 shedding *E. faecalis* 19RS strains are more fit during competition with a

phage susceptible *E. faecalis* strain. Competitive fitness of *E. faecalis* V583 and 19RS

strains relative to *E. faecalis* OG1RF or an *E. faecalis* OG1RF derivative with a deletion in the

epaX homolog *epaOX*.

Figure 7. IS-Seq of a longitudinal series stool samples from a human patient with a

vancomycin resistant enterococcal infection. A) Insertions found via IS-Seq plotted on a

concatenated "metagenome" of contiguous enterococcal sequences from stool genomic DNA.

IS256 sequences were mapped against individual contigs to identify IS256 insertion locations.

Then all contigs were concatenated together to build a linearized representation of the

metagenome. **B)** Volcano plot of differentially abundant insertions during

Daptomycin+Vancomycin co-therapy and Oritavancin therapy. Enriched insertions (dark red)

were overabundant during Oritavancin therapy. **C)** Gene organization of loci where elevated

IS256 insertions occurred are indicated by arrows. Lollipops show the location of IS256 insertion

and bar graphs are read depth abundances found for each insertion. (* $p < 0.05$, **, $p < 0.01$, ***,

$p < 0.001$, determined with DESeq2) **D)** IS256 circle formation for each treatment group inferred

from IS-Seq.

Figure S1. Enrichment of IS256-chrmosomal junctions during IS-Seq. A) Schematic of the

IS-Seq PCR amplification step. **B)** Trimming site and binning strategy used to identify IS256-

termini reads.

Figure S2. Steady state IS256 insertion sites in the native *E. faecalis* V583 plasmids. A)

pTEF1, **B)** pTEF2, and **C)** pTEF3.

Figure S3. IS256 insertions in *E. faecalis* V583 plasmids in vitro and in vivo. A-C) Insertion

sties from in vitro cultured *E. faecalis* 19RS strains; **A)** pTEF1, **B)** pTEF2, and **C)** pTEF3. **D-F)**

Insertion sites from *E. faecalis* isolated from the murine intestine **D)** pTEF1, **E)** pTEF2, and **F)**

pTEF3.

Figure S4. IS-256 insertions in *epa* and *vex/vnc* genes in vitro and in vivo. Insertion sties

from in vitro cultured *E. faecalis* 19RS strains in the **A)** *epa* locus and **B)** *vex/vnc* locus. Insertion

sites from E. faecalis isolated from the murine intestine in the **C)** *epa* locus and **D)** *vex/vnc*

locus.

Figure S5. Murine model and intestinal *E. faecalis* colonization and oral phi19 treatment.

A) Schematic of the *E. faecalis* intestinal colonization model used to identify in vivo IS256

mobilization. **B)** Colony forming units of *E. faecalis* isolated from mouse feces. Bacteria were

enumerated on growth media supplemented with gentamicin with and without the addition of

phi19 to measure the frequency of phage resistant colonies. Arrows indicate time points where

IS-Seq was performed.

Figure S6. Serial passage of phi19 shedding *E. faecalis* 19RS isolates.

Figure S5. IS256 read mapping to enterococcal genomic contigs assembled from stool

samples. Each data point indicates the read mapping abundance per assembled contig. Data

points indicated as *E. faecalis* or *E. faecium* lacked sufficient data to definitively categorize

these contigs as either *E. faecalis* or *E. faecium*.

Figure S8. IS256 insertions from human stool enterococcal isolates in genes involved in

aminoglycoside resistance.

References

1.   Fiore E, Van Tyne D, Gilmore MS. Pathogenicity of enterococci. *Microbiol Spectr* **7**,
(2019).

679
680 2.   Shah ASV*, et al.* Incidence, microbiology, and outcomes in patients hospitalized with
681   infective endocarditis. *Circulation* **141**, 2067-2077 (2020).

682
683 3.   Khan A, Aslam A, Satti KN, Ashiq S. Infective endocarditis post-transcatheter aortic
684   valve implantation (TAVI), microbiological profile and clinical outcomes: A systematic
685   review. *PLoS One* **15**, e0225077 (2020).

686
687 4.   CDC. 2019 AR Threats Report.) (2019).

688
689 5.   Markwart R*, et al.* The rise in vancomycin-resistant *Enterococcus faecium* in Germany:
690   data from the German Antimicrobial Resistance Surveillance (ARS). *Antimicrob Resist*
691   *Infect Control* **8**, 147 (2019).

692
693 6.   Cetinkaya Y, Falk P, Mayhall CG. Vancomycin-resistant enterococci. *Clin Microbiol Rev*
694   **13**, 686-707 (2000).

695
696 7.   Partridge SR, Kwong SM, Firth N, Jensen SO. Mobile genetic elements associated with
697   antimicrobial resistance. *Clin Microbiol Rev* **31**, (2018).

698
699 8.   Schwarz FV, Perreten V, Teuber M. Sequence of the 50-kb conjugative multiresistance
700   plasmid pRE25 from *Enterococcus faecalis* RE25. *Plasmid* **46**, 170-187 (2001).

702 9.   Hodel-Christian SL, Murray BE. Characterization of the gentamicin resistance
703   transposon Tn5281 from *Enterococcus faecalis* and comparison to staphylococcal
704   transposons Tn4001 and Tn4031. *Antimicrob Agents Chemother* **35**, 1147-1152 (1991).

705
706 10. Arthur M, Molinas C, Depardieu F, Courvalin P. Characterization of Tn1546, a Tn3-
707   related transposon conferring glycopeptide resistance by synthesis of depsipeptide
708   peptidoglycan precursors in *Enterococcus faecium* BM4147. *J Bacteriol* **175**, 117-127
709   (1993).

710
711 11. Quintiliani R, Jr., Courvalin P. Characterization of Tn1547, a composite transposon
712   flanked by the IS16 and IS256-like elements, that confers vancomycin resistance in
713   *Enterococcus faecalis* BM4281. *Gene* **172**, 1-8 (1996).

714
715 12. Kortright KE, Chan BK, Koff JL, Turner PE. Phage therapy: a renewed approach to
716   combat antibiotic-resistant bacteria. *Cell Host Microbe* **25**, 219-232 (2019).

717
718 13. Chan BK, Turner PE, Kim S, Mojibian HR, Elefteriades JA, Narayan D. Phage areatment
719   of an aortic graft infected with *Pseudomonas aeruginosa*. *Evol Med Public Health* **2018**,
720   60-66 (2018).

721
722 14. Schooley RT*, et al.* Development and use of personalized bacteriophage-based
723   therapeutic cocktails to treat a patient with a disseminated resistant *Acinetobacter*
724   *baumannii* infection. *Antimicrob Agents Chemother* **61**, (2017).

725
726 15. Dedrick RM*, et al.* Engineered bacteriophages for treatment of a patient with a
727   disseminated drug-resistant *Mycobacterium abscessus*. *Nat Med* **25**, 730-733 (2019).

728
729 16. Aslam S*, et al.* Lessons learned from the first 10 consecutive cases of intravenous
730   bacteriophage therapy to treat multidrug-resistant bacterial infections at a single center
731   in the United States. *Open Forum Infect Dis* **7**, ofaa389 (2020).

732
733 17. Petrovic Fabijan A, Lin RCY, Ho J, Maddocks S, Ben Zakour NL, Iredell JR. Safety of
734   bacteriophage therapy in severe *Staphylococcus aureus* infection. *Nat Microbiol* **5**, 465-
735   472 (2020).

736
737 18. Chatterjee A*, et al.* Bacteriophage resistance alters antibiotic-mediated intestinal
738   expansion of enterococci. *Infect Immun* **87**, (2019).

739
740 19. Duerkop BA, Huo W, Bhardwaj P, Palmer KL, Hooper LV. Molecular basis for lytic
741   bacteriophage resistance in enterococci. *mBio* **7**, (2016).

742
743 20. Markwitz P, Lood C, Olszak T, van Noort V, Lavigne R, Drulis-Kawa Z. Genome-driven
744   elucidation of phage-host interplay and impact of phage resistance evolution on bacterial
745   fitness. *ISME J* **16**, 533-542 (2022).

747   21.   Hesse S*, et al.* Phage resistance in multidrug-resistant *Klebsiella pneumoniae* ST258
748   evolves via diverse mutations that culminate in impaired adsorption. *mBio* **11**, (2020).

750   22.   Woodford N, Ellington MJ. The emergence of antibiotic resistance by mutation. *Clin*
751   *Microbiol Infect* **13**, 5-18 (2007).

753   23.   Mangalea MR, Duerkop BA. Fitness trade-offs resulting from bacteriophage resistance
754   potentiate synergistic antibacterial strategies. *Infect Immun* **88**, (2020).

756   24.   Gordillo Altamirano F*, et al.* Bacteriophage-resistant *Acinetobacter baumannii* are
757   resensitized to antimicrobials. *Nat Microbiol* **6**, 157-161 (2021).

759   25.   Castledine M*, et al.* Parallel evolution of Pseudomonas aeruginosa phage resistance
760   and virulence loss in response to phage treatment in vivo and in vitro. *eLife* **11**, e73679
761   (2022).

763   26.   Paulsen IT*, et al.* Role of mobile DNA in the evolution of vancomycin-resistant
764   *Enterococcus faecalis. Science* **299**, 2071-2074 (2003).

766   27.   Johnson CN, Sheriff EK, Duerkop BA, Chatterjee A. Let me upgrade you: impact of
767   mobile genetic elements on enterococcal adaptation and evolution. *J Bacteriol* **203**,
768   e0017721 (2021).

770   28.   Clewell DB, Weaver KE, Dunny GM, Coque TM, Francia MV, Hayes F.
771   Extrachromosomal and mobile elements in enterococci: transmission, maintenance, and
772   epidemiology. In: *Enterococci: From Commensals to Leading Causes of Drug Resistant*
773   *Infection* (eds Gilmore MS, Clewell DB, Ike Y, Shankar N). Massachusetts Eye and Ear
774   Infirmary (2014).

776   29.   Werner G, Fleige C, Geringer U, van Schaik W, Klare I, Witte W. IS element IS16 as a
777   molecular screening tool to identify hospital-associated strains of *Enterococcus faecium.*
778   *BMC Infect Dis* **11**, 80 (2011).

780   30.   Loessner I, Dietrich K, Dittrich D, Hacker J, Ziebuhr W. Transposase-dependent
781   formation of circular IS256 derivatives in *Staphylococcus epidermidis* and
782   *Staphylococcus aureus. J Bacteriol* **184**, 4709-4714 (2002).

784   31.   Valle J, Vergara-Irigaray M, Merino N, Penadés JR, Lasa I. sigmaB regulates IS256-
785   mediated *Staphylococcus aureus* biofilm phenotypic variation. *J Bacteriol* **189**, 2886-
786   2896 (2007).

788   32.   Kozitskaya S, Cho SH, Dietrich K, Marre R, Naber K, Ziebuhr W. The bacterial insertion
789   sequence element IS256 occurs preferentially in nosocomial *Staphylococcus*

790   *epidermidis* isolates: association with biofilm formation and resistance to
791   aminoglycosides. *Infect Immun* **72**, 1210-1215 (2004).

792
793   33.   Kleinert F*, et al.* Influence of IS256 on genome variability and formation of small-colony
794   variants in *Staphylococcus aureus*. *Antimicrob Agents Chemother* **61**, (2017).

795
796   34.   McEvoy CR*, et al.* Decreased vancomycin susceptibility in *Staphylococcus aureus*
797   caused by IS256 tempering of WalKR expression. *Antimicrob Agents Chemother* **57**,
798   3240-3249 (2013).

799
800   35.   Nagel M, Reuter T, Jansen A, Szekat C, Bierbaum G. Influence of ciprofloxacin and
801   vancomycin on mutation rate and transposition of IS256 in *Staphylococcus aureus*. *Int J*
802   *Med Microbiol* **301**, 229-236 (2011).

803
804   36.   Shankar N, Baghdayan AS, Gilmore MS. Modulation of virulence within a pathogenicity
805   island in vancomycin-resistant *Enterococcus faecalis*. *Nature* **417**, 746-750 (2002).

806
807   37.   Manson JM, Hancock LE, Gilmore MS. Mechanism of chromosomal transfer of
808   *Enterococcus faecalis* pathogenicity island, capsule, antimicrobial resistance, and other
809   traits. *Proc Natl Acad Sci U S A* **107**, 12269-12274 (2010).

810
811   38.   Leavis HL, Willems RJ, van Wamel WJ, Schuren FH, Caspers MP, Bonten MJ. Insertion
812   sequence-driven diversification creates a globally dispersed emerging multiresistant
813   subspecies of *E. faecium*. *PLoS Pathog* **3**, e7 (2007).

814
815   39.   Pöntinen AK*, et al.* Apparent nosocomial adaptation of *Enterococcus faecalis* predates
816   the modern hospital era. *Nat Commun* **12**, 1523 (2021).

817
818   40.   Dabul ANG*, et al.* Molecular basis for the emergence of a new hospital endemic
819   tigecycline-resistant *Enterococcus faecalis* ST103 lineage. *Infect Genet Evol* **67**, 23-32
820   (2019).

821
822   41.   Schell CM*, et al.* Detection of β-lactamase-producing *Enterococcus faecalis* and
823   vancomycin-resistant *Enterococcus faecium* isolates in human invasive infections in the
824   public hospital of Tandil, Argentina. *Pathogens* **9**, (2020).

825
826   42.   Zaheer R*, et al.* Surveillance of *Enterococcus* spp. reveals distinct species and
827   antimicrobial resistance diversity across a One-Health continuum. *Sci Rep* **10**, 3937
828   (2020).

829
830   43.   Lebreton F*, et al.* Emergence of epidemic multidrug-resistant *Enterococcus faecium* from
831   animal and commensal strains. *mBio* **4**, (2013).

833   44.   Chilambi GS, *et al.* Evolution of vancomycin-resistant *Enterococcus faecium* during
834   colonization and infection in immunocompromised pediatric patients. *Proc Natl Acad Sci*
835   *U S A* **117**, 11703-11714 (2020).

836
837   45.   Mikalsen T, *et al.* Investigating the mobilome in clinically important lineages of
838   *Enterococcus faecium* and *Enterococcus faecalis*. *BMC Genomics* **16**, 282 (2015).

839
840   46.   Perez M, *et al.* IS256 abolishes gelatinase activity and biofilm formation in a mutant of
841   the nosocomial pathogen *Enterococcus faecali*s V583. *Can J Microbiol* **61**, 517-519
842   (2015).

843
844   47.   Palmer KL, Daniel A, Hardy C, Silverman J, Gilmore MS. Genetic basis for daptomycin
845   resistance in enterococci. *Antimicrob Agents Chemother* **55**, 3345-3356 (2011).

846
847   48.   Lossouarn J, *et al. Enterococcus faecalis* countermeasures defeat a virulent picovirinae
848   bacteriophage. *Viruses* **11**, (2019).

849
850   49.   Dale JL, *et al.* Comprehensive functional analysis of the *Enterococcus faecalis* core
851   genome using an ordered, sequence-defined collection of insertional mutations in strain
852   OG1RF. *mSystems* **3**, (2018).

853
854   50.   Wright MS, Mountain S, Beeri K, Adams MD. Assessment of insertion sequence
855   mobilization as an adaptive response to oxidative stress in *Acinetobacter baumannii*
856   using IS-seq. *J Bacteriol* **199**, (2017).

857
858   51.   Reyes A, *et al.* IS-seq: a novel high throughput survey of in vivo IS6110 transposition in
859   multiple *Mycobacterium tuberculosis* genomes. *BMC Genomics* **13**, 249 (2012).

860
861   52.   Ellis MJ, Trussler RS, Haniford DB. A cis-encoded sRNA, Hfq and mRNA secondary
862   structure act independently to suppress IS200 transposition. *Nucleic Acids Res* **43**,
863   6511-6527 (2015).

864
865   53.   Michaux C, Hansen EE, Jenniches L, Gerovac M, Barquist L, Vogel J. Single-nucleotide
866   RNA maps for the two major nosocomial pathogens *Enterococcus faecalis* and
867   *Enterococcus faecium*. *Front Cell Infect Microbiol* **10**, 600325 (2020).

868
869   54.   Prudhomme M, Turlan C, Claverys JP, Chandler M. Diversity of Tn4001 transposition
870   products: the flanking IS256 elements can form tandem dimers and IS circles. *J*
871   *Bacteriol* **184**, 433-443 (2002).

872
873   55.   Haas W, Sublett J, Kaushal D, Tuomanen EI. Revising the role of the pneumococcal
874   *vex-vncRS* locus in vancomycin tolerance. *J Bacteriol* **186**, 8463-8471 (2004).

876 56. Lee S, *et al.* Induction of the pneumococcal *vncRS* operon by lactoferrin is essential for
877   pneumonia. *Virulence* **9**, 1562-1575 (2018).

878
879 57. Bolocan AS, *et al.* Evaluation of phage therapy in the context of *Enterococcus faecalis*
880   and its associated diseases. *Viruses* **11**, (2019).

881
882 58. Mäntynen S, Laanto E, Oksanen HM, Poranen MM, Díaz-Muñoz SL. Black box of
883   phage-bacterium interactions: exploring alternative phage infection strategies. *Open Biol*
884   **11**, 210188 (2021).

885
886 59. Liu Y, *et al.* Chronic release of tailless phage particles from *Lactococcus lactis*. *Appl*
887   *Environ Microbiol* **88**, e0148321 (2022).

888
889 60. Liu J, *et al.* The diversity and host interactions of *Propionibacterium acnes*
890   bacteriophages on human skin. *ISME J* **9**, 2078-2093 (2015).

891
892 61. Dale JL, Cagnazzo J, Phan CQ, Barnes AM, Dunny GM. Multiple roles for *Enterococcus*
893   *faecalis* glycosyltransferases in biofilm-associated antibiotic resistance, cell envelope
894   integrity, and conjugative transfer. *Antimicrob Agents Chemother* **59**, 4094-4105 (2015).

895
896 62. Touchon M, Rocha EP. Causes of insertion sequences abundance in prokaryotic
897   genomes. *Mol Biol Evol* **24**, 969-981 (2007).

898
899 63. Drevinek P, *et al.* Oxidative stress of *Burkholderia cenocepacia* induces insertion
900   sequence-mediated genomic rearrangements that interfere with macrorestriction-based
901   genotyping. *J Clin Microbiol* **48**, 34-40 (2010).

902
903 64. Ohtsubo Y, Genka H, Komatsu H, Nagata Y, Tsuda M. High-temperature-induced
904   transposition of insertion elements in *Burkholderia multivorans* ATCC 17616. *Appl*
905   *Environ Microbiol* **71**, 1822-1828 (2005).

906
907 65. Pasternak C, Ton-Hoang B, Coste G, Bailone A, Chandler M, Sommer S. Irradiation-
908   induced *Deinococcus radiodurans* genome fragmentation triggers transposition of a
909   single resident insertion sequence. *PLoS Genet* **6**, e1000799 (2010).

910
911 66. Siguier P, Gourbeyre E, Varani A, Ton-Hoang B, Chandler M. Everyman's guide to
912   bacterial insertion sequences. *Microbiol Spectr* **3**, MDNA3-0030-2014 (2015).

913
914 67. Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their genomic
915   impact and diversity. *FEMS Microbiol Rev* **38**, 865-891 (2014).

917 68. Chain PS, *et al.* Insights into the evolution of *Yersinia pestis* through whole-genome
918   comparison with *Yersinia pseudotuberculosis. Proc Natl Acad Sci U S A* **101**, 13826-
919   13831 (2004).

920
921 69. Gilmore MS, Lebreton F, van Schaik W. Genomic transition of enterococci from gut
922   commensals to leading causes of multidrug-resistant hospital infection in the antibiotic
923   era. *Curr Opin Microbiol* **16**, 10-16 (2013).

924
925 70. Pasternak C, *et al.* ISDra2 transposition in *Deinococcus radiodurans* is downregulated
926   by TnpB. *Mol Microbiol* **88**, 443-455 (2013).

927
928 71. Rigottier-Gois L, *et al.* The surface rhamnopolysaccharide epa of *Enterococcus faecalis*
929   is a key determinant of intestinal colonization. *J Infect Dis* **211**, 62-71 (2015).

930
931 72. Miskinyte M, *et al.* The genetic basis of *Escherichia coli* pathoadaptation to
932   macrophages. *PLoS Pathog* **9**, e1003802 (2013).

933
934 73. Yang Y, *et al.* Within-host evolution of a gut pathobiont facilitates liver translocation.
935   *Nature* **607**, 563-570 (2022).

936
937 74. Van Tyne D, Manson AL, Huycke MM, Karanicolas J, Earl AM, Gilmore MS. Impact of
938   antibiotic treatment and host innate immune pressure on enterococcal adaptation in the
939   human bloodstream. *Sci Transl Med* **11**, (2019).

940
941 75. Kachroo P, *et al.* Integrated analysis of population genomics, transcriptomics and
942   virulence provides novel insights into *Streptococcus pyogenes* pathogenesis. *Nat Genet*
943   **51**, 548-559 (2019).

944
945 76. Stålhammar-Carlemalm M, Stenberg L, Lindahl G. Protein rib: a novel group B
946   streptococcal cell surface protein that confers protective immunity and is expressed by
947   most strains causing invasive infections. *J Exp Med* **177**, 1593-1603 (1993).

948
949 77. Watanakunakorn C, Bakie C. Synergism of vancomycin-gentamicin and vancomycin-
950   streptomycin against enterococci. *Antimicrob Agents Chemother* **4**, 120-124 (1973).

951
952 78. Moellering RC, Jr., Weinberg AN. Studies on antibiotic syngerism against enterococci. II.
953   Effect of various antibiotics on the uptake of 14 C-labeled streptomycin by enterococci. *J*
954   *Clin Invest* **50**, 2580-2584 (1971).

955
956 79. Pericas JM, *et al.* Changes in the treatment of *Enterococcus faecalis* infective
957   endocarditis in Spain in the last 15 years: from ampicillin plus gentamicin to ampicillin
958   plus ceftriaxone. *Clin Microbiol Infect* **20**, O1075-1083 (2014).

960   80.   Gawryszewska I*, et al.* Distribution of antimicrobial resistance determinants, virulence-
961   associated factors and clustered regularly interspaced palindromic repeats loci in
962   isolates of *Enterococcus faecalis* from various settings and genetic lineages. *Pathog Dis*
963   **75**, (2017).

964
965   81.   Zhanel GG, Schweizer F, Karlowsky JA. Oritavancin: mechanism of action. *Clin Infect*
966   *Dis* **54 Suppl 3**, S214-219 (2012).

967
968   82.   Di Gregorio S, Fernandez S, Perazzi B, Bello N, Famiglietti A, Mollerach M. Increase in
969   IS256 transposition in invasive vancomycin heteroresistant *Staphylococcus aureus*
970   isolate belonging to ST100 and its derived VISA mutants. *Infect Genet Evol* **43**, 197-202
971   (2016).

972
973   83.   Lee MD. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* **35**, 4162-
974   4164 (2019).

975
976   84.   Mai U, Mirarab S. TreeShrink: fast and accurate detection of outlier long branches in
977   collections of phylogenetic trees. *BMC Genomics* **19**, 272 (2018).

978
979   85.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
980   developments. *Nucleic Acids Res* **47**, W256-W259 (2019).

981
982   86.   Shen W, Le S, Li Y, Hu F. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file
983   manipulation. *PLoS One* **11**, e0163962 (2016).

984
985   87.   Seemann T. mlst. *Github - https://githubcom/tseemann/mlst*.

986
987   88.   Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at the
988   population level. *BMC Bioinformatics* **11**, 595 (2010).

989
990   89.   Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic
991   gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119
992   (2010).

993
994   90.   Gish W, States DJ. Identification of protein coding regions by database similarity search.
995   *Nat Genet* **3**, 266-272 (1993).

996
997   91.   Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic
998   platform with an interactive web interface. *Nucleic Acids Res* **47**, D687-D692 (2018).

999
1000   92.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
1001   reads. *EMBnetjournal* **17**, 3 (2011).

1002

1003   93.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
1004   357-359 (2012).

1005
1006   94.   Li H*, et al.* The sequence alignment/map format and SAMtools. *Bioinformatics* **25**, 2078-
1007   2079 (2009).

1008
1009   95.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
1010   features. *Bioinformatics* **26**, 841-842 (2010).

1011
1012   96.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
1013   RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1014
1015   97.   Perez-Casal J, Caparon MG, Scott JR. Mry, a trans-acting positive regulator of the M
1016   protein gene of *Streptococcus pyogenes* with similarity to the receptor proteins of two-
1017   component regulatory systems. *J Bacteriol* **173**, 2617-2624 (1991).

1018

1019

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7