Targeted IS-element sequencing uncovers transposition dynamics during selective

pressure in enterococci

Joshua M. Kirsch[1], Shannon Ely[1], Madison E. Stellfox[2], Karthik Hullahalli[3], Phat Luong[1], Kelli L.

Palmer[3], Daria Van Tyne[2], Breck A. Duerkop[1*]

[1]Department of Immunology and Microbiology, University of Colorado – Anschutz Medical

Campus, School of Medicine, Aurora, CO 80045, USA

[2]Department of Medicine, Division of Infectious Diseases, University of Pittsburgh, Pittsburgh,

PA 15214, USA

[3]Department of Biological Sciences, University of Texas at Dallas, Richardson, TX 75080, USA

*Correspondence: Breck A. Duerkop breck.duerkop@cuanschutz.edu

Abstract

Insertion sequences (IS) are simple transposons implicated in the genome evolution of diverse

pathogenic bacterial species. Enterococci have emerged as important human intestinal

pathogens with newly adapted virulence potential and antibiotic resistance. These genetic

features arose in tandem with large-scale genome evolution mediated by mobile elements.

Pathoadaptation in enterococci is thought to be mediated in part by the IS element IS256

through gene inactivation and recombination events. However, the regulation of IS256 and the

mechanisms controlling its activation are not well understood. Here, we adapt an IS256-specfic

deep sequencing method to describe how chronic lytic phage infection drives widespread

diversification of IS256 in *E. faecalis* and how antibiotic exposure is associated with IS256

diversification in both *E. faecalis* and *E. faecium* during a clinical human infection. We show

through comparative genomics that IS256 is primarily found in hospital-adapted enterococcal

isolates. Analyses of IS256 transposase gene levels reveal that IS256 mobility is regulated at

the transcriptional level by multiple mechanisms in *E. faecalis*, indicating tight control of IS256

activation in the absence of selective pressure. Our findings reveal that stressors such as

phages and antibiotic exposure drives rapid genome-scale transposition in the enterococci.

IS256 diversification can therefore explain how selective pressures mediate evolution of the

enterococcal genome, ultimately leading to the emergence of dominant nosocomial lineages

that threaten human health.

Author Summary

Insertion sequence (IS) elements are simple transposons that are ubiquitous in bacteria. In the

enterococci, which includes medically relevant species such as *Enterococcus faecalis* and

Enterococcus faecium, the IS element IS256 is widespread and has been implicated in

pathogenesis and antibiotic resistance. Despite the importance of IS256 to the biology of the

enterococci, we know very about how this element is regulated and diversifies enterococcal

genomes. Here, we show that IS256 is preferentially found in hospital-adapted and virulent

strains of the enterococci. In *E. faecalis* V583, a vancomycin resistant blood isolate, IS256 is

regulated by multiple transcriptional mechanisms. To understand how IS256 is mobilized, we

adapted an Illumina-based deep sequencing method called IS-Seq to find novel IS256

insertions when applying the selective pressure of bacteriophage (phage) predation. Using this

method, we found that chronic phage infection drives IS256 diversification of the *E. faecalis*

V583 genome. Additionally, we tracked IS256 insertional activity during a human *E. faecium*

infection and found that increased IS256 diversity was associated with specific antibiotic usage.

Together, our results demonstrate that the enterococci control IS256 activity to diversify their

genomes which may lead to the emergence of hospital-adapted strains that threaten human

health.

Introduction

Enterococci, including the human commensals *Enterococcus faecalis* and *Enterococcus*

faecium, are opportunistic pathogens of public health concern due to their acquisition of

antibiotic resistance and virulence traits[1]. Enterococcal infections are a leading cause of

infective endocarditis [2, 3] and 30% of nosocomial infections are vancomycin resistant [4, 5].

Antibiotic resistance in enterococci is often encoded on mobile genetic elements, including

plasmids and composite transposons [6]. Plasmids, especially those containing Inc18, Rep_3,

and RepA-N replicon-containing plasmids, frequently encode resistance genes to diverse

antibiotics, including chloramphenicol, erythromycin, and gentamicin [7, 8]. Composite

transposons are typically organized with a cargo gene (such as an antibiotic resistance gene)

flanked by insertion sequence (IS) elements. Examples in the enterococci include Tn4001 or

Tn4031, which carries gentamicin resistance genes [9] and Tn1546 and Tn1547, which carry

the vancomycin resistance *vanA* and *vanB* operons, respectively [10, 11].

To combat clinically relevant enterococcal strains, new therapeutics are urgently

needed. Recently, the concept of using bacterial viruses (bacteriophages or phages) to treat

multi-drug resistant (MDR) bacteria is being revisited [12]. Phage therapy case studies show

that these viruses can be efficacious against refractory MDR bacteria in human patients [13-17].

However, bacterial resistance to phage infection develops quickly [18-21] and the lessons

learned from the past century of antibiotic use suggest that bacteria will gain resistance to

phage infection after widespread phage treatment [22]. Acquired phage resistance, although

beneficial to bacteria in the face of phage pressure, can come with fitness costs that dampen

antibiotic resistance and virulence [12, 23-25]. Therefore, it is imperative to fully understand the

mechanisms that promote phage resistance in bacteria, and the phenotypic outcomes of phage

resistance. With this knowledge it will be possible to use phage resistance as a tool that can be

leveraged against bacteria to enhance current antibacterial therapeutics.

Enterococci have remarkably diverse genomes containing numerous mobile elements

that contribute to their adaptation and evolution [26-28]. These include plasmids, prophages,

and transposable insertion sequence (IS) elements. Pathogenic enterococci frequently contain

multiple IS elements, and IS elements appear to be a feature of newly-adapted nosocomial

strains [29]. In the widely studied nosocomial type-strain *E. faecalis* V583, there are 38 IS

elements consisting of 11 different types [26]. IS256 is the most abundant IS element in the *E.*

faecalis V583 genome, with six chromosomal copies and four plasmid copies. IS256 is also

common in other gram-positive bacteria, including staphylococci, where it has been widely

studied [30-32]. In the staphylococci, insertion of IS256 into different genes causes clinically-

relevant phenotypes, such as small colony variation [33], biofilm formation [31], and antibiotic

resistance [34]. The transcription factor σ-B negatively regulates IS256 transposition in the

staphylococci through the production of a 3' antisense RNA [31, 35]. IS256 is a crucial

component for enterococcal genome adaptation. In *E. faecalis* V583, the virulence factor

cytolysin is attenuated by IS256 and related IS905 insertions [36]. Additionally, copies of IS256

in the *E. faecalis* V583 pheromone-responsive conjugative plasmids pTEF1 and pTEF2

recombine with chromosomal IS256 copies to mobilize broad regions of the *E. faecalis* genome

[37]. A related IS element, IS16, has been shown to be highly abundant in hospital-adapted *E.*

faecium isolates, suggesting that IS16 may aid in the success of *E. faecium* as a nosocomial

pathogen [38]. Considering IS elements are likely involved in the pathoadaptation of the

enterococci, little is known about how these elements are regulated and what events lead to

their activation.

In this work, we investigated both the regulation and dynamics of IS256 transposition in

E. faecalis and *E. faecium*. We found that IS256 is present in multiple, genetically disparate

isolates of both species, and is enriched in hospital adapted lineages. At steady state, IS256

produces a wealth of low abundance insertions throughout the chromosome and is regulated at

the transcriptional levels to tightly control activation. We discovered that phage infection and

antibiotic usage in different biological settings were associated with increased diversity of IS256

insertions. In the case of phage infection, cell populations with diversified IS256 insertions

maintained phage genomes for an extended period of time and a subpopulation of these cells

chronically shed phage particles throughout growth. Phage shedding provided a competitive

advantage during co-culture with phage-susceptible enterococci, but this advantage is lost when

co-cultured with phage-resistant enterococci. This suggests that phage genome carriage is a

strong selective pressure that drives IS256 diversification and can be used to occupy an

environmental niche when competing with phage-susceptible bacterial strains. Lastly, we

investigated how IS256 diversifies enterococcal genomes in a chronically infected human, and

found that IS256 insertion abundances in vivo coincided with specific antibiotic usage. Together,

this work sheds light on how IS elements are regulated and expanded in enterococcal genomes

during both phage predation and clinical infection, and provides evidence for phage carriage as

an important selective pressure that promotes IS256 diversification. Furthermore, this work

suggests that therapeutic use of both phages and antibiotics could cause rapid and widespread

enterococcal genome evolution for which the physiological consequences are unknown.

Results

IS256 is common within enterococcal lineages associated with hospital adaptation and

pathogenesis

IS256 is an important factor driving the pathoadaptation of *E. faecalis* [36], yet it is

unknown how widely IS256 is distributed among *E. faecalis* strains. To answer this, we

searched all available *E. faecalis* genomes from NCBI RefSeq (2065 genomes) for IS256

transposases with 100% amino acid similarity to the IS256 transposase copies found in *E.*

faecalis V583. A total of 232 genomes contained one or more IS256 sequences (Fig. 1A). The

sporadic distribution of IS256 within the *E. faecalis* phylogenetic tree suggests that IS256 is a

recent addition to many *E. faecalis* genomes and has arisen within multiple discrete lineages of

this species. To further characterize the distribution of IS256 across *E. faecalis* strains, we

performed *in silico* multilocus sequence typing (MLST) on all *E. faecalis* genomes and

compared the proportions of genomes with and without IS256 in each sequence type (ST) (Fig.

1B). IS256 occurs within a variety of ST clades and is most abundant in in ST6, ST103, ST778,

and ST388 genomes, all of which are associated with hospital-adapted, opportunistic

pathogenic lineages (Table S1A) [39-42]. We compared the abundances of different virulence

factors between genomes with and without IS256 (Fig. 1C & Table S1C). IS256-containing

genomes are enriched for multiple virulence factors compared to genomes lacking IS256,

demonstrating that IS256 preferentially occurs in hospital-acquired and virulent *E. faecalis*

strains. We next expanded this analysis to include all *E. faecium* genomes from RefSeq (2306

genomes) and similar to *E. faecalis*, *E. faecium* IS256 is primarily found in the nosocomial

lineages ST17, ST664, ST736, and ST18 (Fig. 1D-E and Table S1B) [43, 44]. Although IS256 is

significantly enriched in virulent *E. faecium* genomes, this was less than *E. faecalis* (Fig. 1F and

Table S1D). Finally, IS256 is more frequently found in *E. faecium* genomes compared to *E.*

faecalis genomes, confirming prior research (Fig. 1G) [45]. Together these data show that IS256

is widely distributed within the enterococci and is preferentially found in nosocomial and virulent

isolates.

IS-Seq identifies widespread movement of IS256 in E. faecalis that is transcriptionally

and translationally controlled

Previously, IS256 sequences were identified as hot-spots in *E. faecalis* V583 that

facilitated the integration of the endogenous plasmids pTEF1 and pTEF2, leading to the

mobilization of the pathogenicity island and other chromosomal regions [37]. Instances of IS256

insertions leading to the inactivation of *E. faecalis* genes involved in diverse phenotypes have

been described [19, 46-48]. However, we lack a complete understanding of IS256 mobility

across the *E. faecalis* genome during selective pressures that would potentially drive IS256

activation and genome diversification. To assess genome-wide IS256 insertions in *E. faecalis*

we adapted a next-generation sequencing (NGS) enrichment technique for use with IS256. This

technique, IS-Seq, includes IS256 amplicon enrichment during NGS library construction

followed by read mapping of IS256-chromosomal junctions to identify specific IS256 insertion

locations (Fig. S1) [49]. Similar techniques have been used to identify IS element insertions in

Acinetobacter [50], and *Mycobacterium* [51]. IS-Seq of wild-type (WT) *E. faecalis* V583 under

steady state conditions identified the known IS256 locations in the chromosome and in the three

pTEF plasmids (Fig. 2A & S2). One of the IS256 copies on pTEF1 lacks a canonical left inverted

repeat (IR), preventing binning of the IS-Seq reads originating from this locus. Numerous low

abundance IS256 insertions are observed throughout the *E. faecalis* genome, indicating

promiscuous movement of the element within subpopulations of cells (Fig. 2A). The three

biological replicate cultures tested in Fig. 2A show that each *E. faecalis* population analyzed has

a unique repertoire of IS256 insertions. We hypothesize that these low abundance steady-state

insertion events provide a mechanism for rapid genome diversification following exposure to

selective pressures.

To understand how IS256 is regulated in *E. faecalis* under steady state conditions, we

investigated possible mechanisms of repression of the IS element. First, we assessed the

transcription of the IS256 transposase (*tnp*) gene. During growth in rich media, IS256 gene

transcription modestly increases during exponential growth, and reaches $1/10^{th}$ of the levels of

the reference 16S rRNA gene (Fig. 2B). Considering there are 10 copies of IS256 and 4 copies

of 16S in *E. faecalis* V583, each IS256 copy achieves $1/25^{th}$ the transcriptional activity of a

single 16S gene if all copies are expressed similarly. Thus, the relatively weak IS256 promoter

may limit over-activation of the element. Next, we investigated whether IS256 is further

transcriptionally or translationally regulated. Other IS elements have been reported to be

controlled by antisense small RNAs (asRNA) [52]. These asRNAs repress transposase

translation through diverse mechanisms, including binding to sense transcripts at or before the

start codon and ribosome binding site to prevent translation of the sense transcript or targeting

the *tnp* transcript for RNase III cleavage [53]. We reanalyzed a publicly available stranded RNA-

Seq dataset of *E. faecalis* V583 [54] by aligning reads to an IS256 element with delineation of

the sense and antisense alignments (Fig. 2C). The transcriptional start site (TSS) of IS256 has

not been experimentally confirmed in any bacterial species and is predicted to reside in the 5'

terminus of the element [55]. We demonstrate that IS256 has two TSSs indicated by high sense

read mapping coverage in distinct peaks. Additionally, each sense peak is overlayed by a higher

antisense peak, demonstrating that antisense inhibition is present at both TSSs. If translation

initiates at the site-2 TSS this would produce a truncated IS256 Tnp, here after referred to as

IS256 Tnp site 2. Upon further investigation of these TSSs and their asRNAs, we found that the

site-1 asRNA is likely under the control of a canonical σ-70 promoter. To identify if this asRNA

controls IS256 Tnp translation, we constructed an IS256 Tnp-GFP translational reporter. This

construct consists of an IS256 Tnp coding sequence fused in frame at its C-terminus to a green

fluorescence protein (GFP) gene. GFP fluorescence serves as a proxy for translation of the

IS256 Tnp. In addition, we built a version of the IS256 Tnp-GFP fusion lacking the predicted -10

element of the site-1 asRNA. We transformed these constructs into *E. faecalis* OG1RF, a strain

that lacks native IS256 elements and IS256-derived asRNAs in its genome, and measured GFP

fluorescence in these cells (Fig. 2D). In cells lacking the site-1 asRNA, there is significantly

higher GFP fluorescence compared to cells with an intact site-1 asRNA promoter,

demonstrating that these asRNAs repress IS256 Tnp translation. This phenotype was fully

complemented. We also measured the RNA levels of the IS256 *tnp* gene from the same

constructs and found that there were significantly more *tnp* transcripts in the IS256-GFP

ΔasRNA cells (Fig. 2E). These results show that in *E. faecalis* IS256 is regulated by multiple

mechanisms at the transcriptional level.

To extend our findings, we compared the abundance of IS256 circular intermediates

between different IS256 constructs in *E. faecalis* OG1RF (Fig. 2F), a strain which lacks any

IS256 elements. IS256 elements circularize during transposition, which can be used to measure

active transposition [30, 31]. Using a multi-copy plasmid backbone, we compared the rate of

IS256 circularization in *E. faecalis* OG1RF carrying the following constructs: WT IS256, an

inactive IS256 gene lacking the coding sequences for the two catalytic aspartic acid residues of

the Tnp's DDE motif (ΔDDE), an IS256 lacking the promoter for the site 1 asRNA (IS256

ΔasRNA), and an IS256 lacking the 5' UTR and coding sequences upstream of the site 2

promoter (IS256 site 2). The IS256 site 2 construct retains the 5' IR. Diagrams of these

constructs are shown in Fig. S3. Surprisingly, we found that all constructs produced significantly

less IS256 circles compared to WT IS256. This suggests that the site 2 Tnp product and IS256

ΔasRNA are inactive for transposition, even though loss of the asRNA increases IS256 Tnp

transcription. To confirm these findings, we complemented the asRNA and found that this

restored the phenotype and increased IS256 circular intermediates above WT IS256 levels.

Additionally, we confirmed that IS256 circles require an active IS256 element as there were

significantly less IS256 circles produced by the inactive IS256 element.

IS256 selection during phage predation

Our previous work suggested that phage predation increases IS256 transposition in *E.*

faecalis V583 and is used as a mechanism of phage resistance [18]. To determine if phage

selective pressure results in the diversification of IS256 elements in *E. faecalis*, we challenged

E. faecalis V583 with the phage phi19 [18]. Following phi19 exposure, we isolated single

colonies that were resistant to phi19. Whole genome sequencing (WGS) of these phage-

resistant isolates predicted new IS256 insertions in phi19 resistant isolates (referred to as 19RS

strains) (Fig. 3A). Among the new IS256 insertions in 19RS strains, we found an insertion in the

3' end of *epaX*, a glycosyltransferase gene involved in the biosynthesis of the enterococcal

polysaccharide antigen that is required for infection by phi19 [18]. To verify this insertion and its

orientation, we performed PCR using genomic DNA (gDNA) from four 19RS strains and WT *E.*

faecalis V583 (Fig. 3B). We found that all 19RS strains harbored an IS256 insertion in *epaX* in

both the forward and reverse complement orientations. This IS256 insertion likely prevents

functional EpaX from being expressed blocking phage infection. Recently, Lossouarn et al.

identified IS256 insertions inactivating *epaX*, confirming that an IS256 insertion in *epaX* is

sufficient to prevent phage infection [48]. These results demonstrate that 19RS strains contain a

novel IS256 insertion not found in WT *E. faecalis* V583. These results also suggest that 19RS

strains, even though they were isolated as individual colonies, are heterogeneous populations

with varied abundances of IS256 insertions.

To test if IS256 provides an advantage to *E. faecalis* against phage infection, we used

phi19 to infect *E. faecalis* OG1RF carrying either an empty vector (EV), WT IS256, or the

inactive version of IS256 (ΔDDE) (Fig. 3C). We found that *E. faecalis* cells containing a

functional IS256 gene were more resistant to phage infection compared to cells containing the

catalytically dead Tnp or the empty vector, indicating that IS256 carriage increases mutation

frequency. To understand if IS256 activation was initiated by phi19 infection, we quantified the

number of IS256 circular intermediates. During the course of phi19 infection, IS256 circular

intermediates significantly increased relative to a mock-infected cells (Fig. 3D).

WGS analysis indicates that phage predation increases IS256 across the *E. faecalis*

genome. However, this method is low resolution. To quantify the exact locations and

abundances of IS256 insertions following phage pressure, we performed IS-Seq on 19RS

strains and compared these to WT *E. faecalis* V583 (Fig. 4A). We expected that each strain

would have substantial IS256 insertional variability, given that 19RS strains contained an IS256

insertion in multiple orientations (Fig. 3B). To provide a rigorous and reproducible assessment

of IS256 insertions in these strains, we performed IS-Seq on gDNA collected from three

biological replicates for both WT *E. faecalis* V583 and 19RS strains. The three replicates were

used to calculate the average peak intensity at each insertion site. IS-Seq revealed a genome-

wide diversification of IS256 insertions at new locations in 19RS strains (Fig. 4A & Table S2).

Pairwise comparisons determined that 19RS strains are significantly enriched in the number of

novel IS256 insertions compared WT *E. faecalis* V583. These comparisons were determined

using the DESeq2 [56], which accounts for over-dispersion and provides a rigorous multiple

testing statistic. We used raw sequencing read counts of each insertion for this analysis (Fig.

4B). The results of this analysis are detailed in Tables S2 and S3. New IS256 insertions in the

pTEF plasmids were also found in the 19RS strains (Fig. S4A-C). Additionally, we measured

IS256 circular intermediates in our IS-Seq dataset by aligning IS256 reads (which originate from

the left terminus of IS256) to the right terminus of an IS256 element. We found that 19RS

strains have a significantly greater number of IS256 circular intermediates compared to WT *E.*

faecalis V583 (Fig. 4C).

We focused on two genomic regions that experienced a high level of IS256 insertions in

the 19RS strains. The first was the *epa* locus. In addition to numerous novel insertions in *epaX*,

we found insertions within other *epa* genes including *epaR*, *epaS*, *epaAA* and *epaAB*. (Fig.

S5A). IS256 insertions in these *epa* genes would alter the teichoic acid decoration of the core

rhammnose-containing Epa, likely altering the surface chemistry of *E. faecalis* resulting in phage

resistance [18]. We also identified multiple insertions in the *vex/vnc* operon (Fig. S5B), which is

purported to have both virulence and antibiotic resistance functions [57, 58].

E. faecalis is a native member of the intestinal microbiota and can outgrow and cause

disease during intestinal dysbiosis. Phages have been proposed as a treatment option for

difficult to control enterococcal intestinal blooms [59]. To determine if phage predation

supported IS256 diversification in the intestine, we colonized mice with *E. faecalis* V583 and

orally challenged these animals with phi19 (Fig. S6). We performed IS-Seq on fecal gDNA from

these animals and found that phi19 exposure in the intestine also increased new IS256

insertions, albeit to a lesser extent than in vitro (Fig. 4D-F, Fig. S4D-F, & Table S3). We found

that similar to cells exposed to phi19 in vitro, IS256 insertions within the same location in *epaX*

and the *vex*/*vnc* operons are observed in the mouse intestine following phage administration

(Fig. S5C-D). This suggests that phage selection plays a role in IS256 diversification in the

context of the intestine.

Enterococci with expanded IS256 genomes chronically shed phi19 phages.

Phage genomes can be maintained in their host bacterium in a variety of configurations,

including chromosomally integrated or as cytosolic episomes [60]. In phage-bacteria

communities, sensitive hosts, resistant hosts, and hosts containing phage genomes coexist at a

consistent ratio [60]. Sensitive hosts are infected with phage particles and either perish,

increasing the number of phage particles, become resistant, or become lysogenized. Over time,

a subset of lysogenized and/or resistant bacteria desensitize to the phage infection and the

cycle repeats. This phenomenon is termed carrier state [60]. Phage genomes can also be in an

intermediate state between lytic replication and lysogenic conversion referred to as

pseudolysogeny [60]. Such phages can enter the lytic cycle spontaneously or following cell

stress, and reports indicate that lysogenic phages can be continuously shed from cells with little

to no reduction in viability of the overall bacterial population [61, 62]. During the analysis of

bacterial colonies originating from *E. faecalis* 19RS strains, we discovered that the phi19

genome is maintained in these phage-resistant bacteria and the population can grow at a stable

rate. This carriage of the phi19 genome may explain its ability to influence IS256 transposition.

We found that 19RS strains continually release infectious phage particles during different stages

of growth (Fig. 5A). These particles were confirmed to be phi19 using PCR (Fig. 5B). We

discovered that ~10% of 19RS colonies produced infectious phages (Fig. 5C). This suggests

that a minority of each population exerts a selective pressure to maintain phage resistance by

producing infectious phage progeny. Loss of phage resistance in this population would lead to

phage infection and population collapse. Colonies that produced zones of clearing were isolated

and serially passaged. Cells that initially produced zones of clearing continued to produce zones

for at least three passages (Fig. S7), demonstrating that phi19 is stably carried in this

subpopulation. Next, we reanalyzed reads from our WGS experiments and found that we could

recover the phi19 genome with high coverage from 19RS strains (Fig. 5D). WT *E. faecalis* V583

did not have any detectable phi19 genomic DNA. To determine if the phage genome was

integrated into the *E. faecalis* chromosome or one of its three endogenous plasmids, we

performed long-read Oxford Nanopore sequencing on the 19RS strains. This technique

provides reads up to 20kb and can be used to find large genomic insertions and

rearrangements. We first searched for reads with phage genomic DNA content and then

identified if any of these reads contained host genomic DNA. Every 19RS strain had reads

containing phage and host hybrid genomic reads. This ranged from 0.2-10% of total phi19 reads

(Fig. 5E). While this may be interpreted evidence of phi19 integration, Nanopore reads have a

low rate of chimeric read formation. Chimeric reads are reads containing two sequences

combined as a result of a technical artifact during library preparation. Chimeric read formation in

Nanopore libraries is reported as 1.7% [63] of total reads, which is in agreement with the

proportion of phi19 reads containing both phage and host DNA. Thus, we believe that the phage

genome is not integrated into the host genome in 19RS cells but is otherwise maintained as an

independent genome that could be considered temperate. Collectively, these results show that

19RS strains release phages from a subpopulation of cells, which may impose stress on *E.*

faecalis influencing IS256 mobility.

Considering that 19RS strains have IS256-rich genomes and release infectious phages,

we hypothesized that phage carriage may increase the competitiveness of 19RS strains when

co-cultured with phage sensitive strains. We performed competition assays where WT *E.*

faecalis or 19RS strains were competed against *E. faecalis* OG1RF or an OG1RF Δ*epaOX*

mutant strain, without the addition of exogenous phages to the cultures (Fig. 6A-B). The *E.*

faecalis OG1RF *epaOX* gene is homologous to *E. faecalis* V583 *epaX* [64] and *E. faecalis*

OG1RF Δ*epaOX* is resistant to phi19 infection. 19RS strains that were shedding phi19

outcompeted *E. faecalis* OG1RF. However, they were unable to outcompete *E. faecalis* OG1RF

Δ*epaOX*. These results demonstrate that phi19 carriage provides a selective advantage to *E.*

faecalis cells when competing with related bacteria that are phage susceptible but if competitors

are phage resistant, phi19 carriage comes with a significant fitness cost. To better understand

the fitness cost of phage carriage, we compared the growth rate of E. faecalis 19RS strains,

V583, and OG1RF in monoculture. *E. faecalis* 19RS strains have a growth defect compared to

V583 and OG1RF when measuring optical density (Fig. 6C) and doubling time (Fig. 6D).

IS256 mobilization occurs in enterococcal genomes during human infection.

IS elements contribute to genome diversification and adaptive plasticity that is important

for bacterial evolution [65]. Environmental pressure is a strong driver of IS element mobility [66-

68], and the work described here expands this knowledge to phage predation. Antibiotics are a

strong selective pressure that can promote bacterial evolution, and IS256 insertions have been

tied to pathoadaptation in enterococci during human infections [36]. Furthermore, antibiotics can

increase IS256 transposition [69] To assess if environmental cues such as antibiotics may also

be driving IS element diversification, we investigated whether enterococcal populations within a

single individual infected with *E. faecium* experiences IS256 mobility during therapeutic

intervention. Stools samples were collected at various timepoints over 109 days, during which

the individual was treated with regimens containing daptomycin, vancomycin, and the

vancomycin analogue oritavancin for recurrent *E. faecium* bacteremia. IS-Seq was performed

on these samples across this time series (Fig. 7A). Nearly all of the insertions found using IS-

Seq occurred in contigs which were generated from *E. faecium*, rather than *E. faecalis* (Fig. S8).

We found that these *E. faecium* populations had a larger proportion of IS256 insertions following

oritavancin therapy compared to daptomycin and/or vancomycin therapy. Comparing

differentially abundant insertions during oritavancin treatment revealed multiple enriched

insertions in the coding sequences of a putative Rib transcriptional regulator, and the *vex/vncRS*

operon also identified as a IS256 insertion hotspot in *E. faecalis* 19RS strains (Fig. 7B-C).

These two operons are associated with antibiotic resistance and virulence in *E. faecium* and

IS256 insertions likely render these genes nonfunctional. These insertions increase in

sequencing depth during the course of treatment and peak during oritavancin administration.

The *vex/vncRS* operon in *E. faecalis* V583 contains an IS element belonging to the ISL3 family

[26], indicating that the genome evolution of *E. faecium* during blood stream infection may follow

a similar trajectory to *E. faecalis* V583 [26]. We also found that IS256 regularly interrupts an

aminoglycoside resistance operon (Fig. S9). Oscillating insertions both in coding and non-

coding regions in this operon suggest that this region may be an IS256 insertion hotspot.

Prophage abundance also increased during oritavancin treatment (Fig. S10). Overabundant

prophages had nucleotide sequence homology to the prophage genomes from *E. faecium*

strains, but lacked homology to enterococcal prophages known to carry platelet binding protein

genes [70] or to enterococcal prophages associated with increased colonization ability [71].

These results suggest that antibiotic treatment can also drive the diversification of multiple

classes of mobile elements in the human host. Lastly, we found that oritavancin-treated samples

had an increase in IS256 circle formation indicating that in addition to novel insertion events,

active IS256 transposition is occurring during blood stream infection (Fig. 7D). Overall, these

findings suggest that IS256 transposition is frequent in the intestine during *E. faecium* infection,

and this transposition likely impacts the antibiotic resistance and virulence profiles of the

infecting bacterial population.

Discussion

IS elements directly shape bacterial physiological responses involved in antibiotic

resistance and virulence, and are key drivers of genome evolution [72-74]. Considering the

importance of IS element biology to bacterial fitness, the selective pressures that guide IS

element regulation and mobility remain poorly defined. Using the Gram-positive opportunistic

pathogen *E. faecalis,* we extend our knowledge of IS elements by detailing the population

biology, regulation, and transposition dynamics of IS256. IS256 is a widely distributed IS

element in Gram-positive pathogens. Through the use of bacterial genetics, comparative

genomics, and IS-Seq we have revealed that IS256 insertion events are strongly dictated by

phage predation and the mammalian host environment. Together these results suggest that

IS256 mobility creates a genetically flexible genome that enables enterococci to rapidly adapt to

diverse environmental conditions.

IS256 is common in *E. faecalis* genomes and is found more frequently in hospital

adapted lineages (Fig. 1). Consistent with this idea, IS256 element abundance is a defining

feature of hospital adapted *E. faecium* isolates [43, 44]. Movement of IS elements between cells

requires other mobile elements, such as phages and plasmids, as IS elements lack any

transforming or transducing ability [27]. The pattern of intermittent IS256 abundance in the

enterococci is likely due to periodic exposure to IS256-containing mobile elements. In other

words, genetically-related strains may interact with IS256-containing mobile elements at

different rates, resulting in the dissemination of IS256 to only a subset of the population. The

enrichment of IS256 among hospital adapted strains may reflect the need for increased

genomic plasticity in these strains as they adapt to an altered host environment [44, 75].

Pathogens can utilize IS elements to form pseudogenes and to condense genome size through

IS-mediated recombination, which can inhibit immune detection through loss of virulence factors

and reduces the genome's metabolic burden [74, 75].

Regulation of IS256 has not been investigated in the enterococci and is key to

understanding this element's biology. Here we show that IS256 is basally active in *E. faecalis*

V583 and is regulated by multiple mechanisms at the transcriptional level (Fig. 2B-E). This is

important because overactive IS256 could be catastrophic to *E. faecalis* viability and layers of

regulation likely temper the elements activity until appropriate conditions require IS256

activation. It has been shown previously that transposase protein levels are correlated to

transposition frequency in *E. coli* [76, 77]. Additionally, we show that IS256 has two different

TSSs as part of this regulation and that both sites are regulated by asRNAs. The protein product

occurring from the site-2 TSS is predicted to be truncated. Other IS elements encode two or

more protein coding frames and these different protein products can contribute to IS element

regulation through inhibition [73, 78]. To better understand this, we built a truncated version of

the IS256 Tnp and showed that it is inactive (Fig. 2F). Previous work has shown that similarly

truncated IS256 variants are able to bind to the element's IRs, but structural studies indicate

that this truncation likely loses key dimerization domains essential for transposition [79, 80]. We

hypothesize that IS256 Tnp site 2 binds to the IRs to prevent full-length IS256 Tnp from binding

and activating transposition.

asRNAs are known to regulate transposase protein levels in the *E. coli*, but to our

knowledge, these have not yet been investigated in the enterococci. Here, we show that the

asRNA produced from site1 in IS256 negatively regulates the RNA levels of the IS256 Tnp.

asRNAs can repress transcription through formation of transcriptional repressors or through

targeting of the mRNA for degradation by RNAse III [53, 81]. Additionally, loss of this asRNA led

to a dramatic decrease in IS256 circular intermediates (Fig. 2F). We hypothesize that this

asRNA plays a crucial role in controlling IS256 transposition.

Using IS-Seq as a readout of basal IS256 diversity, we show that IS256 insertions

function as a mechanism of genome-wide mutation (Fig. 2A). This is the highest resolution IS-

Seq map created, to our knowledge, and demonstrates that this technique can identify low-

abundance insertions with high confidence [50, 51]. Baseline IS256 movement likely adds

mutational diversity to an enterococcal population which allows for adaption to changing

environmental conditions through selection.

Previous work from our group determined that *E. faecalis* infection by the phage phi19

was associated with elevated of IS256 insertions [18] (Fig. 3A). This was found using WGS that

was based on less than 10 reads per insertion. We extended this observation by performing IS-

Seq to accurately identify these novel insertion sites both in vitro and in vivo. We found that *E.*

faecalis challenged with phi19 (19RS strains) had a major increase in new IS256 insertions (Fig.

4A-B). 19RS strains were chronically infected by phi19 and cells within the population carried

the phi19 genome and this chronic infection may provide a continuous selective pressure that

aids in IS256 mobility. This correlated with 19RS strains harboring elevated IS256 circular

intermediates (Fig. 4C). These circles are formed during IS256 copy-paste transposition and

have been studied extensively [30].

We also observed that in a mouse model of *E. faecalis* V583 intestinal colonization,

exposure to phi19 resulted in a similar diversification of IS256 (Fig. 4D-F). IS256 movement was

less robust in the mouse intestine compared to the levels seen in vitro. This may be for several

reasons, including a lower frequency of insertion events per location in vivo and a lower number

of samples sequenced. An additional explanation is that Epa is crucial for bacterial colonization

and *epa* mutations may lead to these mutants being outcompeted in the intestine [18, 82].

Although we found evidence for phage-mediated IS256 activation in *E. faecalis* 19RS

strains, we also determined that selection is important for the location of IS256 insertions (Fig.

4G-H). Cells that were not exposed to phi19 had lower IS256 read density in *epaX. EpaX* is a

glycosyltransferase involved in teichoic acid decoration of the core rhamnose containing Epa

exopolysaccharide backbone. Mutations in *epaX* result in poor phage adsorption [18]. We found

no other insertions in genes known to be involved in phage infection indicating that *epaX* and

other *epa* genes are likely hotspots for IS256 integration following phage pressure. In light of

these discoveries, we hypothesize that initial phage infection may select for *epa*-IS256

insertions and that prolonged phage exposure though chronic infection leads to increased and

non-specific IS256 transposition as a stress response. *epa* mutations have been shown to

sensitize *E. faecalis* to cell wall targeting antibiotics and blood killing and reduces virulence in

animal models [18, 83, 84]. This suggests that even though phage carrier state rapidly evolves

the *E. faecalis* population, it could also reduce the populations virulence potential. We suspect

that phage carriage and release will continually pressure intermittently resistant cells and force

fully resistant cells to maintain high levels of resistance. Carrier state is also associated with

genetic adaption in the *Bacteroides*, a prevalent member of the gut microbiota. *Bacteroides*

coexist with crAss-phages in a constant cycle of infection, resistance, and re-sensitization. To

counter crAss-phage, *Bacteroides* will reversibly invert the genetic sequences of capsule genes

to prevent phage adsorption and attachment [85].

IS elements are involved in pseudogene formation during pathoadaption. Examples

include IS256 insertions in the cytolysin operon of *E. faecalis* V583 [36] and a dependence on

IS insertions in *E. coli* to evolve resistance to macrophage killing [86]. Additionally, how

enterococci evolve during human infection is not well understood and may involve a variety of

genetic changes in cell wall modifying [87] and cellular respiration [88] genes. While IS elements

are directly involved in enterococcal pathoadaptation [36], an in-depth analysis of IS-mediated

mutations during human infection had never been performed. Previously, such dynamics were

studied by laboriously sequencing individual isolates. While this approach can successfully

identify new IS element insertion sites, it overlooks IS element mobility within the entirety of the

bacterial population. Here, we show that IS-Seq can be used to study bacterial adaptation at the

population level and can reveal multiple insertions within such a population (Fig. 7). Following

oritavancin therapy an increase in IS256 insertions occurred in the *vex*/*vncRS* operon, which

has been implicated in both virulence [58] and vancomycin resistance [57]. Additionally, there is

an increase in insertions in a putative transcriptional regulator that may regulate a Rib/Alpha

adhesin [89, 90]. Loss of these potential virulence factors may lead to modulation of virulence

and avoidance of immune activation or detection. We predict that these insertions arise from an

increase in IS256 mobility given the increase in IS256 circular intermediates (Fig. 7D), followed

by selection for beneficial insertions. This is likely similar to phage selection for insertions in the

epa operon.

Additionally, we found highly abundant IS256 insertions in an aminoglycoside

acetyltransferase. Vancomycin is reported to act synergistically with gentamicin against

enterococci [91], likely due to increased gentamicin transport through the vancomycin-

permeabilized cell wall [92]. In recent years, enterococci with aminoglycoside resistance have

emerged [93] with acquisition of resistance genes carried in an IS256 composite transposon

[94]. Our results suggest that genetic modulation of aminoglycoside resistance fluctuates in the

host during infection, with different highly abundant insertions prominent at different timepoints.

This data suggests that aminoglycoside resistance is repeatedly modulated and the intestinal

enterococcal population regularly samples different versions of this aminoglycoside resistance

operon. Future studies to identify if glycopeptide treatment potentiates aminoglycoside

sensitivity through IS256 insertions is warranted. We additionally show that oritavancin

treatment is associated with IS256 diversification in this patient. Oritavancin is an analog of

vancomycin, but is also reported to disrupt cell membrane integrity and possess additional

mechanisms for inhibiting cell wall synthesis [95]. This may explain the difference in IS256

expandability, as this antibiotic may impose strong selective pressure resulting in a broader

range of mutations. Likewise, oritavancin treatment was associated with an increase in IS256

circles. In *Staphylococcus aureus*, vancomycin treatment increases IS256 transposition [35, 69],

reflecting a similar glycopeptide-induced IS256 activation.

In summary, we show how the widespread and clinically-relevant IS element IS256

creates genome diversification during phage predation and human infection using sensitive

NGS IS-Seq. This work provides a high-resolution picture of IS256 movement during both

steady-state and stress inducing conditions. Understanding the outcomes of these IS256

insertion events will be critical to the study of the evolution of enterococcal pathogenesis in

response to both phage and antibiotic therapies. We also show that IS256 is regulated at the

transcriptional level in *E. faecalis* and have discovered that IS256 mobility is biased for highly

mobilizable insertion events in nosocomial and virulent enterococcal isolates, further

demonstrating that IS elements are domesticated and specialized in their choice of bacterial

host.

Materials and Methods

Bacteria and bacteriophages.

E. faecalis strains were grown in Brain Heart Infusion (BHI) broth with aeration at 37°C.

A list of bacterial and bacteriophage strains can be found in Table S4. *Escherichia coli* cultures

were grown in Lennox L broth at 37°C with aeration. 15 µg/mL or 5 µg/mL chloramphenicol was

used for selection of transformed *E. coli* and *E. faecalis*, respectively. The generation of phage-

resistant 19RS strains was recently described [18].

Enterococcal phylogeny and Identification of IS256 in enterococcal genomes

All available *E. faecalis* and *E. faecium* genomes were downloaded from NCBI RefSeq

on 3/21/2022. Phylogenetic trees were constructed with GToTree v1.6.31 using default

parameters [96], pruned using treeshrink v1.3.9 [97], and visualized using the iTOL viewer [98].

IS256 transposase genes were identified using GToTree with the IS256 pfam "PF00872" and

were further refined by selecting matches with 100% amino acid matches to the IS256 Tnp from

E. faecalis V583 using seqkit v0.16.1 [99].

MLST and virulence factor abundance

550   *Enterococcal* genomes obtained from RefSeq were assigned an MLST type using mlst

v2.19.0 [100] with the *E. faecalis* and *E. faecium* MLST scheme from pubmlst [101]. Virulence

factors were identified by calling all predicted ORFs in all downloaded RefSeq genomes using

prodigal v2.6.3 [102] and running diamond blastp v0.9.21 [103] against this database with

known *E. faecalis* and *E. faecium* virulence factors from the Virulence Factor Database [104]. All

further analyses and statistics were performed in R.

DNA extraction

558   Bacterial genomic DNA was extracted from overnight cultures grown in 5 mL BHI broth

559   using the Zymo BIOmics DNA Miniprep Kit (Zymo, Irvine, CA).

561   **Whole genome sequencing, IS256 identification from whole genome sequencing, and**

562   **sequencing read alignment to the phi19 genome**

563   Whole genome sequencing of *E. faecalis* V583 and 19RS strains was performed using

Illumina HiSeq 2500 with 150 cycle paired-end sequencing. These genomes were previously

described [18], and the reads have been deposited at the European Nucleotide Archive (ENA)

as project accession number PRJEB30526. Novel IS256 insertions were found using CLC

Workbench (Qiagen, Hilden, Germany) by first mapping reads to an *E. faecalis* V583 reference

genome where all IS256 elements were removed manually. Unmapped reads were collected

and mapped against a database of known IS256 elements with mapping parameters that

allowed for partial matches (0.4 length fraction and 0.9 fraction similarity). Reads that mapped

against IS256 were binned, the section of the read containing the IS256 sequence was

removed, and the remaining read was realigned against the *E. faecalis* V583 reference genome

(GCA_000007785.1) to identify IS256 insertion sites.

Reads aligning to the phi19 genome were found by first aligning whole genome

sequencing reads to the *E. faecalis* V583 reference genome using Bowtie2 v2.3.5.1[105],

collecting unmapped reads, and aligning these reads to the phi19 genome.

Oxford Nanopore Technologies (ONT) sequencing

High molecular weight genomic DNA was isolated from *E. faecalis* V583 and 19RS

strains after overnight growth as described above. ONT sequencing was performed by

SeqCenter (Pittsburgh, PA) using PCR-free V14 chemistry ligation library preparation and

sequenced on a GridION (ONT, Oxford, UK). 113,000 ± 22,400 reads were generated with an

average read length of 3.7 ± 0.8 kb . phi19 sequences in reads were found using blastn [106]

with a minimum alignment length of 1 kb. Reads passing this threshold were compared to the *E.*

faecalis V583 genome with blastn and positive hits were filtered with a minimum alignment

length of 1 kb. The reads have been deposited at the ENA as project accession number

PRJEB30526.

IS-Seq analysis

For library preparation, 200 ng of genomic DNA was used as input in the Illumina DNA

Prep Kit following the manufacturer's instructions (Illumina, San Diego, CA). For the in vitro and

in vivo experiments (Fig. 4), genomic DNA was isolated from 5×10^9 *E. faecalis* cells. Following

library construction, IS256 enrichment was performed as follows. First, 10 ng of library DNA was

amplified with the p7 primer and the IS-Seq Step1 primer (Table S5) for 13 cycles using Q5 2X

Master Mix (New England Biolabs, Ipswich, MA). These reaction products were diluted 1:100

and 10 μL was added to a PCR reaction with the p7 primer and the IS-Seq Step2 primer for 9

cycles using Q5 2X Master Mix. The products of this second reaction were sequenced on an

Illumina NovaSEQ 6000 by the University of Colorado Anschutz Medical Center Genomics Core

with paired-end 150 cycle chemistry. 23.3 ± 3.8 M reads, 17.9 ± 2.5 M reads, and 16.3 ± 2.6 M

reads were obtained from sequencing the human, mouse, and *in vitro* IS-Seq libraries,

respectively. IS-seq reads have been deposited at the ENA under project accession number

PRJEB55280.

Paired end reads obtained from IS-Seq were partitioned into read1 and read2 files, and

reads from read1 files were used in the downstream analysis. First, cutadapt v1.18 was used to

bin reads with a full IS256 5' terminus. The IS256 5' terminus was then trimmed from these

reads using cutadapt with the following flags: '-O 6 -g

"^CGTAAAAGGACTGTTATATGGCCTTTTTACTTTTACACAATTATACGGACTTTATC" ' [107].

These trimmed reads were aligned to the *E. faecalis* V583 reference genome using Bowtie2 and

the location of each insertion was determined using samtools v1.13 [108] and bedtools

genomecov v2.30.0 [109].

For data generated in Figure 7, contigs were co-assembled from Illumina sequencing

reads from each sample using megahit v. 1.2.7 [110]. These Illumina reads were generated

from libraries prepared using the Nextera prep kit (Illumina) and sequenced on an Illumina

NextSeq 500 with 150 cycle paired-end sequencing. Duplicate contigs were removed using

gclust v1.0 [111] and these representative contigs were used as a reference for aligning IS-Seq

reads as described above.

All further downstream analysis was performed in R. Insertion site abundances were

normalized using DESeq2 estimateSizeFactors [56] before plotting to account for abnormalities

during library preparation and sequencing. Significantly enriched unique insertion sites were

found using DESeq2 by providing raw counts to the program and then computing general

linearized models for each insertion site with at least five reads in three replicates. Models were

fit using either a negative binomial fit, or when appropriate, a local fit. To account for fold change

variance during each pairwise comparison, log fold change values were shrunk using the

ApeGLM model [112] in DESeq2's *lfcshrink* function. Each pairwise comparison was performed

using a separate call to the *DESeq* function in order to improve sensitivity, as recommended by

the developers [56]. Both significantly enriched and insignificant insertions with at least 100

reads were reported in tables S2, S3 and figures 4B, 4D, 7B.

Stranded RNA-Seq analysis

The *E. faecalis* V583 RNA-Seq data used for stranded IS256 gene regulation analysis

was obtained from SRA experiments SRR7229491, SRR7229489, and SRR7229487 [54]. The

reads were aligned to an *E. faecalis* V583 IS256 element sequence using Bowtie2 and forward

and reverse aligning reads were found by parsing the SAM file. Only reads that perfectly aligned

to the IS256 reference were retained. The depth of read alignment at each position in IS256

was calculated using samtools *depth* and the data was visualized in R.

E. faecalis phi19 infection in the mouse intestine

All animal protocols were approved by the Institutional Animal Care and Use Committee

of the University of Colorado School of Medicine (protocol number 00253). Conventional 6-

week-old C57BL/6 mice were divided into two groups (Control: 2 female/ 2 male, Experimental:

2 female/ 2 male). All mice were treated with an antibiotic cocktail (streptomycin [1 mg/ml],

gentamicin [1 mg/ml], erythromycin [200 µg/ml]) by first dosing via oral gavage with 100 µl of

the cocktail and replacing their drinking water with the same antibiotic cocktail for 1 week. On

day 7, antibiotic water was replaced with antibiotic free-drinking water. After 24 hours mice were

colonized with *E. faecalis* V583 suspended in phosphate buffered saline (PBS) by oral gavage

(10^9 CFU in 100 µl). After 24 hours mice were orally gavaged daily as follows: 100 µl of 1M

sodium bicarbonate (all mice) and 100 µl PBS per control mouse or 10^9 PFU phi19 in 100 µl per

experimental mouse. Feces were collected daily and homogenized in 1ml PBS. 10µl of the fecal

slurry was serially diluted and plated on BHI agar containing 100 µg/ml gentamicin or 100 µg/ml

gentamicin and 10^8 pfu/ml phi19. From the experimental samples, 5 µl of slurry was mixed with

45µl chloroform, spun at 16,363 RCF for 1 min and the supernatant was enumerated for phage

particles by agar overlay plaque assay as described previously [19]. *E. faecalis* colonies were

isolated on BHI agar containing 100 µg/ml gentamicin. These cells were grown to stationary

phase in 5 mL of BHI broth, after which genomic DNA was isolated and IS-Seq was performed

on these samples.

RNA extraction and qRT-PCR

RNA was isolated from exponentially growing (OD=0.3) *E. faecalis* using a modified

protocol for the RNeasy Mini-Prep Kit (Qiagen). First, cell pellets were collected by

centrifugation of 5 mL of cell culture at 8,228 RCF for 5 minutes. Cell pellets were resuspended

in 1 mL RNAlater and centrifuged again at 8,228 RCF for 10 minutes. The cell pellets were

stored at -80 °C until sample processing. To isolate RNA, the pellets were thawed and

resuspended in 100 µL TE buffer with 15 mg/mL lysozyme and incubated at room temperature

for 30 minutes. Following this, 700 µL Buffer RLT with 0.01% beta-mercaptoethanol was added

and the sample was transferred to a Lysing Matrix B tube (MPBio, Irvine, CA). The tubes were

bead-beaten in a Mini-Beadbeater-16 (Biospec Products, Bartlesville, OK) on the fastest setting

at 30-second intervals with 1-minute rests on ice between cycles. Following bead beating, the

supernatant was removed after centrifugation at 16,363 RCF for 30 seconds and processed

following the manufacturer's instructions. RNA was eluted from the column in RNase/DNase

free H_2O and any residual DNA contamination was removed with a 1 hour off-column DNase

treatment (Qiagen). The RNA was purified following the RNeasy Mini-Prep Kit following

manufacturer's instructions. cDNA was synthesized using Qscript Master Mix (Quantabio,

Beverley, MA) and 1 µg RNA. For qPCR of gDNA targets, 1 ng of gDNA template was used per

reaction. qPCR was run with PowerUp SYBR Green Master Mix (Thermo Fisher, Waltham, MA)

on an Applied Biosystems QuantStudio 7 Flex qRT-PCR system. qRT-PCR primers are listed in

Table S5. Total 16S rRNA gene transcripts were used for qRT-PCR normalization and *clpX*

was used for DNA qPCR normalization associated with IS circle quantification.

DNA manipulation and cloning

All cloning primers and DNA constructs used in this study are listed in Table S5. All

plasmid constructs used the pLZ12 shuttle vector backbone [113]. For cloning, all inserts and

vectors were amplified using 2X Q5 DNA polymerase Master Mix and assembled using 2X

Gibson Assembly Master Mix (New England Biolabs).

Phage carriage identification and serial passaging

To enumerate extracellular viral particles, we collected supernatant from cultures after

centrifugation for 1 minute at 10,000 RCF, treated with 1/10 volume chloroform, centrifuged for

1 min at 21,000 RCF and extracted the aqueous phase. The resulting supernatant was serially

diluted in SM+ buffer (50 mM Tris-HCl, 100 mM NaCl, 8 mM $MgSO_4$, 5 mM $CaCl_2$ [pH 7.4]),

and 10 µL was mixed with 130 µL of a 1:10 dilution of *E. faecalis* V583 cells grown overnight.

This mixture was combined with 5 mL of 0.35% Todd Hewitt agar (THA) supplemented with

10mM $MgSO_4$, and poured onto THA+10mM $MgSO_4$ plates.

To identify 19RS *E. faecalis* cells actively shedding phages, glycerol stocks of phage-

resistant cultures were streaked on BHI agar and incubated at 37°C. After overnight growth, 182

µL of a 1:10 dilution in SM+ of an overnight wild type *E. faecalis* V583 culture was mixed with 7

mL of 0.35% THA and 10mM $MgSO_4$ and solidified on THA+10mM $MgSO_4$ plates for 30

minutes. Individual colonies were picked from the overnight plate patched onto a plate

containing *E. faecalis* V583 embedded in 0.35% THA, with care not to break the surface of the

agar, and incubated at 37°C overnight.

To serially passage phage-positive colonies, colonies producing zones of clearing were

patched onto both THA+10mM $MgSO_4$ plates and THA+10mM $MgSO_4$ with an *E. faecalis* V583

top agar layer as described above. The next day, the colony on the THA+10mM $MgSO_4$ plate

was patched onto both types of plates and this was continued for two subsequent passages.

Competition assays

E. faecalis cultures were grown overnight in BHI broth and diluted to an OD_{600} of 1 in PBS. 30

µL of each strain was added to 3 mL BHI and incubated at 37°C for 24 hours. The cultures were

plated on both BHI gentamicin 100 µg/mL to select for *E. faecalis* V583 strains and BHI

rifampicin 25 µg/mL, fusidic acid 50 µg/mL to select for *E. faecalis* OG1RF or *E. faecalis*

OG1RF Δ*epaOX*. Competitive fitness was calculated by dividing the final concentration of *E.*

faecalis OG1RF or *E. faecalis* OG1RF Δ*epaOX* by the final concentration of *E. faecalis* V583 or

E. faecalis 19RS and logarithmically scaling this quotient.

Growth curves

E. faecalis cultures were grown overnight in BHI broth and diluted to an OD_{600} of 1 in PBS. 50

µL of each strain was added to 50 mL BHI and incubated at 37°C. OD_{600} readings were taken

every 30 minutes on a Tecan Infinite M Plex (Tecan, Männedorf, Switzerland).

Enterococcal isolation from stool samples and gDNA extraction

Enterococcal stool populations were obtained over a span of 109 days from a patient

being treated for recurrent *E. faecium* bacteremia. Stool samples were diluted and plated onto

bile esculin azide – Enterococcosel agar (Becton Dickinson, Franklin Lakes, NJ), and 100-1,000

colonies were collected and pooled into BHI containing 16.7% glycerol and stored at -80°C. One

mL aliquots of the frozen population stocks were thawed and washed with BHI. Populations

were allowed to briefly expand at 37°C at 170 rpm for 5.5 hours. The cultures were pelleted and

genomic DNA was extracted into DNase free water using the DNeasy Blood and Tissue Kit

(Qiagen). gDNA concentrations were determined with the Qubit 1X dsDNA High-Sensitivity Kit

and fluorometer (Invitrogen, Thermo Fisher).

Phage analysis from clinical enterococcal blood isolates

Viral contigs were found in assembled contigs using VIBRANT v1.2.1 [114] and reads

from clinical samples were mapped against these contigs using Bowtie2. Multiple comparison

analysis was performed using DESeq2 as described above.

Statistics and Visualization

Graphs were made in Graphpad Prism or in R using dplyr and ggplot2. t-tests and one-

way ANOVAs were calculated in Graphpad Prism. Chi-squared tests and insertion site

differential abundance and proportional tests (prop.test) were calculated in R. Significantly

abundant insertions from the IS-Seq experiments were found with DESeq as described above.

Acknowledgements

This work was supported by National Institutes of Health grants R01AI141479 (B.A.D.),

R01AI165519 (D.V.T.), R01AI116610 (K.L.P.), T32AI052066 (J.M.K), F31AI169976 (J.M.K.),

and T32AI138954 (M.E.S.).

Figure Legends

Figure 1. Distribution of IS256 in *E. faecalis* and *E. faecium.* A) Phylogenetic tree of *E.*

faecalis genomes. Genomes with IS256 are labeled in red and genomes without IS256 are

labeled in black. **B)** Presence or absence of IS256 in the genomes of hospital adapted *E.*

faecalis MLST types. **C)** Comparison of *E. faecalis* virulence factor abundances in the genomes

of hospital adapted *E. faecalis* MLST types. **D)** Phylogenetic tree of *E. faecium genomes.*

Genomes with IS256 are labeled in red and genomes without IS256 are labeled in black. **E)**

Presence or absence of IS256 in the genomes of hospital adapted *E. faecium* MLST types. **F)**

Comparison of *E. faecium* virulence factor abundances in the genomes of hospital adapted *E.*

faecium MLST types. **G)** Percent of *E. faecalis* and *E. faecium* genomes from RefSeq that

contain IS256 (* p<0.05, ** p<10^{-5}, *** p<10^{-25}, determined with chi-squared test).

Figure 2. Regulation of IS256 Tnp activity. A) IS256 sequencing (IS-Seq) from three separate

biological replicates of *E. faecalis* V583. Peaks in black represent the six endogenous IS256

insertions identified in the V583 chromosome by whole genome sequencing. **B)** RT-qPCR of

IS256 *tnp* gene levels relative to the housekeeping ribosomal gene 16s (three biological

replicates performed in triplicate). **C)** Stranded RNA-Seq of IS256 tnp gene levels in *E. faecalis*

V583 (three biological replicates). **D)** IS256 Tnp protein levels indicated by measuring GFP

fluorescence of an IS256-GFP and IS256-GFP ΔasRNA translational fusions. (** p<0.01,

determined with t-test from three biological replicates). **E**) RT-qPCR of IS256 Tnp RNA levels of

WT IS256-GFP and IS256-GFP ΔasRNA (* p<0.05, determined with t-test from three biological

replicates). **F**) qPCR of IS256 circular intermediates in *E. faecalis* OG1RF carrying either an

empty vector (EV), IS256, catalytically inactive IS256, IS256 ΔasRNA, or IS256 site 2 Only (****

p<0.0001, determined with one-way ANOVA. Every treatment was compared to the IS256

treatment and corrected for multiple comparisons using the Dunnett method).

Figure 3. IS256 activity confers phage resistance and is elevated in *E. faecalis* isolates

resistant to phi19. A) Enumeration of predicted IS256 insertions determined by whole genome

sequencing of a single wild type *E. faecalis* V583 isolate and four independent phage resistant

19RS strains. A value of 10 indicates the total number of IS256 copies found in the *E. faecalis*

V583 reference genome (GCA_000007785.1). **B)** Representative PCR showing that IS256

insertions occur in both the positive and negative strand orientations in the *epaX* gene 19RS

strains. **C)** Emergence of phage resistant *E. faecalis* OG1RF with IS256 or a catalytically dead

(ΔDDE) IS256. EV indicates *E. faecalis* OG1RF carrying the empty vector. (* $p<0.05$, ** $p<0.01$,

determined by multiple comparison t-test using five biological replicates). **D)** qPCR

quantification of IS256 circular intermediates normalized to the single copy gene *clpX* following

phi19 infection or mock infection (MI). (* $p<0.05$, determined with multiple comparison t-test

from three biological replicates performed in triplicate).

Figure 4. IS-Seq to identify IS256 insertion sites in *E. faecalis* phage resistant isolates in

vitro and in vivo. A) IS-Seq of *E. faecalis* V583, and 19RS strains. Each peak in this panel is

an average of three biological replicates per strain. **B)** Quantification of significantly enriched

insertions between pairwise comparisons of each group in A. In each square, the top value is

the number of insertions overabundant in the column group and the bottom value is number of

insertions overabundant in the row group. **C)** IS256 circle formation determined with IS-Seq

from samples shown in A. (* $p<0.05$, ** $p<0.01$, determined with multiple comparison t-test using

three biological replicates). **D)** IS-Seq of *E. faecalis* V583 and phage resistant colonies isolated

from the feces of mice colonized with (Mouse 5 & 6) and without (Mouse 1 & 2) oral phi19

inoculation. Each peak in this figure is an average of two biological replicates. **E)** Quantification

of significantly enriched insertions between pairwise comparisons of each group in D. In each

square, the top value is the number of insertions overabundant in the column group and the

bottom value is number of insertions overabundant in the row group. **F)** IS256 circle formation

determined with IS-Seq from samples shown in D. Insertions considered significant in panels A

and D have an adjusted $p < 0.05$, a fold change > 4, and were supported by at least 100 reads

in one group during pairwise comparison. Statistics were computed using the DESeq2

generalized linear model algorithm.

Figure 5. *E. faecalis* 19RS strains chronically shed phi19 virus and harbor the phage

genome intracellularly. A) Quantification of shed infectious phi19 phage particles. **B)** PCR

confirmation of phi19 recovered from culture fluid of *E. faecalis* 19RS strains. **C)** Proportion of

cells actively replicating and releasing infectious phi19 phage particles. **D)** WGS read mapping

alignments to the phi19 genome from *E. faecalis* V583 and 19RS strains. Data represent three

biological replicates. **E)** Percentage of phi19 nanopore reads from the 19RS strains that contain

host sequences.

Figure 6. phi19 shedding *E. faecalis* 19RS strains are more fit during competition with a

phage susceptible *E. faecalis* strain. A) Competitive fitness of *E. faecalis* V583 and 19RS

strains relative to *E. faecalis* OG1RF or an *E. faecalis* OG1RF derivative with a deletion in the

epaX homolog *epaOX*. Data represent three biological replicates. **B)** Number of generations of

V583, 19RS, and OG1RF strains during the competition assay in **A**. **C)** Growth curves of V583,

19RS, and OG1RF strains. **D)** Doubling time in hours of Growth curves of V583, 19RS, and

OG1RF strains. (* $p<0.05$, determined with t-tests in comparison to V583 from three biological

replicates. Multiple testing correction was performed with FDR method).

Figure 7. IS-Seq of a longitudinal series of stool samples from a human patient with a

vancomycin resistant enterococcal infection. A) Insertions found via IS-Seq plotted on a

concatenated "metagenome" of contiguous enterococcal sequences from stool genomic DNA.

IS256 sequences were mapped against individual contigs to identify IS256 insertion locations.

Then all contigs were concatenated together to build a linearized representation of the

metagenome. Each insertion is derived from one biological replicate per time point from the

patient's stool. **B)** Volcano plot of differentially abundant insertions during

Daptomycin+Vancomycin co-therapy and Oritavancin therapy. Enriched insertions (dark red)

were overabundant during Oritavancin therapy. Significant insertions have an adjusted $p < 0.05$,

a fold change > 4, and were supported by at least 100 reads in one group during pairwise

comparisons. Statistics were computed using the DESeq2 generalized linear model algorithm.

C) Gene organization of loci where elevated IS256 insertions occurred are indicated by arrows.

Lollipops show the location of IS256 insertion and bar graphs are read depth abundances found

for each insertion. (* $p< 0.05$, **, $p< 0.01$, ***, $p< 0.001$, determined with DESeq2) **D)** IS256

circle formation for each treatment group inferred from IS-Seq. (** $p< 0.01$, determined with

unpaired t-test using 3-4 biological replicates).

Supplementary Figure Legends

Figure S1. Enrichment of IS256-chromosomal junctions during IS-Seq. A) Schematic of the

IS-Seq PCR amplification step. **B)** Trimming site and binning strategy used to identify IS256-

termini reads.

Figure S2. Steady state IS256 insertion sites in the native *E. faecalis* V583 plasmids. A)

pTEF1, **B)** pTEF2, and **C)** pTEF3. Three individual biological replicates are shown in each

panel.

Figure S3. Diagram of IS256 mutant constructs.

Figure S4. IS256 insertions in *E. faecalis* V583 plasmids in vitro and in vivo. A-C) Insertion

sties from in vitro cultured *E. faecalis* 19RS strains; **A)** pTEF1, **B)** pTEF2, and **C)** pTEF3. **D-F)**

Insertion sites from *E. faecalis* isolated from the murine intestine **D)** pTEF1, **E)** pTEF2, and **F)**

pTEF3. Each peak in panels A, B, and C are the average of three biological replicates per

strain, and in panels D, E, and F are an average of two biological replicates.

Figure S5. IS-256 insertions in *epa* and *vex/vnc* genes in vitro and in vivo. Insertion sties

from in vitro cultured *E. faecalis* 19RS strains in the **A)** *epa* locus and **B)** *vex/vnc* locus. Insertion

sites from *E. faecalis* isolated from the murine intestine in the **C)** *epa* locus and **D)** *vex/vnc*

locus. Each peak in panels A and B are the average of three biological replicates per strain, and

in panels C and D are an average of two biological replicates.

Figure S6. Murine model and intestinal *E. faecalis* colonization and oral phi19 treatment.

A) Schematic of the *E. faecalis* intestinal colonization model used to identify in vivo IS256

mobilization. The bacterial population was isolated during the phage treatment phase (between

Day 9-22) by plating on selective media and genomic DNA was isolated from these cells. **B)**

Colony forming units of *E. faecalis* isolated from mouse feces. Bacteria were enumerated on

growth media supplemented with gentamicin with and without the addition of phi19 to measure

the frequency of phage resistant colonies. Arrows indicate time points where IS-Seq was

performed. Four mice were used in both the phi19 treated and untreated groups. Bacteria used

for IS-Seq characterization were isolated on BHI gentamicin media.

Figure S7. Serial passage of phi19 shedding *E. faecalis* 19RS isolates.

Figure S8. IS256 read mapping to enterococcal genomic contigs assembled from stool

samples. Each data point indicates the read mapping abundance per assembled contig. Data

points indicated as *E. faecalis* or *E. faecium* lacked sufficient data to definitively categorize

these contigs as either *E. faecalis* or *E. faecium*.

Figure S9. IS256 insertions from human stool enterococcal isolates in genes involved in

aminoglycoside resistance. Each insertion is derived from one biological replicate per time

point from the patient's stool.

Figure S10. Differential abundance of predicted prophages in clinical samples.

Supplementary Table Legends

Table S1. MLST abundances and virulence factor abundances in IS256 and non-IS256 *E.*

***faecalis* and *E. faecium* genomes.**

Table S2. Read counts and differential abundance analysis of abundant IS256 insertions

in *E. faecalis* V583 and 19RS strains.

Table S3. Read counts and differential abundance analysis of abundant IS256 insertions

in *E. faecalis* V583 isolated from mouse stool.

Table S4. List of bacteria and bacteriophages used in this study.

Table S5. List of primers and plasmids used in this study.

References

1. Fiore E, Van Tyne D, Gilmore MS. Pathogenicity of enterococci. Microbiol Spectr. 2019;7(4).

doi: 10.1128/microbiolspec.GPP3-0053-2018. PMID: 31298205.

2. Shah ASV, McAllister DA, Gallacher P, Astengo F, Rodríguez Pérez JA, Hall J, et al.

Incidence, microbiology, and outcomes in patients hospitalized with infective endocarditis.

Circulation. 2020;141(25):2067-77. doi: 10.1161/circulationaha.119.044913. PMID: 32410460.

3. Khan A, Aslam A, Satti KN, Ashiq S. Infective endocarditis post-transcatheter aortic valve

implantation (TAVI), microbiological profile and clinical outcomes: A systematic review. PLoS

One. 2020;15(1):e0225077. doi: 10.1371/journal.pone.0225077. PMID: 31951610.

4. CDC. AR Threats Report. 2019.

5. Markwart R, Willrich N, Haller S, Noll I, Koppe U, Werner G, et al. The rise in vancomycin-

resistant *Enterococcus faecium* in Germany: data from the German Antimicrobial Resistance

Surveillance (ARS). Antimicrob Resist Infect Control. 2019;8(1):147. doi: 10.1186/s13756-019-

0594-3. PMID: 31485325.

6. Cetinkaya Y, Falk P, Mayhall CG. Vancomycin-resistant enterococci. Clin Microbiol Rev.

2000;13(4):686-707. doi: 10.1128/cmr.13.4.686. PMID: 11023964.

7. Partridge SR, Kwong SM, Firth N, Jensen SO. Mobile genetic elements associated with

antimicrobial resistance. Clin Microbiol Rev. 2018;31(4). doi: 10.1128/cmr.00088-17. PMID:

30068738.

8. Schwarz FV, Perreten V, Teuber M. Sequence of the 50-kb conjugative multiresistance

plasmid pRE25 from *Enterococcus faecalis* RE25. Plasmid. 2001;46(3):170-87. doi:

10.1006/plas.2001.1544. PMID: 11735367.

9. Hodel-Christian SL, Murray BE. Characterization of the gentamicin resistance transposon

Tn5281 from *Enterococcus faecalis* and comparison to staphylococcal transposons Tn4001 and

Tn4031. Antimicrob Agents Chemother. 1991;35(6):1147-52. doi: 10.1128/aac.35.6.1147.

PMID: 1656854.

10. Arthur M, Molinas C, Depardieu F, Courvalin P. Characterization of Tn1546, a Tn3-related

transposon conferring glycopeptide resistance by synthesis of depsipeptide peptidoglycan

precursors in *Enterococcus faecium* BM4147. J Bacteriol. 1993;175(1):117-27. doi:

10.1128/jb.175.1.117-127.1993. PMID: 8380148.

11. Quintiliani R, Jr., Courvalin P. Characterization of Tn1547, a composite transposon flanked

by the IS16 and IS256-like elements, that confers vancomycin resistance in *Enterococcus*

faecalis BM4281. Gene. 1996;172(1):1-8. doi: 10.1016/0378-1119(96)00110-2. PMID: 8654967.

12. Kortright KE, Chan BK, Koff JL, Turner PE. Phage therapy: a renewed approach to combat

antibiotic-resistant bacteria. Cell Host Microbe. 2019;25(2):219-32. doi:

10.1016/j.chom.2019.01.014. PMID: 30763536.

13. Chan BK, Turner PE, Kim S, Mojibian HR, Elefteriades JA, Narayan D. Phage areatment of

an aortic graft infected with *Pseudomonas aeruginosa*. Evol Med Public Health.

2018;2018(1):60-6. doi: 10.1093/emph/eoy005. PMID: 29588855.

14. Schooley RT, Biswas B, Gill JJ, Hernandez-Morales A, Lancaster J, Lessor L, et al.

Development and use of personalized bacteriophage-based therapeutic cocktails to treat a

patient with a disseminated resistant *Acinetobacter baumannii* infection. Antimicrob Agents

Chemother. 2017;61(10). doi: 10.1128/aac.00954-17. PMID: 28807909.

15. Dedrick RM, Guerrero-Bustamante CA, Garlena RA, Russell DA, Ford K, Harris K, et al.

Engineered bacteriophages for treatment of a patient with a disseminated drug-resistant

Mycobacterium abscessus. Nat Med. 2019;25(5):730-3. doi: 10.1038/s41591-019-0437-z.

PMID: 31068712.

16. Aslam S, Lampley E, Wooten D, Karris M, Benson C, Strathdee S, et al. Lessons learned

from the first 10 consecutive cases of intravenous bacteriophage therapy to treat multidrug-

resistant bacterial infections at a single center in the United States. Open Forum Infect Dis.

2020;7(9):ofaa389. doi: 10.1093/ofid/ofaa389. PMID: 33005701.

17. Petrovic Fabijan A, Lin RCY, Ho J, Maddocks S, Ben Zakour NL, Iredell JR. Safety of

bacteriophage therapy in severe *Staphylococcus aureus* infection. Nat Microbiol. 2020;5(3):465-

72. doi: 10.1038/s41564-019-0634-z. PMID: 32066959.

18. Chatterjee A, Johnson CN, Luong P, Hullahalli K, McBride SW, Schubert AM, et al.

Bacteriophage resistance alters antibiotic-mediated intestinal expansion of enterococci. Infect

Immun. 2019;87(6). doi: 10.1128/iai.00085-19. PMID: 30936157.

19. Duerkop BA, Huo W, Bhardwaj P, Palmer KL, Hooper LV. Molecular basis for lytic

bacteriophage resistance in enterococci. mBio. 2016;7(4). doi: 10.1128/mBio.01304-16. PMID:

27578757.

20. Markwitz P, Lood C, Olszak T, van Noort V, Lavigne R, Drulis-Kawa Z. Genome-driven

elucidation of phage-host interplay and impact of phage resistance evolution on bacterial

fitness. ISME J. 2022;16(2):533-42. doi: 10.1038/s41396-021-01096-5. PMID: 34465897.

21. Hesse S, Rajaure M, Wall E, Johnson J, Bliskovsky V, Gottesman S, et al. Phage

resistance in multidrug-resistant *Klebsiella pneumoniae* ST258 evolves via diverse mutations

that culminate in impaired adsorption. mBio. 2020;11(1). doi: 10.1128/mBio.02530-19. PMID:

31992617.

22. Woodford N, Ellington MJ. The emergence of antibiotic resistance by mutation. Clin

Microbiol Infect. 2007;13(1):5-18. doi: 10.1111/j.1469-0691.2006.01492.x. PMID: 17184282.

23. Mangalea MR, Duerkop BA. Fitness trade-offs resulting from bacteriophage resistance

potentiate synergistic antibacterial strategies. Infect Immun. 2020;88(7). doi: 10.1128/iai.00926-

19. PMID: 32094257.

24. Gordillo Altamirano F, Forsyth JH, Patwa R, Kostoulias X, Trim M, Subedi D, et al.

Bacteriophage-resistant *Acinetobacter baumannii* are resensitized to antimicrobials. Nat

Microbiol. 2021;6(2):157-61. doi: 10.1038/s41564-020-00830-7. PMID: 33432151.

25. Castledine M, Padfield D, Sierocinski P, Soria Pascual J, Hughes A, Mäkinen L, et al.

Parallel evolution of *Pseudomonas aeruginosa* phage resistance and virulence loss in response

to phage treatment in vivo and in vitro. eLife. 2022;11:e73679. doi: 10.7554/eLife.73679. PMID:

35188102.

26. Paulsen IT, Banerjei L, Myers GS, Nelson KE, Seshadri R, Read TD, et al. Role of mobile

DNA in the evolution of vancomycin-resistant *Enterococcus faecalis*. Science.

2003;299(5615):2071-4. doi: 10.1126/science.1080613. PMID: 12663927.

27. Johnson CN, Sheriff EK, Duerkop BA, Chatterjee A. Let me upgrade you: impact of mobile

genetic elements on enterococcal adaptation and evolution. J Bacteriol.

2021;203(21):e0017721. doi: 10.1128/jb.00177-21. PMID: 34370561.

28. Clewell DB, Weaver KE, Dunny GM, Coque TM, Francia MV, Hayes F. Extrachromosomal

and mobile elements in enterococci: transmission, maintenance, and epidemiology. In: Gilmore

MS, Clewell DB, Ike Y, Shankar N, editors. Enterococci: From Commensals to Leading Causes

of Drug Resistant Infection. Boston: Massachusetts Eye and Ear Infirmary; 2014.

29.   Werner G, Fleige C, Geringer U, van Schaik W, Klare I, Witte W. IS element IS16 as a

molecular screening tool to identify hospital-associated strains of *Enterococcus faecium*. BMC

Infect Dis. 2011;11:80. doi: 10.1186/1471-2334-11-80. PMID: 21453466.

30.   Loessner I, Dietrich K, Dittrich D, Hacker J, Ziebuhr W. Transposase-dependent formation

of circular IS256 derivatives in *Staphylococcus epidermidis* and *Staphylococcus aureus*. J

Bacteriol. 2002;184(17):4709-14. doi: 10.1128/jb.184.17.4709-4714.2002. PMID: 12169594.

31.   Valle J, Vergara-Irigaray M, Merino N, Penadés JR, Lasa I. sigmaB regulates IS256-

mediated *Staphylococcus aureus* biofilm phenotypic variation. J Bacteriol. 2007;189(7):2886-96.

doi: 10.1128/jb.01767-06. PMID: 17277051.

32.   Kozitskaya S, Cho SH, Dietrich K, Marre R, Naber K, Ziebuhr W. The bacterial insertion

sequence element IS256 occurs preferentially in nosocomial *Staphylococcus epidermidis*

isolates: association with biofilm formation and resistance to aminoglycosides. Infect Immun.

2004;72(2):1210-5. doi: 10.1128/IAI.72.2.1210-1215.2004. PMID: 14742578.

33.   Kleinert F, Kallies R, Hort M, Zweynert A, Szekat C, Nagel M, et al. Influence of IS256 on

genome variability and formation of small-colony variants in *Staphylococcus aureus*. Antimicrob

Agents Chemother. 2017;61(8). doi: 10.1128/AAC.00144-17. PMID: 28584147.

34.   McEvoy CR, Tsuji B, Gao W, Seemann T, Porter JL, Doig K, et al. Decreased vancomycin

susceptibility in *Staphylococcus aureus* caused by IS256 tempering of WalKR expression.

Antimicrob Agents Chemother. 2013;57(7):3240-9. doi: 10.1128/AAC.00279-13. PMID:

23629723.

35.   Nagel M, Reuter T, Jansen A, Szekat C, Bierbaum G. Influence of ciprofloxacin and

vancomycin on mutation rate and transposition of IS256 in *Staphylococcus aureus*. Int J Med

Microbiol. 2011;301(3):229-36. doi: 10.1016/j.ijmm.2010.08.021. PMID: 21115395.

36. Shankar N, Baghdayan AS, Gilmore MS. Modulation of virulence within a pathogenicity

island in vancomycin-resistant *Enterococcus faecalis*. Nature. 2002;417(6890):746-50. doi:

10.1038/nature00802. PMID: 12066186.

37. Manson JM, Hancock LE, Gilmore MS. Mechanism of chromosomal transfer of

Enterococcus faecalis pathogenicity island, capsule, antimicrobial resistance, and other traits.

Proc Natl Acad Sci U S A. 2010;107(27):12269-74. doi: 10.1073/pnas.1000139107. PMID:

20566881.

38. Leavis HL, Willems RJ, van Wamel WJ, Schuren FH, Caspers MP, Bonten MJ. Insertion

sequence-driven diversification creates a globally dispersed emerging multiresistant subspecies

of *E. faecium*. PLoS Pathog. 2007;3(1):e7. doi: 10.1371/journal.ppat.0030007. PMID:

17257059.

39. Pöntinen AK, Top J, Arredondo-Alonso S, Tonkin-Hill G, Freitas AR, Novais C, et al.

Apparent nosocomial adaptation of *Enterococcus faecalis* predates the modern hospital era. Nat

Commun. 2021;12(1):1523. doi: 10.1038/s41467-021-21749-5. PMID: 33750782.

40. Dabul ANG, Avaca-Crusca JS, Navais RB, Merlo TP, Van Tyne D, Gilmore MS, et al.

Molecular basis for the emergence of a new hospital endemic tigecycline-resistant

Enterococcus faecalis ST103 lineage. Infect Genet Evol. 2019;67:23-32. doi:

10.1016/j.meegid.2018.10.018. PMID: 30393188.

41. Schell CM, Tedim AP, Rodríguez-Baños M, Sparo MD, Lissarrague S, Basualdo JA, et al.

Detection of β-lactamase-producing *Enterococcus faecalis* and vancomycin-resistant

Enterococcus faecium isolates in human invasive infections in the public hospital of Tandil,

Argentina. Pathogens. 2020;9(2). doi: 10.3390/pathogens9020142. PMID: 32093230.

42. Zaheer R, Cook SR, Barbieri R, Goji N, Cameron A, Petkau A, et al. Surveillance of

Enterococcus spp. reveals distinct species and antimicrobial resistance diversity across a One-

Health continuum. Sci Rep. 2020;10(1):3937. doi: 10.1038/s41598-020-61002-5. PMID:

32127598.

43. Lebreton F, van Schaik W, McGuire AM, Godfrey P, Griggs A, Mazumdar V, et al.

Emergence of epidemic multidrug-resistant *Enterococcus faecium* from animal and commensal

strains. mBio. 2013;4(4). doi: 10.1128/mBio.00534-13. PMID: 23963180.

44. Chilambi GS, Nordstrom HR, Evans DR, Ferrolino JA, Hayden RT, Marón GM, et al.

Evolution of vancomycin-resistant *Enterococcus faecium* during colonization and infection in

immunocompromised pediatric patients. Proc Natl Acad Sci U S A. 2020;117(21):11703-14. doi:

10.1073/pnas.1917130117. PMID: 32393645.

45. Mikalsen T, Pedersen T, Willems R, Coque TM, Werner G, Sadowy E, et al. Investigating

the mobilome in clinically important lineages of *Enterococcus faecium* and *Enterococcus*

faecalis. BMC Genomics. 2015;16:282. doi: 10.1186/s12864-015-1407-6. PMID: 25885771.

46. Perez M, Calles-Enríquez M, del Rio B, Ladero V, Martín MC, Fernández M, et al. IS256

abolishes gelatinase activity and biofilm formation in a mutant of the nosocomial pathogen

*Enterococcus faecali*s V583. Can J Microbiol. 2015;61(7):517-9. doi: 10.1139/cjm-2015-0090.

PMID: 25966618.

47. Palmer KL, Daniel A, Hardy C, Silverman J, Gilmore MS. Genetic basis for daptomycin

resistance in enterococci. Antimicrob Agents Chemother. 2011;55(7):3345-56. doi:

10.1128/aac.00207-11. PMID: 21502617.

48. Lossouarn J, Briet A, Moncaut E, Furlan S, Bouteau A, Son O, et al. *Enterococcus faecalis*

countermeasures defeat a virulent picovirinae bacteriophage. Viruses. 2019;11(1). doi:

10.3390/v11010048. PMID: 30634666.

49. Dale JL, Beckman KB, Willett JLE, Nilson JL, Palani NP, Baller JA, et al. Comprehensive

functional analysis of the *Enterococcus faecalis* core genome using an ordered, sequence-

defined collection of insertional mutations in strain OG1RF. mSystems. 2018;3(5). doi:

10.1128/mSystems.00062-18. PMID: 30225373.

50. Wright MS, Mountain S, Beeri K, Adams MD. Assessment of insertion sequence

mobilization as an adaptive response to oxidative stress in *Acinetobacter baumannii* using IS-

seq. J Bacteriol. 2017;199(9). doi: 10.1128/jb.00833-16. PMID: 28193905.

51. Reyes A, Sandoval A, Cubillos-Ruiz A, Varley KE, Hernández-Neuta I, Samper S, et al. IS-

seq: a novel high throughput survey of in vivo IS6110 transposition in multiple *Mycobacterium*

tuberculosis genomes. BMC Genomics. 2012;13:249. doi: 10.1186/1471-2164-13-249. PMID:

22703188.

52. Ellis MJ, Trussler RS, Haniford DB. A cis-encoded sRNA, Hfq and mRNA secondary

structure act independently to suppress IS200 transposition. Nucleic Acids Res.

2015;43(13):6511-27. doi: 10.1093/nar/gkv584. PMID: 26044710.

53. Case CC, Simons EL, Simons RW. The IS10 transposase mRNA is destabilized during

antisense RNA control. EMBO J. 1990;9(4):1259-66. doi: 10.1002/j.1460-2075.1990.tb08234.x.

PMID: 1691096.

54. Michaux C, Hansen EE, Jenniches L, Gerovac M, Barquist L, Vogel J. Single-nucleotide

RNA maps for the two major nosocomial pathogens *Enterococcus faecalis* and *Enterococcus*

faecium. Front Cell Infect Microbiol. 2020;10:600325. doi: 10.3389/fcimb.2020.600325. PMID:

33324581.

55. Prudhomme M, Turlan C, Claverys JP, Chandler M. Diversity of Tn4001 transposition

products: the flanking IS256 elements can form tandem dimers and IS circles. J Bacteriol.

2002;184(2):433-43. doi: 10.1128/jb.184.2.433-443.2002. PMID: 11751820.

56. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biol. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8. PMID:

25516281.

57. Haas W, Sublett J, Kaushal D, Tuomanen EI. Revising the role of the pneumococcal *vex-*

vncRS locus in vancomycin tolerance. J Bacteriol. 2004;186(24):8463-71. doi:

10.1128/jb.186.24.8463-8471.2004. PMID: 15576796.

58. Lee S, Ghosh P, Kwon H, Park SS, Kim GL, Choi SY, et al. Induction of the pneumococcal

vncRS operon by lactoferrin is essential for pneumonia. Virulence. 2018;9(1):1562-75. doi:

10.1080/21505594.2018.1526529. PMID: 30246592.

59. Bolocan AS, Upadrasta A, Bettio PHA, Clooney AG, Draper LA, Ross RP, et al. Evaluation

of phage therapy in the context of *Enterococcus faecalis* and its associated diseases. Viruses.

2019;11(4). doi: 10.3390/v11040366. PMID: 31010053.

60. Mäntynen S, Laanto E, Oksanen HM, Poranen MM, Díaz-Muñoz SL. Black box of phage-

bacterium interactions: exploring alternative phage infection strategies. Open Biol.

2021;11(9):210188. doi: 10.1098/rsob.210188. PMID: 34520699.

61. Liu Y, Alexeeva S, Bachmann H, Guerra Martínez JA, Yeremenko N, Abee T, et al. Chronic

release of tailless phage particles from *Lactococcus lactis*. Appl Environ Microbiol.

2022;88(1):e0148321. doi: 10.1128/aem.01483-21. PMID: 34705552.

62. Liu J, Yan R, Zhong Q, Ngo S, Bangayan NJ, Nguyen L, et al. The diversity and host

interactions of *Propionibacterium acnes* bacteriophages on human skin. ISME J.

2015;9(9):2078-93. doi: 10.1038/ismej.2015.47. PMID: 25848871.

63. White R, Pellefigues C, Ronchese F, Lamiable O, Eccles D. Investigation of chimeric reads

using the MinION. F1000Res. 2017;6:631. doi: 10.12688/f1000research.11547.2. PMID:

28928943.

64. Dale JL, Cagnazzo J, Phan CQ, Barnes AM, Dunny GM. Multiple roles for *Enterococcus*

faecalis glycosyltransferases in biofilm-associated antibiotic resistance, cell envelope integrity,

and conjugative transfer. Antimicrob Agents Chemother. 2015;59(7):4094-105. doi:

10.1128/aac.00344-15. PMID: 25918141.

65. Touchon M, Rocha EP. Causes of insertion sequences abundance in prokaryotic genomes.

Mol Biol Evol. 2007;24(4):969-81. doi: 10.1093/molbev/msm014. PMID: 17251179.

66. Drevinek P, Baldwin A, Lindenburg L, Joshi LT, Marchbank A, Vosahlikova S, et al.

Oxidative stress of *Burkholderia cenocepaci*a induces insertion sequence-mediated genomic

rearrangements that interfere with macrorestriction-based genotyping. J Clin Microbiol.

2010;48(1):34-40. doi: 10.1128/jcm.01433-09. PMID: 19889907.

67. Ohtsubo Y, Genka H, Komatsu H, Nagata Y, Tsuda M. High-temperature-induced

transposition of insertion elements in *Burkholderia multivorans* ATCC 17616. Appl Environ

Microbiol. 2005;71(4):1822-8. doi: 10.1128/aem.71.4.1822-1828.2005. PMID: 15812007.

68. Pasternak C, Ton-Hoang B, Coste G, Bailone A, Chandler M, Sommer S. Irradiation-

induced *Deinococcus radiodurans* genome fragmentation triggers transposition of a single

resident insertion sequence. PLoS Genet. 2010;6(1):e1000799. doi:

10.1371/journal.pgen.1000799. PMID: 20090938.

69. Di Gregorio S, Fernandez S, Perazzi B, Bello N, Famiglietti A, Mollerach M. Increase in

IS256 transposition in invasive vancomycin heteroresistant *Staphylococcus aureus* isolate

belonging to ST100 and its derived VISA mutants. Infect Genet Evol. 2016;43:197-202. doi:

10.1016/j.meegid.2016.05.001. PMID: 27154328.

70. Matos RC, Lapaque N, Rigottier-Gois L, Debarbieux L, Meylheuc T, Gonzalez-Zorn B, et al.

Enterococcus faecalis prophage dynamics and contributions to pathogenic traits. PLoS Genet.

2013;9(6):e1003539. doi: 10.1371/journal.pgen.1003539. PMID: 23754962.

71. Duerkop BA, Clements CV, Rollins D, Rodrigues JL, Hooper LV. A composite

bacteriophage alters colonization by an intestinal commensal bacterium. Proc Natl Acad Sci U S

A. 2012;109(43):17621-6. doi: 10.1073/pnas.1206136109. PMID: 23045666.

72. Siguier P, Gourbeyre E, Varani A, Ton-Hoang B, Chandler M. Everyman's guide to bacterial

insertion sequences. Microbiol Spectr. 2015;3(2):MDNA3-0030-2014. doi:

10.1128/microbiolspec.MDNA3-0030-2014. PMID: 26104715.

73. Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their genomic impact

and diversity. FEMS Microbiol Rev. 2014;38(5):865-91. doi: 10.1111/1574-6976.12067. PMID:

24499397.

74. Chain PS, Carniel E, Larimer FW, Lamerdin J, Stoutland PO, Regala WM, et al. Insights

into the evolution of *Yersinia pestis* through whole-genome comparison with *Yersinia*

pseudotuberculosis. Proc Natl Acad Sci U S A. 2004;101(38):13826-31. doi:

10.1073/pnas.0404012101. PMID: 15358858.

75. Gilmore MS, Lebreton F, van Schaik W. Genomic transition of enterococci from gut

commensals to leading causes of multidrug-resistant hospital infection in the antibiotic era. Curr

Opin Microbiol. 2013;16(1):10-6. doi: 10.1016/j.mib.2013.01.006. PMID: 23395351.

76. Ross JA, Wardle SJ, Haniford DB. Tn10/IS10 transposition is downregulated at the level of

transposase expression by the RNA-binding protein Hfq. Mol Microbiol. 2010;78(3):607-21. doi:

10.1111/j.1365-2958.2010.07359.x. PMID: 20815820.

77. Ellis MJ, Trussler RS, Haniford DB. Hfq binds directly to the ribosome-binding site of IS10

transposase mRNA to inhibit translation. Mol Microbiol. 2015;96(3):633-50. doi:

10.1111/mmi.12961. PMID: 25649688.

78. Pasternak C, Dulermo R, Ton-Hoang B, Debuchy R, Siguier P, Coste G, et al. ISDra2

transposition in *Deinococcus radiodurans* is downregulated by TnpB. Mol Microbiol.

2013;88(2):443-55. doi: 10.1111/mmi.12194. PMID: 23461641.

79. Hennig S, Ziebuhr W. Characterization of the transposase encoded by IS256, the prototype

of a major family of bacterial insertion sequence elements. J Bacteriol. 2010;192(16):4153-63.

doi: 10.1128/jb.00226-10. PMID: 20543074.

80. Kosek D, Hickman AB, Ghirlando R, He S, Dyda F. Structures of ISCth4 transpososomes

reveal the role of asymmetry in copy-out/paste-in DNA transposition. EMBO J.

2021;40(1):e105666. doi: 10.15252/embj.2020105666. PMID: 33006208.

81. Brantl S, Birch-Hirschfeld E, Behnke D. RepR protein expression on plasmid pIP501 is

controlled by an antisense RNA-mediated transcription attenuation mechanism. J Bacteriol.

1993;175(13):4052-61. doi: 10.1128/jb.175.13.4052-4061.1993. PMID: 8320221.

82. Rigottier-Gois L, Madec C, Navickas A, Matos RC, Akary-Lepage E, Mistou MY, et al. The

surface rhamnopolysaccharide epa of *Enterococcus faecalis* is a key determinant of intestinal

colonization. J Infect Dis. 2015;211(1):62-71. doi: 10.1093/infdis/jiu402. PMID: 25035517.

83. Teng F, Jacques-Palaz KD, Weinstock GM, Murray BE. Evidence that the enterococcal

polysaccharide antigen gene (*epa*) cluster is widespread in *Enterococcus faecalis* and

influences resistance to phagocytic killing of *E. faecalis*. Infect Immun. 2002;70(4):2010-5. doi:

10.1128/iai.70.4.2010-2015.2002. PMID: 11895965.

84. Smith RE, Salamaga B, Szkuta P, Hajdamowicz N, Prajsnar TK, Bulmer GS, et al.

Decoration of the enterococcal polysaccharide antigen EPA is essential for virulence, cell

surface charge and interaction with effectors of the innate immune system. PLoS Pathog.

2019;15(5):e1007730. doi: 10.1371/journal.ppat.1007730. PMID: 31048927.

85. Shkoporov AN, Khokhlova EV, Stephens N, Hueston C, Seymour S, Hryckowian AJ, et al.

Long-term persistence of crAss-like phage crAss001 is associated with phase variation in

Bacteroides intestinalis. BMC Biology. 2021;19(1):163. doi: 10.1186/s12915-021-01084-3.

PMID: 34407825.

86. Miskinyte M, Sousa A, Ramiro RS, de Sousa JA, Kotlinowski J, Caramalho I, et al. The

genetic basis of *Escherichia coli* pathoadaptation to macrophages. PLoS Pathog.

2013;9(12):e1003802. doi: 10.1371/journal.ppat.1003802. PMID: 24348252.

87. Yang Y, Nguyen M, Khetrapal V, Sonnert ND, Martin AL, Chen H, et al. Within-host

evolution of a gut pathobiont facilitates liver translocation. Nature. 2022;607(7919):563-70. doi:

10.1038/s41586-022-04949-x. PMID: 35831502.

88. Van Tyne D, Manson AL, Huycke MM, Karanicolas J, Earl AM, Gilmore MS. Impact of

antibiotic treatment and host innate immune pressure on enterococcal adaptation in the human

bloodstream. Sci Transl Med. 2019;11(487). doi: 10.1126/scitranslmed.aat8418. PMID:

30971455.

89. Kachroo P, Eraso JM, Beres SB, Olsen RJ, Zhu L, Nasser W, et al. Integrated analysis of

population genomics, transcriptomics and virulence provides novel insights into *Streptococcus*

pyogenes pathogenesis. Nat Genet. 2019;51(3):548-59. doi: 10.1038/s41588-018-0343-1.

PMID: 30778225.

90. Stålhammar-Carlemalm M, Stenberg L, Lindahl G. Protein rib: a novel group B

streptococcal cell surface protein that confers protective immunity and is expressed by most

strains causing invasive infections. J Exp Med. 1993;177(6):1593-603. doi:

10.1084/jem.177.6.1593. PMID: 8496678.

91. Watanakunakorn C, Bakie C. Synergism of vancomycin-gentamicin and vancomycin-

streptomycin against enterococci. Antimicrob Agents Chemother. 1973;4(2):120-4. doi:

10.1128/aac.4.2.120. PMID: 4790933.

92. Moellering RC, Jr., Weinberg AN. Studies on antibiotic syngerism against enterococci. II.

Effect of various antibiotics on the uptake of 14 C-labeled streptomycin by enterococci. J Clin

Invest. 1971;50(12):2580-4. doi: 10.1172/jci106758. PMID: 5001959.

93. Pericas JM, Cervera C, del Rio A, Moreno A, Garcia de la Maria C, Almela M, et al.

Changes in the treatment of *Enterococcus faecalis* infective endocarditis in Spain in the last

15 years: from ampicillin plus gentamicin to ampicillin plus ceftriaxone. Clin Microbiol Infect.

2014;20(12):O1075-83. doi: 10.1111/1469-0691.12756. PMID: 25040215.

94. Gawryszewska I, Malinowska K, Kuch A, Chrobak-Chmiel D, Trokenheim LL, Hryniewicz

W, et al. Distribution of antimicrobial resistance determinants, virulence-associated factors and

clustered regularly interspaced palindromic repeats loci in isolates of *Enterococcus faecalis* from

various settings and genetic lineages. Pathog Dis. 2017;75(2). doi: 10.1093/femspd/ftx021.

PMID: 28334141.

95. Zhanel GG, Schweizer F, Karlowsky JA. Oritavancin: mechanism of action. Clin Infect Dis.

2012;54 Suppl 3:S214-9. doi: 10.1093/cid/cir920. PMID: 22431851.

96. Lee MD. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics.

2019;35(20):4162-4. doi: 10.1093/bioinformatics/btz188. PMID: 30865266.

97. Mai U, Mirarab S. TreeShrink: fast and accurate detection of outlier long branches in

collections of phylogenetic trees. BMC Genomics. 2018;19(Suppl 5):272. doi: 10.1186/s12864-

018-4620-2. PMID: 29745847.

98. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.

Nucleic Acids Res. 2019;47(W1):W256-W9. doi: 10.1093/nar/gkz239. PMID: 30931475.

99. Shen W, Le S, Li Y, Hu F. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file

manipulation. PLoS One. 2016;11(10):e0163962. doi: 10.1371/journal.pone.0163962. PMID:

27706213.

100. Seemann T. mlst. Github - https://githubcom/tseemann/mlst.

101. Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at the

population level. BMC Bioinformatics. 2010;11(1):595. doi: 10.1186/1471-2105-11-595. PMID:

21143983.

102. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic

gene recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:119.

doi: 10.1186/1471-2105-11-119. PMID: 20211023.

103. Gish W, States DJ. Identification of protein coding regions by database similarity search.

Nat Genet. 1993;3(3):266-72. doi: 10.1038/ng0393-266. PMID: 8485583.

104. Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic platform

with an interactive web interface. Nucleic Acids Res. 2018;47(D1):D687-D92. doi:

10.1093/nar/gky1080. PMID: 30395255.

105. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012;9(4):357-9. doi: 10.1038/nmeth.1923. PMID: 22388286.

106. McGinnis S, Madden TL. BLAST: at the core of a powerful and diverse set of sequence

analysis tools. Nucleic Acids Res. 2004;32(Web Server issue):W20-5. doi: 10.1093/nar/gkh435.

PMID: 15215342.

107. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnetjournal. 2011;17(1):3. doi: 10.14806/ej.17.1.200.

108. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence

alignment/map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. doi:

10.1093/bioinformatics/btp352. PMID: 19505943.

109.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics. 2010;26(6):841-2. doi: 10.1093/bioinformatics/btq033. PMID: 20110278.

110.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution

for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.

2015;31(10):1674-6. doi: 10.1093/bioinformatics/btv033. PMID: 25609793.

111.   Li R, He X, Dai C, Zhu H, Lang X, Chen W, et al. Gclust: A parallel clustering tool for

microbial genomic data. Genomics, Proteomics & Bioinformatics. 2019;17(5):496-502. doi:

10.1016/j.gpb.2018.10.008. PMID: 31917259.

112.   Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data:

removing the noise and preserving large differences. Bioinformatics. 2019;35(12):2084-92. doi:

10.1093/bioinformatics/bty895. PMID: 30395178.

113.   Perez-Casal J, Caparon MG, Scott JR. Mry, a trans-acting positive regulator of the M

protein gene of *Streptococcus pyogenes* with similarity to the receptor proteins of two-

component regulatory systems. J Bacteriol. 1991;173(8):2617-24. doi: 10.1128/jb.173.8.2617-

2624.1991. PMID: 1849511.

114.   Kieft K, Zhou Z, Anantharaman K. VIBRANT: automated recovery, annotation and curation

of microbial viruses, and evaluation of viral community function from genomic sequences.

Microbiome. 2020;8(1):90. doi: 10.1186/s40168-020-00867-0. PMID: 32522236.

1270

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7