C$_4$ Grasses Employ Various Strategies to Acclimate Rubisco Activase to Heat Stress

Sarah C Stainbrook*[1,3], Lindsey N Aubuchon[1], Amanda Chen[2], Emily Johnson[2], Audrey Si[2],
Laila Walton[2], Angela Ahrendt[2], Daniela Strenkert[3], Joseph Jez[1]
[1] Department of Biology, Washington University in St Louis, St Louis, MO
[2] Illinois Mathematics and Science Academy, Aurora, IL
[3] Plant Research Laboratory, Michigan State University, East Lansing, MI
* Correspondence: Sarah Stainbrook stainbr1@msu.edu
ORCIDs: Sarah Stainbrook: 0000-0002-0124-8230, Daniela Strenkert: 0000-0002-3420-1332,
Joseph Jez: 0000-0001-7911-9108

1 Abstract

C$_4$ crops such as *Zea mays* (maize) and *Sorghum bicolor* (sorghum) are crucial producers of
food and bioenergy. In these crops, carbon assimilation is limited at higher temperatures because the
enzyme rubisco activase (RCA), which is responsible for proper function of the CO$_2$ fixing protein
rubisco, has been shown to be thermolabile. We present a holistic approach to understand carbon
assimilation and RCA function in maize, sorghum, and the C$_4$ model grass *Setaria viridis* (setaria)
during heat stress acclimation. Gas exchange data confirms that assimilation is limited by Ribulose
1,5-bisphosphate (RuBP) carboxylation during heat. Plants express a variety of RCA isoforms and
we show that each species changes the isoform expression and proteoform abundances of these
variants in different ways. In addition, to understand how changing conditions in the chloroplast
stroma affect RCA function during heat, we examined the regulation of RCA activity by thioredoxin
F, magnesium ions and adenosine diphosphate. As expected, the activity of RCA is modulated by a
combination of these variables, but surprisingly, how these biochemical environment factors affect
RCA function differs vastly between species. Unexpected differences are even apparent within a
single species: different cultivars of maize show differences in assimilation, proteoform abundance
and regulation. Our surprising observation that each grass expresses and regulates RCA differently
suggests that even closely related plants use different strategies to maintain RCA function and we
conclude that a single reference system cannot accurately reflect protein function in every cultivar.

2 Keywords

28 C$_4$ photosynthesis, rubisco activase, heat stress, assimilation, abiotic stress, acclimation

3 Conflict of Interest

30 The authors declare that the research was conducted in the absence of any commercial or financial
31 relationships that could be construed as a potential conflict of interest.

4 Author contributions

Conceptualization: SCS. Methodology: SCS, LNA, AC, EJ, AS, LW, AA, JJ. Validation:
SCS, LNA, AC, EJ, LW, AA. Investigation: SCS, LNA. Analysis: SCS, LNA, AC, EJ, AS, LW.
Writing: SCS, LNA. Review and editing: SCS, LNA, AC, EJ, AS, LW, AA, DS, JJ. Funding: JJ, DS,
SCS. Supervision: JJ, DS.

5   Data Availability

All data is available upon request to the corresponding author at stainbr1@msu.edu.

6   Acknowledgments

We thank Ru Zhang for the initial inspiration of this work, Tom Sharkey, Oliver Mueller-Cajar and
Kevin Wyss for feedback on the manuscript, Leonardo Chavez for assistance with the LI-6400XT
machine and Mike Dyers for assistance with plant growth. This work was supported by the Amgen
summer scholar program (LNA) and an NSF Postdoctoral Research Fellowship in Biology #1907288
(SCS).

7   Introduction

Although ribulose-1,5-bisphosphate carboxylase/oxygenase (rubisco) is responsible for almost all carbon fixation on Earth, its lack of specificity between carbon (CO_2) and oxygen (O_2) results in photorespiration and loss of biomass. Both photorespiratory products and the ribulose-1,5-bisphosphate (RuBP) substrate itself can bind to the active site of rubisco and inhibit its function. Rubisco activase (RCA) has evolved in all land plants to release these inhibitory molecules from the active site and restore the activity of rubisco, rendering this protein essential for maintaining rubisco activity. RCA is an ATPase which functions *in vivo* in a hexameric ring structure (Portis 2003). However, RCA has a low thermotolerance and its activity is inhibited by even moderate heat (32.5^{o}C in maize) (Busch and Sage 2017; Steven J. Crafts-Brandner and Salvucci 2002). RCA's inability to function properly at higher temperatures has implications for crop yield given the rise in global temperatures.

In C_3 plants, photosynthesis takes place entirely within the mesophyll in the presence of O_2 produced by Photosystem II, leading to photorespiration and costly corrections. Photorespiration, and its associated metabolic burden, increase with rising temperatures (Sage 2013). This has led some to propose that the thermolability of RCA is in fact a protective mechanism to disable photosynthesis under unfavorable conditions, similar to a fuse (Degen, Worrall, and Carmo-Silva 2020).

In C_4 plants, rubisco functions exclusively in the bundle sheath cells where O_2 is not produced. This adaptation improves the efficiency of photosynthesis by limiting photorespiration to less than a fifth of that of comparable C_3 grasses (Kanai and Edwards 1999). The decrease in photorespiratory products lessens the inhibition of rubisco. The kinetics of rubisco improve with increasing temperature in the absence of O_2, potentially allowing C_4 plants to photosynthesize more efficiently at higher temperatures (S J Crafts-Brandner and Salvucci 2000). However, the RCA expressed in C_4 plants is also thermolabile and is thought to limit photosynthesis unnecessarily. The RCA of C_4 plants is therefore an attractive target for protein engineering with the goal to improve its thermostability and to increase CO_2 assimilation when plants experience high temperatures (Parry et al. 2013).

Most plant species express two RCA isoforms, α and β, which arise from alternative splicing of a single gene in some species and from dedicated genes in others. The β isoform is approximately 43 kDa and can undergo post-translational proteolytic cleavage to form a 41 kDa proteoform (Z. Yin et al. 2014; Vargas-Suárez et al. 2004). In most species, the β isoform is constitutively expressed. The α isoform is approximately 45kDa and is mostly identical to the β isoform with the addition of a C-terminal domain (CTD). The CTD contains two cysteines capable of forming a disulfide bond which regulates RCA function according to the redox state of the chloroplast. Redox state dependent regulation of the α isoform couples the activity of RCA to illumination (N Zhang and Portis 1999). The α isoform has also been reported to undergo inhibitory phosphorylation at threonine 78 near the N-terminus (Kim et al. 2019), but this threonine is not conserved in all species and is not present in the β isoform. The α and β monomers can differ in thermostability, with the α isoform frequently being more thermostable (Steven J Crafts-Brandner et al. 1997).

The thermolability of RCA may be due to structural changes such as denaturation and aggregation (Barta et al. 2010). However, RCA may also be inactivated by other unfavorable conditions inside the chloroplast. The ATP/ADP ratio in the chloroplast affects RCA function in some species (A. E. Carmo-Silva and Salvucci 2013; Barta et al. 2010; Kallis, Ewy, and Portis 2000). This is likely due to competitive binding of ADP to the ATP hydrolysis site that interferes with RCA's function, but the presence of ADP can also improve the thermostability of RCA (Henderson et al. 2013). RCA is also sensitive to the concentration of Mg^{2+} ions in the stroma (Hazra et al. 2015; Barta et al. 2010), possibly through the function of Mg^{2+} in coordinating the binding of ATP. The sensitivity of RCA to ADP and to redox regulation are attributed to the CTD in the α isoform, where

redox regulation occurring at the disulfide bond favors the binding of ADP while destabilizing the
binding of ATP (Wang and Portis 2006; Scafaro, De Vleesschauwer, et al. 2019). Because α and β
RCA isoforms form hetero-oligomeric complexes, the thermostability and activity of the entire
functional complex is regulated by the conditions affecting the α isoform (Keown and Pearce 2014)
although denatured isoforms are excluded from complexes under high temperature conditions
(Scafaro, Bautsoens, et al. 2019).
Efforts to find or create thermostable RCA isoforms have been advocated widely and have
already shown moderate success (Davidi et al. 2020; Kurek et al. 2007; Scafaro, Bautsoens, et al.
2019; Qu, Mueller-Cajar, and Yamori 2023). To date, these efforts have not considered possible
effects of stromal conditions or post-translational modifications on either native or novel enzymes,
making it likely that potential pitfalls and promising avenues for improvement have been missed.
Protein engineering efforts are currently hampered by an incomplete understanding of RCA
regulation in its native environment: the dynamically changing chloroplast stroma. Toward this end,
we examined whether assimilation is truly limited by RCA in different cultivars of the C_4 food crops
Zea mays (maize) and in *Sorghum bicolor* (sorghum) and in the common C_4 model plant *Setaria*
viridis (setaria). RCA isoform expression was determined in each species, and we examined how
stromal conditions regulate RCA function in these plants before and during heat treatment. Our
results expose potential avenues for improvement of RCA with the long-term goal to bioengineer
crop species that maintain photosynthetic performance despite rising global temperature.

8   Materials and Methods

8.1   Plant Material

S. viridis A10 was a gift from the lab of Dr. Ivan Baxter at the Donald Danforth Plant Science
Center. Seeds were sown 1cm deep in 18-well trays and thinned to one plant per cell. Plants were
grown in the Washington University in St Louis greenhouse with 28°C/22°C temperatures and a 16-
hour photoperiod.
Z. mays B73 was obtained from the greenhouse at Washington University in St Louis. All
other maize accessions were obtained from the USDA GRIN catalog. *S. bicolor* BTx623 was a gift
from the lab of Dr. Malia Gehan at the Donald Danforth Plant Science Center. Sorghum and maize
were planted 2cm deep in a 4-inch pot in a 50% mixture of Berger Bark Mix and Turface and grown
in the Washington University in St Louis greenhouse with 28°C/22°C temperatures and a 16-hour
photoperiod. Plants were grown to the two- or three-leaf stage, then transplanted into 6-inch pots in
the same soil mix.

8.2   Heat Treatment

Greenhouse-grown plants were moved into a growth chamber of the appropriate temperature
(42°C for 42°C measurements, etc.). Chambers were kept at 50% humidity and plants were
maintained with the bottom of the pot in standing water to avoid drought stress. For 1-hour
measurements, the leaf was acclimated in the gas exchange chamber of the LI-6400XT instrument
with the instrument controlling the leaf temperature. For longer measurements, the gas exchange
chamber was attached to the leaf twenty minutes prior to the measurement to allow stomata to
equilibrate. For 48-hour measurements, plants were maintained at the same photoperiod as the
greenhouse with 42°C day and 38°C night temperatures. All heat treatments were initiated and all gas
exchange measurements were taken at least two hours from either end of the photoperiod.

8.3   Gas exchange measurements

Gas exchange measurements were performed on the most recently fully expanded leaf. The
leaf measured was from the main stem of each plant, and all measurements were taken prior to
panicle emergence. CO_2 concentration within the gas exchange chamber was controlled at 400ppm.
Gas exchange measurements were performed at saturating illumination of 1400 µmol m^{-2} s^{-1} (*S.*
viridis) or 2000 µmol m^{-2}s^{-1} (*Z. mays*, *S. bicolor*). Humidity was controlled to maintain stomatal
conductance above 0.3 for all measurements. The same plants were measured at the 1, 6 and 48 hour
time points whenever feasible.
A/Ci curve measurements were taken according to the standard method (Long and Bernacchi
2003) and were fitted using the Excel tool from (Zhou, Akçay, and Helliker 2018), which
implemented the model of (X. Yin et al. 2011) with the following modifications: changes to the
constants found in cells P22, Q22, D126 and D127 (calculations of variables x_1 and x_2 for the
electron transport limited state) were corrected to match the equations from (X. Yin et al. 2011), and
α (the fraction of O_2 evolution occurring in the bundle sheath) was assigned the value of 0 as PSII
activity is negligible in bundle sheath cells in NADP-ME type monocot C_4 photosynthesis (Von
Caemmerer and Furbank 1999). For all measurements, the first measurement at 400µM CO_2 was
checked for stomatal limitation according to the equation in (Long and Bernacchi 2003)and any
dataset in which stomatal conductance limited photosynthesis by more than 30% was discarded. This
initial measurement was also compared to the A/Ci curve fit to determine the limitation to
photosynthesis. Representative A/Ci curves and fits can be found in Figure S2, and all A/Ci curve fit
variables may be seen in Table S1.

8.4   Tissue harvest and protein extraction

Leaf tissue from plants subjected to heat in growth chambers was harvested during the middle
of the photoperiod, flash-frozen in liquid nitrogen, ground with mortar and pestle and pooled prior to
storage at -80°C. All tissue samples contained leaves from at least 3 plants. Care was taken to
exclude the ligule, the midrib, and leaves that were not yet fully expanded or which were senescent.
RCA was extracted by ammonium sulfate precipitation according to the protocol of (Barta, Carmo-
Silva, and Salvucci 2011a) with the modification that the 4.1M ammonium sulfate was adjusted to
pH 4.6 instead of 7.0. Purified protein was resuspended in Leaf Resuspension Buffer (50 mM
HEPES-KOH, pH 7.0, 2 mM $MgCl_2$, 100 mM KCl, 5 mM dithiothreitol, 1 mM ATP, 1 mM PMSF,
and 10 mM leupeptin.) Aliquots were stored at -80°C until use. When compared to samples purified
using the complete protocol of (Barta, Carmo-Silva, and Salvucci 2011a), samples using only the
modified ammonium sulfate precipitation method were considerably cleaner and gave similar thermal
curve results (Figure S5). Tissue was harvested from sorghum at 7 hours of heat treatment rather than
6 due to a gun-related lockdown at the ideal time of harvest.

8.5   Western blotting

Crude protein extracts from the first step of purification were resolved by SDS-PAGE on a
12.5% gel and transferred onto PVDF membrane using a semidry transfer apparatus according to the
manufacturer's instructions. 20µg of crude extract was used for maize, setaria and sorghum, and 5ug
was used for tobacco, Chlamydomonas, Arabidopsis and spinach. The membrane was blocked with
5% w/v nonfat dry milk in PBS supplemented with 0.1% v/v Tween-20 (PBST) with shaking at 4°C.
Primary antibody (anti-RCA, rabbit, 1:10000 Agrisera) was added and incubated for one hour. The
membrane was thoroughly washed with PBST and bound with secondary goat anti-rabbit-HRP
(MilliPore) at 1:50,000 for 1 hour. The membrane was thoroughly washed and visualized with the
FemtoGlow Western Plus chemiluminescent diagnostic kit (Michigan Diagnostics) according to the
manufacturer's instructions.

8.6 Recombinant protein production

The coding sequence of Mesembryanthemum crystallinum thioredoxin F was codon-optimized for expression in *Escherichia coli*, synthesized and cloned into pET28a with an N-terminal 6xHis tag. Transformed BL21-DE3 was grown to OD_{600} 0.6-0.8, induced with 1mM IPTG and incubated at 12°C overnight. TrxF was purified by Ni-NTA affinity chromatography in buffer containing 50mM HEPES pH 8.0, 0.5M NaCl, 5% glycerol, 10mM β-mercaptoethanol and 5-300mM imidazole, then dialyzed into the same buffer without imidazole and stored in aliquots at -80°C. TrxF activity was verified using the insulin reduction assay (Holmgren 1979).

8.7 Biochemical assays

RCA thermostability was measured with an NADH-linked ATPase enzymatic assay in microtiter plates based on the protocol of (Barta, Carmo-Silva, and Salvucci 2011b). Briefly, 20µL of purified RCA was aliquoted into 0.2mL tubes and incubated in a gradient thermocycler for 1 hour at temperatures ranging from 30-66°C and returned to ice until measurement, while a control aliquot was maintained on ice. RCA sample (5µL) was added to 120µL of reaction mixture in microtiter plates and the change in A_{340} was monitored for 10 minutes using a Tecan plate reader. ATPase activity was calculated as the initial slope of the line (change in absorbance per minute). T_{50} was calculated by fitting a 4-parameter logistic equation to the dataset using the calculator at (AAT BioQuest Inc, 2022), see example in Figure S4. Temperature response data that showed two distinct regions of response (sorghum and tobacco) were divided into two separate curves and each curve was fitted separately to produce two T_{50} values. The two plateaus are unlikely to represent a true response of RCA to heat. Previous work in spinach has shown that mixing of isoforms of different thermal stabilities confers an overall RCA response intermediate between the individual isoforms (Keown and Pearce 2014). Rather, the lower of the two T_{50} values is likely another ATPase present which was not fully removed by our purification technique. The portion of total ATPase activity due to the low-thermal-stability ATPase was calculated and subtracted from total ATPase activity results.

Response of ATPase activity to thioredoxin F, Mg^{2+} and ADP:ATP ratio was measured according to the method of (Chifflet et al. 1988; N Zhang and Portis 1999). Briefly, 40ug of purified enzyme was incubated in 100mM Tris-HCl pH 8.0 with 5mM DL Dithiothreitol with or without 2.6ug of TrxF as indicated for at least 20 minutes at room temperature to allow reduction of RCA. ATPase activity was then assayed in a total volume of 50uL of reaction buffer (50mM tricine pH 8, indicated concentration of $MgCl_2$, 20mM KCl, 4mM total of ATP and ADP according to ratio) for 1 hour. Sodium dodecyl sulfate and ammonium molybdate reagent were added and allowed to react for 5 minutes. Color development was terminated by addition of sodium meta-arsenite reagent and A_{850nm} was read. Because the amount of ATP and the presence or absence of TrxF contributed different amounts to the background, no-RCA controls were performed for all treatments and the background was subtracted from all samples prior to statistical analysis. Measurements were normalized to ATP:ADP=1:0 without TrxF for all time points.

8.8 Statistical Analysis

Outliers were removed by Grubbs' test, $p < 0.05$, and significant differences between treatments were determined by T-test with Bonferroni correction. A two-way ANOVA was used to examine the interaction between ATP:ADP ratio and redox regulation.

9 Results

9.1 Assimilation

We tested how pretreatment with elevated temperature affected assimilation in different crop
species. Assimilation was optimal in all species when plants were exposed to 32°C and 35°C for 1 h
prior to measurement but it decreased significantly at temperatures above 42°C (Figure S1).
Individual plants were then measured during the course of acclimation over 48 hours. At 0 hours
(before heat treatment) and 48 hours (after acclimation), all replicates gave similar values, because
the plants' physiology and biochemistry were in a stable state. However, at the more transient time
points (after 1 hour and 6 hours of heat treatment), the plants' physiology and biochemistry were
transitioning between the two stable states, and individual plants produced different assimilation
values. Although all plants experienced the same condition for the same time, there is variability in
how dramatically and how quickly each plant responded to the heat. The assimilation values at 1
hour showed the highest variability, which we attribute to the inherent variability in measuring a
transient phenomenon

236
Fig 1: assimilation and C_i during heat acclimation. **A**. Assimilation of various cultivars of maize. **B**. Assimilation of
maize, sorghum and setaria. **C**. Ci of maize, sorghum and setaria. D. Vcmax of maize, sorghum and setaria. * $p<0.05$, **
$p<0.01$ vs. 0-hour control. Error bars are standard error of at least 3 biological replicates.

To examine how different cultivars of the same species respond to heat pretreatment, we
measured the CO_2 assimilation of a total of 9 cultivars of maize in greenhouse conditions and after 1
and 48 hours at 42°C (Figure 1A). The common maize reference cultivar B73 showed an early
decrease in assimilation at 1 hour but recovered within 6 hours. This response was consistent
between four other cultivars (KI3, Tzi8, MR25, Tx303), albeit with less significance (Figure 1A).
More importantly, we observed cultivar-specific differences within the same species. For example,
one cultivar, MR19, retained high assimilation after 1 hour but displayed significantly reduced
assimilation after 48 hours of heat exposure. This cultivar was chosen for further study because of its
failure to acclimate to heat.
We also examined C_4 species sorghum and setaria on their ability to assimilate CO_2 in
response to heat pretreatment to observe inter-species variation. We measured one additional time

point 6 hours after heat exposure to increase temporal resolution of the experiment. We included the
common maize cultivar B73 as well as MR19 because of its failure to acclimate to heat. Assimilation
values in sorghum and setaria were not significantly affected by 42°C heat at any time point (Figure
1B). Intercellular CO_2 concentration (C_i) was measured for all species at normal and elevated
temperature at 400ppm CO_2 (Figure 1C). All species exhibited elevated Ci at high temperatures. In
both maize cultivars, the Ci more than doubled within 1 hour and remained elevated. In sorghum, Ci
remained low until the 48-hour time point. In setaria, the increase in Ci was transient, and Ci returned
to pre-treatment values within 6 hours.

9.2 Limitations to Photosynthesis

The response of assimilation to changes in intercellular CO_2 concentration (Ci) can be
measured to determine the limitations to photosynthesis. At low C_i, photosynthesis is limited
primarily by the speed of the carboxylation reaction carried out by rubisco. At higher C_i,
photosynthesis is primarily limited by the ability of the CBB cycle to regenerate the RuBP substrate
of rubisco, which in turn is limited by the rate of electron transport to regenerate reducing equivalents
for this process (Long and Bernacchi 2003). In C_4 plants, which use phosphoenolpyruvate (PEP) to
transport CO_2 from the mesophyll cells to the bundle sheath cells, the speed of PEP carboxylation or
regeneration can also limit assimilation. We used A/Ci curve measurements to assess which
limitations applied in our plants under heat-exposed conditions. Briefly, four different mathematical
models are used to determine the limitations to assimilation posed by the C_4 cycle and by the CBB
cycle. These models are RcPc (limited by RuBP carboxylation and PEP carboxylation), RcPr (limited
by RuBP carboxylation and PEP regeneration), RrPc (limited by RuBP regeneration and PEP
carboxylation), and RrPr (limited by RuBP regeneration and PEP regeneration.)
We measured A/Ci curves in triplicate for each of the four cultivars after 0, 1, 6 and 48 hours
of heat exposure and fit the measurements for each plant to the model of C_4 photosynthesis to
determine the limitation to photosynthesis at atmospheric CO_2 levels (400ppm) as described in
(Zhou, Akçay, and Helliker 2018)(Figure S2 and Table S1). For both maize cultivars at all time
points, assimilation was limited by RuBP carboxylation and PEP carboxylation (model RcPc).
Setaria was also limited by RuBP carboxylation at all time points, and all plants except one were
limited by PEP carboxylation (model RcPc.) In sorghum, RuBP carboxylation was limiting at 1 and 7
hours of heat treatment (model RcPc), but by 48 hours RuBP regeneration became limiting (model
RrPc). Prior to heat treatment, different sorghum plants showed limitation by RuBP carboxylation or
by RuBP regeneration (models RcPc and RrPc.) Our data implies that the carbon fixation reaction
catalyzed by rubisco is the rate-limiting step for all plants during the first several hours of heat
exposure prior to acclimation.
The A/Ci curve fits five parameters for photosynthesis, including Vcmax, the maximum
carboxylation rate of rubisco. We found that during heat treatment, Vcmax was decreased during the
1 hour time point in all species, though not significantly (Figure 1D). In maize cultivar MR19, which
failed to recover assimilation at the 48 hour time point, Vcmax was also significantly compromised at
48 hours. The close match of assimilation and Vcmax trends, as well as the limitation of RuBP
carboxylation in all plants at 1 and 6 hours, indicates that rubisco has suboptimal activity at these
time points, likely due to thermal limitations to RCA.

9.3 RCA Proteoform Abundance

As expected, heat inactivation of RCA seems to be a major bottleneck in maintaining
photosynthetic performance during heat acclimation in all plants investigated in this study. To
determine how plants overcome this limitation, we assessed RCA abundance and proteoform

composition in maize, sorghum and setaria. Immunodetections were performed on crude leaf tissue
extracts from these species before and after heat treatment (Figure 2). Additional samples from
reference species with known RCA isoform composition were examined for comparison. The C_3
plants examined here (*Nicotiana tabacum, Spinacia oleracea* and *Arabidopsis thaliana,* Figure 2)
contained more RCA per mg total protein than Setaria did prior to heat stress, in agreement with
previously published results (Kanai and Edwards 1999). The lower band intensity for the alga
Chlamydomonas reinhardtii could indicate a lower RCA abundance allowed by its known carbon
concentrating mechanism (CCM), or may indicate lower antibody affinity for this more distant
homolog. RCA abundance was reduced in all C_4 species after 48 hours compared to the peak RCA
abundance. Both maize cultivars showed a transient increase in RCA abundance after 6 hours, before
RCA abundance was reduced substantially after 48 hours. RCA was initially much more abundant in
maize MR19 than in maize B73 prior to heat treatment (Figure 2). Similar to maize, sorghum and
setaria showed a slight depletion in RCA abundance at the 48-hour time point, although with
sorghum there was first a transient increase.

Fig 2: RCA abundance and proteoform distribution during heat acclimation. Immunodetections were done on 20µg of
crude protein extract for C_4 species and 5µg of crude protein extract for controls. Cr *C. reinhardtii*, So *S. oleracea*, At *A.*
thaliana, Nt *N. tabacum*. Times are indicated for each sample in hours. Complete blots may be viewed in the supplement.
Black line indicates 37kDa marker; proteoforms are indicated by shaded arrows according to the legend
315

We confirmed the presence of proteoforms that have been described previously in the
literature for Arabidopsis and spinach samples (α at 45kDa, β at 43kDa and a proteolytic product of β
at 41kDa). We know that Tobacco and Chlamydomonas express solely the β isoform, and in our
detections both species lack the higher molecular weight bands corresponding to the α
isoform. While Chlamydomonas shows only a single proteoform, tobacco displays several
proteoforms of β, including a band at a slightly lower molecular weight of about 39kDa that we
suggest to be another proteoform of β (see complete blots in Figure S3). In both maize cultivars and
in sorghum, the phosphothreonine proteoform of α appeared after heat treatment. This threonine is
not conserved in Setaria, and the higher molecular weight form of α is not observed in our
immunodetections. Other changes in proteoform abundance were specific to the species or cultivar.
In Setaria the α isoform is depleted and the 41kDa proteolytic product is transiently enriched. The α
isoform was transiently depleted in MR19 after 1 hour but recovered during acclimation.

9.4   Regulation by variable components within the chloroplast stroma

We next characterized the effects of the native setting within the chloroplast on RCA's
enzymatic ATPase activity. Recombinantly produced proteins are commonly used for biochemical
assays due to the ease of sample preparation and the high sample purity. However, since multiple
RCA proteoforms are present in varying ratios *in vivo*, we decided to purify RCA directly from plant
tissue for our assays. This is particularly important given that the dynamically changing,
combinatorial interaction of different proteoforms in hetero-oligomers determines the activity of
RCA, (Keown and Pearce 2014) and this mixture of proteoforms is impossible to accurately
reproduce with recombinant protein. In addition, post-translational modifications that are known to
affect RCA activity are also not reliably reproduced in a recombinant expression system. For these
reasons, we used RCA purified from plant tissue for biochemical characterization.

9.4.1 RCA activity

We next examined how changes in RCA abundance caused by heat stress affect RCA
activity. We measured the ATPase activity of 40μg purified RCA from maize, sorghum and setaria
and each time point to determine the activity per μg of RCA protein. We then scaled the amount of
protein for each reaction by the abundance of RCA relative to the 0 hour control as determined from
our immunodetections. This analysis allows us to distinguish whether differences in RCA activity are
due to changes in the enzyme activity per μg of RCA, or simply the result of increased RCA protein
abundance. In maize cultivar B73, RCA activity was increased but we attribute that increase to the
observed increase in RCA protein expression and not to increased RCA activity (Figure 3A). This
contrasts with the other maize cultivar (MR19), in which RCA activity and abundance both
decreased in response to the heat treatment (Figure 3B). In sorghum, the increase in overall RCA
activity at the 1- and 6-hour time points were more than accounted for by the increase in activity per
μg of RCA (Figure 3C). In setaria, activity per μg of RCA was dramatically increased at the 48-hour
time point, leading to a modest increase in overall RCA activity (Figure 3D).

Fig 3. ATPase activity of RCA. **A**. Maize B73 **B**. maize MR19 **C**. sorghum **D**. setaria. Black = activity per 40ug; white = total activity. Letters denote significance within data series of p<0.05. Error bars are standard deviation of n=5

9.4.2 RCA thermostability

Our data indicate that different plant species (and even different cultivars of the same species) show enormous flexibility in how RCA abundance and activity change during heat acclimation. We next sought to determine how these changes affect RCA thermostability. We measured RCA thermostability by pre-incubating purified enzyme for 1 hour at temperatures ranging from 30°C to 66°C, then measuring activity using a spectroscopic ATPase assay based on NADH consumption. The resulting heat response curves were fitted to determine the temperature at which half of the activity was lost (T_{50}). As expected, spinach RCA had a lower T_{50} than tobacco RCA (37.6°C vs. 48.0°C). The thermostability of RCA extracted from both maize cultivars varied with time but never increased beyond the 0-hour control. In contrast, both Setaria and sorghum RCA displayed increased T_{50} at various times, with Setaria increasing from 43.6°C (0-hour) to 49.0°C (6 hours) and sorghum increasing from 47.9°C (0 hour) to 50.3°C (48 hours).

Fig 4. temperature sensitivity of ATPase activity. * p<0.05, ** p<0.01 relative to 0 hours, error bars are standard
deviation of n=3 technical replicates

9.4.3 Magnesium ion sensitivity

Another aspect that affects RCA activity is the availability of magnesium. We found that
different species responded differently to Mg^{2+} concentration in the reaction buffer (Figure 5). RCA
from Arabidopsis, tobacco, spinach and Chlamydomonas were significantly more active with 5-
10mM Mg^{2+} as compared to background (0.1mM Mg^{2+}, Figure 5A), which is consistent with what
has been previously reported for tobacco (Hazra et al. 2015). The same is true for RCA activity in
Sorghum: supplementing 10mM Mg^{2+} increased RCA activity by 43% (Figure 5D). However, heat
treatment transiently dampened the response to Mg^{2+} in sorghum (<10% increase at 1 and 7 hours of
heat treatment). By 48 hours of heat treatment, Mg^{2+} stimulation had largely recovered (23%
increase).

To our surprise, both maize B73 and setaria showed inhibition by magnesium concentrations
greater than background (Figure 5B, C and E). For maize, the inhibition was dramatic (34% activity
inhibited by 10mM Mg^{2+}, p<0.01), but the response to magnesium was eliminated after 6 hours of
heat treatment. For setaria, the inhibition was less pronounced (9% at 10mM Mg^{2+}, not significant)
and only rose to the level of significance after 48 hours of heat treatment (16% inhibition at 10mM
Mg^{2+}, p<0.01). Unlike B73 RCA, which was strongly inhibited by magnesium, RCA from MR19 is
insensitive to magnesium at all points of heat treatment.

Fig 5. response of ATPase activity to changes in Mg^{2+} concentration. All measurements performed with ATP:ADP=4:0. **A**. Controls **B**. Maize B73 **C**. maize MR19 **D**. sorghum **E**. setaria. * p<0.05, ** p<0.01 relative to 0.1mM Mg^{2+}, error bars are standard deviation of n=5

9.4.4 ADP sensitivity

Previous work has established that the ATPase activity of RCA is inhibited by increasing concentrations of ADP (Ning Zhang, Schürmann, and Portis 2001). We were able to replicate these results with RCA from tobacco, Arabidopsis, Chlamydomonas and spinach (Figure 6A). RCA from setaria was similarly inhibited by ADP but became insensitive to ADP after 48 hours of heat treatment (Figure 6E). Maize B73 RCA was initially insensitive to ADP, but after heat treatment was mildly stimulated (15% increase, ATP:ADP of 1:3, p<0.01, Figure 6B). Similar to B73, MR19 was

stimulated by ADP, but to a greater extent (64% increase, ATP:ADP = 1:3, p<0.01, Figure 6C).
Unlike B73, however, the increase in activity was present at all time points of heat treatment.
Sorghum RCA was initially stimulated by ADP (12% increase, ATP:ADP of 1:3, p<0.01, Figure
6D), but lost sensitivity to ADP with heat treatment.

Fig 6. response of ATPase activity to changes in ATP:ADP ratio. All measurements performed with 10mM Mg^{2+}. **A**.
Controls **B**. maize B73 **C**. maize MR19 **D**. sorghum **E**. setaria * p<0.05, ** p<0.01 relative to 4:0 ratio, error bars are
standard deviation of n=5

9.4.5 Redox sensitivity

Fig 7: response of ATPase activity to TrxF relative to 4:0 ratio. ATP:ADP ratio was 3:1 and 10mM Mg^{2+}. **A**. Controls **B**.
maize B73 **C**. maize MR19 **D**. sorghum **E**. setaria. * $p<0.05$, ** $p<0.01$ vs. no TrxF, error bars are standard deviation of
n=5

The α isoform of RCA is redox regulated by a disulfide bond only present in the CTD of the α
isoform. This regulation may happen through the enzyme thioredoxin F (TrxF) (N Zhang and Portis
1999) or directly from Photosystem I (Kale et al. 2022; Bhaduri et al. 2020). We tested RCA activity
in the presence and absence of TrxF. We reproduced published stimulation of Arabidopsis and

spinach RCA ATPase activity by TrxF, and the expected lack of a response by tobacco and
Chlamydomonas, both of which lack an α isoform (Figure 7A). Maize B73 RCA was slightly
stimulated by treatment with TrxF (15%, $p<0.05$, Figure 7B). This stimulation was transiently
eliminated at 1 and 6 hours of heat treatment but reappeared after 48 hours of heat treatment. Maize
MR19 was initially insensitive to regulation by TrxF, but by 48 hours showed a 32% increase in
activity with TrxF treatment ($p<0.01$, Figure 7C). Surprisingly, we found that treatment with TrxF
slightly inhibited RCA from sorghum (decreased by 6%, $p<0.01$, Figure 7D) at 0 and 6 hours of heat
treatment. Setaria RCA appeared insensitive to TrxF regardless of heat treatment (Figure 7E).

10   Discussion

10.1 Assimilation and intercellular CO_2

Our data generally do not show a significant decrease of CO_2 assimilation in response to heat
stress as compared to untreated control plants. However, because the activity of purified, fully active
rubisco is more than doubled at 42°C compared to 28°C (S J Crafts-Brandner and Salvucci 2000), the
lack of a comparable increase in assimilation should be interpreted as a dramatic impairment of
assimilation at 42°C in all plants examined.
Among cultivars of maize, some acclimate to 42°C while maintaining stable assimilation
(such as MR15 and M162W) while others transiently decrease assimilation (B73, Ki3, and Tx303)
(Figure 1c). The variability of this phenotype, presumably due to genetic diversity within the species,
indicates that traditional breeding approaches may be fruitful in improving high-temperature
acclimation in maize. Among all cultivars examined, maize MR19 uniquely failed to acclimate to
higher temperatures at all and exhibited reduced assimilation after 48 hours of heat treatment. This
cultivar may be useful in future studies to determine the mechanism(s) underlying thermal
acclimation. Variability in the thermal tolerance of assimilation and RCA stability between cultivars
of the same species has previously been reported in rice (Scafaro et al. 2012), but the causal genetic
variant has not yet been identified in maize.
Recent studies (Anderson et al. 2021; Kim, Slattery, and Ort 2021) report decreased
assimilation in setaria after 1 hour of treatment at 40°C and in setaria and sorghum after 1 hour of
treatment at 42°C respectively. These trends are not present in our results. As we have demonstrated
significant variability in assimilation response to heat among maize cultivars, these differences could
be explained by differences in the cultivars of setaria and sorghum used between these studies and
ours. Alternatively, increasing the number of plants measured and improving temporal resolution
may reveal these differences to be significant.
The elevated C_i we observed at many time points can be caused by a variety of biochemical
and physiological factors. Increased stomatal opening can elevate C_i while also allowing
transpirational cooling of the leaves. Stomatal conductance was significantly elevated during all heat
treatments for maize B73, matching elevated C_i (Figures 1 and S1). However, stomatal conductance
but does not correlate with C_i in maize MR19 or setaria. In these plants, biochemical factors must
also be considered. Decreased efficiency of carbon fixation at rubisco (e.g., by inactivation of RCA)
or at the initial C_4 fixation steps in the mesophyll cells (e.g., PEPC) may cause an increased
equilibrium concentration of C_i. Correspondingly, our data indicate that during heat acclimation
plants are limited by carbon fixation at rubisco and by PEP regeneration. High temperature also
decreases the mesophyll conductance in C_3 plants such as tobacco (Bernacchi et al. 2002) and
spinach (Yamori et al. 2006), which could limit transfer of CO_2 from the intercellular space into the
cytosol of mesophyll cells, but this has not been directly examined in C_4 plants.
In sorghum, one plant at 0 hours and all three plants at 48 hours were limited by RuBP
regeneration rather than by RuBP carboxylation. At 48 hours we also observed a strong increase in

the C_i which was not present at other time points and which cannot be explained by changes in
stomatal conductance (Figure S1). Vcmax is also not significantly increased at 48 hours (Figure 1D).
As the expression and activity of RCA in sorghum at 48 hours are very similar to the 0-hour control
(Figure 2), this may indicate that the limit of rubisco carboxylation speed is being overcome by
increasing the concentration of CO_2 available to rubisco by limiting the speed of RuBP regeneration.
Others have demonstrated that increased C_i improves the thermotolerance of photosynthesis in C_3
plants (Hikosaka et al. 2006; Kattge and Knorr 2007), specifically by increasing the concentration of
CO_2 at rubisco and improving its kinetics. Thus, increased C_i may compensate for the thermal
limitation of RCA to some extent. However, it is apparent that merely increasing Ci is not sufficient
to overcome rubisco carboxylation limits in all circumstances, as both maize cultivars showed
limitation by RuBP carboxylation despite elevated C_i (B73 at 1 hour, MR19 at 48 hours).
In our work, plants were kept well-watered so that water limitation would not convolute the
heat response; however, in most real-world scenarios elevated heat often co-occurs with water
limitation. In such cases, stomatal opening is limited to retain moisture, and C_i may be
correspondingly decreased. Water limitation may thus decrease rubisco efficiency and exacerbate the
effects of heat on assimilation. (A. Elizabete Carmo-Silva et al. 2012).

10.2 Proteoform abundance

Molecular and biochemical assays also highlight the differences among species. Recent
transcriptomics work by (Anderson et al. 2021) in setaria found induction of RCAα in response to
heat stress, but no change in RCAβ expression. Our immunodetections showed a decrease in the
abundance of α; however, our higher temperature of 42°C and different cultivar (A10 vs. ME034)
may account for these differences. Similarly, (Kim, Slattery, and Ort 2021) examined RCA
expression in sorghum after 6 hours of heat treatment at 42°C using qRT-PCR. They report induction
of RCAα, which we confirm. However, we caution that post-transcriptional regulation creates
proteoforms with distinct molecular weights and activity profiles but is not detectable by
transcriptomic analysis. Using Western blotting, we demonstrate that heat treatment changes the
phosphorylation state of the α isoform and also alters amounts of limited proteolysis of the β isoform
resulting in the 41kDa proteoform. Further study on the exact post-translational modifications used
by each species under various conditions should be pursued.

10.3 Effects of the internal chloroplast environment

Changes in the external environment also affect the plant's internal environment. For example,
the presence of light decreases the pH of the lumen, increases the available pool of ATP (increasing
the ATP:ADP ratio) and allows electron flow through TrxF to reduce target proteins (Nikkanen and
Rintamäki 2014). Similarly, Mg^{2+} concentration in the stroma is known to increase with illumination
(Ishijima et al. 2003). We speculated that a combination of the changes in proteoform abundance and
the thylakoid environment may combine to determine RCA activity.
Previous studies report that magnesium is important both for coordinating binding of ATP to
RCA, and for coordinating binding and cooperativity of RCA monomers in the hexameric state
(Hazra et al. 2015). Furthermore, Mg^{2+} destabilizes RCA in the presence of ADP but stabilizes it in
the presence of ATP (Henderson et al. 2013). Tobacco RCA is known to be activated by increasing
concentrations of magnesium, a result which we reproduced (Figure 5). Maize B73 was uniquely
inhibited by magnesium ions, even at the lowest concentration studied and in the absence of ADP.
This inhibition was modulated by heat treatment. Notably, the 1-hour time point at which maize B73
RCA was most strongly inhibited by magnesium was also the time at which assimilation was
significantly compromised and thermostability of RCA was decreased. We are unaware of any

mechanism reported in literature that could explain inhibition by magnesium ions. The correlation
between increased magnesium sensitivity and decreased RCA function at high temperature should be
examined more closely in maize. However, this mechanism is unlikely to translate to other C_4
grasses, which each show different responses to magnesium.
Responses of maize and sorghum to ADP concentration (Figure 6) differed from those
reported for other species in the literature and may have regulatory significance. The inhibition of
RCA by ADP has been explained as competition for the same binding pocket and is a regulatory
mechanism that prevents use of ATP by the CCB Cycle until the plastid triphosphate pool can be
replenished. Others have shown that ATP induces oligomerization while ADP encourages formation
of monomers in spinach RCA (Keown and Pierce 2014). Increased activity stimulated by ADP has
not previously been reported but is seen in sorghum and in both cultivars of maize. We are unaware
of any known mechanism by which the presence of ADP would stimulate the ATPase activity of
RCA. Furthermore, the stimulation of RCA by ADP is altered during heat treatment, with different
species showing different responses. Others have previously speculated that the inactivity of RCA at
high temperature in C_3 plants is due to unfavorable ATP:ADP ratios (S J Crafts-Brandner and
Salvucci 2000), but have noted that this effect is not present in maize (Steven J. Crafts-Brandner and
Salvucci 2002). Our result corroborates this explanation and begs further study into the mechanism
and significance of ADP stimulation of RCA activity.
Redox regulation of RCA by TrxF is postulated to couple the electron transport chain to
carbon fixation. In past work, the effect of TrxF reduction of RCAα has been shown to be stimulatory
for Arabidopsis and spinach. We here show that this stimulatory effect is present in two maize
cultivars and is modulated by heat treatment. Setaria and sorghum appear to be insensitive to redox
regulation, despite the expression of the α isoform with an intact C-terminal disulfide domain. The
lack of redox sensitivity may be due to incompatibility with the TrxF isoform used in this study or
may reflect a true difference in regulation among the C_4 species which could be used to more deeply
understand the mechanisms and residues involved in redox regulation.

11   Conclusions

Heat acclimation is a dynamic process involving changes in both thylakoid environment and
proteoform abundance, and changes in plant physiology including stomatal conductance and
intercellular CO_2 concentration. All these parameters must be considered to create an holistic
understanding of the mechanisms contributing to thermal limitation in each plant at each time point.
The increased activity of RCA in setaria and sorghum is not correlated with increased
abundance of RCA, and conversely, increased abundance of RCA in MR19 does not correlate with
increased activity. This indicates that the strategy of overexpressing RCA is unlikely to improve
assimilation at high temperatures in these plants, although it has shown promise in other species (Qu
et al. 2021). Furthermore, the thermostability of the enzyme itself is not predictive of its activity, so
attempts to engineer a more thermostable protein may not improve activity *in planta*. Rather, it is
clear that RCA activity is regulated in response to heat by small molecules, by post-translational
modifications and by the redox state of the chloroplast. The magnitude and direction of these effects
is dependent on the duration of the heat treatment of the plant and the corresponding changes made to
the proteoforms present.
Assimilation, isoform expression, thermostability and sensitivity to TrxF, Mg^{2+} and ADP vary
among C_4 grass species and even among cultivars of maize. Thus, we caution that characterizations
performed on the model species setaria cannot reliably be assumed to be true of other species. Nor
can the effects of the biochemical environment be safely extrapolated even among cultivars of the
same species. Future work must therefore be specific to the species and cultivar.

We conclude that complex interactions occur between changes in proteoform abundance and
changes in the thylakoid environment to create a more permissive environment for RCA function,
enabling recovery of Vcmax and of assimilation. We caution that reductionist approaches to studying
RCA, such as transgenic expression or characterization in non-physiological biochemical settings,
will inevitably overlook these complex interactions. Rather, future work should seek mechanistic
explanations for the differences in RCA activity caused by heat, post-translational modifications,
redox environment and small molecules. A more mechanistic understanding will likely generate
actionable predictions of mutations that could improve assimilation in these plants during heat
exposure.
563

12   References

Anderson, Cheyenne M., Erin M. Mattoon, Ningning Zhang, Eric Becker, William McHargue, Jiani
Yang, Dhruv Patel, et al. 2021. "High Light and Temperature Reduce Photosynthetic Efficiency
through Different Mechanisms in the C4 Model Setaria Viridis." *Communications Biology* 4 (1).
https://doi.org/10.1038/s42003-021-02576-2.

Barta, Csengele, A. Elizabete Carmo-Silva, and Michael E. Salvucci. 2011a. "Purification of Rubisco
Activase from Leaves or after Expression in Escherichia Coli." *Methods in Molecular Biology
(Clifton, N.J.)* 684: 363–74. https://doi.org/10.1007/978-1-60761-925-3_28.

———. 2011b. "Rubisco Activase Activity Assays." In *Photosynthesis Research Protocols*,
684:375–82. https://doi.org/10.1007/978-1-60761-925-3.

Barta, Csengele, Alison M. Dunkle, Rebekka M. Wachter, and Michael E. Salvucci. 2010.
"Structural Changes Associated with the Acute Thermal Instability of Rubisco Activase."
Archives of Biochemistry and Biophysics 499 (1–2): 17–25.
https://doi.org/10.1016/j.abb.2010.04.022.

Bernacchi, Carl J., Archie R. Portis, Hiromi Nakano, Susanne Von Caemmerer, and Stephen P. Long.
2002. "Temperature Response of Mesophyll Conductance. Implications for the Determination of
Rubisco Enzyme Kinetics and for Limitations to Photosynthesis in Vivo." *Plant Physiology* 130
(4): 1992–98. https://doi.org/10.1104/pp.008250.

Bhaduri, Satarupa, Sandeep K. Singh, Whitaker Cohn, S. Saif Hasan, Julian P. Whitelegge, and
William A. Cramer. 2020. "A Novel Chloroplast Super-Complex Consisting of the ATP
Synthase and Photosystem I Reaction Center." *PLoS ONE*.
https://doi.org/10.1371/journal.pone.0237569.

Busch, Florian A., and Rowan F. Sage. 2017. "The Sensitivity of Photosynthesis to O2 and CO2
Concentration Identifies Strong Rubisco Control above the Thermal Optimum." *New
Phytologist* 213 (3): 1036–51. https://doi.org/10.1111/nph.14258.

Caemmerer, Susanne Von, and Robert T. Furbank. 1999. "Modeling C4 Photosynthesis." In *C4 Plant
Biology*, 173–211.

Carmo-Silva, A. E., and M. E. Salvucci. 2013. "The Regulatory Properties of Rubisco Activase
Differ among Species and Affect Photosynthetic Induction during Light Transitions." *Plant
Physiology* 161 (4): 1645–55. https://doi.org/10.1104/pp.112.213348.

Carmo-Silva, A. Elizabete, Michael A. Gore, Pedro Andrade-Sanchez, Andrew N. French, Doug J.
Hunsaker, and Michael E. Salvucci. 2012. "Decreased CO 2 Availability and Inactivation of
Rubisco Limit Photosynthesis in Cotton Plants under Heat and Drought Stress in the Field."
Environmental and Experimental Botany 83: 1–11.
https://doi.org/10.1016/j.envexpbot.2012.04.001.

Chifflet, S., A. Torriglia, R. Chiesa, and S. Tolosa. 1988. "A Method for the Determination of
Inorganic Phosphate in the Presence of Labile Organic Phosphate and High Concentrations of
Protein: Application to Lens ATPases." *Analytical Biochemistry* 168 (1): 1–4.
https://doi.org/10.1016/0003-2697(88)90002-4.

Crafts-Brandner, S J, and M E Salvucci. 2000. "Rubisco Activase Constrains the Photosynthetic
Potential of Leaves at High Temperature and CO2." *Proceedings of the National Academy of
Sciences of the United States of America* 97 (24): 13430–35.
https://doi.org/10.1073/pnas.230451497.

Crafts-Brandner, Steven J., and Michael E. Salvucci. 2002. "Sensitivity of Photosynthesis Heat Stress
in a C4 Plant, Maize, to Heat Stress." *Plant Physiology* 129 (4): 1773–80.
https://doi.org/10.1104/pp.002170.or.

Crafts-Brandner, Steven J, Frank J Van De Loo, Michael E. Salvucci, Frank J van de Loo, and
Michael E. Salvucci. 1997. "The Two Forms of Ribulose-1,5-Bisphosphate
Carboxylase/Oxygenase Activase Differ in Sensitivity to Elevated Temperature." *Plant
Physiology* 114: 439–44.

Davidi, Dan, Melina Shamshoum, Zhijun Guo, Yinon M Bar-On, Noam Prywes, Aia Oz, Jagoda
Jablonska, et al. 2020. "Highly Active Rubiscos Discovered by Systematic Interrogation of
Natural Sequence Diversity." *The EMBO Journal* 39 (18): 1–11.
https://doi.org/10.15252/embj.2019104081.

Degen, Gustaf E., Dawn Worrall, and Elizabete Carmo-Silva. 2020. "An Isoleucine Residue Acts as
a Thermal and Regulatory Switch in Wheat Rubisco Activase." *Plant Journal* 103 (2): 742–51.
https://doi.org/10.1111/tpj.14766.

Hazra, Suratna, J. Nathan Henderson, Kevin Liles, Matthew T. Hilton, and Rebekka M. Wachter.
2015. "Regulation of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase (Rubisco) Activase:
Product Inhibition, Cooperativity, and Magnesium Activation." *Journal of Biological Chemistry*
290 (40): 24222–36. https://doi.org/10.1074/jbc.M115.651745.

Henderson, J. Nathan, Suratna Hazra, Alison M. Dunkle, Michael E. Salvucci, and Rebekka M.
Wachter. 2013. "Biophysical Characterization of Higher Plant Rubisco Activase." *Biochimica et
Biophysica Acta - Proteins and Proteomics* 1834 (1): 87–97.
https://doi.org/10.1016/j.bbapap.2012.09.006.

Hikosaka, Kouki, Kazumasa Ishikawa, Almaz Borjigidai, Onno Muller, and Yusuke Onoda. 2006.
"Temperature Acclimation of Photosynthesis: Mechanisms Involved in the Changes in
Temperature Dependence of Photosynthetic Rate." *Journal of Experimental Botany* 57 (2 SPEC.
ISS.): 291–302. https://doi.org/10.1093/jxb/erj049.

Holmgren, A. 1979. "Thioredoxin Catalyzes the Reduction of Insulin Disulfides by Dithiothreitol

and Dihydrolipoamide." *Journal of Biological Chemistry* 254 (19): 9627–32.
https://doi.org/10.1016/s0021-9258(19)83562-7.

Inc, AAT BioQuest. 2022. "Quest Graph^{TM} Four Parameter Logistic (4PL) Curve Calculator."
https://www.aatbio.com/tools/four-parameter-logistic-4pl-curve-regression-online-calculator.

Ishijima, Sumio, Atsuhito Uchibori, Hiroki Takagi, Ryoji Maki, and Masatake Ohnishi. 2003.
"Light-Induced Increase in Free Mg2+ Concentration in Spinach Chloroplasts: Measurement of
Free Mg2+ by Using a Fluorescent Probe and Necessity of Stromal Alkalinization." *Archives of*
Biochemistry and Biophysics 412 (1): 126–32. https://doi.org/10.1016/S0003-9861(03)00038-9.

Kale, Ravindra S, Larry Sallans, Laurie K Frankel, and Terry M Bricker. 2022. "High-Resolution
Tandem Mass Spectrometry Indicates Rubisco Activase Is Associated with PS I-LHC I-LHC II
Membranes." *BioRxiv* 70803: 1–16.

Kallis, Russell P., Robert G. Ewy, and Archie Jr R. Jr Portis. 2000. "Alteration of the Adenine
Nucleotide Response and Increased Rubisco Activation Activity of Arabidopsis Rubisco
Activase by Site-Directed Mutagenesis." *Plant Physiology* 123 (3): 1077–86.
https://doi.org/10.1104/pp.123.3.1077.

Kanai, Ryuzi, and Gerald E Edwards. 1999. "The Biochemistry of C4 Photosynthesis." In *C4 Plant*
Biology, 49–87. https://doi.org/10.5860/choice.36-6275.

Kattge, Jens, and Wolfgang Knorr. 2007. "Temperature Acclimation in a Biochemical Model of
Photosynthesis: A Reanalysis of Data from 36 Species." *Plant, Cell and Environment* 30 (9):
1176–90. https://doi.org/10.1111/j.1365-3040.2007.01690.x.

Keown, Jeremy R., and Frederick Grant Pearce. 2014. "Characterization of Spinach Ribulose-1,5-
Bisphosphate Carboxylase/Oxygenase Activase Isoforms Reveals Hexameric Assemblies with
Increased Thermal Stability." *Biochemical Journal* 464 (3): 413–23.
https://doi.org/10.1042/BJ20140676.

Kim, Sang Yeol, Christopher M. Harvey, Jonas Giese, Ines Lassowskat, Vijayata Singh, Amanda P.
Cavanagh, Martin H. Spalding, Iris Finkemeier, Donald R. Ort, and Steven C. Huber. 2019. "In
Vivo Evidence for a Regulatory Role of Phosphorylation of Arabidopsis Rubisco Activase at the
Thr78 Site." *Proceedings of the National Academy of Sciences of the United States of America*
116 (37): 18723–31. https://doi.org/10.1073/pnas.1812916116.

Kim, Sang Yeol, Rebecca A. Slattery, and Donald R. Ort. 2021. "A Role for Differential Rubisco
Activase Isoform Expression in C4 Bioenergy Grasses at High Temperature." *GCB Bioenergy*
13 (1): 211–23. https://doi.org/10.1111/gcbb.12768.

Kurek, Itzhak, Thom Kai Chang, Sean M. Bertain, Alfredo Madrigal, Lu Liu, Michael W. Lassner,
and Genhai Zhu. 2007. "Enhanced Thermostability of *Arabidopsis* Rubisco Activase Improves
Photosynthesis and Growth Rates under Moderate Heat Stress." *The Plant Cell* 19 (10): 3230–
41. https://doi.org/10.1105/tpc.107.054171.

Long, S. P., and C. J. Bernacchi. 2003. "Gas Exchange Measurements, What Can They Tell Us about
the Underlying Limitations to Photosynthesis? Procedures and Sources of Error." *Journal of*
Experimental Botany 54 (392): 2393–2401. https://doi.org/10.1093/jxb/erg262.

Nikkanen, Lauri, and Eevi Rintamäki. 2014. "Thioredoxin-Dependent Regulatory Networks in
Chloroplasts under Fluctuating Light Conditions." *Philosophical Transactions of the Royal*
Society B: Biological Sciences. https://doi.org/10.1098/rstb.2013.0224.

Parry, Martin A.J., P. John Andralojc, Joanna C. Scales, Michael E. Salvucci, Elizabete Carmo-Silva,
Hernan Alonso, and Spencer M. Whitney. 2013. "Rubisco Activity and Regulation as Targets
for Crop Improvement." *Journal of Experimental Botany* 64 (3): 717–30.
https://doi.org/10.1093/jxb/err313.

Portis, Archie R. 2003. "Rubisco Activase - Rubisco's Catalytic Chaperone." *Photosynthesis*
Research 75 (1): 11–27. https://doi.org/10.1023/A:1022458108678.

Qu, Yuchen, Oliver Mueller-Cajar, and Wataru Yamori. 2023. "Improving Plant Heat Tolerance
through Modification of Rubisco Activase in C3 Plants to Secure Crop Yield and Food Security
in a Future Warming World." *Journal of Experimental Botany* 74 (2): 591–99.
https://doi.org/10.1093/jxb/erac340.

Qu, Yuchen, Kazuma Sakoda, Hiroshi Fukayama, Eri Kondo, Yuji Suzuki, Amane Makino, Ichiro
Terashima, and Wataru Yamori. 2021. "Overexpression of Both Rubisco and Rubisco Activase
Rescues Rice Photosynthesis and Biomass under Heat Stress." *Plant Cell and Environment* 44
(7): 2308–20. https://doi.org/10.1111/pce.14051.

Sage, R. F. 2013. "Photorespiratory Compensation: A Driver for Biological Diversity." *Plant Biology*
15 (4): 624–38. https://doi.org/10.1111/plb.12024.

Scafaro, Andrew P., Nadine Bautsoens, Bart Den Boer, Jeroen Van Rie, and Alexander Gallé. 2019.
"A Conserved Sequence from Heat-Adapted Species Improves Rubisco Activase
Thermostability in Wheat." *Plant Physiology* 181 (1): 43–54.
https://doi.org/10.1104/pp.19.00425.

Scafaro, Andrew P., David De Vleesschauwer, Nadine Bautsoens, Matthew A. Hannah, Bart Den
Boer, Alexander Gallé, Jeroen Van Rie, et al. 2019. "A Single Point Mutation in the C-Terminal
Extension of Wheat Rubisco Activase Dramatically Reduces ADP Inhibition via Enhanced ATP
Binding Affinity." *Journal of Biological Chemistry* 294 (47): 17931–40.
https://doi.org/10.1074/jbc.RA119.010684.

Scafaro, Andrew P., Wataru Yamori, A. Elizabete Carmo-Silva, Michael E. Salvucci, Susanne von
Caemmerer, and Brian J. Atwell. 2012. "Rubisco Activity Is Associated with Photosynthetic
Thermotolerance in a Wild Rice (Oryza Meridionalis)." *Physiologia Plantarum* 146 (1): 99–
109. https://doi.org/10.1111/j.1399-3054.2012.01597.x.

Vargas-Suárez, Martín, Alfredo Ayala-Ochoa, Jessica Lozano-Franco, Itzhel García-Torres, Alberto
Díaz-Quiñonez, Vianney F. Ortíz-Navarrete, and Estela Sánchez-De-Jiménez. 2004. "Rubisco
Activase Chaperone Activity Is Regulated by a Post-Translational Mechanism in Maize
Leaves." *Journal of Experimental Botany* 55 (408): 2533–39.
https://doi.org/10.1093/jxb/erh268.

Wang, Dafu, and Archie R. Portis. 2006. "Increased Sensitivity of Oxidized Large Isoform of
Ribulose-1,5- Bisphosphate Carboxylase/Oxygenase (Rubisco) Activase to ADP Inhibition Is
Due to an Interaction between Its Carboxyl Extension and Nucleotide-Binding Pocket." *Journal*

of Biological Chemistry 281 (35): 25241–49. https://doi.org/10.1074/jbc.M604756200.

Yamori, Wataru, Ko Noguchi, Yuko T. Hanba, and Ichiro Terashima. 2006. "Effects of Internal
Conductance on the Temperature Dependence of the Photosynthetic Rate in Spinach Leaves
from Contrasting Growth Temperatures." *Plant and Cell Physiology* 47 (8): 1069–80.
https://doi.org/10.1093/pcp/pcj077.

Yin, Xinyou, Zhouping Sun, Paul C. Struik, Peter E.L. Van Der Putten, Wim Van Ieperen, and
Jeremy Harbinson. 2011. "Using a Biochemical C4 Photosynthesis Model and Combined Gas
Exchange and Chlorophyll Fluorescence Measurements to Estimate Bundle-Sheath
Conductance of Maize Leaves Differing in Age and Nitrogen Content." *Plant, Cell and
Environment* 34 (12): 2183–99. https://doi.org/10.1111/j.1365-3040.2011.02414.x.

Yin, Z., Z. Zhang, D. Deng, M. Chao, Q. Gao, Y. Wang, Z. Yang, Y. Bian, D. Hao, and C. Xu. 2014.
"Characterization of Rubisco Activase Genes in Maize: An Alpha-Isoform Gene Functions
alongside a Beta-Isoform Gene." *Plant Physiology* 164 (4): 2096–2106.
https://doi.org/10.1104/pp.113.230854.

Zhang, N, and a R Portis. 1999. "Mechanism of Light Regulation of Rubisco: A Specific Role for the
Larger Rubisco Activase Isoform Involving Reductive Activation by Thioredoxin-F."
Proceedings of the National Academy of Sciences of the United States of America 96 (16):
9438–43. https://doi.org/10.1073/pnas.96.16.9438.

Zhang, Ning, Peter Schürmann, and Archie R. Portis. 2001. "Characterization of the Regulatory
Function of the 46-KDa Isoform of Rubisco Activase from Arabidopsis." *Photosynthesis
Research* 68 (1): 29–37. https://doi.org/10.1023/A:1011845506196.

Zhou, Haoran, Erol Akçay, and Brent R. Helliker. 2018. "Estimating C4 Photosynthesis Parameters
by Fitting Intensive A/C i Curves." *Photosynthesis Research* 141 (2): 181–94.
https://doi.org/10.1007/s11120-019-00619-8.