Impaired intracellular Ca^{2+} signaling contributes to age-related cerebral small vessel disease in *Col4a1* mutant mice

Evan Yamasaki[1], Pratish Thakore[1], Sher Ali[1], Alfredo Sanchez Solano[1], Xiaowei Wang[2], Xiao Gao[3,4], Cassandre Labelle-Dumais[2], Myriam M. Chaumeil[3,4], Douglas B. Gould[2,5], and Scott Earley[1*]

[1]Department of Pharmacology, Center for Molecular and Cellular Signaling in the Cardiovascular System, University of Nevada, Reno School of Medicine, Reno, NV 89557-0318, USA

[2]Department of Ophthalmology, UCSF School of Medicine, San Francisco, CA 94158, USA

[3]Department of Physical Therapy and Rehabilitation Science, UCSF School of Medicine, San Francisco, CA 94143, USA

[4]Department of Radiology and Biomedical Imaging, UCSF School of Medicine, San Francisco, CA 94143, USA

[5]Department of Anatomy, Institute for Human Genetics, Cardiovascular Research Institute, Bakar Aging Research Institute, UCSF School of Medicine, San Francisco, CA 94158, USA

Keywords: *COL4A1*, collagen, cerebral small vessel diseases, basement membrane, myogenic tone, TRPM4, IP$_3$R, Calcium, Ca^{2+}, ER Stress, SR Stress, ion channels.

[*]Corresponding author

Address Correspondence To: Scott Earley, Ph.D.
University of Nevada, Reno School of Medicine
Manville Health Sciences Building, Room 8
MS-0318
Reno, NV 89557-0318, USA
Phone: (775) 784-4117
Fax: (775) 784-1620
Email: searley@med.unr.edu

Abstract

Humans and mice with mutations in *COL4A1* and *COL4A2* manifest hallmarks of cerebral small vessel disease (cSVD), but the pathogenic mechanisms are largely unknown. Mice with a missense mutation in *Col4a1* at amino acid 1344 (*Col4a1$^{+/G1344D}$*) exhibited age-dependent intracerebral hemorrhage (ICH) and brain lesions. Here we report that this pathology was associated with the loss of myogenic vasoconstriction, an intrinsic vascular response essential for the autoregulation of cerebral blood flow. Electrophysiological analyses showed that the loss of myogenic constriction resulted from blunted pressure-induced smooth muscle cell (SMC) membrane depolarization. Further, we found that dysregulation of membrane potential was associated with impaired Ca^{2+}-dependent activation of large-conductance Ca^{2+}-activated K^+ (BK) and transient receptor potential melastatin 4 (TRPM4) cation channels linked to disruptions in sarcoplasmic reticulum (SR) Ca^{2+} signaling. Treating *Col4a1$^{+/G1344D}$* mice with 4-phenylbutyrate, a compound that promotes the trafficking of misfolded proteins and alleviates SR stress, restored SR Ca^{2+} signaling, BK and TRPM4 channel activity, prevented loss of myogenic tone, and reduced ICH. We conclude that alterations in SR Ca^{2+} handling that impair membrane potential regulating ion channel activity result in dysregulation of SMC membrane potential and loss of myogenic tone contributing to age-related cSVD in *Col4a1$^{+/G1344D}$* mice.

56 **Graphical Abstract**

Introduction

Cerebral small vessel diseases (cSVDs) are a group of related pathologies that damage arteries, arterioles, venules, and capillaries in the brain. cSVDs constitute a significant cause of stroke and, after Alzheimer's disease, are the second leading cause of adult cognitive impairment and dementia [1,2]. The underlying pathogenic mechanisms remain largely unknown, and no specific treatment options exist. Sporadic and familial forms of cSVD share clinical and radiological features, including white matter hyperintensities, dilated perivascular spaces, lacunar infarcts, microbleeds, and intracerebral hemorrhages (ICHs) [3-6]. Individuals with Gould syndrome, a rare multisystem disorder caused by autosomal dominant mutations in collagen IV alpha 1 (*COL4A1*) and alpha 2 (*COL4A2*), manifest these hallmarks [7-9]. The majority of pathogenic *COL4A1* and *COL4A2* mutations substitute critical glycine (G) residues within triple helical domains [10]. These mutations have several deleterious effects, including impaired secretion of mature COL4A1 and COL4A2 heterotrimers [$\alpha 1\alpha 1\alpha 2$(IV)] that form basement membranes and intracellular retention of mutant collagen that can lead to endoplasmic reticulum (ER) stress [11-13]. However, it is unclear how these defects influence the pathology of cSVD. *Col4a1* and *Col4a2* mutations demonstrate position-dependent effects contributing to phenotypic heterogeneity [13]. For the current study, we investigated the pathogenesis of *Col4a1*-associated cSVD using mice with a missense mutation that changes the G residue at amino acid position 1344 to aspartate (D). These animals, designated *Col4a1$^{+/G1344D}$*, model the most common type of mutation in humans and develop severe cSVD pathology compared to other

mutant mice carrying glycine missense mutations in a murine allelic series of *Col4a1*

and *Col4a2* mutations [10,13].

Distinctive attributes of the cerebral vasculature serve the unique requirements of

the brain. For example, cerebral arteries and arterioles exhibit graded constriction in

response to changes in intravascular pressure. This vital autoregulatory process, known

as the vascular myogenic response, protects downstream capillary beds from potentially

harmful increases in perfusion pressure when cardiac output is transiently elevated

during exercise or other stimuli [14]. The myogenic response is intrinsic to vascular

smooth muscle cells (SMCs) and requires the coordinated activity of several ion

channels. Increases in intraluminal pressure activate transient receptor potential

melastatin 4 (TRPM4) channels, allowing an influx of Na^+ ions that depolarizes the

plasma membrane to increase voltage-dependent Ca^{2+} inflow through Cav1.2 Ca^{2+}

channels and engage the contractile apparatus [15,16]. Knockdown studies of TRPM4

show that *in vivo* suppression of TRPM4 decreases cerebral artery myogenic

constriction and impairs cerebral blood flow autoregulation [17]. Pressure-induced

membrane depolarization is balanced by hyperpolarizing currents conducted by several

K^+-permeable ion channels, including large-conductance Ca^{2+}-activated K^+ (BK)

channels, voltage-dependent K^+ (K_V) channels, and others [18-20]. The interplay of

depolarizing and hyperpolarizing currents controls the resting membrane potential of

SMCs and maintains cerebral arteries in a state of partial contraction. BK and TRPM4

channels require high levels of intracellular Ca^{2+} for activation and, under native

conditions, are stimulated by Ca^{2+} released from the sarcoplasmic reticulum (SR)

through ryanodine receptors (RyRs) and inositol trisphosphate receptors (IP_3Rs),

respectively [18,21]. Impairment of the myogenic response has been implicated in many

types of cerebrovascular disease, including familial cSVDs associated with cerebral

autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy

(CADASIL) caused by mutations in *NOTCH3* [22]. However, the impact of the

Col4a1[G1344D] mutation on the myogenic response is not known.

Here, we report that *Col4a1*[+/G1344D] mice exhibit age-dependent loss of myogenic

tone that is associated with spontaneous intracerebral hemorrhage (ICH) and brain

lesions detected by susceptibility-weighted magnetic resonance imaging (SWI). The

deficit in myogenic tone resulted from diminished stretch-induced activation of TRPM4

channels and subsequent impairment of SMC membrane depolarization. We also

found that subcellular and global Ca^{2+} signals generated by releasing Ca^{2+} from the SR

through RyRs and IP$_3$Rs were disrupted in SMCs from aged *Col4a1*[+/G1344D] mice.

Treating *Col4a1*[+/G1344D] mice with 4-phenylbutyrate (4PBA), a small molecule with

chaperone properties that improves the secretion of misfolded proteins from the ER/SR

[10,13,23-26], prevented SR Ca^{2+} signaling defects, preserved TRPM4 channel activity,

attenuated the loss of myogenic tone, and largely protected mice from ICH. These data

provide evidence that accumulation of misfolded α1α1α2(IV) collagen in the SR of

SMCs from *Col4a1*[+/G1344D] mice causes disruptions in intracellular Ca^{2+} signaling,

stretch-induced activation of TRPM4 channels, pressure-induced SMC membrane

potential depolarization, and myogenic constriction. Our data also suggests these

defects contribute to spontaneous age-related ICH in *Col4a1*[+/G1344D] mice. Thus, the

findings of this study reveal a novel age-dependent facet of cSVD pathogenesis.

Results

Col4a1$^{+/G1344D}$ mice exhibit age-dependent pathology.

All studies utilized *Col4a1*$^{+/G1344D}$ mice and *Col4a1*$^{+/+}$ littermates as controls. *Col4a1*$^{+/G1344D}$ mice carry missense mutation at amino acid 1344 near the C-terminus of the triple helical domain (Figure 1A). Age is the most critical risk factor for cSVD and vascular dementia [27,28]. To investigate age-related aspects of cSVD pathology, mice were studied at 3 and at least 12 months (M) of age, representing the young adult and middle age stages of life, respectively. Male and female mice were used throughout the study, and no sex-specific differences were detected.

Spontaneous ICH is a highly penetrant and consequential manifestation of *Col4a1* mutations that increases in severity with age [10,29,30]. ICH severity was measured in brain sections from *Col4a1*$^{+/+}$ and *Col4a1*$^{+/G1344D}$ mice stained with Prussian blue. ICH was not detected in 3 M-old *Col4a1*$^{+/+}$ mice. Low-level, punctate Prussian blue staining was present in brain sections from 3 M-old *Col4a1*$^{+/G1344D}$ mice (Figure 1B and C). 12 M-old *Col4a1*$^{+/+}$ mice did not exhibit ICH, but brain sections from 12 M-old *Col4a1*$^{+/G1344D}$ mice displayed large areas of Prussian blue staining mainly localized to the subcortical regions (Figure 1D and E). The total area of Prussian blue staining in 12 M-old *Col4a1*$^{+/G1344D}$ mice was nearly 20-fold greater than in 3 M-old mutants (1.9% vs. 0.1% of brain area, respectively), indicating that the severity of spontaneous ICH increases dramatically with age.

Neuroimaging is frequently used to diagnose cSVD in human patients [31]. In this study, 12 M-old *Col4a1*$^{+/+}$ and *Col4a1*$^{+/G1344D}$ mice were evaluated using SWI. 12 M-old

Col4a1+/+ mice did not exhibit SWI-detectable brain lesions, but brains from age-

matched *Col4a1+/G1344D* mice had multiple lesions, mainly localized to the thalamus

(Figure 1F-I). The number, total volume, and localization of brain lesions detected with

SWI are consistent with ICH detected with Prussian blue staining.

Diminished pressure-induced constriction of cerebral arteries from 12 M-old

***Col4a1+/G1344D* mice is due to impaired SMC membrane depolarization.**

To better understand the underlying causes of ICH in *Col4a1+/G1344D* mice, we

used *ex vivo* pressure myography to study the myogenic response of cerebral arteries

[32]. Vasoconstriction in response to stepwise increases in intraluminal pressure (from 5

to 120 mmHg) was evaluated by measuring the active steady-state luminal diameter of

cannulated cerebral arteries bathed in standard physiological saline solution at each

pressure. This procedure was then repeated using a Ca^{2+}-free bathing solution to

determine the passive response to pressure, and myogenic tone was calculated as the

difference in active versus passive diameter normalized to the passive diameter. The

myogenic tone of cerebral arteries isolated from 3 M-old *Col4a1+/+* and *Col4a1+/G1344D*

mice did not differ (Figure 2A and B). However, cerebral arteries from 12 M-old

Col4a1+/G1344D mice barely constricted in response to pressure and had significantly

lower levels of myogenic tone than those from age-matched *Col4a1+/+* mice (Figure 2C

and D).

Increases in intraluminal pressure cause SMCs in the vascular wall to depolarize

[33], triggering voltage-dependent Ca^{2+} influx and muscular contraction [33,34]. To

determine if the loss of myogenic tone in 12 M-old $Col4a1^{+/G1344D}$ mice was associated

with impaired pressure-induced depolarization, we recorded the membrane potentials of

SMCs in intact cerebral arteries using intracellular microelectrodes. At low intraluminal

pressure (20 mmHg), the membrane potential of SMCs in cerebral arteries from 3 M-old

$Col4a1^{+/+}$ and $Col4a1^{+/G1344D}$ mice did not differ (Figure 2E and F). Increasing

intraluminal pressure to physiological levels (80 mmHg) depolarized the SMC

membrane potential in arteries from both groups to the same extent (Figure 2E and F).

When vessels from 12 M-old animals were investigated, SMC membrane potential did

not differ between $Col4a1^{+/+}$ and $Col4a1^{+/G1344D}$ when arteries were pressurized to 20

mmHg (Figure 2G and H), but when intraluminal pressure was increased to 80 mmHg,

the depolarization of SMCs in cerebral arteries from $Col4a1^{+/G1344D}$ mice was

significantly blunted compared with controls (Figure 2G and H). These data suggest that

cerebral arteries from 12 M-old $Col4a1^{+/G1344D}$ mice lose myogenic tone because

pressure-induced SMC membrane depolarization is impaired.

Age-dependent declines in BK and TRPM4 channel activity in SMCs from

$Col4a1^{+/G1344D}$ mice.

To determine if changes in ion channel activity account for impaired pressure-

induced SMC depolarization, we used patch-clamp electrophysiology to compare

hyperpolarizing K_V and BK currents and depolarizing TRPM4 currents in native SMCs

from control and $Col4a1^{+/G1344D}$ mice.

Elevated SMC K_V channel current density is responsible for diminished myogenic

tone in cerebral arteries from CADASIL cSVD mice [22]. We measured K_V currents in

freshly isolated SMCs to determine if a similar mechanism accounts for the loss of

myogenic tone in cerebral arteries from 12 M-old *Col4a1+/G1344D* mice. K_V currents were

recorded using the amphotericin B perforated patch-clamp configuration to maintain the

integrity of intracellular signaling cascades. Whole-cell currents were evoked by

applying voltage steps from -60 to +60 mV from a holding potential of -80 mV in the

presence of selective BK channel blocker paxilline (1 µM). The voltage step protocol

was repeated in the presence of K_V channel blocker 4-aminopyridine (5 mM), and K_V

current amplitude was determined at each potential by subtraction. We found that K_V

current amplitudes did not differ between SMCs from 12 M-old *Col4a1+/+* and

Col4a1+/G1344D mice at all applied command potentials (Supplemental Figure 1A and B).

The activity of BK channels in cerebral artery SMCs under native conditions is

driven by the transient release of Ca^{2+} ions through RyRs on the SR into restricted

subcellular microdomains proximal to the plasma membrane [35]. These signaling events,

known as Ca^{2+} sparks, activate clusters of BK channels on the plasma membrane to

generate large-amplitude spontaneous transient outward currents (STOCs) [18,19]. We

utilized the amphotericin B perforated patch-clamp configuration to measure STOCs in

cerebral artery SMCs from 3 M-old *Col4a1+/+* and *Col4a1+/G1344D* mice over a range of

command potentials (-60 to 0 mV) and saw no significant difference in STOC frequency

or amplitude (Figure 3A-C). However, the frequency and amplitude of STOCs were

significantly lower in cerebral artery SMCs from 12 M-old *Col4a1+/G1344D* mice compared

with age-matched controls (Figure 3D-F). To determine if decreases in STOCs result

from changes in gene expression, we utilized ddPCR to measure transcript levels of

Kcnma1 (BK subunit α), *Kcnmb1 (*BK subunit β1), and *Ryr2* (ryanodine receptor 2). No

differences in mRNA expression levels were detected (Supplemental Figure 2).

Reduced BK channel activity depolarizes the SMC membrane potential and

increases contractility [36]. Thus, diminished BK channel activity in SMCs from 12 M-old

Col4a1+/G1344D mice cannot account for impaired pressure-induced membrane

depolarization.

TRPM4 is a Ca^{2+}-activated, nonselective monovalent cation channel required for

pressure-induced SMC depolarization and the development of myogenic tone in

cerebral arteries [15,16,37,38]. At the membrane potentials of SMCs in the vascular wall

under physiological conditions (-60 to -30 mV), TRPM4 channels conduct inward Na^+

currents that depolarize the plasma membrane in response to increases in intraluminal

pressure [15,39]. Patch-clamp electrophysiology was used to determine if TRPM4 activity

is diminished in *Col4a1+/G1344D* mice. We assessed channel function and availability in

freshly isolated cerebral artery SMCs using the conventional whole-cell patch-clamp

configuration with a high [Ca^{2+}] intracellular solution to directly activate TRPM4.

Currents were recorded as voltage ramps (-100 to +100 mV) were applied. Voltage

ramps were repeated in the presence of TRPM4 blocker 9-phenanthrol (30 µM) [15], and

TRPM4 currents were isolated by subtraction. Prior studies using ion substitution

protocols have shown that Ca^{2+}-activated, 9-phenanthrol currents recorded using these

methods are carried by Na^+ ions, providing substantial evidence that these are *bona*

fide TRPM4 currents [16,37,40]. We found that the amplitudes of whole-cell TRPM4 currents

recorded from cerebral artery SMCs did not differ between *Col4a1+/+* and *Col4a1+/G1344D*

mice at 3 and 12 M of age (Figure 3G-J). These data indicate that the collagen mutation

studied here has no direct effects on TRPM4 channel function or the trafficking of

channel protein to the plasma membrane.

TRPM4 channels in SMCs are activated by a signal transduction cascade

initiated by angiotensin II type 1 receptors (AT$_1$Rs) [41-44]. Activation of Gq protein-

coupled AT$_1$Rs stimulates phospholipase C (PLC), leading to the cleavage of

phosphatidylinositol 4,5-bisphosphate (PIP$_2$) into diacylglycerol and inositol triphosphate

(IP$_3$). IP$_3$ stimulates Ca^{2+} release from the SR through IP$_3$Rs to transiently activate

TRPM4 channels [21,38,41]. AT$_1$Rs are directly activated by the stretch of the plasma

membrane (independent of their ligand) and are a critical mechanosensor in SMCs [41-44].

To measure the activity of TRPM4 channels in response to mechanical force, we patch-

clamped SMCs in the amphotericin B perforated patch configuration and applied

negative pressure (-20 mmHg) through the patch pipet to stretch the plasma membrane.

Stretch-induced transient inward cation currents (TICCs) evoked in this manner are

blocked by 9-phenanthrol and abolished by the knockdown of TRPM4 expression [15,21].

Stretch-activated TICC activity in SMCs from 3 M-old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$*

(Figure 3K and L) did not differ. In contrast, stretch-induced TICC activity in SMCs from

12 M-old *Col4a1$^{+/G1344D}$* mice was significantly lower than that of age-matched *Col4a1$^{+/+}$*

mice (Figure 3M and N). To determine if altered gene expression underlies the

reduction in TICC activity, we utilized ddPCR to measure transcript levels of *Trpm4*

(TRPM4), *Itpr1* (IP$_3$ receptor 1), and *Itpr2* (IP$_3$ receptor 2) in cerebral arteries from 12 M-

old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice. No differences in mRNA copy numbers were

observed (Supplemental Figure 2). These data suggest that blunted pressure-induced

SMC depolarization and impaired myogenic tone result from diminished stretch-induced

activation of TRPM4 channels.

Defective intracellular Ca^{2+} signaling in cerebral artery SMCs from 12 M-old

***Col4a1$^{+/G1344D}$* mice.**

Ca^{2+} released from the SR drives BK and TRPM4 channel activity in cerebral

artery SMCs [18,21]. Therefore, we investigated the possibility that these vital Ca^{2+}

signaling pathways are disrupted in SMCs from 12 M-old *Col4a1$^{+/G1344D}$* mice. Freshly

isolated SMCs were loaded with the Ca^{2+}-sensitive fluorophore Fluo-4 AM and imaged

using high-speed, high-resolution spinning disk confocal microscopy. Spontaneous Ca^{2+}

sparks were present in SMCs isolated from 3 M-old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice.

The frequency, amplitude, duration, rise time, and decay rate did not differ between

groups (Figure 4A-C). Spontaneous Ca^{2+} sparks were also present in SMCs isolated

from both groups of 12 M-old mice. No significant differences in Ca^{2+} spark amplitude,

duration, rise time, and decay rate were observed, but notably, Ca^{2+} spark frequency

was significantly lower in SMCs from 12 M-old *Col4a1$^{+/G1344D}$* mice (Figure 4D-F). Thus,

diminished Ca^{2+} spark frequency likely accounts for decreased STOC frequency in

SMCs from 12 M-old *Col4a1* mutant mice.

Fluo-4 AM-loaded SMCs were challenged with the RyR agonist caffeine (10 mM)

to investigate how RyR function was affected by the *Col4a1^{G1344D}* mutation [45]. Ca^{2+}

signals arising from intracellular release were isolated from Ca^{2+} influx events by acutely

removing Ca^{2+} from the extracellular solution immediately before recording. Caffeine-

evoked global cytosolic [Ca^{2+}] increases did not differ between SMCs from 3 M-old

Col4a1$^{+/+}$ and *Col4a1$^{+/G1344D}$* mice (Figure 4G and H). In contrast, caffeine-induced

global Ca^{2+} signals were smaller in amplitude in SMCs from 12 M-old *Col4a1$^{+/G1344D}$*

mice than in controls (Figure 4I and J). These data indicate that the release of SR Ca^{2+}

from RyRs is impaired in the SMCs of *Col4a1$^{+/G1344D}$* mice in an age-dependent manner.

Fluo-4 AM loaded-SMCs were also treated with the G_q protein-coupled

thromboxane A2 receptor agonist U46619 (100 nM) to investigate IP$_3$R-mediated Ca^{2+}

signaling. U46619-induced Ca^{2+} signals did not differ between cerebral artery SMCs

from 3 M-old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice (Figure 4K and L). In contrast, Ca^{2+}

signals induced by U46619 were smaller in amplitude in SMCs from 12 M-old

Col4a1$^{+/G1344D}$ mice compared to controls (Figure 4M and N). These data suggest that

the release of SR Ca^{2+} from IP$_3$Rs is also impaired in the SMCs of *Col4a1$^{+/G1344D}$* mice

in an age-dependent manner.

RyRs and IP$_3$Rs share a common SR Ca^{2+} pool [46]. Impairment of Ca^{2+} release

through both receptors suggests that the SR [Ca^{2+}] of SMCs from 12 M-old

Col4a1$^{+/G1344D}$ mice is lower than that of corresponding controls. Further, diminished SR

Ca^{2+} levels likely account for decreased Ca^{2+} spark frequency and diminished STOC

and TICC activity.

4PBA prevents age-related cerebral artery defects in *Col4a1$^{+/G1344D}$* mice.

Col4a1 mutations impair collagen α1α1α2(IV) secretion *in vitro* and *in vivo* and

can be alleviated by treatment with 4PBA [10,13,26]. The chemical chaperone 4PBA

facilitates the trafficking of misfolded mutant proteins and reduces ER/SR stress [47]. To

investigate a potential link between impaired collagen secretion and the cerebral

vascular pathology of *Col4a1*[+/G1344D] mice, 4PBA (50 mM) was added to the drinking

water of mutant and control animals from birth (postnatal day 0).

We found that 4PBA treatment prevented the diminishment of caffeine-evoked

(Figure 5A and B) and U46619-induced (Figure 5C and D) Ca^{2+} signals in cerebral

artery SMCs from 12 M-old *Col4a1*[+/G1344D] mice. Ca^{2+} spark frequency was also

normalized to control levels in 4PBA-treated *Col4a1*[+/G1344D] mice (Supplemental Figure

3A-C). We also found that 4PBA treatment averted the loss of STOC (Supplemental

Figure 3D-E) and TICC (Figure 5E and F) activity. These data demonstrate that age-

related defects in intracellular Ca^{2+} signaling that control ion channel activity and

membrane potential were prevented by treating *Col4a1*[+/G1344D] mice with 4PBA. 4PBA

treatment also prevented loss of myogenic tone (Figure 5G and H) and significantly

diminished spontaneous ICH in 12 M-old *Col4a1*[+/G1344D] mice (Figure 5I and J). These

data suggest that 4PBA protects against age-dependent defects in myogenic tone and

ICH in *Col4a1*[+/G1344D] mice.

Discussion

Despite the enormous impact of the disease, the molecular pathogenesis of

cSVD is essentially unknown. Here we utilized the *Col4a1*[+/G1344D] mouse model of Gould

syndrome to investigate how this monogenic form of cSVD damages small arteries in

the brain during aging. Our findings show that these mice exhibit pathological changes

with age, including the loss of myogenic tone, high levels of spontaneous ICH, and brain

lesions. Electrophysiology experiments revealed that the loss of myogenic tone is

associated with diminished pressure-induced SMC depolarization and decreased

activities of BK and TRPM4 channels. In addition, we found that fundamental RyR- and

IP$_3$R-dependent Ca^{2+} signaling pathways are disrupted in SMCs from 12 M-old

Col4a1$^{+/G1344D}$ mice, likely due to a reduction in SR Ca^{2+} levels. Treating Col4a1$^{+/G1344D}$

mice with the chemical chaperone, 4PBA negated Ca^{2+} signaling disruptions, increased

BK and TRPM4 channel activity, prevented loss of myogenic tone, and reduced

spontaneous ICH. We conclude that the age-dependent cSVD pathogenesis of

Col4a1$^{+/G1344D}$ mice results from defects in SMC SR function that lead to reduced Ca^{2+}

store load and disrupted intracellular Ca^{2+} signaling. Defects in subcellular Ca^{2+} signals

diminish BK and TRPM4 channel activity, disrupting membrane potential regulation and

impeding pressure-induced constriction of cerebral arteries. These findings reveal a

novel mechanism that contributes to age-dependent cSVD.

Myogenic constriction protects delicate capillaries in the brain from pressure

overload during increases in cardiac output and perfusion associated with exercise,

stress, and other stimuli [14]. This vital response is deficient in several monogenetic forms

of cSVD [22,30,48]. However, the defects responsible for the loss of myogenic tone differ

between disease models. For example, Dabertrand et al. linked the loss of myogenic

tone in the CADASIL model with blunted pressure-induced SMC membrane potential

depolarization and connected the defect in membrane potential regulation with

increased surface expression of K$_V$ channels and elevated hyperpolarizing K$^+$ current

density [22]. Notably, depletion of the minor membrane phospholipid PIP$_2$, leading to the

loss of TRPM4 channel activity, is the fundamental defect responsible for the loss of

myogenic constriction in a distinct mouse model of Gould syndrome caused by a

Col4a1^{G394V} mutation [30]. The current study uncovered a distinctive flaw in the SMCs of

Col4a1$^{+/G1344D}$ cSVD mice – impaired Ca^{2+} signaling. Our data show that the release of

Ca^{2+} from the SR through RyRs and IP$_3$Rs became faulty in *Col4a1$^{+/G1344D}$* mice by 12

M of age. Impaired SR Ca^{2+} release disrupted the activity of BK and TRPM4 channels.

BK and TRPM4 channels play opposing roles in regulating SMC membrane potential

and contractility. TRPM4 channels depolarize the membrane and promote contractility,

whereas BK channels provide negative feedback to limit the magnitude of this response

[49]. When pressure-induced membrane depolarization is impaired, the BK pathway is

disengaged and has minimal influence on membrane potential [50].

Diminished levels of Ca^{2+} in the SR likely cause Ca^{2+} signaling and ion channel

defects in SMCs from 12 M-old *Col4a1$^{+/G1344D}$*. This concept is supported by prior

studies showing that a ~16% decrease in SR [Ca^{2+}] diminished STOC frequency by

70% and that a >20% decrease in SR [Ca^{2+}] completely abolished IP$_3$R-mediated Ca^{2+}

release in native SMCs [51,52]. In many types of cells, store-operated Ca^{2+} entry (SOCE)

is protective against decreases in ER/SR [Ca^{2+}] [53]. However, SOCE is virtually

undetectable in native, contractile vascular SMCs, promoting their susceptibility to

pathogenic SR Ca^{2+} depletion [50,54,55]. We conclude that SR Ca^{2+} store depletion and the

resultant dysregulation of BK, TRPM4, and SMC membrane potential is responsible for

the age-related loss of myogenic tone in *Col4a1$^{+/G1344D}$* cSVD mice.

Steady-state free Ca^{2+} levels in the ER/SR are maintained by the release of Ca^{2+}

into the cytosol through RyRs and IP$_3$Rs, mobile and stationary Ca^{2+}-binding proteins in

the SR lumen, and transport of Ca^{2+} from the cytosol into the SR by sarco/endoplasmic

reticulum ATPase (SERCA) enzymes [46]. Disruption of one or more of these elements

could lead to SR Ca^{2+} defects in *Col4a1^{+/G1344D}* mice. Interestingly, *Col4a1^{+/G1344D}* mice

display impaired secretion and increased intracellular accumulation of collagen

α1α1α2(IV), presumably due to difficulties associated with the trafficking of misfolding of

mutant proteins out of the ER/SR [13]. The retention of misfolded proteins causes a

pathological condition known as ER/SR stress [56] that is associated with the depletion of

Ca^{2+} stores [57]. For example, Yamato *et al.* [58] demonstrated that tunicamycin-induced

ER stress promotes RyR-mediated Ca^{2+} leak, potentially due to increased reactive

oxygen species (ROS) production. Our data showed that treatment with a molecular

chaperone prevented SR Ca^{2+} handling defects and loss of myogenic tone and

significantly attenuated spontaneous ICH. These data provide evidence that ER/SR

stress contributes to SMC Ca^{2+} signaling defects and cSVD pathology of *Col4a1^{+/G1344D}*

mice. The age dependency of the disease process may reflect the normal functional

decline in SR protein homeostasis that is exacerbated by mutant collagen in the SR.

We conclude that age-dependent disruption of Ca^{2+} signaling in SMCs due to SR

stress and diminished SR Ca^{2+} store load contributes to cSVD pathology in

Col4a1^{+/G1344D} mice. Future investigations into links between SMC SR stress and Ca^{2+}

handling defects in other genetic and sporadic forms of cSVD may provide insight into

novel pathogenic mechanisms.

Materials and Methods

Chemical and reagents: All chemicals and other reagents were obtained from Sigma-Aldrich, Inc. (St. Louis, MO, USA) unless otherwise specified.

Animals: Young adult (~3M) and middle-aged (~12M) male and female littermate *Col4a1*$^{+/+}$ and *Col4a1*$^{+/G1344D}$ mice were used in this study. The *Col4a1*^{G1344D} mutation was backcrossed to C57BL/6J mice for over 20 generations [59]. Animals were maintained in individually ventilated cages (≤5 mice/cage) with *ad libitum* access to food and water in a room with controlled 12-hour light and dark cycles. All animal care procedures and experimental protocols involving animals complied with the National Institutes of Health (NIH) *Guide for the Care and Use of Laboratory Animals* and were approved by the Institutional Animal Care and Use Committees at the University of Nevada, Reno, and University of California, San Francisco.

4PBA treatment: Mice were provided with 4-phenylbutyrate (4PBA; 50 mM; Scandinavian Formulas Inc., Sellersville, PA, USA) from birth in drinking water that was refreshed weekly as described previously [26].

Histological analysis: Isoflurane anesthetized mice were transcardially perfused with ice-cold phosphate-buffered saline (PBS) followed by ice-cold 4% paraformaldehyde (Fisher, Waltham, MA, USA) in PBS. Brains were postfixed with 4% paraformaldehyde for 24 hours at 4°C, cryoprotected in 30% sucrose in PBS, and embedded in optimal cutting temperature compound (Fisher). Coronal cryosections (35 μm) regularly spaced (500 μm) along the rostrocaudal axis were stained with a Prussian blue and nuclear fast red stain kit (Abcam, Cambridge, UK) following the manufacturer's protocol. Images

were acquired with a BZ-X700 microscope using BZ-X Viewer 1.3.0.5 software and

stitched with BZ-X Analyzer 1.3.0.3 software (Keyence, Osaka, Japan). The percentage

of brain area with Prussian blue staining for each section was calculated with ImageJ

software (v2.3.0/1.53f; NIH, Bethesda, MD, USA). Hemorrhage severity was expressed

as the average percentage of hemosiderin surface area on ~21 sections for each brain.

***In vivo* magnetic resonance imaging:** All *in vivo* magnetic resonance (MR)

experiments were conducted on a 14.1 Tesla vertical MR system (Agilent Technologies,

Palo-Alto, CA) equipped with 100 G/cm gradients and a single tuned millipede ^{1}H proton

coil ($\emptyset_I$ = 40 mm). For each imaging session, mice were anesthetized using isoflurane

(1-1.5% in O_2), positioned in a dedicated cradle maintaining constant anesthesia, and

placed in the MR bore; respiration and temperature were continuously monitored during

all acquisitions to ensure animal well-being and data reproducibility. Susceptibility

weighted imaging (SWI) was performed to acquire high-resolution images of the mouse

brain using the following parameters: gradient-echo scheme, field of view (FOV) = 20x

20 mm^2, matrix = 256 x 256, 16 slices, 0.4 mm slice thickness, 0.1 mm interslice gap,

number of averages = 16, echo time (TE)/repetition time (TR) = 4.60/140ms, flip angle =

10 degrees. T2-weighted (T2W) images were also acquired in a fast-spin-echo scheme

to evaluate anatomical brain structures, using the same FOV geometry as SWI and the

following parameters: number of averages = 8, TE/TR = 21.38/2500ms, flip angle = 90

degrees. The fast-spin-echo based T2W imaging is less sensitive to distortion artifacts

caused by B_0 inhomogeneity than SWI and is thus used for the atlas-based brain

structure registration as described below.

Using in-house software, hypointense lesions corresponding to ICH were

quantified based on manual segmentation of the SWI images. To investigate the

number and distribution of hypointense SWI lesions across the brain, an open-sourced

atlas-based imaging data analysis pipeline (AIDAmri) [60] was customized to register T2W

brain images to the Allen Brain Reference Atlas [61]. Based on the registration result, the

normalized lesion volumes for each brain region were summed up across all animals,

and an intra-group region-wise volumetric analysis was performed.

Isolation of cerebral arteries and SMCs: Mice were euthanized by decapitation under

isoflurane anesthesia. Cerebral arteries were carefully isolated from the brain in Ca^{2+}-

free Mg^{2+} physiological saline solution (Mg^{2+}-PSS) consisting of 140 mM NaCl, 5 mM

KCl, 2 mM $MgCl_2$, 10 mM HEPES, and 10 mM glucose (pH 7.4, adjusted with NaOH),

with 0.5% bovine serum albumin (BSA). Native SMCs for patch-clamp experiments

were obtained by initially digesting isolated arteries in 1 mg/mL papain (Worthington

Biochemical Corporation, Lakewood, NJ, USA), 1 mg/mL dithiothreitol (DTT), and 10

mg/mL BSA in Ca^{2+}-free PSS at 37°C for 12 minutes, followed by a 14-minute

incubation with 1 mg/mL type II collagenase (Worthington Biochemical Corporation). A

single-cell suspension was prepared by washing digested arteries three times with

Mg^{2+}-PSS and triturating with a fire-polished glass pipette. All cells used for this study

were freshly dissociated on the day of experimentation.

Pressure myography: The current best practice guidelines for pressure myography

experiments were followed [32]. Arteries were mounted between two glass cannulas

(outer diameter, 40–50 µm) in a pressure myograph chamber (Living Systems

Instrumentation, St Albans City, VT, USA) and secured by a nylon thread. Intraluminal

pressure was controlled using a servo-controlled peristaltic pump (Living Systems

Instrumentation), and preparations were visualized with an inverted microscope (Accu-

Scope Inc., Commack, NY, USA) coupled to a USB camera (The Imaging Source LLC,

Charlotte, NC, USA). Changes in luminal diameter were assessed using IonWizard

software (version 7.2.7.138; IonOptix LLC, Westwood, MA, USA). Arteries were bathed

in warmed (37°C), oxygenated (21% O_2, 6% CO_2, 73% N_2) PSS (119 mM NaCl, 4.7 mM

KCl, 21 mM $NaHCO_3$, 1.17 mM $MgSO_4$, 1.8 mM $CaCl_2$, 1.18 mM KH_2PO_4, 5 mM

glucose, 0.03 mM EDTA) at an intraluminal pressure of 5 mmHg. Following equilibration

for 15 min, intraluminal pressure was increased to 110 mmHg, and vessels were

stretched to their approximate *in vivo* length, after which pressure was reduced back to

5 mmHg for an additional 15 min. Vessel viability was assessed for each preparation by

evaluating vasoconstriction response to high extracellular [K^+] PSS, made isotonic by

adjusting the [NaCl] (60 mM KCl, 63.7 mM NaCl). Arteries that showed less than 10%

constriction in response to elevated [K^+] were excluded from further investigation.

Myogenic tone was assessed by raising the intraluminal pressure stepwise from

5 mmHg to 120 mmHg in 20 mmHg increments (5 to 20 mmHg for the first step). The

active diameter was obtained by allowing vessels to equilibrate for at least 5 minutes at

each pressure or until a steady-state diameter was reached. Following completion of the

pressure-response study, intraluminal pressure was lowered to 5 mmHg, and arteries

were superfused with Ca^{2+}-free PSS supplemented with EGTA (2 mM) and the voltage-

dependent Ca^{2+} channel blocker diltiazem (10 μM) to inhibit SMC contraction, after

which passive diameter was obtained by repeating the stepwise increase in intraluminal

pressure. Myogenic tone at each pressure step was calculated as myogenic tone (%) =

[1 − (active lumen diameter/passive lumen diameter)] × 100.

Membrane potential: For measurements of SMC membrane potential, cerebral arteries

were isolated, cannulated, and viability confirmed as described above. SMC membrane

potential was recorded from intact pressurized arteries at intraluminal pressures of 20

mmHg and 80 mmHg. SMCs were impaled through the adventitia with glass intracellular

microelectrodes (100-200 MΩ) backfilled with 3M KCl. Membrane potential was

recorded using an Electro 705 amplifier (World Precision Instruments, Sarasota, FL,

USA). Analog output from the amplifier was recorded using Ionwizard software (version

7.5.1.162; IonOptix). Criteria for acceptance of membrane potential recordings were 1)

an abrupt negative deflection of potential as the microelectrode was advanced into a

cell, 2) stable membrane potential for at least 30 seconds, and 3) an abrupt change in

the potential to ~0 mV after the electrode was retracted from the cell.

Electrophysiological recordings: Enzymatically isolated native SMCs were

transferred to a recording chamber (Warner Instruments, Holliston, MA, USA) and

allowed to adhere to glass coverslips for 20 minutes at room temperature. Recording

electrodes (3–5 MΩ) were pulled on a model P-87 micropipette puller (Sutter

Instruments, Novato, CA, USA) and polished using a MF-830 MicroForge (Narishige

Scientific Instruments Laboratories, Long Island, NY, USA). Currents were recorded at

room temperature using an Axopatch 200B amplifier (Molecular Devices, Sunnyvale,

CA, USA) equipped with an Axon CV 203BU headstage and Digidata 1440A digitizer

(Molecular Devices) for all patch-clamp electrophysiology experiments. Clampex and

Clampfit (version 10.2; Molecular Devices) were used for data acquisition and analysis.

The bathing solution composition for perforated-patch recordings of Kv currents,

STOCs, and TICCs was 134 mM NaCl, 6 mM KCl, 1 mM $MgCl_2$, 2 mM $CaCl_2$, 10 mM

HEPES, and 10 mM glucose (pH 7.4, adjusted with NaOH). The patch pipette solution

contained 110 mM K-aspartate, 1 mM $MgCl_2$, 30 mM KCl, 10 mM NaCl, 10 mM HEPES,

and 5 µM EGTA (pH 7.2, adjusted with KOH). Amphotericin B (200 µM) was included in

the pipette solution to gain electrical access. Whole-cell K^+ currents were recorded

using a step protocol (-60 to +60 mV in 10 mV, 250 ms steps) from a holding potential of

-80 mV. The BK channel blocker paxilline (1 µM) was included in the bath solution when

Kv currents were recorded. Whole-cell K^+ currents were recorded in the absence and

presence of K_V channel blocker 4-aminopyridine (5 mM), and the Kv component was

determined by subtraction. Summary current-voltage (I-V) plots were generated using

values obtained from the last 50 ms of each step. STOCs were recorded from SMCs

voltage-clamped over a range of membrane potentials (-60 to 0 mV; 10 mV steps).

TICCs were recorded from SMCs voltage-clamped at -70 mV; membrane stretch was

delivered by applying negative pressure through the recording electrode using a

pressure clamp system (HSPC-1; ALA Scientific Instruments Inc., Farmingdale, NY,

USA). TICC activity was calculated as the sum of the open channel probability (NP_o) of

multiple open states of 1.75 pA [62].

Whole-cell TRPM4 currents were recorded in a bath solution consisting of 156

mM NaCl, 1.5 mM $CaCl_2$, 10 mM glucose, 10 mM HEPES, and 10 mM TEACl (pH 7.4,

adjusted with NaOH). The patch pipette solution contained 156 mM CsCl, 8 mM NaCl,

1 mM $MgCl_2$, 10 mM HEPES (pH 7.4, adjusted with NaOH), and a free Ca^{2+}

concentration of 200 µM, adjusted using the appropriate amount of $CaCl_2$ and EGTA, as

determined with Max-Chelator software (WEBMAXC standard, available at

https://somapp.ucdmc.ucdavis.edu/pharmacology/bers/maxchelator/webmaxc/webmaxc

S.htm. Whole-cell cation currents were evoked by applying 400 ms voltage ramps from -

100 to +100 mV from a holding potential of -60 mV. Voltage ramps were repeated every

2 s for 300 s. The selective TRPM4 inhibitor 9-phenanthrol (30 μM) was applied after

peak TRPM4 current was recorded (~100 s). Whole-cell TRPM4 current amplitude was

expressed as the 9-phenanthrol-sensitive current at +100 mV.

Quantitative droplet digital PCR: Total RNA was extracted from isolated cerebral

arteries by homogenization in TRIzol reagent (Invitrogen, Waltham, MA, USA), followed

by purification using a Direct-zol RNA microprep kit (Zymo Research, Irvine, CA, USA)

with on-column DNAse treatment. RNA concentration was determined using an RNA

6000 Pico Kit run on a Bioanalyzer 2100 running Agilent 2100 Expert Software

(B.02.11; Agilent Technologies, Santa Clara, CA, USA). RNA was converted to cDNA

using iScript cDNA Supermix (Bio-Rad, Hercules, CA, USA). Quantitative droplet digital

PCR (ddPCR) was performed using QX200 ddPCR EvaGreen Supermix (Bio-Rad),

custom-designed primers (Supplementary Table 1), and cDNA templates. Generated

droplet emulsions were amplified using a C100 Touch Thermal Cycler (Bio-Rad), and

the fluorescence intensities of individual droplets were measured using a QX200

Droplet Reader (Bio-Rad) running QuantaSoft (version 1.7.4; Bio-Rad). Analysis was

performed using QuantaSoft Analysis Pro (version 1.0596; Bio-Rad).

Ca^{2+} imaging: Images were acquired using an iXon 897 EMCCD camera (Andor; 16 ×

16 µm pixel size) coupled to a spinning-disk confocal head (CSU-X1; Yokogawa) with a

100× oil-immersion objective (Olympus; NA 1.45) at an acquisition rate of 33 frames per

second (fps). Custom software (SparkAn; https://github.com/vesselman/SparkAn)

provided by Dr. Adrian D. Bonev (University of Vermont) was used to analyze the

properties of Ca^{2+} sparks and whole-cell RyR and IP_3R Ca^{2+} signals.

To image Ca^{2+} sparks, a suspension of freshly isolated SMCs was placed in a glass-

bottom 35 mm dish, and cells were allowed to adhere to the glass coverslip for 20 min

at room temperature. SMCs were then loaded with the Ca^{2+}-sensitive fluorophore, Fluo-

4-AM (1 µM; Invitrogen), in the dark for 20 minutes at room temperature in Mg^{2+}-PSS.

Cells were subsequently washed three times with Ca^{2+}-containing PSS and incubated

at room temperature for 20 min in the dark to allow sufficient time for Fluo-4 de-

esterification. The threshold for Ca^{2+} spark detection was defined as local increases in

fluorescence ≥0.2 $\Delta F/F_0$.

To image RyR and IP_3R Ca^{2+} signals, SMCs were plated and loaded with Fluo-4-AM as

stated above, but immediately before imaging, cells were washed three times and

imaged in Mg^{2+}-PSS to eliminate signals resulting from the influx of Ca^{2+}. Caffeine (10

mM) or U46619 (100 nM; Enzo Biochem, Farmingdale, NY) were applied to the bath.

Statistical analysis: All summary data are presented as means ± SEM. Statistical

analyses were performed, and graphs were constructed using Prism version 9.3.1

(GraphPad Software, San Diego, CA, USA). The value of n refers to the number of cells

for patch-clamp electrophysiology and Ca^{2+} imaging experiments, the number of

arteries for pressure myography and membrane potential experiments, and the number

of animals for histological analyses, magnetic resonance imaging, and ddPCR

experiments. Statistical analyses were performed using unpaired Student's t-tests and

one-way and two-way analysis of variance (ANOVA). A P value < 0.05 was considered

statistically significant for all analyses.

Grant support: This study was supported by grants from the National Institutes of

Health (NHLBI R35HL155008, R01HL091905, R01HL137852, R01HL139585,

R01146054, and R01HL122770, NIGMS P20GM130459 to S.E., and NINDS

RF1NS110044 and R33NS115132 to M.M.C, D.B.G, and S.E.; The Transgenic

Genotyping and Phenotyping Core at the COBRE Center for Molecular and Cellular

Signaling in the Cardiovascular System, University of Nevada, Reno, is maintained by a

grant from NIH/NIGMS (P20GM130459 Sub#5451), as is the High Spatial and

Temporal Resolution Imaging Core at the COBRE Center for Molecular and Cellular

Signaling in the Cardiovascular System, University of Nevada, Reno (P20GM130459

Sub#5452). The UCSF Department of Ophthalmology is supported by P30EY002162

and an unrestricted grant from Research to Prevent Blindness. This study was also

supported by a grant from the American Heart Association (23CDA1054627 to P.T.).

Author contributions: S.E. and D.B.G. initiated and supervised the project. S.E.

designed the experiments. E.Y. performed brain histology, pressure myography, sharp

electrode and patch-clamp electrophysiology, Ca^{2+} imaging, and ddPCR experiments.

X.W. and X.G. performed magnetic resonance imaging experiments. S.A. and A.S.S.

performed patch-clamp electrophysiology experiments. P.T. performed pressure

myography experiments. C.L.D. performed animal colony management. E.Y., S.A.,

A.S.S., P.T., X.G., and S.E. analyzed the data. E.Y. and S.E. drafted the manuscript

and prepared the figures. E.Y., C.L.D, M.M.C., D.B.G., and S.E. revised the manuscript.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: All data needed to evaluate the conclusions in the paper are present in the paper or the Supplementary Materials

Figures

Figure 1. *Col4a1+/G1344D* mice exhibit age-dependent pathologies. (A) Schematic

representation of the *Col4a1* point mutation that leads to *Col4a1G1344D*. **(B)** Intracerebral

hemorrhage (ICH) was assessed using Prussian blue staining. Representative images

of brain sections from 3 M-old *Col4a1*[+/+] and *Col4a1*[+/G1344D] mice stained with Prussian

blue. **(C)** Summary data presented as percentage of brain area with Prussian blue

staining in brain sections (500 μm intervals). N = 3 animals per group. *$P \leq 0.05$,

unpaired t-test. **(D)** Representative images of brain sections from 12 M-old *Col4a1*[+/+]

and *Col4a1*[+/G1344D] mice stained with Prussian blue. **(E)** Summary data presented as

percentage of brain area with Prussian blue staining in brain sections (500 μm

intervals). N = 3 animals per group. *$P \leq 0.05$, unpaired t-test. **(F)** Brain lesions were

evaluated *in vivo* using high-field susceptibility-weighted magnetic resonance imaging

(SWI). Representative SWI images from 12 M-old *Col4a1*[+/+] and *Col4a1*[+/G1344D] mice.

Red arrowheads highlight hypointense pixels that indicate hemorrhagic lesions.

Quantification of **(G)** the number of lesions and **(H)** the total volume of lesions detected

by SWI. N = 4 animals per group. *$P \leq 0.05$, unpaired t-test. **(I)** Region-wise volumetric

distribution of lesions detected by SWI from 12 M-old *Col4a1*[+/G1344D] mice. Scale bars =

1 mm. Inset scale bars = 250 μm.

Figure 2. Diminished pressure-induced constriction of cerebral arteries from 12

M-old *Col4a1+/G1344D* mice is due to impaired SMC membrane depolarization. (A)

Representative recordings of the inner diameter of isolated cerebral arteries from 3 M-

old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice show the myogenic response to increased

pressure (active) and the dilation of the same arteries when extracellular Ca^{2+} has been

removed (passive). **(B)** Summary data of myogenic tone as a function of intraluminal

pressure. n = 5-10 vessels from 5 or 6 animals per group. ns = not significant, two-way

ANOVA. **(C)** Representative recordings of the inner diameter of isolated cerebral

arteries from 12 M-old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice. **(D)** Summary data of

myogenic tone as a function of intraluminal pressure. n = 13 vessels from 6 or 9 animals

per group. *P ≤ 0.05, two-way ANOVA. **(E)** Representative membrane potential (mV)

recordings of smooth muscle cells (SMCs) in pressurized cerebral arteries isolated from

3 M-old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice at 20 mmHg and 80 mmHg intraluminal

pressure. **(F)** Summary data showing membrane potential of SMCs in pressurized

cerebral arteries at 20 mmHg and 80 mmHg intraluminal pressure. n = 4-5 vessels from

3 or 4 animals per group. ns = not significant, unpaired t-test. **(G)** Representative

membrane potential recordings of SMCs in pressurized cerebral arteries isolated from

12 M-old *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice at 20 mmHg and 80 mmHg intraluminal

pressure. **(H)** Summary data showing membrane potential of SMCs in pressurized

cerebral arteries at 20 mmHg and 80 mmHg intraluminal pressure. n = 7-9 vessels from

5 or 6 mice per group. *P ≤ 0.05, unpaired t-test.

Figure 3. Age-dependent declines in BK and TRPM4 channel activity in SMCs

from _Col4a1+/G1344D_ mice. (A) Representative traces of spontaneous transient outward

currents (STOCs) in freshly isolated cerebral artery smooth muscle cells (SMCs) from 3

M-old *Col4a1*[+/+] and *Col4a1*[+/G1344D] mice over a range of membrane potentials (-60 to 0

mV). Summary data showing STOC frequency **(B)** and amplitude **(C)** at each command

potential. n = 8-12 cells from 5 animals per group. ns = not significant, two-way ANOVA.

(D) Representative traces of STOCs in cerebral artery SMCs from 12 M-old *Col4a1*[+/+]

and *Col4a1*[+/G1344D] mice. Summary data showing STOC frequency **(E)** and amplitude

(F) at each command potential. n = 12 cells from 5 animals per group. *P ≤ 0.05, two-

way ANOVA. **(G)** Representative I-V plots of whole-cell patch-clamp TRPM4 current

recordings during voltage ramps (-100 to 100 mV) in cerebral artery SMCs from 3 M-old

Col4a1[+/+] and *Col4a1*[+/G1344D] mice. Currents were evoked by including 200 µM free Ca^{2+}

in the intracellular solution, and the TRPM4 portion of the current was determined by

adding TRPM4 blocker 9-phenanthrol (30 µM). **(H)** Summary of whole-cell TRPM4

current density at +100 mV. n = 8-10 cells from 3 or 4 animals per group. ns = not

significant, unpaired t-test. **(I)** Representative I-V plots of whole-cell patch-clamp

TRPM4 current recordings in cerebral artery SMCs from 12 M-old *Col4a1*[+/+] and

Col4a1[+/G1344D] mice. **(J)** Summary of whole-cell TRPM4 current density at +100 mV. n =

10-12 cells from 4 animals per group. ns = not significant, unpaired t-test. **(K)**

Representative traces of transient inward cation currents (TICCs) evoked by applying

negative pressure (-20 mmHg) to stretch the plasma membrane of voltage-clamped (-70

mV) cerebral artery SMCs isolated from 3 M-old *Col4a1*[+/+] and *Col4a1*[+/G1344D] mice. **(L)**

Summary data of TICC activity. n = 5-6 cells from 3 animals per group. ns = not

significant, unpaired t-test. **(M)** Representative traces of TICCs in cerebral artery SMCs

isolated from 12 M-old *Col4a1*[+/+] and *Col4a1*[+/G1344D] mice. **(N)** Summary data of TICC

activity. n = 8-10 cells from 4 or 5 animals per group. *P ≤ 0.05, unpaired t-test.

Figure 4. Defective intracellular Ca^{2+} signaling in cerebral artery SMCs from 12 M-old _Col4a1$^{+/G1344D}$_ mice. (A) Representative time-course spinning disk confocal images exhibiting Ca^{2+} spark events in Fluo-4-AM loaded cerebral artery smooth muscle cells (SMCs) from 3 M-old _Col4a1$^{+/+}$_ and _Col4a1$^{+/G1344D}$_ mice. Time = milliseconds, scale bar = 5 μm. **(B)** Representative traces of Ca^{2+} spark events in cerebral artery SMCs isolated from 3 M-old _Col4a1$^{+/+}$_ and _Col4a1$^{+/G1344D}$_ mice presented as changes in fractional fluorescence ($\Delta F/F_0$) vs. time. **(C)** Summary data showing Ca^{2+} spark frequency, amplitude, duration, rise time, and decay. n = 4-6 cells from 3 animals per group. ns = not significant, unpaired t-test. **(D)** Representative time-course images exhibiting Ca^{2+} spark events in cerebral artery SMCs from 12 M-old _Col4a1$^{+/+}$_ and _Col4a1$^{+/G1344D}$_. Time = milliseconds, scale bar = 5 μm. **(E)** Representative traces of Ca^{2+} spark events in cerebral artery SMCs isolated from 12 M-old _Col4a1$^{+/+}$_ and _Col4a1$^{+/G1344D}$_ mice presented as $\Delta F/F_0$ vs. time. **(F)** Summary data showing Ca^{2+} spark frequency, amplitude, duration, rise time, and decay. n = 7-8 cells from 3 or 4 animals per group. *P ≤ 0.05, ns = not significant, unpaired t-test. **(G)** Representative traces showing whole-cell F/F_0 changes in response to caffeine (10 mM) in Fluo-4-AM loaded cerebral artery SMCs from 3 M-old _Col4a1$^{+/+}$_ and _Col4a1$^{+/G1344D}$_ mice. **(H)** Summary data of the $\Delta F/F_0$ in response to caffeine. n = 8-9 cells from 3 animals per group. ns = not significant, unpaired t-test. **(I)** Representative traces showing whole-cell F/F_0 changes in response to caffeine in SMCs from 12 M-old _Col4a1$^{+/+}$_ and _Col4a1$^{+/G1344D}$_ mice. **(J)** Summary data of the $\Delta F/F_0$ in response to caffeine. n = 6-7 cells from 3 animals per group. *P ≤ 0.05, unpaired t-test. **(K)** Representative traces showing whole-cell F/F_0 changes in response to U46619 (100 nM) in Fluo-4-AM loaded cerebral artery SMCs from 3 M-old _Col4a1$^{+/+}$_

694   and *Col4a1$^{+/G1344D}$* mice. **(L)** Summary data of the $\Delta F/F_0$ in response to U46619. n = 8-9

695   cells from 3 animals per group. ns = not significant, unpaired t-test. **(M)** Representative

696   traces showing whole-cell F/F_0 changes in response to U46619 in SMCs from 12 M-old

697   *Col4a1$^{+/+}$* and *Col4a1$^{+/G1344D}$* mice. **(N)** Summary data of the $\Delta F/F_0$ in response to

698   U46619. n = 6-7 cells from 3 animals per group. *$P \leq 0.05$, unpaired t-test.

Figure 5. 4PBA prevents age-related cerebral artery defects in *Col4a1+/G1344D* mice.

(A) Representative traces showing whole-cell F/F$_0$ changes in response to caffeine (10

mM) in Fluo-4-AM loaded cerebral artery SMCs from 4PBA treated 12 M-old $Col4a1^{+/+}$

and $Col4a1^{+/G1344D}$ and untreated $Col4a1^{+/G1344D}$ mice. **(B)** Summary data of the $\Delta F/F_0$ in

response to caffeine. n = 6-12 cells from 3-4 animals per group. *P ≤ 0.05, ns = not

significant, one-way ANOVA. **(C)** Representative traces showing whole-cell F/F_0

changes in response to U46619 (100 nM) in Fluo-4-AM loaded cerebral artery SMCs

from 4PBA treated 12 M-old $Col4a1^{+/+}$ and $Col4a1^{+/G1344D}$ and untreated $Col4a1^{+/G1344D}$

mice. **(D)** Summary data of the $\Delta F/F_0$ in response to U46619. n = 4-6 cells from 3-4

animals per group. *P ≤ 0.05, ns = not significant, one-way ANOVA. **(E)** Representative

traces of transient inward cation currents (TICCs) evoked by applying negative pressure

(-20 mmHg) to stretch the plasma membrane of voltage-clamped (-70 mV) cerebral

artery SMCs isolated from 4PBA treated 12 M-old $Col4a1^{+/+}$ and $Col4a1^{+/G1344D}$ and

untreated $Col4a1^{+/G1344D}$ mice. **(F)** Summary data of TICC activity. n = 7-11 cells from 3-

4 animals per group. *P ≤ 0.05, ns = not significant, one-way ANOVA. **(G)**

Representative recordings of the inner diameter of isolated cerebral arteries from 4PBA-

treated 12 M-old $Col4a1^{+/+}$ and $Col4a1^{+/G1344D}$ and untreated $Col4a1^{+/G1344D}$ mice

showing the myogenic response to increases in pressure (active) and the dilation of the

same arteries when extracellular Ca^{2+} has been removed (passive). **(H)** Summary data

of myogenic tone as a function of intraluminal pressure. n = 6 or 9 vessels from 3 or 5

mice. *P ≤ 0.05, ns = not significant, two-way ANOVA. **(I)** Representative images of

Prussian blue stained brain sections from 4PBA treated 12 M-old $Col4a1^{+/+}$ and

$Col4a1^{+/G1344D}$ and untreated $Col4a1^{+/G1344D}$ mice. Scale bar = 1 mm. **(J)** Summary data

presented as percentage of brain area with Prussian blue staining in brain sections (500

µm intervals). N = 3 animals per group. *P ≤ 0.05, ns = not significant, one-way ANOVA.

References Cited

1.   Markus HS. Genes, endothelial function and cerebral small vessel disease in man. *Exp Physiol*. 2008;93:121-127. doi: 10.1113/expphysiol.2007.038752

2.   Pantoni L. Cerebral small vessel disease: from pathogenesis and clinical characteristics to therapeutic challenges. *Lancet Neurol*. 2010;9:689-701. doi: 10.1016/S1474-4422(10)70104-6

3.   Choi JC. Genetics of cerebral small vessel disease. *J Stroke*. 2015;17:7-16. doi: 10.5853/jos.2015.17.1.7

4.   Joutel A, Faraci FM. Cerebral small vessel disease: insights and opportunities from mouse models of collagen IV-related small vessel disease and cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy. *Stroke*. 2014;45:1215-1221. doi: 10.1161/STROKEAHA.113.002878

5.   Joutel A, Haddad I, Ratelade J, Nelson MT. Perturbations of the cerebrovascular matrisome: A convergent mechanism in small vessel disease of the brain? *J Cereb Blood Flow Metab*. 2016;36:143-157. doi: 10.1038/jcbfm.2015.62

6.   Rutten-Jacobs LC, Traylor M, Adib-Samii P, Thijs V, Sudlow C, Rothwell PM, Boncoraglio G, Dichgans M, Bevan S, Meschia J, et al. Common NOTCH3 Variants and Cerebral Small-Vessel Disease. *Stroke*. 2015;46:1482-1487. doi: 10.1161/STROKEAHA.114.008540

7.   Gould DB, Phalan FC, Breedveld GJ, van Mil SE, Smith RS, Schimenti JC, Aguglia U, van der Knaap MS, Heutink P, John SW. Mutations in Col4a1 cause perinatal cerebral hemorrhage and porencephaly. *Science*. 2005;308:1167-1171. doi: 10.1126/science.1109418

8.   Gould DB, Phalan FC, van Mil SE, Sundberg JP, Vahedi K, Massin P, Bousser MG, Heutink P, Miner JH, Tournier-Lasserve E, et al. Role of COL4A1 in small-vessel disease and hemorrhagic stroke. *N Engl J Med*. 2006;354:1489-1496. doi: 10.1056/NEJMoa053727

9.   Lanfranconi S, Markus HS. COL4A1 mutations as a monogenic cause of cerebral small vessel disease: a systematic review. *Stroke*. 2010;41:e513-518. doi: 10.1161/STROKEAHA.110.581918

10.   Jeanne M, Jorgensen J, Gould DB. Molecular and Genetic Analyses of Collagen Type IV Mutant Mouse Models of Spontaneous Intracerebral Hemorrhage Identify Mechanisms for Stroke Prevention. *Circulation*. 2015;131:1555-1565. doi: 10.1161/CIRCULATIONAHA.114.013395

11.   Jeanne M, Labelle-Dumais C, Jorgensen J, Kauffman WB, Mancini GM, Favor J, Valant V, Greenberg SM, Rosand J, Gould DB. COL4A2 mutations impair COL4A1 and COL4A2 secretion and cause hemorrhagic stroke. *Am J Hum Genet*. 2012;90:91-101. doi: 10.1016/j.ajhg.2011.11.022

12.   Gould DB, Marchant JK, Savinova OV, Smith RS, John SW. Col4a1 mutation causes endoplasmic reticulum stress and genetically modifiable ocular dysgenesis. *Hum Mol Genet*. 2007;16:798-807. doi: 10.1093/hmg/ddm024

13.   Kuo DS, Labelle-Dumais C, Mao M, Jeanne M, Kauffman WB, Allen J, Favor J, Gould DB. Allelic heterogeneity contributes to variability in ocular dysgenesis,

770 myopathy and brain malformations caused by Col4a1 and Col4a2 mutations.
771 *Hum Mol Genet*. 2014;23:1709-1722. doi: 10.1093/hmg/ddt560

14. Davis MJ, Hill MA. Signaling mechanisms underlying the vascular myogenic
773 response. *Physiol Rev*. 1999;79:387-423. doi: 10.1152/physrev.1999.79.2.387

15. Gonzales AL, Garcia ZI, Amberg GC, Earley S. Pharmacological inhibition of
775 TRPM4 hyperpolarizes vascular smooth muscle. *Am J Physiol Cell Physiol*.
776 2010;299:C1195-1202. doi: 10.1152/ajpcell.00269.2010

16. Earley S, Waldron BJ, Brayden JE. Critical role for transient receptor potential
778 channel TRPM4 in myogenic constriction of cerebral arteries. *Circ Res*.
779 2004;95:922-929. doi: 10.1161/01.RES.0000147311.54833.03

17. Reading SA, Brayden JE. Central role of TRPM4 channels in cerebral blood flow
781 regulation. *Stroke*. 2007;38:2322-2328. doi: 10.1161/STROKEAHA.107.483404

18. Nelson MT, Cheng H, Rubart M, Santana LF, Bonev AD, Knot HJ, Lederer WJ.
783 Relaxation of arterial smooth muscle by calcium sparks. *Science*. 1995;270:633-
784 637. doi: 10.1126/science.270.5236.633

19. Knot HJ, Standen NB, Nelson MT. Ryanodine receptors regulate arterial
786 diameter and wall [Ca2+] in cerebral arteries of rat via Ca2+-dependent K+
787 channels. *J Physiol*. 1998;508 (Pt 1):211-221. doi: 10.1111/j.1469-
788 7793.1998.211br.x

20. Jackson WF. Potassium Channels in Regulation of Vascular Smooth Muscle
790 Contraction and Growth. *Adv Pharmacol*. 2017;78:89-144. doi:
791 10.1016/bs.apha.2016.07.001

21. Gonzales AL, Amberg GC, Earley S. Ca2+ release from the sarcoplasmic
793 reticulum is required for sustained TRPM4 activity in cerebral artery smooth
794 muscle cells. *Am J Physiol Cell Physiol*. 2010;299:C279-288. doi:
795 10.1152/ajpcell.00550.2009

22. Dabertrand F, Kroigaard C, Bonev AD, Cognat E, Dalsgaard T, Domenga-Denier
797 V, Hill-Eubanks DC, Brayden JE, Joutel A, Nelson MT. Potassium
798 channelopathy-like defect underlies early-stage cerebrovascular dysfunction in a
799 genetic model of small vessel disease. *Proc Natl Acad Sci U S A*.
800 2015;112:E796-805. doi: 10.1073/pnas.1420765112

23. Murray LS, Lu Y, Taggart A, Van Regemorter N, Vilain C, Abramowicz M, Kadler
802 KE, Van Agtmael T. Chemical chaperone treatment reduces intracellular
803 accumulation of mutant collagen IV and ameliorates the cellular phenotype of a
804 COL4A2 mutation that causes haemorrhagic stroke. *Hum Mol Genet*.
805 2014;23:283-292. doi: 10.1093/hmg/ddt418

24. Malo A, Kruger B, Goke B, Kubisch CH. 4-Phenylbutyric acid reduces
807 endoplasmic reticulum stress, trypsin activation, and acinar cell apoptosis while
808 increasing secretion in rat pancreatic acini. *Pancreas*. 2013;42:92-101. doi:
809 10.1097/MPA.0b013e318259f6ca

25. Perlmutter DH. Chemical chaperones: a pharmacological strategy for disorders
811 of protein folding and trafficking. *Pediatr Res*. 2002;52:832-836. doi:
812 10.1203/00006450-200212000-00004

26. Hayashi G, Labelle-Dumais C, Gould DB. Use of sodium 4-phenylbutyrate to
define therapeutic parameters for reducing intracerebral hemorrhage and

myopathy in Col4a1 mutant mice. *Dis Model Mech*. 2018;11. doi:
10.1242/dmm.034157

27.   Cannistraro RJ, Badi M, Eidelman BH, Dickson DW, Middlebrooks EH, Meschia
JF. CNS small vessel disease: A clinical review. *Neurology*. 2019;92:1146-1156.
doi: 10.1212/WNL.0000000000007654

28.   Hilal S, Mok V, Youn YC, Wong A, Ikram MK, Chen CL. Prevalence, risk factors
and consequences of cerebral small vessel diseases: data from three Asian
countries. *J Neurol Neurosurg Psychiatry*. 2017;88:669-674. doi: 10.1136/jnnp-
2016-315324

29.   Meuwissen ME, Halley DJ, Smit LS, Lequin MH, Cobben JM, de Coo R, van
Harssel J, Sallevelt S, Woldringh G, van der Knaap MS, et al. The expanding
phenotype of COL4A1 and COL4A2 mutations: clinical data on 13 newly
identified families and a review of the literature. *Genet Med*. 2015;17:843-853.
doi: 10.1038/gim.2014.210

30.   Yamasaki E, Ali S, Sanchez Solano A, Thakore P, Smith M, Wang X, Labelle-
Dumais C, Gould DB, Earley S. Faulty TRPM4 channels underlie age-dependent
cerebral vascular dysfunction in Gould syndrome. *Proc Natl Acad Sci U S A*.
2023;120:e2217327120. doi: 10.1073/pnas.2217327120

31.   Shi Y, Wardlaw JM. Update on cerebral small vessel disease: a dynamic whole-
brain disease. *Stroke Vasc Neurol*. 2016;1:83-92. doi: 10.1136/svn-2016-000035

32.   Wenceslau CF, McCarthy CG, Earley S, England SK, Filosa JA, Goulopoulou S,
Gutterman DD, Isakson BE, Kanagy NL, Martinez-Lemus LA, et al. Guidelines for
the measurement of vascular function and structure in isolated arteries and
veins. *Am J Physiol Heart Circ Physiol*. 2021;321:H77-H111. doi:
10.1152/ajpheart.01021.2020

33.   Harder DR. Pressure-dependent membrane depolarization in cat middle cerebral
artery. *Circ Res*. 1984;55:197-202. doi: 10.1161/01.res.55.2.197

34.   Knot HJ, Nelson MT. Regulation of arterial diameter and wall [Ca2+] in cerebral
arteries of rat by membrane potential and intravascular pressure. *J Physiol*.
1998;508 (Pt 1):199-209. doi: 10.1111/j.1469-7793.1998.199br.x

35.   Pritchard HAT, Gonzales AL, Pires PW, Drumm BT, Ko EA, Sanders KM, Hennig
GW, Earley S. Microtubule structures underlying the sarcoplasmic reticulum
support peripheral coupling sites to regulate smooth muscle contractility. *Sci*
Signal. 2017;10. doi: 10.1126/scisignal.aan2694

36.   Brayden JE, Nelson MT. Regulation of arterial tone by activation of calcium-
dependent potassium channels. *Science*. 1992;256:532-535. doi:
10.1126/science.1373909

37.   Launay P, Fleig A, Perraud AL, Scharenberg AM, Penner R, Kinet JP. TRPM4 is
a Ca2+-activated nonselective cation channel mediating cell membrane
depolarization. *Cell*. 2002;109:397-407. doi: 10.1016/s0092-8674(02)00719-5

38.   Gonzales AL, Yang Y, Sullivan MN, Sanders L, Dabertrand F, Hill-Eubanks DC,
Nelson MT, Earley S. A PLCgamma1-dependent, force-sensitive signaling
network in the myogenic constriction of cerebral arteries. *Sci Signal*. 2014;7:ra49.
doi: 10.1126/scisignal.2004732

39.   Earley S, Straub SV, Brayden JE. Protein kinase C regulates vascular myogenic
tone through activation of TRPM4. *Am J Physiol Heart Circ Physiol*.
2007;292:H2613-2622. doi: 10.1152/ajpheart.01286.2006

40.   Nilius B, Prenen J, Droogmans G, Voets T, Vennekens R, Freichel M,
Wissenbach U, Flockerzi V. Voltage dependence of the Ca2+-activated cation
channel TRPM4. *J Biol Chem*. 2003;278:30813-30820. doi:
10.1074/jbc.M305127200

41.   Pires PW, Ko EA, Pritchard HAT, Rudokas M, Yamasaki E, Earley S. The
angiotensin II receptor type 1b is the primary sensor of intraluminal pressure in
cerebral artery smooth muscle cells. *J Physiol*. 2017;595:4735-4753. doi:
10.1113/JP274310

42.   Yamasaki E, Thakore P, Krishnan V, Earley S. Differential expression of
angiotensin II type 1 receptor subtypes within the cerebral microvasculature. *Am
J Physiol Heart Circ Physiol*. 2020;318:H461-H469. doi:
10.1152/ajpheart.00582.2019

43.   Schleifenbaum J, Kassmann M, Szijarto IA, Hercule HC, Tano JY, Weinert S,
Heidenreich M, Pathan AR, Anistan YM, Alenina N, et al. Stretch-activation of
angiotensin II type 1a receptors contributes to the myogenic response of mouse
mesenteric and renal arteries. *Circ Res*. 2014;115:263-272. doi:
10.1161/CIRCRESAHA.115.302882

44.   Yasuda N, Miura S, Akazawa H, Tanaka T, Qin Y, Kiya Y, Imaizumi S, Fujino M,
Ito K, Zou Y, et al. Conformational switch of angiotensin II type 1 receptor
underlying mechanical stress-induced activation. *EMBO Rep*. 2008;9:179-186.
doi: 10.1038/sj.embor.7401157

45.   McPherson PS, Kim YK, Valdivia H, Knudson CM, Takekura H, Franzini-
Armstrong C, Coronado R, Campbell KP. The brain ryanodine receptor: a
caffeine-sensitive calcium release channel. *Neuron*. 1991;7:17-25. doi:
10.1016/0896-6273(91)90070-g

46.   Wray S, Burdyga T. Sarcoplasmic reticulum function in smooth muscle. *Physiol
Rev*. 2010;90:113-178. doi: 10.1152/physrev.00018.2008

47.   Iannitti T, Palmieri B. Clinical and experimental applications of sodium
phenylbutyrate. *Drugs R D*. 2011;11:227-249. doi: 10.2165/11591280-
000000000-00000

48.   Joutel A, Monet-Lepretre M, Gosele C, Baron-Menguy C, Hammes A, Schmidt S,
Lemaire-Carrette B, Domenga V, Schedl A, Lacombe P, et al. Cerebrovascular
dysfunction and microcirculation rarefaction precede white matter lesions in a
mouse genetic model of cerebral ischemic small vessel disease. *J Clin Invest*.
2010;120:433-445. doi: 10.1172/JCI39733

49.   Nelson MT, Quayle JM. Physiological roles and properties of potassium channels
in arterial smooth muscle. *Am J Physiol*. 1995;268:C799-822. doi:
10.1152/ajpcell.1995.268.4.C799

50.   Krishnan V, Ali S, Gonzales AL, Thakore P, Griffin CS, Yamasaki E, Alvarado
MG, Johnson MT, Trebak M, Earley S. STIM1-dependent peripheral coupling
governs the contractility of vascular smooth muscle cells. *Elife*. 2022;11. doi:
10.7554/eLife.70278

51. McCarron JG, Craig JW, Bradley KN, Muir TC. Agonist-induced phasic and tonic
responses in smooth muscle are mediated by InsP(3). *J Cell Sci*. 2002;115:2207-
2218. doi: 10.1242/jcs.115.10.2207
52. Shmygol A, Wray S. Modulation of agonist-induced Ca2+ release by SR Ca2+
load: direct SR and cytosolic Ca2+ measurements in rat uterine myocytes. *Cell*
Calcium. 2005;37:215-223. doi: 10.1016/j.ceca.2004.10.002
53. Parekh AB, Putney JW, Jr. Store-operated calcium channels. *Physiol Rev*.
2005;85:757-810. doi: 10.1152/physrev.00057.2003
54. Fernandez RA, Wan J, Song S, Smith KA, Gu Y, Tauseef M, Tang H, Makino A,
Mehta D, Yuan JX. Upregulated expression of STIM2, TRPC6, and Orai2
contributes to the transition of pulmonary arterial smooth muscle cells from a
contractile to proliferative phenotype. *Am J Physiol Cell Physiol*. 2015;308:C581-
593. doi: 10.1152/ajpcell.00202.2014
55. Potier M, Gonzalez JC, Motiani RK, Abdullaev IF, Bisaillon JM, Singer HA,
Trebak M. Evidence for STIM1- and Orai1-dependent store-operated calcium
influx through ICRAC in vascular smooth muscle cells: role in proliferation and
migration. *FASEB J*. 2009;23:2425-2437. doi: 10.1096/fj.09-131128
56. Schroder M, Kaufman RJ. The mammalian unfolded protein response. *Annu Rev*
Biochem. 2005;74:739-789. doi: 10.1146/annurev.biochem.73.011303.074134
57. Lebeau PF, Platko K, Byun JH, Austin RC. Calcium as a reliable marker for the
quantitative assessment of endoplasmic reticulum stress in live cells. *J Biol*
Chem. 2021;296:100779. doi: 10.1016/j.jbc.2021.100779
58. Yamamoto WR, Bone RN, Sohn P, Syed F, Reissaus CA, Mosley AL, Wijeratne
AB, True JD, Tong X, Kono T, et al. Endoplasmic reticulum stress alters
ryanodine receptor function in the murine pancreatic beta cell. *J Biol Chem*.
2019;294:168-181. doi: 10.1074/jbc.RA118.005683
59. Favor J, Gloeckner CJ, Janik D, Klempt M, Neuhauser-Klaus A, Pretsch W,
Schmahl W, Quintanilla-Fend L. Type IV procollagen missense mutations
associated with defects of the eye, vascular stability, the brain, kidney function
and embryonic or postnatal viability in the mouse, Mus musculus: an extension of
the Col4a1 allelic series and the identification of the first two Col4a2 mutant
alleles. *Genetics*. 2007;175:725-736. doi: 10.1534/genetics.106.064733
60. Pallast N, Diedenhofen M, Blaschke S, Wieters F, Wiedermann D, Hoehn M,
Fink GR, Aswendt M. Processing Pipeline for Atlas-Based Imaging Data Analysis
of Structural and Functional Mouse Brain MRI (AIDAmri). *Front Neuroinform*.
2019;13:42. doi: 10.3389/fninf.2019.00042
61. Dong HW. *Allen reference atlas : a digital color brain atlas of the C57Black/6J*
male mouse. Hoboken, N.J.: Wiley; 2008.
62. Gonzales AL, Amberg GC, Earley S. Ca2+ release from the sarcoplasmic
reticulum is required for sustained TRPM4 activity in cerebral artery smooth
muscle cells. *American journal of physiology Cell physiology*. 2010;299:C279-
288.