Detection of Domestication Signals through the Analysis of the Full

Distribution of Fitness Effects using Simulations

David Castellano[1†], Ioanna-Theoni Vourlaki[2,3†], Ryan N. Gutenkunst[1*], Sebastian E. Ramos-Onsins[2*]

1. Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ, USA.
2. Centre for Research in Agricultural Genomics (CRAG), Autonomous University of Barcelona (UAB), Vall de
la Moronta s/n, Cerdanyola del Valles, Barcelona, 08193, Spain.
3. Animal Breeding and Genetics Program, IRTA, Torre Marimón, 08140 Caldes de Montbui, Barcelona, Spain
9
† Equal contribution
*Co-corresponding authors

ABSTRACT

The process of domestication involves a complex interplay between concurrent demographic

and selective changes. While we can readily observe the phenotypic effects of domestication,

the genetic consequences of domestication often remain elusive. Artificial selection can alter

the selection coefficients of both new and pre-existing genetic variation within domesticated

populations. To investigate this, we conducted simulations using a combination of population

genomic parameters designed to reflect the domestication process observed in large livestock

mammals. Our study uses forward-in-time simulations to examine the 1D and 2D site

frequency spectra (SFS) of mutations in two populations that have diverged since the

domestication split. In total, we examined eighteen different scenarios, varying the strength of

selection acting on beneficial mutations and the proportion of mutations with altered selection

coefficients post-domestication. First, we re-evaluate how linked selection and fluctuating

selection coefficients affect the accuracy of inferred demographic histories. Second, we find

that certain aspects of the full distribution of fitness effects (DFE), such as the shape and

strength of the deleterious DFE, can be accurately estimated in both populations using only the

1D-SFS. However, the accurate characterization of the beneficial DFE remains a challenge,

even when using the 2D-SFS. Third, using a novel joint DFE model, we are able to quantify

the fraction of mutations that have experienced a change in their selection coefficient (p_C)

during domestication. Interestingly, classic hard selective sweeps can mimic a substantial

increase in p_C, even when the simulated p_C was initially zero. In summary, our work highlights

the strengths and limitations of detecting changes in the DFE using a novel joint DFE model

and emphasizes the risks of over-interpreting demographic histories across a range of realistic

domestication scenarios.

Keywords: Domestication, DFE, demography, selective sweeps, linked selection, 2D-SFS,

forward-in-time simulations.

INTRODUCTION

The increase in human population and the emergence of modern society are linked to the domestication of plants and animals (Purugganan and Fuller 2009; Driscoll *et al.* 2009; Larson and Burger 2013; Amills *et al.* 2017). Human civilization it was made possible by the domestication of surrounding life forms, where humans began to domesticate plants and animals such as wheat, dogs, pigs, or chickens (Dayan 1994; Zeder *et al.* 2006; Zeder 2012, 2015; Redding 2015; Avni *et al.* 2017). Domestication is a process that allows humans and other species to establish a long-term mutualistic relationship that provides benefits to both species (Zeder *et al.* 2006). Human domestication of fauna and flora began about 10-15 thousand years ago and is still ongoing (Larson and Burger 2013; Zeder 2015). Although human civilization is based on domestication, we still lack a complete genomic and evolutionary understanding of domestication. Domestication is a rapid process on the evolutionary time scale, but it is not a discrete event and implies the gradual improvement of domesticated traits. It is believed that human-induced artificial selection during domestication can be expected to be relatively stronger and therefore faster than natural selection. However, in plants it has been shown that the evolutionary rate of domesticated varieties is similar to that of wild plants, suggesting a process similar to natural selection (Purugannan and Fuller 2010). In addition, domestication tends to be associated with bottlenecks; only a small number of individuals from the wild population are domesticated, which is expected to reduce the efficiency of natural selection (Wright *et al.* 2005). Another important difference between natural and artificial selection is that modern breeders typically use truncation selection, which is the selection of the top percentage of individuals for the desired trait (Granleese *et al.* 2019). The prevalence of truncation selection in nature, or prior to industrialization, is unknown. Truncation selection is known to be a simple and efficient form of directional selection (Crow and Kimura 1979), and no severe accumulation of genetic load is expected in outcrossing species (Kondrashov 1988; Ohta 1989) if the population size remains large enough (Marsden *et al.* 2016). A recent comprehensive meta-analysis of the genetic costs of domestication (Moyers *et al.* 2018) found that deleterious variants are more numerous (or segregate at higher frequencies) in domesticated populations compared to their wild relatives. However, this pattern may not be general, as seen in sorghum (Lozano et al. 2021). This pattern is likely driven by a number of processes that collectively reduce the effectiveness of selection in domesticated populations, as first suggested in rice genomes (Lu *et al.* 2006).

Selection, both natural and artificial, can occur through either a few loci with strong effects or
many loci with small effects (Jain and Stephan 2017a; b), depending on the genetic architecture
of the trait and the strength of selection on it. Different patterns of genetic diversity around
selected loci are expected in response to these two models of selection (Stephan and John
2020). Classic hard selective sweeps have been reported at a few candidate loci among
important domesticated traits (Andersson 2012), such as the IGF2 gene region associated with
lean domesticated pigs (Van Laere *et al.* 2003), the thyroid-stimulating hormone receptor
(TSHR) in domesticated chickens (Rubin *et al.* 2010), or the sh4 and qSW5 loci (Shomura *et*
al. 2008; Li *et al.* 2018) involved in the traits of seed shattering and grain width, respectively,
in domesticated rice (Huang *et al.* 2012). These examples are consistent with a simple
Mendelian genetic architecture, in which a few loci account for most of the variance in the
domesticated trait (for a comprehensive list of genes important for domestication, see Courtier-
Orgogozo and Martin 2020).

Polygenic adaptation, on the other hand, describes a process in which a constellation of small
changes in allele frequencies modify differences in the trait under selection, where a trait is a
phenotypic trait but also can be the fitness itself. A wide range of population genetic models
and simulations have been examined to describe polygenic adaptation (*e.g.,* de Vladar and
Barton 2014; Stephan 2016). Some models analyze the polygenic response of a trait in the
presence of mutation and stabilizing selection (de Vladar and Barton 2014; Stephan and John
2020), while others capture the response of a trait under mutation and stabilizing or directional
selection following an environmental change in a finite size population (Jain and Stephan
2017a; Devi and Jain 2023). In practice, polygenic adaptation using genetic data is harder to
detect than classic selective sweeps (Pritchard et al. 2010), but the combined use of phenotypic
data together with genetic data can detect the selective effect of quantitative traits (e.g., Chen
et al. 2021, Berg and Coop 2014). Polygenic adaptation has been detected in some specific
studies in wild, domesticated and experimental evolution populations (Barghi *et al.* 2019; Reid
et al. 2023; He *et al.* 2023).

In this study, we ask to what extent we can detect a genomic signal of domestication using a
different approach: comparing the full distribution of fitness effects (DFE) on new and standing
variation. To do this, we inferred the full DFE of new non-synonymous mutations in wild and
domesticated populations. The DFE of new deleterious mutations has previously been
estimated by contrasting the 1D-SFS of synonymous and non-synonymous mutations from a

variety of species, assuming that beneficial mutations contribute only to divergence but not to polymorphism due to their rapid fixation in the population (Keightley and Eyre-Walker 2007; Boyko *et al.* 2009; Kim *et al.* 2017; Tataru *et al.* 2017; Barton and Zeng 2018). Tataru *et al.* (2017) proposed a model, polyDFE, to infer the full DFE and the proportion of adaptive substitutions (α) using only polymorphism data. Castellano *et al.* (2019) applied polyDFE to compare the full DFE of new amino acid mutations across great apes and found that the shape of the deleterious DFE is constant across this set of closely related species. More recently, a new method using 2D-SFS has been proposed to jointly estimate the deleterious DFE between two populations that recently diverged and share many polymorphisms (Huang *et al.* 2021). Inference on 1D-SFS indicates that the inference of the parameters for a given model are made considering the SFS of a single population. Instead, the joint inference on 2D-SFS indicates that the inference is made considering the joint SFS distribution of two populations together. Note that the inference of parameters in case considering two populations can also be performed separately (which implicates the inference of parameters for each separated population -that is, the marginal distribution-), which is less informative.

However, few studies have compared the DFE between domesticated and wild populations (Leno-Colorado *et al.* 2022). First, the analysis of differences between wild and domesticated individuals has been focused on finding large effects that determine the phenotypic differences between these two groups. Therefore, DFE does not appear to be an adequate method for detecting candidate regions, as DFE considers the entire distribution of selective effects and appears to be blind to single events. Second, the split between wild and domesticated populations is very recent (expected to be no more than 10K years), suggesting that the DFE of these two populations would be very similar or identical. Third, methods to detect subtle differences in DFE between two closely related populations have not been developed. For example, polyDFE (Tataru et al. 2017) does not infer a joint DFE, but rather infers and tests the marginal DFEs.

On the other hand, the study of the demographic processes involved in domestication has been addressed in several species (e.g., Morell Miranda et al. 2023, Arnoux et al. 2020, Murray et al. 2010), where the split between wild and domesticated populations, bottlenecks, and gene flow events between these two populations have been inferred, also considering multiple selective sweeps (Caicedo et al. 2007). It has also been shown (Torres et al. 2020) that ignoring deleterious selective effects in the study of demographic patterns, can lead to biased estimates

of silent variability patterns (see also Comeron 2017, Beissinger et al 2016). Thus, some efforts

have been driven to infer the DFE and demography together to avoid deviations from the true

value (Johri et al. 2020).

Several studies have been focusing on the significant distortions in the DFE shape caused by

the effect of linked selection (e.g., Gilbert et al. 2021, Johri et al 2021, Shrider et al. 2016).

Particularly, Johri et al. (2021) studied how direct and indirect negative selection affect the

inference of population demographic history. Their results point out the bias caused by

background selection even after masking functional genomic regions and propose an ABC

method to jointly infer demography and the deleterious DFE. On the other hand, Gilbert et al.

(2021), utilized SLiM simulations (Haller and Messer 2019) to evaluate the effect of linked

selection and inbreeding. They observed a marked effect on the inference of demography and

on DFE parameters caused by extreme inbreeding patterns, but more accurate inference on

outbreeding populations. They also observed severe deviations of real parameters caused by

linkage selection, especially in regions with low recombination or high gene density. In

scenarios where they included beneficial mutations, the inferred negative DFE was

significantly deviated to an excessive proportion of mutations at moderate-low negative effects.

In another interesting study Shrider et al. (2016) show that positive selection can lead to the

mis-inference of parameters and even infer changes in the population size when no change

occurred, pointing to the effect of linked selection as the cause of the misinference.

Here, we use forward-in-time simulations that consider the domestication process under

different demographic and selective models to study the capacity to detect the selective

differences between these two (domesticated and wild) populations. Several combinations of

genetic architectures and selective effects (both beneficial and deleterious) have been

simulated, ranging from one that considers a relatively small number of loci that change their

selective effects to another that considers polygenic adaptation (considering fitness as a trait)

where many loci have divergent selective effects. Importantly, we release a new methodology

that includes an additional parameter that is crucial to distinguish populations in processes of

rapid selective change; the selective effects of a fraction of the existing variants can change

(from deleterious to beneficial and vice versa) in the domesticated population. This method

jointly infers the full DFE parameters of wild and domesticated populations, including the

inference of changes (i.e., beneficial to detrimental or vice versa) between them. We also

describe and discuss how linked selection and changing the DFE deform the recovered

demographic histories.

MATERIALS AND METHODS

Simulation of the Domestication Process

A simulation analysis of the domestication process is developed using the forward-in-time simulator SLiM4 (Haller and Messer 2023). The general model of the domestication process is developed in the SLiM script available in github (https://github.com/CastellanoED/domesticationDFE). Eighteen different domestication scenarios are analyzed, and the parameters for each scenario are shown in Table 1. All options (flags) used to run the SLiM script are also available in github (https://github.com/CastellanoED/domesticationDFE). We aim to model a general domestication process that resembles the genomic configuration, generation time, mutation and recombination landscape relevant to large domesticated mammals used as livestock. Note that we are not considering the recent processes of genetic improvement performed by commercial companies in the last decades. The constructed model assumes a genome containing a single chromosomal "chunk" or window, with 10,000 loci/exons of 120 base pairs in length, and each locus/exon with one-third (4-fold) neutral synonymous positions and two-thirds (0-fold) selected non-synonymous positions scattered along the locus.

The simulation parameters for each scenario (Figure 1A) are as follows: the initial population at time 0 run for $10*N_e$ generations to reach mutation-selection-drift equilibrium, then splits into the domesticated and wild populations. Hereafter we refer to the Wild and Domesticated populations. We aim to mimic a realistic but still general domestication process in large livestock mammals where ancestral N_e estimates are on average around 10,000 (Murray *et al.* 2010; Groenen *et al.* 2012; Larson *et al.* 2014; Frantz *et al.* 2015; Yang *et al.* 2016; Librado *et al.* 2021; Todd *et al.* 2022) and the domestication process, according to archeological records, started around 10,000 years ago (Ahmad *et al.* 2020). The average generation time in these large domesticated mammals is about 5 years per generation (Pacifici *et al.* 2013). Note that in this study we had to reduce the population size and related population parameters below from 10,000 diploid individuals to N_e=5,000 diploid individuals for computational reasons.

Genomic parameters: The mutation rate per site (μ) and generation is 2.5×10^{-7}, and the population size (N_e) is 5,000 diploid individuals, thus the expected θ under neutrality is **0.005**.

Each locus is separated from its neighbors by 3×10^{-6} recombination events per generation. The

recombination rate per site and generation within the loci is fixed to a rate of 1.5×10^{-7}

recombination events per site. Note that the higher recombination between loci aims to mimic

their real genetic distance separation (assuming a functional site density of 5%) - this greatly

speeds up the simulation as non-coding sites do not need to be simulated. In other words, we

simulate 120 Kb of coding sequence in each run, which is equivalent to simulating a 2.4 Mb

chromosome window with 5% coding sites. We perform 100 independent runs for each of the

eighteen scenarios.

Demographic parameters: The Domesticated populations of 5,000 diploid individuals suffer

a bottleneck, reducing their population size temporarily to 200 diploid individuals, to recover

again to 5,000 diploid individuals after the bottleneck. The bottleneck lasted 100 generations.

The simulation finishes 900 generations after the bottleneck. In nine of the eighteen simulated

scenarios we allow that 1% of the Wild individuals migrate to the Domesticated population

during the 100 generations of the bottleneck. Thus, during the bottleneck 25% of the

domesticated population is wild. In the other nine combinations there is no exchange of

individuals between the Wild and Domesticated populations. This demographic history is

equivalent to a 1,000 years long bottleneck followed by a 9,000 years long recovery in an

ancestral population with 10,000 diploid individuals and a generation time of 5 years.

DFE parameters. The selective effects produced by domestication are modeled by changing the

fitness values of a proportion of the existing and new mutations in the domesticated population

(at the time of the split) (Table 1). We call this probability of change p_c and it can be 0% (our

negative control), 5% or 25%. Domesticated and Wild populations show different proportions

of beneficial and deleterious new mutations depending on the scenario. The negative effects in

all scenarios and populations follow a gamma distribution with a shape value of 0.3 and a mean

of S_d = -100 ($2N_e s_d$ in the heterozygote, N_e = 5,000 diploid individuals and s_d = -1%), which

is in the range of values inferred using empirical data (Boyko et al. PLoS Genet 2008, Galtier

PLoS Genet 2016), while variants with positive effects follow an exponential distribution. We

investigate three combinations of parameters for the positive DFE: pervasive & nearly neutral

(with mean selective effects S_b of 1 and probability of being beneficial p_b of 10%), common

& weak (S_b = 10 and p_b = 1%) and rare & strong (S_b = 100 and p_b = 0.1%) (Table 1).

Types of Sites

The sites are initially divided into seven different types (named m_1 to m_7), being m_1 neutral

(synonymous) and m_2 to m_7 functional (non-synonymous) sites having a different selective

effect when mutated (see Table 2 and Figure 1C). Mutations at m_5, m_6 and m_7 sites generate

deleterious variants in the Wild population, and mutations at m_2, m_3 and m_4 sites generate

beneficial mutations in the Wild population. The selection coefficient of mutations generated

at m_2 (beneficial) or at m_5 (deleterious) sites are invariant for the Wild and Domesticated

populations. However, the mutations at m_3, m_4, m_6 and m_7 sites will change their selective

effect in the Domesticated populations relative to the Wild populations. That is, the new

selective effect is drawn from the corresponding DFE section (positive or negative),

independently of their value in the wild population. The selection coefficient of a given

beneficial mutation at m_3 sites will remain beneficial in the Domesticated population, but it

will be different from the original beneficial effect at Wild. A mutation at m_4 sites will change

its selection coefficient from beneficial in the Wild to deleterious in the Domesticated

population. Equivalently, the selection coefficient of a deleterious mutation at m_6 sites will

remain negative in the Domesticated population but it will be different from that found at Wild.

A mutation at m_7 sites will change its selection coefficient from deleterious in the Wild to

beneficial in the Domesticated population. For each of the 18x100 independent simulation runs,

we randomly precalculate the location of each site type (except for the permanent disposition

of two non-synonymous sites followed by a synonymous site within codons) using an ad hoc

R script. This hard-coding of selective effects on different sites allows us to gain insight into

the relative importance of each mutation type for the domestication process.

Distribution of fitness effects (DFE): Two complementary approaches

polyDFE: 1D-SFS and 1D-DFE

We use the polyDFEv2.0 framework (Tataru and Bataillon 2019) to estimate and compare the

DFE across Wild-Domesticated population pairs by means of likelihood ratio tests (LRTs). We

use the R function compareModels (from https://github.com/paula-

tataru/polyDFE/blob/master/postprocessing.R) to compare pairs of models. The inference is

performed only on the unfolded SFS data (divergence counts to the outgroup are not fitted),

and unfolded SFS data are fitted using a DFE model comprising both deleterious (gamma

distributed) and beneficial (exponentially distributed) mutations. The DFE of each Wild-

Domesticated population pair is inferred using the 1D-SFS of each population. polyDFE

assumes that new mutations in a genomic region arise as a Poisson process with an intensity
that is proportional to the length of the region and the mutation rate per nucleotide (μ). We
assume that μ remains constant across simulations (as it is the case). Both an ancestral SNP
misidentification error (ε) and distortion parameters (r_i) are estimated. However, we notice that
the exclusion of ε does not affect the rest of estimated parameters because under the simulation
conditions used here no sites are expected to be misidentified. The r_i parameters are fitted
independently for each frequency bin (from n = 1 to n = 19), and they are able to correct any
distortion that affects equally the SFS of synonymous and non-synonymous variants (such as,
in principle, demography or linked selection). Model averaging provides a way to obtain honest
estimates that account for model uncertainty. To produce the model average estimates of the
full DFE we weight each competing model according to their AIC following the equation 6.1
shown in the polyDFEv2 tutorial ("polyDFE/tutorial.pdf at master · paula-tataru/polyDFE").
We use the R function getAICweights (from https://github.com/paula-
tataru/polyDFE/blob/master/postprocessing.R to do the model averaging R) to obtain the AIC
values.

dadi: 2D-SFS and 2D-DFE

dadi (Gutenkunst *et al.* 2009) is employed to infer the joint distribution of fitness effects
(Jerison *et al.* 2014; Ragsdale *et al.* 2016; Huang *et al.* 2021) and the demographic history of
all simulated population pairs. Our new model for the joint DFE between the two populations
is a mixture of multiple components designed to mimic the selected mutation types in the
simulations (Table 2; Figure 1D). The major exception is that beneficial mutations are
modelled to have a single fixed selection coefficient, rather than arising from an exponential
distribution. Let p_{+w} be the fraction of mutations that are positively selected in the Wild
population, p_C be the fraction of mutations that change selection coefficient in the
Domesticated population, and p_{C+} be the fraction of those mutations that become beneficial in
the Domesticated population (note in our simulations $p_{+w} = p_{C+}$). To model mutation types
m2 and m3, a proportion p_{+w} x (1-p_C) + p_{+w} x p_C x p_{C+} of mutations are assumed to have the
same fixed positive selection coefficient in both populations. To model m4, a proportion p_{+w}
x p_C x (1-p_{C+}) is assumed to have a fixed positive selection coefficient in the Wild population
and a gamma-distributed negative selection coefficient in the Domesticated population. To
model m5, a proportion (1-p_{+w}) x (1-p_C) of mutations are assumed to have equal negative

gamma-distributed selection coefficients in the two populations. To model m_6, a proportion $(1\text{-}p_{+w})$ x p_C x $(1\text{-}p_{C+})$ is assumed to have independent gamma-distributed selection coefficients in the two populations. To model m_7, a proportion $(1\text{-}p_{+w})$ x p_C x p_{C+} mutations is assumed to have a gamma-distributed negative selection coefficient in the Wild population and a fixed positive selection coefficient in the Domesticated population. All gamma distributions are assumed to have the same shape and scale. This model is implemented in *dadi* as the function **dadi.DFE.Vourlaki_mixture**. Note in our simulations the marginal 1D-DFEs of Wild and Domesticated populations are exactly the same; the difference is that in Domesticated populations a given fraction of sites (some already polymorphic, some still monomorphic) can change their selection coefficient relative to the Wild population.

Figure 1: Joint demographic and DFE models simulated and fit. **A:** Illustration of the joint demographic model used in SLiM simulations. N_a: Effective population size of the Ancestral population. N_{ew}: Effective population

size of the Wild population. N_{e1d}: Effective population size of the Domesticated population during the bottleneck.
N_{e2d}: Effective population size of the Domesticated population after the bottleneck. T_1: Number of generations
in the bottleneck period. T_2: Number of generations from the bottleneck to the present. m: Wild to Domesticated
migration rate (migration occurs along T_1). **B:** Illustration of a more general joint demographic model used in the
dadi inferences. N_a: Effective population size of the Ancestral population. N_{pre}: Effective population size before
the domestication split. N_{1div}: Effective population size of the Wild population after the split. N_{1F}: Effective
population size of the Wild population at the end of the simulation. N_{2div}: Effective population size of the
Domesticated population after the split. N_{2F}: Effective population size of the Domesticated population at the end
of the simulation. T_{pre}: Number of generations before the domestication split. T_{div}: Number of generations after
the bottleneck. T_{1F}: Number of generations under N_{1F}. T_{2F}: Number of generations under N_{2F}. Note that T_{1F}
and T_{2F} are estimated independently and that T_{1F} can be the same, longer or shorter than T_{2F}. m_d: Wild to
Domesticated migration rate. m_w: Domesticated to Wild migration rate. Both migration rates occur after the
domestication split. **C:** Illustration of the joint DFE model used in the SLiM simulations, with mutation types
illustrated. **D:** Illustration of the joint DFE model used in the *dadi* inferences, in which a fixed positive selection
coefficient is assumed.

For inference, a new, more general demographic model with branch-independent population
size changes is first fit to the synonymous mutations from each simulation, and then the newly
proposed joint DFE model is fit to the non-synonymous mutations. The parameters of the
demographic model (Figure 1B) are estimated by running 100 optimizations per inference unit.
The 2D-SFS for selected sites are precomputed conditional on the demography for 104^2 values
of γ_1 and γ_2 (*$2N_a s$*, a population scaled selection coefficient for the heterozygote where N_a is
the ancestral population size), 102 negative and 2 positives. For the negative part of the DFE,
γ values were logarithmically equally spaced between -2000 and -10^{-4}. The expected DFE for
selected sites can then be computed as a weighted sum over these cached spectra (Kim *et al.*
2017). The DFE parameters shape, scale, p_{+w}, p_c, and p_{c+} are then estimated by maximizing
the Poisson likelihood of the simulated data, with the non-synonymous rate of mutation influx
fixed to twice that inferred for neutral sites in the demographic history fit. For the DFE
inference, optimization is repeated until the best three results are within 0.5 log-likelihood
units. Ancestral state misidentification is modelled, however in our simulations no sites are
expected to be misidentified.

For the purpose of this work, *dadi* software is downloaded and installed according to the
instructions   provided   at   the   following   link:

https://bitbucket.org/gutenkunstlab/dadi/src/master/. Since *dadi* operates as a module of

Python, the Anaconda3 and Spyder (Python 3.7, Rossum and Drake 2009; Anaconda 2016;

Raybaut 2009) versions are used in this study.

340   Inference units, and confidence intervals in demographic and DFE parameters

To obtain the sampling variance of parameter estimates and approximate confidence intervals,

we use a bootstrap approach. We resample with replacement 100 times 20 independent

simulation runs or chromosomal "chunks" (from a pool of 100 "chunks") and concatenate

them. Hence, each concatenated unit (or inference unit) is made of 24 Mb of coding sequence

(as comparison, the human genome contains ~26 Mb of coding sequence). Uncertainties of

DFE parameter inferences in polyDFE and *dadi* are calculated by this conventional

bootstrapping, but in *dadi* we hold the demographic model fixed. In polyDFE the distortion

introduced by demography (and linked selection) is not estimated but corrected with the r_i

parameters. Note that our procedure with *dadi* does not propagate uncertainty in demographic

parameters through to the DFE parameters. To obtain the sampling variance of demographic

parameter estimates with *dadi* we use the Godambe approach as described in Coffman *et al.*

2016. A final consideration on the factor of two differences across simulation and inference

tools. In polyDFE, *s* is defined as the selection coefficient on the heterozygote (as in *dadi*), and

the scaled selection coefficient is defined as $4N_e s$, while in *dadi* it is defined as $2N_a s$ (in the

ancestral population). In SLiM2, *s* is defined as the selection coefficient on the homozygote.

RESULTS AND DISCUSSION

Studying the effect of domestication on the DFE of natural populations can be very challenging, especially when available methods for inferring and comparing the DFE have not been compared using exactly the same dataset. In the present work, we conduct a simulation study using different combinations of parameters relevant to the domestication process. Note that the main difference between domestication demographic model used here and demographic models used in speciation studies is the time scale since the split occurred. Our simulated domesticated populations experience a large or small change in the number and selective effects of loci under domestication after a bottleneck period, with or without migration. Hereafter, we refer to the Wild and Domesticated populations. In this study we are interested in the evolutionary process of domestication from the split to domesticated lineages. We are not considering the programs of genetic improvement that have been performed in the last decades for some domesticated animals, which can highly increase the levels of inbreeding (e.g., Makanjuola et al. 2020 estimated up to 40% levels of inbreeding in some cattle breeds submitted to intense genomic selection). The models simulated in this work include strong selection as well as reduction in population sizes, which together could also moderately increase the level of inbreeding in populations. In any case, Gilbert et al. (2021) observed that only high levels of inbreeding (>80%) affected severely the inference of the DFE.

The Wild populations have a constant DFE and constant population size, but limited recombination across loci to mimic a realistic recombination landscape. Beneficial mutations arise at Wild populations following an exponential distribution, while deleterious mutations are drawn from a gamma distribution with shape 0.3 and mean $S_d = -100$ (where $S_d = 2N_a s_d$, the selection coefficient s_d in the heterozygote is -1%, and $N_a = 5,000$ diploid individuals is the ancestral effective population size, see Material and Methods: Simulating the Domestication Process). All mutations, beneficial and deleterious, are co-dominant. The Domesticated population originates from the Wild population through a bottleneck and a concomitant change in selective effects at a fraction of non-synonymous sites (Figure 1; Table 1). The recombination and mutation landscapes are drawn from the same distribution in the Domesticated and Wild populations.

Table 1

Variable demographic and selective parameters across scenarios

Positive DFE	Migration (W>D)	p_c	Scenario ID
		0%	1
	0%	5%	2
$p_b = 10\%$ & $S_b = 1$		25%	3
		0%	4
Pervasive and nearly neutral	1%	5%	5
		25%	6
		0%	7
	0%	5%	8
$p_b = 1\%$ & $S_b = 10$		25%	9
		0%	10
Common and weak	1%	5%	11
		25%	12
		0%	13
	0%	5%	14
$p_b = 0.1\%$ & $S_b = 100$		25%	15
		0%	16
Rare and strong	1%	5%	17
		25%	18

The eighteen simulated combinations of parameters in this study. The first column refers to the DFE of new beneficial mutations, the second column represents the migration rate from the Wild to the Domesticated population and the third column shows the percentage of sites that change their selection coefficients in the Domesticated population (p_c). Last column shows the ID we use to quickly label scenarios along the manuscript.

The change in selective effects affects both new mutations that arise within the Domesticated population and existing variants that existed before the domestication event. Put simply, not only can mutations that were deleterious (or beneficial) before the population split become

beneficial (or deleterious) within the domesticated population, but even if the direction of the selective effect remains the same, the intensity of selection can change. Table 2 shows all the combinations of changes in selective effects between Wild and Domesticated populations. Our simulated scenarios aim to cover a variety of possible changes in the genetic architecture (number of loci) and the strength of selection (selection coefficients) of the trait/s under domestication. Three DFEs for beneficial mutations are assumed: (i) pervasive and nearly neutral, where a large fraction of new mutations (10%) are on average nearly neutral ($S_b = 1$), (ii) common and weak, where beneficial mutations are still fairly common (1%) but weakly selected ($S_b = 10$) and (iii) rare and strong, where very few mutations (0.1%) are strongly beneficial ($S_b = 100$). Depending on the scenario, a selective change occurs only at a small (5%) or at a substantial proportion (25%) of sites in the Domesticated population (Table 1, "p_c" column). We leave six scenarios as negative controls; the selection coefficients of new and standing variation in the Domesticated and Wild populations are exactly the same. Finally, demographic changes affect only the Domesticated population; the Wild population evolves under a constant population size. Two versions of the same demographic model (Figure 1A) are simulated: (i) one with migration, and (ii) another without migration. When there is migration, it only occurs from the Wild to the Domesticated population during the domestication bottleneck.

Table 2

Types of sites in simulated scenarios

Site	Wild	Domesticated
m_1	Neutral	No change, remain Neutral
m_2	Beneficial	No change, remain Beneficial
m_3	Beneficial	Change to a different Beneficial Effect
m_4	Beneficial	Change to Deleterious
m_5	Deleterious	No change, remain Deleterious
m_6	Deleterious	Change to a different Deleterious Effect
m_7	Deleterious	Change to Beneficial

Estimation of demographic parameters in Wild and Domesticated populations

In this study, we investigate the effects of natural selection, both in a general sense and specifically the change in selection coefficient in shared variation due to artificial selection, on our ability to reconstruct the demographic history of domestication and the DFE. We do this using two commonly used inference tools (polyDFE and *dadi*) that assume free recombination across loci. Note that *dadi* first infers the demographic history and then infers the DFE assuming those inferred demographic parameters, whereas polyDFE operates independently of specific demographic histories and is designed to correct for distortions that affect both synonymous and non-synonymous site frequency spectra equally (Tataru and Bataillon 2019). Figure 1 A and B show the simulated joint demographic model and the joint demographic model used in the *dadi* inferences, respectively. We have increased the complexity of the inference model by introducing additional parameters, allowing it to account not only for "simulated" or true demographic changes, but also for more complex and unknown demographic histories and the potential influence of linked selection on synonymous SFS. The diagnostic plots can be found in Supplementary Figure 1; there is good agreement between the model fits and the data.

Our findings indicate that when positive selection is relatively weak ($S_b = 1$ or $S_b = 10$), the estimated onset of domestication tends to be approximately twice as old as the actual simulated starting point. Additionally, the bottleneck appears slightly shallower but considerably longer than the simulated value (see Figure 2 and Supplementary Table 1 for the confidence intervals). This suggests that the influence of linked selection, likely driven primarily by background selection when $S_b <= 10$, has the effect of elongating the timeline. Consequently, it makes the domestication divergence and bottleneck appear more ancient and extended, respectively. For the Wild populations we always detect a larger population expansion than for the Domesticated populations, but without a bottleneck. This signal of a recent expansion in the Wild population is expected because when we consider how linked selection affects the SFS, there are more rare synonymous polymorphisms compared to what we would expect if there was free recombination under a constant population size (Nielsen 2005). Remarkably, when positive selection is strong ($S_b = 100$), the temporal stretch becomes even more pronounced and the demographic history of both populations overlap extensively. The domestication divergence shifts to approximately 50,000 years ago, whereas the actual simulated split occurred 10,000 years ago. Additionally, the bottleneck appears significantly longer and less severe, while there

is an inferred large population expansion in both Wild and Domesticated populations. Although in Figure 2 there appears to be a change in population size before the domestication split, only three scenarios (with IDs 5, 6, and 7) are statistically significant (Supplementary Table 1 and Supplementary Figure 2). Interestingly, we find the migration rate from Wild to Domesticated (m_{w2d}) and from Domesticated to Wild (m_{d2w}) are overestimated in all scenarios (Supplementary Table 1 and Supplementary Figure 3). We observe that neither migration nor an increase in p_c appears to significantly change the inferred demographic histories that we have just described.

Torres et al. (2020) and Johri et al. (2022) detected a distortion in the patterns of diversity due to deleterious effects. Instead, we detected a distortion in the demographic inference that differs depending on positive selective effects. This distortion has been already described by Shrider et al. (2016) using a single population simulation study that included strong beneficial selection, indicating the important role of linked selection. Our findings indicate that when linked selection is at play, the reconstructed demographic history captures certain elements of the actual simulated history. For instance, if positive selection is not strong, it successfully identifies a bottleneck in Domesticated populations compared to Wild populations. However, when positive selection is strong ($S_b = 100$), it tends to "erase" the demographic history through indirect selection effects and recreate large recent population expansions in both populations. Nevertheless, and more importantly, in all scenarios it is not possible to accurately determine the timing of the onset of domestication, the duration of the domestication bottleneck, or to distinguish between the presence and absence of migration between populations. We believe that these aspects are crucial for contextualizing the role of domestication in human history. Unfortunately, either the 2D-SFS or our modeling assumptions (or both) do not seem to be useful in this context. Thus, the next question is to what extent can the nuisance r_i parameters from polyDFE or this distorted inferred demography from *dadi* help to recover the simulated DFE parameters?

Figure 2. Solid lines showing the inferred demographic histories for the eighteen simulated scenarios. In salmon the Wild population and in light-blue the Domesticated population. The dashed line shows the true simulated demography in Domesticated populations. The true Wild population is not shown but it is a constant population size with relative $Ne = 1$. The 95% confidence intervals calculated using the Godambe approximation can be found in Supplementary Table 1.

Can domestication be detected as an artificial change in the marginal full DFE between populations? No.

Next, we investigate whether polyDFE captures differences in the marginal (or 1D) full DFE of Domesticated and Wild populations across the eighteen domestication scenarios (Table 1). We run five nested models (Table 3) and compare them using likelihood ratio tests (LRTs) (Supplementary Table 2). It is important to note that in all our simulations, the marginal full DFE for new mutations in both Domesticated and Wild populations is the same within a given scenario (as detailed in Table 1). This means that the selection coefficients for sites, whether they are monomorphic or polymorphic, are drawn from the same full DFE. In simpler terms, the proportion of new mutations that are advantageous or detrimental is identical for both Domesticated and Wild populations within a given scenario. The key distinction lies in the fact that when $p_C > 0\%$, Domesticated populations might have a higher number of advantageous mutations as polymorphisms. This is because some of these beneficial mutations were already present at intermediate or high frequencies as nearly neutral polymorphisms in the ancestral population, and we expect that migration after the domestication split can also re-introduce beneficial mutations from the Wild to the Domesticated population.

Table 3

List of nested polyDFE models and (co)estimated parameters.

Model ID	Negative DFE		Positive DFE	
	shape	S_d	p_b	S_b
M1	Var	Var	-	-
M10	Fix	Var	-	-
M2	Var	Var	Var	Var
M20	Fix	Var	Var	Var
M30	Fix	Var	Fix	Fix

Independently estimated parameters for the Domesticated and Wild populations (Var). Jointly estimated parameters for the Domesticated and Wild populations (Fix). S_d is always independently estimated to accommodate potential changes in N_e between populations. The population mutation rate (Θ), the nuisance

parameters (r_i) and the mispolarization parameter (ε) are all independently estimated across Wild and Domesticated populations.

LRTs between different nested models allow us to address important questions about the DFE, without assuming any prior knowledge of our datasets. First, we assess whether the inferred shape of the negative DFE is similar in both populations while also examining if the estimation of the shape parameter is influenced by the presence of advantageous mutations. When comparing models that do not consider beneficial mutations (models M1 versus M10 in the second column of Supplementary Table 2), the model with a distinct shape for Domesticated and Wild populations is accepted only in two, rather unrelated, scenarios (scenarios 7 and 14). This indicates that an artificial alteration in the shape of the deleterious DFE between Domesticated and Wild populations can be inferred. Fortunately, when comparing models that take into account beneficial mutations (models M2 vs M20, third column in Supplementary Table 2), all scenarios show a shared shape of the deleterious DFE, which is expected based on the simulation parameters. These findings suggest that disregarding beneficial mutations can cause an artificial change in the inferred shape of the marginal deleterious DFE between populations, as noted previously by Tataru et al. in 2017. Second, incorporating the positive DFE yields statistically significant results in all scenarios (see Supplementary Table 2, fourth and fifth columns). Hence, polyDFE appears to effectively detect beneficial mutations, regardless of the strength of positive selection. Third, we investigate whether Domesticated and Wild populations could exhibit an artificial change in the beneficial DFEs as a consequence of domestication. When comparing the M20 and M30 models (refer to the last column in Supplementary Table 2), polyDFE invokes changes in the positive DFE between populations in some scenarios. This is applicable when there is no migration between the populations and only a minimal amount (5%) or when none of the sites change their selection coefficient, regardless of the mean strength of positive selection. These artificial changes occur in scenarios 2, 7, 8, 13, and 14, with a marginal p-value in scenario 1. It is noteworthy that the polyDFE analysis shows no significant difference in the positive DFE between populations with the presence of migration and when p_c equals 25%. Our initial expectation was that in scenarios with a large fraction of sites changing selection coefficients and migration from the Wild population, would result in an increase in the load of advantageous polymorphisms in the Domesticated population (due to the re-introduction of beneficial mutations from the Wild to the Domesticated population), leading to a higher inferred rate of new advantageous mutations (p_b) compared to the Wild population. We do not observe this result. We suspect that the

absence of this result could be due to linkage between selected mutations and synonymous
mutations, which may lead to an overcorrection of the excess of non-synonymous
polymorphisms at high frequency via the r_i parameters.

Tartaru et al. (2016) polyDFE suggests that specific domestication scenarios, particularly those
lacking migration and featuring minimal changes in selection coefficients, may artificially alter
the marginal full DFE between populations, particularly in its positive side. In the next section,
we find that the artificial change in the marginal full DFE is due to the detection of a higher
proportion of new, effectively neutral, advantageous mutations (>10%) in Domesticated
populations compared to Wild populations. Hence, this finding has no significant impact on
the marginal full DFE differences between the populations when the DFE is represented in
discrete intervals. We demonstrate that domestication does not significantly impact polyDFE's
ability to detect a false difference in the marginal full DFE among populations. We conclude
that if a significant change is detected in the discretized marginal full DFE, it must be
considered valid.

Estimation of DFE parameters in Wild and Domesticated populations

Under the polyDFE framework, we begin by extracting the Akaike Information Criterion (AIC)
from every model (Table 3) and then computing the AIC-weighted parameters for all models
(Tataru and Bataillon 2019; Castellano *et al.* 2019). This approach is used because the true
model generating real data in both Wild and Domesticated populations is unknown. Instead,
under *dadi*'s framework, we adopt an alternative methodology that utilizes very general,
parameter rich and versatile joint demographic and DFE models to fit the 2D-SFS. The
diagnostic plots of the new joint DFE model is shown in Supplementary Figure 1, again there
is good agreement between the model fits and the data.

Figure 3. Sampling distributions of estimated parameters for the deleterious DFE are obtained using 100 bootstrap replicates. Dotted vertical lines indicate the actual simulated parameter values. A) Shape parameter estimated with polyDFE in dark gray the Wild population and in light gray the Domesticated population, B) shape parameter estimated with *dadi*, C) mean s_d estimated with polyDFE in dark gray the Wild population and in light gray the Domesticated population and D) mean s_d estimated with *dadi*. To calculate s_d from inferred S_d values, we divided S_d by 4 times the N_e estimate in polyDFE and by 2 times the N_a estimate in *dadi*. To obtain the N_e (and N_a) we divide π at synonymous sites by the true simulated mutation rate (2.5×10^{-7} per site and generation). Note that *dadi* uses the joint demographic model parameters to compute π at synonymous sites, while polyDFE assumes a constant population size.

Figure 3 depicts the distribution of parameters related to the deleterious DFE that are estimated
by performing bootstrap analysis using polyDFE and *dadi*. We observe that both tools have a
tendency to marginally overestimate the shape parameter of the gamma distribution employed
to model the deleterious DFE. The overestimation is particularly significant in polyDFE, when
positive selection is rare and strong. In such scenarios, *dadi*'s shape estimation is sometimes
rather noisy. Regarding the mean of the deleterious (s_d), we observe quite accurate inferences
with both tools across domestication scenarios. This finding indicates that regardless of the
inference method used, the estimation of the deleterious DFE is resilient to demographic and
selective changes, as well as the pervasive impact of linked selection. In contrast, our previous
inference with *dadi* on the demographic parameters concluded that linked selection
significantly complicates the process of obtaining accurate demographic histories. Therefore,
although it is generally believed that demographic changes should be considered to infer the
underlying DFE, we found that inferring the deleterious DFE is "easier" than inferring the true
demographic history. We conclude that correcting the non-synonymous SFS through nuisance
r_i parameters or using a demographic history that fits the data well, even if it is incorrect,
appears sufficient for obtaining an accurate depiction of the deleterious DFE.

The distribution of parameters associated with the beneficial DFE, estimated by bootstrap
analysis using polyDFE and *dadi*, is shown in Supplementary Figure 4 and Supplementary
Table 3. Depending on the scenario, we simulate an average increase in relative fitness (s_b) of
1%, 0.1%, and 0.01%. Positive selection's strength is usually substantially underestimated by
polyDFE and *dadi*, but only polyDFE consistently overestimates the proportion of new
advantageous mutations (p_b), regardless of the true simulated value. Supplementary Figure 5
displays the discretized full DFE. Given the distribution of inferred values of p_b and s_b, we
reason that a peak of effectively neutral advantageous mutations is being measured by
polyDFE. The overall excess of effectively neutral advantageous mutations measured by
polyDFE is generally balanced by the defect of effectively neutral deleterious mutations.
Consequently, polyDFE seems to have limited power in identifying effectively beneficial
mutations on the 1D-SFS (under these simulation conditions). As suggested before, the r_i
parameters might be overcorrecting for the increase in high-frequency non-synonymous
polymorphisms in the SFS expected from beneficial mutations. By the same logic, *dadi's*
inferred demographic history might be operating in a similar way to explain the uptick of

synonymous polymorphisms at high frequency due to linkage to beneficial mutations. Thus,
we conclude that both tools struggle to infer the positive DFE and tend to be overconservative
and identify weaker positive selection than what has been simulated. However, it is noteworthy
that both tools typically yield comparable (and reasonably accurate) discretized deleterious
DFEs upon considering the tendency of polyDFE to infer a peak of effectively neutral
beneficial mutations. Noteworthy that this finding is in agreement with what was already point
our by Tataru et al. (2017) and Booker et al (2020) using a single population. They focus
attention on the challenge to infer parameters of positive selection when counting for weak and
strong selected mutations. Indeed, Booker et al. (2020) enphasizes that, in case of having rare
and strong positive selection, the study of the site frequency spectrum can be very noisy, where
linked sites play an important role that hinders the inference of the full DFE.

617 Estimation of the fraction of mutations with divergent selective effects (p_C) between
618 Domesticated and Wild populations

One of the main goals of this study is to determine the proportion of new and standing non-
synonymous mutations with differing selection coefficients in Wild and Domesticated
populations. The usage of our new joint DFE model is not limited to the current study. Our
new model, created by mixing multiple distributions to mimic mutation types in our
simulations (Table 2; Figure 1C-D), is suitable for usage in any recently diverged populations.
Hence, while we acknowledge that our simulation and inference pipeline has the potential to
provide insights into recent parapatric and allopatric speciation events, our primary focus in
this work is on assessing our ability to identify the impact of domestication on the full DFE
within domesticated populations. This emphasis is due to the availability of independent
archaeological evidence that can be used to determine the timing of domestication onset.

Figure 4 displays the distribution of the inferred p_C for three different positive DFEs, along
with simulated p_C values. When positive selection is not strong, it becomes apparent that
scenarios with a significant fraction of mutations with dissimilar selective effects ($p_C = 25\%$)
can readily be differentiated from those where a small ($p_C = 5\%$) or nonexistent ($p_C = 0\%$)
number of sites alter their selection coefficient. However, differentiating our negative control
from a positive control proves difficult when only 5% of the sites show a difference in their

selection coefficients. Notably, we overestimate p_C significantly in cases of strong positive

selection, indicating that hard selective sweeps may mimic divergent selection in a substantial

amount of non-synonymous mutations. We observe no major impact of migration on the

inferred p_C values across scenarios.

Figure 4. Sampling distributions of inferred p_C are obtained using 100 bootstrap replicates. In

light green scenarios without migration and in dark green scenarios with migration.

The overestimation of p_C when positive selection is strong is not surprising, since mutations

with stable selection coefficients between populations may be in close recombinational

proximity and can hitchhike with strongly beneficial mutations that are population-specific.

This will exacerbate the apparent fraction of mutations with divergent selective effects. In

contrast, if positive selection is weaker, recombination will be able to disentangle beneficial

mutations from the rest of mutation types and simplify our estimation of p_C. One way to

ameliorate this problem would be to remove genomic windows with evidence of recent

population-specific, complete or partial, selective sweeps and rerun our inference pipeline. For

example, these could be regions with low neutral genetic diversity. However, we find this

heuristic solution might be difficult to implement in practice.

Supplementary Figure 6 shows the observed level of neutral genetic diversity (measured using

Watterson's theta and synonymous sites, θ_S) and the selective constraint (i.e., the ratio of non-

synonymous polymorphisms to synonymous polymorphisms per site, P_n/P_S) for each

independent simulation run. Note the large decrease in the observed θ_S, driven entirely by

linked selection in Wild populations, relative to the expected level of neutral genetic diversity

(expected $\theta_S = 0.005$ under free recombination). Particularly important is the reduction in the

average θ_S across independent simulation runs in Wild populations when positive selection is

rare and strong (θ_S is ~20% of the expected value), whereas when positive selection is weaker

(and perhaps absent if $S_b = 1$) the observed level of genetic diversity is ~40% of the expected

value. In strong positive selection scenarios, there may be no heuristic correction or genomic

region that escapes *genetic draft* (Gillespie 2000), and our current definition and interpretation

of p_C would be misleading. We also observe that although synonymous genetic diversity and

the selective constraint is higher in Wild populations than in Domesticated populations, Wild

populations show more variability in θ_S and P_n/P_S across independent simulation runs. This is

likely explained by both the higher chance of sampling strongly beneficial mutations and the

greater selection efficacy expected in the constant population size of Wild populations.

Moreover, relative to Domesticated populations, we find that a higher p_C always corresponds

to a lower θ_S and selective constraint (all else being equal). The largest difference in the level

of genetic diversity and P_n/P_S between Domesticated and Wild populations occurs when the

simulated p_C is substantial (25%), and S_b is less than 100. Thus, for a given joint demographic

history, there appears to be a positive relationship between the simulated p_C and the difference

in genetic diversity and selective constraint between Wild and Domesticated populations. This

observation only applies when positive selection is not strong. When positive selection is

strong, genetic diversity and selective pressure are significantly further reduced in both

Domesticated and Wild populations, causing the two distributions to largely overlap. As

described above for the reconstruction of demographic history, when selective sweeps are

strong, the recovered demographic history also tends to overlap between Wild and

Domesticated populations. The overlap of demographic histories and neutral genetic diversity

distributions could be used as a caution signal and as an indicator of strong positive selection

and widespread genetic draft. Finally, migration appears to cause a minor reduction in P_n/P_S

and increase genetic diversity within Domesticated populations. Thus, migration acts slightly

diminishing the P_n/P_S discrepancy between Wild and Domesticated populations.

Implications for empirical analysis of populations

Although the time of separation between wild and domesticated is recent, the external

environments in which they live are very different. This means that an unspecified proportion

of environmentally influenced variants may have changed their fitness effect in the

domesticated population, which may have altered their frequencies. In this work, we simulated

differential effects between wild and domesticated populations, and we observed that selective

effects affect the inference of demographic parameters by linked selection, but to different

extents depending on the DFE. Linked selection caused by positive effects is responsible for

strong differences from the true parameters, even if they are very rare but strong. Nevertheless,

the DFE can be estimated quite accurately, suggesting that methods that infer the entire DFE

can be used first and then estimate the demographic parameters using this information.

Another point of interest for empirical geneticists is the development of a new method to jointly

infer the DFE between wild and domesticated and their differences in the positive part of the

distribution. The 2D dadi extension algorithm allows to infer differences in p+w (the fraction

of mutations that are positively selected in the wild population), pc (the fraction of mutations

that change the coefficient of selection in the domesticated population), pc+ (the fraction of

those mutations that become beneficial in the domesticated population).

CONCLUSIONS

In summary, our use of forward-in-time simulations has provided valuable insights into the complex genetic demographic history and distribution of fitness effects (DFE) for both new and standing amino acid mutations in the context of domestication. Through a comparative analysis of two methods, polyDFE and *dadi*, and the new implementation of a full 2D-SFS full inference of DFE, we have uncovered the impact of linked selection on the reconstructed demographic history of both wild and domesticated populations. Despite biases in the timelines of domestication events and bottleneck characteristics, the estimation of deleterious DFE remains remarkably reliable, demonstrating the robustness of these analytical approaches in the studied conditions. In particular, the underestimation of effectively beneficial mutations in the DFE highlights the influence of linkage between beneficial and neutral mutations, which requires careful consideration in model interpretation. In addition, our results shed light on distinguishing scenarios of divergent selective effects between populations under weak and strong positive selection, providing a nuanced understanding of the interplay of evolutionary forces. As we navigate the complex landscape of domestication, these methodological approaches contribute significantly to unraveling the evolutionary dynamics and adaptive processes that shape the genomes of domesticated organisms, and provide a foundation for future research in this critical area of study.

Supplementary Information

We have added the Diagnostic plots 2D-SFS for synonymous and Nonsynonymous positions obtained with *dadi* for all the scenarios analyzed, plus six additional Figures and three additional Tables in the Supplementary Material. This Supplementary Information is available Github at https://github.com/CastellanoED/domesticationDFE/blob/main/Supp_Material.zip, and the scripts used in the analyses of this work are available at the link https://github.com/CastellanoED/domesticationDFE.

730
731

Acknowledgments

SERO. was supported by grants AGL2016-78709-R (MEC, Spain) and PID2020-119255GB-I00 (MICINN, Spain), by the CERCA Programme/Generalitat de Catalunya and acknowledges financial support from the Spanish Ministry of Economy and Competitiveness, through the Severo Ochoa Programme for Centres of Excellence in R&D 2016-2019 and 2020-2023 (SEV-2015-0533, CEX2019-000917) and the European Regional Development Fund (ERDF). RNG and DC were supported by the National Institute of General Medical Sciences of the National Institutes of Health (R35GM149235 and R01GM127348 to RNG). ITV was supported by a predoctoral fellowship funded by MCIN/AEI/ 10.13039/501100011033 through the Grant BES-2017-081139 and by "ESF Investing in your future". This research was also supported by NSF Grant No. PHY-1748958 and the Gordon and Betty Moore Foundation Grant No. 2919.02.

LITERATURE CITED

Ahmad H. I., M. J. Ahmad, F. Jabbir, S. Ahmar, N. Ahmad, *et al.*, 2020 The Domestication Makeup: Evolution, Survival, and Challenges. Front. Ecol. Evol. 8.

Amills M., H. J. W. C. Megens, A. Manunza, S. E. Ramos-Onsins, and M. Groenen, 2017 A genomic perspective about wild boar demography and evolution, pp. 376–387 in *Ecology, Conservation and Management of Wild Pigs and Peccaries*, Cambridge University Press.

Anaconda Software Distribution. Computer software. Vers. 2-2.4.0. Anaconda, Nov. 2016. Web.

Andersson L., 2012 Genetics of Animal Domestication, pp. 260–274 in *Biodiversity in Agriculture: Domestication, Evolution, and Sustainability*, edited by Damania A. B., Qualset C. O., McGuire P. E., Gepts P., Bettinger R. L., *et al.* Cambridge University Press, Cambridge.

Arnoux S, Fraïsse C, Sauvage C. Genomic inference of complex domestication histories in three Solanaceae species. J Evol Biol. 2021 Feb;34(2):270-283. doi: 10.1111/jeb.13723. Epub 2020 Nov 10. PMID: 33107098.

Avni R., M. Nave, O. Barad, K. Baruch, S. O. Twardziok, *et al.*, 2017 Wild emmer genome architecture and diversity elucidate wheat evolution and domestication. Science 357: 93–97. https://doi.org/10.1126/science.aan0032

Barghi N., R. Tobler, V. Nolte, A. M. Jakšić, F. Mallard, *et al.*, 2019 Genetic redundancy fuels polygenic adaptation in Drosophila. PLOS Biol. 17: e3000128. https://doi.org/10.1371/journal.pbio.3000128

Barton H. J., and K. Zeng, 2018 New Methods for Inferring the Distribution of Fitness Effects for INDELs and SNPs. Mol. Biol. Evol. 35: 1536–1546. https://doi.org/10.1093/molbev/msy054

Beissinger TM, Wang L, Crosby K, Durvasula A, Hufford MB, Ross-Ibarra J. Recent demography drives changes in linked selection across the maize genome. Nat Plants. 2016 Jun 13;2:16084. doi: 10.1038/nplants.2016.84. PMID: 27294617.

Berg J. J., and G. Coop, 2014 A population genetic signal of polygenic adaptation. PLoS Genet.

10: e1004412. https://doi.org/10.1371/journal.pgen.1004412

Booker T.R. 2020 Inferring Parameters of the Distribution of Fitness Effects of New Mutations

When Beneficial Mutations Are Strongly Advantageous and Rare. G3 (Bethesda). 10(7):2317-

2326. doi: 10.1534/g3.120.401052.

Boyko A. R., R. H. Boyko, C. M. Boyko, H. G. Parker, M. Castelhano, et al., 2009 Complex

population structure in African village dogs and its implications for inferring dog domestication

history.   Proc.   Natl.   Acad.   Sci.   U.   S.   A.   106:   13903–13908.

https://doi.org/10.1073/pnas.0902129106

Caicedo AL, Williamson SH, Hernandez RD, Boyko A, Fledel-Alon A, York TL, Polato NR,

Olsen KM, Nielsen R, McCouch SR, Bustamante CD, Purugganan MD. Genome-wide patterns

of nucleotide polymorphism in domesticated rice. PLoS Genet. 2007 Sep;3(9):1745-56. doi:

10.1371/journal.pgen.0030163. Epub 2007 Aug 6.

Castellano D., M. C. Macià, P. Tataru, T. Bataillon, and K. Munch, 2019 Comparison of the

Full Distribution of Fitness Effects of New Amino Acid Mutations Across Great Apes.

Genetics 213: 953–966. https://doi.org/10.1534/genetics.119.302494

Chen Q, Samayoa LF, Yang CJ, Olukolu BA, York AM, Sanchez-Gonzalez JJ, Xue W,

Glaubitz JC, Bradbury PJ, Romay MC, Sun Q, Buckler ES, Holland JB, Doebley JF. A

conserved genetic architecture among populations of the maize progenitor, teosinte, was

radically   altered   by   domestication.   Proc   Natl   Acad   Sci   U   S   A.   2021   Oct

26;118(43):e2112970118.   doi: 10.1073/pnas.2112970118.   PMID: 34686607;   PMCID:

PMC8639367.

Coffman A. J., P. H. Hsieh, S. Gravel, and R. N. Gutenkunst, 2016 Computationally Efficient

Composite Likelihood Statistics for Demographic Inference. Mol. Biol. Evol. 33: 591–593.

https://doi.org/10.1093/molbev/msv255

Comeron JM. Background selection as null hypothesis in population genomics: insights and

challenges from Drosophila studies. Philos Trans R Soc Lond B Biol Sci. 2017 Dec

19;372(1736):20160471.   doi: 10.1098/rstb.2016.0471.   PMID: 29109230;   PMCID:

PMC5698629.

Courtier-Orgogozo V., and A. Martin, 2020 The coding loci of evolution and domestication:
current knowledge and implications for bio-inspired genome editing, (M. H. Dickinson, 27 L.
B. Vosshall, and J. A. T. Dow, Eds.). J. Exp. Biol. 223: jeb208934.
https://doi.org/10.1242/jeb.208934

Crow J. F., and M. Kimura, 1979 Efficiency of truncation selection*. Proc. Natl. Acad. Sci. U.
S. A. 76: 396–399. Dayan T., 1994 Early Domesticated Dogs of the Near East. J. Archaeol.
Sci. 21: 633–640. https://doi.org/10.1006/jasc.1994.1062

Devi A., and K. Jain, 2023 Polygenic adaptation dynamics in large, finite populations.
2023.01.25.525607. Driscoll C. A., D. W. Macdonald, and S. J. O'Brien, 2009 From wild
animals to domestic pets, an evolutionary view of domestication. Proc. Natl. Acad. Sci. 106:
9971–9978. https://doi.org/10.1073/pnas.0901586106

Frantz L. A. F., J. G. Schraiber, O. Madsen, H.-J. Megens, A. Cagan, *et al.*, 2015 Evidence of
long-term gene flow and selection during domestication from analyses of Eurasian wild and
domestic pig genomes. Nat. Genet. 47: 1141–1148. https://doi.org/10.1038/ng.3394

Galtier N. Adaptive Protein Evolution in Animals and the Effective Population Size
Hypothesis. PLoS Genet. 2016 Jan 11;12(1):e1005774. doi: 10.1371/journal.pgen.1005774.
PMID: 26752180; PMCID: PMC4709115.

Gilbert KJ, Zdraljevic S, Cook DE, Cutter AD, Andersen EC, Baer CF. The distribution of
mutational effects on fitness in Caenorhabditis elegans inferred from standing genetic
variation. Genetics. 2022 Jan 4;220(1):iyab166. doi: 10.1093/genetics/iyab166.

Gillespie J. H., 2000 Genetic drift in an infinite population. The pseudohitchhiking model.
Genetics 155: 909–919. https://doi.org/10.1093/genetics/155.2.909

Granleese T., S. A. Clark, and J. H. J. van der Werf, 2019 Genotyping strategies of selection
candidates in livestock breeding programmes. J. Anim. Breed. Genet. 136: 91–101.
https://doi.org/10.1111/jbg.12381

Groenen M. A. M., A. L. Archibald, H. Uenishi, C. K. Tuggle, Y. Takeuchi, *et al.*, 2012
Analyses of pig genomes provide insight into porcine demography and evolution. Nature 491:
393–398. https://doi.org/10.1038/nature11622

Gutenkunst R. N., R. D. Hernandez, S. H. Williamson, and C. D. Bustamante, 2009 Inferring

the Joint Demographic History of Multiple Populations from Multidimensional SNP Frequency

Data. PLOS Genet. 5: e1000695. https://doi.org/10.1371/journal.pgen.1000695

Haller B. C., and P. W. Messer, 2023 SLiM 4: Multispecies Eco-Evolutionary Modeling. Am.

Nat. 201: E127–E139. https://doi.org/10.1086/723601

He Y., Y. Guo, W. Zheng, T. Yue, H. Zhang, *et al.*, 2023 Polygenic adaptation leads to a higher

reproductive fitness of native Tibetans at high altitude. Curr. Biol. CB 33: 4037-4051.e5.

https://doi.org/10.1016/j.cub.2023.08.021

Huang X., A. L. Fortier, A. J. Coffman, T. J. Struck, M. N. Irby, *et al.*, 2021 Inferring Genome-

Wide Correlations of Mutation Fitness Effects between Populations. Mol. Biol. Evol. 38:

4588–4602. https://doi.org/10.1093/molbev/msab162

Huang X., N. Kurata, X. Wei, Z.-X. Wang, A. Wang, *et al.*, 2012 A map of rice genome

variation reveals the origin of cultivated rice. Nature 490: 497–501.

https://doi.org/10.1038/nature11532

Jain K., and W. Stephan, 2017a Rapid Adaptation of a Polygenic Trait After a Sudden

Environmental Shift. Genetics 206: 389–406. https://doi.org/10.1534/genetics.116.196972

Jain K., and W. Stephan, 2017b Modes of Rapid Polygenic Adaptation. Mol. Biol. Evol. 34:

3169–3175. https://doi.org/10.1093/molbev/msx240

Jerison E. R., S. Kryazhimskiy, and M. M. Desai, 2014 Pleiotropic consequences of adaptation

across gradations of environmental stress in budding yeast

Johri P., Riall K., Becher H., Excoffier L., Charlesworth B., Jensen J.D., 2021 The Impact of

Purifying and Background Selection on the Inference of Population History: Problems and

Prospects. Mol Biol Evol. 38(7):2986-3003. doi: 10.1093/molbev/msab050.

Keightley P. D., and A. Eyre-Walker, 2007 Joint inference of the distribution of fitness effects

of deleterious mutations and population demography based on nucleotide polymorphism

frequencies. Genetics 177: 2251–2261. https://doi.org/10.1534/genetics.107.080663

Kim B. Y., C. D. Huber, and K. E. Lohmueller, 2017 Inference of the Distribution of Selection

Coefficients for New Nonsynonymous Mutations Using Large Samples. Genetics 206: 345–

361. https://doi.org/10.1534/genetics.116.197145

Kondrashov A. S., 1988 Deleterious mutations and the evolution of sexual reproduction.

Nature 336: 435–440. https://doi.org/10.1038/336435a0

Larson G., and J. Burger, 2013 A population genetics view of animal domestication. Trends

Genet. TIG 29: 197–205. https://doi.org/10.1016/j.tig.2013.01.003

Larson G., D. R. Piperno, R. G. Allaby, M. D. Purugganan, L. Andersson, *et al.*, 2014 Current

perspectives and the future of domestication studies. Proc. Natl. Acad. Sci. 111: 6139–6146.

https://doi.org/10.1073/pnas.1323964111

Leno-Colorado J., S. Guirao-Rico, M. Pérez-Enciso, and S. E. Ramos-Onsins, 2022 Pervasive

selection pressure in wild and domestic pigs. bioRxiv 2020.09.09.289439; doi:

https://doi.org/10.1101/2020.09.09.289439

Li N., R. Xu, P. Duan, and Y. Li, 2018 Control of grain size in rice. Plant Reprod. 31: 237–

251. https://doi.org/10.1007/s00497-018-0333-6

Librado P., N. Khan, A. Fages, M. A. Kusliy, T. Suchan, *et al.*, 2021 The origins and spread of

domestic horses from the Western Eurasian steppes. Nature 598: 634–640.

https://doi.org/10.1038/s41586-021-04018-9

Lozano R, Gazave E, Dos Santos JPR, Stetter MG, Valluru R, Bandillo N, Fernandes SB,

Brown PJ, Shakoor N, Mockler TC, Cooper EA, Taylor Perkins M, Buckler ES, Ross-Ibarra J,

Gore MA. Comparative evolutionary genetics of deleterious load in sorghum and maize. Nat

Plants. 2021 Jan;7(1):17-24. doi: 10.1038/s41477-020-00834-5. Epub 2021 Jan 15. PMID:

33452486.

Lu J., T. Tang, H. Tang, J. Huang, S. Shi, *et al.*, 2006 The accumulation of deleterious

mutations in rice genomes: a hypothesis on the cost of domestication. Trends Genet. 22: 126–

131. https://doi.org/10.1016/j.tig.2006.01.004

Makanjuola BO, Miglior F, Abdalla EA, Maltecca C, Schenkel FS, Baes CF. Effect of genomic
selection on rate of inbreeding and coancestry and effective population size of Holstein and
Jersey cattle populations. J Dairy Sci. 2020 Jun;103(6):5183-5199. doi: 10.3168/jds.2019-
18013.

Marsden C. D., D. Ortega-Del Vecchyo, D. P. O'Brien, J. F. Taylor, O. Ramirez, *et al.*, 2016
Bottlenecks and selective sweeps during domestication have increased deleterious genetic
variation in dogs. Proc. Natl. Acad. Sci. 113: 152–157.
https://doi.org/10.1073/pnas.1512501113

Morell Miranda P, Soares AER, Günther T. Demographic reconstruction of the Western sheep
expansion from whole-genome sequences. G3 (Bethesda). 2023 Nov 1;13(11):jkad199. doi:
10.1093/g3journal/jkad199. PMID: 37675574.

Moyers, Morrell, and McKay, 2018 Genetic Costs of Domestication and Improvement. J.
Hered. 109. https://doi.org/10.1093/jhered/esx069 Murray C., E. Huerta-Sanchez, F. Casey,
and D. G. Bradley, 2010 Cattle demographic history modelled from autosomal sequence
variation. Philos. Trans. R. Soc. B Biol. Sci. 365: 2531–2539.
https://doi.org/10.1098/rstb.2010.0103

Murray C, Huerta-Sanchez E, Casey F, Bradley DG. Cattle demographic history modelled from
autosomal sequence variation. Philos Trans R Soc Lond B Biol Sci. 2010 Aug
27;365(1552):2531-9. doi: 10.1098/rstb.2010.0103. PMID: 20643743; PMCID:
PMC2935105.

Nielsen R., 2005 Molecular signatures of natural selection. Annu. Rev. Genet. 39: 197–218.
https://doi.org/10.1146/annurev.genet.39.073003.112420

Ohta T., 1989 The mutational load of a multigene family with uniform members. Genet. Res.
53: 141–145. https://doi.org/10.1017/s0016672300028020

Pacifici M., L. Santini, M. D. Marco, D. Baisero, L. Francucci, *et al.*, 2013 Generation length
for mammals. Nat. Conserv. 5: 89–94. https://doi.org/10.3897/natureconservation.5.5734

Pritchard J. K., J. K. Pickrell, and G. Coop, 2010 The Genetics of Human Adaptation: Hard
Sweeps, Soft Sweeps, and Polygenic Adaptation. Curr. Biol. 20: R208–R215.
https://doi.org/10.1016/j.cub.2009.11.055

Purugganan M. D., and D. Q. Fuller, 2009 The nature of selection during plant domestication.
Nature 457: 843–848. https://doi.org/10.1038/nature07895

Ragsdale A. P., A. J. Coffman, P. Hsieh, T. J. Struck, and R. N. Gutenkunst, 2016 Triallelic
Population Genomics for Inferring Correlated Fitness Effects of Same Site Nonsynonymous
Mutations. Genetics 203: 513–523. https://doi.org/10.1534/genetics.115.184812

Raybaut, P. (2009). Spyder-documentation. *Available Online at: Pythonhosted. Org*.
Redding R. W., 2015 The Pig and the Chicken in the Middle East: Modeling Human
Subsistence Behavior in the Archaeological Record Using Historical and Animal Husbandry
Data. J. Archaeol. Res. 23: 325–368. https://doi.org/10.1007/s10814-015-9083-2

Reid B. N., B. Star, and M. L. Pinsky, 2023 Detecting parallel polygenic adaptation to novel
evolutionary pressure in wild populations: a case study in Atlantic cod (Gadus morhua). Philos.
Trans. R. Soc. B Biol. Sci. 378: 20220190. https://doi.org/10.1098/rstb.2022.0190

Rossum G. V., and F. L. Drake, 2009 *Python 3 Reference Manual: (Python Documentation*
Manual Part 2). CreateSpace Independent Publishing Platform.

Rubin C.-J., M. C. Zody, J. Eriksson, J. R. S. Meadows, E. Sherwood, *et al.*, 2010 Whole-
genome resequencing reveals loci under selection during chicken domestication. Nature 464:
587–591. https://doi.org/10.1038/nature08832

Schrider D.R., Shanku A.G., Kern A.D. 2016 Effects of Linked Selective Sweeps on
Demographic Inference and Model Selection. Genetics. 204(3):1207-1223. doi:
10.1534/genetics.116.190223.

Shomura A., T. Izawa, K. Ebana, T. Ebitani, H. Kanegae, *et al.*, 2008 Deletion in a gene
associated with grain size increased yields during rice domestication. Nat. Genet. 40: 1023–
1028. https://doi.org/10.1038/ng.169

Stephan W., 2016 Signatures of positive selection: from selective sweeps at individual loci to
subtle allele frequency changes in polygenic adaptation. Mol. Ecol. 25: 79–88.
https://doi.org/10.1111/mec.13288

Stephan W., and S. John, 2020 Polygenic Adaptation in a Population of Finite Size. Entropy
22: 907. https://doi.org/10.3390/e22080907

Tataru P., and T. Bataillon, 2019 polyDFEv2.0: testing for invariance of the distribution of
fitness effects within and across species. Bioinforma. Oxf. Engl. 35: 2868–2869.
https://doi.org/10.1093/bioinformatics/bty1060

Tataru P., M. Mollion, S. Glémin, and T. Bataillon, 2017 Inference of Distribution of Fitness
Effects and Proportion of Adaptive Substitutions from Polymorphism Data. Genetics 207:
1103–1119. https://doi.org/10.1534/genetics.117.300323

Tataru P.2018. polyDFE/tutorial.pdf at master. https://github.com/paula-tataru/polyDFE

Todd E. T., L. Tonasso-Calvière, L. Chauvey, S. Schiavinato, A. Fages, *et al.*, 2022 The
genomic history and global expansion of domestic donkeys. Science 377: 1172–1180.
https://doi.org/10.1126/science.abo3503

Torres R, Stetter MG, Hernandez RD, Ross-Ibarra J. The Temporal Dynamics of Background
Selection in Nonequilibrium Populations. Genetics. 2020 Apr;214(4):1019-1030. doi:
10.1534/genetics.119.302892. Epub 2020 Feb 18. PMID: 32071195; PMCID: PMC7153942.

Van Laere A.-S., M. Nguyen, M. Braunschweig, C. Nezer, C. Collette, *et al.*, 2003 A regulatory
mutation in IGF2 causes a major QTL effect on muscle growth in the pig. Nature 425: 832–
836. https://doi.org/10.1038/nature02064

Vladar H. P. de, and N. Barton, 2014 Stability and response of polygenic traits to stabilizing
selection and mutation. Genetics 197: 749–767. https://doi.org/10.1534/genetics.113.159111

Wright S. I., I. V. Bi, S. G. Schroeder, M. Yamasaki, J. F. Doebley, *et al.*, 2005 The effects of
artificial selection on the maize genome. Science 308: 1310–1314.
https://doi.org/10.1126/science.1107891

Yang J., W.-R. Li, F.-H. Lv, S.-G. He, S.-L. Tian, *et al.*, 2016 Whole-Genome Sequencing of
Native Sheep Provides Insights into Rapid Adaptations to Extreme Environments. Mol. Biol.
Evol. 33: 2576–2592. https://doi.org/10.1093/molbev/msw129

Zeder M. A., 2012 The Domestication of Animals. J. Anthropol. Res. 68: 161–190.
https://doi.org/10.3998/jar.0521004.0068.201

Zeder M. A., 2015 Core questions in domestication research. Proc. Natl. Acad. Sci. 112: 3191–
3198. https://doi.org/10.1073/pnas.1501711112

Zeder M. A., E. Emshwiller, B. D. Smith, and D. G. Bradley, 2006 Documenting
domestication: the intersection of genetics and archaeology. Trends Genet. TIG 22: 139–155.
https://doi.org/10.1016/j.tig.2006.01.007