1 **SARS-CoV-2 Omicron BA.2.75 variant may be much more infective than**
2 **preexisting variants**
3

4 Aki Sugano [a,b], Yutaka Takaoka [b,c,d,e,f,h,] *, Haruyuki Kataguchi [c,d],
5 Minoru Kumaoka [b,d], Mika Ohta [b,c,d,f], Shigemi Kimura [b], Masatake Araki [h,],
6 Yoshitomo Morinaga [g], Yoshihiro Yamamoto [i]
7

[a] Center for Clinical Research, Toyama University Hospital, Toyama, Toyama
930-0194, Japan
[b] Department of Medical Systems, Kobe University Graduate School of
Medicine, Kobe, Hyogo 650-0017, Japan
[c] Department of Computational Drug Design and Mathematical Medicine,
Graduate School of Medicine and Pharmaceutical Sciences, University of
Toyama, Toyama, Toyama 930-0194, Japan
[d] Data Science Center for Medicine and Hospital Management, Toyama
University Hospital, Toyama, Toyama 930-0194, Japan
[e] Center for Advanced Antibody Drug Development, University of Toyama,
Toyama, Toyama 930-0194, Japan
[f] Life Science Institute, Kobe Tokiwa University, Kobe, Hyogo 653-0838, Japan
[g] Department of Microbiology, Toyama University Graduate School of Medicine
and Pharmaceutical Sciences, University of Toyama, Toyama 930-0194, Japan
[h] Division of Genomics, Institute of Resource Development and Analysis,
Kumamoto University, Kumamoto 860-0811, Japan
[i] Department of Clinical Infectious diseases, Toyama University Graduate
School of Medicine and Pharmaceutical Sciences, University of Toyama,
Toyama 930-0194, Japan
27

Running head: BA.2.75 may be much more infective than preexisting variants

Keywords: SARS-CoV-2; COVID-19; Infectivity; Spike protein; Docking affinity;
Mathematical model

*Corresponding author:
Phone: +81-76-434-7311, Fax: +81-76-434-5129
e-mail address: ytakaoka@med.u-toyama.ac.jp (Y. Takaoka)

Abbreviations: spike protein gene, S gene; angiotensin-converting enzyme 2,
ACE2; receptor-binding domain, RBD

ABSTRACT

Objectives: In our previous research, we have reported mathematical model with molecular simulation analysis to predict the infectivity of seven SARS-CoV-2 variants. In this report, we aimed to predict the relative risk of the recent new variants of SARS-CoV-2 based on our previous research.

Methods: We subjected Omicron BA.4/5 or BA.2.75 variant of SARS-CoV-2 to the analysis for the evolutionary distance of the spike protein gene (S gene) to appreciate the changes of the spike protein. We performed the molecular docking simulation analyses of the spike protein with human angiotensin-converting enzyme 2 (ACE2) to understand the docking affinity in these variants. We then compared the evolutionary distance and the docking affinity of these variants with that of the seven variants which were analyzed in our previous research.

Results: The evolutionary distance of the S gene in BA.4/5 or BA.2.75 from the Wuhan variant were longer than the other variants. The highest docking affinity of the spike protein with ACE2 (ratio per Wuhan variant) was observed in BA.2.75.

Conclusion: It is important to note that BA.2.75 has both the highest docking affinity and the longest evolutionary distance of the S gene. These results suggest that the infection of BA.2.75 can be spread further than preexisting variants.

Introduction

In Japan, Omicron BA.5 variants of SARS-CoV-2 is being epidemic as of July in 2022. In addition, Omicron BA.2.75 is discovered and concerned as the coming epidemic. We have previously constructed a mathematical model to predict the infectivity of SARS-CoV-2—Wuhan, Alpha, Beta, Gamma, Delta, Omicron BA.1and BA.2 variants (Takaoka et al., 2022). In this research, we report the predicted risk of Omicron BA.4/5, and BA.2.75 which are recently being epidemic, by using our methodology in the previous report (Takaoka et al., 2022).

Materials and methods

Determination of the evolutionary distance between Wuhan variant and variant S genes, docking affinity of each spike protein with ACE2

We analyzed the evolutionary distance of S gene between the Wuhan variant and Omicron BA.4/5, or BA.2.75 according to our previous method (Takaoka et al., 2022). The sequences of the S gene were obtained by searching EpiCoV database of GISAID for the one with complete sequence in S gene: Accession IDs for Omicron BA.4/5, EPI_ISL_13410397; BA.2.75, EPI_ISL_13438754 respectively (Elbe et al., 2017; Khare et al., 2021).

The docking affinity of the receptor binding domain (RBD) of each spike protein

78 of these variants with ACE2 was analyzed by docking simulation and cluster

79 analysis as in our previous report (Takaoka et al., 2022). The docking affinity is

80 calculated by the following expression:

81
$$\frac{A^{\left(\frac{B}{B_{Wuhan}}\right)}}{A_{Wuhan}}$$

where A is the average score of the most stable cluster and B is the score of the

most stable complex in the docking result.

The amino acid sequences of spike proteins were obtained from CoVariants

website which classifies the variants by Nextstrain clades: 22A (Omicron, BA.4),

22B (Omicron, BA.5) and 22D (Omicron, BA.2.75) (Hadfield et al., 2018; Hodcroft

2021). BA.4 and BA.5 have the identical spike proteins and thus the sequence of

the S gene or amino acid sequence were grouped together in this research.

Results

Evolutionary distances for S gene variants and results of docking of RBD with

ACE2 protein

Table 1 shows the S gene evolutionary distances between the Wuhan variant and

each of the other variants, as well as the docking affinities of RBD of the spike

protein with ACE2 (ratio per Wuhan variant) which are resulted from the docking

simulation. The variants with longer evolutionary distance from the Wuhan variant

had a tendency toward more epidemic. Omicron BA.2.75 variant was observed

the highest docking affinity of the spike protein with ACE2 protein.

In comparison with BA.2, BA.4/5 has longer S gene evolutionary distance but

showed lower docking affinity. Meanwhile, BA.2.75 showed not only longer

evolutionary distance but also much higher docking affinity than that of BA.2.

Discussion

In this report, we analyzed the evolutionary distance of the S gene and the

docking affinity with ACE2 of the recent new variants, BA.4/5 and BA.2.75. We

focused on only the S gene sequence to calculate the evolutionary distance, not

using genome sequence as in the phylogeny provided by Nextstrain (Hadfield et

al., 2018).

We consider that the following two factors play important roles for the infectivity:

the ability of virus to enter into human cells and the effect of neutralizing antibody

in human or the effect of vaccines. The former is shown by the docking affinity of

each RBD in the spike protein with ACE2, which is the higher, the greater. The

latter is shown by the evolutionary distance of the S gene: The longer the distance

is, the weaker the effect would be. It is based on the fact that the currently

available vaccines are developed based on Wuhan variant (Hassine 2022).

However, it should be kept in mind that the risk for the exacerbation of SARS-

CoV-2 cannot be appreciated in this method.

Omicron BA.2.75 showed the highest docking affinity of the spike protein with

ACE2 protein among the other seven variants (ratio per Wuhan). In addition, the

S gene evolutionary distance of BA.2.75 was the longest from the Wuhan variant.

Therefore, BA.2.75 has high ability to enter into human cells and simultaneously

the current vaccines are much less effective to this variant. These results suggest

that the infection of BA.2.75 can be spread further than preexisting variants and

the need for a strong note of caution.

Conclusion

We demonstrated that Omicron BA.2.75 variant of SARS-CoV-2 is very high by

our results of the evolutionary distance of S gene and the docking simulation for

spike protein with ACE2. Our results indicates that Omicron BA.2.75 has a

greater risk to global health than the other ten variants. and there is a need for

paying close attention to the infection trends of Omicron BA.2.75.

Data availability

Data that support the findings of this study are available from the corresponding

author upon reasonable request, except publicly available data source.

Conflict of Interest Disclosures

Authors declare no conflict of interest.

Funding

This work was supported by JSPS Grant-in-Aid for Scientific Research [grant

numbers 21K12110 to Y.T. and 22K12261 to A.S].

Acknowledgements

We gratefully acknowledge all data contributors, i.e., the Authors and their

Originating laboratories responsible for obtaining the specimens, and their

Submitting laboratories for generating the genetic sequence and metadata and

sharing via the GISAID Initiative and all data provided by CoVariants, on which

this research is based.

Author contributions

159 Y.T. conceived and designed this research. Y.T. and A.S. preformed the analyses

160 and acquired the data. Y.T., A.S., and M.O. interpreted the data. Y.T. and A.S.

161 wrote the draft, and all authors reviewed and approved the manuscript.

Ethical approval statement

165 This research is not applicable because we performed computer analyses by

166 using sequence data obtained from public database.

References

References

Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution to

global health. Global Challenges 2017; 1:33-46. DOI: 10.1002/gch2.1018PMCID: 31565258.

Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al. Nextstrain: real-time

tracking of pathogen evolution. Bioinformatics. 2018;34(23):4121-4123. doi:

10.1093/bioinformatics/bty407. Available from: https://nextstrain.org/. [Accessed August 24,

2022].

Hassine IH. Covid-19 vaccines and variants of concern: a review. Rev Med Virol 2022;32(4):

e2313. https://doi.org/10.1002/rmv.2313.

Hodcroft EB. CoVariants: SARS-CoV-2 Mutations and Variants of Interes. 2021. Available from:

https://covariants.org/. [Accessed August 24, 2022].

Khare S, Gurry C, Freitas L, Schultz MB, Bach G, Diallo G, et al. GISAID's Role in Pandemic

Response. China CDC Weekly 2021;3(49): 1049-1051. doi:10.46234/ccdcw2021.255. Available

from: https://gisaid.org/. [Accessed August 24, 2022].

Takaoka Y, Sugano A, Morinaga Y, Ohta, M, Miura K, Kataguchi H, et al. Prediction of infectivity

of SARS-CoV2: Mathematical model with analysis of docking simulation for spike proteins and

angiotensin-converting enzyme 2. Microb Risk Anal 2022.

https://doi.org/10.1016/j.mran.2022.100227.

Table 1 The evolutionary distance of the S gene and the docking affinity of spike protein with ACE2 (ratio per Wuhan variant)

	Wuhan	Alpha	Beta	Gamma	Delta	Omicron BA.1	Omicron BA.2	Omicron BA.4/5	Omicron BA.2.75
Evolutionary Distance of S gene (from Wuhan) x 10^-3	-	4.2	4.5	10.8	12.1	14.3	25.6	41.2	46.8
Docking affinity (Ratio per Wuhan)	1	1.15	1.23	1.32	2.40	3.00	7.91	4.82	21.05