1

Abiotic niche predictors of long-term trends in body mass and survival of Eastern Himalayan birds

Akshay Bharadwaj[1*], Ritobroto Chanda[1], Aman Biswakarma[1], Bharath Tamang[1], Binod Munda[1], Dambar K Pradhan[1], Mangal K Rai[1], Shambu Rai[1], Umesh Srinivasan[1]

1. Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India 560 012

*Email of corresponding author: bakshay@iisc.ac.in

Keywords:
niche shift, selective logging, tropical birds, arthropod community, habitat modification, climate change impacts

Number of words in abstract:
Number of words in main text:
Number of references:
Number of figures, tables and textboxes:

Statement of authorship
AB and UM designed the study. UM collected the mark-recapture dataset from 2011 to 2021 and the temperature-humidity dataset in 2021. RC collected the arthropod community data. AB performed the survival, temperature-humidity and arthropod community analysis. AB wrote the first draft of the manuscript, and UM aided in its revision. UM supervised this work.

Data accessibility statement:
If accepted, all data supporting the results will be archived in an appropriate public repository and the data DOI will be included at the end of the article.

Abstract:

The synergistic impacts of climate change and habitat degradation threaten tropical species worldwide. However, how species' abiotic niches affect their demographic vital rates and phenotypic changes under anthropogenic change remains poorly understood. Using an 11-year mark-recapture dataset from primary and selectively logged forest in the Eastern Himalayas, we investigated how temperature-humidity niche characteristics predicted body mass and survival trends in understorey insectivorous birds over time in each habitat. Our results show that logged forest is hotter and drier than primary forest, and the arthropod community shows dramatic shifts in composition upon selective logging. In understorey insectivores, the degree of dissimilarity between species-specific primary and logged forest niches was strongly and negatively correlated with survival and body mass trends in logged forest. Here, we show that temperature-humidity niche shifts in response to anthropogenic habitat modification can impact demographic vital rates and body condition crucial for population persistence. This work has the potential to inform prompt, targeted conservation efforts toward species that are the most threatened in a warmer and more degraded world.

Introduction:

Climate change and habitat degradation are the two greatest threats to tropical biodiversity today (Pimm, 2008; Pringle et al., 2012). Not only are global temperatures rising rapidly due to climate change, the anthropogenic modification of tropical forest (e.g., by selective logging and conversion to agriculture) further compounds climate impacts; modified habitats reach far higher temperatures than natural forests (Jucker et al., 2020) and lose the buffering effects of the forest canopy (De Frenne et al., 2019). There is also a general consensus on the expected effects of climate change and forest modification on the structure and function of ecological communities — large-bodied, k-selected species and those at higher trophic levels decline in abundance with selective logging and habitat fragmentation (Burivalova et al., 2015; Duffy, 2003; Gray et al., 2007; Hamer et al., 2015; Petchey et al., 1999).

While the impacts of these two factors on biodiversity have received great attention, these impacts have been mostly considered separately, notwithstanding the fact that species are simultaneously exposed to, and therefore respond to, both climate change and habitat degradation concurrently (Srinivasan & Wilcove, 2021). Further, most research on the impacts of climate change and forest loss often examines biodiversity responses in terms of range shifts (Freeman et al., 2018), changes in the population sizes of species (Kim et al., 2022; Şekercioğlu et al., 2012), and the structure of ecological communities (Newmark, 2006). Little is known about the compounding effects of climate change and habitat modification on demographic vital rates such as survival, and phenotypic characteristics, such as body mass (Cosset et al., 2019; Messina et al., 2021; Srinivasan, 2019). Such an understanding, however, remains crucial because: (a) vital rates determine the long-term persistence and viability of populations; (b) studying phenotypic changes over time can help us understand the underlying drivers behind shifts in community structure and vital rates; and (c) quantification of an objective, reliable estimate of a species' vulnerability to extinction — using niche parameters—enables effective use of resources to best conserve biodiversity.

All species occupy distinct ecological niches, defined as the sum of the habitat requirements, environmental conditions and behaviours that enable a species to persist at a location (Jirinec et al., 2021; Ruegg et al., 2021). All else being equal, species with large fundamental abiotic niches, quantified as a function of environmental parameters such as temperature and humidity, should be more adaptable to habitat modifications that alter the abiotic environment (Moritz & Agudo, 2013). In contrast, niche specialists with narrow temperature-humidity niches are likely to be more sensitive to environmental change (Clavel et al., 2011). Temperature and humidity have especially been shown to determine species, and even individual, responses to habitat change and the resulting alteration in climatic variables (Bay et al., 2021; Frishkoff et al., 2016;

Kim et al., 2022; Neate-Clegg et al., 2021). For instance, outright habitat loss from
conversion to agriculture leads to an average 7.6°C increase in temperature and a
reduction in humidity relative to primary forest (Fick & Hijmans, 2017). Even relatively
minor forest degradation (for instance, through selective logging) raised mean
temperatures by 2°C and maximum temperatures by 5.8°C (Srinivasan & Wilcove
2021).

Selective logging is the predominant logging regime in the tropics (Edwards &
Laurance, 2013), and results in logged forest occuring in a continuum of varying tree
densities throughout the world (Burivalova et al., 2015). Although not leading to outright
habitat loss, selective logging leads to habitat degradation and the associated impacts
on plant and animal communities (Halaj et al., 2008; Padmawathe et al., 2004; Velho et
al., 2012). Logging can directly affect species; for instance via the felling of trees leading
to reduced availability of nest spaces and refugia and increased predation risks (Tuff et
al., 2016). Logging can also impact species indirectly through changes in microclimate,
an important determinant of species presence across plant and animal taxa (Boyle et
al., 2021; Zellweger et al., 2020). Microclimatic characteristics and diversity are in turn
linked to the physical structure of the tree community (Han et al., 2022) and
consequently, sensitive to selective logging (Aleixo, 1999; Thiollay, 1992).

Altered thermal environments can also affect phenotypes. A morphological change that
has been postulated to occur under climate warming is the reduction of body size (Watt
et al., 2010). This follows from Bergmann's rule, based on the fact that a higher surface
area:volume ratio translates to better heat loss, leading to the expectation that warming
climates should favour smaller body sizes (Buskirk et al., 2010; Gardner et al., 2011).
The evidence, however, shows mixed support for this 'rule' with some species showing
reductions in body size, while others showed increased sizes with rising temperatures
(Jirinec et al., 2021; Teplitsky & Millien, 2014). This may be because the adaptability of
species to climate change and the determinants of this adaptability are yet to be fully
understood. An understanding of the drivers of adaptability of a species to climate
change and habitat degradation is, therefore, crucial to forecast the vulnerability of
various species present in an ecosystem. Furthermore, the ability of a species to adapt
to altered abiotic conditions via reduction in body size is likely to also be linked to long-
term survival in modified habitats (Møller & Szép, 2002; Rioux Paquette et al., 2014).

We studied year-round resident bird species in the Eastern Himalaya Global
Biodiversity Hotspot for an 11-year period, using mark-recapture in both primary and
logged forest plots. The Eastern Himalayas is among the world's most biodiverse
regions (Grenyer et al., 2006; Pandit et al., 2014), and presently faces the twin threats
of selective logging and climatic warming. Furthermore, Eastern Himalayan bird
species, especially the understorey insectivores, are adapted to narrow thermal ranges

and microhabitats, potentially making them extremely vulnerable to anthropogenic
change (Jetz et al., 2007; Powell et al., 2015; Srinivasan et al., 2018).

In this study, we asked:
(a) Are there abiotic and biotic differences between logged and primary forest
environments?
We expected logged forest to be hotter and drier than primary forest
habitats. As a consequence, we also expected shifts in the biotic
communities of arthropods in selectively logged forests, due to their
thermally-sensitive and ectothermal nature (Renault et al., 2022).
(b) Does the degree of dissimilarity between species' primary and logged
forest niches correlate with their body mass and/or survival trends over
time in logged forest?
We expected that the more dissimilar the primary and logged forest
abiotic niche of a species is, the steeper the body mass and survival
declines of that species in logged forest. In other words, we expect
to see a positive correlation between the degree of overlap between
primary and logged forest niches, and species' body mass and
survival trends over time in logged forest.

Methods
Study area and field sampling
We sampled primary and logged montane broadleaf wet evergreen forest at 2000m
ASL in Eaglenest Wildlife Sanctuary, West Kameng district, Arunachal Pradesh, India.
Six forest patches were selected for sampling based on interviews with persons
involved in logging operations (three each in primary and logged forest; a total
of 9ha in each habitat; plot centroids were separated by an average of 721 ∓
301m SD; Fig S1). Variation in tree densities on these plots tallied exactly with semi-
quantitative estimates of timber extraction (Srinivasan, 2013). On average, tree
densities in primary forest plots (~200 trees per ha) were more than double those in
logged forest (~75 trees per ha).

In each plot, a team operated 24-48 mist nets (12m length, 4 shelf, 16mm mesh size;
158 nets in total) from 0500 to 1200h for three consecutive days during the early
breeding season (April-May) each year, from 2011-2021. Sampling was not possible in
2020 because of a nationwide lockdown caused by the COVID-19 pandemic. Mist-
netting was conducted in three cycles, each spanning three days each in every plot.
Nets were set up systematically at the same locations each year within well-defined
study plots, with neighbouring nets placed ~40m apart (Fig S1). No mist nets were
placed along roads or trails, but a narrow line was cleared in the understorey prior to the

deployment of mist-nets. Because sampling was done during the early breeding
season, all captured small passerine birds were adults and were found in small, distinct
territories very little movement between plots (Srinivasan & Wilcove, 2021). All captured
individual birds were ringed with a uniquely numbered aluminium ring, weighed and
then released.

Differences in overall temperature-humidity profiles in primary and logged forest
To measure the temperature and humidity associated with each capture of an individual
bird, a single DS 1923 iButton Hygrochron (Maxim Integrated) temperature-humidity
logger was placed at the location of each mist net (158 logger locations in total).
Temperature-humidity loggers were programmed to record both temperature and
relative humidity at 10-minute intervals. Loggers were deployed along with the mist nets
throughout the mist-netting conducted in 2021, which was conducted in three cycles
spanning three days each in every plot. We returned to mist net in each plot roughly
after every 15 days, a timeframe in which there are no major changes in climate in the
relatively aseasonal north-eastern part of India (Srinivasan et al., 2018). Because the
order of deployment in the plots was randomised, we do not expect any biases in
microclimate characterisation across habitats. A previous behavioural study which also
measured microclimates from the same plots in an earlier period (March-April) in 2021
showed qualitatively the same temperature-humidity differences between primary and
logged forest (Aggarwal et al., 2023).

Each plot had 25-28 mist nets, and correspondingly, 25-28 temperature humidity
loggers (one deployed at each mist net). Each bird capture was then associated with
the closest temperature and humidity reading in time from the mist net of capture. The
bird species we select for our analyses are insectivorous birds found in the
understorey/midstorey (where our temperature-humidity loggers are deployed). They do
not move up vertically into higher forest strata such as the canopy, thereby allowing an
accurate estimation of their abiotic niche use. Unlike capture efforts (which span 2011 to
2021), temperature-humidity data associated with bird captures in primary and logged
forest was only measured in 2021.

To analyse the temperature and humidity profiles of primary and logged forest, we used
a mixed effects model in the lme4 package (Bates et al., 2015) in R (R. Development
Core Team, 2013) parameterised as follows:

$$\text{Temperature} || \text{Humidity}_i \sim N\left(\alpha_{j[i]}, \sigma^2\right)$$
$$\alpha_j \sim N\left(\gamma_0^\alpha + \gamma_1^\alpha(\text{Habitat}_{primary}), \sigma_{\alpha_j}^2\right), \text{ for Plot } j = 1, \ldots, J$$

where habitat was the fixed variable with two levels (primary and logged) and plot was a
random variable with six levels (three plots each in primary and logged forest). From
these models, we report the differences in temperature and humidities between primary
and logged forest (i.e., the effect size) and 95% confidence intervals.

Differences in arthropod community composition between primary and logged forest
Like all ectotherms, arthropods are sensitive to the temperature and humidity changes
(Burraco et al., 2020) and make up the diets of all our study species (Supriya et al.,
2020). Therefore, to better understand biotic changes accompanying selective logging,
we studied the arthropod community compositions in primary and logged forest.

We used three arthropod sampling techniques at 25-28 points within each plot to
sample the arthropod community in primary and logged forest (55 sampling stations in
primary forest and 53 sampling stations in logged forest). We accounted for ground-
dwelling, flying and understorey foliage arthropods through the use of sticky traps, pitfall
traps and branch-beats.

For ground-dwelling arthropods, we placed pitfall traps containing detergent water for 48
hours at each sampling station. For flying arthropods, we used sticky traps - a
cardboard rectangle 150 x 200mm with an adhesive coating on both sides. These non-
toxic sticky traps contained an aromatic substance to attract insects, and were hung
approximately 1.5m from the ground for 48 hours at each location. Lastly, we used
branch beating for sampling foliage arthropods. For branch beating, we stitched a white
linen cloth in the shape of a funnel around a metal ring 1m in diameter at the larger end
of the funnel and with a container attached to the thinner end of the funnel. At each
station, we placed a randomly selected branch within this funnel, beat it ten times with a
standard-sized stick and sprayed commercially available insecticide spray. This
momentarily paralyzed the arthropods, allowing us to collect them in the container
affixed to the narrow end of the funnel.

All collected arthropod samples were photographed with scale using a Nikon P900
camera. Each arthropod within the sample was identified to the order level.

Selection of bird species for analysis
We selected species suitable for analyses based on the following criteria:
1) Selected species were understorey insectivores that breed on our sampling plots
(migrants were excluded).

2) Selected species had sufficient recapture rates to provide robust estimates of
species survival in both primary and logged forest.

3) Selected species had sufficient temperature-humidity data associated with their
captures to allow the modelling of the abiotic niches.

The list of all selected species is reported in Table S1.

Temperature-Humidity niche analysis
To estimate the temperature-humidity niches of different species we used the
adeHabitat package (Calenge, 2006) in Program R (Core Team 2022), to model the
temperature-humidity niche using a Utilisation Distribution (UD) kernel approach
(Calenge, 2006; Fig 1). While this method was initially developed to analyse the
utilisation of various parts of the home ranges of radio-tracked animals using a Kernel
Density Estimate (KDE), we use it to model the use of the temperature-humidity space
for each species, in primary and logged forest (Fig 1). Through this approach, for each
species, we generated separate temperature-humidity KDEs for primary and logged
forest separately, and subsequently calculated the Utilisation Density Overlap Index
(UDOI) — a standard measure of UD overlap based on the Hurlbert's index of niche
overlap (Hurlbert, 1978; Fig 1). UDOI (henceforth 'niche overlap') is an individual
covariate associated with the capture history of each individual, and is calculated
separately for each species. All individuals of a particular species are therefore
assigned the same UDOI value in the mark-recapture models (see Srinivasan et al.,
2015).

Figure 1: Example of the temperature-humidity niches of Yellow-throated Fulvetta
(*Pseudominla cinerea*) in primary and logged forest, visualised as a Utilisation

Distribution kernel. The abiotic niche for this species is generally more moist and slightly cooler in primary than in logged forest. The intensity of use of a given region in temperature-humidity space is indicated with a gradient of shading, opaque representing region of most use and transparent representing regions of no use.

Body mass analysis

First, we standardised all body masses to a species-wise z-score to account for the fact that species which are larger can potentially lose more absolute body mass than species which are inherently small. We then used a generalised linear model to examine the relationship between body mass, time and niche overlap.

We used generalised linear models to examine how the average body mass of birds of each species changed over time in primary and in logged forest. This was modelled as follows:

Z-score ~ Year * Habitat * Niche_Overlap, where

```
    1) Z-score = Z-score of the body mass (continuous variable)
    2) Year = Year of study (continuous variable)
    3) Habitat = habitat type (categorical variable with two levels; primary
       and logged)
    4) Niche_Overlap = Niche overlap between primary and logged forest niches
       (continuous variable)
```

To visualise our results we created a heatmap whose axes was time and niche overlap, and the color of the grid cells indicated the Z-score predicted by our model. Niche overlap, which is a continuous variable, was split into 10 different equally-spaced values within the naturally occurring range of values.

To visualise the difference in body mass trends between species with highly overlapping and highly disjunct primary and logged forest niches, we plotted the predicted Z-score over time for these species only.

Survival analysis

We estimated the apparent survival (phi) for each species in each type of habitat using the Cormack-Jolly Seber (CJS) mark-recapture model (Cormack, 1964; Jolly, 1965; Seber, 1965), to estimate adult apparent survival in primary and logged forests separately (Fig S1). Apparent survival is an amalgam of true survivorship and site fidelity (i.e., death and emigration are indistinguishable; Barbour et al., 2013). Therefore, an annual apparent survival estimate of 0.60 indicates a 60% probability of an individual in the population surviving *and* remaining in the same location from one year to the next. For each species, to analyse how apparent survival changed over time, we first created a capture history for each individual bird spanning eleven years. Multiple

captures of the same individual within a single year (e.g., on consecutive sampling
days) were not considered recaptures (i.e., all captures within a single sampling season
were collapsed into a single event in the capture histories). We then pooled capture
histories from the three primary forest plots to represent populations sampled in primary
forest and pooled capture histories from all logged plots to represent populations
sampled in logged forest. We pooled the data to estimate apparent survival in primary
and logged forests for each species because plot-level data was insufficient to
adequately and reliably estimate survival for each species in each plot.
284
The CJS model is an open-population model that estimates apparent survival while
accounting for imperfect detection (i.e., capture probability < 1). Using the *R2ucare*
package (Gimenez et al., 2018) in Program R (Core Team, 2013), we first ran tests to
ensure that the CJS models fitted the capture history data well. We ran two models
separately — one in which recapture probability was constant throughout the study
period (across all sampling occasions), and another in which recapture probability was
allowed to vary with sampling occasion. In both models, apparent survival was
constrained to vary unidirectionally over time (but the slope of survival over time was
allowed to be zero). We chose the best model (between time-variant and time-invariant
recapture probabilities) by using the small sample size-corrected Akaike's Information
Criterion (AICc). The best-performing model was used, and it was as follows:

Phi(~Year * Habitat * Niche_Overlap)p(~1), where

1) Phi() = apparent survival (continuous variable between 0 and 1)
2) Year = Year of study (continuous variable)
3) Habitat = habitat type (categorical variable with two levels; primary
300   and logged)
301   4) Niche_overlap = niche overlap between primary and logged forest niches
302   (continuous variable)
303   5) p(~1) = capture probability (remains constant in the best performing
304   model)

Using the estimates from this model, we created a heatmap whose axes was time and
niche overlap, and the color of the grid cells indicated the survival predicted by our
model. Niche overlap, which is a continuous variable, was split into 10 different equally-
spaced values within the naturally occurring range of values.

To visualise the difference in body mass trends between species with highly overlapping
and highly disjunct primary and logged forest niches, we plotted the predicted survival
over time for these species only.

Results

Temperature and humidity in primary and logged forest

Over the entire 24-hr daily period, primary forest was -0.87 degC cooler than logged forest plots (95% CI = [-2.17, 1.97]; marginal R2 = 0.0001; conditional R2 = 0.10). However, while the mean temperature does not differ significantly between the two, it hides differences in the daily temperature regimes in primary and logged forest. Logged forest was more variable than primary forest with lower temperatures at night, warming more rapidly during the day, up to 4.4°C warmer than primary forest at mid-day (mean difference in temperatures between primary and logged forest at 1300hrs = 0.83°C; Fig 2a).

Regardless of the time of the day, on average, logged forest was less humid than primary forest (difference in average humidity between primary and logged forest = 8.73%; 95% CI = [-3.73, 21.20]; marginal R^2 = 0.08; conditional R^2 = 0.39; Fig 2b).

Figure 2: (a) Daily temperature (in ℃) and (b) daily relative humidity (%) trends in primary (green) and logged forest (brown) over a twenty-four hour period. Logged forest has, on average, more variable temperatures and is consistentely and considerably drier than primary forest. Solid lines indicate trendlines from locally estimated scatterplot smoothening (LOESS), and the polygons around the lines represent the 95% confidence interval around the trendline .

Arthropod Community Composition

Primary and logged forest differed significantly in athropod community composition
(NMDS, MoristaHorn index, stress = 0.20; perMANOVA: $F_{1,106}$ = 24.69, p < 0.01, R^2 =
0.82). In general, primary forests harboured more arachnids, lepidopterans and
hemipterans while logged forest insect communities had more flies (Fig 3). The total
abundance of arthropods did not differ significantly between primary and logged forest
(Poisson ANOVA, $z_{1,106}$ = 1.77; p = 0.08), although there were trap-specific differences
observed. Foliage arthropod density was higher in primary forest (branch beats;
ANOVA, $t_{1,106}$ = 4.22; p < 0.01) while density of arthropods in flight was higher in logged
forest (sticky traps; $Z_{1,106}$ = -5.12, p < 0.01; Aggarwal et al., 2023). No difference was
observed between terrestrial arthopod abundances in primary and logged forest (pitfall
traps; $Z_{1,106}$ = 0.33, p = 0.75; Aggarwal et al., 2023).

Figure 3: (a) An NMDS visualisation of the arthropod community composition in
primary and logged forests at our study site. Primary forest plots had more
arachnids, lepidopterans and hemipterans while logged forest insect arthropod
communities have a greater proportion of dipterans; (b) The abundance of
arthropods in both logged and primary forest are not significantly different
(Poisson ANOVA, $z_{1,106}$ = 1.77; p = 0.08). However, there are trap-wise differences
between primary and logged forest arthropod communities, corresponding to the
higher abundance of arachnids and dipterans in primary and logged forest,
respectively (Aggarwal et al., 2023).

Preliminary mist-netting results

During the 11-year study period, we recorded 9,316 captures of 6,340 individuals of 135
species. Based on three selection criteria (breeding, enough data for capture-recapture
analysis and enough data for temperature-humidity niche modelling) we selected 17
species for further analyses. In total, we recorded 5,428 captures of 3,519 individuals
belonging to these 17 species.

Body mass trends
Our analysis suggests a significant effect of climate and land-use change on the body
mass trends of birds over time. In logged forest, body mass trends over time showed a
positive relationship with niche overlap ($\beta_{time*niche\_overlap}$ = 0.03, 95%CI = [-0.01, 0.08];
$\beta_{time*niche\_overlap*habitat\_primary}$ = -0.04, 95%CI = [-1.10, 0.02]; Fig 4a-b). Species with greater
niche dissimilarities between their primary and logged forest niches also had steeper
survival declines over time in a logged forest (Fig 4b). The model parameters for the
estimates and confidence intervals are reported in Table S2.

Figure 4: (*Top row*) A heat map showing the relationship between body mass, time and niche overlap in (a) primary and (b) logged forest. Cells with darker hues of blue indicate lower survival, while those with hues of green have higher

**survival; (*bottom row*) The relationship between survival and time for species
with high (blue) and low (orange) niche overlap between their primary forest and
logged forest niches.**

Survival analysis

Individuals that were captured on more than one habitat type (primary or logged) were
excluded from further analyses (3.9% of all captured individuals). Models with constant
recapture probability performed better for all species, in comparison to one with time-
varying capture probability. The model parameters for the estimates and confidence
intervals are reported in Table S3.

Niche dissimilarity (i.e., the degree to which a species' temperature-humidity niche
changes with logging) is strongly related to survival trends in logged forest ($\beta_{time*niche\_overlap}$
= 0.23, 95%CI = [0.001, 0.47]; $\beta_{time*niche\_overlap*habitat\_primary}$ = -0.37, 95%CI = [-0.70, -0.03];
Fig 5a-b). Species with greater niche dissimilarities between their primary and logged
forest niches also had steeper survival declines over time in a logged forest (Fig 5b).

Figure 5: (*Top row*) A heat map showing the relationship between survival, time and niche overlap in (a) primary and (b) logged forest. Cells with darker hues of blue indicate lower survival, while those with hues of green have higher survival;

(*bottom row*) **The relationship between survival and time for species with high**
(blue) and low (orange) niche overlap between their primary forest and logged
forest niches.

Discussion
We examined differences in the abiotic and biotic environments for 17 species of
understory insectivorous birds in primary and selectively logged forest in the Eastern
Himalaya. We analysed how the temperature-humidity niche sizes of understorey bird
species were related to body mass and survival trends over an 11-year period in
primary and logged forest. Primary and logged forest had different temperature-humidity
profiles and differed in the composition of arthropod communities. Importantly, less than
four percent of all captured birds were captured in two different plots, indicative of their
small territory sizes and site fidelity during the breeding season, irrespective of the
habitat type that the individual is found in (Ibarra-Macias et al., 2011; Srinivasan &
Wilcove, 2021).

As hypothesized, we found a strong positive relationship between the degree of niche
dissimilarity (i.e., the degree to which a species' temperature-humidity niche changes
with logging) and survival rates and body mass trends over time. In other words,
species with very different temperature-humidity niches in primary and logged forest
experienced survival and body mass declines in logged forest, while those with high
degree of niche overlap between primary and logged forest have near-constant or
increasing survival and body mass over time (Fig 4, Fig 5), even after logging.

Body size is an important and widely-studied life-history trait that greatly influences
thermoregulation by individuals/species (James, 1970). Habitat-specific differences in a
species' body size (specifically, body mass) can be the result of two possibly concurrent
processes: (a) the consistent reduction in body mass of each individual of the species
and/or (b) the accumulation of lower quality (in this case, lighter) individuals in logged
forest.

Heat dissipation theory postulates that tropical animals are limited energetically by the
rate at which body heat is lost (Speakman & Król, 2010), and purely based on
thermodynamic considerations, smaller body sizes should be selected for in
progressively warmer conditions (Prokosch et al., 2019). The Eastern Himalaya has a
relatively aseasonal climate which selects for thermal niche specialist species
(Srinivasan et al., 2018). Not only is the Eastern Himalaya warming three times as fast
as the global average (Shrestha et al., 2012), but forest degradation is concomitantly
removing the buffering effects of the primary forest canopy, making logged forests more
thermally variable, and significantly hotter and drier than primary forest (De Frenne et
al., 2019; Figs 2a-b). Indeed, our *a priori* expectation was that thermally-stressed

specialist species would undergo steeper declines in body mass over time, in keeping
with the thermoregulatory capacity hypothesis. In other words, species with more
different primary and logged forest niches would undergo steeper declines in body mass
over the study period. Indeed, this is what we observe - body mass over time is
negatively correlated with niche overlap (Fig 4a).

Several bird species elsewhere are indeed undergoing other morphological changes
over time, likely because of climate-driven thermal constraints (Jirinec et al., 2021;
Messina et al., 2021). Specifically, birds can undergo changes in the morphometry of
organs and appendages that serve thermoregulatory (Tattersall et al., 2017) and
locomotory functions (Weeks et al., 2020), without drastically changing overall body
mass. In the rainforests of the Amazon, for example, climate change appears to have
selected for progressively longer wings in comparison to body mass, which suggests
that birds are adapting toward relatively longer appendages to offset thermal stress
(Jirinec et al., 2021). Therefore, in addition to progressively lighter body mass in logged
forests, birds within primary and logged forests would likely exhibit differences in their
mass:wing ratio, a hypothesis that remains to be tested.

In addition to thermal stressors, resource availability also plays a role in determining
body size (McNab, 2010). Since we were interested in insectivorous understorey birds,
we also measured biotic factors determining body sizes such as arthropod composition
and abundance (Cox et al., 2019). Arthropod community composition showed a
considerable shift with selective logging. This result highlights that a possible
confounding variable hitherto unexplored at our study site is the dietary niche breadth of
birds (Bond & Lavers, 2014; Hansen et al., 2019). An assessment of dietary niche
space (i.e., its size and shift in response to selective logging) would help in better
understanding the various mechanisms underlying body mass declines with logging
(Edwards et al., 2013; Hamer et al., 2015; Messina et al., 2021). Furthermore, we
investigate only the abiotic changes brought about by selective logging. Investigating
the other impacts of logging, such as an altered availability of nest spaces and predator-
prey interactions will shed more light on the mechanisms impacting species over time
(Tuff et al., 2016).

We also acknowledge that the trends we observe in body mass declines over time in
the logged forest can occur through the movement of lighter, "low-quality" individuals to
the logged forest plots (Calsbeek & Sinervo, 2002). Such a mechanism, arising from a
despotic distribution, would be expected to result in a situation where the body masses
of individuals (within a species) in logged forest would, on average, be lower than for
individuals in primary forest (Andren, 1990). In the absence of long-term processes
such as climate change, body mass differences between primary and logged forest
would be expected to be stable. However, we report consistent long-term declines in

body mass of in logged forest of species that show little niche overlap between the two
habitats, likely as a result of warming-induced morhphological change (Jirinec et al.,
2021).

The characteristics of a species' niche determines how it interacts with the larger
ecosystem and shapes its individual and population-level responses to ecological
disturbances (Bellwood et al., 2019; Neate-Clegg et al., 2021; Voigt et al., 2007).
Previous studies have suggested that species with larger niches (i.e., generalists) can
inhabit a greater range of environments by drawing from a wider pool of resources. This
makes them more resilient to climate change and less likely to face population declines
in comparison to species with smaller niches (Clavel et al., 2011; or, specialists; Swihart
et al., 2003). Specialist species use a narrow range of resources and/or habitats, and
changes in the availability of both can have drastic effects on their populations since
they are unable to exploit alternative habitats or resources (Graham et al., 2011).

Studies investigating the effect of niche characterstics on sensitivity to climate change
have thus far used occupancy data and theoretical models to estimate niche sizes
(Loiselle et al., 2010; Rinnan & Lawler, 2019). The direct impact of climatic niche
parameters on the long-term survival of species has seldom been tested in the field.
Previous studies on tropical rainforest species show that vulnerability is strongly
predicted by ecological specialisation (Sekercioglu, 2011), and that understory
insectivores are in particular peril due to anthropogenic change because of their limited
dispersal ability (Moore et al., 2008; Powell et al., 2015) and sensitivity to the opening
up of the canopy (Patten & Smith-Patten, 2012; Pollock et al., 2015).

We found that the degree of niche dissimilarity between primary and logged forests was
strongly correlated with survival trends over time — the more dissimilar a species'
primary and logged forest niches are, the steeper the survival declines in logged forest.
A recent paper from the same study system and study species found that the
elevational ranges of species (and potential upslope movement; Girish & Srinivasan,
2022) explained changes in survival rates of species in primary but not in logged forest
(Srinivasan & Wilcove, 2021). In logged forest therefore, rather than elevational ranges
and range shifts, mismatches in the abiotic niche between primary and logged forest
might better explain trends in survival. Although a reduction in the apparent survival of a
species is possible through emigration alone, it is unlikely that the trends we observe
arise from the increased emigration of individuals from the logged forest (see Srinivasan
& Wilcove, 2021).

Such a relationship can occur in two ways. Firstly, an altered abiotic environment can
interact directly with species' physiology, especially on stress response mechanisms. It
is likely that birds with more dissimilar temperature-humidity niche spaces (between

primary and logged forest) face elevated levels of stress and, therefore, chronically
higher levels of stress hormones over time (Wingfield et al., 2017). Heat-shock proteins
(HSPs), which indicate long-term thermal stress, can be used to study this effect (Maak
et al., 2003; Pusch et al., 2018). HSPs are also well suited for a mark-recapture
methodology because other stress indicator hormones are shorter-lived and capture
itself can lead to biased readings arising from temporary capture-related stress. Apart
from HSPs, glucocorticoids (corticosterone) are hormones produced in response to
stress in birds and can induce a wide range of metabolic cascades, such as the
utilisation of stored nutrients for usage in physiological functions, changes in immunity
and the prevention of oxidative damage (Marasco et al., 2017). A meta-analysis of avian
response to logging revealed that species also have lowered immunity in logged forests
in comparison with old-growth forests (Messina et al., 2018), and were likely to carry
higher disease loads. Furthermore, glucocorticoid levels have been shown to have
carry-over effects on population dynamics in many bird species (W. K. Hansen et al.,
2016; Harms et al., 2015; Messina et al., 2020).

A change in abiotic conditions can also have indirect impacts on bird species; for
instance, changes in temperature and humidity can impact the abundances and
diversity of arthropod prey, which we demonstrate here. While one particular study of
altered prey abundances of understory insectivores yielded little support for the prey-
limitation hypothesis (Sekercioglu, 2011), the same species of birds often forage on
arthropods at different trophic levels in primary versus logged forest (Hamer et al.
2015). The method of prey capture has been shown to be a consistent predictor of
sensitivity to disturbance; for instance, sallying insectivores —those that capture flying
prey while themselves in flight— were more sensitive to logging in African forests
(Arcilla et al., 2015).

Our results suggest that the degree of niche dissimilarity is a strong predictor of survival
declines of Eastern Himalayan birds in degraded habitats facing climate change. By
using temperature and humidity loggers, this study demonstrates the applicability of
niche estimation and comparison to reliably estimate the vulnerability of species to
anthropogenic change. The IUCN Red List, widely used to set species-level
conservation priorities, has been criticised for not accounting for climate-related threats
to species (Thuiller et al., 2005). More generally, formal assessments of slow-acting
threats such as climate change is a necessary but challenging step to kickstarting
conservation action. Most often, however, by the time existing criteria such as
population or range-size declines are observable at population levels, it may well be too
late for effective conservation action (Hannah, 2012).

For effective conservation, it is essential that vulnerable species are identified as early
as possible to maximise the chances of conservation success (Stanton et al., 2015). To

this end, Climatic Niche Factor Analysis (CNFA) has been suggested as a tool to predict
the differential vulnerability of various species to extinction (Rinnan & Lawler, 2019).
However, the real-world applicability of niche modelling approaches to assess risk,
especially in terms of demographic rates, is hitherto unexamined.

Our methodology of estimating temperature-humidity niche shifts that correlate with
long-term survival declines in Eastern Himalayan bird species is one that potentially has
widespread use in conservation management and planning. By providing managers and
scientists with an inexpensive quantification method of the relative vulnerabilities of
various species found in bird communities, such CNFA methodologies would enable
targeted, prompt and judicious use of limited conservation resources for the species
which are at the greatest risk of extinction due to global change drivers. Further
research is needed to understand the physiological impacts of climate change and
forest degradation/loss on tropical bird species. The integration of dietary and climatic
requirements of species would also yield a more holistic understanding of the grave
threat of anthropogenic change on tropical biodiversity worldwide.

Acknowledgements:
We thank S. Rai, B. Tamang, D.K. Pradhan, and M. Rai for their enthusiasm and
professionalism in the field. We are grateful to the Arunachal Pradesh Forest
Department for providing permission to conduct this study, and to Mr. Millo Tasser
(IFS), Divisional Forest Officer of Shergaon Forest Division, for his continued support of
our work. I. Glow, N. Tsering Monpa, and A. Pradhan helped greatly with logistics.
Funding for this study came from the National Centre for Biological Sciences,
International Foundation for Science, Princeton University, Princeton University's Global
Collaborative Network on Himalayan Biodiversity, and the High Meadows Foundation.

References:

Aggarwal, K., Chanda, R., Rai, S., Rai, M., Pradhan, D. K., Munda, B., Tamang, B.,

Biswakarma, A., & Srinivasan, U. (2023). The effect of selective logging on

microclimates, arthropod abundance and the foraging behaviour of Eastern Himalayan

birds. *Forest Ecology and Management*, *541*, 121076.

https://doi.org/10.1016/j.foreco.2023.121076

Aleixo, A. (1999). Effects of selective logging on a bird community in the Brazilian Atlantic

Forest. *The Condor*, *101*(3), 537–548.

Andren, H. (1990). Despotic Distribution, Unequal Reproductive Success, and Population

Regulation in the Jay Garrulus Glandarius L. *Ecology*, *71*(5), 1796–1803.

https://doi.org/10.2307/1937587

Arcilla, N., Holbech, L. H., & O'Donnell, S. (2015). Severe declines of understory birds follow

illegal logging in Upper Guinea forests of Ghana, West Africa. *Biological Conservation*,

188, 41–49.

Barbour, A. B., Ponciano, J. M., & Lorenzen, K. (2013). Apparent survival estimation from

continuous mark–recapture/resighting data. *Methods in Ecology and Evolution*, *4*(9),

846–853. https://doi.org/10.1111/2041-210X.12059

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, *67*, 1–48.

https://doi.org/10.18637/jss.v067.i01

Bay, R. A., Karp, D. S., Saracco, J. F., Anderegg, W. R. L., Frishkoff, L. O., Wiedenfeld, D.,

Smith, T. B., & Ruegg, K. (2021). Genetic variation reveals individual-level climate

tracking across the annual cycle of a migratory bird. *Ecology Letters*, *24*(4), 819–828.

https://doi.org/10.1111/ele.13706

Bellwood, D. R., Streit, R. P., Brandl, S. J., & Tebbett, S. B. (2019). The meaning of the term

'function'in ecology: A coral reef perspective. *Functional Ecology*, *33*(6), 948–961.

Billerman, S. M., Keeney, B. K., Rodewald, P. G., & Schulenberg, T. S. (Eds.). (2022). *Birds of

the World*. https://birdsoftheworld.org/bow/home

Bond, A. L., & Lavers, J. L. (2014). Climate change alters the trophic niche of a declining apex

marine predator. *Global Change Biology*, *20*(7), 2100–2107.

https://doi.org/10.1111/gcb.12554

Boyle, M. J., Bishop, T. R., Luke, S. H., Van Breugel, M., Evans, T. A., Pfeifer, M., Fayle, T. M.,

Hardwick, S. R., Lane-Shaw, R. I., & Yusah, K. M. (2021). Localised climate change

defines ant communities in human-modified tropical landscapes. *Functional Ecology*,

35(5), 1094–1108.

Burivalova, Z., Lee, T. M., Giam, X., Şekercioğlu, Ç. H., Wilcove, D. S., & Koh, L. P. (2015).

Avian responses to selective logging shaped by species traits and logging practices.

Proceedings of the Royal Society B: Biological Sciences, *282*(1808), 20150164.

https://doi.org/10.1098/rspb.2015.0164

Burraco, P., Orizaola, G., Monaghan, P., & Metcalfe, N. B. (2020). Climate change and ageing

in ectotherms. *Global Change Biology*, *26*(10), 5371–5381.

Buskirk, J. V., Mulvihill, R. S., & Leberman, R. C. (2010). Declining body sizes in North

American birds associated with climate change. *Oikos*, *119*(6), 1047–1055.

https://doi.org/10.1111/j.1600-0706.2009.18349.x

Calenge, C. (2006). The package "adehabitat" for the R software: A tool for the analysis of

space and habitat use by animals. *Ecological Modelling*, *197*(3–4), 516–519.

Calsbeek, R., & Sinervo, B. (2002). An Experimental Test of the Ideal Despotic Distribution.

Journal of Animal Ecology, *71*(3), 513–523.

Clavel, J., Julliard, R., & Devictor, V. (2011). Worldwide decline of specialist species: Toward a

global functional homogenization? *Frontiers in Ecology and the Environment*, *9*(4), 222–

228. https://doi.org/10.1890/080216

Core Team, R. (2013). R: A language and environment for statistical computing. *R Foundation*

for Statistical Computing, Vienna.

Cormack, R. M. (1964). Estimates of survival from the sighting of marked animals. *Biometrika*,

51(3/4), 429–438.

Cosset, C. C. P., Gilroy, J. J., & Edwards, D. P. (2019). Impacts of tropical forest disturbance on

species vital rates. *Conservation Biology*, *33*(1), 66–75.

https://doi.org/10.1111/cobi.13182

Cox, A. R., Robertson, R. J., Lendvai, Á. Z., Everitt, K., & Bonier, F. (2019). Rainy springs linked

to poor nestling growth in a declining avian aerial insectivore (Tachycineta bicolor).

Proceedings of the Royal Society B: Biological Sciences, *286*(1898), 20190018.

https://doi.org/10.1098/rspb.2019.0018

De Frenne, P., Zellweger, F., Rodríguez-Sánchez, F., Scheffers, B. R., Hylander, K., Luoto, M.,

Vellend, M., Verheyen, K., & Lenoir, J. (2019). Global buffering of temperatures under

forest canopies. *Nature Ecology & Evolution*, *3*(5), 744–749.

https://doi.org/10.1038/s41559-019-0842-1

Duffy, J. E. (2003). Biodiversity loss, trophic skew and ecosystem functioning. *Ecology Letters*,

6(8), 680–687. https://doi.org/10.1046/j.1461-0248.2003.00494.x

Edwards, D. P., & Laurance, W. F. (2013). Biodiversity despite selective logging. *Science*,

339(6120), 646–647.

Edwards, D. P., Woodcock, P., Newton, R. J., Edwards, F. A., Andrews, D. J. R., Docherty, T.

D. S., Mitchell, S. L., Ota, T., Benedick, S., Bottrell, S. H., & Hamer, K. C. (2013).

Trophic Flexibility and the Persistence of Understory Birds in Intensively Logged

Rainforest. *Conservation Biology*, *27*(5), 1079–1086. https://doi.org/10.1111/cobi.12059

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315.

https://doi.org/10.1002/joc.5086

Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018). Climate change

causes upslope shifts and mountaintop extirpations in a tropical bird community.

Proceedings of the National Academy of Sciences, *115*(47), 11982–11987.

Frishkoff, L. O., Karp, D. S., Flanders, J. R., Zook, J., Hadly, E. A., Daily, G. C., & M'Gonigle, L.

K. (2016). Climate change and habitat conversion favour the same species. *Ecology*

Letters, *19*(9), 1081–1090. https://doi.org/10.1111/ele.12645

Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body

size: A third universal response to warming? *Trends in Ecology & Evolution*, *26*(6), 285–

291.

Gimenez, O., Lebreton, J.-D., Choquet, R., & Pradel, R. (2018). R2ucare: An r package to

perform goodness-of-fit tests for capture–recapture models. *Methods in Ecology and*

Evolution, *9*(7), 1749–1754. https://doi.org/10.1111/2041-210X.13014

Girish, K. S., & Srinivasan, U. (2022). Community science data provide evidence for upward

elevational range shifts by Eastern Himalayan birds. *Biotropica*.

Graham, N. A., Chabanet, P., Evans, R. D., Jennings, S., Letourneur, Y., Aaron MacNeil, M.,

McClanahan, T. R., Öhman, M. C., Polunin, N. V., & Wilson, S. K. (2011). Extinction

vulnerability of coral reef fishes. *Ecology Letters*, *14*(4), 341–348.

Gray, M. A., Baldauf, S. L., Mayhew, P. J., & Hill, J. K. (2007). The Response of Avian Feeding

Guilds to Tropical Forest Disturbance. *Conservation Biology*, *21*(1), 133–141.

https://doi.org/10.1111/j.1523-1739.2006.00557.x

Grenyer, R., Orme, C. D. L., Jackson, S. F., Thomas, G. H., Davies, R. G., Davies, T. J., Jones,

K. E., Olson, V. A., Ridgely, R. S., Rasmussen, P. C., Ding, T.-S., Bennett, P. M.,

Blackburn, T. M., Gaston, K. J., Gittleman, J. L., & Owens, I. P. F. (2006). Global

distribution and conservation of rare and threatened vertebrates. *Nature*, *444*(7115),

Article 7115. https://doi.org/10.1038/nature05237

Halaj, J., Halpern, C. B., & Yi, H. (2008). Responses of litter-dwelling spiders and carabid

beetles to varying levels and patterns of green-tree retention. *Forest Ecology and*

Management, *255*(3–4), 887–900.

Hamer, K. C., Newton, R. J., Edwards, F. A., Benedick, S., Bottrell, S. H., & Edwards, D. P.

(2015). Impacts of selective logging on insectivorous birds in Borneo: The importance of

trophic position, body size and foraging height. *Biological Conservation*, *188*, 82–88.

https://doi.org/10.1016/j.biocon.2014.09.026

Han, D., Wang, C., Sun, Z., She, J., Yin, L., Bian, Q., & Han, W. (2022). Microhabitat

preferences of butterflies in urban parks: Both vegetation structure and resources are

decisive. *Urban Forestry & Urban Greening*, *71*, 127552.

Hannah, L. (2012). Are a million species at risk? In *Saving a Million Species* (pp. 3–9). Springer.

Hansen, B. B., Lorentzen, J. R., Welker, J. M., Varpe, Ø., Aanes, R., Beumer, L. T., &

Pedersen, Å. Ø. (2019). Reindeer turning maritime: Ice-locked tundra triggers changes

in dietary niche utilization. *Ecosphere*, *10*(4), e02672. https://doi.org/10.1002/ecs2.2672

Hansen, W. K., Bate, L. J., Landry, D. W., Chastel, O., Parenteau, C., & Breuner, C. W. (2016).

Feather and faecal corticosterone concentrations predict future reproductive decisions in

harlequin ducks (Histrionicus histrionicus). *Conservation Physiology*, *4*(1), cow015.

Harms, N. J., Legagneux, P., Gilchrist, H. G., Bêty, J., Love, O. P., Forbes, M. R., Bortolotti, G.

R., & Soos, C. (2015). Feather corticosterone reveals effect of moulting conditions in the

autumn on subsequent reproductive output and survival in an Arctic migratory bird.

Proceedings of the Royal Society B: Biological Sciences, *282*(1800), 20142085.

Hurlbert, S. (1978). *The Measurement of Niche Overlap and Some Relatives*.

https://doi.org/10.2307/1936632

Ibarra-Macias, A., Robinson, W. D., & Gaines, M. S. (2011). Experimental evaluation of bird

movements in a fragmented Neotropical landscape. *Biological Conservation*, *144*(2),

703–712. https://doi.org/10.1016/j.biocon.2010.08.006

James, F. C. (1970). Geographic Size Variation in Birds and Its Relationship to Climate.

Ecology, *51*(3), 365–390. https://doi.org/10.2307/1935374

Jetz, W., Wilcove, D. S., & Dobson, A. P. (2007). Projected Impacts of Climate and Land-Use

Change on the Global Diversity of Birds. *PLOS Biology*, *5*(6), e157.

https://doi.org/10.1371/journal.pbio.0050157

Jirinec, V., Burner, R. C., Amaral, B. R., BierregaardJr, R. O., Fernández-Arellano, G.,

Hernández-Palma, A., Johnson, E. I., Lovejoy, T. E., Powell, L. L., Rutt, C. L., Wolfe, J.

D., & Stouffer, P. C. (2021). Morphological consequences of climate change for resident

birds in intact Amazonian rainforest. *Science Advances*.

https://doi.org/10.1126/sciadv.abk1743

Jolly, G. M. (1965). Explicit Estimates from Capture-Recapture Data with Both Death and

Immigration-Stochastic Model. *Biometrika*, *52*(1/2), 225–247.

https://doi.org/10.2307/2333826

Jucker, T., Jackson, T. D., Zellweger, F., Swinfield, T., Gregory, N., Williamson, J., Slade, E. M.,

Phillips, J. W., Bittencourt, P. R. L., Blonder, B., Boyle, M. J. W., Ellwood, M. D. F.,

Hemprich-Bennett, D., Lewis, O. T., Matula, R., Senior, R. A., Shenkin, A., Svátek, M., &

Coomes, D. A. (2020). A Research Agenda for Microclimate Ecology in Human-Modified

Tropical Forests. *Frontiers in Forests and Global Change*, *2*.

https://www.frontiersin.org/article/10.3389/ffgc.2019.00092

Kim, H., McComb, B. C., Frey, S. J. K., Bell, D. M., & Betts, M. G. (2022). Forest microclimate

and composition mediate long-term trends of breeding bird populations. *Global Change*

Biology. https://doi.org/10.1111/gcb.16353

Loiselle, B. A., Graham, C. H., Goerck, J. M., & Ribeiro, M. C. (2010). Assessing the impact of

deforestation and climate change on the range size and environmental niche of bird

species in the Atlantic forests, Brazil. *Journal of Biogeography*, *37*(7), 1288–1301.

https://doi.org/10.1111/j.1365-2699.2010.02285.x

Maak, S., Melesse, A., Schmidt, R., Schneider, F., & Von Lengerken, G. (2003). Effect of long-

term heat exposure on peripheral concentrations of heat shock protein 70 (Hsp70) and

hormones in laying hens with different genotypes. *British Poultry Science*, *44*(1), 133–

138. https://doi.org/10.1080/0007166031000085319

Marasco, V., Stier, A., Boner, W., Griffiths, K., Heidinger, B., & Monaghan, P. (2017).

Environmental conditions can modulate the links among oxidative stress, age, and

longevity. *Mechanisms of Ageing and Development*, *164*, 100–107.

https://doi.org/10.1016/j.mad.2017.04.012

McNab, B. K. (2010). Geographic and temporal correlations of mammalian size reconsidered: A

resource rule. *Oecologia*, *164*(1), 13–23.

Messina, S., Costantini, D., Tomassi, S., Cosset, C. C. P., Benedick, S., Eens, M., & Edwards,

D. P. (2021). Selective logging reduces body size in omnivorous and frugivorous tropical

forest birds. *Biological Conservation*, *256*, 109036.

https://doi.org/10.1016/j.biocon.2021.109036

Messina, S., Edwards, D. P., Eens, M., & Costantini, D. (2018). Physiological and

immunological responses of birds and mammals to forest degradation: A meta-analysis.

Biological Conservation, *224*, 223–229. https://doi.org/10.1016/j.biocon.2018.06.002

Messina, S., Edwards, D. P., Marasco, V., Canoine, V., Cosset, C. C. P., Tomassi, S.,

Benedick, S., Eens, M., & Costantini, D. (2020). Glucocorticoids link forest type to local

abundance in tropical birds. *Functional Ecology*, *34*(9), 1814–1825.

https://doi.org/10.1111/1365-2435.13586

Møller, A., & Szép, T. (2002). Survival rate of adult barn swallows Hirundo rustica in relation to

sexual selection and reproduction. *Ecology*, *83*(8), 2220–2228.

Moore, R. P., Robinson, W. D., Lovette, I. J., & Robinson, T. R. (2008). Experimental evidence

for extreme dispersal limitation in tropical forest birds. *Ecology Letters*, *11*(9), 960–968.

Moritz, C., & Agudo, R. (2013). The future of species under climate change: Resilience or

decline? *Science*, *341*(6145), 504–508.

Neate-Clegg, M. H., Stanley, T. R., Şekercioğlu, Ç. H., & Newmark, W. D. (2021). Temperature-

associated decreases in demographic rates of Afrotropical bird species over 30 years.

Global Change Biology, *27*(10), 2254–2268.

Newmark, W. D. (2006). A 16-Year Study of Forest Disturbance and Understory Bird

Community Structure and Composition in Tanzania. *Conservation Biology*, *20*(1), 122–

134. https://doi.org/10.1111/j.1523-1739.2005.00273.x

Padmawathe, R., Qureshi, Q., & Rawat, G. S. (2004). Effects of selective logging on vascular

epiphyte diversity in a moist lowland forest of Eastern Himalaya, India. *Biological*

Conservation, *119*(1), 81–92.

Pandit, M. K., Manish, K., & Koh, L. P. (2014). Dancing on the roof of the world: Ecological

transformation of the Himalayan landscape. *BioScience*, *64*(11), 980–992.

Patten, M. A., & Smith-Patten, B. D. (2012). Testing the microclimate hypothesis: Light

environment and population trends of Neotropical birds. *Biological Conservation*, *155*,

85–93. https://doi.org/10.1016/j.biocon.2012.06.004

Petchey, O. L., McPhearson, P. T., Casey, T. M., & Morin, P. J. (1999). Environmental warming

alters food-web structure and ecosystem function. *Nature*, *402*(6757), Article 6757.

https://doi.org/10.1038/47023

Pimm, S. L. (2008). Biodiversity: Climate change or habitat loss—Which will kill more species?

Current Biology, *18*(3), R117–R119.

Pollock, H. S., Cheviron, Z. A., Agin, T. J., & Brawn, J. D. (2015). Absence of microclimate

selectivity in insectivorous birds of the Neotropical forest understory. *Biological

Conservation*, *188*, 116–125. https://doi.org/10.1016/j.biocon.2014.11.013

Powell, L. L., Cordeiro, N. J., & Stratford, J. A. (2015). Ecology and conservation of avian

insectivores of the rainforest understory: A pantropical perspective. *Biological

Conservation*, *188*, 1–10. https://doi.org/10.1016/j.biocon.2015.03.025

Pringle, C. S., Martin, T. G., & Rhodes, J. R. (2012). Interactions between climate and habitat

loss effects on biodiversity: A systematic review and meta-analysis. *Global Change

Biology*, *18*, 1239–1252.

Prokosch, J., Bernitz, Z., Bernitz, H., Erni, B., & Altwegg, R. (2019). Are animals shrinking due

to climate change? Temperature-mediated selection on body mass in mountain wagtails.

Oecologia, *189*(3), 841–849. https://doi.org/10.1007/s00442-019-04368-2

Pusch, E. A., Bentz, A. B., Becker, D. J., & Navara, K. J. (2018). Behavioral phenotype predicts

physiological responses to chronic stress in proactive and reactive birds. *General and

Comparative Endocrinology*, *255*, 71–77.

R. Development Core Team. (2013). *R: A language and environment for statistical computing*.

Renault, D., Leclerc, C., Colleu, M.-A., Boutet, A., Hotte, H., Colinet, H., Chown, S. L., &

Convey, P. (2022). The rising threat of climate change for arthropods from Earth's cold

regions: Taxonomic rather than native status drives species sensitivity. *Global Change*

Biology, *28*(20), 5914–5927.

Rinnan, D. S., & Lawler, J. (2019). Climate-niche factor analysis: A spatial approach to

quantifying species vulnerability to climate change. *Ecography*, *42*(9), 1494–1503.

https://doi.org/10.1111/ecog.03937

Rioux Paquette, S., Pelletier, F., Garant, D., & Bélisle, M. (2014). Severe recent decrease of

adult body mass in a declining insectivorous bird population. *Proceedings of the Royal*

Society B: Biological Sciences, *281*(1786), 20140649.

https://doi.org/10.1098/rspb.2014.0649

Ruegg, K., Anderson, E. C., Somveille, M., Bay, R. A., Whitfield, M., Paxton, E. H., & Smith, T.

B. (2021). Linking climate niches across seasons to assess population vulnerability in a

migratory bird. *Global Change Biology*, *27*(15), 3519–3531.

https://doi.org/10.1111/gcb.15639

Seber, G. A. (1965). A note on the multiple-recapture census. *Biometrika*, *52*(1/2), 249–259.

Sekercioglu, C. H. (2011). Functional extinctions of bird pollinators cause plant declines.

Science, *331*(6020), 1019–1020.

Şekercioğlu, Ç. H., Primack, R. B., & Wormworth, J. (2012). The effects of climate change on

tropical birds. *Biological Conservation*, *148*(1), 1–18.

https://doi.org/10.1016/j.biocon.2011.10.019

Shrestha, U. B., Gautam, S., & Bawa, K. S. (2012). Widespread Climate Change in the

Himalayas and Associated Changes in Local Ecosystems. *PLOS ONE*, *7*(5), e36741.

https://doi.org/10.1371/journal.pone.0036741

Speakman, J. R., & Król, E. (2010). Maximal heat dissipation capacity and hyperthermia risk:

Neglected key factors in the ecology of endotherms. *Journal of Animal Ecology*, *79*(4),

726–746.

Srinivasan, U. (2013). A slippery slope: Logging alters mass–abundance scaling in ecological

communities. *Journal of Applied Ecology*, *50*(4), 920–928. https://doi.org/10.1111/1365-

2664.12123

Srinivasan, U. (2019). Morphological and Behavioral Correlates of Long-Term Bird Survival in

Selectively Logged Forest. *Frontiers in Ecology and Evolution*, *7*.

https://www.frontiersin.org/article/10.3389/fevo.2019.00017

Srinivasan, U., Elsen, P. R., Tingley, M. W., & Wilcove, D. S. (2018). Temperature and

competition interact to structure Himalayan bird communities. *Proceedings of the Royal*

Society B: Biological Sciences, *285*(1874), 20172593.

https://doi.org/10.1098/rspb.2017.2593

Srinivasan, U., Hines, J. E., & Quader, S. (2015). Demographic superiority with increased

logging in tropical understorey insectivorous birds. *Journal of Applied Ecology*, *52*(5),

1374–1380.

Srinivasan, U., & Wilcove, D. S. (2021). Interactive impacts of climate change and land-use

change on the demography of montane birds. *Ecology*, *102*(1), e03223.

https://doi.org/10.1002/ecy.3223

Stanton, J. C., Shoemaker, K. T., Pearson, R. G., & Akçakaya, H. R. (2015). Warning times for

species extinctions due to climate change. *Global Change Biology*, *21*(3), 1066–1077.

Supriya, K., Price, T. D., & Moreau, C. S. (2020). Competition with insectivorous ants as a

contributor to low songbird diversity at low elevations in the eastern Himalaya. *Ecology*

and Evolution, *10*(10), 4280–4290. https://doi.org/10.1002/ece3.6196

Swihart, R. K., Gehring, T. M., Kolozsvary, M. B., & Nupp, T. E. (2003). Responses of

'resistant'vertebrates to habitat loss and fragmentation: The importance of niche breadth

and range boundaries. *Diversity and Distributions*, *9*(1), 1–18.

Tattersall, G. J., Arnaout, B., & Symonds, M. R. E. (2017). The evolution of the avian bill as a

thermoregulatory organ. *Biological Reviews*, *92*(3), 1630–1656.

https://doi.org/10.1111/brv.12299

Teplitsky, C., & Millien, V. (2014). Climate warming and Bergmann's rule through time: Is there

any evidence? *Evolutionary Applications*, *7*(1), 156–168.

https://doi.org/10.1111/eva.12129

Thiollay, J.-M. (1992). Influence of Selective Logging on Bird Species Diversity in a Guianan

Rain Forest. *Conservation Biology*, *6*(1), 47–63.

Thuiller, W., Lavorel, S., & Araújo, M. B. (2005). Niche properties and geographical extent as

predictors of species sensitivity to climate change. *Global Ecology and Biogeography*,

14(4), 347–357. https://doi.org/10.1111/j.1466-822X.2005.00162.x

Tuff, K. T., Tuff, T., & Davies, K. F. (2016). A framework for integrating thermal biology into

fragmentation research. *Ecology Letters*, *19*(4), 361–374.

https://doi.org/10.1111/ele.12579

Velho, N., Ratnam, J., Srinivasan, U., & Sankaran, M. (2012). Shifts in community structure of

tropical trees and avian frugivores in forests recovering from past logging. *Biological*

Conservation, *153*, 32–40. https://doi.org/10.1016/j.biocon.2012.04.028

Voigt, W., Perner, J., & Hefin Jones, T. (2007). Using functional groups to investigate

community response to environmental changes: Two grassland case studies. *Global*

Change Biology, *13*(8), 1710–1721. https://doi.org/10.1111/j.1365-2486.2007.01398.x

Watt, C., Mitchell, S., & Salewski, V. (2010). Bergmann's rule; a concept cluster? *Oikos*, *119*(1),

89–100. https://doi.org/10.1111/j.1600-0706.2009.17959.x

Weeks, B. C., Willard, D. E., Zimova, M., Ellis, A. A., Witynski, M. L., Hennen, M., & Winger, B.

M. (2020). Shared morphological consequences of global warming in North American

migratory birds. *Ecology Letters*, *23*(2), 316–325.

Wingfield, J. C., Pérez, J. H., Krause, J. S., Word, K. R., González-Gómez, P. L., Lisovski, S., &

Chmura, H. E. (2017). How birds cope physiologically and behaviourally with extreme

climatic events. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

372(1723), 20160140. https://doi.org/10.1098/rstb.2016.0140

Zellweger, F., De Frenne, P., Lenoir, J., Vangansbeke, P., Verheyen, K., Bernhardt-

Römermann, M., Baeten, L., Hédl, R., Berki, I., Brunet, J., Van Calster, H.,

Chudomelová, M., Decocq, G., Dirnböck, T., Durak, T., Heinken, T., Jaroszewicz, B.,

Kopecký, M., Máliš, F., … Coomes, D. (2020). Forest microclimate dynamics drive plant

responses to warming. *Science*, *368*(6492), 772–775.

https://doi.org/10.1126/science.aba6880

Supplementary Material:

Figure S1: A map of the study area in Eaglenest Wildlife Sanctuary, Arunachal
Pradesh, India. Sampling plots in primary forest are outlined in green and plots in
logged forest are outlined in red. White arrows show the locations of mist nets
along with their corresponding temperature-humidity loggers.

Table S1: The common names and scientific names (Billerman et al., 2022) of the species selected for our analysis. All chosen species are breeders at our study site and had sufficient data for the modelling of survival trends and their temperature-humidity niches.

Common name	Scientific name
Black-throated Parrotbill	*Suthora nipalensis*
Rufous-capped babbler	*Cyanoderma ruficeps*
Yellow-throated Fulvetta	*Schoeniparus cinerea*
Rufous-winged Fulvetta	*Schoeniparus castaneceps*
Golden-breasted Fulvetta	*Lioparus chrysotis*
Snowy-browed Flycatcher	*Ficedula hyperythra*
White-gorgetted Flycatcher	*Anthipes monileger*
Grey-cheeked Warbler	*Phylloscopus poliogenys*
Rusty-fronted Barwing	*Actinodura egertoni*
Golden Babbler	*Cyanoderma chrysaeum*
White-spectacled Warbler	*Phylloscopus intermedius*
Blyth's Leaf Warbler	*Phylloscopus reguloides*
Chestnut-crowned Laughingthrush	*Trochalopteron erythrocephalum*
Black-faced Warbler	*Abroscopus schisticeps*
Large Niltava	*Niltava grandis*
White-tailed Robin	*Myiomela leucura*
White-browed Piculet	*Sasia ochracea*

865 **Table S2: Model parameters for the estimate and 95% confidence intervals around**
866 **the estimate for the generalised linear model to study the relationship between**
867 **body mass, time, habitat and niche overlap.**

Model:	Z-score ~ Year * Habitat * Niche_Overlap		
Parameter	Estimate	Lower CI bound	Upper CI bound
(Intercept)	0.34	0.01	0.66
Year	-0.04	-0.08	0.01
Niche_Overlap	-0.33	-0.68	0.03
Habitat_primary	-0.53	-0.98	-0.09
Year:Niche_Overlap	0.03	-0.01	0.08
Year:Habitat_primary	0.04	-0.02	0.1
Niche_overlap:Habitat_primary	0.52	0.04	1
Year:Niche_Overlap:Habitat_primary	-0.04	-0.11	0.03

868 **Table S3: Model parameters for the estimate and 95% confidence intervals around**
869 **the estimate for the generalised linear model to study the relationship between**
870 **body mass, time, habitat and niche overlap.**

Model:	Phi(~Year * Habitat * Niche_Overlap) p(~1)		
Parameter	Estimate	Lower CI bound	Upper CI bound
Phi:(Intercept)	0.57	-0.46	1.59
Phi:Year	-0.19	-0.41	0.02
Phi:Niche_Overlap	-0.87	-1.99	0.24
Phi:Habitat_primary	-1.98	-3.41	-0.55
Phi:Year:Niche_Overlap	0.24	0	0.47
Phi:Year:Habitat_primary	0.36	0.05	0.67
Phi:Niche_Overlap:Habitat_primary	2.27	0.71	3.83
Phi:Year:Niche_Overlap:Habitat_primary	-0.37	-0.71	-0.04
p:(Intercept)	-0.47	-0.61	-0.33