***Beet curly top Iran virus* Rep and V2 gene work as suppressors of post-transcriptional gene**
silencing through separate mechanisms

Saeideh Ebrahimi[1,2], Omid Eini[2,3], Alexandra Baßler[1], Zeynep Yildirim[1], Michael Wassenegger[1], Gabi
Krczal[1], Veli Vural Uslu[1,4,*]

[1] RLP AgroScience GmbH, Breitenweg 71, 67435, Neustadt an der Weinstraße, Germany
[2] Department of Plant Protection, University of Zanjan, 313 Zanjan, Iran
[3] Department of Phytopathology; Institute for Sugar Beet Research; 37079 Göttingen,
Germany
[4] MAPS, Heidelberg University, Heidelberg, Germany

Corresponding author: velivural.uslu@agroscience.rlp.de

ABSTRACT

Beet curly top Iran Virus (BCTIV) is a yield-limiting geminivirus belonging to the *becurtovirus*
genus. The genome organization of BCTIV is unique such that the complementary strand of BCTIV
resembles mastreviruses, whereas the virion strand organization is close to curtoviruses.
Geminiviruses are known to avoid the plant defense system by suppressing the RNA interference
mechanisms both at the transcriptional gene silencing (TGS) and post-transcriptional gene silencing
(PTGS) level. Multiple geminivirus genes have been identified as viral suppressors of RNA silencing
(VSR) but VSR activity remains elusive in becurtoviruses. By screening all verified open reading
frames in the BCTIV genome, we found that only V2 and Rep were able to suppress specific PTGS
mechanisms, triggered by the expression of a partial or full-length sense-strand transcript of the target
gene (S-PTGS). BCTIV-V2 could suppress S-PTGS more efficiently than BCTIV-Rep when then the
target GFP gene is transiently expressed. On the other hand, S-PTGS is suppressed by Rep but not V2
when target GFP is only stably expressed. Deletional mutagenesis of BCTIV-Rep implicated that
multiple domains are required for its VSR activity. Furthermore, neither V2 nor Rep could fully
suppress local PTGS induced by inverted repeat targeting GFP (GFP-IR). Also, in a closer look at the
spread of local silencing by GFP-IR, we observed that V2 or Rep are not able to suppress the
movement of sRNAs. Nevertheless, Rep suppressed the systemic silencing induced by GFP-IR in 16C
plants. Northern blot analyses showed that BCTIV-Rep inhibits silencing by mitigating sRNA
production, whereas BCTIV-V2 does not alter sRNA levels. In summary, both the silencing
phenotype and the molecular signatures of silencing implicate distinct modes of VSR activity of
BCTIV-Rep and -V2.

KEYWORDS

Geminivirus, RNAi, VSR, PTGS, S-PTGS, Rep, V2, p19, TBSV, Nicotiana benthamiana

BACKGROUND

Beet curly top Iran virus (BCTIV) has a monopartite single-stranded DNA (ssDNA) genome and

belongs to *becurtovirus* genus of the *geminiviridae* family (Heydarnejad *et al.* 2013). BCTIV is a

yield-limiting virus, causing a serious leaf curl disease in economically important crops such as sugar

beet, tomato, pepper, and beans (Yazdi *et al.* 2008, Heydarnejad *et al.* 2013). The leaf curl symptom

induced by BCTIV is very similar to those caused by other members of the geminiviridae, such as

beet curly top viruses (BCTV) of curtovirus genus (Anabestani et al. 2017). However, BCTIV

transmission relies on a distinct vector, *Circulifer haematoceps* (Taheri et al. 2012), and the genome

organization of BCTIV differs from the other members of the *geminividae* (Yazdi et al. 2008). Yet the

functions of the BCTIV encoded proteins have not been addressed rigorously.

Geminiviruses encode four to eight proteins in their mono- or bi-partite genomes. The transcription

from the geminivirus genome takes place in a bidirectional fashion: complementary strand (C) and

virion strand (V) (Teixeira et al. 2021). Geminiviruses contain a TAGTATTAC sequence at the

replication origin (*ori*), where the replication is initiated by the viral replication initiator protein,

known as the complementary strand protein 1 (C1) (Teixeira et al. 2021). Other proteins in the

complementary strand are transcriptional activator protein C2, replication enhancer protein C3, and

pathogenicity determinant protein C4 (Kumar 2019). In addition, in *mastrevirus*, *capulavirus*, and

grablovirus genera, due to a splicing event between C1 and C2, a new open reading frame (ORF),

called Rep, emerges containing the N terminus of C1 and a frame-shifted C terminus of C2. Rep

protein of geminiviruses is a multifunctional protein, which is indispensable for the viral genome

replication (Ruhel and Chakraborty 2019). On the other hand, the virion strand encodes the coat

protein (V1), movement protein (V2), and a poorly characterized gene (V3) (Kotlizky et al. 2000).

The C and V genes are organized in unique combinations for each genus of the geminiviridae family.

For example, BCTIV contains C1, C2, and Rep on the complementary-sense strand, similar to

Mastreviruses, whereas V1, V2, and V3 on the virion-sense strand, similar to curtoviruses (Bozorgi et

al. 2017, Zerbini et al. 2017) (Supplementary Figure 1a).

Geminivirus pathology is influenced by the efficiency of virus accumulation, which predominantly

depends on viral suppression of RNA interference (RNAi) in plants (Hanley-Bowdoin et al. 2013).

RNAi is a conserved defense mechanism, acting through transcriptional (TGS) or post-transcriptional

(PTGS) repression of gene expression in a sequence-specific manner (Baulcombe 2004). Bi-

directional transcription from the geminivirus genome creates double-stranded RNA (Vanitharani et

al. 2005). In addition, aberrant geminivirus transcripts can be used as templates for dsRNA production

by plant RNA-dependent RNA polymerases (RDRs) (Bisaro 2006). dsRNAs are cleaved into small

RNAs (sRNAs) of distinct lengths by Type III endonucleases, called DICER-LIKE (DCL) proteins

(Baulcombe 2004). sRNAs are incorporated into ARGONAUTE (AGO) proteins and form the RNA-

induced silencing complex (RISC) for the cleavage of the target sequence. Besides cleavage, RISC

can also control PTGS by recruiting RDRs for amplification of the RNAi or inhibiting target gene

translation facultatively (Yoshikawa et al. 2021, Uslu et al. 2021, Wu et al. 2020). In addition,

auxiliary RNAi proteins like HUA ENHANCER 1 (HEN1) RNA methyltransferase, which protects

sRNAs from degradation, contribute to viral defense in plants (Boutet et al. 2016, Muangsan et al.

2004, Jin et.al 2022)

There are many identified viral PTGS suppressors in plant viruses. For example, helper component-

proteinase (HC-Pro) encoded by *Potyviridae* inhibits RDR6 (Zhang et al. 2008) and HEN1 (Jamous et

al. 2011, Sanobar et al. 2021) among other components of the RNAi pathway, whereas, P19 protein

(P19) of *Tomato bushy stunt virus* (TBSV) sequesters ds-siRNAs inhibiting RNAi based protection in

plants (Voinnet et al. 2003, Ye et al. 2003). In addition, the P6 protein of *Strawberry vein banding*

virus (Feng et al. 2018), the P20, P23, and capsid proteins of *Citrus tristeza virus* (Lu et al. 2004), the

P6 protein of *Cauliflower mosaic virus* (Haas et al. 2008), P38 protein of *Turnip crinkle virus* (Iki et

al. 2017), P1 of *Sweet potato mild mottle virus* (Giner et al. 2010) are among those, which are have

been shown to function as viral suppressors of RNA silencing (VSR). In *Geminiviridae*, several

reports have implicated that C2, C4, Rep, V2, and V3 proteins (Liu et al. 2014, Wang et al. 2014,

Hanley-Bowdoin et al. 2013) function as viral VSRs, involved in the suppression of transcriptional

gene silencing (TGS) or post-transcriptional gene silencing (PTGS) in plants. Yet, the silencing

suppressor activity of the BCTIV gene products remains elusive.

In this study, we induced PTGS using specific silencing inducers, including full-length GFP, partial

sense-strand GFP, and inverted-repeat GFP to silence GFP expression. By combining the given

silencing inducers with each of the BCTIV open reading frames on *Nicotiana benthamiana*, we

investigated the efficiency of PTGS in local and systemic contexts to find out the BCTIV genes,

responsible for PTGS suppression.

METHODS

Plasmid constructs for transient expression

Sense-PTGS silencing inducers were cloned into pGreen (pG104) vector containing full length (FL-

GFP) or only 139nt long 5´-GFP sequence (5´-GFP) (Dalakouras et al. 2019). Strong PTGS was

induced by the inverted repeat of the very same 139nt sequence (GFP-IR) with a 90nt long spacer as

described in Dalakouras et al. 2019. The positive VSR control P19 of TBSV was expressed in

pPCV702SM binary vector (Saeed et al. 2015). All silencing inducers are expressed under the control
of 35S promoter.

For unbiased investigation of the genes potentially responsible for VSR all verified ORFs of BCTIV
(C1, C2, C4, Rep, V1, V2, V3) in pDIVA and/or pG104 binary vectors (Ebrahimi et al. 2022) were
examined on 16C plants (Supplementary Figure 1). The coding sequences of the C1, C2, Rep, and V2
genes were amplified with primers containing *BamHI* and *XhoI* restriction cut sites (Table 1), using
Q5 polymerase (NEB). The PCR products were first cloned into pJET1.2/blunt cloning vector
(Thermo Fischer Scientific, Germany) and they were sub-cloned using *Xho*I and *BamHI* sites of
pG104 vector to obtain pG-C1, pG-C2, pG-Rep and pG-V2, to obtain the same vector backbone for
all silencing inducers. The inserts were confirmed by Sanger sequencing. Of note, we found out that
the vector backbone did not influence the silencing or VSR phenotypes.

12 truncations from the Rep gene were PCR-amplified using corresponding primers (Table1) and
cloned into pJET1.2. Then, the plasmids with Rep truncations were digested with the *BamHI* and
XhoI enzymes they were cloned into the pG104 vector backbone to obtain pG-Rep1, pG-Rep2, pG-
Rep3, pG-Rep4, pG-Rep5, pG-Rep6, pG-Rep7, pG-Rep8, pG-Rep9, pG-Rep10, pG-Rep11, and pG-
Rep12 (Supplementary Figure 2) . *Escherichia coli* (strain INVαF, ThermoFisher) was used for
cloning plasmid constructs and *A. tumefaciens* strain ATHV was used for transient transformation of
the plants. pSoup helper plasmid was used together with pG104 for transforming agrobacterium by
electroporation. In addition to the plasmid constructs obtained from INVαF, plasmids in the
agrobacterium clones were also sequenced to ensure the insert integrity.

Agroinfiltration

The BCTIV genes were co-infiltrated with *Agrobacterium* ATHV containing silencing inducer
constructs into wildtype *Nicotiana benthamiana* (WT) and GFP-expressing *N.benthamiana* (16C) at
the four/six-leaf-stage. *Agrobacteria* containing the constructs were grown in LB medium (containing
Kanamycin and Rifampicin) at 28.0 °C for 16 h with continuous shaking at 135 rpm, then re-
suspended in MES buffer (10 mM MgCl2, 10 mM MES (pH 5.6), and 100 μM acetosyringone) to
reach $OD_{600}= 1$ for agroinfiltration. For negative control, *N. benthamiana* plants were separately
infiltrated with FL-GFP, 5´-GFP, and GFP-IR. In order to equalize the strength of silencing induction
in single silencing inducer infiltrations, agrobacterium containing empty pG104 ($OD_{600}= 1$) has been
added for mock co-infiltrations. Unless otherwise stated, three plants were used for each treatment in
each experiment, and each experiment was repeated at least three times at different times of the year
to avoid any effect of temperature fluctuations in the greenhouse.

GFP imaging and quantification

The inoculated plants were maintained in a growth chamber at 25 °C with 16-h light/ 8-h dark period.
After 3 days post inoculation (dpi) the control plants were monitored regularly under UV light (Blak-

Ray B-100 AP Lamp, www.uvp.com) to visualize the silencing phenotype. At 6dpi all the plants were

recorded using a Canon camera (Canon EOS700D, 18-5mm Lens) under UV light and leaves were

scanned using a laser scanner (Molecular Imager® PharosFX™ Systems, BioRad) at 50µm per pixel,

in GFP (excitation at 488nm, emission at 530nm) and in Chlorophyll (excitation at 488nm, emission

at 695nm) channels.

The GFP and the Chlorophyll images were merged and then GFP expression was quantified in the

limited area of infiltrated patches. GFP expression was measured by FIJI image analysis

(www.imagej.net) in treated, non-necrotic area and the acquired values were normalized using non

treated areas. The final values were plotted in bar charts (BoxPlotR, Shiny Lab).

148   Protein Extraction and Western blot

Total protein was extracted from 6 dpi infiltrated leaf patches of WT plants using protein extraction

buffer (1M Tris-HCl, PH 8.8, 10% Glycerol, 10% SDS, and H2O) together with cOmplete™ Protease

inhibitor solution (Roche). SDS-PAGE was performed for Coomassie staining and Western blotting

(Wb), polyclonal goat anti-GFP antibody (Sicgen, AB0066-200, 1:1000) as primary, Donkey anti-

goat IgG-HRP antibody (sc-2020, Santa Cruz Biotechnology, (1:5000) as secondary antibody were

used for western blot. Pierce™ ECL Western Blotting Substrate (ThermoFisher) was used to detect

the signal. 16C *N.benthamiana* were not included for the western blot analysis due to the fact that the

vasculature of 16C is barely affected by silencing and depending on how much vasculature is included

in the western blot sample collection from silenced areas, the results fluctuate dramatically.

158   RNA extraction, RT-PCR, and Northern blot

Total RNA was extracted from infiltrated patches of 16C leaves at 6 dpi using TRIzol® Reagent

(Qiagen). RNA concentration and RNA purity were measured using Nanodrop (ND-1000) and RNA

quality was assessed by agarose gel electrophoresis. RNAs are treated with Lucigen Baseline-ZERO

DNase and purified by acid-Phenol-Chloroform pH:4.5 (Invitrogen™, Thermofisher). RT-PCR was

performed using the SuperScript™ III One-Step RT-PCR System with Platinum™ Taq DNA

Polymerase (Invitrogen™, ThermoFisher) and gene-specific primers (Table 2) to amplify BCTIV

genes to confirm transient expression of the constructs upon agroinfiltration. For the Northern blot of

GFP mRNA and GFP siRNA 10µg and 5µg of total RNA were used, respectively, as previously

described (Dalakouras et al. 2019). The siRNA Northern blots were stripped and rehybridized with a

U6 snoRNA probe. For GFP probes [alpha-P32] CTP is used for labeling, for U6 snoRNA probe

[gamma-p32] ATP is used from Hartman Analytic, Germany. The detection of the Northern blot

signals was performed by the Molecular Imager® PharosFX™ Systems (BioRad).

171   Statistical Analysis

Statistical tests have been performed by One-way ANOVA using JMP Software

(https://www.jmp.com/). Unless otherwise stated, $p < 0.05$ is taken as the significance threshold.

Phylogenetic Tree

Nucleotide sequences of the Rep (or homologous) proteins of different geminiviruses were obtained

from NCBI. MEGA (Molecular Evolutionary Genetics Analysis) was used for the phylogenetic tree,

based on the neighbor-joining method.

RESULTS

Phenotypic analysis of transient BCTIV gene expression and silencing inducers on *N.*

benthamiana

Before we began investigating the VSR activity of BCTIV genes, we tested the expression and the

effect of the BCTIV genes upon agroinfiltration into 16C *N. benthamiana* leaves. C1, C2, Rep, V1,

V2, and V3 genes, which were previously shown to be expressed by BCTIV, and C4 was shown to be

expressed in BCTV. (Supplementary Figure 1a) (Eini et al. 2016, Yazdi et al. 2008, Teng et al. 2010).

Using RT-PCR, we confirmed the expression of the viral genes in *N. benthamiana* leaves

(Supplementary Figure 1b). Notably, C1 and Rep expression also led to necrotic speckles starting

from 5dpi and the necrotic phenotype gradually got stronger. As sense-PTGS (S-PTGS) inducer, the

5'-GFP construct was agroinfiltrated into GFP expressing 16C leaves. The silencing phenotype

induced by 5'-GFP was most consistently visible earliest at 6 dpi. Therefore, the effect of BCTIV

genes on local PTGS was examined at 6 dpi.

BCTIV-Rep suppresses local sense-PTGS in *N.benthamiana*

First, we aimed at investigating the role of BCTIV genes in the Sense-PTGS (S-PTGS) pathway. For

this assay, 139nt long sequence mapping to GFP (5'-GFP) was transiently expressed together with the

BCTIV genes in 16C *N. benthamiana*. Due to S-PTGS induction, agroinfiltrated leaf sectors turn red

under UV light due to loss of GFP at 6 dpi. However, if a BCTIV gene functions as a viral suppressor

of RNA silencing (VSR) the agroinfiltrated sector should remain green under UV light (Figure 1a).

Based on the GFP expression level under the UV light at 6 dpi, we found out that only Rep protein

could suppress the S-PTGS, while none of the other constructs, including the positive control TBSV-

P19, could suppress S-PTGS (Figure 1b). For the exact quantification of GFP levels after silencing

induction and silencing suppression, the leaves were imaged in a high-resolution laser scanner (Figure

1c). The GFP intensity measurements in exclusively non-necrotic areas showed that GFP expression

was significantly ($P < 0.05$) higher in Rep infiltrated 16C *N. benthamiana* leaves in comparison to

TBSV-P19 and other BCTIV gene infiltration plants. (Figure 1d).

To reveal the mode of action of VSR activity of BCTIV-Rep, we investigated the GFP mRNA and
small RNA populations upon co-infiltration in the leaf sections of 16C plants using Northern blot. To
compare with Rep, we used a restricted set of BCTIV-genes, including C1 and C2 because of
overlapping sequences with Rep, and V2 genes due to the VSR activity of V2 homologs in other
geminiviruses (Luna et al. 2017, Zhang et al. 2012, Zrachya et al. 2006). Despite loading the same
amounts of material according to nanodrop measurements, the RNA quality of different co-
infiltrations differed observably based on rRNA bands on the agarose gel electrophoresis of the RNA
samples. (Figure 1e). Especially, due to the necrotic areas caused by Rep, the RNA integrity was the
lowest in Rep + 5´-GFP co-infiltrated sample. Nevertheless, the GFP mRNA level in Rep + 5´-GFP
was the highest when compared to other co-infiltrated samples (Figure 1e).

In addition, we performed sRNA Northern Blot on the very same co-infiltrated samples. End labeled
U6 snoRNA probe was used as a loading control, which did not reveal any major fluctuations of
sRNA concentration (Figure 1f). Samples from untreated 16C did not show any sRNAs, while 5´-GFP
treated 16C samples had a clear accumulation of sRNAs, suggesting that the experimental setup was
valid. Rep + 5´-GFP sample showed a very faint sRNA band at the level of 21-24 nucleotides (nt),
suggesting that Rep interfered with the silencing pathway, before the production of sRNAs by DCLs
(Figure 1f). P19 and V2 co-infiltration with 5´-GFP led to only a limited decrease in sRNA production
in concordance with the previous literature (Saeed et al. 2015).

BCTIV-V2 suppresses local S-PTGS only when the target GFP is transiently expressed.

In order to detect the changes in GFP expression and silencing levels in a clean background with no
GFP fluorescence over the whole leaf surface, full-length GFP (GFP-FL) was co-infiltrated together
with BCTIV-genes in WT *N. benthamiana*. GFP-FL under the control of the 35S promoter induces
GFP expression. However, due to the S-PTGS mediated self-silencing of the transient GFP-FL
expression, the GFP signal remains low. When a BCTIV gene increases the GFP expression, upon co-
infiltration with GFP-IR, this should be due to the VSR activity of the gene, which inhibits S-PTGS,
hence, the self-silencing of GFP-FL (Figure 2a).

Plants under the UV light at 6 dpi showed high GFP expression in the BCTIV-V2 and BCTIV-Rep
treated plants compared to only GFP-FL infiltrated plants (Figure 2b). The strongest green
fluorescence was observed in TBSV-P19 treated plants, in accordance with the previous literature
(Figure 2b) (Ahn et al. 2011, Saeed et al. 2015, Zhang 2015). The quantification based on high-
resolution images of infiltrated leaves also corroborated with the observations under the UV lamp, in
which P19, V2, and Rep showed higher GFP signal than the GFP-FL only and GFP-FL with C1 or C2
(Figure 2c, d).

Since WT plants provided a blank background for GFP analysis, we addressed whether the
quantifications on the images based on GFP intensity in non-necrotic areas matched with the GFP
amount in the non-necrotic areas obtained by the Western blot of the infiltrated areas. The Western
Blot results showed that co-infiltration of GFP-FL with P19, V2, and Rep led to higher GFP protein
expression than GFP-FL alone at 6dpi, in the same order of intensity (Figure 2e). Moreover, C1 and
C2 co-infiltration with GFP-FL did not increase the GFP protein level in Western blot in accordance
with the image analysis.

We performed the GFP-FL co-infiltration with P19, V2, and Rep on GFP expressing 16C
N.benthamiana. GFP-FL increases GFP expression but also functions as an S-PTGS inducer in the
later stages. At 5.25dpi, we found out that the net contribution of GFP-FL on the GFP level is zero
(Figure 2f, g). At this stage, P19 works as a silencing suppressor, more efficiently than V2. Although
V2+GFP-FL tends to have a stronger silencing suppressor effect than Rep+GFP-FL on average, the
difference is not statistically significant (Figure 2g).

Neither Rep nor V2 can suppress PTGS induced by a hairpin construct targeting GFP.
Since Rep is capable of decreasing sRNA production, necessary for S-PTGS, we addressed whether
Rep could repress PTGS induced by hairpin RNA (GFP-IR) expressed under the control of the 35S
promoter. Similar to the S-PTGS infiltrations using GFP-FL and 5´GFP, the same agrobacterium
batch containing BCTIV genes (C1, C2, Rep and V2) were co-infiltrated with GFP-IR in stably GFP
expressing 16C plants (Figure 3a). GFP-IR is a very strong inducer of silencing and at 6dpi, none of
C1-, C2-, Rep-, V2-, P19- co-infiltrated leaves showed loss of the silencing phenotype (Figure 3b).
Nevertheless, C1 and Rep co-infiltrated with GFP-IR had a slightly higher level of GFP than GFP-IR
only (Figure 3c, d). Therefore, we investigated the molecular fingerprints of BCTIV genes on PTGS
by checking GFP mRNA and sRNA levels using Northern blot on stably GFP expressing 16C plants.
Due to lower necrotic areas in Rep + GFP-IR sample, the RNA quality was comparable to the other
treatments and controls. In addition, the abundance of GFP mRNA in 16C and the low level of GFP in
GFP-IR treatment validated the experimental set-up. None of the treatments, including Rep, could
block the decrease of GFP mRNA level upon GFP-IR induction (Figure 3e).
Next, we performed an sRNA Northern blot on GFP-IR induced PTGS samples. U6 snoRNA loading
control shows comparable amounts of total sRNAs among different treatments and controls (Figure
3f). In addition, GFP-IR only shows very strong sRNA accumulation, whereas no sRNAs are visible
in 16C control, validating the experimental set-up. Interestingly, we have seen that the sRNAs
mapping to GFP are almost absent in Rep + GFP-IR and dramatically reduced in C1+GFP-IR at 6dpi.
On the other hand, V2 and P19 co-expression with GFP-IR did not have any effect on sRNA
accumulation at 6dpi (Figure 3f).
274
can suppress systemic silencing 2Rep but not V

To find out whether the decrease in sRNA amounts upon expression of BCTIV genes suppresses
systemic silencing in 16C plants (Figure 3f), the GFP-IR treated plants were observed and recorded
regularly under UV light until 14dpi (Figure 3g). Systemic silencing emerged at 10 dpi in plants
infiltrated with GFP-IR only, upon C2+GFP-IR and V2+GFP-IR co-infiltration the systemic silencing
phenotype spread in the newly emerging leaves at 14 dpi, without any exception (Figure 3g, h).
However, none of the eighteen plants co-infiltrated with Rep+GFP-IR showed systemic silencing
(Figure 3h). These results indicate Rep protein suppression activity in systemic PTGS. It is
noteworthy that twelve out of eighteen C1+GFP-IR treated plants did not show any systemic silencing
and the remaining six plants showed rather restricted systemic spread of the silencing signal (Figure
3g, h).

287 Rep and V2 do not interfere with the spread of silencing through plasmodesmata

Local silencing induced by GFP-IR and/or GFP-FL creates a visible area encircling the infiltration
zone, which shows stronger silencing than the infiltration zone (Figure 4a, Himber et al. 2003). It is
postulated that the silencing signal reaches this area via the short-range spreading of sRNAs through
plasmodesmata (Dunoyer et al. 2005, Kalantidis et al. 2008). It is possible to quantify the length and
GFP intensity (Gray value) of this area by plotting the profile (using FIJI) along a line in the GFP
channel. Our data showed that the GFP-IR+GFP-FL co-infiltration created a silenced zone of 0.7 mm
around the infiltration zone, which roughly corresponds to 15 cells (Figure 4b). When GFP-FL, GFP-
IR are co-infiltrated with p19, V2 or Rep on 16C *N.benthamiana*, the silencing phenotype is partially
repressed at 5.25dpi (Figure 4c, e, g, i). P19 abolishes the silenced ring around the infiltrated area
(Figure 4d). However, the silenced area is preserved when BCTIV-V2 and BCTIV-Rep are co-
infiltrated with the GFP-IR and GFP-FL (Figure 4f, g).

299 Multiple Rep domains are indispensable for VSR activity

To figure out if there is any specific domain responsible for suppression activity in Rep protein,
different truncations of the Rep gene were constructed based on C1 and frameshift C2 motifs
(Supplementary Fig 2). The Rep truncations were co-expressed together with 5´-GFP in 16C *N.*
benthamiana plants (Figure 5a). None of the 12 Rep truncations recapitulated the Rep-VSR activity at
6 dpi (Figure 5b) which correlates with GFP quantification in infiltrated leaf patches (Figure 5 c,d).
Noteworthy that the 30 amino acid long deletion at the C-terminus of Rep partially kept the VSR
activity, suggesting that the C terminus could have minor functions for S-PTGS suppression. Besides,
N-terminus deletions but not C-terminus deletions of Rep truncations showed necrotic phenotypes
comparable to Full-length Rep protein. These results suggest that multiple domains BCTIV-Rep
domains except the very end of C-terminus, are indispensable for silencing function, but the N-

terminus of the Rep protein is essential for necrosis, which is similar to wheat dwarf virus (WDV)

Rep protein (Liu et al. 2014)

DISCUSSION

RNA silencing is a cellular defense mechanism conserved across kingdoms, which is triggered by ds-

RNA in plants (Zhao *et al.* 2022). In spite of being ssDNA viruses with no dsRNA phase in their

replication cycle, geminiviruses have been shown to be targeted by plants via post-transcriptional

gene silencing by producing bi-directional transcription from opposite polarity or the strong fold-back

structure of transcripts (Vanitharani et al. 2005). In addition, viral gene expression can lead to

aberrant transcripts, which are further processed by RDRs to form dsRNAs. Therefore, geminiviruses

must confront posttranscriptional gene silencing (PTGS) to achieve successful infections (Luna et al.

2017). Several open reading frames were identified as PTGS suppressors in geminiviruses but still,

there is no report of silencing suppressors in BCTIV (Ramesh et al. 2017). In this study, we used

distinct approaches to dissect the role of BCTIV genes in the suppression of PTGS.

When the GFP-FL is agroinfiltrated on WT plants, it induces GFP expression, but it also silences

itself due to the production of aberrant RNAs, triggering S-PTGS (Parent et al. 2015). When GFP-FL

is transiently expressed, we showed that BCTIV-V2 works as an effective silencing suppressor

showing overlapping features with TBSV-P19 VSR activity whereas Rep also represses the self-

silencing of FL-GFP but only very mildly (Figure 2). However, when the GFP is solely expressed in

the genome in 16C *N. benthamiana* (without transient GFP-FL expression) and the S-PTGS is

induced by transient expression of 5´-GFP, TBSV-P19 and BCTIV-V2 have no repressive effect on

the silencing, whereas BCTIV-Rep substantially suppresses the S-PTGS (Figure 1). The Northern

blots of leaves in which S-PGTS was induced by infiltration with 5´-GFP, TBSV-P19, and BCTIV-

V2 co-infiltrated samples showed accumulation of sRNAs, whereas BCTIV-Rep co-infiltrated sample

was almost fully devoid of sRNAs (Figure 1). The sRNA signature indicates that P19 and V2 do not

interfere with the sRNA production substantially. On the other hand, Rep interferes with sRNA

production. These observations implicate that BCTIV-Rep has a distinct mechanism of action from

BCTIV-V2 and TBSV-P19.

GFP-IR is a very strong silencing inducer, which can lead to systemic silencing within 10 dpi. Upon

transient expression of GFP-IR, due to the hairpin structure, dsRNA immediately forms and is

subsequently processed by the DCL proteins. In this situation, none of the silencing suppressors,

including Rep could repress the local silencing induced by GFP-IR (Figure 3). Nevertheless, the

sRNA Northern blot showed that Rep and C1 could decrease the amount of sRNA produced upon

GFP-IR expression. Although the decrease did not lead to suppression of local silencing, the systemic

silencing phenotype was consequently affected. Contrary to Rep, V2 did not have any effect on

systemic silencing. It has recently been proposed that the AGO proteins sequester the sRNAs and

mobile sRNAs can induce systemic silencing only when their concentration is above a certain

threshold, determined by the concentration of AGO proteins (Vionnet 2022). Therefore, the reduction

in sRNA level by Rep which does not affect local silencing is low enough to avoid induction of

systemic silencing. Besides, it is worth mentioning that the necrotic phenotype of the Rep and C1

could further decrease the production of sRNAs after 6dpi, which contributes at least partly to the lack

of systemic silencing phenotype in these co-infiltrations.

Nevertheless, it was still puzzling that P19 did not lead to any silencing suppression when 5´-GFP or

GFP-IR was expressed on stably GFP expressing 16C plants (Supplementary Figure 3a, b). The vast

majority of the characterization of the P19 activity was performed under transient expression of GFP

in 16C or WT backgrounds (Martin-Hernandez and Baulcombe, 2008). Confirming the previous

literature, when GFP was transiently expressed, we could recapitulate the silencing suppression

phenotype of p19 (and V2) on WT or 16C plants (Vargason et al. 2003, Silhavy et al. 2002)

(Supplementary Figure 3c-f). The underlying mechanism of the VSR activity of TBSV-P19 on S-

PTGS depending on the presence of transient GFP-FL expression remains enigmatic (Supplementary

Figure 3a-f). In order to understand the phenomenon better, we transiently expressed P19 two and one

day before the GFP-IR expression on 16C samples. In three independent experiments, we consistently

found out that P19 can act as a silencing suppressor in the absence of GFP-FL, if it is expressed not

one day but at least two days before the silencing inducer GFP-IR or 5'-GFP is expressed on 16C

plants (Supplementary Fig 4a-e). Although p19 suppresses silencing when co-infiltrated with GFP-

FL, when GFP-FL is expressed one or two days before p19, p19 cannot suppress the silencing

(Supplementary Figure 4f-i). It has previously been demonstrated that P19 binds to ds-siRNA to

inhibit the silencing derived from ds-siRNA (Vargason et al. 2003). Also, in line with P19-binding

specificity, P19 failed to inhibit the activation of RISC by ss-siRNA, similarly, P19 was not able to

suppress the target cleavage of the already activated RISC (Lakatos et al. 2004). Furthermore, another

study showed CymRSV infection was not able to revert the already established GFP RNA silencing.

(Silhavy et al. 2002). In summary, our experiments as well as the given literature indicate that the

time of p19 expression relative to the GFP-FL is critical for its function. The difference in timing may

create a difference in p19 and siRNA concentrations at a given time, which is critical for the VSR

activity.

Although this is the first report of silencing suppressor gene of BCTIV-Rep protein, Rep was

previously identified as a PTGS suppressor in different geminiviruses like *Wheat dwarf virus* (WDV)

from *Mastrevirus* (Kasschau and Carrington 1998, Liu et al. 2014). Yet, BCTIV-Rep protein is only

very distantly related to WDV-Rep proteins based on nucleic acid sequence (Supplementary Figure

5). Therefore, the precise identification of the RNAi pathways that Rep proteins involve in may shed

light on uncharted functions of Rep proteins in geminiviruses. Besides, Rep proteins from TYLCV,

TGMV, and ACMV were shown to interfere with the plant DNA methylation machinery and suppress

transcriptional gene silencing (Rodríguez-Negrete et al. 2013). However, the role of BCTIV-Rep

protein on TGS remains unexplored. Moreover, V2 and its homolog AV2 have previously been
identified as a suppressor in Geminiviruses, with a weaker silencing outcome than P19 (Sharma et al.
2010), but it had not been reported for BCTIV before this study.

To highlight the mode of action of PTGS suppressor activity of BCTIV, we propose that BCTIV-Rep
protein interferes with RDR6 (Figure 7). In S-PTGS, where WT and 16C *N.benthamiana* inoculated
with FL-GFP and 5´GFP, respectively, RDR6 is recruited to produce dsRNA from the aberrant GFP
transcripts (Figure 5a). Rep protein suppressed the local silencing derived from RDR6 mediated
dsRNA production. Therefore, 5´-GFP infiltrations do not lead to sRNA production when co-
infiltrated with Rep. Unlike during S-PTGS, in GFP-IR induced PTGS, the hairpin transcript is
already dsRNA, therefore RDR6 is not required for the production of primary siRNAs, which leads to
local silencing (Figure 5b) and in consequence BCTIV-Rep does not suppress IR-induced PGTS. It is
noteworthy, however, that RDR6 is also indispensable for systemic silencing (Harmoko et al. 2013,
Vaistij and Jones 2009, Dalakouras et al. 2020). The model where Rep interferes with RDR6 activity
can explain our current observations, but further experiments are needed to prove the direct or indirect
interaction of Rep and RDR6.

The Rep protein of WDV has been functionally characterized. Truncations of WDV-Rep at the N
terminus (9 amino acids) and the C-terminus (30 amino acids) did not affect the silencing suppression
function and the N-terminus was critical for the necrotic phenotype (Wang et al. 2014). On the other
hand, we have further characterized BCTIV-Rep with truncating domains on both the N and the C
terminus. In the line with previous findings in WDV, the pathogenicity determinant domain of
BCTIV-Rep is also located on the N-terminus. However, we found that all truncated Rep proteins we
produced, did not maintain the suppression of the silencing phenotype of BCTIV-Rep, except that C-
terminus is shown to have a limited effect on the silencing suppression of Rep.

CONCLUSION
In this study, we determined the BCTIV-Rep and -V2 as viral suppressors of PTGS. We showed that
Rep reduces the siRNA accumulation upon transient and stable expression of the target gene. On the
other hand, BCTIV-V2 protein also has S-PTGS suppression activity only if GFP-FL is also
transiently expressed, which is similar to its homologous geminivirus-V2 proteins and TBSV-P19.

Data Availability Statement
The coding sequences of the BCTIV genes are provided in the text. In addition, plasmids used in this
study are available upon request.

Statement of Conflict
The authors declare that they have no competing interests.

Funding

This study was financially supported by the Federal Ministry of Education and Research (BMBF)
Grant 03180531. SE was supported by the University of Zanjan and the Ministry of Science, Research
and Technology/Iran. ZY was financially supported by Erasmus+ Student Mobility for Traineeships
fellowship.

Author Contributions

OE, GK, MW, and VVU designed the project. SE, AB, ZY conducted experiments. SE, VVU, and OE
analyzed the data. VVU, GK, and SE wrote the manuscript. All authors analyzed the data and
reviewed the manuscript. VVU supervised the study.

Acknowledgements

We thank Arvid Hanke and the other Epigenetics Lab members at RLP AgroScience GmbH for the
careful revision of the data and the manuscript.

REFERENCES:

1.   Ahn JW, Lee JS, Davarpanah SJ, et al. Host-dependent suppression of RNA silencing
       mediated by the viral suppressor p19 in potato. *Planta*. 2011;234(5):1065-1072.
       doi:10.1007/S00425-011-1465-1/FIGURES/6

2.   Anabestani A, Behjatnia SAA, Izadpanah K, Tabein S, Accotto GP. Seed
       Transmission of Beet Curly Top Virus and Beet Curly Top Iran Virus in a Local
       Cultivar of Petunia in Iran. *Viruses*. 2017;9(10). doi:10.3390/V9100299

3.   Baulcombe D. RNA silencing in plants. *Nat 2004 4317006*. 2004;431(7006):356-363.
       doi:10.1038/nature02874

4.   Bisaro DM. Silencing suppression by geminivirus proteins. *Virology*.
       2006;344(1):158-168. doi:10.1016/J.VIROL.2005.09.041

5.   Bozorgi N, Heydarnejad J, Kamali M, Massumi H. Splicing features in the expression
       of the complementary-sense genes of Beet curly top Iran virus. *Virus Genes*.
       2017;53(2):323-327. doi:10.1007/S11262-016-1422-Y

6.   Dalakouras A, Lauter A, Bassler A, Krczal G, Wassenegger M. Transient expression
of intron-containing transgenes generates non-spliced aberrant pre-mRNAs that are
processed into siRNAs. *Planta*. 2019;249(2):457-468. doi:10.1007/S00425-018-3015-
6/FIGURES/8

7.   Dunoyer P, Himber C, Voinnet O. DICER-LIKE 4 is required for RNA interference
and produces the 21-nucleotide small interfering RNA component of the plant cell-to-
cell silencing signal. *Nat Genet*. 2005;37(12):1356-1360. doi:10.1038/NG1675

8.   Ebrahimi S, Eini O, Koolivand D, Varrelmann M. The Rep and C1 of Beet curly top
Iran virus represent pathogenicity factors and induce hypersensitive response in
Nicotiana benthamiana plants. *Virus Genes*. Published online August 12, 2022.
doi:10.1007/S11262-022-01927-3

9.   Eini O, Sahraei GE, Ali S, Behjatnia A. Molecular characterization and construction of
an infectious clone of a pepper isolate of Beet curly top Iran virus. *Mol Biol Res*
Commun. 2016;5(2):101. Accessed August 10, 2022. /pmc/articles/PMC5219900/

10.   Feng M, Zuo D, Jiang X, et al. Identification of Strawberry vein banding virus encoded
P6 as an RNA silencing suppressor. *Virology*. 2018;520:103-110.
doi:10.1016/J.VIROL.2018.05.003

11.   Giner A, Lakatos L, García-Chapa M, López-Moya JJ, Burgyán J. Viral protein
inhibits RISC activity by argonaute binding through conserved WG/GW motifs. *PLoS*
Pathog. 2010;6(7):1-13. doi:10.1371/JOURNAL.PPAT.1000996

12.   Haas G, Azevedo J, Moissiard G, et al. Nuclear import of CaMV P6 is required for
infection and suppression of the RNA silencing factor DRB4. *EMBO J*.
2008;27(15):2102-2112. doi:10.1038/EMBOJ.2008.129

13.   Hanley-Bowdoin L, Bejarano ER, Robertson D, Mansoor S. Geminiviruses: masters at
redirecting and reprogramming plant processes. *Nat Rev Microbiol*. 2013;11(11):777-
788. doi:10.1038/NRMICRO3117

14.   Harmoko R, Fanata WID, Yoo JY, et al. RNA-Dependent RNA Polymerase 6 Is

477   Required for Efficient hpRNA-Induced Gene Silencing in Plants. *Mol Cells*.

478   2013;35(3):202. doi:10.1007/S10059-013-2203-2

479   15.   Heydarnejad J, Keyvani N, Razavinejad S, Massumi H, Varsani A. Fulfilling Koch's

480   postulates for beet curly top Iran virus and proposal for consideration of new genus in

481   the family Geminiviridae. *Arch Virol*. 2013;158(2):435-443. doi:10.1007/S00705-012-

482   1485-6

483   16.   Himber C, Dunoyer P, Moissiard G, Ritzenthaler C, Voinnet O. Transitivity-dependent

484   and -independent cell-to-cell movement of RNA silencing. *EMBO J*.

485   2003;22(17):4523-4533. doi:10.1093/EMBOJ/CDG431

486   17.   Iki T, Tschopp MA, Voinnet O. Biochemical and genetic functional dissection of the

487   P38 viral suppressor of RNA silencing. *RNA*. 2017;23(5):639-654.

488   doi:10.1261/RNA.060434.116

489   18.   Jin L, Chen M, Xiang M, Guo Z. RNAi-Based Antiviral Innate Immunity in Plants.

490   *Viruses 2022, Vol 14, Page 432*. 2022;14(2):432. doi:10.3390/V14020432

491   19.   Kalantidis K, Schumacher HT, Alexiadis T, Helm JM. RNA silencing movement in

492   plants. *Biol Cell*. 2008;100(1):13-26. doi:10.1042/BC20070079

493   20.   Kasschau KD, Carrington JC. A counterdefensive strategy of plant viruses:

494   suppression of posttranscriptional gene silencing. *Cell*. 1998;95(4):461-470.

495   doi:10.1016/S0092-8674(00)81614-1

496   21.   Kotlizky G, Boulton MI, Pitaksutheepong C, Davies JW, Epel BL. Intracellular and

497   Intercellular Movement of Maize Streak Geminivirus V1 and V2 Proteins Transiently

498   Expressed as Green Fluorescent Protein Fusions. *Virology*. 2000;274(1):32-38.

499   doi:10.1006/VIRO.2000.0415

500   22.   Kumar RV. Plant antiviral immunity against geminiviruses and viral counter-defense

501   for survival. *Front Microbiol*. 2019;10(JUN):1460.

502   doi:10.3389/FMICB.2019.01460/BIBTEX

23.   Liu Y, Jin W, Wang L, Wang X. Replication-associated proteins encoded by Wheat
dwarf virus act as RNA silencing suppressors. *Virus Res*. 2014;190:34-39.
doi:10.1016/J.VIRUSRES.2014.06.014

24.   Lu R, Folimonov A, Shintaku M, et al. Three distinct suppressors of RNA silencing
encoded by a 20-kb viral RNA genome. *Proc Natl Acad Sci U S A*.
2004;101(44):15742-15747. doi:10.1073/PNAS.0404940101

25.   Luna AP, Rodríguez-Negrete EA, Morilla G, et al. V2 from a curtovirus is a
suppressor of post-transcriptional gene silencing. *J Gen Virol*. 2017;98(10):2607-2614.
doi:10.1099/JGV.0.000933

26.   Martín-Hernández AM, Baulcombe DC. Tobacco rattle virus 16-kilodalton protein
encodes a suppressor of RNA silencing that allows transient viral entry in meristems. *J
Virol*. 2008;82(8):4064-4071. doi:10.1128/JVI.02438-07

27.   Muangsan N, Beclin C, Vaucheret H, Robertson D. Geminivirus VIGS of endogenous
genes requires SGS2/SDE1 and SGS3 and defines a new branch in the genetic
pathway for silencing in plants. *Plant J*. 2004;38(6):1004-1014. doi:10.1111/J.1365-
313X.2004.02103.X

28.   Parent JS, Jauvion V, Bouché N, et al. Post-transcriptional gene silencing triggered by
sense transgenes involves uncapped antisense RNA and differs from silencing
intentionally triggered by antisense transgenes. *Nucleic Acids Res*. 2015;43(17):8464-
8475. doi:10.1093/NAR/GKV753

29.   Ramesh S V., Sahu PP, Prasad M, Praveen S, Pappu HR. Geminiviruses and Plant
Hosts: A Closer Examination of the Molecular Arms Race. *Viruses*. 2017;9(9).
doi:10.3390/V9090256

30.   Ruhel R, Chakraborty S. Multifunctional roles of geminivirus encoded replication
initiator protein. *Virusdisease*. 2019;30(1):66-73. doi:10.1007/S13337-018-0458-0

31.   Saeed M, Briddon RW, Dalakouras A, Krczal G, Wassenegger M. Functional Analysis
of Cotton Leaf Curl Kokhran Virus/Cotton Leaf Curl Multan Betasatellite RNA

Silencing Suppressors. *Biol 2015, Vol 4, Pages 697-714*. 2015;4(4):697-714.

doi:10.3390/BIOLOGY4040697

32. Sanobar N, Lin PC, Pan ZJ, et al. Investigating the Viral Suppressor HC-Pro Inhibiting

Small RNA Methylation through Functional Comparison of HEN1 in Angiosperm and

Bryophyte. *Viruses 2021, Vol 13, Page 1837*. 2021;13(9):1837.

doi:10.3390/V13091837

33. Silhavy D, Molnár A, Lucioli A, et al. A viral protein suppresses RNA silencing and

binds silencing-generated, 21- to 25-nucleotide double-stranded RNAs. *EMBO J*.

2002;21(12):3070-3080. doi:10.1093/EMBOJ/CDF312

34. Stanley J, Latham JR. A symptom variant of beet curly top geminivirus produced by

mutation of open reading frame C4. *Virology*. 1992;190(1):506-509.

doi:10.1016/0042-6822(92)91243-N

35. Teixeira RM, Ferreira MA, Raimundo GAS, Fontes EPB. Geminiviral Triggers and

Suppressors of Plant Antiviral Immunity. *Microorganisms*. 2021;9(4).

doi:10.3390/MICROORGANISMS9040775

36. Teng K, Chen H, Lai J, et al. Involvement of C4 Protein of Beet Severe Curly Top

Virus (Family Geminiviridae) in Virus Movement. *PLoS One*. 2010;5(6).

doi:10.1371/JOURNAL.PONE.0011280

37. Vaistij FE, Jones L. Compromised Virus-Induced Gene Silencing in RDR6-Deficient

Plants. *Plant Physiol*. 2009;149(3):1399. doi:10.1104/PP.108.132688

38. Vargason JM, Szittya G, Burgyán J, Tanaka Hall TM. Size Selective Recognition of

siRNA by an RNA Silencing Suppressor. *Cell*. 2003;115(7):799-811.

doi:10.1016/S0092-8674(03)00984-X

39. Voinnet O. Revisiting small RNA movement in plants. *Nat Rev Mol Cell Biol*.

2022;23(3):163-164. doi:10.1038/s41580-022-00455-0

40.   Voinnet O, Rivas S, Mestre P, Baulcombe D. An enhanced transient expression system
in plants based on suppression of gene silencing by the p19 protein of tomato bushy
stunt virus. *Plant J*. 2003;33(5):949-956. doi:10.1046/J.1365-313X.2003.01676.X

41.   Wang Y, Dang M, Hou H, Mei Y, Qian Y, Zhou X. Identification of an RNA silencing
suppressor encoded by a mastrevirus. *J Gen Virol*. 2014;95(Pt 9):2082-2088.
doi:10.1099/VIR.0.064246-0

42.   Wu H, Li B, Iwakawa H oki, et al. Plant 22-nt siRNAs mediate translational repression
and stress adaptation. *Nat 2020 5817806*. 2020;581(7806):89-93. doi:10.1038/s41586-
020-2231-y

43.   Yazdi HRB, Heydarnejad J, Massumi H. Genome characterization and genetic
diversity of beet curly top Iran virus: a geminivirus with a novel nonanucleotide. *Virus
Genes*. 2008;36(3):539-545. doi:10.1007/S11262-008-0224-2

44.   Ye K, Malinina L, Patel DJ. Recognition of small interfering RNA by a viral
suppressor of RNA silencing. *Nature*. 2003;426(6968):874.
doi:10.1038/NATURE02213

45.   Yoshikawa M, Han YW, Fujii H, Aizawa S, Nishino T, Ishikawa M. Cooperative
recruitment of RDR6 by SGS3 and SDE5 during small interfering RNA amplification
in Arabidopsis. *Proc Natl Acad Sci U S A*. 2021;118(34):e2102885118.
doi:10.1073/PNAS.2102885118/SUPPL_FILE/PNAS.2102885118.SAPP.PDF

46.   Zerbini FM, Briddon RW, Idris A, et al. ICTV virus taxonomy profile: Geminiviridae.
J Gen Virol. 2017;98(2):131-133. doi:10.1099/JGV.0.000738/CITE/REFWORKS

47.   Zhang J, Dong J, Xu Y, Wu J. V2 protein encoded by Tomato yellow leaf curl China
virus is an RNA silencing suppressor. *Virus Res*. 2012;163(1):51-58.
doi:10.1016/J.VIRUSRES.2011.08.009

48.   Zhang X-F, Guo J, Zhang X, et al. Random Plant Viral Variants Attain Temporal
Advantages During Systemic Infections and in Turn Resist other Variants of the Same
Virus OPEN. *Nat Publ Gr*. Published online 2015. doi:10.1038/srep15346

49.   Zhang X, Du P, Lu L, et al. Contrasting effects of HC-Pro and 2b viral suppressors
from Sugarcane mosaic virus and Tomato aspermy cucumovirus on the accumulation
of siRNAs. *Virology*. 2008;374(2):351-360. doi:10.1016/J.VIROL.2007.12.045

50.   Zrachya A, Glick E, Levy Y, Arazi T, Citovsky V, Gafni Y. Suppressor of RNA
silencing encoded by Tomato yellow leaf curl virus-Israel. Published online 2006.
doi:10.1016/j.virol.2006.08.016

Figure Legends

Figure 1| Screening of BCTIV genes for VSR activity a) GFP expressing 16C *N.benthamiana*
leaves were co-infiltrated with S-PTGS inducer 5'-GFP and a single BCTIV gene for each
experiment. At 6dpi, if 5'-GFP leads to red sectors under the UV lamp, it suggests that the VSR
activity of the co-infiltrated gene is negligible. However, if the infiltrated sector remains green under
UV lamp, it shows VSR activity for the gene of interest. b) Whole plant photos of the co-infiltrated
plants were taken under UV light by a DLSR camera at 6dpi. Representative images of at least three
biological replicates are provided. c) High-resolution images of the leaves infiltrated by 5'-GFP and
C1, C2, C4, Rep, V1, V2, and V3 are obtained at 50µm/pixel using green and far-red channels of
PharosFX™ imager. Green (GFP) and far-red channels (chlorophyll) are merged d) Non-vasculature
sectors of the high-resolution leaf images are quantified based on GFP. At least three leaves and
multiple areas per leaf were quantified. One-way ANOVA statistical test shows significance at $p<0.05$
e) Top panel shows northern blot results using a radioactively labelled GFP probe. 5'-GFP serves as a
low GFP expression control and a 16C-untreated sample shows high-abundance GFP mRNA control.
rRNA is a loading control for the upper panel, obtained by agarose gel electrophoresis using ethidium
bromide staining. f) Top panel shows small RNA Northern blot using a radioactively labelled GFP
probe. The bands correspond to 21 and 24 nt long siRNAs. 5'-GFP serves as a highly abundant
siRNA control and 16C-untreated serves as a clean negative control. The lower panel is the small
RNA Northern blot by using U6 RNA labeling as a control for RNA loading. U6 RNA labelling took
place on the same membrane given in the panel above, after stripping.

Figure 2|BCTIV-Rep and BCTIV-V2 genes repress S-PTGS upon transient expression of GFP
a) S-PTGS inducing FL-GFP is co-infiltrated with C1, C2, Rep, and V2 of BCTIV on WT
N.benthamiana. Upon VSR activity, GFP-FL agroinfiltration is expected to show high GFP
fluorescence under UV light. b) Whole plant photos of the co-infiltrated plants taken by a DLSR
camera at 6dpi are shown. TBSV-P19 co-infiltration with FL-GFP is used as a positive control for

very high GFP expression and FL-GFP single infiltration serves as a low GFP expression control. c)

High-resolution images of the leaves infiltrated by GFP-FL and virus genes are obtained at

50µm/pixel using green and far-red channels of PharosFX™ imager. Green (GFP) and far-red

channels (chlorophyll) are merged using FIJI. d) Infiltrated areas in the leaves are quantified based on

GFP intensity. At least three leaves and multiple areas per leaf were quantified. One-way ANOVA

statistical test shows significance at p<0.05. e) The top panel is the Western blot result, showing the

abundance of GFP protein in the co-infiltrated sections of the very same leaves shown in c. the lower

panel is the Rubisco band in the Coomassie staining as a loading control for the Western blot. Since

the assay is performed on WT *N.benthamiana,* it allows quantification of absolute changes in GFP

level, and therefore a reliable comparison of image quantifications and protein abundance in Western

blot f) FL-GFP is co-infiltrated with p19, V2, and Rep. Representative images from at least two plants

and multiple leaves are provided. g) The GFP level quantification of the co-infiltrated sectors

normalized to the non-infiltrated sectors of the treated leaves. One-Way ANOVA test has been

conducted with a threshold of p<0.05.

Figure 3|Neither BCTIV-Rep nor BCTIV-V2 can suppress PTGS induced by GFP-IR. a) PTGS-

inducing GFP-IR is co-infiltrated with C1, C2, Rep, and V2 of BCTIV on 16C *N.benthamiana*. Upon

VSR activity, GFP-IR agroinfiltrated sectors are expected to retain high GFP fluorescence under UV

light. b) Whole plant photos of the co-infiltrated plants taken under UV light by a DLSR camera at

6dpi are shown. c) High-resolution images of the leaves infiltrated by GFP-IR and C1, C2, Rep, V2,

and p19 are obtained at 50µm/pixel using green and far-red channels d) Non-vasculature sectors of

the high-resolution leaf images are quantified based on GFP intensity. At least three leaves and

multiple areas per leaf were quantified. e) Top panel shows northern blot results using a radioactively

labelled GFP probe. GFP-IR serves as a low GFP expression control and a 16C-untreated sample

shows high-abundance GFP mRNA control. rRNA is a loading control for the upper panel, obtained

by agarose gel electrophoresis using ethidium bromide staining. f) Top panel shows small-RNA-

Northern blot using a radioactively labelled GFP probe. The bands correspond to 21 and 24 nt long

siRNAs. GFP-IR serves as a highly abundant siRNA control and 16C-untreated serves as a negative

control. The lower panel is the small RNA Northern blot by using U6 RNA labelling as a control for

RNA loading. U6 RNA labelling took place on the same membrane given in the panel above, after

stripping. g) A minimum of twelve 16C *N.benthamiana* used for GFP-IR co-infiltration with VSR

candidates are followed for until 14 dpi. The images of the whole plant are taken using a DSLR

Camera under UV h) The total number of co-infiltrated plants, remaining healthy at 14dpi is provided

on the top row, and the number of plants showing systemic silencing in at least two leaves is given the

bottom row.

Figure 4| BCTIV-Rep or -V2 does not block the short-range spreading of silencing upon IR induced PTGS. a-g) 16C plants are infiltrated with GFP-FL and GFP-IR only(a), GFP-FL, GFP-IR, and TBSV-p19(b), GFP-FL, GFP-IR, and BCTIV-V2(c), GFP-FL, GFP-IR, and BCTIV-Rep(d). The images are overlay of green and far-red channels, acquired by PharosFX™ imager at 5.25dpi. The close-up on the right highlights the red sector around the infiltration area. b-h) GFP intensity profile of a 10pixel-wide line depicted on the close-up image on the right. Y-axis gives the intensity of the GFP in gray value, X-axis is the distance (in mm) of the spot to the * end of the line given. The left end of the x-axis is the # end of the line. The red dotted line indicates the distance of the short-term spread of local silencing in mm. i) The GFP intensity measurement of infiltrated non-vasculature areas is normalized to their non-infiltrated areas in the same leaf. 16C + GFP-FL infiltrated measurements are taken from Figure 1g for comparison. The corresponding measurement number is given below the bar. One-Way ANOVA test has been conducted with a threshold of $p<0.05$.

Figure 5| Multiple domains are needed for the VSR activity of BCTIV-Rep. a) GFP expressing 16C *N.benthamiana* leaves were co-infiltrated with S-PTGS inducer 5'-GFP and a single BCTIV-Rep truncation in each experiment. At 6dpi, if 5'-GFP leads to red sectors under the UV lamp, it suggests that the VSR activity of the co-infiltrated gene is negligible. However, if the infiltrated sector remains green under UV lamp, it shows VSR activity for the gene of interest. b) Whole plant photos of the co-infiltrated plants were taken under UV light by a DLSR camera at 6dpi. Representative images of the two biological replicates are provided for each co-infiltration. c) High-resolution images of the co-infiltrated leaves are obtained at 50µm/pixel using green and far-red channels of PharosFX™ imager. Green (GFP) and far-red channels (chlorophyll) are merged d) The GFP intensity of the non-vasculature sectors of the high-resolution leaf images are normalized to the non-infiltrated side of the same leaf. At least two leaves and multiple areas per leaf were quantified. The values are plotted by the online BoxPlotR tool by Shiny Lab. One-way ANOVA statistical test shows significance at $p<0.05$.

Figure 6| Proposed model for the mode of action of BCTIV-Rep VSR activity. a) S-PTGS is triggered by the RDR6-driven dsRNA synthesis using aberrant RNA (abRNA) as a template. Both 5'-GFP and GFP-FL dsRNAs are cleaved into sRNAs by DCLs. The sRNAs are loaded onto AGO proteins for cleavage of the target GFP mRNA. In the presence of Rep, S-PTGS mechanism does not function. GFP mRNA Northern blot shows that Rep represses the cleavage of GFP mRNA by S-PTGS inducer. Moreover, sRNA Northern Blot showed that Rep also suppresses the production of sRNAs. Therefore, Rep-VSR activity should be associated with the inhibition of DCL or RDR6

685   activity b) PTGS induced by GFP-IR shows that Rep cannot block the local silencing, triggered by

686   DCL cleave of the GFP-IR, suggesting that Rep does not interfere with the DCL activity, which

687   leaves RDR6 as the only possible target of Rep. Moreover, Rep still reduces the sRNAs produced

688   GFP-IR, suggesting that the propagation of the sRNA production (positive feedback loop) by RDR6

689   activity is mitigated by Rep, pointing out the interaction between Rep and RDR6.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

a Sense-PTGS

b IR-driven PTGS

702
703
704
705
706