Genome-wide peptidoglycan profiling of *Vibrio cholerae.*

Sara B. Hernandez[1, a #], Laura Alvarez[1, #], Barbara Ritzl-Rinkenberger[1, #], Bastian

Schiffthaler[2], Alonso R. Serrano[2, b] and Felipe Cava[1, *].

 # Contributed equally

* For correspondence: felipe.cava@umu.se

Affiliations:

[1] The laboratory for Molecular Infection Medicine Sweden (MIMS), Department of

Molecular Biology, Umeå University, Umeå, Sweden.

[a] Current address: Instituto de Bioquímica Vegetal y Fotosíntesis, Consejo Superior

de Investigaciones Científicas and Universidad de Sevilla, Seville, Spain.

[2] Umeå Plant Science Centre, Department of Plant Physiology, Umeå University,

Umeå, Sweden.

[b] Current address: Codon Consulting AB, Stockholm, Sweden.

Lead contact: Felipe Cava

Running title: Genome-wide peptidoglycan profiling

Keywords: cell wall, peptidoglycan, high throughput, *Vibrio cholerae,* penicillin

binding proteins.

SUMMARY

Most bacteria cells are protected by a peptidoglycan cell wall. Defining the chemical structure of the peptidoglycan has been instrumental to characterize cell wall associated proteins and to illuminate the mode of action of cell wall-acting antibiotics. However, a major roadblock for a comprehensive understanding of peptidoglycan homeostasis has been the lack of methods to conduct large-scale, systematic studies. Here we have developed and applied an innovative high throughput peptidoglycan analytical pipeline to analyze the entire non-essential, arrayed mutant library of *Vibrio cholerae*. The unprecedented breadth of these analyses revealed that peptidoglycan homeostasis is preserved by a large percentage of the genome organized in complex networks that functionally link peptidoglycan features with genetic determinants. As an example, we discovered a novel bifunctional penicillin-binding protein in *V. cholerae*. Collectively, genome-wide peptidoglycan profiling provides a fast, easy, and unbiased method for systematic identification of the genetic determinants of peptidoglycan synthesis and remodeling.

INTRODUCTION

Most bacteria present a strong, yet elastic, cell wall surrounding the cytosolic

membrane that maintains cell turgor, determines the bacterial shape and serves as

scaffold for the attachment of other cellular components (Dufresne and Paradis-

Bleau, 2015). The main component of the bacterial cell wall is the peptidoglycan

(PG), a polymer of glycan chains of the alternating *N*-acetyl-glucosamine (NAG) and

N-acetyl-muramic acid (NAM) disaccharide that are crosslinked through stem

peptides. PG composition and structure can vary between bacterial species and also

in response to environmental changes (i.e., growth conditions) (Vollmer et al., 2008).

PG is a unique and essential component in bacteria, and hence the enzymes that

synthesize and remodel this polymer are commonly used as antibacterial targets.

Consequently, extensive research has been performed during decades to provide a

more comprehensive understanding of the genetic and environmental factors that

govern PG homeostasis (Dorr, 2021; Hernandez and Cava, 2021; Kumar et al.,

2022; Porfirio et al., 2019; Rohs and Bernhardt, 2021). Unfortunately, conventional

methods for the fine analysis of the chemical structure of the PG are tedious, time-

consuming and require a significant culture size as starting material, making them

largely incompatible with large-scale, systematic screenings.

Current knowledge about the bacterial PG chemical structure has been mainly

obtained by analytical methods proposed by Glauner *et al.* more than 30 years ago

(Glauner et al., 1988), which involves: purification of the mature PG sacculus,

removal of proteins covalently linked to the PG, digestion with muramidase (i.e.,

lysozyme) to generate soluble disaccharide peptides (muropeptides) and separation

of the muropeptides by liquid chromatography (LC) (Alvarez et al., 2016; Desmarais

et al., 2013). These analyses have significantly improved during the last decade

thanks to the use of a more advanced ultra-performance liquid chromatography

(UPLC) technology replacing high pressure liquid chromatography (HPLC) systems,

and the use of in-line LC-mass spectrometry (MS) systems for PG analysis (Alvarez

et al., 2016; Bern et al., 2017; Desmarais et al., 2013; Kuhner et al., 2014; Patel et

al., 2021). Additionally, the PG isolation protocol has also been simplified and

accelerated by replacing the use of the ultra-centrifuge by bench centrifuges, and the

reduction or removal of detergents used during sacculi isolation (Alvarez et al., 2016;

Kuhner et al., 2014). Yet, current protocols are still far from being truly high

throughput to be used in large-scale studies.

Here, we have developed a high throughput (HT) method for PG isolation and LC-

based analysis appropriate for hundreds to thousands of samples of both Gram-

negative and Gram-positive bacteria. This approach has empowered an

unprecedented breadth and detail in PG homeostasis genetic determination and PG

functional networks. As a proof of concept, we have screened the muropeptide

profiles of the non-redundant mutant library of the Gram-negative pathogen *Vibrio*

cholerae. *V. cholerae* genome-wide PG profiling revealed a much broader

interrelation between PG homeostasis and other cellular processes/pathways than

anticipated. Paradoxically, multiple mutants in cell wall genes did not show significant

changes in the chemical structure of the PG thus underscoring the existence of

functional redundancies and/or conditional determinants. Correlative analyses have

allowed to globally visualize the degree of interdependence between PG features as

well as potential compensatory mechanisms.

In-depth scrutiny of the *V. cholerae* genome-wide PG profiling uncovered *vc1321*.

Despite its annotation as a hypothetical protein and the apparent lack of phenotype,

our analysis revealed that VC1321 is a novel high molecular weight penicillin-binding

protein (PBP1V, for Penicillin Binding Protein 1 of Vibrio) that contributes to *V.*

cholerae PG amount and crosslink. Collectively, these results highlight the capability

of our innovative HT PG profiling method to perform genome-wide PG analysis to

overcome phenotypic and genomic annotation hurdles towards unveiling the full

repertoire of cell wall determinants in bacteria.

RESULTS

Development of a high throughput method for peptidoglycan isolation

The most commonly used PG isolation protocols were developed in the 60's

(Kolenbrander and Ensign, 1968; Porfirio et al., 2019). These methods capitalize on

the separation of SDS-insoluble PG (i.e., murein) from the rest of the cell

components and require an individual processing of the samples incompatible with

large-scale analyses (**Fig. 1a,b**). First, the bacterial suspension is added dropwise

into boiling SDS and stirred for at least one hour to guarantee the complete lysis.

Prior to the enzymatic treatment that renders the soluble muropeptides from the

macromolecular murein, the detergent is washed away by repeated

ultracentrifugation. To overcome these roadblocks, we explored alternative methods

compatible with HT sample processing (**Supplementary Fig. 1a**). To this end, we

boiled bacterial cells with detergent using different heat transmission systems: water

bath, in a heat block (dry) or autoclave. Autoclaving performed the best compared to

the other methods, yielding the highest PG amount from the same starting material.

Since autoclaving can be easily scaled up to 96-well plates, we selected this

approach for the first step.

Removing the SDS by series of ultracentrifugation steps is time consuming. Further,

since a considerable amount of the sample is lost it requires sufficient cells to

produce good quality PG analysis. Traditionally, cultures ranging from 10-100 ml (ca.

10^{11} cells for *E. coli*) are typically used as starting material. This is an obvious

bottleneck that limits not just the scale and throughput of the studies, but also the

type of samples that can be successfully processed by this protocol, i.e., low-

concentrated environmental or *in vivo* samples. As purified sacculus retains the

shape of the cell (Formanek and Formanek, 1970) we reasoned that filtration

methods used for bacteria could also work for sacculi isolation. Using 0.2 µm pore-

size hydrophilic polypropylene (GHP) filters we washed away the SDS from sacculi

of different bacteria species (**Supplementary Fig. 1b**) without significant sample

loss, thereby allowing the analysis of smaller starting cultures. In fact, SDS-sacculi

from only 500 µl of a saturated bacterial culture ($\sim10^9$ total cells) was successfully

processed.

Since the sacculi are retained by the filters we reasoned that the subsequent

enzymatic digestion steps could be performed in-a-pot. To this end, after protease

treatment, we discarded the flow-through and the retained sacculi were subjected to

digestion with muramidase to finally elute the soluble muropeptides. Additional

modifications of the original protocol were implemented to further cut down the total

processing time: i) the final SDS concentration was lowered (from 5 to 1.5% (w/v)) to

reduce the number of washes required, ii) the time of the enzymatic reactions was

adjusted to ensure their best performance in the shortest time iii) and the reaction

buffer was substituted by water since the muramidase performs equally in both

solutions (**Supplementary Fig. 1c,d**). Finally, we also adapted for automated liquid

handling the sodium borohydride treatment that reduces NAM residues to NA-

muraminitol to avoid double peaks due to α- and β- anomeric configurations. All in

all, these modifications resulted in the development of an isolation procedure

compatible with HT methods using multiwell plates and automated pipetting systems

to process simultaneously large number of samples more than ten times faster than

the traditional methods (**Fig. 1b,c**). This novel filter-based PG isolation protocol

together with the higher speed and sensitivity of UPLC systems (Alvarez et al., 2020;

Alvarez et al., 2016) enables the production of large chromatographic datasets for a

great range of bacterial species, including both Gram-negative and Gram-positive

bacteria (**Fig. 1d**), that can be further analyzed using a variety of bioinformatic tools

(Kumar et al., 2017).

Validation of the high throughput muropeptide profiling

As proof of concept, we used the model organism *V. cholerae*. The muropeptides of

the wildtype strain were first identified by UPLC-MS (**Supplementary Fig. 2a,b**). PG

analysis of the wildtype strain and over 900 samples grown in duplicates resulted in

highly correlated relative abundances for each muropeptide (Pearson correlation

score 0.994) (**Fig. 2a**) and demonstrated the robustness and reproducibility of the

method. To assess the potential of the HT PG isolation method in discovering novel

genetic determinants involved in PG homeostasis, we first analyzed as controls well-

characterized *V. cholerae* mutants in cell wall biosynthetic genes: the bifunctional

PBP1A (Dorr et al., 2014b), the LD-transpeptidase LdtA (Cava et al., 2011) and the

DD-endopeptidase DacA1 (Moll et al., 2015), all of them known to present specific

structural changes in their PG. To facilitate comparative analysis between samples,

the muropeptide profiles (**Fig. 2b**) were transformed into numeric data. The relative

abundance for each muropeptide was calculated by integrating the area under

individual peaks divided by the total area of the whole chromatogram. The Log2 fold

change (Log2FC) was calculated for every muropeptide using the wildtype values as

control (**Supplementary Fig. 2c**). Replicas of each strain properly clustered together

and separated from the rest in a principal component analysis (PCA) (**Fig. 2c**).

Expected differences between the wildtype PG and that of the three mutant strains

were clearly highlighted: the PBP1A$^-$ mutant showed a defect in PG amount (**Fig.**

2d); LdtA$^-$ a decrease of LD-crosslinked muropeptides, and the DacA1$^-$ mutant an

increase in pentapeptide-containing muropeptides (**Fig. 2d-f**). Variations in the PG

profile of the LdtA- and DacA1- mutants, correspond to muropeptide changes of ca.

2% of the total area demonstrating that HT PG profiling is reliable to uncover subtle

PG variations.

Genome-wide peptidoglycan profiling of *V. cholerae*

To test the potential of the HT PG isolation method in large-scale PG screenings we

processed the whole transposon-mutant library of *V. cholerae* (Cameron et al., 2008)

(**Fig. 3a**). After discarding inconclusive samples (e.g., mutants that fail to grow), we

analysed the PG profile of 3,030 mutants (96% of the library) which altogether

generated a comprehensive quantitative dataset for each muropeptide and the main

PG-features (**Extended Data 1**).

The comparison of the standard deviation (SD) of the relative abundance of each

muropeptide in the library dataset with the variation between replicas demonstrated

that, except for peaks of very low abundance (i.e., D43 or T344), technical variability

is lower than that between biological samples (i.e., mutants), thus supporting the

robustness of our method to identify mutants with altered PG (**Supplementary Fig.**

3a). Additionally, we studied the variability of each muropeptide with respect to their

relative abundance (**Supplementary Fig. 3b**). Most of the muropeptides adjusted to

a linear regression model, except for the M2 and D34 peaks, which show a higher

variation degree, indicating that homeostasis of these muropeptides in *V. cholerae* is

safeguarded by a high number of genetic determinants.

Relative abundance is not a reliable metric for sample discrimination (**Fig. 3b and**

Supplementary Fig. 3c-d). The scree plot and biplot show that the most abundant

muropeptides, M4, D44 and D44^{Anh}, are the major contributors to the sample

variability while fluctuations in less abundant peaks have a more limited impact. On

the contrary, Log2FC provided a deeper statistical analysis where the contribution of

all peaks had similar weight to define sample's variability (**Supplementary Fig. 3e**).

Hierarchical clustering of the Log2FC values was used to identify sets of genes with

similar PG profiles that may belong to the same biological pathway (**Fig. 3c**). We

focused on the main PG features: PG amount, type and degree of crosslink, and

length of the PG chains. As expected, mutants in genes organized in the same

operon (based on multiple RNA-seq datasets) clustered together based on their PG

profile (**Fig. 3d-f and Extended Data 2**). As an example, mutation of the loci *vc0975*

and *vc0976* results in a significant decrease in the relative amount of PG and an

increase in LD-crosslink (**Fig. 3d**). These genes encode two conserved predicted

proteases (Teoh et al., 2015) whose function has not been described yet. Similar

changes in the PG were observed for the *vc0377* and *vc0378* mutants (**Fig. 3e**),

encoding a CheX-like protein proposed to be involved in motility regulation (Altinoglu

et al., 2022) and the zinc uptake regulator Zur (Sheng et al., 2015) respectively,

which influences PG endopeptidase activity in *V. cholerae* (Murphy et al., 2019).

Conversely, mutation of *vc0694* or *vc0693* presented a significant increase of LD-

crosslink (**Fig. 3f**). These encode an uncharacterized two component system that

has been previously related to intestinal colonization in infant mice (Cheng et al.,

2015).

The Log2FC-based PCA for all 20 *V. cholerae* muropeptides revealed outliers

representing those mutants with the most defective PGs at a general level (**Fig. 3g**).

Even when the PCA is a multiparametric spatial classification of the samples, these

outliers grouped in 6 major clusters (**Fig. 3g-h and Extended Data 3**) that shared

variability in specific muropeptides (e.g., M2, M4^M and D45 muropeptides). The 50

outlier candidates presenting the highest PCA scores included genes of a wide

variety of COG (**SI3**) suggesting that multiple cellular pathways preserve bacterial

PG homeostasis. As an example, we identified a mutant in *vc0245* (**Fig. 3h**), a gene

of unknown function but predicted as an LPS O-antigen polymerase (Manning et al.,

1995). This result is in agreement with recent reports pointing the cooperative

crosstalk between PG and LPS to maintain envelope integrity (Boll et al., 2016;

Goodall et al., 2021; Simpson et al., 2021).

Genome-wide peptidoglycan profiling reveals correlation between different

peptidoglycan features in *V. cholerae*.

Multivariate analysis (i.e., PCA) has low resolution to identify genetic determinants of

PG homeostasis because while it identifies major general alterations of cell wall's

chemical structure, it fails to discriminate more subtle changes. To solve this

bottleneck, we performed univariate data analyses of single PG-features and

observed that mutants in known PG-related loci appeared as outliers in the

corresponding analyses (**Fig. 4a**). For example, mutation of genes involved in PG

synthesis are typically characterized by a defective (sedimentable) PG amount.

Accordingly, both the mutant in the PG synthase PBP1A (encoded by *mrcA*), and its

cognate regulator CsiV (Dorr et al., 2014a) show up as outliers with low PG amount.

Further, the mutant in *vc1718*, *V. cholerae* homolog of *E. coli*'s envelope biogenesis

factor ElyC also displayed low PG amount, in agreement with a previous report

(Paradis-Bleau et al., 2014).

Additional examples can be found in relation to PG chain length. PG strands

terminate in anhydro NAM (Anh) caps generated by the action of lytic

transglycosylases (Irazoki et al., 2019). Therefore, quantification of the

anhydromuropeptide levels is a proxy to calculate average chain length: high

anhydromuropeptide level correlates with shorter PG chains and is indicative of high

autolytic activities, while low anhydromuropeptide content is indicative of longer

chains typically associated with defects in PG turnover pathways. As expected, most

of the mutants in genes encoding lytic transglycosylases presented longer PG chains

(**Fig. 4a**). These results are in agreement with recent studies that demonstrated non-

redundant roles for some of these lytic transglycosylases in *V. cholerae* (Weaver et

al., 2022; Weaver et al., 2019).

The type and degree of the PG crosslinking is another critical aspect of the

functionality of the cell wall. *V. cholerae* uses two types of crosslinks: the DD and LD

type. DD-crosslinks (AKA 4-3 type) are catalyzed by the DD-transpeptidase activity

of certain PBPs such as the bifunctional high molecular weight class A PBPs or the

monofunctional class B PBPs. Alternatively, the LD-crosslinks (so called 3-3 type)

depend on the activity of LD-transpeptidases (Aliashkevich and Cava, 2021). As for

most bacteria, *V. cholerae* primary crosslinking type is DD. This is clearly illustrated

in **Fig. 4b** by the direct correlation between DD-crosslink and total crosslink values

across mutants. Consistently, mutations in *mrcA* (the bifunctional PBP1A) or its

activators *lpoA* or *csiV* resulted in both decreased PG levels and crosslinkage. As

mutually inclusive features, PG amount and DD-crosslinking share most of their

outliers. However, there are also a number of exceptions (e.g., *elyC* mutant in PG

amount but not in crosslinking) thereby suggesting the existence of highly complex

functional networks governing PG homeostasis.

In relation with LD-crosslinkage, our data show that the mutant lacking the LD-

transpeptidase A (LdtA) and its regulator RpoS are deficient in this crosslinkage type

(Cava et al., 2011). Also consistent with recent reports (Hernandez et al., 2020), the

mutant in the recycling LD-carboxypeptidase LdcV shows elevated LD-crosslink.

Interestingly, a mutant in *vc1834*, which encodes the homolog of *E. coli's* coordinator

of PG synthesis and outer membrane constriction CpoB (Gray et al., 2015), also

showed high levels of LD-crosslink. In *E. coli* CpoB is associated to PBP1B, its main

PG-synthetase, but in *V. cholerae* PBP1A plays a more prominent role in cell wall

synthesis than PBP1B and deletion of PBP1B in *V. cholerae* has not produced any

PG phenotype (Dorr et al., 2014b). In *V. cholerae* the two bifunctional PBPs PBP1A

and PBP1B have been shown to play distinct roles than in *E. coli,* the PG-phenotype

observed here for the CpoB homolog mutant also suggests a different role for this

PG-synthesis coordinator in *V. cholerae,* compared to *E. coli.*

Finally, our genome-wide PG analysis also revealed a clear inverse correlation

between chain length and total crosslink (**Fig. 4b**). This is consistent with previous

observations (Quintela et al., 1995) supporting that a higher number of peptide-

crosslinks is required to maintain the mesh integrity of sacculi consisting of shorter

glycan chains in average. Collectively these data show that univariate analysis

complements the discovery potential of the genome-wide PG profiling.

The normal distribution of the data (**Fig. 4b**) classified as outliers those samples that

deviate 1, 2 or 3 SD for at least one muropeptide or PG feature. These data revealed

an unexpectedly high genomic contribution to PG homeostasis (**Supplementary Fig.**

3g) that spans beyond traditionally implicated cell wall enzymes and regulators.

Interestingly, a Venn diagram representing the outliers that deviate more than 1SD

revealed that more than half of the identified mutants (57.4%) displays changes in

more than one PG feature. In this line, the aforementioned contribution of DD-

crosslink to the total crosslink is supported by the high number of mutants with a

phenotype for both features (159), compared to the features alone (27 and 66

respectively). Collectively, genome-wide PG analysis permitted to appreciate a high

degree of genomic contribution and cell wall structural interdependence.

Identification of a new high molecular weight penicillin binding protein in _V._

cholerae

To identify biological processes related with the change of the different PG features,

we performed gene-ontology (GO) enrichment analyses using the sets of mutants

with significantly altered PG features (i.e., low or high PG amount, anhydro

muropeptides, and different kind of crosslink) **(Supplementary Fig. 4 and Extended**

Data 4). We focused on the samples with lower PG amount, since determinants of

PG synthesis are a common target for antibiotics. The biological processes

"peptidoglycan metabolism" and "peptidoglycan-based cell wall biogenesis" appeared

significantly enriched (p-val <0.05) (**Fig. 5a**). In addition to genes encoding known

PG-related proteins (e.g., the synthetases PBP1A and PBP1B, or the PG

degradative enzymes MltA, MltF and DacA2) (Alvarez et al., 2021), we found

_vc1321_, which was annotated as a hypothetical protein in the KEGG database (**Fig.**

5b). We constructed the deletion mutant _Δvc1321_ and its complemented strain

derivative. Analysis of the PG confirmed that the disruption of this gene not only

impairs the PG amount, but also decreases the amount of total crosslink (**Fig.**

5c,d). We confirmed that VC1321 can be specifically labelled by the fluorescent

penicillin Bocillin-FL, strongly suggesting that this protein is a new type of class A

PBP and so we named it as PBP1V (for Penicillin Binding Protein 1 of Vibrio) (**Fig.**

5e). Sequence similarity and identity was higher between PBP1A and PBP1B than

between these PBPs and PBP1V (**Fig. 5f**). *In silico* analysis predicted that PBP1V

contains an N-terminal transglycosylase domain (TG) and a putative C-terminal

transpeptidase (TP) domain that is split by a region with no homology to previously

characterized class A PBPs (**Fig. 5f**). Based on sequence alignments with other high

molecular weight PBPs (Sauvage et al., 2008) (**Supplementary Fig. 5**), we selected

and mutated three residues: the E172 as the putative TG active site; and S504 and

S610 as the potential TP active serines in the SxxK and SxN conserved motifs,

respectively (**Fig. 5f**). Mutation of S504 but not of S610 impaired PBP1V binding of

Bocillin-FL suggesting that S504 is the TP catalytic serine. Interestingly mutation of

E172 also prevented Bocillin-FL binding suggesting a regulatory role of the TG

domain in PBP1V transpeptidation activity (**Fig. 5g**).

As PBP1V's molecular weight was larger than the theoretical value, we hypothesized

it could be present in a dimeric form. We noticed that the sequence of PBP1V has

two cysteine residues that could be involved in the formation of disulfide bridges

between subunits (C139 and C1013). Effectively, *V. cholerae* wildtype membrane

fraction treated with the reducing agent DTT resulted in a band shift which coincided

with the monomeric size (116 KDa). This result was further confirmed by mutating

any of these cysteines (**Fig. 5h**). PG analysis of these mutants demonstrated that the

oligomeric state of PVP1V does not affect its role in PG synthesis under the condition

tested (**Supplementary Fig. 6**).

Interestingly, unlike the PBP1A mutant, the absence of PBP1V did not impair proper

morphology or growth under tested conditions (**Fig. 5i**) reinforcing the idea that

genome-wide PG profiling is a powerful tool to identify novel genetic determinants on

PG homeostasis, some of which might lack functional annotation or play relevant

biological roles in specific conditions. Finally, while PBP1V is only conserved in *V.*

cholerae and its close relative *V. mimicus* within the genus of *Vibrio*, orthologs of this

PBP are also encoded by other species frequently associated with marine lifestyles

(**Fig. 5j**).

DISCUSSION

We have developed a filtration-based method for PG isolation adapted to 96-well

plates and automated pipetting systems that enables HT PG biochemical analyses.

We have proved that the new protocol is applicable to different bacterial species and

provides PG profiles comparable to those obtained using the traditional procedure.

The use of filters drastically reduces the initial number of bacteria required for PG

isolation by ca. 2 orders of magnitude compared to the minimum amount required by

protocols using ultracentrifugation. Therefore, in addition to enabling larger-scale

analyses, this method opens up the possibility to characterize the PG chemical

structure of low-PG challenging samples such as intracellular non-proliferating

bacteria. Moreover, as filtration prevents PG loss, the limiting factor with this method

is no longer the sample size (amount of bacteria) but rather the detection limit of the

analytical system. In this regard, the accuracy and sensitivity of current MS has been

improving during the last decades and can facilitate the muropeptide characterization

of challenging low-abundant PG samples.

Developing a HT PG isolation method has been instrumental to analyse the cell wall

of the non-redundant mutant library of *V. cholerae,* i.e., the first genome-wide PG

profiling of a bacterium. Analysis of the data confirmed the role of known PG-related

genes and established new links between previously unrelated genes with cell wall

homeostasis, including genes encoding for proteins with previously described or

unknown functions. The unprecedented width of the omic-scale PG profiling has

made possible to witness a remarkably high genomic contribution to PG

homeostasis. For instance, almost 98% of the mutants of the *V. cholerae* transposon

library present a statistically significant alteration of their PG profile (deviate at least

1SD from the average), and more than the 50% of the mutants presented a variation

higher than 2SD of the average of the whole library for at least one muropeptide or a PG-feature) that included loci from all COG categories. These results are in agreement with studies that linked the cell wall with seemingly unrelated pathways/processes (e.g.,(Bancroft et al., 2020; Campbell et al., 2021; Masser et al., 2021)). Conversely, certain mutants in cell wall-annotated genes did not show an altered PG profile suggesting the existence of functional redundancies or condition-specific activities. Future research should uncover the functional relationships within these complex genetic networks and what regulatory rewiring occurs in response to environmental changes.

Additionally, we observed that variations in PG features and muropeptides are in general mutually inclusive: i.e., a significant proportion of the mutants present changes in more than one PG feature. This result is consistent with the idea of PG structure being safeguarded by compensatory mechanisms that preserve the cell wall integrity. This balance can be explained following a "substrate-product" logic as it is for the inverse relationship between monomers and dimers or between pentapeptides and tetrapeptides, but often seems to be the result of more complex regulatory associations (e.g., crosslink vs chain length) implicating multiple variables.

A main strength of this global study is its potential to identify novel cell wall genetic determinants even when lacking a growth phenotype under the conditions studied. As an example, we discovered PBP1V, a third high molecular weight PBP in *V. cholerae* which differs from the previously characterized PBP1A and PBP1B of this bacterium and is also not homolog to PBP1C proteins found in other bacteria. Deletion of *pbp1V* did not produce any growth defect in our analyses, but it decreased the total amount and the degree of crosslink of the PG. Although only the TG domain was predicted for this protein, sequence alignment with other high

molecular weight PBPs suggested a potential TP domain consisting of the canonical

active site sequences separated by a 186 amino acids domain that was not

previously described in any bi-functional PBPs so far studied. Mutation of the S504

residue of PBP1V that forms part of a conserved S*xxK TP motif, prevented binding

of PBP1V to Bocillin FL, strongly suggesting that this is the catalytic serine of the TP

domain. Surprisingly, mutation of E172, which is not found into a conserved

EDxxFxxHxG sequence typical of TG domains, also affected binding of Bocillin FL,

suggesting that the TG activity might also influence the TP activity of PBP1V's

transpeptidation, similar as previously reported (Zijderveld et al., 1991). PBP1V was

found to form dimers by intermolecular disulfide bonds between cysteines C139 and

C1013. Dimerization of high molecular weight PBPs by disulfide bonds has been

previously reported as a mechanism of redox control of their activity (Bukowska-

Faniband and Hederstedt, 2017). However, the monomeric PBP1V retains its TP

activity suggesting that dimerization might rather affect other aspects such as

localization or protein interactions.

The HT method presented here for PG isolation and genome-wide PG profiling

represents a promising tool for discovering new genetic determinants of PG

biosynthesis and remodeling (i.e., PBP1V) and their genetic networks, the functional

relations between the PG features and the structural constraints that define a

"healthy" PG sacculus. By profiling mutant libraries of diverse bacterial species

(including some of the most important bacterial pathogens), the publicly available

experimental platforms and databases generated will have an even wider impact due

to their utility to uncover general and species-specific mechanisms underpinning PG

homeostasis. Additionally, by using diverse growth conditions we can learn how

bacteria adapt their cell wall to relevant environments (e.g. the host) and which

constraints cannot be surpassed in the PG structure to preserve viability and shape.

Interlinking this PG-biochemical information with phenotypic analyses of the same

mutant libraries (Nichols et al., 2011) will permit to create highly informative PG

profile-phenotype-gene clusters to better understand the biological consequences of

PG synthesis and remodeling. Finally, since the cell wall is one of the major "Achilles

heels" of bacteria, this systematic survey of new players in the synthesis and

regulation of the PG structure will be particularly relevant on antibiotic resistant

pathogenic bacteria and infection-mimicking conditions as it has a great potential to

promote the discovery of new pathways that can be targeted in antimicrobial

therapies.

ACKNOWLEDGEMENTS

Research in the Cava lab is supported by The Swedish Research Council (VR), The
Knut and Alice Wallenberg Foundation (KAW), The Laboratory of Molecular Infection
Medicine Sweden (MIMS) and The Kempe Foundation. We thank John J. Mekalanos
for providing the *V. cholerae* transposon mutant library and the Cava Lab and Miguel
A. de Pedro for their helpful discussions.

AUTHOR CONTRIBUTION

SBH, LAM and FC designed the experiments. SBH, LAM and BRR performed
research. SBH, LAM, BRR, BS and AS analysed data. SBH, LAM and FC wrote the
paper with input from the rest of the authors.

COMPETING INTEREST

None

  FIGURE LEGENDS

Fig. 1. Development of a high throughput method for peptidoglycan sample

preparation and analysis.

(a) Schematic of the comparison between the traditional and the novel high

throughput (HT) method for PG isolation and examples of the potential types of

analyses that can be carried out with large-scale muropeptide profile datasets. **(b)**

Comparative pipeline for the HT and the traditional PG isolation method. **(c)**

Comparative timescales for the isolation of PG from 96 samples following the

traditional or the HT sample preparation method. **(d)** Representative PG profiles

obtained for diverse bacteria using the HT PG isolation method. **See also**

Supplementary Fig. 1.

Fig. 2. Validation of the high throughput method for peptidoglycan sample

preparation.

(a) Pairwise scatter plot comparing the relative abundance of every muropeptide

between replicates of 900 samples. Mean Pearson correlation coefficient is shown.

(b-f) Validation of the HT method for detection of minor changes in the PG structure

by analysis of mutants known to have a phenotype in their muropeptide profile:

Pbp1A (PG synthetase), LdtA (LD-transpeptidase) and DacA1 (DD-

carboxypeptidase); **(b)** representative chromatogram of each strain; **(c)** PCA using

the Log2FC of peak areas of all the muropeptides of each sample; **(d)** analysis of PG

features; **(e)** heatmap presenting the relative peak area of each muropeptide

numbered in panel **b**; **(f)** heatmap of the Log2FC values (relative to the wildtype)

calculated for each peak numbered in **b.** In **e** and **f**, peaks expected to change in

LdtA⁻ and DacA1⁻ mutants are highlighted in green and blue respectively. Error bars

in **d** correspond to the standard deviation of triplicates. **See also Supplementary**

Fig. 2.

Fig. 3. Peptidoglycan profiling of the *V. cholerae* non-redundant mutant library.

(a) Total number of mutant strains available, processed and analyzed, and their

classification into main COG groups. **(b)** Muropeptide abundance and variation (SD)

across the whole dataset. **(c)** Hierarchical clustering and heat-map representing the
Log2FC calculated for different PG-features and muropeptides. **(d-f)** Analysis of the
PG profile of genes belonging to the same transcriptional unit: PG features, heatmap
and clustering of the PG profiles of indicated mutants and the expression data of
each gene observed in 307 different RNAseq experiments are shown. **(g)** PCA
obtained for the mutants of the library based on the Log2FC values observed for the
20 identified muropeptides. Outliers are colour-coded based on their PG-similarities
clustering (see **h**). **(h)** Hierarchical clustering of the 50 outliers selected in **g. See**
also Supplementary Fig. 3 and Extended Data 1-3.

Fig. 4. Analysis of the main peptidoglycan features of the *V. cholerae* mutant
library.
(a) Violin plots showing the distribution of samples for the indicated PG-features in
the 3030 analyzed mutants. Mutants expected to present a distinctive feature are
labelled in each group. **(b)** Correlation scatterplot matrix for the different PG features
in the *V. cholerae* mutant library. Histograms showing the distribution of samples for
every PG feature are represented in the diagonal.

Fig. 5. *vc1321* encodes a new high molecular weight PBP in *V. cholerae*.
(a) Over-represented biological processes associated to those mutants presenting low
relative amount of PG (Log2FC <-0.5). The most representative biological processes with a
fold enrichment higher than 2 are represented. Dot size indicates the number of loci/category
and colour indicates the significance of the enrichment (-Log10(*P*-values)). **(b)** Plot showing
the Log2FC of the PG amount for the complete dataset. Genes belonging to the
"peptidoglycan metabolism" category in **a** are labelled. **(c)** Representative PG
chromatograms of the wildtype, the *Δvc1321* mutant strain and the *Δvc1321* strain
overexpressing *vc1321* from the arabinose-inducible promoter of the pBAD plasmid. **(d)**
Quantification of the relative PG amount and DD-crosslink of the wildtype, *Δvc1321* mutant
strain, and the *Δvc1321* carrying the pBAD empty plasmid as control or overexpressing
vc1321 from the arabinose-inducible promoter. **(e)** Bocillin gel of indicated strains showing
the band corresponding to the product of *vc1321* (named PBP1 of *V. cholerae*, PBP1V) **(f)**
Schematic representation of the transglycosylase (TG) and transpeptidase (TP) domains
predicted in the sequence of VC1321. **(g-h)** Bocillin gels of *vc1321* point mutants in different

putative catalytic residues **(h)** and in candidate cysteines involved in the dimerization of the

protein. **(i)** Deletion of *pbp1V* does not result in growth or morphological defect. Scale bar = 2

µm. **(j)** PBP1V homologs across different species. **See also Supplementary Figs. 4-6 and**

Extended Data 4.

SUPPLEMENTARY FIGURES TITLES AND LEGENDS

Supplementary Fig. 1. High throughput method optimization, Related to Fig. 1.
(a) Comparison of the efficiency of PG isolation methods. Samples corresponding to different cell numbers were collected, supplemented with SDS 1.5% (w/v) final concentration, and solubilized by boiling, autoclaving or using a heat-block during different times. Samples were processed and analyzed by UPLC. Yield is quantified as maximum intensity of absorbance at 204 nm for the M4 muropeptide, in arbitrary units (AU). **(b)** Phase contrast microscopy of different bacteria and their purified sacculi processed using the traditional or the HT purification method. Scale bar = 2 μm. **(c)** Relative muramidase activity at different time points. **(d)** Efficiency of the muramidase digestion in buffer or water on sacculi from two different concentrated starting cultures at two time points. Yield is quantified as maximum intensity of absorbance at 204 nm for the M4 muropeptide, in arbitrary units (AU).

Supplementary Fig. 2. Muropeptide profile of *V. cholerae*'s peptidoglycan obtained using the high throughput sample preparation method and analysis of data, Related to Figs. 2-3.
(a) Representative chromatogram obtained for *V. cholerae* wildtype strain using the HT sample preparation method. Peaks of interest are listed. **(b)** Table of identified muropeptides in **a**. Identity was confirmed by MS/MS analysis, theoretical and observed neutral mass in Da are indicated. GlcNAc: N-acetyl glucosamine; MurNAc: N-acetyl muramic acid; (1-6anhydro) MurNAc: 1-6 anhydro N-acetyl muramic acid, terminal muropeptide; L-Ala: L-alanine; D-Glu: D-glutamic acid; DAP: *meso*-diaminopimelic acid; D-Ala: D-alanine; D-Met: D-methionine; Gly: glycine; Lpp: Braun's lipoprotein. **(c)** Pipeline for the data transformation and analysis: i) abundance was calculated by integration of the area of the peaks, ii) relative area of each muropeptide was calculated by dividing the peak area by the total area of the chromatogram, iii) finally, fold change (FC) relative to the control sample was calculated and Log2FC was used for representation in heat maps.

Supplementary Fig. 3. Analysis of the peptidoglycan profiling of the *V. cholerae* non-redundant mutant library, Related to Fig. 3.
(a) Biological versus technical variability: comparison of the standard deviation of the relative abundance of each muropeptide across all samples in the *V. cholerae* mutant library versus the average standard deviation across replicas of 300 samples. **(b)** Correlation between each muropeptide relative abundance and its standard deviation across the library dataset. **(c)** Scree plot of the main components used for the PCA using relative abundance of

muropeptides and Log2FC calculated values. **(d)** Biplot showing the contribution of the different variables (muropeptides) to the data variability in the dataset of relative amounts**. (e)** Biplot showing the contribution of the different variables (muropeptides) to the data variability in the Log2FC dataset. **(f)** Violin plots showing the distribution of the samples for each muropeptide. Log2FC data was used. Median (continued lines) and quartiles (dashed lines) are represented. **(g)** Table representing the number and percentage of mutants with a significant phenotype for at least one muropeptide or a PG feature. A significant phenotype is considered when the sample is further than 1, 2 or 3 standard deviations (SD) from the average. **(h)** Venn diagram showing the number of outliers with a phenotype (using 1SD as threshold) in the main PG features.

Supplementary Fig. 4. Enrichment analysis of mutants presenting differential PG features, Related to Fig. 5.
Over-represented biological processes of mutant candidates presenting low (left, red dots) and high (right, blue dots) relative amounts of: **(a)** chain length (Log2FC <-0.2 and >0.18); **(b)** total crosslink (Log2FC <-0.15 and >0.15); **(c)** DD-crosslink (Log2FC <-0.2 and >0.2); and **(d)** LD-crosslink (Log2FC <-0.7 and >0.7). The most representative biological processes with a fold enrichment higher than 2 are represented. The dot size in the balloon plots indicates the number of loci included in each category and the colour the significance of the enrichment (-Log10(p-values)).

Supplementary Fig. 5. Multiple sequence alignment of PBP1V with other high molecular weight penicillin-binding proteins, Related to Fig. 5.
Sequence alignment of *V. cholerae* PBP1V (Vc-PBP1V, Uniprot: Q9KSD7), *V. cholerae* PBP1A (Vc-PBP1A, Uniprot: Q9KNU5), *V. cholerae* PBP1B (Vc-PBP1B, Uniprot: Q9KUC0), *E. coli* PBP1A (Ec-PBP1A, Uniprot: P02918), *E. coli* PBP1B (Ec-PBP1B, Uniprot: P02919) and *E. coli* PBP1C (Ec-PBP1C, Uniprot: A0A093EN65), performed with T-COFFEE Expresso. Background residue colour indicates degree of conservation. TG domain is highlighted in yellow, TP domain is highlighted in light blue, PBP1C-binding domain is highlighted in green. PBPs conserved motifs are indicated in the red boxes. Vc-PBP1V new domain in red letters.

Supplementary Fig. 6. Peptidoglycan analysis of PBP1V cysteine point mutants, **Related to Fig. 5.**
Quantification of the relative PG amount and DD-crosslink of the wildtype, Δ*pbp1V* mutant strain, and indicated cysteine mutants. (ns: not significant).

EXTENDED DATA FILES AND LEGENDS

Extended Data 1. PG dataset from the *V. cholerae* Tn-library, **Related to Fig. 3.**

Extended Data 2. SRA numbers used for the study of gene expression analysis, **Related to Fig. 3 d-f.**

Extended Data 3. Data 50 selected PCA outliers, **Related to Fig. 3 g-h.**

Extended Data 4. GO-enrichments, **Related to Fig. 5.**

Extended Data 5. Bacterial strains list, **Related to Methods.**

Extended Data 6. List of primers, **Related to Methods.**

METHODS

- Lead Contact and Materials Availability

• All the data obtained from the genome-wide HTPGP of *V. cholerae* non-redundant mutant collection is available in the Extended Data files.

• Further information and requests for resources and data should be directed and will be fulfilled by the Lead Contact, Felipe Cava (felipe.cava@umu.se).

- Bacterial strains, media, and antibiotics

All the bacterial strains used in this study are listed in **Extended Data 5**. Lysogeny broth (LB) containing 10 or 5 mg/ml of NaCl (for *V. cholerae* or the other bacterial species respectively) was used as the standard growth medium. Agar 1.5% (w/v) was used in solid plates. Unless otherwise specified, antibiotics were used at the following concentrations: streptomycin 200 μg/ml, kanamycin 50 μg/ml.

Plasmids were constructed by standard DNA cloning techniques. *V. cholerae* mutant strains were constructed using the primers listed in **Extended Data 6,** and standard allele exchange techniques with derivatives of the suicide plasmid pCVD442 as described previously (Donnenberg and Kaper, 1991). Plasmid used for complementation are derivatives of pBAD18 (Guzman et al., 1995), where expression is controlled by the P_{BAD} promoter. The fidelity of the DNA regions generated by PCR amplification was confirmed by DNA sequencing.

- Growth conditions

Bacterial species were grown in LB at 30 or 37 ºC for sacculi isolation. The culture volume used for high throughput sacculi preparation ranged between 0.5 and 2 ml and was adjusted to obtain the best PG-profile (highest absorbance, UPLC signal)

without saturating the filters during the purification. For the analysis of the *V.*

cholerae mutant library, cells from glycerol stocks were first inoculated in 96-well

plates containing 200 µl of LB medium, grown overnight at 37 °C and further diluted

1000-fold into 500 µl of fresh medium in 96-well deep-well plates, where they were

finally grown at 37 °C with shaking overnight. For induction of the P_{BAD} promoter,

0.2% (w/v) of arabinose (final concentration) was added to bacterial cultures.

- High Throughput (HT) sacculi preparation and PG isolation and digestion

For optimization of the method, different sacculi isolation protocols were tested,

shown in **Supplementary Fig. 1**. Bacterial cultures grown overnight in 96-well deep-

well plates were pelleted in the same culture plates at 5,250 *x g* for 30 min. The

supernatant was discarded, and the pellets were resuspended with 50 µl of fresh

medium. After addition of 25 µl of 5% SDS, samples were tightly sealed using

silicone mats and tape, and autoclaved (15 min at 120 °C 1atm). Samples were then

transferred to 96-well-filter plates (AcroPrep Advance 96-well 0.2 µm GHP

membrane filter plates, 2 ml volume, PALL). SDS was removed by washing twice

with 1 ml Milli-Q water (5,250 *x g* for 15 min). SDS-free sacculi samples were treated

with 200 µl proteinase K (40 µg/ml in 100 mM TrisHCl pH 8.0, 30 min at 37°C) for

removal of the Braun's lipoprotein. After an additional 1 ml wash with water, samples

were digested with 50 µl muramidase (100 µg/ml in water, overnight at 37°C).

Enzymatic reactions were performed directly on the filter, sealing the plates with

aluminium foil and using a humidity chamber to reduce evaporation.

After muramidase digestion, soluble muropeptides were eluted into a new 96-well

deep-well plate by centrifuging the filter plate at 5,250 *x g* for 5 min. Finally, eluted

muropeptides were reduced using 10 µl borate buffer (0.5 M pH 9.0) and 25 µl

NaBH$_4$ (10 mg/ml) for 30 min at room temperature. The pH of the samples was

adjusted to 3.5 using 8 µl orthophosphoric acid 25% (v/v) and the samples were

stored frozen at -20 °C until analysis by UPLC.

- Data acquisition by liquid chromatography

PG samples were analyzed by liquid chromatography, as previously described

(Alvarez et al., 2020). Briefly, muropeptides were separated using an UPLC system

(Waters) equipped with a trapping cartridge precolumn (SecurityGuard ULTRA

Cartridge UHPLC C18 2.1 mm, Phenomenex) and an analytical column (Waters

ACQUITY UPLC BEH C18, 130Å, 1.7 µm, 2.1 mm×150 mm), using as solvents 0.1%

formic acid in Milli-Q water (buffer A) and 0.1% formic acid (v/v) in acetonitrile (buffer

B) in a 14 min run. Muropeptides were detected by measuring the absorbance at 204

nm. Identity of the peaks was first assigned by LC coupled to a QTOF-MS instrument

and then by comparison of their retention time. The QTOF–MS instrument was

operated in positive ionization mode. Detection of muropeptides was performed by

MS^E to allow for the acquisition of precursor and product ion data simultaneously,

using the following parameters: capillary voltage at 3.0 kV, source temperature to

120 ºC, desolvation temperature to 350 ºC, sample cone voltage to 40 V, cone gas

flow 100 l/h, desolvation gas flow 500 l/h and collision energy (CE): low CE: 6 eV and

high CE ramp: 15-40 eV. Mass spectra were acquired at a speed of 0.25 s/scan. The

scan was in a range of 100–2000 *m/z.* Data acquisition and processing was

performed using UNIFI software package (Waters Corp.).

For the study of the PG chemical variability between samples, collected UPLC

chromatographic data were analyzed using the PG-metrics pipeline (Kumar et al.,

2017). In brief, raw data were pre-processed by trimming out irrelevant segments

and by baseline correction subtracting a synthetically created baseline. Trimmed and

baseline corrected datasets were then aligned by segments using the COW

algorithm by selecting appropriate reference samples and combinations of the

segment length and slack parameters. The peak area of aligned chromatograms was

calculated using a MATLAB routine based on the trapezoidal integration approach

wherein the boundaries of the integration were manually selected.

The relative area of each muropeptide was calculated dividing its peak area by the

total area of the chromatogram (**Supplementary Fig. 2b**). The different PG features

were calculated as described previously (Alvarez et al., 2020; Vigouroux et al.,

2020). When relative peak areas of each mutant were used, the most abundant

muropeptides presented the greatest impact on the results of PCA analyses

(**Supplementary Fig. 3a-c**), thus the log2 fold change (Log2FC) was calculated for

more comprehensive analysis of the data. Log2FCs were calculated by dividing each

value (peak area or PG-feature) by the one of the control sample when available or

by the mean value of all samples from the same sample set (i.e., 96-well plates)

(**Supplementary Fig. 2c**).

- Data analysis and representation

Prism 8.0 (GraphPad Software) was used to plot and analyze numerical data.

Statistical tests in Prism calculated significance of measurements as reported in the

corresponding figure legends.

Statistical analyses (correlation, standard deviation, PCA, biplots, scree plots) were

carried out using R version 4.1.2.

Venn diagrams were created selecting as outliers those mutants with a PG

phenotype (more than 1, 2 or 3 standard deviations) in at least one muropeptide or

PG feature.

Heatmaps and hierarchical clustering analyses (agglomeration method: ward;

distance metric: euclidean) were performed using the NG-CHM Heat Map Builder,

version 2.20.2 (Ryan et al., 2019).

- Gene expression analysis

In order to calculate gene expression estimates, 307 RNA-Sequencing data

annotated to *V. cholerae* and performed on an Illumina sequencing machine were

downloaded from the NCBI's sequence read archive (SRA) (Leinonen et al., 2011)

(See **Extended Data 2**). We then used salmon/v1.3.0 (Patro et al., 2017) with

parameters --libType A --seqBias and --posBias to quantify expression against a

decoy-aware index generated from the GCF_000829215.1_ASM82921v1 reference

(Okada et al., 2014) as described in Srivastava et al. (Srivastava et al., 2020). Post

quantification, we filtered experiments where less than 50% of reads could be

matched to a reference gene and imported the estimated counts into R/v4.0 using

the tximport package (Soneson et al., 2015). After importing, we used the

varianceStabilizingTransformation procedure implemented in the DESeq2 (Love et

al., 2014) R package to generate a blind, homoscedastic, and pseudo-log2 count set.

- Functional enrichment analysis

The Protein ANalysis THrough Evolutionary Relationships (PANTHER) Classification

System and the PANTHER Statistical Overrepresentation Test (released 20200407)

were used to categorize the selected datasets by Gene Ontology (GO) and to

determine the overrepresented GO terms (Mi et al., 2019a; Mi et al., 2019b). For this,

selected datasets were searched using the "GO biological process complete"

annotation classification available in PANTHER to find functional classes statistically

overrepresented when compared to the reference list containing the 3030 proteins

corresponding to the Tn-mutants analysed by HT-PG profiling. GO terms were

considered significantly overrepresented only when the P-value (Fisher's exact test)

was lower than 0.05. The resulting list for each dataset was summarized (similar

terms including the same list of genes were removed to avoid redundancy) and

visually represented as described by Bonnot *et* al. (Bonnot et al., 2019).

- **Microscopy imaging**

Bacteria were immobilized on LB pads containing 1% agarose (w/v). Phase contrast

microscopy was performed using a Zeiss Axio Imager .Z2 microscope (Zeiss,

Germany) equipped with a Plan-Apochromat 63X phase contrast objective lens and

an ORCA-Flash 4.0 LT digital CMOS camera (Hamamatsu Photonics, Japan), using

the Zeiss Zen Blue software. Image analysis and processing were performed using

ImageJ software (Rasband, W.S., ImageJ, U. S. National Institutes of Health,

Bethesda, Maryland, USA, https://imagej.nih.gov/ij/, 1997-2018).

- **Membrane preparation and detection of penicillin-binding proteins**

For the preparation of membranes of *V. cholerae*, 100 ml cell cultures were grown

over night and harvested by centrifugation at 15,000 $x\,g$ for 15 min at 4 °C (Beckman

JLA-16.250 rotor, Beckman Avanti J-25, Beckman Coulter). Pellets were washed

with 20 mM potassium phosphate (pH 7.5) and 140 mM sodium chloride buffer.

Following another centrifugation for 30 min at 360 $x\,g$ at 4 °C, cells were disrupted

twice using a French press (Pressure Cell Homogenizer FPG12800, Stansted Fluid

Power Ltd). The resulting cell lysates were subjected to centrifugation at 360 x g for

10 min at 4 °C. The supernatant fractions were collected and centrifuged at 75,600 x

g for 90 min at 4 °C (Beckman JA25.50 rotor, Beckman Avanati J-25, Beckman

Coulter). The resulting pellets were washed again and resuspended in 500 µl

potassium phosphate buffer. The concentration of the membrane preparations was

measured using Bradford reagent (Bio-Rad) with bovine serum albumin as standard

(Sigma).

For the detection of the PBPs, membranes were incubated for 30 min at 37 °C with a

fluorescent labelling agent, Bocillin FL Penicillin (Invitrogen, Thermo Fisher Scientific)

at a final concentration of 0.05 mM. Finally, the samples were denatured with 4x

Laemmli sample buffer (Bio-Rad) at 95 °C for 5 min, followed by centrifugation to

remove membrane debris. 20 µl of the reaction mixture were subjected to SDS-

PAGE analysis on an 8% polyacrylamide gel. Penicillin-binding proteins were

visualised with a laser scanner, Typhoon FLA 9500 (GE Healthcare Life Sciences) at

473 nm with a 530DF20 emission filter.

- Multiple sequence alignments

Sequence from *V. cholerae* PBP1V (Vc-PBP1V, Uniprot: Q9KSD7), *V. cholerae*

PBP1A (Vc-PBP1A, Uniprot: Q9KNU5), *V. cholerae* PBP1B (Vc-PBP1B, Uniprot:

Q9KUC0), *E. coli* PBP1A (Ec-PBP1A, Uniprot: P02918), *E. coli* PBP1B (Ec-PBP1B,

Uniprot: P02919) and *E. coli* PBP1C (Ec-PBP1C, Uniprot: A0A093EN65) were

aligned using T-COFFEE Expresso Version 11.00 (Notredame et al., 2000) and

visualized with JalView (Waterhouse et al., 2009).

We run BLAST to study the homology between *V. cholerae* HMW-PBPs (Altschul et

al., 1997).

- Protein structure prediction

PBP1V protein 3D structural model was built with Alphafold, using both monomer

and multimer modes (Evans et al., 2022; Jumper et al., 2021). Only the best model

among the five best given by default was examined in detail and represented in the

figures. Molecular graphics and analyses were performed with ChimeraX (Goddard

et al., 2018; Pettersen et al., 2021).

REFERENCES

Aliashkevich, A., and Cava, F. (2021). LD-transpeptidases: the great unknown among the peptidoglycan cross-linkers. The FEBS journal.

Altinoglu, I., Abriat, G., Carreaux, A., Torres-Sanchez, L., Poidevin, M., Krasteva, P.V., and Yamaichi, Y. (2022). Analysis of HubP-dependent cell pole protein targeting in Vibrio cholerae uncovers novel motility regulators. PLoS Genet *18*, e1009991.

Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res *25*, 3389-3402.

Alvarez, L., Cordier, B., van Teeffelen, S., and Cava, F. (2020). Analysis of Gram-negative bacteria peptidoglycan by Ultra-Performance Liquid Chromatography. Bio-protocol *10*, e3780.

Alvarez, L., Hernandez, S.B., and Cava, F. (2021). Cell Wall Biology of Vibrio cholerae. Annual review of microbiology *75*, 151-174.

Alvarez, L., Hernandez, S.B., de Pedro, M.A., and Cava, F. (2016). Ultra-sensitive, high-resolution liquid chromatography methods for the high-throughput quantitative analysis of bacterial cell wall chemistry and structure. Methods in molecular biology *1440*, 11-27.

Bancroft, P.J., Turapov, O., Jagatia, H., Arnvig, K.B., Mukamolova, G.V., and Green, J. (2020). Coupling of peptidoglycan synthesis to central metabolism in Mycobacteria: post-transcriptional control of CwlM by aconitase. Cell Rep *32*, 108209.

Bern, M., Beniston, R., and Mesnage, S. (2017). Towards an automated analysis of bacterial peptidoglycan structure. Anal Bioanal Chem *409*, 551-560.

Boll, J.M., Crofts, A.A., Peters, K., Cattoir, V., Vollmer, W., Davies, B.W., and Trent, M.S. (2016). A penicillin-binding protein inhibits selection of colistin-resistant, lipooligosaccharide-deficient Acinetobacter baumannii. Proc Natl Acad Sci U S A *113*, E6228-e6237.

Bonnot, T., Gillard, M.B., and Nagel, D.H. (2019). A simple protocol for informative visualization of enriched gene ontology terms. Bio-protocol *9*, e3429.

Bukowska-Faniband, E., and Hederstedt, L. (2017). Transpeptidase activity of penicillin-binding protein SpoVD in peptidoglycan synthesis conditionally depends on the disulfide reductase StoA. Molecular microbiology *105*, 98-114.

Cameron, D.E., Urbach, J.M., and Mekalanos, J.J. (2008). A defined transposon mutant library and its use in identifying motility genes in Vibrio cholerae. Proc Natl Acad Sci U S A *105*, 8736-8741.

Campbell, C., Fingleton, C., Zeden, M.S., Bueno, E., Gallagher, L.A., Shinde, D.,
Ahn, J., Olson, H.M., Fillmore, T.L., Adkins, J.N.*, et al.* (2021). Accumulation of
succinyl coenzyme A perturbs the Methicillin-Resistant Staphylococcus aureus
(MRSA) succinylome and is associated with increased susceptibility to beta-lactam
antibiotics. mBio *12*, e0053021.

Cava, F., de Pedro, M.A., Lam, H., Davis, B.M., and Waldor, M.K. (2011). Distinct
pathways for modification of the bacterial cell wall by non-canonical D-amino acids.
EMBO J *30*, 3442-3453.

Cheng, A.T., Ottemann, K.M., and Yildiz, F.H. (2015). Vibrio cholerae response
regulator VxrB controls colonization and regulates the Type VI Secretion System.
PLoS Pathog *11*, e1004933.

Desmarais, S.M., De Pedro, M.A., Cava, F., and Huang, K.C. (2013). Peptidoglycan
at its peaks: how chromatographic analyses can reveal bacterial cell wall structure
and assembly. Molecular microbiology *89*, 1-13.

Donnenberg, M.S., and Kaper, J.B. (1991). Construction of an eae deletion mutant of
enteropathogenic Escherichia coli by using a positive-selection suicide vector.
Infection and immunity *59*, 4310-4317.

Dorr, T. (2021). Understanding tolerance to cell wall-active antibiotics. Ann N Y Acad
Sci *1496*, 35-58.

Dorr, T., Lam, H., Alvarez, L., Cava, F., Davis, B.M., and Waldor, M.K. (2014a). A
novel peptidoglycan binding protein crucial for PBP1A-mediated cell wall biogenesis
in Vibrio cholerae. PLoS Genet *10*, e1004433.

Dorr, T., Moll, A., Chao, M.C., Cava, F., Lam, H., Davis, B.M., and Waldor, M.K.
(2014b). Differential requirement for PBP1a and PBP1b in in vivo and in vitro fitness
of Vibrio cholerae. Infect Immun *82*, 2115-2124.

Dufresne, K., and Paradis-Bleau, C. (2015). Biology and assembly of the bacterial
envelope. Advances in experimental medicine and biology *883*, 41-76.

Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A.,
Bates, R., Blackwell, S., Yim, J.*, et al.* (2022). Protein complex prediction with
AlphaFold-Multimer. 2021.2010.2004.463034.

Formanek, H., and Formanek, S. (1970). Specific staining for electron microscopy of
murein sacculi of bacterial cell walls. European journal of biochemistry *17*, 78-84.

Glauner, B., Holtje, J.V., and Schwarz, U. (1988). The composition of the murein of
Escherichia coli. The Journal of biological chemistry *263*, 10088-10095.

Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H.,
and Ferrin, T.E. (2018). UCSF ChimeraX: Meeting modern challenges in
visualization and analysis. Protein science : a publication of the Protein Society *27*,
14-25.

Goodall, E.C.A., Isom, G.L., Rooke, J.L., Icke, C., Pullela, K., Zhang, B., Rae, J.,
Tan, W.B., Winkle, M., Delhaye, A.*, et al.* (2021). YhcB coordinates peptidoglycan
and LPS biogenesis with phospholipid synthesis during Escherichia coli
cell growth. 2021.2004.2016.440158.

Gray, A.N., Egan, A.J., Van't Veer, I.L., Verheul, J., Colavin, A., Koumoutsi, A.,
Biboy, J., Altelaar, A.F., Damen, M.J., Huang, K.C.*, et al.* (2015). Coordination of
peptidoglycan synthesis and outer membrane constriction during Escherichia coli cell
division. Elife *4*.

Guzman, L.M., Belin, D., Carson, M.J., and Beckwith, J. (1995). Tight regulation,
modulation, and high-level expression by vectors containing the arabinose PBAD
promoter. Journal of bacteriology *177*, 4121-4130.

Hernandez, S.B., and Cava, F. (2021). New approaches and techniques for bacterial
cell wall analysis. Curr Opin Microbiol *60*, 88-95.

Hernandez, S.B., Dorr, T., Waldor, M.K., and Cava, F. (2020). Modulation of
peptidoglycan synthesis by recycled cell wall tetrapeptides. Cell Rep *31*, 107578.

Irazoki, O., Hernandez, S.B., and Cava, F. (2019). Peptidoglycan Muropeptides:
Release, Perception, and Functions as Signaling Molecules. Front Microbiol *10*, 500.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A.*, et al.* (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.

Kolenbrander, P.E., and Ensign, J.C. (1968). Isolation and chemical structure of the
peptidoglycan of Spirillum serpens cell walls. Journal of bacteriology *95*, 201-210.

Kuhner, D., Stahl, M., Demircioglu, D.D., and Bertsche, U. (2014). From cells to
muropeptide structures in 24 h: peptidoglycan mapping by UPLC-MS. Sci Rep *4*,
7494.

Kumar, K., Espaillat, A., and Cava, F. (2017). PG-Metrics: A chemometric-based
approach for classifying bacterial peptidoglycan data sets and uncovering their
subjacent chemical variability. PLoS One *12*, e0186197.

Kumar, S., Mollo, A., Kahne, D., and Ruiz, N. (2022). The bacterial cell wall: From
lipid II flipping to polymerization. Chem Rev *122*, 8884-8910.

Leinonen, R., Sugawara, H., Shumway, M., and International Nucleotide Sequence
Database, C. (2011). The sequence read archive. Nucleic Acids Res *39*, D19-21.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Manning, P.A., Stroeher, U.H., Karageorgos, L.E., and Morona, R. (1995). Putative
O-antigen transport genes within the rfb region of Vibrio cholerae O1 are
homologous to those for capsule transport. Gene *158*, 1-7.

Masser, E.A., Burby, P.E., Hawkins, W.D., Gustafson, B.R., Lenhart, J.S., and
Simmons, L.A. (2021). DNA damage checkpoint activation affects peptidoglycan
synthesis and late divisome components in Bacillus subtilis. Molecular microbiology
116, 707-722.

Mi, H., Muruganujan, A., Ebert, D., Huang, X., and Thomas, P.D. (2019a). PANTHER
version 14: more genomes, a new PANTHER GO-slim and improvements in
enrichment analysis tools. Nucleic Acids Res *47*, D419-D426.

Mi, H., Muruganujan, A., Huang, X., Ebert, D., Mills, C., Guo, X., and Thomas, P.D.
(2019b). Protocol Update for large-scale genome and gene function analysis with the
PANTHER classification system (v.14.0). Nat Protoc *14*, 703-721.

Moll, A., Dorr, T., Alvarez, L., Davis, B.M., Cava, F., and Waldor, M.K. (2015). A D,
D-carboxypeptidase is required for Vibrio cholerae halotolerance. Environ Microbiol
17, 527-540.

Murphy, S.G., Alvarez, L., Adams, M.C., Liu, S., Chappie, J.S., Cava, F., and Dörr, T.
(2019). Endopeptidase regulation as a novel function of the Zur-dependent Zinc
starvation response. mBio *10*.

Nichols, R.J., Sen, S., Choo, Y.J., Beltrao, P., Zietek, M., Chaba, R., Lee, S.,
Kazmierczak, K.M., Lee, K.J., Wong, A.*, et al.* (2011). Phenotypic landscape of a
bacterial cell. Cell *144*, 143-156.

Notredame, C., Higgins, D.G., and Heringa, J. (2000). T-Coffee: A novel method for
fast and accurate multiple sequence alignment. Journal of molecular biology *302*,
205-217.

Okada, K., Na-Ubol, M., Natakuathung, W., Roobthaisong, A., Maruyama, F.,
Nakagawa, I., Chantaroj, S., and Hamada, S. (2014). Comparative genomic
characterization of a Thailand–Myanmar isolate, MS6, of Vibrio cholerae O1 El Tor,
which is phylogenetically related to a "US gulf coast" clone. PLOS ONE *9*, e98120.

Paradis-Bleau, C., Kritikos, G., Orlova, K., Typas, A., and Bernhardt, T.G. (2014). A
genome-wide screen for bacterial envelope biogenesis mutants identifies a novel
factor involved in cell wall precursor metabolism. PLoS Genet *10*, e1004056.

Patel, A.V., Turner, R.D., Rifflet, A., Acosta-Martin, A.E., Nichols, A., Awad, M.M.,
Lyras, D., Gomperts Boneca, I., Bern, M., Collins, M.O.*, et al.* (2021). PGFinder, a
novel analysis pipeline for the consistent, reproducible, and high-resolution structural
analysis of bacterial peptidoglycans. Elife *10*.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon
provides fast and bias-aware quantification of transcript expression. Nat Methods *14*,
417-419.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I.,
Morris, J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for
researchers, educators, and developers. Protein science : a publication of the Protein
Society *30*, 70-82.

Porfirio, S., Carlson, R.W., and Azadi, P. (2019). Elucidating peptidoglycan structure:
An analytical toolset. Trends in microbiology *27*, 607-622.

Quintela, J.C., Caparros, M., and de Pedro, M.A. (1995). Variability of peptidoglycan
structural parameters in gram-negative bacteria. FEMS Microbiol Lett *125*, 95-100.

Rohs, P.D.A., and Bernhardt, T.G. (2021). Growth and Division of the Peptidoglycan
Matrix. Annual review of microbiology *75*, 315-336.

Ryan, M.C., Stucky, M., Wakefield, C., Melott, J.M., Akbani, R., Weinstein, J.N., and
Broom, B.M. (2019). Interactive Clustered Heat Map Builder: An easy web-based tool
for creating sophisticated clustered heat maps. F1000Res *8*.

Sauvage, E., Kerff, F., Terrak, M., Ayala, J.A., and Charlier, P. (2008). The penicillin-
binding proteins: structure and role in peptidoglycan biosynthesis. FEMS Microbiol
Rev *32*, 234-258.

Sheng, Y., Fan, F., Jensen, O., Zhong, Z., Kan, B., Wang, H., and Zhu, J. (2015).
Dual Zinc transporter systems in Vibrio cholerae promote competitive advantages
over gut microbiome. Infect Immun *83*, 3902-3908.

Simpson, B.W., Nieckarz, M., Pinedo, V., McLean, A.B., Cava, F., and Trent, M.S.
(2021). Acinetobacter baumannii can survive with an outer membrane lacking
lipooligosaccharide due to structural support from elongasome peptidoglycan
synthesis. mBio *12*, e0309921.

Soneson, C., Love, M.I., and Robinson, M.D. (2015). Differential analyses for RNA-
seq: transcript-level estimates improve gene-level inferences. F1000Res *4*, 1521.

Srivastava, A., Malik, L., Sarkar, H., Zakeri, M., Almodaresi, F., Soneson, C., Love,
M.I., Kingsford, C., and Patro, R. (2020). Alignment and mapping methodology
influence transcript abundance estimation. Genome Biol *21*, 239.

Teoh, W.P., Matson, J.S., and DiRita, V.J. (2015). Regulated intramembrane
proteolysis of the virulence activator TcpP in Vibrio cholerae is initiated by the tail-
specific protease (Tsp). Molecular microbiology *97*, 822-831.

Vigouroux, A., Cordier, B., Aristov, A., Alvarez, L., Ozbaykal, G., Chaze, T.,
Oldewurtel, E.R., Matondo, M., Cava, F., Bikard, D.*, et al.* (2020). Class-A penicillin
binding proteins do not contribute to cell shape but repair cell-wall defects. Elife *9*.

Vollmer, W., Blanot, D., and de Pedro, M.A. (2008). Peptidoglycan structure and
architecture. FEMS Microbiol Rev *32*, 149-167.

Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M., and Barton, G.J. (2009).
Jalview Version 2--a multiple sequence alignment editor and analysis workbench.
Bioinformatics *25*, 1189-1191.

Weaver, A.I., Alvarez, L., Rosch, K.M., Ahmed, A., Wang, G.S., van Nieuwenhze,
M.S., Cava, F., and Dorr, T. (2022). Lytic transglycosylases mitigate periplasmic
crowding by degrading soluble cell wall turnover products. Elife *11*.

Weaver, A.I., Jimenez-Ruiz, V., Tallavajhala, S.R., Ransegnola, B.P., Wong, K.Q.,
and Dorr, T. (2019). Lytic transglycosylases RlpA and MltC assist in Vibrio cholerae
daughter cell separation. Molecular microbiology *112*, 1100-1115.

Zijderveld, C.A., Aarsman, M.E., den Blaauwen, T., and Nanninga, N. (1991).
Penicillin-binding protein 1B of Escherichia coli exists in dimeric forms. Journal of
bacteriology *173*, 5740-5746.

Fig. 1. Development of a high throughput method for peptidoglycan sample preparation and analysis. (a) Schematic of the comparison between the traditional and the novel high throughput (HT) method for PG isolation and examples of the potential types of analyses that can be carried out with large-scale muropeptide profile datasets. **(b)** Comparative pipeline for the HT and the traditional PG isolation method. **(c)** Comparative timescales for the isolation of PG from 96 samples following the traditional or the HT sample preparation method. **(d)** Representative PG profiles obtained for diverse bacteria using the HT PG isolation method.
See also Supplementary Fig. 1.

Fig. 2. Validation of the high throughput method for peptidoglycan sample preparation.
(a) Pairwise scatter plot comparing the relative abundance of every muropeptide between replicates of 900 samples. Mean Pearson correlation coefficient is shown. **(b-f)** Validation of the HT method for detection of minor changes in the PG structure by analysis of mutants known to have a phenotype in their muropeptide profile: Pbp1A (PG synthetase), LdtA (LD-transpeptidase) and DacA1 (DD-carboxypeptidase); **(b)** representative chromatogram of each strain; **(c)** PCA using the Log2FC of peak areas of all the muropeptides of each sample; **(d)** analysis of PG features; **(e)** heatmap presenting the relative peak area of each muropeptide numbered in panel **b**; **(f)** heatmap of the Log2FC values (relative to the wildtype) calculated for each peak numbered in **b**. In **e** and **f**, peaks expected to change in LdtA⁻ and DacA1⁻ mutants are highlighted in green and blue respectively. Error bars in **d** correspond to the standard deviation of triplicates.
See also Supplementary Fig. 2.

Fig. 3. Peptidoglycan profiling of the *V. cholerae* non-redundant mutant library.
(a) Total number of mutant strains available, processed and analyzed, and their classification into main COG groups.
(b) Muropeptide abundance and variation (SD) across the whole dataset. **(c)** Hierarchical clustering and heat-map representing the Log2FC calculated for different PG-features and muropeptides. **(d-f)** Analysis of the PG profile of genes belonging to the same transcriptional unit: PG features, heatmap and clustering of the PG profiles of indicated mutants and the expression data of each gene observed in 307 different RNAseq experiments are shown. **(g)** PCA obtained for the mutants of the library based on the Log2FC values observed for the 20 identified muropeptides. Outliers are colour-coded based on their PG-similarities clustering (see **h**). **(h)** Hierarchical clustering of the 50 outliers selected in **g**.
See also Supplementary Fig. 3 and Extended Data 1-3.

Fig. 4. Analysis of the main peptidoglycan features of the *V. cholerae* mutant library
(a) Violin plots showing the distribution of samples for the indicated PG-features in the 3030 analyzed mutants. Mutants expected to present a distinctive feature are labelled in each group. **(b)** Correlation scatterplot matrix for the different PG features in the *V. cholerae* mutant library. Histograms showing the distribution of samples for every PG feature are represented in the diagonal.

Fig. 5. *vc1321* encodes a new high molecular weight PBP in *V. cholerae*
(a) Over-represented biological processes associated to those mutants presenting low relative amount of PG (Log2FC <-0.5). The most representative biological processes with a fold enrichment higher than 2 are represented. Dot size indicates the number of loci/category and colour indicates the significance of the enrichment (-Log10(*P*-values)). **(b)** Plot showing the Log2FC of the PG amount for the complete dataset. Genes belonging to the "peptidoglycan metabolism" category in **a** are labelled. **(c)** Representative PG chromatograms of the wildtype, the *Δvc1321* mutant strain and the *Δvc1321* strain overexpressing *vc1321* from the arabinose-inducible promoter of the pBAD plasmid. **(d)** Quantification of the relative PG amount and DD-crosslink of the wildtype, *Δvc1321* mutant strain, and the *Δvc1321* carrying the pBAD empty plasmid as control or overexpressing *vc1321* from the arabinose-inducible promoter. **(e)** Bocillin gel of indicated strains showing the band corresponding to the product of *vc1321* (named PBP1 of *V. cholerae*, PBP1V) **(f)** Schematic representation of the transglycosylase (TG) and transpeptidase (TP) domains predicted in the sequence of VC1321. **(g-h)** Bocillin gels of *vc1321* point mutants in different putative catalytic residues **(h)** and in candidate cysteines involved in the dimerization of the protein. **(i)** Deletion of *pbp1V* does not result in growth or morphological defect. Scale bar = 2 μm. **(j)** PBP1V homologs across different species.
See also Supplementary Fig. 4-6 and Extended Data 4.