Syntaxin-6 delays prion protein fibril formation and prolongs presence of toxic

aggregation intermediates

Daljit Sangar, Elizabeth Hill, Kezia Jack, Mark Batchelor, Beenaben Mistry, Juan M. Ribes,

Graham S. Jackson, Simon Mead, Jan Bieschke*

MRC Prion Unit at UCL, Institute of Prion Diseases, Courtauld Building, London W1W 7FF,

United Kingdom

*to whom correspondence should be addressed: j.bieschke@ucl.ac.uk

Abstract

Prions replicate via the autocatalytic conversion of cellular prion protein (PrPC) into fibrillar

assemblies of misfolded PrP. While this process has been extensively studied in vivo and in

vitro, non-physiological reaction conditions of fibril formation in vitro have precluded the

identification and mechanistic analysis of cellular proteins, which may alter PrP self-

assembly and prion replication. Here, we have developed a fibril formation assay for

recombinant murine and human PrP (23-231) under near-native conditions (NAA) to study

the effect of cellular proteins, which may be risk factors or potential therapeutic targets in

prion disease. Genetic screening suggests that variants that increase syntaxin-6 expression

in the brain (gene: STX6) are risk factors for sporadic Creutzfeldt-Jakob disease (CJD).

Analysis of the protein in NAA revealed counterintuitively that syntaxin-6 is a potent inhibitor

of PrP fibril formation. It significantly delayed the lag phase of fibril formation at highly sub-

stoichiometric molar ratios. However, when assessing toxicity of different aggregation time

points to primary neurons, syntaxin-6 prolonged the presence of neurotoxic PrP species.

Electron microscopy and super-resolution fluorescence microscopy revealed that, instead of

highly ordered fibrils, in the presence of syntaxin-6 PrP formed less-ordered aggregates

containing syntaxin-6. These data strongly suggest that the protein can directly alter the

initial phase of PrP self-assembly and, uniquely, can act as an 'anti-chaperone', which

promotes toxic aggregation intermediates by inhibiting fibril formation.

Introduction

Prion replication and prion-like mechanisms are believed to drive dozens of human neurodegenerative and systemic diseases, as well as scrapie, bovine spongiform encephalopathy and chronic wasting disease in animals (1-3). All of these disorders are caused by protein misfolding, but the details of the self-replication mechanism, the cellular factors involved in prion replication and the molecular basis for neurotoxicity are far from clear despite decades of research (4). Prions are fibrillar assemblies of misfolded cellular prion protein (PrP^C), which form highly ordered parallel in register intermolecular β-sheet amyloid structures (prion rods, (5,6)). While purified prion rods are not directly neurotoxic, prion-associated toxicity can be blocked by anti-PrP antibodies (7,8) suggesting that non-fibrillar PrP assemblies, which form after prion titres plateau (9-11), could be the toxic agents, as is the case for other amyloid forming proteins (12,13).

Prions can be highly infectious. However, accidental exposure only accounts for a tiny percentage of CJD cases. Similarly, mutations in the *PRNP* gene, which codes for the prion protein PrP, cause only 10-15% of CJD cases (14). The majority of prion disease cases are sporadic in nature, where PrP is thought to spontaneously misfold. It is likely that hidden genetic risk factors modulate either the susceptibility to prion replication, prion-associated neurotoxicity, or the initial prion formation. A recent genome-wide association study to discover non-*PRNP* risk factors of sporadic CJD (sCJD) identified two highly significant risk loci including one in and near to *STX6* (15). Elevated *STX6* mRNA in the caudate and putamen nuclei, two regions implicated in CJD pathology (16), correlated with CJD risk.

STX6 encodes syntaxin-6, an eight exon, 255 amino-acid protein that predominantly localises to the trans-Golgi network (TGN), and is mainly involved in recycling of cargo between the TGN and early endosomes (17). Syntaxin-6 is thought to form part of the t-SNARE complex involved in the decision of a target membrane to accept the fusion of a vesicle (18). Misfolded prion protein in infected cells localizes at the plasma membrane, the likely site of prion conversion (19), as well as in perinuclear compartments. Early and late endosomes, the endocytic recycling compartment and multivesicular bodies have also been implicated as prion replication sites (20,21). These observations raise two alternative mechanistic hypotheses for the role of syntaxin-6 in prion disease: (1) A direct interaction with PrP in the course of misfolding and prion replication, or (2) an indirect effect through the cellular processing of PrP^C or prions.

STX6 is also a risk gene for progressive supranuclear palsy (22), where the protein interacts directly with tau through its transmembrane domain (23). STX6 has been identified in a proteome-wide association study (PWAS) in Alzheimer's disease (24). Downregulation of

syntaxin-6 expression in a murine neuroblastoma scrapie cell model (PK1) did not alter PrP^C

expression and had no obvious effect on PrP^C trafficking (15), suggesting that it may directly

alter prion replication by a yet undiscovered chaperone-like activity.

An array of chaperone proteins aid and tightly control protein folding and protein

homeostasis in the cell (25,26). Chaperones, such as the heat shock protein HSP70 inhibit

protein misfolding and can delay amyloid formation *in vitro* (27) and in model organisms (28).

It seems likely that more proteins with chaperone-like functions are yet to be discovered.

Assessing the role of proteins with chaperone-like activity in PrP misfolding, fibril formation

and toxicity requires assays under native or near-native conditions. The established fibril

formation assays for PrP *in vitro* require non-physiological conditions, such as high

temperatures (29), denaturants (30), detergents or low pH (31). These conditions facilitate

the unfolding and conversion of PrP^C, but preclude the analysis of other folded proteins in

the context of PrP fibril formation. Here, we have developed a fibril formation assay for

recombinant murine and human PrP^C (23-231) under near-native conditions to study the

effect of syntaxin-6 and other cellular factors, which may be risk factors or potential

therapeutic targets in prion disease.

Results

Full-length prion protein forms amyloid fibrils under near-native conditions

In order to study PrP fibril formation under near-native conditions *in vitro,* full-length murine PrP^C 23-231 (mPrP23) and human PrP^C 23-231 (hPrP23) were expressed recombinantly (Figure 1 – figure supplement 1A), purified via Ni-NTA chromatography and refolded into their native conformation (Figure 1 – figure supplement 1E, F). Proteins were incubated in a non-binding 96-well plate under agitation, and amyloid formation was monitored via Thioflavin T (ThT) fluorescence (Figure 1A, Figure 1 – figure supplement 2). We tested different temperatures and pH values to assess whether PrP could form amyloid fibrils *in vitro* under near-native conditions (Figure 1 – figure supplement 2A, B). While PrP was slow to form aggregates *de novo* without the addition of seeds, elevated temperatures accelerated fibril formation (Figure 1 – figure supplement 2A) as had been previously observed (32). Aggregation at 42°C, pH 6.8 yielded highly reproducible conditions for the formation of fibrils (Figures 1A-C and Figure 1 – figure supplement 2B, C) when agitated by glass or silanized Zirconia beads. Fibrils had diameters of 8-12 nm and morphologies typical of amyloid fibrils (Figures 1C, Figure 1 – figure supplement 2G). The addition of preformed fibrillar aggregates as seeds accelerated fibril formation in a concentration-dependent manner by decreasing lag-times (t_{50}, Figures 1A, B and Figure 1 – figure supplements 2D and 3) but had no significant influence on rates of fibril elongation (Figure 1 – figure supplement 2E). Lag phases and elongation rates of mPrP23 *in vitro* were indistinguishable when using synthetic seed formed from recombinant PrP^C or using authentic prion rods, isolated from mice infected with scrapie strains RML or ME7 in (Figure 1 – figure supplements 2D, E, and 3). Interestingly, fibrils formed from synthetic seeds displayed increased ThT fluorescence when compared to RML and ME7 seeded fibrils, hinting at a difference in fibril structure imprinted by the different seed templates (Figures 1G, Figure 1 – figure supplement 2F). For safety reasons, seeding of hPrP23 was restricted to synthetic PrP fibrils.

The lag phase of hPrP23 aggregation depended only weakly on PrP monomer concentration (Figure 1D, E), while elongation rates even showed a slight inverse reaction to monomer concentration (Figure 1F). Therefore, PrP refolding rather than elongation may be rate-limiting under the assay conditions. ThT amplitude at the end of growth was proportional to monomer concentration (Figure 1G). However, the lag phase increased and ThT amplitude dropped at monomer concentrations above 16 µM, suggesting formation of off-pathway PrP aggregates at high monomer concentrations, which do not contribute to fibril formation (33).

Syntaxin-6 inhibits PrP fibril formation

The *STX6* gene is a proposed risk factor for sCJD, suggesting it may alter PrP fibril

formation. We therefore tested what effect the syntaxin-6 protein had on PrP fibril formation

under native conditions and compared it to two reference proteins: a well characterized

amyloid inhibitor, the heat shock protein HSPA1A, which is the human inducible form of

HSP70, and the α-helical microtubule-associated protein stathmin 1 (STMN1) as positive

and negative controls, respectively (Figure 1 – figure supplement 1A). Circular dichroism

confirmed that all proteins retained their native, mostly α-helical folds under assay conditions

(Figure 1 – figure supplement 1B-F).

To our surprise, syntaxin-6 delayed the lag phase of hPrP23 fibril formation even at highly

sub-stoichiometric molar ratios of 1:100, and prevented fibril formation entirely at equimolar

ratio (STX6, Figure 2A, B). At the same time, it only slightly lowered fibril elongation rates

(Figure 2C). Syntaxin-6 promoted the formation of aggregate clusters (Figure 2D, E) rather

than distinct amyloid fibrils (Figure 1C). Immuno-gold staining revealed that syntaxin-6 was

incorporated into these aggregates (Figure 2E). Similarly, HSPA1A prevented fibril formation

in favour of amorphous PrP aggregates (Figure 2 – figure supplement 1A), while hPrP23

formed fibrillar aggregates in the presence of control protein STMN1 (Figure 2 – figure

supplement 1B). However, sedimentation after 0-90 h incubation revealed, that syntaxin-6

delayed but did not prevent the formation of hPrP23 aggregates. In contrast, a substantial

fraction of hPrP23 remained soluble when incubated with HSPA1A at 1:10 molar ratio

(Figure 2 – figure supplement 2).

To further probe the mechanism of syntaxin-6, we tested the influence of NaCl concentration

on fibril formation in the presence and absence of syntaxin-6 (1 in 10 molar ratio) in the NAA

(Figure 2 – figure supplement 3, Figure 3 – figure supplements 2 and 3). We compared

kinetics in the presence of syntaxin-6 to those in the presence of HSPA1A and STMN1,

respectively, at the same molar ratio (Figure 2 – figure supplement 3A). HSPA1A delayed

fibril formation to a similar degree as syntaxin-6, whilst, as expected, STMN1 had no

significant effect on fibril formation kinetics. In the absence of syntaxin-6, the lag phase of

PrP fibril formation decreased only weakly with NaCl concentration while the elongation rate

increased (Figure 2 – figure supplement 3B, C). Syntaxin-6 prolonged the lag phase at low

and physiological salt concentrations, but lag phases decreased with increasing NaCl

concentration, suggesting that ionic interactions may mediate the binding of syntaxin-6 to

PrP. A similar salt dependence was observed for HSPA1A (Figure 2 – figure supplement

3D), while STMN1 did not inhibit fibril formation at any salt concentration (Figure 2 – figure

supplement 3E). Syntaxin-6 did not significantly reduce fibril elongation when compared

with untreated PrP independently of salt concentration (Figure 2 – figure supplement 3C).

To visualize the interaction of syntaxin-6 with PrP aggregates and fibrils we imaged NAA

endpoints by dSTORM super-resolution microscopy (Figure 3 – figure supplement 1)

revealing that hPrP23 and syntaxin-6 co-aggregated into large aggregate clusters. Super-

resolution microscopy indicated a fibrillar substructure of these aggregates (Figure 3 – figure

supplement 1A, B) similar to that observed in TEM (Figure 2D, E). When, conversely,

syntaxin-6 labelled with AlexaFluor647 was added to pre-formed fibrillar hPrP23 formed

under NAA conditions, it preferentially bound to specific 'hot-spots' on the PrP fibril (Figure

3A, B, indicated by arrows). In this experiment, hPrP23 fibrils were unlabelled and were

visualized by transient amyloid binding (TAB) imaging using the amyloidophilic dye Nile red

(34). We repeated the experiment with syntaxin-6-AlexaFluor488 (Figure 3E) and with

unlabelled proteins using immuno-TEM (Figure 3F) to exclude that the interaction was

influenced by the choice of fluorescent dye. Both assays confirmed the presence of

interaction hot-spots. Interestingly, interaction sites were often located at fibril ends or kinks

in the fibril. Since syntaxin-6 only weakly affected apparent fibril elongation rates in bulk

assays, this may suggest a role of syntaxin-6 in fibril breakage or secondary nucleation.

PrP formed longer, unbranched fibrils with increasing salt concentration. As had been seen

in TEM, syntaxin-6 induced the formation of aggregate clusters when visualized by TAB

imaging (Figure 3 – figure supplements 2 and 3) when compared to untreated fibrils,

supporting the hypothesis that the protein may alter secondary nucleation / branching of

fibrils.

Syntaxin-6 interacts with PrP in cell models of prion disease

Syntaxin-6 and PrP need to interact in order to directly affect prion replication and / or prion-

associated toxicity *in vivo*. We therefore probed the interaction of syntaxin-6 with PrP via

Förster resonance energy transfer (FRET) imaging in the PK1 cell model (35), which can be

persistently infected with RML prions. Proteins were stained with anti-syntaxin-6 antibody

(C34B2) / anti-rabbit-Alx647 and three monoclonal anti-PrP antibodies (5B2, 6D11, 8H4) /

anti-mouse-RhX, which bound different regions of PrP (Figure 4A, and Figure 4 – figure

supplements 1 and 2). PixFRET analysis indicated a perinuclear interaction between both

proteins, both in non-infected (PK1) and infected (iS7) cells (Figure 4A, wide arrows). An

additional FRET signal on the plasma membrane of infected cells suggests that syntaxin-6

may be recruited into misfolded PrP assemblies (Figure 4A, narrow arrows). This interaction

resulted in FRET signals with all three anti-PrP antibodies, whereas only the two antibodies

directed to the unfolded N-terminal domain of PrP (5B2 and 6D11) displayed FRET in

membrane-associated compartments (Figure 4 – figure supplement 1). This may suggest a

different binding mode of syntaxin-6 to PrP species in both compartments.

Syntaxin-6 knockout does not affect the replication of prions in vitro

Infectious prions can be replicated in cell-free conditions using protein misfolding cyclic

amplification (PMCA), whereby the conversion of PrP^C into PK-resistant PrP^{Sc} is enhanced

by cyclic bursts of sonication in a brain homogenate substrate (36). Here, we employed

PMCA to explore whether the presence or absence of syntaxin-6 in the substrate affected

prion conversion. To this end, we seeded PMCA substrates derived from $Stx6^{+/+}$ and $Stx6^{-/-}$

mice, respectively, (37) with RML prions. Both substrates generated comparable amounts of

PK-resistant PrP (Figure 4B), indicating that syntaxin-6 does not directly alter prion

replication *in vitro* under PMCA conditions, which are dominated by fibril fragmentation /

elongation.

Syntaxin-6 prolongs the presence of toxic PrP aggregation intermediates

The previous data would suggest a protective rather than deleterious effect of syntaxin-6 in

prion disease. However, the oligomer toxicity hypothesis posits that mature fibrils may not be

the toxic species in prion nor in other amyloid diseases (11,12). Correspondingly, purified,

highly infectious prion rods were not toxic to primary neurons (7). To test whether the same

logic applied to the effect of syntaxin-6 on PrP aggregation, we assessed the toxicity of PrP

to mouse primary neurons at different stages of the aggregation kinetics by neurite length

(Figure 5) and by counting the number of viable neurons (Figure 5 – figure supplement 1).

PrP aggregates were highly toxic to primary neurons during lag and early growth phases (20

h and 40 h incubation), but neurotoxicity was diminished at the plateau phase of fibril

formation, so that end-point aggregates (90 h) were no more toxic than PrP monomers or

buffer controls (Figure 5A, C and Figure 5 – figure supplement 1). Notably, PrP toxicity

preceded the formation of seeding-competent assemblies (Figure 5 – figure supplement 2),

supporting the hypothesis that the toxic PrP species is a pre-fibrillar assembly.

In contrast, PrP aggregated in the presence of syntaxin-6 (1:0.1 molar ratio) retained its

neurotoxicity significantly longer than PrP incubated in the absence of syntaxin-6 (Figure 5B,

C and Figure 5 – figure supplement 1). While PrP species formed early in the lag phase of

aggregation at 20 h were neurotoxic both in the presence and absence of syntaxin-6 when

compared to t = 0 h, PrP was significantly more toxic in the presence of syntaxin-6 at later

time-points when PrP would form fibrils in the absence of syntaxin-6 (Figures 5C and Figure

5 – figure supplement 1).   Thus, in delaying or preventing PrP fibril formation, syntaxin-6

prolonged the presence of toxic aggregation intermediates and exacerbated neurotoxicity.

Discussion

Development of a novel native aggregation assay

We developed a fibril formation assay for full length mouse and human PrP[C] to be able to

study the mechanistic effect of cellular proteins such as the proposed sCJD risk factor

syntaxin-6 on PrP misfolding and self-assembly under near-native conditions. A multitude of

in vitro fibril formation assays have been developed previously (38). While the generation of

authentic prion infectivity was reported for several assays (30,38), fibrils generally form

under highly non-physiological conditions in the presence of high concentrations of

Guanidine or Urea (30), at low pH (39), in the presence of SDS (40), or at very high

temperatures (41). None of these condition provide a native environment for protein-protein

interaction, and many cellular proteins will be unfolded under these assay conditions.

Full length PrP formed fibrils with morphology and β-sheet secondary structure characteristic

of amyloid under conditions of the native aggregation assay (NAA) (42). Fibril formation

could be tracked by binding of the amyloidophilic dye Thioflavin T and displayed sigmoid

aggregation kinetics typical of a nucleated polymerization mechanism (43). As had been

observed for other amyloidogenic proteins, solution conditions such as pH and temperature

affected aggregation kinetics (44,45). Aggregation proceeded faster at neutral pH, while

monomeric protein was stable at pH of 6.5 and below.

Seeding by preformed fibrils reduced the lag phase of aggregation (46,47). The limiting

dilution for seeding was ~ 10^{-10} (w/w), which corresponds to approximately 2000 PrP

molecules per well or 5 prion rods of a typical length of 200 nm (48), similar to other seeding

assays (40,41,49). In NAA, both lag phase t_{50} and elongation rate constant k displayed little

dependence on monomer concentration, which suggests that monomer refolding was rate

limiting for fibril elongation. While in general aggregation kinetics scale with monomer

concentration, elongation saturates at high monomer concentration as conformational

change of PrP becomes rate-limiting (50,51). ThT fluorescence amplitudes scaled with PrP

concentration between 2 – 16 μM, it then dropped and lag phase increased. A competing

non-fibrillar aggregation pathway at high protein concentration could account for this

observation (33,52). High salt concentration shields charges, which promotes hydrophobic

interactions, lowers the critical concentration of nucleated polymerization often leading to

formation of amyloid fibril clusters (53). Notably, while PrP aggregated faster in the NAA at

high salt concentration, it tended to form longer, more isolated fibrils.

Inhibition of fibril formation by syntaxin-6

Syntaxin-6 delayed the formation of PrP fibrils at highly sub-stoichiometric ratios and co-

aggregated with PrP into aggregate clusters. On the face, its effect was similar to that of the

heat shock protein family A (HSP70) member 1A (Figure 2 and Figure 2 – figure supplement

3), which would suggest a role of syntaxin-6 as a potent aggregation inhibitor with

chaperone-like activity. Chaperones assist folding of newly synthesised proteins, sequester

soluble misfolded polypeptides and target them for degradation, which makes them

promising therapeutic targets in protein misfolding diseases (26,54,55).

HSP70 has been described as a 'holdase', which inhibits amyloid formation of various

proteins at substoichiometric concentrations even in the absence of ATP by binding to early

stage aggregation intermediates (56,57). Overexpression or exogeneous addition of HSP70

rescued cytotoxicity in cell cultures and model organisms (58-60). Sequestering aggregation

intermediates into inert off-pathway aggregates can be a powerful strategy against amyloid

toxicity (61,62). However, our data indicate that, far from detoxifying PrP aggregation

intermediates, the delay of PrP fibril formation by syntaxin-6 exacerbated PrP toxicity by

allowing toxic aggregation intermediates to persist for longer (Figure 5).

Conversely, natural and chemical chaperones can detoxify aggregation intermediates by

promoting fibril formation and by stabilizing mature amyloid fibrils (47,63,64). Syntaxin-6

seems to present the flip side of this benign activity in delaying PrP fibril formation and thus

prolonging the presence of toxic aggregation intermediates.

The nature of the toxic species in prion disease is under active investigation. Brain

homogenates from scrapie-infected mice have a specific toxic component, which can be

blocked by the addition of anti-PrP antibody (7). However, highly purified infectious prion

rods were shown not to be directly neurotoxic (7), suggesting that toxicity may be caused by

non-prion assemblies, which accumulate after the infectious titer has peaked and which,

unlike prions, are sensitive to sarkosyl (8-10) or by transient aggregation intermediates,

whose formation may be catalysed by prions or by other fibrillar PrP assemblies (4).

The chaperone Brichos breaks generation of toxic Aβ oligomers, which are formed through

surface catalysed secondary nucleation (65). Conceivably, syntaxin-6 could have the

opposite effect on PrP fibril formation, in which toxic oligomer formation is promoted, by

delaying competing fibril formation pathways. PrP itself binds to and inhibits the elongation

of Aβ amyloid fibrils (66,67) oligomers and nanotubes. The presence of PrP during

aggregation led to the formation of larger numbers of shorter fibrils and increased Aβ

neurotoxicity in a dose-dependent manner (66). This suggests that PrP not only acts as a

receptor for toxic amyloid oligomers (68,69), but may exacerbate neurotoxicity in a

mechanism similar to syntaxin-6. Notably, syntaxin-6, when added to preformed fibrils,

preferentially bound to fibril ends and other hotspots (Figure 3). Previous studies found only

weak binding of syntaxin-6 to monomeric PrP^C (15), which together with the inhibition at sub-

stoichiometric concentrations strongly suggests an interaction with early aggregation

intermediates and with fibrillary assemblies, possibly promoting secondary nucleation.

While variants at the *STX6* locus are known risk factors for sCJD (15), its deletion only

modestly delayed the incubation period in RML prion infected mice, an observation that is

open to different interpretations (37). Rather than directly altering prion replication kinetics,

the protein may confer risk of disease by either facilitating the initial generation of prions in

sporadic disease, or by exacerbating prion-associated toxicity. Our results lend support to

the second hypothesis, as syntaxin-6 interacted with early aggregation intermediates of PrP

in vitro and exacerbated their toxicity (Figure 5) but did not accelerate the formation of

seeding-competent PrP species (Figure 5 – figure supplement 2) nor directly affected the

replication of pre-existing prion seeds in PMCA reactions (Figure 4B). This finding argues

against the hypothesis that syntaxin-6 binding to mature PrP fibrils induces fibril breakage.

Correspondingly, an increase in fibrils branching / secondary nucleation as observed in the

presence of syntaxin-6 (Figure 3 – figure supplements 1 and 3) would have little effect on

replication kinetics dominated by fragmentation / elongation.

Localization of syntaxin-6 and PrP

A direct effect of syntaxin-6 on PrP assembly and / or prion-associated toxicity *in vivo*

obviously requires the two proteins to be present in the same sub-cellular compartment.

PrP^C is present mainly in the TGN and on the outer plasma membrane, while mature PrP^C is

tethered to the membrane by a C-terminal glycophosphadityl-inositol (GPI) anchor. At the

plasma membrane, PrP^C is incorporated into lipid rafts and caveolae, which are membrane

microdomains enriched in cholesterol and sphingolipids. However, it has also been detected

in multi-vesicular bodies and, to a small degree, in the cytosol (70). Both lipid rafts and the

endocytic pathway are implicated as sites of prion replication (70).

Interestingly, syntaxin-6 has role in caveolin membrane recycling (71). As a SNARE protein,
syntaxin-6 sits on the outside of vesicles / intracellular compartments and on the inner
plasma membrane, so that a direct interaction with PrP would require mislocation of either
protein. Both, PrP and syntaxin-6 contain a trans-membrane domain and PrP can populate
transmembrane forms in neurons (72,73). It is therefore plausible that, similar to the tau
protein, PrP and prions bound to the membrane bilayer could interact with the
transmembrane domain of syntaxin-6. Alternatively, mislocation of misfolded PrP assemblies
could facilitate interaction with syntaxin-6. Amyloid assemblies can disrupt membrane
integrity through mechanical rupture, pore formation or altering membrane curvature (74,75).
Disruption of membranes on the cell surface and in vesicles of the endocytic pathway, is a
central step in the internalization of other misfolded polypeptides, such as Aβ (76,77).
Similarly, it has been hypothesized to be necessary for cellular prion propagation (21).
These processes could bring misfolded PrP into direct contact with syntaxin-6. FRET
imaging suggests a direct contact between PrP and syntaxin-6 in a membrane-associated
compartment in prion infected cells (Figure 4A and Figure 4 – figure supplement 1B),
supporting this hypothesis.

Conclusions

We developed an aggregation assay of murine and human prion protein under near-native
conditions. PrP forms amyloid fibrils under the assay conditions, which bind ThT and which
can seed further fibril formation. Unlike natively unfolded proteins, however, PrP fibril
formation does not scale strongly with monomer and seed concentrations, suggesting that
structural conversion of native PrP into the amyloid fold is rate limiting. This is consistent
with the observation that the time constant of exponential prion replication *in vivo* is
independent of PrP expression (9).

In the NAA both the PrP and potential protein modulators of aggregation are in their native
conformation, which means that mechanisms by which protein and small modulators alter
amyloid formation can be assessed. We analysed the effect of syntaxin-6, a recently
proposed risk factor for sCJD, on PrP self-assembly. To our surprise, we discovered that the
protein acts as an 'anti-chaperone', which, by delaying PrP fibril formation, prolonged the
persistence of toxic aggregation intermediates *in vitro*. Genetic variants in or near to *STX6*
that enhance brain expression of the protein might therefore modify risk of CJD by direct
interaction with PrP and changing aggregation pathways, including the possibility of
favouring more toxic aggregation intermediates.

While at this point we cannot assess whether the toxicity of PrP aggregation intermediates
formed *in vitro* recapitulates the authentic PrP species responsible for neurotoxicity in prion
disease *in vivo*, our data highlight a new mechanism by which protein modulators of amyloid
formation can have counter-intuitive deleterious effects.

Acknowledgements

Research was supported by the National Institute of Neurological Disorders and Stroke of
the National Institutes of Health grant number 1R21NS101588-01A1 and by MRC grant
MC_UU_00024/6 to JB.

We thank Adam Wenborn and Dr. Jonathan Wadsworth, MRC Prion Unit, for providing
mouse scrapie material and mass spectrometry, Dr. Peter Klöhn for providing PK1 S7 and
iS7 cell lines, Prof. Erich Wanker, MDC-Berlin, Germany for providing *E. coli* clones for
protein expression and the staff of the MRC Prion Unit at UCL Biological Services Facility for
animal husbandry and care. We thank Dr. Georg Meisl, University of Cambridge, for helpful
discussions.

The *Stx6*[-/-] mice were obtained from the MRC Harwell Institute, which distributes these mice
on behalf of the European Mouse Mutant Archive (https://www.infrafrontier.eu/emma/). The
MRC Harwell Institute is also a member of the International Mouse Phenotyping Consortium
(IMPC) and has received funding from the Medical Research Council for generating and/or
phenotyping the *Stx6*[-/-] mice. The research reported in this publication is solely the
responsibility of the authors and does not necessarily represent the official views of the
Medical Research Council. Funding and associated primary phenotypic information may be
found at www.mousephenotype.org.

Methods and Materials

Protein Expression and Purification:

Human PrP (23-231), Mouse PrP (23-231)

The open reading frame of the human PrP gene (*PRNP*) (residues 23-231), containing methionine at residue 129 and the mouse PrP gene (*Prnp*) (residues 23-231, including S231), was synthesised de novo by Eurofins MWG Operon, with a thrombin-cleavable His-Tag added to the PrP N-terminus. The ligated pTrcHisB/PRNP and pTrcHisB/Prnp constructs were used to transform the Escherichia coli host strain BL21(DE3) (Novagen), genotype F′ ompT hsdSB (rB- mB-) gal dcm (DE3), which was then plated onto Luria-Bertoni (LB) agar plates containing 100 μg / mL carbenicillin. Cultures were grown for purification using a modification of protocols previously described (78). Briefly, following harvesting, cells were sonicated and their inclusion bodies containing PrP resolubilised in 6 M Guanidine Hydrochloride (GuHCl), 50 mM Tris-HCl, 0.8% β-mercaptoethanol, pH 8.0. These were loaded onto a Ni-NTA column equilibrated in 6 M GuHCl, 10 mM Tris-HCl, 100 mM Na_2PO_4, 10 mM glutathione pH 8.0, and eluted from the column using 10 mM Tris-HCl, 100 mM Na_2PO_4, 2 M Imidazole pH 5.8. Residual GuHCl was removed through dialysis against 20 mM Bis Tris.HCl pH 6.5, $CaCl_2$ added to a final concentration of 2.5 mM, and the N-terminal His-tag cleaved by thrombin for 16 h at room temperature (0.1 U thrombin (*Novagene*) / 1 mg of PrP added). The cleaved protein was loaded onto a second Ni NTA column equilibrated with 20 mM Bis-Tris pH 6.5, 25 mM imidazole pH 6.5 and the eluted PrP peak was collected and dialysed against 10 mM Bis Tris pH 6.5, and aliquoted and stored at -80°C. Protein concentrations were determined by UV absorption at 280 nm using a calculated molar extinction of 56667 M^{-1} cm^{-1} and 62268 M^{-1} cm^{-1} for human and mouse PrP, respectively (*https://web.expasy.org/protparam*).

SEC of Syntaxin-6 / hPrP23 / mPrP23

2 mL Pre-concentrated, pre-cleared protein was loaded via injection valve onto a Sephacryl S100HR column (GE Healthcare) (26/60: 320 mL bed volume) that had been pre-equilibrated with 20 mM Tris-HCl, 0.2 M NaCl, pH 8. A flow-rate of 2 mL / min was used; all eluted peaks were checked on a silver stained NuPage Bis Tris 12% gel, using the protocol according to the silver staining kit (SilverQuest Silver Staining Kit-1, LC6070, Life Technologies Ltd) and the purified protein was aliquoted and stored at –80°C until use.

Stathmin 1 (STMN1) / Heat-Shock Protein 70 (HSPA1A)

Glycerol stocks of E. Coli BL21 strains expressing His-tagged STMN1 and HSPA1A were a gift by E. Wanker, MDC-Berlin, Germany. Small scale cultures of STMN1 and HSPA1A were cultured in LB/Amp (100 µg / mL) by inoculating with stabs from glycerol stocks. Cells were pelleted and then inclusion bodies lysed in 1 mL lysis buffer (containing 50 mM Tris, 0.2 M NaCl, pH8 containing 0.1% Tween20, 0.5% NP40, 50 U / mL Benzonase, 10 µg / mL Lysozyme and 1 mM PMSF) placed on ice for 1 h with gentle vortexing every 15 min. (All centrifugations were done at 4°C to further prevent non-specific cleavage by proteases and samples were kept on ice.) The lysate was cleared by centrifugation (1 h, 16100 rpm) and then loaded onto 2 mL NiNTA resin bed packed in mini spin columns, pre-equilibrated with buffer containing 50 mM NaH_2PO_4, 0.3 M NaCl, 20 mM Imidazole at pH 8. The cleared lysate was loaded onto the column, unbound was eluted off the column first with gentle centrifugation (2 min, 2000 rpm); the resin was washed to thoroughly remove unbound protein by twice loading 700 µl pre-equilibration buffer (see above) and centrifugation for 30 s, 2000 rpm and then 2 min, 2000 rpm for the second wash. Finally, the pure protein was eluted in elution buffer (containing 50 mM NaH_2PO_4, 0.5 M NaCl, 0.5 M imidazole, pH 8). This was done by loading 100 µl elution buffer with a 30 s centrifugation at 2000 rpm followed by a second elution with 150 µl elution buffer and a 2 min centrifugation at 2000 rpm. Both fractions were pooled and checked for purity as below.

The purified STMN1 and HSP70 recombinant proteins were checked for purity by coomassie staining after PAGE on NuPage BisTris 12% gels. Protein identity was confirmed by mass spectrometry following a standard trypsinization protocol on a Waters Xevo-XS spectrometer as described in (5).

Syntaxin-6 (STX6)

Syntaxin-6 (residues 38-318; Accession number KU144824) was prepared according to (79) with modifications. Briefly, the DNA sequences encoding the syntaxin-6 protein in pQTEV were transformed into BL21 (DE3). BL21 cultures were grown in LB medium in the presence of 100 µg / mL Ampicillin. Expression of the protein was induced using 1 mM IPTG and was purified from inclusion bodies under denaturing conditions using Nickel superflow resin with an AKTA Pure (GE health care Life Sciences). The protein was refolded on NiNTA

resin and eluted from the column using an imidazole gradient. The eluted material was
extensively dialysed against 20 mM Tris, 2 mM EDTA, 10 mM DTT, 200 mM NaCl pH 8.0.
DTT concentration was reduced to 2 mM in the final storage buffer. Syntaxin-6 was further
purified by size exclusion chromatography as detailed above resulting in a protein band with
an apparent molecular weight of ~28 kD (Figure 1 – figure supplement 1A).   The final
concentration of the syntaxin-6 protein was determined by absorption measurement at 280
nm, ε = 31970/M/cm. Aliquots were stored at –80°C until use.
463
464

Circular Dichroism (CD) spectroscopy

466 CD data were recorded on a Jasco J-715 spectrophotometer equipped with a thermoelectric
467 temperature controller. A 1 mm path length cuvette was used for all CD spectroscopy
468 measurements. Wavelength measurements were recorded from 195 nm to 260 nm at 20□C,
469 using 0.36 mg / mL syntaxin-6 in 10 mM Na-phosphate, pH 8.
470

Native Aggregation Assay (NAA)

PrP[C] was filtered through a 100 kD membrane filter in 10 mM Bis Tris pH 6.5 (Amicon
ULTRA 0.5 mL 100K 96PK, UFC510096, Merck Life Science UK Ltd) to remove aggregates
and then diluted into reaction buffer (50 mM Na-phosphate, pH 6.8, 150 mM NaCl, 0.01%
Na-Azide, 20 µM ThT and 5 mM Bis-Tris pH 6.5) with the addition of 0.1% seed (w/w),
unless indicated otherwise. The standard final concentrations of PrP and NaCl were 2.5 µM
and 150 mM, pH 6.8 unless indicated otherwise. The seed consisted of aggregated PrP
material from an unseeded reaction under the same conditions as above. It was diluted to
10% (w/w) in 5 mM Bis-Tris, pH 6.5 and sonicated in a water bath sonicator (GRANT,
XUBA1) for 15 min prior to addition to the reaction mix. Aggregation assays for neuronal
toxicity were set up in parallel to the standard assays for kinetic analysis, but here the ThT
and Na-Azide was omitted from the reaction mixes.

The seed and post aggregation samples were handled with pre-silanised tips, in pre-
silanised non-binding tubes (Repel-silane ES, Sigma, 17-1332-01). The reaction mix (94 µl
per well) was dispensed into low binding 96 well COSTAR (#3651) plates and three silanised
zirconium beads (0.5 mm diameter, Figures 1-4; 1 mm diameter, Figure 5) were added to
each well. A single glass bead (2 mm) was added instead of the Zr beads where indicated.
ThT kinetics were recorded on a BMG ClarioStar plate reader at 42°C with a shaking speed

of 700 rpm, set to shake for 100 s with an incubation time of 20 s between agitation. Focal

height was set to 20 mm, top read, with excitation at 440 nm and emission set to 485 nm.

Aggregation kinetics were fitted in Matlab E2021b using the following equation:

$$F = f_0 + \frac{A}{1 + e^{-k(t - t_{50})}}$$

$$t_{lag} = t_{50} - \frac{2}{k} \qquad\qquad \text{Equation 1}$$

EM grid preparation

End-point samples (5-6 µL) from native aggregation assays sonicated in a water bath for

10-15 s   were loaded onto carbon-coated 300 mesh copper grids (Electron microscopy

Sciences) that had been glow discharged for 40 s using an PELCO easiGLOW™ glow

discharge unit (Ted Pella Inc., USA). Samples were left to bind for 30 min, blotted dry,

washed in water (1 x 50 µL), blotted, and then stained with 10 µL Nano-W (methylamine

tungstate) stain (Nanoprobes) for 1 min followed by 30 s (with blotting in between stain

times). Images were acquired on a Talos electron microscope (FEI, Eindhoven, NL now

ThermoFisher).

Immunogold labelling of PrP fibrils with NAPTA precipitation

40 µL of fibrils (completion at 115 h) were dispensed into an RNase free pre-silanised tube

with silanised tips and centrifuged for 1.5 h (4°C, 16100 rpm) after which the pellet was

resuspended in 500 µL TBS containing 0.1% (w/v) sarkosyl with sonication for 30 s in a

water bath sonicator (GRANT, XUBA1), and incubated with syntaxin-6 antibody (C34B2)

Rabbit mAb 2869 (Cell Signaling Technology Europe B.V, 2869S; 1:100) for 16 h at 25 °C

with gentle agitation.

The following day, the fibrils were precipitated with sodium phospho-tungstate (NaPTA) by

addition of 40.5 µL of 4% (w/v) NaPTA (prepared in H_2O; pH 7.4) and centrifuged at top

speed for 30 min to recover a pellet which was resuspended in 10 µL total volume of 1:20

(v/v) Goat anti-Rabbit IgG conjugated to colloidal gold (10 nm) (Insight Biotechnology Ltd,

GA1014) in TBS containing 5% (v/v) glycerol and incubated at 25 °C for 3 h with gentle

agitation. The sample was pulse centrifuged for 5 s and 5 µL labelled sample was loaded

onto a glow-discharged carbon coated grids and stained with Nano-W as previously

described without the 10-15 s pre-sonication step.

SR sample preparation

8-well chamber slides (IBL Baustoff, 2 20.140.082, C8-1.5-H-N, 8 well chambered cover

glass with #1.5 high performance cover glass, 0.170 +/- 0.005 mm) were cleaned by soaking

them overnight in a 2% solution of Hellmanex II detergent diluted in ultrapure water and then

washed thoroughly in ultrapure water before rinsing in 100% methanol and then a final

thorough rinse in ultrapure water and then allowing the slides to dry.

Once dry, the slides were glow discharged for 40 s using a PELCO easiGLOW™ glow

discharge unit (Ted Pella Inc., USA). 10 µL sample was added to the centre of a well and

allowed to incubate for 1 h for the fibrils to adhere to the glass. The well was then washed

5X with 500 µL of HPLC purified water before 200 µL buffer (which consists of   10nM Nile

Red (Sigma-Aldrich, 72485-1G in PBS or GLOX buffer). The wells were then sealed either

with Nescofilm or TWINSIL before imaging on a custom built TAB/dSTROM super-resolution

microscope (80).

Enzymatic oxygen scavenger (GLOX, glucose oxidase with catalase) buffer consisted of two

solutions. Solution A: Tris (50 mM, pH 8.3), NaCl (10 mM), glucose (10% w/v), and β-

mercaptoethylamine (Sigma-Aldrich, 30070, 10 mM). Solution B: glucose oxidase (Sigma-

Aldrich, G2133, 8 mg), and catalase (Sigma-Aldrich, C100, 38 µL, 21 mg / mL) in PBS (160

µL). Solutions A and B were mixed at the ratio of 99:1 (v/v) immediately before use.

Labelling recombinant protein with AlexaFluor dyes

AlexaFluor Dyes (AlexaFluor ^{TM} 488 NHS Ester, A20000, AlexaFluor ^{TM} 647 NHS Ester,

A20006, AlexaFluor™ 488 C5 Maleimide, A10254, thermofisher.com) were diluted in DMSO

to a stock concentration of 10 mg / mL. The dyes were mixed with the recombinant protein

constructs at a molar ratio of 2:1 dye:protein construct. The mixture of dye and protein

constructs were protected from light by covering the vessel with foil and left to mix on a

rotator overnight at 4°C. The following day, the unbound dye was removed by dialysis in 2.5

L for 2 x 1 h, with 8 kD dialysis membrane (SpectraPor 7 8000 Dalton MWCO; 11425919,

Fisher Scientific UK). For syntaxin-6 the buffer used for dialysis was 10 mM Na-

phosphate,150 mM NaCl, 2 mM EDTA, 2 mM DTT, pH 8; for PrP, the buffer used for dialysis

was 10 mM Bis-Tris, pH 6.5. Once dialysis was complete, the labelled protein was recovered

from the dialysis membrane and checked on a 4-12% Bis-Tris gel. To check labelling

efficiency, an absorption spectrum measurement (200-700 nm) was taken and the final

concentrations of dye and protein were calculated according to the manufacturer protocol.

The labelled proteins were then aliquoted and stored at -80°C.

Primary Neuronal Culture and Toxicity Analysis

Primary neuronal cultures were derived from brains of unmodified inbred FVB/N mice.

Hippocampi of male and female E17 mouse brains from a single litter (7-9 embryos) were

dissected in HBSS (ThermoFisher Scientific) supplemented with 1% L-glutamine, 1%

HEPES and 1% pen-strep. Cells were dissociated using 0.25% trypsin+0.04% benzonase,

triturated mechanically and counted using a Neubauer haemocytometer. Cells were plated in

DMEM supplemented with 10% horse serum (#26050-88, Invitrogen) at 10K / well to the

inner 60 wells of poly-L-lysine coated 96 well plates (Greiner, 655936). At 1 h post-plating,

DMEM medium was aspirated and exchanged for Neuralbasal medium (21103049,

ThermoFisher Scientific) supplemented with 0.25% Glutamax (35050061, ThermoFisher

Scientific) 2% Gibco B27 supplement (17504044, ThermoFisher Scientific) and incubated at

37°C (20% O2, 5% CO2). FVB/N neurons were maintained in culture for 11-12 days prior to

a 96 h treatment. Images of live cells were taken on an IncuCyte S3 reader (Sartorius) with a

20x objective in phase contrast. 4 views were captured in each well every 4 h. PrP samples

were diluted 1:10 into fresh media and added at the 24 h mark.

Neurite lengths were evaluated using the NeuroTrack module of the IncuCyte S3 software

package (rev 2019A) using the following parameters: cell body cluster segmentation 0.7;

cleanup 0; cluster filter 0; neurite sensitivity 0.25; neurite width 1 μm. Detected neurite

masks are highlighted in pink in images. Neurite length data were normalized to the initial (0

h) value for each well and means ± standard deviation were calculated from quintuplicate

sample wells. Average neurite lengths at the 80-100 h time interval were visualized in a box

plot.

The number of live neurons was counted after 0 d and 4 d incubation as a secondary toxicity

assay. Data represent averages from four images under each condition and time point,

normalized to the number of neurons present at the time of protein addition in each field of

view. ANOVA statistical analysis was performed in OriginPro 2019.

Förster Resonance Energy Transfer (FRET) imaging

500 μL of 50,000 cells / mL (uninfected S7 and infected iS7 subclones of PK1 N2a

neuroblastoma cells) were plated in 8-well chambered glass coverslips (ThermoScientific

(155411)) and incubated at 37°C / 5% CO2 for 3 days. Cells were then fixed for 15 min at

RT with 3.7% formaldehyde diluted in Dulbecco's phosphate-buffered saline (DPBS; Gibco,
(14190-094)). After washing once with DPBS, cells were treated with ice-cold acetone for 1
min. Cells were washed before treatment with 3.5 M GdnSCN for 10 min. Following five
washes the cells were incubated with anti-syntaxin-6 (Cell Signalling Technologies (#2869),
clone C34B2, 1:300) and/or anti-PrP (Biolegend (808001) clone 6D11, 1:10,000; Santa Cruz
(sc-47730), 5B2, 1:500; Sigma (P0110), clone 8H4, 1:500) in sterile-filtered 0.25X
SuperBlock in PBS overnight at 4°C, followed by secondary antibodies (Alexa Fluor 647-
AffiniPure Goat Anti-Rabbit IgG (H+L) and/or Rhodamine Red™-X (RRX) AffiniPure Goat
Anti-Mouse IgG (H+L), 1:1000) and a DNA counterstain (DAPI; 1:10,000) in 0.25X
SuperBlock overnight at 4°C. Antibodies were removed and following one wash imaging
was performed with a Zeiss LSM 710 laser-scanning microscope with oil-immersion 63x/1.4
NA objective.

Four images were acquired for each sample: a) donor (excitation 561 nm, emission 580-610
nm); b) acceptor (633 nm excitation, 670-800 nm emission, c) FRET excitation 561 nm,
emission 670-800 nm, all using the MBS 488/561/633 triple dichroic mirror, and d) DAPI
(405 nm excitation, 440-460 nm emission). Laser power, photomultiplier gain and pinhole
size were adjusted as to not exceed the PMT dynamic range. Identical parameters were
used for all image acquisition. Three biological replicates were imaged, with n = 3-6 images
each.

FRET data were analysed using the ImageJ PixFRET plugin according to the developers'
manual (81). Syntaxin-6-only stained iS7 cells were used as acceptor bleed-though control
and 5B2 / 6D11 / 8H4 only stained cells were used as donor bleed-through controls,
respectively. FRET images were calculated at 1 pixel Gaussian blur and rendered using the
Parula HDR lookup table at a range of 0-500.

Cell lines

Neuro2a (N2a) cell line and derivatives - N2a cells (male donor) were sourced from ATCC
(CCL-131). Prion susceptible subclones of N2a cells were derived as described in (35,82)
and authenticated by transcriptomic analysis, where gene expression differences between
original N2a cells and prion-susceptible subclones (PK1 S7, iS7) were investigated and
documented (82). All actively used N2a sublines are tested for mycoplasma contamination
at least once every 2 years. No mycoplasma contaminations have been reported in the MRC
Prion Unit in the last 10 years.

Protein Misfolding Cyclic Amplification (PMCA) using Stx6+/+ and Stx6-/- Mouse Brain
Homogenate

Tissue Collection - Brains for the PMCA substrate were derived from juvenile $Stx6^{+/+}$ and
$Stx6^{-/-}$ male C57BL/6N mice at three months of age. Animals were sacrificed by CO_2
asphyxiation and immediately perfused with 20 mL ice cold perfusion buffer (1x DPBS +
5mM EDTA). The perfused brain was then removed and stored frozen at -70°C until use.

Substrate Preparation - 9% (w/v) mouse brain homogenates were prepared in ice cold
conversion buffer (1x DPBS, 150 mM NaCl, 1% (v/v) Triton X-100) with protease inhibitors
(1x Protease Complete with EDTA) with a dounce, glass homogeniser. Debris was removed
by centrifuging the homogenates at 1200 g for 2 min, with the supernatant subsequently
being collected and stored at -70°C until use.

PMCA - 0.8 µL of seed (I6200 10% (w/v) brain homogenate from terminal RML-infected CD-
1 mice) was spiked into 0.2 mL PCR tubes containing 79.2 µL of $Stx6^{+/+}$ or $Stx6^{-/-}$ substrate
containing three 0.5 mm diameter zirconium beads. Samples were subjected to PMCA in an
automated sonication bath (QSONICA) at 35°C for 96 cycles (50% amplitude, 30 sec
sonication every 30 min) for a total of 48 min sonication over 48 h. Following sonication the
samples were briefly centrifuged before storage at -80°C. Control reactions were also
prepared which were directly frozen.

Immunoblotting - Immunoblotting was performed as previously described with minor
modifications 1. Briefly, each reaction was incubated with proteinase K at 50 µg/mL for 1 h at
37°C. 20 µL of PK digested material was electrophoresed and gels electroblotted.
Membranes were probed with 200 ng/ml ICSM35 anti-PrP antibody (D-Gen Ltd) in PBST for
1 h at RT or overnight at 4°C. After washing the membranes were probed with a 1:10,000
dilution of alkaline-phosphatase-conjugated goat anti-mouse IgG secondary antibody
(Sigma-Aldrich, A2179) in PBST. Blots were incubated for 5 min with CDP-Star™
Chemiluminescent Substrate (Thermo Scientific, T2147) and visualized on Biomax MR film
(Kodak).

Work with animals was performed under the licence granted by the UK Home Office (Project
Licences 70/6454 and 70/7274) and conformed to University College London institutional
and ARRIVE guidelines.

Declarations

Ethics approval

Work with animals was performed under the licence granted by the UK Home Office (Project

Licences 70/6454 and 70/7274) and conformed to University College London institutional

and ARRIVE guidelines.

Consent for publication

All authors have consented to the publication in its present form.

Availability of data and materials

Original data available under doi: 10.17632/yggpkrgnx8.1. All bacterial strains and genetic
constructs are available from the authors upon request.

Competing interests

The authors declare no competing interests.

Funding

Research was supported by the National Institute of Neurological Disorders and Stroke of

the National Institutes of Health grant number 1R21NS101588-01A1, by MRC grant

MC_UU_00024/6 to JB and by the MRC Prion Unit at UCL graduate programme.

References

680   1.   Prusiner, S. B. (1998) Prions. *Proc Natl Acad Sci U S A* **95**, 13363-13383
681   2.   Brundin, P., Melki, R., and Kopito, R. (2010) Prion-like transmission of protein aggregates in
682   neurodegenerative diseases. *Nat Rev Mol Cell Biol* **11**, 301-307
683   3.   Frost, B., and Diamond, M. I. (2010) Prion-like mechanisms in neurodegenerative diseases.
684   *Nat Rev Neurosci* **11**, 155-159
685   4.   Collinge, J., and Clarke, A. R. (2007) A general model of prion strains and their pathogenicity.
686   *Science* **318**, 930-936
687   5.   Manka, S. (2022) 2.7 Å cryo-EM structure of ex vivo RML prion fibrils. *Nature*
688   *Communications* **13**
689   6.   Kraus, A., Hoyt, F., Schwartz, C. L., Hansen, B., Artikis, E., Hughson, A. G., Raymond, G. J.,
690   Race, B., Baron, G. S., and Caughey, B. (2021) High-resolution structure and strain
691   comparison of infectious mammalian prions. *Mol Cell* **81**, 4540-4551 e4546
692   7.   Benilova, I., Reilly, M., Terry, C., Wenborn, A., Schmidt, C., Marinho, A. T., Risse, E., Al-
693   Doujaily, H., Wiggins De Oliveira, M., Sandberg, M. K., Wadsworth, J. D. F., Jat, P. S., and
694   Collinge, J. (2020) Highly infectious prions are not directly neurotoxic. *Proc Natl Acad Sci U S*
695   *A* **117**, 23815-23822
696   8.   Reilly, M., Benilova, I., Khalili-Shirazi, A., Schmidt, C., Ahmed, P., Yip, D., Jat, P. S., and
697   Collinge, J. (2022) A high-content neuron imaging assay demonstrates inhibition of prion
698   disease-associated neurotoxicity by an anti-prion protein antibody. *Sci Rep* **12**, 9493
699   9.   Sandberg, M. K., Al-Doujaily, H., Sharps, B., Clarke, A. R., and Collinge, J. (2011) Prion
700   propagation and toxicity in vivo occur in two distinct mechanistic phases. *Nature* **470**, 540-
701   542
702   10.   Sandberg, M. K., Al-Doujaily, H., Sharps, B., De Oliveira, M. W., Schmidt, C., Richard-Londt,
703   A., Lyall, S., Linehan, J. M., Brandner, S., Wadsworth, J. D., Clarke, A. R., and Collinge, J.
704   (2014) Prion neuropathology follows the accumulation of alternate prion protein isoforms
705   after infective titre has peaked. *Nat Commun* **5**, 4347
706   11.   Collinge, J. (2016) Mammalian prions and their wider relevance in neurodegenerative
707   diseases. *Nature* **539**, 217-226
708   12.   Haass, C., and Selkoe, D. J. (2007) Soluble protein oligomers in neurodegeneration: lessons
709   from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* **8**, 101-112
710   13.   Corbett, G. T., Wang, Z., Hong, W., Colom-Cadena, M., Rose, J., Liao, M., Asfaw, A., Hall, T. C.,
711   Ding, L., DeSousa, A., Frosch, M. P., Collinge, J., Harris, D. A., Perkinton, M. S., Spires-Jones, T.
712   L., Young-Pearse, T. L., Billinton, A., and Walsh, D. M. (2020) PrP is a central player in toxicity
713   mediated by soluble aggregates of neurodegeneration-causing proteins. *Acta Neuropathol*
714   **139**, 503-526
715   14.   Mead, S., Lloyd, S., and Collinge, J. (2019) Genetic Factors in Mammalian Prion Diseases.
716   *Annu Rev Genet* **53**, 117-147
717   15.   Jones, E., Hummerich, H., Vire, E., Uphill, J., Dimitriadis, A., Speedy, H., Campbell, T.,
718   Norsworthy, P., Quinn, L., Whitfield, J., Linehan, J., Jaunmuktane, Z., Brandner, S., Jat, P.,
719   Nihat, A., How Mok, T., Ahmed, P., Collins, S., Stehmann, C., Sarros, S., Kovacs, G. G.,
720   Geschwind, M. D., Golubjatnikov, A., Frontzek, K., Budka, H., Aguzzi, A., Karamujic-Comic, H.,
721   van der Lee, S. J., Ibrahim-Verbaas, C. A., van Duijn, C. M., Sikorska, B., Golanska, E., Liberski,
722   P. P., Calero, M., Calero, O., Sanchez-Juan, P., Salas, A., Martinon-Torres, F., Bouaziz-Amar,
723   E., Haik, S., Laplanche, J. L., Brandel, J. P., Amouyel, P., Lambert, J. C., Parchi, P., Bartoletti-
724   Stella, A., Capellari, S., Poleggi, A., Ladogana, A., Pocchiari, M., Aneli, S., Matullo, G., Knight,
725   R., Zafar, S., Zerr, I., Booth, S., Coulthart, M. B., Jansen, G. H., Glisic, K., Blevins, J., Gambetti,
726   P., Safar, J., Appleby, B., Collinge, J., and Mead, S. (2020) Identification of novel risk loci and

causal insights for sporadic Creutzfeldt-Jakob disease: a genome-wide association study.
Lancet Neurol **19**, 840-848

16.   Meissner, B., Kallenberg, K., Sanchez-Juan, P., Ramljak, S., Krasnianski, A., Heinemann, U.,
Eigenbrod, S., Gelpi, E., Barsic, B., Kretzschmar, H. A., Schulz-Schaeffer, W. J., Knauth, M.,
and Zerr, I. (2009) MRI and clinical syndrome in dura mater-related Creutzfeldt-Jakob
disease. *J Neurol* **256**, 355-363

17.   Jung, J. J., Inamdar, S. M., Tiwari, A., and Choudhury, A. (2012) Regulation of intracellular
membrane trafficking and cell dynamics by syntaxin-6. *Biosci Rep* **32**, 383-391

18.   Wendler, F., and Tooze, S. (2001) Syntaxin 6: the promiscuous behaviour of a SNARE protein.
Traffic **2**, 606-611

19.   Goold, R., Rabbanian, S., Sutton, L., Andre, R., Arora, P., Moonga, J., Clarke, A. R., Schiavo, G.,
Jat, P., Collinge, J., and Tabrizi, S. J. (2011) Rapid cell-surface prion protein conversion
revealed using a novel cell system. *Nat Commun* **2**, 281

20.   Yamasaki, T., Suzuki, A., Hasebe, R., and Horiuchi, M. (2018) Retrograde Transport by
Clathrin-Coated Vesicles is Involved in Intracellular Transport of PrP(Sc) in Persistently Prion-
Infected Cells. *Sci Rep* **8**, 12241

21.   Yim, Y. I., Park, B. C., Yadavalli, R., Zhao, X., Eisenberg, E., and Greene, L. E. (2015) The
multivesicular body is the major internal site of prion conversion. *J Cell Sci* **128**, 1434-1443

22.   Hoglinger, G. U., Melhem, N. M., Dickson, D. W., Sleiman, P. M., Wang, L. S., Klei, L.,
Rademakers, R., de Silva, R., Litvan, I., Riley, D. E., van Swieten, J. C., Heutink, P., Wszolek, Z.
K., Uitti, R. J., Vandrovcova, J., Hurtig, H. I., Gross, R. G., Maetzler, W., Goldwurm, S., Tolosa,
E., Borroni, B., Pastor, P., Group, P. S. P. G. S., Cantwell, L. B., Han, M. R., Dillman, A., van der
Brug, M. P., Gibbs, J. R., Cookson, M. R., Hernandez, D. G., Singleton, A. B., Farrer, M. J., Yu,
C. E., Golbe, L. I., Revesz, T., Hardy, J., Lees, A. J., Devlin, B., Hakonarson, H., Muller, U., and
Schellenberg, G. D. (2011) Identification of common variants influencing risk of the
tauopathy progressive supranuclear palsy. *Nat Genet* **43**, 699-705

23.   Lee, W. S., Tan, D. C., Deng, Y., van Hummel, A., Ippati, S., Stevens, C., Carmona-Mora, P.,
Ariawan, D., Hou, L., Stefen, H., Tomanic, T., Bi, M., Tomasetig, F., Martin, A., Fath, T.,
Palmer, S., Ke, Y. D., and Ittner, L. M. (2021) Syntaxins 6 and 8 facilitate tau into secretory
pathways. *Biochem J* **478**, 1471-1484

24.   Wingo, A. P., Liu, Y., Gerasimov, E. S., Gockley, J., Logsdon, B. A., Duong, D. M., Dammer, E.
B., Robins, C., Beach, T. G., Reiman, E. M., Epstein, M. P., De Jager, P. L., Lah, J. J., Bennett, D.
A., Seyfried, N. T., Levey, A. I., and Wingo, T. S. (2021) Integrating human brain proteomes
with genome-wide association data implicates new proteins in Alzheimer's disease
pathogenesis. *Nat Genet* **53**, 143-146

25.   Hartl, F. U. (2011) Chaperone-assisted protein folding: the path to discovery from a personal
perspective. *Nat Med* **17**, 1206-1210

26.   Balch, W. E., Morimoto, R. I., Dillin, A., and Kelly, J. W. (2008) Adapting proteostasis for
disease intervention. *Science* **319**, 916-919

27.   Glover, J. R., and Lindquist, S. (1998) Hsp104, Hsp70, and Hsp40: a novel chaperone system
that rescues previously aggregated proteins. *Cell* **94**, 73-82

28.   Thackray, A. M., Lam, B., McNulty, E. E., Nalls, A. V., Mathiason, C. K., Magadi, S. S., Jackson,
W. S., Andreoletti, O., Marrero-Winkens, C., Schatzl, H., and Bujdoso, R. (2022) Clearance of
variant Creutzfeldt-Jakob disease prions in vivo by the Hsp70 disaggregase system. *Brain*

29.   Atarashi, R., Sano, K., Satoh, K., and Nishida, N. (2011) Real-time quaking-induced
conversion: a highly sensitive assay for prion detection. *Prion* **5**, 150-153

30.   Legname, G., Baskakov, I. V., Nguyen, H. O., Riesner, D., Cohen, F. E., DeArmond, S. J., and
Prusiner, S. B. (2004) Synthetic mammalian prions. *Science* **305**, 673-676

31.   Post, K., Pitschke, M., Schafer, O., Wille, H., Appel, T. R., Kirsch, D., Mehlhorn, I., Serban, H.,
Prusiner, S. B., and Riesner, D. (1998) Rapid acquisition of beta-sheet structure in the prion
protein prior to multimer formation. *Biol Chem* **379**, 1307-1317

32.   Ziaunys, M., Sakalauskas, A., Mikalauskaite, K., Snieckute, R., and Smirnovas, V. (2021)
779   Temperature-Dependent Structural Variability of Prion Protein Amyloid Fibrils. *International*
780   *Journal of Molecular Sciences* **22**

781   33.   Powers, E. T., and Powers, D. L. (2008) Mechanisms of protein fibril formation: nucleated
782   polymerization with competing off-pathway aggregation. *Biophys J* **94**, 379-391

783   34.   Spehar, K., Ding, T., Sun, Y., Kedia, N., Lu, J., Nahass, G. R., Lew, M. D., and Bieschke, J. (2018)
784   Super-resolution Imaging of Amyloid Structures over Extended Times by Using Transient
785   Binding of Single Thioflavin T Molecules. *Chembiochem* **19**, 1944-1948

786   35.   Klohn, P. C., Stoltze, L., Flechsig, E., Enari, M., and Weissmann, C. (2003) A quantitative,
787   highly sensitive cell-based infectivity assay for mouse scrapie prions. *Proc Natl Acad Sci U S A*
788   **100**, 11666-11671

789   36.   Bieschke, J., Weber, P., Sarafoff, N., Beekes, M., Giese, A., and Kretzschmar, H. (2004)
790   Autocatalytic self-propagation of misfolded prion protein. *Proc Natl Acad Sci U S A* **101**,
791   12207-12211

792   37.   Jones, E., Hill, E., Linehan, J., Nazari, T., Caulder, A., Codner, G. F., Hutchison, M., Mackenzie,
793   M., Wiggins De Oliveira, M. A.-D., Huda , Sandberg, M., Viré, E., Cunningham, T. J., Asante, E.
794   A. B., Sebastian , Collinge, J., and Mead, S. (2023) Knockout of Sporadic Creutzfeldt-Jakob
795   Disease Risk Gene Stx6 in Mice Extends Prion Disease Incubation Time. *bioRxiv*

796   38.   Schmidt, C., Fizet, J., Properzi, F., Batchelor, M., Sandberg, M. K., Edgeworth, J. A., Afran, L.,
797   Ho, S., Badhan, A., Klier, S., Linehan, J. M., Brandner, S., Hosszu, L. L., Tattum, M. H., Jat, P.,
798   Clarke, A. R., Klohn, P. C., Wadsworth, J. D., Jackson, G. S., and Collinge, J. (2015) A
799   systematic investigation of production of synthetic prions from recombinant prion protein.
800   *Open Biol* **5**, 150165

801   39.   Swietnicki, W., Morillas, M., Chen, S. G., Gambetti, P., and Surewicz, W. K. (2000)
802   Aggregation and fibrillization of the recombinant human prion protein huPrP90-231.
803   *Biochemistry* **39**, 424-431

804   40.   Stohr, J., Weinmann, N., Wille, H., Kaimann, T., Nagel-Steger, L., Birkmann, E., Panza, G.,
805   Prusiner, S. B., Eigen, M., and Riesner, D. (2008) Mechanisms of prion protein assembly into
806   amyloid. *Proc Natl Acad Sci U S A* **105**, 2409-2414

807   41.   Atarashi, R., Moore, R. A., Sim, V. L., Hughson, A. G., Dorward, D. W., Onwubiko, H. A., Priola,
808   S. A., and Caughey, B. (2007) Ultrasensitive detection of scrapie prion protein using seeded
809   conversion of recombinant prion protein. *Nat Methods* **4**, 645-650

810   42.   Willbold, D., Strodel, B., Schroder, G. F., Hoyer, W., and Heise, H. (2021) Amyloid-type
811   Protein Aggregation and Prion-like Properties of Amyloids. *Chem Rev* **121**, 8285-8307

812   43.   Cohen, S. I., Linse, S., Luheshi, L. M., Hellstrand, E., White, D. A., Rajah, L., Otzen, D. E.,
813   Vendruscolo, M., Dobson, C. M., and Knowles, T. P. (2013) Proliferation of amyloid-beta42
814   aggregates occurs through a secondary nucleation mechanism. *Proc Natl Acad Sci U S A* **110**,
815   9758-9763

816   44.   Buell, A. K., Galvagnion, C., Gaspar, R., Sparr, E., Vendruscolo, M., Knowles, T. P., Linse, S.,
817   and Dobson, C. M. (2014) Solution conditions determine the relative importance of
818   nucleation and growth processes in alpha-synuclein aggregation. *Proc Natl Acad Sci U S A*
819   **111**, 7671-7676

820   45.   Milto, K., Michailova, K., and Smirnovas, V. (2014) Elongation of Mouse Prion Protein
821   Amyloid-Like Fibrils: Effect of Temperature and Denaturant Concentration. *Plos One* **9**

822   46.   Jarrett, J. T., and Lansbury, P. T., Jr. (1993) Seeding "one-dimensional crystallization" of
823   amyloid: a pathogenic mechanism in Alzheimer's disease and scrapie? *Cell* **73**, 1055-1058

824   47.   Cohen, E., Bieschke, J., Perciavalle, R. M., Kelly, J. W., and Dillin, A. (2006) Opposing activities
825   protect against age-onset proteotoxicity. *Science* **313**, 1604-1610

826   48.   Terry, C., Wenborn, A., Gros, N., Sells, J., Joiner, S., Hosszu, L. L., Tattum, M. H., Panico, S.,
827   Clare, D. K., Collinge, J., Saibil, H. R., and Wadsworth, J. D. (2016) Ex vivo mammalian prions
828   are formed of paired double helical prion protein fibrils. *Open Biol* **6**

49. Du, D., Murray, A. N., Cohen, E., Kim, H. E., Simkovsky, R., Dillin, A., and Kelly, J. W. (2011) A
kinetic aggregation assay allowing selective and sensitive amyloid-beta quantification in cells
and tissues. *Biochemistry* **50**, 1607-1617

50. Jain, S., and Udgaonkar, J. B. (2008) Evidence for stepwise formation of amyloid fibrils by the
mouse prion protein. *J Mol Biol* **382**, 1228-1241

51. Honda, R. P., and Kuwata, K. (2017) The native state of prion protein (PrP) directly inhibits
formation of PrP-amyloid fibrils in vitro. *Scientific Reports* **7**

52. Bieschke, J., Zhang, Q., Powers, E. T., Lerner, R. A., and Kelly, J. W. (2005) Oxidative
metabolites accelerate Alzheimer's amyloidogenesis by a two-step mechanism, eliminating
the requirement for nucleation. *Biochemistry* **44**, 4977-4983

53. Klement, K., Wieligmann, K., Meinhardt, J., Hortschansky, P., Richter, W., and Fandrich, M.
(2007) Effect of different salt ions on the propensity of aggregation and on the structure of
Alzheimer's abeta(1-40) amyloid fibrils. *J Mol Biol* **373**, 1321-1333

54. Hartl, F. U. (1996) Molecular chaperones in cellular protein folding. *Nature* **381**, 571-579

55. Muchowski, P. J., and Wacker, J. L. (2005) Modulation of neurodegeneration by molecular
chaperones. *Nat Rev Neurosci* **6**, 11-22

56. Evans, C. G., Wisen, S., and Gestwicki, J. E. (2006) Heat shock proteins 70 and 90 inhibit early
stages of amyloid beta-(1-42) aggregation in vitro. *J Biol Chem* **281**, 33182-33191

57. Wacker, J. L., Zareie, M. H., Fong, H., Sarikaya, M., and Muchowski, P. J. (2004) Hsp70 and
Hsp40 attenuate formation of spherical and annular polyglutamine oligomers by partitioning
monomer. *Nat Struct Mol Biol* **11**, 1215-1222

58. Rosas, P. C., Nagaraja, G. M., Kaur, P., Panossian, A., Wickman, G., Garcia, L. R., Al-Khamis, F.
A., and Asea, A. A. (2016) Hsp72 (HSPA1A) Prevents Human Islet Amyloid Polypeptide
Aggregation and Toxicity: A New Approach for Type 2 Diabetes Treatment. *PLoS One* **11**,
e0149409

59. Bongiovanni, M. N., Aprile, F. A., Sormanni, P., and Vendruscolo, M. (2018) A Rationally
Designed Hsp70 Variant Rescues the Aggregation-Associated Toxicity of Human IAPP in
Cultured Pancreatic Islet beta-Cells. *Int J Mol Sci* **19**

60. Fernandez-Funez, P., Sanchez-Garcia, J., de Mena, L., Zhang, Y., Levites, Y., Khare, S., Golde,
T. E., and Rincon-Limas, D. E. (2016) Holdase activity of secreted Hsp70 masks amyloid-
beta42 neurotoxicity in Drosophila. *Proc Natl Acad Sci U S A* **113**, E5212-5221

61. Bieschke, J., Russ, J., Friedrich, R. P., Ehrnhoefer, D. E., Wobst, H., Neugebauer, K., and
Wanker, E. E. (2010) EGCG remodels mature alpha-synuclein and amyloid-beta fibrils and
reduces cellular toxicity. *Proc Natl Acad Sci U S A* **107**, 7710-7715

62. Ehrnhoefer, D. E., Bieschke, J., Boeddrich, A., Herbst, M., Masino, L., Lurz, R., Engemann, S.,
Pastore, A., and Wanker, E. E. (2008) EGCG redirects amyloidogenic polypeptides into
unstructured, off-pathway oligomers. *Nat Struct Mol Biol* **15**, 558-566

63. Bieschke, J., Herbst, M., Wiglenda, T., Friedrich, R. P., Boeddrich, A., Schiele, F., Kleckers, D.,
Lopez del Amo, J. M., Gruning, B. A., Wang, Q., Schmidt, M. R., Lurz, R., Anwyl, R., Schnoegl,
S., Fandrich, M., Frank, R. F., Reif, B., Gunther, S., Walsh, D. M., and Wanker, E. E. (2012)
Small-molecule conversion of toxic oligomers to nontoxic beta-sheet-rich amyloid fibrils. *Nat*
Chem Biol **8**, 93-101

64. Lam, H. T., Graber, M. C., Gentry, K. A., and Bieschke, J. (2016) Stabilization of alpha-
Synuclein Fibril Clusters Prevents Fragmentation and Reduces Seeding Activity and Toxicity.
Biochemistry **55**, 675-685

65. Cohen, S. I., Arosio, P., Presto, J., Kurudenkandy, F. R., Biverstal, H., Dolfe, L., Dunning, C.,
Yang, X., Frohm, B., Vendruscolo, M., Johansson, J., Dobson, C. M., Fisahn, A., Knowles, T. P.,
and Linse, S. (2015) A molecular chaperone breaks the catalytic cycle that generates toxic
Abeta oligomers. *Nat Struct Mol Biol* **22**, 207-213

66. Amin, L., and Harris, D. A. (2021) Abeta receptors specifically recognize molecular features
displayed by fibril ends and neurotoxic oligomers. *Nat Commun* **12**, 3451

67. Bove-Fenderson, E., Urano, R., Straub, J. E., and Harris, D. A. (2017) Cellular prion protein
targets amyloid-beta fibril ends via its C-terminal domain to prevent elongation. *J Biol Chem*
292, 16858-16871
68. Freir, D. B., Nicoll, A. J., Klyubin, I., Panico, S., Mc Donald, J. M., Risse, E., Asante, E. A.,
Farrow, M. A., Sessions, R. B., Saibil, H. R., Clarke, A. R., Rowan, M. J., Walsh, D. M., and
Collinge, J. (2011) Interaction between prion protein and toxic amyloid beta assemblies can
be therapeutically targeted at multiple sites. *Nat Commun* **2**, 336
69. Nicoll, A. J., Panico, S., Freir, D. B., Wright, D., Terry, C., Risse, E., Herron, C. E., O'Malley, T.,
Wadsworth, J. D., Farrow, M. A., Walsh, D. M., Saibil, H. R., and Collinge, J. (2013) Amyloid-
beta nanotubes are associated with prion protein-dependent synaptotoxicity. *Nat Commun*
4, 2416
70. Grassmann, A., Wolf, H., Hofmann, J., Graham, J., and Vorberg, I. (2013) Cellular aspects of
prion replication in vitro. *Viruses* **5**, 374-405
71. Choudhury, A., Marks, D. L., Proctor, K. M., Gould, G. W., and Pagano, R. E. (2006) Regulation
of caveolar endocytosis by syntaxin 6-dependent delivery of membrane components to the
cell surface. *Nat Cell Biol* **8**, 317-328
72. Hegde, R. S., Mastrianni, J. A., Scott, M. R., DeFea, K. A., Tremblay, P., Torchia, M.,
DeArmond, S. J., Prusiner, S. B., and Lingappa, V. R. (1998) A transmembrane form of the
prion protein in neurodegenerative disease. *Science* **279**, 827-834
73. Stewart, R. S., and Harris, D. A. (2005) A transmembrane form of the prion protein is
localized in the Golgi apparatus of neurons. *J Biol Chem* **280**, 15855-15864
74. Shi, Z., Sachs, J. N., Rhoades, E., and Baumgart, T. (2015) Biophysics of alpha-synuclein
induced membrane remodelling. *Phys Chem Chem Phys* **17**, 15561-15568
75. Lashuel, H. A., Hartley, D., Petre, B. M., Walz, T., and Lansbury, P. T., Jr. (2002)
Neurodegenerative disease: amyloid pores from pathogenic mutations. *Nature* **418**, 291
76. Jin, S., Kedia, N., Illes-Toth, E., Haralampiev, I., Prisner, S., Herrmann, A., Wanker, E. E., and
Bieschke, J. (2016) Amyloid-beta(1-42) Aggregation Initiates Its Cellular Uptake and
Cytotoxicity. *J Biol Chem* **291**, 19590-19606
77. Friedrich, R. P., Tepper, K., Ronicke, R., Soom, M., Westermann, M., Reymann, K., Kaether,
C., and Fandrich, M. (2010) Mechanism of amyloid plaque formation suggests an
intracellular basis of Abeta pathogenicity. *Proc Natl Acad Sci U S A* **107**, 1942-1947
78. Hosszu, L. L., Wells, M. A., Jackson, G. S., Jones, S., Batchelor, M., Clarke, A. R., Craven, C. J.,
Waltho, J. P., and Collinge, J. (2005) Definable equilibrium states in the folding of human
prion protein. *Biochemistry* **44**, 16649-16657
79. Jackson, G. S., Hill, A. F., Joseph, C., Hosszu, L., Power, A., Waltho, J. P., Clarke, A. R., and
Collinge, J. (1999) Multiple folding pathways for heterologously expressed human prion
protein. *Biochim Biophys Acta* **1431**, 1-13
80. Sun, Y. (2022) Direct Observation of Prion Protein Fibril Elongation Kinetics Reveals
Competing Fibril Populations with Distinct Structural and Dynamic Properties. *BioRxiv*
81. Feige, J. N., Sage, D., Wahli, W., Desvergne, B., and Gelman, L. (2005) PixFRET, an ImageJ
plug-in for FRET calculation that can accommodate variations in spectral bleed-throughs.
Microsc Res Tech **68**, 51-58
82. Marbiah, M. M., Harvey, A., West, B. T., Louzolo, A., Banerjee, P., Alden, J., Grigoriadis, A.,
Hummerich, H., Kan, H. M., Cai, Y., Bloom, G. S., Jat, P., Collinge, J., and Klohn, P. C. (2014)
Identification of a gene regulatory network associated with prion replication. *EMBO J* **33**,
1527-1547
83. Castle, A. R., and Gill, A. C. (2017) Physiological Functions of the Cellular Prion Protein. *Front*
Mol Biosci **4**, 19

Figure 1

Figure 1: Native aggregation assay (NAA) of human PrP (23-231). (A) Seed titration
experiment using seed generated from hPrP23 fibrils formed de novo in NAA, seed fraction
(w/w), 2.5 µM hPrP23, pH 6.8, 42°C. (B) Plot of lag phase vs. fraction of seed, n = 3. (C)
TEM image of hPrP23 fibrils formed after 166 h incubation. (D-G) Titration of PrP
concentration (0.3 µM - 20 µM) in ThT aggregation assays containing 0.01% seed, pH 6.8,
42°C. (D) Plot of ThT fluorescence vs. time. (E) Plot of lag time t_{50} vs. PrP concentration. (F)
Plot of elongation rate constant k vs. PrP concentration. (G) Plot of fluorescence amplitude A
vs. PrP concentration.

Figure 2

Figure 2: Syntaxin-6 (STX6) delays hPrP 23 fibril formation at molar ratios of 1:1 –
1:100; 2.5 µM hPrP23, pH 6.8, 42°C, 0.01% seed. (A) ThT fluorescence vs time. (B) Plot of
lag phase t_{50} vs. molar ratio of STX6 / PrP, n=3. (C) Plot of elongation rate constant k vs.
molar ratio of syntaxin-6 / PrP. (D) TEM image of hPrP23 co-aggregated with syntaxin-6 at
1:10 (STX6 / PrP) molar ratio under standard NAA conditions for 116 h. (E) Immuno-TEM
image of hPrP23 – syntaxin-6 co-aggregate cluster after 100 h aggregation. Syntaxin-6 is
labelled with anti-syntaxin-6 Ab / 10 nm anti-rabbit immunogold beads); scale bars 200 nm.

Figure 3

Figure 3: Imaging of hPrP23 fibrils incubated with syntaxin-6. hPrP23 (2.5 μM) was pre-
aggregated in NAA for 115 h, and incubated with syntaxin-6 (250 nM) for 1 h. hPrP was
visualized by TAB imaging using 10 nM Nile red dye; syntaxin-6 was labelled with
AlexaFluor488 (A-D) or AlexaFluor647 (E) and imaged by dSTORM. (A) SR image overlay
shows syntaxin-6 binding at hotspots and PrP fibril ends. (B) Magnified area from (A). (C)
Widefield image taken with green laser (561 nm) illumination. (D) Widefield image taken
under blue laser (473 nm) illumination. (E) SR image overlay of PrP and syntaxin-6-AF647
images; scale bars 2 μm. (F) TEM Immuno-gold staining of hPrP23 fibrils incubated with
syntaxin-6; scale bar 100 nm.

 Figure 4

Figure 4. Interaction of PrP and syntaxin-6 in vivo. Non-infected PK1 cells and persistently infected PK1 cells (iS7) were immuno-stained with anti-PrP antibodies 6D11 and 5B2 (green), with anti-syntaxin-6 antibody (red) and with DAPI (blue). FRET analysis reveals interaction in perinuclear compartments (wide arrows) and at membranes in infected cells (narrow arrows). Panels show zoomed regions of images in Figure 4 – figure supplement 1. (B, C) In vitro prion replication by PMCA using $Stx6^{+/+}$ and $Stx6^{-/-}$ mouse brains as substrate. PMCA reactions were seeded with RML prions from terminally ill mice and subjected to PMCA for 96 cycles over 48 h. (B) Representative Western blot (ICSM35) after PK digestion. Molecular weight markers are indicated on the left. (C) The PrP^{Sc} signal was quantified using densitometry and normalized to the control unamplified reaction. Bar graphs each represent mean ± SD of biological replicates from three separate mice, each blotted as two technical replicates.

Figure 5

Figure 5: Toxicity in mouse primary neurons incubated with hPrP23 / syntaxin-6 at
different aggregation time points. NAA was performed as in Figure 2 and samples were
diluted 1:10 into cell culture media at the indicated time points (0 h, 20 h, 40 h, 64 h, 90 h).
(A) Plot of effect of PrP (250 nM) on neurite length compared to effect of media or NAA
assay buffer at the same dilution from four independent wells. (B) Plot of effect of hPrP23
co-aggregated with syntaxin-6 (STX6, 1 in 10 molar ratio). (C) Box plot of average relative
neurite lengths between 80-100h incubation of PrP in the cell culture media. P-values are
derived from ANOVA testing between PrP vs PrP/syntaxin-6 samples at each time point.

Figure 1 – figure supplement 1

Figure 1 – figure supplement 1: Protein characterization. (A) Coomassie/silver stained
gels of purified hPrP23, mPrP23 and candidate proteins. (B) Circular dichroism spectrum of
purified syntaxin-6 (STX6) protein showing a mostly alpha helical shape with minima at 208
nm and 223 nm. (C - E) Circular dichroism spectra of Hsp70 (HSPA1A), stathmin 1
(STMN1), human PrP 23-231,129M (hPrP23), and mouse PrP 23-231 (mPrP23).

     Figure 1 – figure supplement 2

Figure 1 – figure supplement 2: Optimisation of NAA. (A) Plot of mPrP23 ThT
fluorescence over time at 37°C, 42°C and 50°C (pH 7.4). (B) mPrP23 NAA aggregation
kinetics at pH 6.6, 6.8, 7.0 and 7.2 at 42°C. (C) NAA aggregation kinetics of hPrP23 (2.5 µM)
with Zr vs glass beads, pH 6.8 at 42°C. (D-F) mPrP23 seed titration experiments comparing
seeding from synthetic PrP fibrils (Sp) and purified prion rods from mouse prion strains ME7
and RML. (D) Plot of time to half-maximal fluorescence (t_{50}) (E) elongation rate constant (k),
and (F) ThT amplitudes as functions of seed concentration, n=5. (G) TEM image of mPrP23
fibrils formed after 47 h incubation at pH 6.8, 42°C; scale bar 100 nm.

1005

1006

1007 **Figure 1 – figure supplement 3**

1008

Figure 1 – figure supplement 3: mPrP23 seeding kinetics. Graphs of NAA on mPrP23
(10 µM= seed titration experiments demonstrate seed concentration-dependent acceleration
of fibril formation. (A) First generation (spontaneous) seed. (B) Seed generated from RML
mouse strain (C) Seed generated from ME7 mouse strain. Solid lines indicate sigmoidal fits
of the data. Post aggregation decay of ThT fluorescence, which was likely due to adsorption
and sedimentation, was not included in the fits.

Figure 2 – figure supplement 1

PrP + HSPA1A

PrP + STMN1

1016

1017   **Figure 2 – figure supplement 1: TEM images of NAA aggregation end-points of hPrP23**
1018   **co-aggregated with HSP70 (HSPA1A) or stathmin 1 (STMN1).** (A) PrP forms amorphous
1019   aggregates in presence of HSPA1A at 1 in 10 molar ratio. (B) PrP forms fibrillar aggregates
1020   in presence of STMN1 at 1 in 10 molar ratio; scale bar 100 nm.

1021

1022   **Figure 2 – figure supplement 2**

1023

1024

1025

1026   **Figure 2 – figure supplement 2: Sedimentation assay of hPrP23.** hPrP23 was incubated
1027   for 0 – 90 h alone or co-aggregated with syntaxin-6 (1 in 10 molar ratio) or HSPA1A (1 in 10
1028   molar ratio). Protein samples were centrifuged; supernatants were collected and analysed by
1029   silver stained SDS-PAGE.

1030

1031   **Figure 2 – figure supplement 3**

1032

Figure 2 – figure supplement 3: Salt-dependent inhibition of fibril formation. Syntaxin-
6, STMN1, HSPA1A 1:10 molar ratio were incubated with hPrP23. (A) Plot of relative ThT
fluorescence profiles vs time in NAA (150 mM NaCl, 42°C, 0.01% seed,). (B) Plot of lag
phase t_{50} vs NaCl concentration for hPrP23 alone vs co-aggregated with syntaxin-6 (STX6)
at 1:10 molar ratio. (C) Plot of elongation rate constant k vs. NaCl concentration. (D) Plot of
lag phase t_{50} vs NaCl concentration for hPrP23 alone vs co-aggregated with HSPA1A at
1:10 molar ratio. (E) Plot of lag phase t_{50} vs NaCl concentration for hPrP 23-231 alone vs co-
aggregated with STMN1 at 1:10 molar ratio.

 Figure 3 – figure supplement 1

1042

Figure 3 – figure supplement 1: Overlay of dSTORM SR images taken at 473 nm and 638 nm excitation. Co-aggregates of hPrP23 and syntaxin-6 formed in NAA under conditions of Figure 3. (A) Co-aggregation of hPrP23 labelled with AlexaFluor 647-NHS (2.5 µM, 3% labelled, 95% unlabelled) with 10 nM syntaxin-6 labelled with AlexaFluor488-NHS (B) Co-aggregation of PrP labelled with AlexaFluor 488-NHS (2.5 µM, 5% labelled, 95% unlabelled) with syntaxin-6-AlexaFluor647 at 1:10 molar ratio (syntaxin-6, 250 nM, 4% labelled, 96% unlabelled). *Insets*: wide field images with 473 nm and 638 nm excitation and composite image of both channels.

1051   **Figure 3 – figure supplement 2**

1052

1053

1054   **Figure 3 – figure supplement 2: TAB SR microscopy images of aggregation endpoints**
1055   **of hPrP23.** hPrP23 was aggregated in NAA with increasing salt concentrations from 0 mM to
1056   600 mM NaCl for 140 h; scale bars 2 μm.

1057   **Figure 3 – figure supplement 3**

1058

1059   **Figure 3 – figure supplement 3: TAB SR microscopy images of aggregation endpoints**
1060   **of hPrP23 co-aggregated with syntaxin-6.** NAA was performed for hPrP23 / syntaxin-6 co-
1061   aggregation (1:0.1 molar ratio) at increasing salt concentrations from 0 mM to 600 mM NaCl
1062   for 140 h; scale bars 2 µm.

1063

     Figure 4 – figure supplement 1

1065

Figure 4 – figure supplement 1: hPrP / syntaxin-6 FRET analysis. PixFRET analysis of
non-infected PK1 cells and persistently infected PK1 cells (iS7). Cells were immuno-stained
with anti-PrP antibodies 6D11, 8H4 or 5B2 (green), with anti-syntaxin-6 antibody (STX6, red)
and with DAPI. (C) Schematic of binding locations of antibodies used in FRET analysis (5B2,

1070 6D11, 8H4). Numbers represent putative epitopes. Figure 4 - figure supplement 1C is
1071 adapted from Figure 2B from Castle and Gill, 2017 (83).

1072 **Figure 4 – figure supplement 2**

1073

1074 **Figure 4 – figure supplement 2: hPrP/syntaxin-6 FRET analysis controls.** PixFRET
1075 analysis of control cells. iS7 Cells were immuno-stained with anti-PrP antibodies 6D11, 8H4
1076 or 5B2 (green) only, with anti-syntaxin-6 antibody (STX6, red) only, or with secondary
1077 antibodies only and with DAPI.

1078   **Figure 5 – figure supplement 1**

1079
1080   **Figure 5 – figure supplement 1: Survival of primary neurons.** Numbers of live primary
1081   neurons after incubation with hPrP23 or hPrP23 co-aggregated with syntaxin-6 (STX6, 1 in
1082   10 molar ratio) as in Figure 5. Relative numbers of neurons after 4 h incubation in four fields
1083   of view per sample were normalized to the same FOV at t = 0 h; * indicates p < 0.05, ** p ,
1084   0.001, *** p < 0.0001, n.s. not significant from ANOVA statistical analysis.

1085

1086

1087 **Figure 5 – figure supplement 2**

1088

1089

Figure 5 – figure supplement 2: Seeding capacity of aggregation intermediates.
Aggregates of hPrP23 (black) or hPrP23 co-aggregated with syntaxin-6 (STX6, 1 in 10 molar
ratio; orange) collected at 20 h (A), 40 h (B), 60 h (C) and 90 h (D) were used to seed a
second aggregation assay using hPrP23 as substrate; seed concentration 0.1% (w/w).
Aggregation time points correspond to those used in toxicity assays. Graphs represent
averaged data of three replicate experiments.