Modeling tissue co-regulation to estimate tissue-specific contributions to disease

Tiffany Amariuta[1,2], Katherine Siewert-Rocks[1,2], Alkes L. Price[1,2,3]

1. Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA
2. Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA
3. Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, USA

Abstract

Integrative analyses of genome-wide association studies (GWAS) and gene expression data across diverse tissues and cell types have enabled the identification of putative disease-critical tissues. However, co-regulation of genetic effects on gene expression across tissues makes it difficult to distinguish biologically causal tissues from tagging tissues. While previous work emphasized the potential of accounting for tissue co-regulation, tissue-specific disease effects have not previously been formally modeled. Here, we introduce a new method, tissue co-regulation score regression (TCSC), that disentangles causal tissues from tagging tissues and partitions disease heritability (or covariance) into tissue-specific components. TCSC leverages gene-disease association statistics across tissues from transcriptome-wide association studies (TWAS), which implicate both causal and tagging genes and tissues. TCSC regresses TWAS chi-square statistics (or products of z-scores) on tissue co-regulation scores reflecting correlations of predicted gene expression across genes and tissues. In simulations, TCSC powerfully distinguishes causal tissues from tagging tissues while controlling type I error. We applied TCSC to GWAS summary statistics for 78 diseases and complex traits (average N = 302K) and gene expression prediction models for 48 GTEx tissues. TCSC identified 27 causal tissue-trait pairs at 10% FDR, including well-established findings, biologically plausible novel findings (e.g. aorta artery and glaucoma), and increased specificity of known tissue-trait associations (e.g. subcutaneous adipose, but not visceral adipose, and HDL). TCSC also identified 30 causal tissue-trait covariance pairs at 10% FDR. For the positive genetic covariance between eosinophil count and white blood cell count, whole blood contributed positive covariance while LCLs contributed negative covariance; this suggests that genetic covariance may reflect distinct tissue-specific contributions. Overall, TCSC is a powerful method for distinguishing causal tissues from tagging tissues, improving our understanding of disease and complex trait biology.

Introduction

Most diseases are driven by tissue-specific or cell-type-specific mechanisms, thus the inference of causal disease tissues is an important goal[1]. For many polygenic diseases and complex traits, disease-associated tissues have previously been identified via the integration of genome-wide association studies (GWAS) with tissue-level functional data characterizing expression quantitative trait loci (eQTLs)[2-5], gene expression[6-8], or epigenetic features[9-16]. However, it is likely that most disease-associated tissues are not actually causal, due to the high correlation of eQTL effects (resp. gene expression or epigenetic features) across tissues; the correlation of eQTL effects across tissues, i.e. tissue co-regulation, can arise due to shared eQTLs or distinct eQTLs in linkage disequilibrium (LD)[2,17,18,5]. One approach to address this involves comparing eQTL-disease colocalizations across different tissues[2]; however, this approach relies on colocalizations with disease that are specific to a single tissue, and may implicate co-regulated tagging tissues that colocalize with disease. Another approach leverages multi-trait fine-mapping methods to simultaneously evaluate all tissues for colocalization with disease[5]; however, this locus-based approach does not produce genome-wide estimates and it remains the case that many (causal or tagging) tissues may colocalize with disease under this framework. To our knowledge, no previous study has formally modeled genetic co-regulation across tissues to statistically disentangle causal from tagging tissues.

Here, we introduce a new method, tissue co-regulation score regression (TCSC), that disentangles causal tissues from tagging tissues and partitions disease heritability (or genetic covariance of two diseases/traits) into tissue-specific components. TCSC leverages gene-disease association statistics across tissues from transcriptome-wide association studies (TWAS)[19,20,17]. A challenge is that TWAS association statistics include the effects of both co-regulated tissues (see above) and co-regulated genes[17,21]. To address this, TCSC regresses TWAS chi-square statistics (or products of z-scores for two diseases/traits) on tissue co-regulation scores reflecting correlations of predicted gene expression across genes and tissues. (TCSC is conceptually related to gene co-regulation score regression (GCSC)[21], a method for identifying disease-enriched gene sets that models gene co-regulation but does not model tissue co-regulation.) Distinct from previous methods that analyze each tissue marginally, TCSC jointly models contributions from each tissue to identify causal tissues (analogous to the distinction in GWAS between marginal association and fine-mapping[22]). We validate TCSC using extensive simulations and apply TCSC to 78 diseases and complex traits (average N = 302K) and 48 GTEx tissues[18], showing that TCSC recapitulates known biology and identifies biologically plausible novel tissue-trait pairs (or tissue-trait covariance pairs) while attaining increased specificity relative to previous methods.

Results

Overview of TCSC regression

TCSC estimates the disease heritability explained by *cis*-genetic components of gene expression in each tissue when jointly modeling contributions from each tissue. We refer to

tissues with nonzero contributions as "causal" tissues (with the caveat that joint-fit effects of
gene expression on disease may not reflect biological causality; see **Discussion**). TCSC leverages
the fact that TWAS χ^2 statistics for each gene and tissue include both causal effects of that gene
and tissue on disease and tagging effects of *co-regulated* genes and tissues. We define co-
regulation based on squared correlations in *cis*-genetic expression, which can arise due to
shared causal eQTLs and/or LD between causal eQTLs[17]. TCSC determines that a tissue is causal
for disease if genes and tissues with high co-regulation to that tissue have higher TWAS χ^2
statistics than genes and tissues with low co-regulation to that tissue.
100
101 In detail, the expected TWAS χ^2 statistic for gene g and tagging tissue t is

$$E[\chi^2_{g,t}] = N \sum_{t'} l(g,t;t') h^2_{ge(t')} / G_{t'} + 1, \tag{1}$$

where N is GWAS sample size, t' indexes causal tissues, $l(g,t;t')$ are tissue co-regulation scores
(defined as $l(g,t;t') = \sum_{g'} r^2(W_{g,t}, W_{g',t'})$, where W denotes the *cis*-genetic component of
gene expression for a gene-tissue pair across individuals and the sum is over genes g' within +/-
1 Mb to gene g), $h^2_{ge(t')}$ is the disease heritability explained by the *cis*-genetic component of
gene expression in tissue t', and $G_{t'}$ is the number of significantly *cis*-heritable genes in tissue t'
(see below). Equation (1) allows us to estimate $h^2_{ge(t')}$ via a multiple linear regression of TWAS
χ^2 statistics (for each gene and tagging tissue) on tissue co-regulation scores (**Figure 1**). To
facilitate comparisons across diseases/traits, we primarily report the proportion of disease
heritability explained by the *cis*-genetic component of gene expression in tissue t'
($\pi_{t'} = h^2_{ge(t')} / h^2_g$), where h^2_g is the common variant SNP-heritability estimated by S-
LDSC[12,23,24].
114
115   TCSC can also estimate the genetic covariance between two diseases explained by *cis*-
116 genetic components of gene expression in each tissue, using products of TWAS z-scores. In
117 detail, the expected product of TWAS z-scores in disease 1 and disease 2 for gene g and tagging
118 tissue t is
119

$$E[z^1_{gt} \times z^2_{gt}] = \sqrt{N_1 N_2} \sum_{t'} l(g,t;t') \omega_{ge(t')} / G_{t'} + \rho N_s / \sqrt{N_1 N_2} \tag{2}$$

121
where N_1 is GWAS sample size for disease 1, N_2 is GWAS sample size for disease 2, t' indexes
causal tissues, $l(g,t;t')$ are tissue co-regulation scores (see above), $\omega_{ge(t')}$ is the genetic
covariance explained by the *cis*-genetic component of gene expression in tissue t', $G_{t'}$ is the
number of significantly *cis*-heritable genes in tissue t' (see below), ρ is the phenotypic
correlation between disease 1 and disease 2, and N_s is the number of overlapping GWAS
samples between disease 1 and disease 2. Equation (2) allows us to estimate $\omega_{ge(t')}$ via a
multiple linear regression of products of TWAS z-scores in disease 1 and disease 2 (for each
gene and tagging tissue) on tissue co-regulation scores. We note that the last term in Equation
(2) is not known a priori but is accounted for via the regression intercept, analogous to previous
work[25]. To facilitate comparisons across diseases/traits, we primarily report the proportion of

genetic covariance explained by the *cis*-genetic component of gene expression in tissue t'
($\zeta_{t'} = \omega_{ge(t')}/|\omega_g|$), where ω_g is the common variant genetic covariance estimated by cross-
trait LDSC[26].
135
We restrict gene expression prediction models and TWAS association statistics for each
tissue to significantly *cis*-heritable genes in that tissue, defined as genes with significantly
positive *cis*-heritability (2-sided $p < 0.01$; estimated using GCTA[27]) and positive adjusted-R^2 in
cross-validation prediction. We note that quantitative estimates of the disease heritability
explained by the *cis*-genetic component of gene expression in tissue t' ($h^2_{ge(t')}$) are impacted
by the number of significantly *cis*-heritable genes in tissue t' ($G_{t'}$), which may be sensitive to
eQTL sample size. For each disease (or pair of diseases), we use a genomic block-jackknife with
200 blocks to estimate standard errors on the disease heritability (or covariance) explained by
cis-genetic components of gene expression in each tissue, and compute 1-sided P-values for
nonzero heritability (or 2-sided P-values for nonzero covariance) and false discovery rates (FDR)
accordingly; we primarily report causal tissues with FDR < 10%. Further details, including
correcting for bias in tissue co-regulation scores arising from differences between *cis*-genetic vs.
cis-predicted expression (analogous to GCSC[21]) and utilizing regression weights to improve
power, are provided in the **Methods** section. We have publicly released open-source software
implementing TCSC regression (see **Code Availability**), as well as all GWAS summary statistics,
TWAS association statistics, tissue co-regulation scores, and TCSC output from this study (see
Data Availability).
153
154   *Simulations*
155
We performed extensive simulations to evaluate the robustness and power of TCSC,
using the TWAS simulator of Mancuso et al.[28] (see **Code Availability**). We used simulated
genotypes to simulate gene expression values (for each gene and tissue) and complex trait
phenotypes, and computed TWAS association statistics for each gene and tissue. In our default
simulations, the number of tissues was set to 10. The gene expression sample size (in each
tissue) varied from 100 to 1,000 (with the value of 300 corresponding most closely to the GTEx
data[18] used in our analyses of real diseases/traits; see below). The number of genes was set to
1,245, distributed across 249 genomic blocks (representing *cis* regions of length 1 Mb on
chromosome 1) with each block uniformly containing 50 SNPs and 5 genes. Each gene had 5
causal *cis*-eQTLs, consistent with the upper range of independent eQTLs per gene detected in
GTEx[18] and others studies[29-32]. The *cis*-eQTL effect sizes for each gene were drawn from a
multivariate normal distribution across tissues to achieve a specified level of co-regulation (see
below), and the *cis*-heritability of each gene was approximated as the sum of squared
standardized *cis*-eQTL effect sizes. In each tissue, the average *cis*-heritability (across genes) was
tuned to 0.08 (sd = 0.05, ranging from 0.01 to 0.40) in order to achieve an average estimated
cis-heritability (*across significantly cis-heritable genes*) varying from 0.11 to 0.31 (across gene
expression sample sizes), matching GTEx[18]; the proportions of expressed genes that were
significantly *cis*-heritable were also matched to GTEx data[18]. The 10 tissues were split into three
tissue categories to mimic biological tissue modules in GTEx[18] (tissues 1-3, tissues 4-6, and

tissues 7-10), and average *cis*-genetic correlations between tissues (averaged across genes)
were set to 0.789 within the same tissue category, 0.737 between tissue categories, and 0.751
overall[33] (**Methods**). The GWAS sample size was set to 100,000. The 10 tissues included one
causal tissue explaining 5% of complex trait heritability and nine non-causal tissues. 10% of the
genes (125 of the 1,245 genes) had nonzero (normally distributed) gene-trait effects in the
causal tissue[34]. The SNP-heritability of the trait was set to 5%, all of which was explained by
gene expression in the causal tissue; we note that TCSC is not impacted by SNP-heritability that
is not explained by gene expression. Other parameter values were also explored, including
other values of the number of causal tissues and the number of tagging tissues. Further details
of the simulation framework are provided in the **Methods** section. We primarily focused our
comparisons between TCSC and other methods on analyses of real diseases/traits (see below),
because the power of TCSC relative to other methods is likely to be highly sensitive to
assumptions about the role of gene expression in disease architectures. However, we included
comparisons to RTC Coloc[2] (which also uses eQTL data) in our simulations.
189
We first evaluated the bias in TCSC estimates of the disease heritability explained by the
cis-genetic component of gene expression in tissue t' ($h^2_{ge(t')}$), for both causal and non-causal
tissues. For causal tissues, TCSC produced conservative estimates of $h^2_{ge(t')}$, particularly at
smaller eQTL sample sizes (**Figure 2A**, **Supplementary Table 1**). As noted above, estimates of
$h^2_{ge(t')}$ are impacted by the number of significantly *cis*-heritable genes in tissue t' ($G_{t'}$), which
may be sensitive to eQTL sample size. Estimates were more conservative when setting $G_{t'}$ to
the number of significantly *cis*-heritable genes, and less conservative when setting $G_{t'}$ to the
number of true *cis*-heritable genes. In the latter case, estimates were unbiased at the largest
eQTL sample size, suggesting that the conservative bias is due to noise in gene expression
prediction models. For non-causal tissues, TCSC produced very slightly negative estimates of
$h^2_{ge(t')}$ (e.g. -7.2×10^{-4}, s.e. = 2.8×10^{-4} at eQTL sample size of 100 when setting $G_{t'}$ to
the number of true *cis*-heritable genes). We did not include a comparison to RTC Coloc[2] in our
analyses of bias, because RTC Coloc does not provide quantitative estimates of $h^2_{ge(t')}$.
203
We next evaluated the type I error of TCSC for non-causal tissues. TCSC was
conservative, with type I error less than 5% at a significance threshold of p = 0.05 (**Figure 2B**,
Supplementary Table 1). The conservative type I error was most pronounced at small eQTL
sample sizes. We determined that the conservative type I error is primarily due to conservative
jackknife standard errors (1.3x-1.6x; see **Supplementary Table 2**) rather than the very slight
negative bias, which is generally two to three orders of magnitude smaller than the jackknife
standard error for a given estimate (**Supplementary Table 3**); we hypothesize that the standard
errors are conservative relative to the empirical standard error due to variation of causal signal
across the genome[35]. For comparison, we also evaluated the type I error of RTC Coloc[2]. We
determined that RTC Coloc was not well-calibrated at lower eQTL sample sizes, with type I error
of 8.4% at eQTL sample size of 100 (**Supplementary Figure 1**, **Supplementary Table 4**).
215
We next evaluated the power of TCSC for causal tissues. We determined that TCSC was
well-powered to detect causal tissues at realistic eQTL sample sizes. When analyzing 300 eQTL

samples, TCSC attained 72% power at a nominal significance threshold of $p < 0.05$ (**Figure 2C**)
and 26% power at a stringent significance threshold of $p < 0.005$ (corresponding to 10% per-
trait FDR across tissues in these simulations, analogous to our analyses of real diseases/traits)
(**Supplementary Table 1**). As expected, power increased at larger eQTL sample sizes, due to
lower standard errors on point estimates of $h^2_{ge(t')}$ (**Figure 2A**). For comparison, we also
evaluated the power of RTC Coloc[2]. We determined that RTC Coloc attained substantially lower
power, which did not vary with eQTL sample size (**Supplementary Figure 1**, **Supplementary**
Table 4). We note that eQTL sample size is more likely to be a limiting factor when
disentangling co-regulated tissues using TCSC, whereas GWAS sample size is more likely to be a
limiting factor in other analyses such as the colocalization analyses used by RTC Coloc.
228
229   We similarly evaluated the robustness and power of TCSC when estimating tissue-
specific contributions to the genetic covariance between two diseases/traits. We employed the
same simulation framework described above and set the genetic correlation of the two
simulated traits to 0.5. We first evaluated the bias in TCSC estimates of the genetic covariance
explained by the *cis*-genetic component of gene expression in tissue t' ($\omega_{ge(t')}$), for both causal
and non-causal tissues. For causal tissues, TCSC produced conservative estimates of $\omega_{ge(t')}$,
particularly at small eQTL sample sizes and when setting $G_{t'}$ to the number of significantly *cis*-
heritable genes (rather than the number of true *cis*-heritable genes) (**Figure 3A**, **Supplementary**
Table 5), analogous to single-trait simulations. For non-causal tissues, TCSC again produced very
slightly negative estimates of $\omega_{ge(t')}$ (**Figure 3A**). We next evaluated the type I error of cross-
trait TCSC for non-causal tissues. TCSC was conservative at lower eQTL sample sizes, with type I
error less than 5% at $P = 0.05$ (**Figure 3B**). Finally, we evaluated the power of cross-trait TCSC
for causal tissues. We determined that cross-trait TCSC was modestly powered at realistic eQTL
sample sizes. When analyzing 300 gene expression samples, TCSC attained 29% power at $p <$
0.05 (**Figure 3C**) and 5.6% power at $p < 0.005$ (**Supplementary Table 5**).
244
245   We performed 6 secondary analyses. First, we set the eQTL sample size of the causal
tissue to 100 individuals and the eQTL sample sizes of the non-causal tissues to range between
100-1,000 individuals. We observed inflated type I error for non-causal tissues (particularly for
non-causal tissues with larger eQTL sample sizes), implying that large variations in eQTL sample
sizes may compromise type I error (**Supplementary Figure 2**). Second, we varied the true values
of $h^2_{ge(t')}$ (or $\omega_{ge(t')}$) for causal tissues. We determined that patterns of bias, type I error, and
power were similar across different parameter values (**Supplementary Figures 3-4**). Third, we
varied the number of causal tissues, considering 1, 2, or 3 causal tissues. We determined that
the power of TCSC decreased with each additional causal tissue (**Supplementary Figures 5-6**).
Fourth, we varied the number of non-causal tissues from 0 to 9. We determined that the
conservative bias in estimates of $h^2_{ge(t')}$ (or $\omega_{ge(t')}$) for causal tissues became less pronounced
as the number of non-causal tissues increased (**Supplementary Figures 7-8**). Fifth, we modified
TCSC to not correct for bias in tissue co-regulation scores arising from differences between *cis*-
genetic and *cis*-predicted expression. We determined that this increased the conservative bias
in estimates of $h^2_{ge(t')}$ for causal tissues and also resulted in anti-conservative jackknife
standard errors and type I error for non-causal tissues (**Supplementary Figure 9**), although the

impact on cross-trait simulations was limited (**Supplementary Figure 10**). Finally, we modified
TCSC to use "cheating" tissue co-regulation scores constructed using true eQTL effects. We
determined that this exacerbated the conservative bias in estimates of $h^2_{ge(t')}$ (or $\omega_{ge(t')}$) for
causal tissues, while also producing anti-conservative jackknife standard errors and type I error
for non-causal tissues (**Supplementary Figures 11-12**), perhaps because correlation patterns in
TWAS χ^2 statistics reflect in-sample correlations of *cis*-predicted expression. Further details of
these secondary analyses are provided in the **Supplementary Note**.
268
269   *Identifying tissue-specific contributions to 78 diseases and complex traits*
270
We applied TCSC to publicly available GWAS summary statistics for 78 diseases and
complex traits (average *N* = 302K; **Supplementary Table 6**) and gene expression data for 48
GTEx tissues[18] (**Table 1**) (see **Data Availability**). The 78 diseases/traits (which include 33
diseases/traits from UK Biobank[36]) were selected to have z-score > 6 for nonzero SNP-
heritability (as in previous studies[12,23,37]), with no pair of traits having squared genetic
correlation > 0.1[26] and substantial sample overlap (**Methods**). The 48 GTEx tissues were
aggregated into 39 *meta-tissues* (average N = 266, range: *N* = 101-320 individuals, 23 meta-
tissues with *N* = 320) in order to reduce variation in eQTL sample size across tissues (**Table 1**
and **Methods**); below, we refer to these as "tissues" for simplicity. We constructed gene
expression prediction models for an average of 3,993 significantly *cis*-heritable protein-coding
genes (as defined above) in each tissue. We primarily report the proportion of disease
heritability explained by the *cis*-genetic component of gene expression in tissue t'
($\pi_{t'} = h^2_{ge(t')}/h^2_g$), as well as its statistical significance (using per-trait FDR).
284
TCSC identified 27 causal tissue-trait pairs with significantly positive contributions to
disease/trait heritability at 10% FDR, spanning 9 distinct tissues and 23 distinct diseases/traits
(**Figure 4** and **Supplementary Table 7**). Many of the significant findings recapitulated known
biology, including associations of whole blood with blood cell traits such as white blood cell
count ($\pi_{t'}$ = 0.21, s.e. = 0.064, *P* = 5.7×10^{-4}); liver with lipid traits such as LDL ($\pi_{t'}$ = 0.20, s.e.
= 0.050, *P* = 2.9×10^{-5}); and thyroid with hypothyroidism ($\pi_{t'}$ = 0.31, s.e. = 0.11, *P* =
1.9×10^{-3}).
292
TCSC also identified several biologically plausible findings not previously reported in the
genetics literature. First, aorta artery was associated with glaucoma ($\pi_{t'}$ = 0.15, s.e. = 0.051, *P* =
1.3×10^{-3}). TCSC also identified aorta artery as a causal tissue for diastolic blood pressure
(DBP) ($\pi_{t'}$ = 0.078, s.e. = 0.024, *P* = 5.1×10^{-4}), which is consistent with DBP measuring the
pressure exerted on the aorta when the heart is relaxed[38]. High blood pressure is a known risk
factor for glaucoma[39-43], explaining the role of aorta artery in genetic susceptibility to glaucoma.
Second, esophagus muscularis was associated with the lung trait FEV1/FVC[44] ($\pi_{t'}$ = 0.15, s.e. =
0.053, *P* = 1.7×10^{-3}), consistent with previous work reporting that acidification of the
esophagus leads to airway constriction[45] and birth defects in the esophagus have consequences
for pulmonary function likely stemming from shared causal mechanisms in Sonic hedgehog[46-49].
FEV1/FVC is computed by dividing the amount of air an individual can expel from their lungs in
one second (FEV1) by total lung capacity (FVC); therefore, the esophagus muscularis could

impact either FEV1 and/or FVC. We sought to assess tissue-specific contributions to FEV1 and
FVC separately. For FVC, no causal tissues were identified at 10% FDR, and the tissues with the
most significant P-values were coronary artery and aorta artery (nominal P = 0.02 and
0.07, respectively; FDR > 10%), not esophagus muscularis. For FEV1, we analyzed summary
statistics from ref.[50] and determined that the tissue with the most significant P-value was
esophagus muscularis (nominal $P = 8.3 \times 10^{-3}$; FDR > 10%), suggesting that esophagus
muscularis may specifically impact FEV1 (we note that FEV1/FVC, FEV1, and FVC had similar z-
scores for nonzero SNP heritability, implying similar power in this analysis). Third, TCSC
identified heart left ventricle (in addition to whole blood) as a causal tissue for platelet count
($\pi_{t'}$ = 0.091, s.e. = 0.031, $P = 1.7 \times 10^{-3}$), consistent with the role of platelets in the formation
of blood clots in cardiovascular disease[51-54]. In cardiovascular disease, platelets are recruited to
damaged heart vessels after cholesterol plaques rupture, resulting in blood clots due to the
secretion of coagulating molecules[55]; antiplatelet drugs have been successful at reducing
adverse cardiovascular outcomes[56]. Moreover, the left ventricle serves as a muscle to pump
blood throughout the body[57], likely modulating platelet counts and other blood cell counts,
creating detectable changes in serum from which platelet counts are measured. Other
significant findings are discussed in the **Supplementary Note**, and numerical results for all
tissues and diseases/traits analyzed are reported in **Supplementary Table 8**.
323
TCSC also increased the specificity of known tissue-trait associations. For high density
lipoprotein (HDL), previous studies reported that deletion of a cholesterol transporter gene in
adipose tissue reduces HDL levels, consistent with the fact that adipose tissues are storage sites
of cholesterol and express genes involved in cholesterol transport and HDL lipidation[58,59]. While
there are three adipose tissues represented in the GTEx data that we analyzed (subcutaneous,
visceral, and breast tissue), TCSC specifically identified subcutaneous adipose ($\pi_{t'}$ = 0.16, s.e. =
0.054, $P = 1.5 \times 10^{-3}$; **Figure 4**), but not visceral adipose or breast tissue ($P > 0.05$;
Supplementary Table 8), as a causal tissue for HDL. Previous studies have established that
levels of adiponectin, a hormone released by adipose tissue to regulate insulin, are significantly
positively correlated with HDL[60-62] and more recently, a study has reported that adiponectin
levels are associated specifically with subcutaneous adipose tissue and not visceral adipose
tissue[63]; thus, the specific role of subcutaneous adipose tissue in HDL may be due to a causal
mechanism related to adiponectin. We note that TCSC did not identify liver as a causal tissue
for HDL (FDR > 10%), which may be due to limited power in liver due to smaller eQTL sample
size. For waist-hip ratio adjusted for BMI (WHRadjBMI), previous studies reported colocalization
of WHRadjBMI GWAS variants with *cis*-eQTLs in subcutaneous adipose, visceral adipose, liver,
and whole blood[64], consistent with WHRadjBMI measuring adiposity in the intraabdominal
space which is likely regulated by metabolically active tissues[65]. TCSC specifically identified
subcutaneous adipose ($\pi_{t'}$ = 0.10, s.e. = 0.037, $P = 2.4 \times 10^{-3}$; **Figure 4**), but not visceral
adipose, breast, liver, or whole blood ($P > 0.05$; **Supplementary Table 8**), as a causal tissue for
WHRadjBMI. The causal mechanism may involve adiponectin secreted from subcutaneous
adipose tissue, which is negatively correlated with WHRadjBMI[66]. We note that the P value
distributions across traits are similar for subcutaneous adipose (median P = 0.42) and visceral
adipose (median P = 0.56) and are comparable to the other 37 analyzed (median P = 0.20 –
0.84, **Supplementary Table 9**). For BMI, previous studies have broadly implicated the central

nervous system, but did not reveal more precise contributions[67,12,68,69,7,70]. TCSC specifically
identified brain cereb. ($\pi_{t'}$ = 0.042, s.e. = 0.015, P = 2.6×10^{-3}), but not brain cortex or brain
limbic (P > 0.05; **Supplementary Table 8**), as a causal tissue for BMI. Our finding is consistent
with brain cerebellum accounting for the majority of the brain's neurons and serving as a key
area of neurogenesis in the developing brain[71,72].
354
We performed a secondary analysis in which we removed tissues with eQTL sample size
less than 320 individuals, as these tissues may often be underpowered (**Figure 2C**). Results are
reported in **Supplementary Figure 13** and **Supplementary Table 10.** The number of causal
tissue-trait pairs with significantly positive contributions to disease/trait heritability (at 10%
FDR) increased from 27 to 33, likely due to a decrease in multiple hypothesis testing burden
from removing underpowered tissues. The 33 significant tissue-trait pairs reflect a gain of 14
newly significant tissue-trait pairs (and a loss of 8 formerly significant tissue-trait pairs, of which
5 were lost because the tissue was removed), but estimates of $\pi_{t'}$ for each significant tissue-
trait pair were not statistically different from our primary analysis (**Supplementary Table 11**).
Notably, among the newly significant tissue-trait pairs, esophagus mucosa was associated with
eczema ($\pi_{t'}$ = 0.085, s.e. = 0.032, P = 4.0×10^{-3}), consistent with the comorbidity of acid reflux
and eczema and hypotheses of a causal mechanism in which acid reflux stimulates the immune
cells in the mucosa, prompting an allergic response such as eczema[73]. Other newly significant
findings are discussed in the **Supplementary Note**, and numerical results for all tissues and
diseases/traits are reported in **Supplementary Table 10**.
370
We also performed a brain-specific analysis in which we applied TCSC to 41 brain traits
(average N = 226K, **Supplementary Table 12**) while restricting to 13 individual GTEx brain
tissues (**Supplementary Table 13**), analogous to previous work[7]. The 41 brain traits reflect a less
stringent squared genetic correlation threshold of 0.25. The 13 GTEx brain tissues were
analyzed without merging tissues into meta-tissues, and irrespective of eQTL sample size
(range: N = 101-189 individuals); we expected power to be limited due to the eQTL small
sample sizes and substantial co-regulation among individual brain tissues. TCSC identified 16
brain tissue-brain trait pairs at 10% FDR (**Supplementary Figure 14, Supplementary Table 14**)—
substantially more than the 2 brain tissue-brain trait pairs in **Figure 4**, although most results are
not directly comparable due to the more fine-grained individual brain tissues analyzed. For
ADHD, TCSC identified brain hippocampus as a causal tissue ($\pi_{t'}$ = 0.28, s.e. = 0.10, P =
2.5×10^{-3}), consistent with the correlation between hippocampal volume and ADHD
diagnosis in children[74]. A recent ADHD GWAS identified a locus implicating the *FOXP2* gene[75],
which has been reported to regulate dopamine secretion in mice[76]; hippocampal activation
results in the firing of dopamine neurons[77]. For BMI, TCSC identified brain amygdala (in addition
to brain cerebellum, a constituent tissue of the meta-tissue implicated in **Figure 4**) as a causal
tissue ($\pi_{t'}$ = 0.54, s.e. = 0.023, P = 8.3×10^{-3}), consistent with previous work linking the
amygdala to obesity and dietary self-control[78], although no previous study has implicated the
amygdala in genetic regulation of BMI. Other significant findings are discussed in the
Supplementary Note, and numerical results for all brain tissues and brain traits analyzed are
reported in **Supplementary Table 14**.
392

393   *Comparisons of TCSC to other methods*
394
We compared TCSC to two previous methods, RTC Coloc[2] and LDSC-SEG[7], that identify
disease-critical tissues using gene expression data. RTC Coloc identifies disease-critical tissues
based on tissue specificity of eQTL-GWAS colocalizations. LDSC-SEG identifies disease-critical
tissues based on heritability enrichment of specifically expressed genes. We note that RTC
Coloc and LDSC-SEG analyze each tissue marginally, whereas TCSC jointly models contributions
from each tissue to identify causal tissues (analogous to the distinction in GWAS between
marginal association and fine-mapping[22]). Thus, we hypothesized that RTC Coloc and LDSC-SEG
may output multiple highly statistically significant associated tissues for a given trait, whereas
TCSC may output a single causal tissue with weaker statistical evidence of causality. To assess
whether TCSC indeed attains higher specificity, we evaluated the results of each method both
for causal tissues identified by TCSC and for the most strongly co-regulated tagging tissue
(based on Spearman ρ for estimated eQTL effect sizes, averaged across genes, from ref.[18]). Our
primary analyses focused on 5 representative traits, defined as the 5 traits with highest z-score
for nonzero SNP-heritability among the 10 diseases/traits with at least one tissue-trait
association for each of the three methods (**Methods**).
410
Results for the 5 representative traits are reported in **Figure 5** and **Supplementary Table
15**; results for all 23 diseases/traits with causal tissue-trait associations identified by TCSC
(**Figure 4**) are reported in **Supplementary Figure 15** and **Supplementary Table 16**, and
complete results for all diseases/traits and tissues included in these comparisons are reported
in **Supplementary Table 17**. We reached three main conclusions. First, for a given disease/trait,
RTC Coloc typically implicates a broad set of tissues (not just strongly co-regulated tissues)
(**Figure 5A**); for example, for BMI, RTC Coloc implicated 6 of 10 tissues in **Figure 5** (and 9 of 17
tissues in **Supplementary Figure 15**). Second, for a given disease/trait, LDSC-SEG typically
implicates a small set of strongly co-regulated tissues (**Figure 5B**); for BMI, LDSC-SEG implicated
2 of 10 tissues in **Figure 5** (and 2 of 17 tissues in **Supplementary Figure 15**), consisting of brain
cereb. and its most strongly co-regulated tagging tissue. Third, for a given disease/trait, TCSC
typically implicates one causal tissue (**Figure 5C**); for BMI, TCSC implicated only brain cereb. as a
causal tissue, with even the most strongly co-regulated tagging tissue reported as non-
significant. For diastolic blood pressure (DBP), RTC Coloc implicated 9 of 10 tissues, LDSC-SEG
implicated 2 of 10 tissues (consisting of aorta artery and its most strongly co-regulated tagging
tissue), and TCSC implicated only aorta artery; as noted above, this result is consistent with DBP
measuring the pressure exerted on the aorta while the heart is relaxed[38]. As expected, the
higher specificity of TCSC in identifying unique causal tissues involves a trade-off of reduced
sensitivity, with less significant (lower $-\log_{10}$P-value and lower $-\log_{10}$FDR) results for causal
tissues in **Figure 5C** than in **Figure 5A** and **Figure 5B**) (**Supplementary Table 16**). We observed
similar patterns for HDL and WHRadjBMI, for which the increased specificity of TCSC is
discussed above (**Supplementary Figure 15, Supplementary Table 17, Supplementary Note**).
We also observed similar patterns when comparing TCSC to RTC Coloc and LDSC-SEG in the
brain-specific analysis of **Supplementary Figure 14** (**Supplementary Figure 16, Supplementary
Table 18, Supplementary Note**).
436

437   *Identifying tissue-specific contributions to the genetic covariance between two diseases/traits*
438

We applied cross-trait TCSC to 256 pairs of disease/traits (**Supplementary Table 19**) and
gene expression data for 48 GTEx tissues[18] (**Table 1**) (see **Data Availability**). Of 3,003 pairs of
the 78 disease/traits analyzed above, the 256 pairs of diseases/traits were selected based on
significantly nonzero genetic correlation ($p < 0.05 / 3{,}003$; see **Methods**). The 48 GTEx tissues
were aggregated into 39 meta-tissues, as before (**Table 1** and **Methods**). We primarily report
the proportion of genetic covariance explained by the *cis*-genetic component of gene
expression in tissue t' ($\zeta_{t'} = \omega_{ge(t')} / |\omega_g|$), as well as its statistical significance (using per-trait
FDR).

TCSC identified 30 causal tissue-trait covariance pairs with significant contributions to
trait covariance at 10% FDR, spanning 21 distinct tissues and 23 distinct trait pairs (**Figure 6A**
and **Supplementary Table 20**). For 27 of the 30 causal tissue-trait covariance pairs, the causal
tissue was non-significant for both constituent traits in the single-trait analysis of
Supplementary Table 8. Findings that recapitulated known biology included both examples
involving a tissue-trait pair that was significant in the single-trait analysis (**Figure 4**) and
examples in which both tissue-trait pairs were non-significant in the single-trait analysis
(**Supplementary Table 8**). Consistent with the significant contribution of liver to LDL heritability
in the single-trait analysis, TCSC identified a positive contribution of liver to the genetic
covariance of LDL and total cholesterol ($\zeta_{t'}$ = 0.090, s.e. = 0.029, $P = 1.0 \times 10^{-3}$), and
consistent with the positive contributions of whole blood to eosinophil count heritability and to
white blood cell count heritability in the single-trait analysis, TCSC identified a positive
contribution of whole blood to the genetic covariance of eosinophil count and white blood cell
count ($\zeta_{t'}$ = 0.32, s.e. = 0.12, $P = 2.4 \times 10^{-3}$). TCSC also identified a negative contribution of
muscle skeletal to the genetic covariance of WHRadjBMI and total protein ($\zeta_{t'}$ = -0.27, s.e. =
0.084, $P = 8.0 \times 10^{-4}$), consistent with reduced muscle mass in individuals with high waist-hip-
ratio[79]; this suggests that cross-trait TCSC can reveal tissue-trait biology distinct from what is
identified in the single-trait analyses.

TCSC also identified several biologically plausible findings not previously reported in the
genetics literature. First, LCLs had a significantly negative contribution to the genetic covariance
of eosinophil count and white blood cell count ($\zeta_{t'}$ = -0.081, s.e. = 0.028, $P = 1.8 \times 10^{-3}$, in
contrast to the positive contribution of whole blood). This is plausible as previous studies have
reported the suppression of proliferation of lymphocytes (the white blood cell hematopoietic
lineage from which LCLs are derived) by molecules secreted from eosinophils[80-82]. The
contrasting results for whole blood and LCLs suggest that genetic covariance may reflect
distinct tissue-specific contributions. Second, brain substantia nigra had a significantly positive
contribution to the genetic covariance of BMI and red blood cell count (RBC count) ($\zeta_{t'}$ = 0.28,
s.e. = 0.084, $P = 4.6 \times 10^{-4}$), while pancreas had a significantly negative contribution ($\zeta_{t'}$ = -
0.25, s.e. = 0.079, $P = 8.7 \times 10^{-4}$). In the brain, energy metabolism is regulated by oxidation
and previous work has shown that red blood cells play a large role in these metabolic processes
as oxygen sensors[83]; in addition, previous studies have reported differences in the level of
oxidative enzymes in red blood cells between individuals with high BMI and low BMI[84,85],

suggesting that genes regulating oxidative processes might have pleiotropic effects on RBC
count and BMI. In the pancreas, pancreatic inflammation (specifically acute pancreatitis) is
associated with reduced levels of red blood cells, or anemia[86], while pancreatic fat is associated
with metabolic disease and increased BMI[87]. Once again, the contrasting results for brain
substantia nigra and pancreas suggest that genetic covariance may reflect distinct tissue-
specific contributions. Third, brain substantia nigra had a significantly negative contribution to
the genetic covariance of age at first birth and height ($\zeta_{t'}$ = -0.11, s.e. = 0.032, $P = 4.5 \times 10^{-4}$).
Previous work in *C. elegans* reported that fecundity is positively regulated by dopamine[88,89],
which is produced in the substantia nigra[90]. Therefore, it is plausible that reproductive
outcomes related to fecundity, such as age at first birth, are also regulated by dopamine via the
substantia nigra. Dopamine also plays a role in regulating the levels of key growth hormones
such as IGF-1 and IGF-BP3[91] and has been previously shown to be associated with height[92].
Other significant findings are discussed in the **Supplementary Note**. Numerical results for all
tissues and disease/trait pairs analyzed are reported in **Supplementary Table 21**.
495
As noted above, for 27 of the 30 causal tissue-trait covariance pairs, the causal tissue
was non-significant for both constituent traits. We sought to formally assess whether
differences in tissue-specific contributions to genetic covariance vs. constituent trait heritability
were statistically significant. Specifically, for each causal tissue-trait covariance pair, we
estimated the differences between the tissue-specific contribution to covariance ($\zeta_{t'}$) and the
tissue-specific contributions to heritability for each constituent trait ($\pi_{t'}$). We identified eight
tissue-trait covariance pairs for which these differences were statistically significant at 10% FDR
for both constituent traits and $\pi_{t'}$ was non-significant for both constituent traits (**Figure 6A** and
Supplementary Table 22). For BMI and RBC count, the positive contribution of brain substantia
nigra and the negative contribution of pancreas to genetic covariance were each significantly
larger than the respective contributions of those tissues to BMI and RBC count heritability,
which were non-significant (**Figure 6B**). Other examples are discussed in the **Supplementary**
Note. Numerical results for all tissues and trait pairs are reported in **Supplementary Table 22**.
509
510 **Discussion**
511
We developed a new method, tissue co-regulation score regression (TCSC), that
disentangles causal tissues from tagging tissues and partitions disease heritability (or genetic
covariance of two diseases/traits) into tissue-specific components. We applied TCSC to 78
diseases and complex traits and 48 GTEx tissues, identifying 27 tissue-trait pairs (and 30 tissue-
trait covariance pairs) with significant tissue-specific contributions. TCSC identified biologically
plausible novel tissue-trait pairs, including associations of aorta artery with glaucoma,
esophagus muscularis with FEV1/FVC, and heart left ventricle with platelet count. TCSC also
identified biologically plausible novel tissue-trait covariance pairs, including a negative
contribution of LCLs to the covariance of eosinophil count and white blood cell count (in
contrast to the positive contribution of whole blood) and a positive contribution of brain
substantia nigra and a negative contribution of pancreas to the covariance of BMI and red
blood cell count; in particular, our findings suggest that genetic covariance may reflect distinct
tissue-specific contributions.

TCSC differs from previous methods in jointly modeling contributions from each tissue
to disentangle causal tissues from tagging tissues (analogous to the distinction in GWAS
between marginal association and fine-mapping[22]). We briefly discuss several other methods
that use eQTL or gene expression data to identify disease-associated tissues. RTC Coloc
identifies disease-associated tissues based on tissue specificity of eQTL-GWAS colocalizations[2];
this study made a valuable contribution in emphasizing the importance of tissue co-regulation,
but did not model tissue-specific effects, such that RTC Coloc may implicate many tissues
(**Figure 5A**). LDSC-SEG identifies disease-critical tissues based on heritability enrichment of
specifically expressed genes[7]; this distinguishes a focal tissue from the set of all tissues
analyzed, but does not distinguish closely co-regulated tissues (**Figure 5B**). MaxCPP models
contributions to heritability enrichment of fine-mapped eQTL variants across tissues or meta-
tissues[4]; although this approach proved powerful when analyzing eQTL effects that were meta-
analyzed across all tissues, it has limited power to identify disease-critical tissues: fine-mapped
eQTL annotations for blood (resp. brain) were significant conditional on annotations
constructed using all tissues only when meta-analyzing results across a large set of blood (resp.
brain) traits (Fig. 4 of ref.[4]). eQTLenrich compares eQTL enrichments of disease-associated
variants across tissues[3]; this approach produced compelling findings for eQTL that were
aggregated across tissues, but tissue-specific analyses often implicated many tissues (Fig. 1d of
ref.[3]). MESC estimates the proportion of heritability causally mediated by gene expression in
assayed tissues[93]; this study made a valuable contribution in its strict definition and estimation
of mediated effects (see below), but did not jointly model distinct tissues and had limited
power to distinguish disease-critical tissues (Fig. 3 of ref.[93]). CAFEH leverages multi-trait fine-
mapping methods to simultaneously evaluate all tissues for colocalization with disease[5];
however, this locus-based approach does not produce genome-wide estimates and it remains
the case that many (causal or tagging) tissues may colocalize with disease under this
framework. Likewise, methods for identifying tissues associated to disease/trait covariance do
not distinguish causal tissues from tagging tissues[94,95].

We note several limitations of our work. First, joint-fit effects of gene expression on
disease may not reflect biological causality; if a causal tissue or cell type is not assayed[96], TCSC
may identify a co-regulated tissue (e.g. a tissue whose cell type composition favors a causal cell
type) as causal. We anticipate that this limitation will become less severe as potentially causal
tissues, cell types and contexts are more comprehensively assayed. Second, TCSC does not
achieve a strict definition or estimation of mediated effects; this is conceptually appealing and
can, in principle, be achieved my modeling non-mediated effects, but may result in limited
power to distinguish disease-critical tissues[93]. Third, TCSC has low power at small eQTL sample
sizes; in addition, TCSC estimates are impacted by the number of significantly *cis*-heritable
genes in a focal tissue, which can lead to conservative bias at small eQTL sample sizes. We
anticipate that these limitations will become less severe as eQTL sample sizes increase. Fourth,
TCSC is susceptible to large variations in eQTL sample size, which may compromise type I error;
therefore, there is a tradeoff between maximizing the number of tissues analyzed and limiting
the variation in eQTL sample size. Fifth, TCSC assumes that causal gene expression-disease
effects are independent across tissues; this assumption may become invalid for tissues and cell

types assayed at high resolution. However, we verified via simulations that in the case of two
causal tissues with identical gene expression-trait effects, TCSC correctly identified both tissues
as causal and split estimates of explained heritability across the tissues (**Supplementary Figure**
17). Sixth, TCSC does not formally model measurement error in tissue co-regulation scores, but
instead applies a heuristic bias correction. We determined that the bias correction generally
performs well in simulations. Seventh, TCSC does not produce locus-specific estimates or
identify causal tissues at specific loci. However, genome-wide results from TCSC may be used as
a prior for locus-based methods (analogous to GWAS fine-mapping with functional priors[97]).
Finally, we did not apply TCSC to single-cell RNA-seq (scRNA-seq) data, which represents a
promising new direction as scRNA-seq sample sizes increase[98-100,32]; we caution that scRNA-seq
data may require new eQTL modeling approaches[98]. Despite these limitations, TCSC is a
powerful and generalizable approach for modeling tissue co-regulation to estimate tissue-
specific contributions to disease.
582
583 **Code Availability**
584
585 TCSC software: https://github.com/TiffanyAmariuta/TCSC/tree/main/analysis.
586 Mancuso Lab TWAS Simulator: https://github.com/mancusolab/twas_sim.
587 FUSION software: http://gusevlab.org/projects/fusion/.
588
589 **Data Availability**
590
We have made 78 GWAS summary statistics and 41 brain-specific summary statistics publicly
available at https://github.com/TiffanyAmariuta/TCSC/tree/main/sumstats, TWAS association
statistics publicly available at
https://github.com/TiffanyAmariuta/TCSC/tree/main/twas_statistics, tissue co-regulation
scores publicly available at
https://github.com/TiffanyAmariuta/TCSC/tree/main/coregulation_scores, and TCSC output
will be publicly available at https://github.com/TiffanyAmariuta/TCSC/tree/main/results.
598

599 **Acknowledgements**
600
601 We thank Huwenbo Shi, Martin Zhang, and Benjamin Strober for helpful discussions. This work
602 was funded by NIH grants U01 HG009379, R01 MH101244, R37 MH107649, R01 HG006399, R01
603 MH115676 and U01 HG012009.

604
605

606   **Online Methods**
607
TCSC regression
TCSC leverages the fact that the TWAS χ^2 statistic for a gene-tissue pair includes the
direct effects of the gene on the disease as well as the tagging effects of co-regulated tissues
and genes with shared eQTLs or eQTLs in LD. Thus, genes that are co-regulated across many
tissues will tend to have higher χ^2 statistics than genes regulated in a single tissue. TCSC
determines that a tissue causally contributes to disease if genes with high co-regulation to the
tissue have higher TWAS χ^2 statistics than genes with low co-regulation to the tissue.
We model the genetic component of gene expression as a linear combination of SNP-
level effects:

617
$$W_{jgt} = \sum_m X_{jm}\beta_{gtm},$$

where W_{jgt} is the *cis*-genetic component of gene expression in individual j for gene g and tissue
t, X_{jm} is the standardized genotype of individual j for SNP m, and β_{gtm} is the effect of the m^{th}
SNP on gene expression of gene g in tissue t.
TCSC assumes that true gene-disease effects are independent and additive across
tissues[1]. However, *cis* eQTLs are highly correlated across tissues, leading to tagging from co-
regulated tissues[2]. We model phenotype as a linear combination of genetic components of
gene expression across genes in different tissues:

625
$$Y_j = \sum_t \sum_g W_{jgt}\, \alpha_{gt} + \epsilon_j,$$

626   where Y_j is the (binary or continuous-valued) phenotype of individual j, α_{gt} is the gene-trait
627   effect size and ϵ_j is the error term.
628   We define the disease heritability explained by *cis*-predicted expression across all
629   tissues as follows:

630
$$h^2_{ge} = \sum_t \sum_g \alpha^2_{gt},$$

631   analogous to the relationship between SNP effect sizes and SNP-heritability[25]. It follows that the
632   disease heritability explained by a particular tissue t' is

633
$$h^2_{ge(t')} = \sum_g \alpha^2_{gt'}.$$

634   Let $\alpha_{gt'}$ be a random variable drawn from a normal distribution with mean zero and tissue-
635   specific variance $var(\alpha_{gt'}) = \tau_{t'}$. Then

636
$$h^2_{ge(t')} = \sum_g var(\alpha_{gt'}) = \sum_g \tau_{t'} G_{t'},$$

637   where $G_{t'}$ is the number of *cis* heritable genes in t'. With this variance term, we can define a
638   polygenic model that relates TWAS χ^2 statistics to co-regulation scores, which explicitly model
639   the covariance structure of the χ^2 statistics. This strategy is analogous to modeling the
640   dependence of GWAS χ^2 statistics on LD scores[25]. We model the expected value of the TWAS χ^2
641   across genes and tissues as follows:

$$642 \qquad\qquad\qquad\qquad E[\chi^2_{g,t}] = N \sum_{t'} \tau_{t'} l(g,t;t') + 1, \qquad\qquad\qquad (1)$$

643   where g indexes genes, t indexes tissues, $\chi^2_{g,t}$ is a vector of TWAS χ^2 statistics across all
644   significantly *cis*-heritable genes and tissues, N is GWAS sample size, $\tau_{t'}$ is the per-gene
645   heritability explained by predicted gene expression in tissue t', and $l(g,t;t')$ are tissue and
646   gene co-regulation scores (see below). From the derivation, the genome-wide tissue-specific
647   contribution to disease heritability is estimated as

$$648 \qquad\qquad\qquad\qquad h^2_{ge(t')} = G_{t'} \tau_{t'}.$$

649   For the analysis of tissue-specific contributions to the covariance between two diseases,
650   we can extend TCSC by using products of TWAS z-scores. Following the polygenic model
651   described above, the expected product of TWAS z-scores in disease 1 and disease 2 for gene g
652   and tagging tissue t is
653

$$654 \qquad E[z^1_{gt} \times z^2_{gt}] = \sqrt{N_1 N_2} \sum_{t'} l(g,t;t') \omega_{ge(t')}/G_{t'} + \rho N_s/\sqrt{N_1 N_2} \qquad (2)$$

655
where N_1 is GWAS sample size for disease 1, N_2 is GWAS sample size for disease 2, t' indexes
causal tissues, $l(g,t;t')$ are tissue co-regulation scores (see below), $\omega_{ge(t')}$ is the genetic
covariance explained by the *cis*-genetic component of gene expression in tissue t', $G_{t'}$ is the
number of significantly *cis*-heritable genes in tissue t' (see below), ρ is the phenotypic
correlation between disease 1 and disease 2, and N_s is the number of overlapping GWAS
samples between disease 1 and disease 2. The last term represents the intercept[26], and while
we use a free intercept in the multivariate regression on co-regulation scores, the estimation of
this term only plays a role in the estimation of regression weights (see below).
664
For estimates of $h^2_{ge(t')}$ and $\omega_{ge(t')}$, we use a free intercept; the estimation of
$\rho N_s \sqrt{N_1 N_2}$ serves only to inform the heteroscedasticity weights (see below) and is not used in
the multivariate TCSC regression to estimate $\omega_{ge(t')}$. To estimate standard errors, we use a
genomic block jackknife over 200 genomic blocks with an equal number of genes in each. The
standard deviation is computed as the square root of the weighted variance across the
jackknife estimates (where the weight of each block is equal to the sum of the regression
weights for the genes in that block) multiplied by 200 blocks. We expect that the jackknife
standard error will be conservative relative to the empirical standard error across estimates due
to variation in causal signal across loci[35].
674
675   *Estimating tissue co-regulation scores and correcting for bias*
676   We define the co-regulation score of gene g with tissues t and t' as
677

$$678 \qquad\qquad\qquad l(g,t;t') = \sum_{g'} r^2(W_{g,t}, W_{g',t'}), \qquad\qquad\qquad (3)$$

where W denotes the *cis*-genetic component of gene expression for a gene-tissue pair across
individuals and genes g' are within +/- 1 Mb of the focal gene g. TCSC corrects for bias in tissue
co-regulation scores arising from differences between *cis*-genetic vs. *cis*-predicted expression
(analogous to GCSC[21]). We apply bias correction to co-regulation scores in the special case
when $t = t'$. While co-regulation scores aim to estimate the squared correlation of *cis*-
predicted gene expression of gene g and tissue t (corresponding to the TWAS $\chi^2_{g,t}$ statistic)
with the true *cis*-genetic component of gene expression of *cis*-genes in tissue t', when $g = g'$
and $t = t'$, the estimated value of $r^2(W_{g,t}, W_{g',t'})$ will always equals one. However, in
practice, the prediction R^2 of a gene expression model will always be less than one, except in
the case of a perfect model, causing the value of $r^2(W_{g,t}, W_{g',t'})$ to be systematically inflated.
Therefore, when $g = g'$ and $t = t'$, we set

$$r^2(W_{g,t}, W_{g',t'}) = R^2/h^2_{GCTA}, \qquad\qquad (4)$$

694 where R^2 is the cross-validation prediction statistic of the gene expression model for gene g in
695 tissue t and h^2_{GCTA} is the GCTA-estimated *cis*-heritability of gene expression for gene g in tissue
696 t. The quotient R^2/h^2_{GCTA} is the accuracy of the gene expression prediction model, which
697 reflects the upper bound on how much the *cis*-predicted expression can be correlated with the
698 true *cis*-genetic component of gene expression.

TCSC regression weights

701 TCSC uses three sets of regression weights to increase power (analogous to GCSC[21]). The
702 first regression weight is inversely proportional to $L(g,t)$, the total co-regulation score of each
703 gene-tissue pair summed across tissues t':

$$L(g,t) = \sum_{t'} \sum_{g'} r^2(W_{g,t}, W_{g',t'})$$

707 (without applying bias correction; see above), which allows TCSC to properly account for
708 redundant contributions of co-regulated genes to TWAS χ^2 statistics.

710 The second regression weight is inversely proportional to $T(g,t)$, the number of tissues
711 in which a gene is significantly *cis*-heritable:

$$T(g,t) = \sum_{t' \ni g,t' significantly\ cis-heritable} 1,$$

thereby up-weighting signal from genes that are regulated in a limited number of tissues and
preventing TCSC from attributing more weight to genes that are co-regulated across many
tissues.

The third regression weight is inversely proportional to $H_{h^2}(g,t)$, the heteroscedasticity
of χ^2 statistics, and is computed differently for estimates of $h^2_{ge(t')}$ than for estimates of $\omega_{ge(t')}$
(analogous to GCSC[21] and cross-trait LDSC[26], respectively).

For estimates of $h^2_{ge(t')}$, we estimate $H_{h^2}(g,t)$ in two steps. First, we make a crude

estimate of heritability explained by predicted expression (h^2_{ge}) as follows:

$$\mu_\chi = N\mu_L h^2_{ge} + 1,$$

726   where μ_χ is the mean χ^2 statistic:

$$\mu_\chi = \frac{\sum_{t'}\sum_{g'} \chi^2_{g',t'}}{\sum_{t'} G_{t'}},$$

728   where N is the GWAS sample size, g' iterates over significantly *cis*-heritable genes and t'

729   iterates over tissues, and μ_L is the mean value of total co-regulation across tissues t',

$$\mu_L = \frac{\sum_{t'}\sum_{g'} L(g',t')}{\sum_{t'} G_{t'}}.$$

733   Then, we compute the heteroscedasticity for each significantly *cis*-heritable gene-tissue pair as

$$H_{h^2}(g,t) = \left(NL(g,t)h^2_{ge} + 1\right)^2.$$

736   Finally, we combine the three regression weights as follows:

$$Weight_{h^2}(g,t) = \frac{1}{L(g,t)T(g,t)H_{h^2}(g,t)}.$$

739   For estimates of $\omega_{ge(t')}$, we estimate $H_\omega(g,t)$ in two steps. First, we regress the

740   products of TWAS z-scores on total tissue co-regulation scores, $L(g,t)$, using regression

741   weights, $Weight_\omega(g,t)$, computed as follows:

$$Weight_\omega(g,t) = \frac{1}{L(g,t)T(g,t)H_\omega(g,t)}$$

743   where $H_\omega(g,t)$ is *first* estimated as follows:

$$H_\omega(g,t) = \left(N_1 L(g,t)h^2_{ge}(trait\ 1) + 1\right)\left(N_2 L(g,t)h^2_{ge}(trait\ 2) + 1\right) +$$

$$\left(\frac{\sqrt{N_1 N_2}\,\omega_{ge}L(g,t)}{\sum_{t'}\frac{G_{t'}}{T'}}\right)^2,$$

748   where $h^2_{ge}(trait\ 1)$ is the crude heritability estimate for trait 1 and $h^2_{ge}(trait\ 2)$ is the crude

749   heritability estimate for trait 2, ω_{ge} is estimated as $\frac{\sum_{t'}\sum_{g'} z^{(1)}_{g't'} z^{(2)}_{g't'}}{\sqrt{N_1 N_2}}$, N_1 is the sample size of the

750   first GWAS, N_2 is the sample size of the second GWAS, and T' is the total number of tissues in

751   the regression.

753   Second, we use the regression intercept to estimate the product ρN_s:

$$\rho N_s = intercept * \sqrt{N_1 N_2},$$

755   where ρ represents the phenotypic correlation between trait 1 and 2 and N_s represents the

756   number of shared samples between GWAS 1 and 2. We also use the coefficient of the

757   regression to update our estimate of ω_{ge}, such that we may update the heteroscedasticity
758   weight as follows:

$$H_\omega(g,t) = \left(N_1 L(g,t) h_{ge}^2(trait\ 1) + 1\right)\left(N_2 L(g,t) h_{ge}^2(trait\ 2) + 1\right)$$

$$+ \left(\frac{\sqrt{N_1 N_2}\,\omega_{ge} L(g,t)}{\sum_{t'} \frac{G_{t'}}{T'}} + \rho N_s \sqrt{N_1 N_2}\right)^2.$$

761
762   Finally, we combine the three regression weights as follows:

$$Weight_\omega(g,t) = \frac{1}{L(g,t)T(g,t)H_\omega(g,t)}.$$

764
Simulating TCSC
We employed a widely used TWAS simulation framework (Mancuso Lab TWAS
Simulator, see **Code Availability**) to assess the power, bias, and calibration of TCSC in the
presence of co-regulation across genes and tissues. We simulated a genome in which there are
1,245 genes, of which 125 (10%) are causal[34]. Each gene belongs to one of 249 independent
genomic blocks, representing approximately 1Mb intervals of chromosome 1, which is 249 Mb
in length. Each block contains 50 variants and 5 genes.
Each primary simulation consists of 10 tissues, of which at least one is causal, defined as
having nonzero gene-trait effect sizes. We create a covariance structure among tissues
mimicking empirical GTEx data. We use a previously published method to estimate the causal
cross-tissue correlation of eQTL effect sizes which is 0.75[33]. We observe that not all GTEx tissues
are equally correlated to one another. We estimate three different cross-tissue eQTL
correlation quantities: (1) average correlation across all pairs of tissues = 0.75, (2) average
correlation across similar tissues = 0.80, e.g. brain (13 in GTEx) or adipose (2 in GTEx) tissues,
and (3) average correlation across dissimilar tissues, e.g. pairs of brain and adipose tissues =
0.74. To represent these biological modules, we let simulated tissues 1-3 have higher
correlation of true eQTL effects to one another than to other tissues; likewise for tissues 4-6
and 7-10. We set covariance parameters, described below, such that the similar tissues had an
average eQTL correlation of 0.789 across genes, dissimilar tissues have an average eQTL
correlation of 0.737, and the average eQTL correlation across any pair of tissues is 0.751. We do
not simulate linkage disequilibrium, which is another source of co-regulation in real data.
Rather, we simulate co-regulation strictly using shared eQTLs across genes and tissues. We do
not expect this to impact the interpretation of our empirical results. We simulate each gene has
5 true eQTLs, based on the upper bound of empirical data from GTEx[18] and others[32], as well as
the value used in other TWAS simulation methods[31]. Between co-regulated tissues, each pair of
genes shares 3 eQTLs. The minimum allowed *cis*-heritability of a gene is 0.01 in our simulations.
Cis-heritability is approximated as the sum of squared true *cis*-eQTL effect sizes, as done
previously[21]; this approximation of *cis*-heritability is more accurate than if we were to model
linkage disequilibrium between *cis*-eQTLs. Effect sizes for the 3 shared eQTLs across tissues are
sampled from a multivariate normal distribution with mean 0 and a variance-covariance matrix.
We define the variance and covariance terms of this matrix such that (1) the proportion of
genes detected as significantly *cis*-heritable by GCTA at a given sample size and (2) the average

cis heritability of detected genes at a given sample size match empirical observations from GTEx
data at sample sizes N = 100, 200, 300 and 500. As a result, the diagonal of the variance-
covariance matrix, e.g. the variance term, is set to 0.015, and the off-diagonal elements are set
to the product of the variance term and the desired correlation for each tissue pair, described
above.
For each of 1,000 independent simulations per analysis, we simulate a GWAS (N =
100,000) by creating a complex trait which is the summation of the genetic components of
causal gene expression (in the causal tissue). We use simulated genotypes drawn from a
standard normal distribution. Gene-trait effect sizes are drawn from a normal distribution with
mean 0 and variance 1. In cross-trait TCSC analysis, effect sizes across genes between the two
traits are correlated with default R_g = 0.5. To simulate a GWAS trait, we first compute the
genetic component of each gene, which is the product of GWAS cohort genotypes and eQTL
effects, such that we have 125 gene-specific traits. We then add noise to each gene-specific
trait such that the total variance of the phenotype explained by the five eQTLs from the causal
tissue is equal to a specified value; the value of $h^2_{ge(t)}$ in primary simulations is 5%. Then, we
multiply each gene-specific trait by the causal gene-trait effect size, consistent with the additive
generative model of gene-level effects on trait (see above). Finally, we take the sum across all
gene-specific traits to make one complex trait, where the total variance of the trait explained
by gene effects from the causal tissue is $h^2_{ge(t)}$, e.g. 5%.
We simulate an eQTL cohort of various gene expression sample sizes (N = 100, 200, 300,
500, 1000) using simulated genotypes drawn from a standard normal distribution. We simulate
total gene expression in the eQTL cohort by adding a desired amount of noise to the genetic
component of gene expression, e.g. the product of individual genotypes and true eQTL effect
sizes, with variance equal to one minus the gene expression heritability, which is the sum of
squared eQTL effects. Next, we fit gene expression prediction models by regressing the total
gene expression on eQTL cohort genotypes of *cis* variants using lasso regularization, a standard
approach used in TWAS. We define significantly *cis*-heritable genes as genes with GCTA
heritability *P* value < 0.01[20] and heritability estimate > 0, and adjusted-R^2 > 0 in cross-validation
prediction.
Then we estimate co-regulation scores at each different eQTL sample size by predicting
gene expression into a cohort of 500 individuals, to approximate the size of the European
sample of 1000 Genomes (*N* = 489). Using significantly *cis*-heritable genes from each tissue at a
given sample size, we estimate gene and tissue co-regulation scores $l(g, t; t')$ as described
above, including bias correction. In simulations, *cis* genes are defined as genes within the same
1 Mb block.
Then we apply TWAS to individual-level simulated GWAS data and gene expression
prediction models. We predict gene expression into each of the 100,000 GWAS cohort
individuals across all significantly *cis*-heritable genes for each tissue. We regress each complex
trait on predicted gene expression to obtain TWAS z-scores. Finally, we run TCSC by regressing
TWAS χ^2 statistics, or products of TWAS z-scores, on bias-corrected gene and tissue co-
regulation scores.
838
Simulating the RTC Coloc method

We simulated the RTC Coloc method[2] by leveraging our existing TCSC simulation
framework such that both methods could be compared via application to same simulated data.
We precisely followed the steps of the RTC Coloc method. We first simulated individual level
genotypes of a GWAS (N = 100,000) using a standard normal distribution, as described. We then
regress the simulated complex traits on each of the GWAS genotypes in turn using a Bonferroni
significance threshold to identify genome-wide significant variants. We then select one null SNP
per GWAS variant for comparison. Next, we simulate an eQTL cohort consisting of total gene
expression and genotypes, using the same underlying true eQTL effect sizes as for TCSC
simulations. Then we perform colocalization analysis of GWAS variants with eQTLs, across 10
tissues at 4 different eQTL sample sizes, to obtain the regulatory trait concordance (RTC) score.
This is repeated for the set of null variants. Next, we perform colocalization analysis of eQTL
variants between pairs of tissues to obtain tissue-sharing RTC scores, and similarly repeat this
for null variants. GWAS-eQTL RTC scores are divided by tissue-sharing RTC scores summed
across variants. Tissue-specific enrichment is computed as the ratio of this quotient to the null
quotient. The enrichment *P* value is obtained using a Wilcox test comparing the values of the
quotient to the values of the null quotient.
856

Gene expression prediction models and tissue co-regulation scores in GTEx data
We downloaded GTEx v8 gene expression data for 49 tissues. We excluded tissues with
fewer than 100 samples, e.g. kidney cortex (n = 69). We retained only European samples for
each tissue, as labeled by GTEx via PCA of genotypes. We constructed gene expression models
for two scenarios: (1) subsampling to 320 individuals including meta-analyzed tissues (**Table 1**)
or (2) using all European samples per tissue. We recommend meta-analyzing gene expression
prediction models across tissues in the case of tissues with low eQTL sample size (e.g. < 320
samples) and high pairwise genetic correlation (e.g. > 0.93). We determined in simulations that
TCSC is sensitive to eQTL sample size differences, such that a tagging tissue with larger sample
size than a causal tissue can produce false positive results; the subsampling approach was
designed to mitigate this issue. For the subsampling procedure, we first set aside tissues with
more than 320 samples; we chose 320 based on the average GTEx tissue sample size (*N* = 271)
and robustness of TCSC in simulations at *N* = 300. Then, we grouped tissues with genetic
correlation, e.g. marginal effect size correlation as reported by GTEx, with R_g > 0.93, an arbitrary
threshold that produced biologically plausible groups of related tissues, separating groups of
brain tissues based on cranial compartment. We meta-analyzed gene expression prediction
models for these grouped tissues in order to achieve a total sample size of 320 individuals
where each tissue contributed an approximately equal number of samples, using an inverse-
variance weighted meta-analysis across genes that were significantly *cis*-heritable in two or
more constituent tissues. The prediction weights of genes that were significantly *cis*-heritable in
a single constituent tissue were left unmodified.
To construct gene expression prediction models, we applied FUSION[20] (**Code
Availability**) to individual-level GTEx data by regressing measured gene expression on
genotypes of common variants (MAF > 0.05) and covariates provided by GTEx[18]. FUSION uses
several different regression models: single eQTL, elastic net, lasso, and BLUP and the following
covariates: sex, 5 genotyping principal components, PEER factors[101], and assay type. We

defined significantly *cis*-heritable genes as protein-coding genes with GCTA heritability *p* <
0.01[20], heritability estimate > 0, and adjusted-R^2 > 0 in cross-validation prediction.
We used gene expression prediction models of significantly *cis*-heritable genes to
predict expression into 489 European individuals from 1000 Genomes[102]. We then estimated
tissue co-regulation scores using Equation (3) and Equation (4), where *cis*-predicted gene
expression is used to estimate the *cis*-genetic component of gene expression.
889
GWAS summary statistics and TWAS association statistics
We collected GWAS summary statistics from 78 independent heritable complex diseases
and traits (average N = 302K) with heritability z-score > 6. We estimated the heritability of all
summary statistics and genetic correlation of all pairs of summary statistics. We excluded traits
with heritability z-score < 6, using S-LDSC with the baseline-LD v2.2 model[12,23,24] and as done
previously[23]. We excluded one of each pair of traits that are *both* genetically correlated and
have significantly overlapping samples. Specifically, for any pair of non-UK Biobank traits with
an estimated sample overlap greater than the following threshold -- squared cross-trait LDSC
intercept / (trait 1 S-LDSC intercept * trait 2 S-LDSC intercept) > 0.1[26] -- the trait with the larger
SNP heritability z-score was retained. For any pair of UK Biobank traits with a squared genetic
correlation > 0.1, the trait with the larger SNP heritability z-score was retained[37]. In total, this
procedure resulted in 78 sets of GWAS summary statistics. For the brain-specific analysis, we
first selected brain-related diseases and complex traits, e.g. psychiatric disorders and behavioral
phenotypes, excluding multi case-control studies and case vs case studies. Then, we applied our
standard filters as described above, but relaxing the threshold of squared genetic correlation to
0.25.
We used FUSION[20] (**Code Availability**) to compute TWAS association statistics for each
pair of signed GWAS summary statistics and each significantly *cis*-heritable gene-tissue pair,
across the two scenarios described above. We further removed genes within the MHC
(chromosome 6, 29 Mb - 33 Mb) and TWAS χ^2 > 80 or χ^2 > 0.001N, where N is the GWAS sample
size, as previously used for quality control in the heritability analysis of GWAS summary
statistics[12].
912
RTC Coloc and LDSC-SEG analysis of GWAS summary statistics and GTEx tissues
We downloaded supplementary tables for the RTC coloc method[2] and for LDSC-SEG[7].
For traits in our set of 78 GWAS summary statistics that were not analyzed by the LDSC-SEG
study and for traits that are inherently brain-related (as these traits require a different
procedure for generating tissue-specific gene sets), we ran LDSC-SEG ourselves. To this end, we
downloaded LD scores for GTEx tissues and specifically expressed gene set SNP-level
annotations (https://alkesgroup.broadinstitute.org/LDSCORE/LDSC_SEG_ldscores/) and ran
LDSC-SEG as previously described[7]. For brain-related traits, we additionally ran a brain-specific
analysis using LDSC-SEG, also as previously described[7]. Briefly, specifically expressed genes were
determined via a t-test of the sentinel brain tissue against all other brain tissues, rather than
against all other non-brain GTEx tissues, as done in the primary analysis of the LDSC-SEG study.
For traits in our set that were not analyzed by the RTC Coloc study, of which there were few, we
did not apply their method, as it was too computationally intensive to apply to real trait data.

Tables
928

Meta-tissue	Constituent Tissue(s)	Sample Size
Adipose Subcutaneous	Adipose Subcutaneous	320
Adipose Visceral Omentum	Adipose Visceral Omentum	320
Adrenal Gland	Adrenal Gland	200
Aorta Artery	Aorta Artery	320
† Brain Basal Ganglia	Putamen, Caudate, Nucleus Accumbens	320
† Brain Cereb.	Cerebellum, Cerebellar Hemisphere	320
† Brain Cortex	Frontal, Anterior, Cingulate	320
† Brain Limbic	Amygdala, Hippocampus, Hypothalamus	320
Brain Spinal Cord	Brain Spinal Cord	115
Brain Substantia Nigra	Brain Substantia Nigra	101
Breast Mammary Gland	Breast Mammary Gland	320
Coronary Artery	Coronary Artery	180
Cultured Fibroblasts	Cultured Fibroblasts	320
Esophagus Mucosa	Esophagus Mucosa	320
Esophagus Muscularis	Esophagus Muscularis	320
Heart Atrial Appendage	Heart Atrial Appendage	320
Heart Left Ventricle	Heart Left Ventricle	320
LCLs	LCLs	116
Liver	Liver	183
Lung	Lung	320
Minor Salivary Gland	Minor Salivary Gland	118
Muscle Skeletal	Muscle Skeletal	320
Ovary	Ovary	140
Pancreas	Pancreas	252
Pituitary	Pituitary	220
Prostate	Prostate	186
Skin (sun exposed)	Skin (sun exposed)	320
Skin (sun unexposed)	Skin (sun unexposed)	320
† Sigmoid Intestine	Sigmoid Colon, Gastroesophageal Junction	320
Spleen	Spleen	185
Stomach	Stomach	269
Tibial Artery	Tibial Artery	320
Tibial Nerve	Tibial Nerve	320
Testis	Testis	277
Thyroid	Thyroid	320
† Transverse Intestine	Transverse Colon, Small Intestine	320
Uterus	Uterus	108
Vagina	Vagina	122
Whole Blood	Whole Blood	320

930 **Table 1. GTEx meta-tissues and constituent tissues analyzed.** For each meta-tissue we list the
931 constituent tissue(s) and total sample size. Daggers denote meta-tissues with more than one
932 constituent tissue; for these meta-tissues, each constituent tissue has equal sample size up to
933 rounding error (an exception is the transverse intestine meta-tissue, which includes 176
934 transverse colon samples and all 144 small intestine samples).
935

937 **Figures**

939
Figure 1. Overview of TCSC regression. (A) Input data to TCSC includes (1) GWAS summary
statistics for a disease and (2) gene expression prediction models for each tissue, which are
used to produce (3) TWAS summary statistics for the disease for each tissue. (B) TCSC computes
tissue co-regulation scores $L(g, t; t')$ for each gene-tissue pair (g, t) with potentially causal
tissues t'. (C) TCSC regresses TWAS chi-squares on tissue co-regulation scores to estimate
tissue-specific contributions to disease.
946

947
Figure 2. Robustness and power of TCSC regression in simulations. (A) Bias in estimates of
disease heritability explained by the *cis*-genetic component of gene expression in tissue t'
($h^2_{ge(t')}$) for causal (light and dark purple) and non-causal (gray) tissues, across 1,000
simulations per eQTL sample size. The dashed line indicates the true value of $h^2_{ge(t')}$ for causal
tissues. Light purple indicates that $G_{t'}$ was set to the number of true *cis*-heritable genes, dark
purple indicates that $G_{t'}$ was set to the number of significantly *cis*-heritable genes. (B)
Percentage of estimates of $h^2_{ge(t')}$ for non-causal tissues that were significantly positive at $p <$
0.05, across 1,000 simulations per eQTL sample size. (C) Percentage of estimates of $h^2_{ge(t')}$ for
causal tissues that were significantly positive at $p < 0.05$, across 1,000 simulations per eQTL
sample size. We note that (B) type I error and (C) power are not impacted by the value of $G_{t'}$.
Error bars denote 95% confidence intervals. Numerical results are reported in **Supplementary**
Table 1.
960

961
Figure 3. Robustness and power of cross-trait TCSC in simulations. (A) Bias in estimates of
genetic covariance explained by the *cis*-genetic component of gene expression in tissue t'
($\omega_{ge(t')}$) for causal (light and dark purple) and non-causal (gray) tissues, across 1,000
simulations per eQTL sample size. The dashed line indicates the true value of $\omega_{ge(t')}$ for causal
tissues. Light purple indicates that $G_{t'}$ was set to the number of true *cis*-heritable genes, dark
purple indicates that $G_{t'}$ was set to the number of significantly *cis*-heritable genes. (B)
Percentage of estimates of $\omega_{ge(t')}$ for non-causal tissues that were significantly positive at $p <$
0.05, across 1,000 simulations per eQTL sample size. (C) Percentage of estimates of $\omega_{ge(t')}$ for
causal tissues that were significantly positive at $p < 0.05$, across 1,000 simulations per eQTL
sample size. We note that (B) type I error and (C) power are not impacted by the value of $G_{t'}$.
Error bars denote 95% confidence intervals. Numerical results are reported in **Supplementary**
Table 5.
974

Figure 4. TCSC estimates tissue-specific contributions to disease and complex trait heritability. We report estimates of the proportion of disease heritability explained by the cis-genetic component of gene expression in tissue t' ($\pi_{t'}$) for 27 causal tissue-trait pairs with significantly positive contributions to disease/trait heritability at 10% FDR, spanning 9 distinct tissues and 23 distinct diseases/traits. Tissues are ordered alphabetically. Daggers denote meta-tissues with more than one constituent tissue. Diseases/traits are ordered with respect to causal tissues. Asterisks denote significance at FDR < 5%. Dashed boxes denote results highlighted in the text. Numerical results are reported in **Supplementary Table 7**. WHRadjBMI: waist-hip-ratio adjusted for body mass index. HDL: high-density lipoprotein. DBP: diastolic blood pressure. BMI: body mass index. FEV1/FVC: forced expiratory volume in one second divided by forced vital capacity. Cereb. Cortex Ar.: cerebral cortex surface area. AST: aspartate aminotransferase. LDL: low-density lipoprotein. WBC Count: white blood cell count. MDD: major depressive disorder. Scz vs BD: schizophrenia versus bipolar disorder.

Figure 5. Comparison of disease-critical tissues identified by RTC Coloc, LDSC-SEG and TCSC. We report -$\log_{10}$FDR values (restricted to FDR ≤ 10%) for (A) RTC Coloc, (B) LDSC-SEG, (C) TCSC, across 5 representative traits (see text) and 10 tissues consisting of the causal tissues identified by TCSC and the most strongly co-regulated tagging tissues, ordered consecutively. In panel (B), results of brain-specific LDSC-SEG analyses are reported for BMI (a brain-related trait). Purple circles in panels (A) and (B) denote the causal tissue-trait pairs found by TCSC. Daggers denote meta-tissues involving more than one constituent tissue. Numerical results are reported in **Supplementary Table 15**.

999
Figure 6. Cross-trait TCSC estimates tissue-specific contributions to the genetic covariance of
two diseases/traits. (A) We report estimates of the proportion of genetic covariance explained
by the *cis*-genetic component of gene expression in tissue t' ($\zeta_{t'}$) for 30 causal tissue-trait
covariance pairs with significant contributions to trait covariance at 10% FDR, spanning 21
distinct tissues and 23 distinct trait pairs. Tissues are ordered alphabetically. Daggers denote
meta-tissues with more than one constituent tissue. Trait pairs are ordered by positive (+) or
negative (-) genetic covariance, and further ordered with respect to causal tissues. Underlined
traits are those for which TCSC identified a causal tissue in **Figure 4**: for eosinophil count, WBC
count, and platelet count the causal tissue was whole blood, and for LDL the causal tissue was
liver. Double asterisks denote trait pairs for which the differences between the tissue-specific
contribution to covariance and the tissue-specific contributions to heritability were significant
for both constituent traits and the tissue-specific contributions to heritability were non-
significant for both constituent traits. Asterisks denote significance at FDR < 5%. Dashed boxes
denote results highlighted in the text. Numerical results are reported in **Supplementary Table**
1013

20. BMI: body mass index. RBC Count: red blood cell count. WBC Count: white blood cell count.
LDL: low-density lipoprotein. Yrs Edu: years of education. WHRadjBMI: waist-hip-ratio adjusted
for body mass index. Accumbens Vol: brain accumbens volume. Caudate Vol: brain caudate
volume. MDD: major depressive disorder. T2D: type 2 diabetes. FVC: forced vital capacity. RA:
rheumatoid arthritis. (B) For BMI and red blood cell count (RBC Count), we report estimates of
the proportion of trait heritability for each trait and proportion of genetic covariance explained
by the *cis*-genetic component of gene expression in pancreas (left panel) and brain substantia
nigra (right panel). Lines with asterisks denote significant differences at 10% FDR between
respective estimates, assessed by jackknifing the differences. Numerical results are reported in
Supplementary Table 22.
1024

References

1025   **References**
1026
1.   Hekselman, I. & Yeger-Lotem, E. Mechanisms of tissue and cell-type specificity in
heritable traits and diseases. *Nat Rev Genet* **21**, 137-150 (2020).
2.   Ongen, H. *et al.* Estimating the causal tissues for complex traits and diseases. *Nat Genet*
49, 1676-1683 (2017).
3.   Gamazon, E.R. *et al.* Using an atlas of gene regulation across 44 human tissues to inform
complex disease- and trait-associated variation. *Nat Genet* **50**, 956-967 (2018).
4.   Hormozdiari, F. *et al.* Leveraging molecular quantitative trait loci to understand the
genetic architecture of diseases and complex traits. *Nat Genet* **50**, 1041-1047 (2018).
5.   Arvanitis, M., Tayeb, K., Strober, B.J. & Battle, A. Redefining tissue specificity of genetic
regulation of gene expression in the presence of allelic heterogeneity. *Am J Hum Genet*
109, 223-239 (2022).
6.   Calderon, D. *et al.* Inferring Relevant Cell Types for Complex Traits by Using Single-Cell
Gene Expression. *Am J Hum Genet* **101**, 686-699 (2017).
7.   Finucane, H.K. *et al.* Heritability enrichment of specifically expressed genes identifies
disease-relevant tissues and cell types. *Nat Genet* **50**, 621-629 (2018).
8.   Bryois, J. *et al.* Genetic identification of cell types underlying brain complex traits yields
insights into the etiology of Parkinson's disease. *Nat Genet* **52**, 482-493 (2020).
9.   Maurano, M.T. *et al.* Systematic localization of common disease-associated variation in
regulatory DNA. *Science* **337**, 1190-5 (2012).
10.   Trynka, G. *et al.* Chromatin marks identify critical cell types for fine mapping complex
trait variants. *Nat Genet* **45**, 124-30 (2013).
11.   Pickrell, J.K. Joint analysis of functional genomic data and genome-wide association
studies of 18 human traits. *Am J Hum Genet* **94**, 559-73 (2014).
12.   Finucane, H.K. *et al.* Partitioning heritability by functional annotation using genome-
wide association summary statistics. *Nat Genet* **47**, 1228-35 (2015).
13.   Roadmap Epigenomics, C. *et al.* Integrative analysis of 111 reference human
epigenomes. *Nature* **518**, 317-30 (2015).
14.   Backenroth, D. *et al.* FUN-LDA: A Latent Dirichlet Allocation Model for Predicting Tissue-
Specific Functional Effects of Noncoding Variation: Methods and Applications. *Am J Hum*
Genet **102**, 920-942 (2018).
15.   Amariuta, T. *et al.* IMPACT: Genomic Annotation of Cell-State-Specific Regulatory
Elements Inferred from the Epigenome of Bound Transcription Factors. *Am J Hum Genet*
104, 879-895 (2019).
16.   Boix, C.A., James, B.T., Park, Y.P., Meuleman, W. & Kellis, M. Regulatory genomic
circuitry of human disease loci by integrative epigenomics. *Nature* **590**, 300-307 (2021).
17.   Wainberg, M. *et al.* Opportunities and challenges for transcriptome-wide association
studies. *Nat Genet* **51**, 592-599 (2019).
18.   Consortium, G.T. The GTEx Consortium atlas of genetic regulatory effects across human
tissues. *Science* **369**, 1318-1330 (2020).
19.   Gamazon, E.R. *et al.* A gene-based association method for mapping traits using
reference transcriptome data. *Nat Genet* **47**, 1091-8 (2015).

20. Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association
studies. *Nat Genet* **48**, 245-52 (2016).
21. Siewert-Rocks, K.M., Kim, S.S., Yao, D.W., Shi, H. & Price, A.L. Leveraging gene co-
regulation to identify gene sets enriched for disease heritability. *Am J Hum Genet* **109**,
393-404 (2022).
22. Schaid, D.J., Chen, W. & Larson, N.B. From genome-wide associations to candidate
causal variants by statistical fine-mapping. *Nat Rev Genet* **19**, 491-504 (2018).
23. Gazal, S. *et al.* Linkage disequilibrium-dependent architecture of human complex traits
shows action of negative selection. *Nat Genet* **49**, 1421-1427 (2017).
24. Gazal, S., Marquez-Luna, C., Finucane, H.K. & Price, A.L. Reconciling S-LDSC and LDAK
functional enrichment estimates. *Nat Genet* **51**, 1202-1204 (2019).
25. Bulik-Sullivan, B.K. *et al.* LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat Genet* **47**, 291-5 (2015).
26. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits.
Nat Genet **47**, 1236-41 (2015).
27. Yang, J., Lee, S.H., Goddard, M.E. & Visscher, P.M. GCTA: a tool for genome-wide
complex trait analysis. *Am J Hum Genet* **88**, 76-82 (2011).
28. Mancuso, N. *et al.* Probabilistic fine-mapping of transcriptome-wide association studies.
Nat Genet **51**, 675-682 (2019).
29. Hormozdiari, F. *et al.* Widespread Allelic Heterogeneity in Complex Traits. *Am J Hum*
Genet **100**, 789-802 (2017).
30. Abell, N.S. *et al.* Multiple causal variants underlie genetic associations in humans.
Science **375**, 1247-1254 (2022).
31. Li, Z. *et al.* METRO: Multi-ancestry transcriptome-wide association studies for powerful
gene-trait association detection. *Am J Hum Genet* **109**, 783-801 (2022).
32. Yazar, S. *et al.* Single-cell eQTL mapping identifies cell type-specific genetic control of
autoimmune disease. *Science* **376**, eabf3041 (2022).
33. Liu, X. *et al.* Functional Architectures of Local and Distal Regulation of Gene Expression
in Multiple Human Tissues. *Am J Hum Genet* **100**, 605-616 (2017).
34. Gazal, S. *et al.* Combining SNP-to-gene linking strategies to identify disease genes and
assess disease omnigenicity. *Nat Genet* **54**, 827-836 (2022).
35. Weissbrod, O. *et al.* Leveraging fine-mapping and multipopulation training data to
improve cross-population polygenic risk scores. *Nat Genet* **54**, 450-458 (2022).
36. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.
Nature **562**, 203-209 (2018).
37. Gazal, S. *et al.* Functional architecture of low-frequency variants highlights strength of
negative selection across coding and non-coding annotations. *Nat Genet* **50**, 1600-1607
(2018).
38. Homan, T.D., Bordes, S. & Cichowski, E. Physiology, Pulse Pressure. in *StatPearls*
(Treasure Island (FL), 2022).
39. Kass, M.A. *et al.* The Ocular Hypertension Treatment Study: a randomized trial
determines that topical ocular hypotensive medication delays or prevents the onset of
primary open-angle glaucoma. *Arch Ophthalmol* **120**, 701-13; discussion 829-30 (2002).

40.   Zhao, D., Cho, J., Kim, M.H. & Guallar, E. The association of blood pressure and primary
1112   open-angle glaucoma: a meta-analysis. *Am J Ophthalmol* **158**, 615-27 e9 (2014).
1113   41.   Levine, R.M., Yang, A., Brahma, V. & Martone, J.F. Management of Blood Pressure in
1114   Patients with Glaucoma. *Curr Cardiol Rep* **19**, 109 (2017).
1115   42.   De Moraes, C.G., Cioffi, G.A., Weinreb, R.N. & Liebmann, J.M. New Recommendations
1116   for the Treatment of Systemic Hypertension and their Potential Implications for
1117   Glaucoma Management. *J Glaucoma* **27**, 567-571 (2018).
1118   43.   Leeman, M. & Kestelyn, P. Glaucoma and Blood Pressure. *Hypertension* **73**, 944-950
1119   (2019).
1120   44.   Soler Artigas, M. *et al.* Sixteen new lung function signals identified through 1000
1121   Genomes Project reference panel imputation. *Nat Commun* **6**, 8658 (2015).
1122   45.   Lang, I.M., Haworth, S.T., Medda, B.K., Forster, H. & Shaker, R. Mechanisms of airway
1123   responses to esophageal acidification in cats. *J Appl Physiol (1985)* **120**, 774-83 (2016).
1124   46.   Cardoso, W.V. & Lu, J. Regulation of early lung morphogenesis: questions, facts and
1125   controversies. *Development* **133**, 1611-24 (2006).
1126   47.   Shu, W. *et al.* Foxp2 and Foxp1 cooperatively regulate lung and esophagus
1127   development. *Development* **134**, 1991-2000 (2007).
1128   48.   Jacobs, I.J., Ku, W.Y. & Que, J. Genetic and cellular mechanisms regulating anterior
1129   foregut and esophageal development. *Dev Biol* **369**, 54-64 (2012).
1130   49.   Jia, X., Min, L., Zhu, S., Zhang, S. & Huang, X. Loss of sonic hedgehog gene leads to
1131   muscle development disorder and megaesophagus in mice. *FASEB J* **32**, 5703-5715
1132   (2018).
1133   50.   Shrine, N. *et al.* New genetic signals for lung function highlight pathways and chronic
1134   obstructive pulmonary disease associations across multiple ancestries. *Nat Genet* **51**,
1135   481-493 (2019).
1136   51.   Gregg, D. & Goldschmidt-Clermont, P.J. Cardiology patient page. Platelets and
1137   cardiovascular disease. *Circulation* **108**, e88-90 (2003).
1138   52.   Coppinger, J.A. *et al.* Characterization of the proteins released from activated platelets
1139   leads to localization of novel platelet proteins in human atherosclerotic lesions. *Blood*
1140   **103**, 2096-104 (2004).
1141   53.   Gawaz, M., Langer, H. & May, A.E. Platelets in inflammation and atherogenesis. *J Clin*
1142   *Invest* **115**, 3378-84 (2005).
1143   54.   Davi, G. & Patrono, C. Platelet activation and atherothrombosis. *N Engl J Med* **357**, 2482-
1144   94 (2007).
1145   55.   Badimon, L., Padro, T. & Vilahur, G. Atherosclerosis, platelets and thrombosis in acute
1146   ischaemic heart disease. *Eur Heart J Acute Cardiovasc Care* **1**, 60-74 (2012).
1147   56.   Meadows, T.A. & Bhatt, D.L. Clinical aspects of platelet inhibitors and thrombus
1148   formation. *Circ Res* **100**, 1261-75 (2007).
1149   57.   Berman, M.N., Tupper, C. & Bhardwaj, A. Physiology, Left Ventricular Function. in
1150   *StatPearls* (Treasure Island (FL), 2022).
1151   58.   Chung, S., Sawyer, J.K., Gebre, A.K., Maeda, N. & Parks, J.S. Adipose tissue ATP binding
1152   cassette transporter A1 contributes to high-density lipoprotein biogenesis in vivo.
1153   *Circulation* **124**, 1663-72 (2011).

1154   59.   McGillicuddy, F.C., Reilly, M.P. & Rader, D.J. Adipose modulation of high-density
1155   lipoprotein cholesterol: implications for obesity, high-density lipoprotein metabolism,
1156   and cardiovascular disease. *Circulation* **124**, 1602-5 (2011).
1157   60.   Zoccali, C. *et al.* Adiponectin, metabolic risk factors, and cardiovascular events among
1158   patients with end-stage renal disease. *J Am Soc Nephrol* **13**, 134-141 (2002).
1159   61.   Ryo, M. *et al.* Adiponectin as a biomarker of the metabolic syndrome. *Circ J* **68**, 975-81
1160   (2004).
1161   62.   Toth, P.P. Adiponectin and high-density lipoprotein: a metabolic association through
1162   thick and thin. *Eur Heart J* **26**, 1579-81 (2005).
1163   63.   Van Linthout, S. *et al.* Impact of HDL on adipose tissue metabolism and adiponectin
1164   expression. *Atherosclerosis* **210**, 438-44 (2010).
1165   64.   Shungin, D. *et al.* New genetic loci link adipose and insulin biology to body fat
1166   distribution. *Nature* **518**, 187-196 (2015).
1167   65.   Emdin, C.A. *et al.* Genetic Association of Waist-to-Hip Ratio With Cardiometabolic Traits,
1168   Type 2 Diabetes, and Coronary Heart Disease. *JAMA* **317**, 626-634 (2017).
1169   66.   Smith, J., Al-Amri, M., Sniderman, A. & Cianflone, K. Leptin and adiponectin in relation to
1170   body fat percentage, waist to hip ratio and the apoB/apoA1 ratio in Asian Indian and
1171   Caucasian men and women. *Nutr Metab (Lond)* **3**, 18 (2006).
1172   67.   Farooqi, I.S. Defining the neural basis of appetite and obesity: from genes to behaviour.
1173   *Clin Med (Lond)* **14**, 286-9 (2014).
1174   68.   Locke, A.E. *et al.* Genetic studies of body mass index yield new insights for obesity
1175   biology. *Nature* **518**, 197-206 (2015).
1176   69.   Medic, N. *et al.* Increased body mass index is associated with specific regional
1177   alterations in brain structure. *Int J Obes (Lond)* **40**, 1177-82 (2016).
1178   70.   Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body
1179   mass index in approximately 700000 individuals of European ancestry. *Hum Mol Genet*
1180   **27**, 3641-3649 (2018).
1181   71.   Wojcinski, A. *et al.* Cerebellar granule cell replenishment postinjury by adaptive
1182   reprogramming of Nestin(+) progenitors. *Nat Neurosci* **20**, 1361-1370 (2017).
1183   72.   Andreotti, J.P. *et al.* Neurogenesis in the postnatal cerebellum after injury. *Int J Dev*
1184   *Neurosci* **67**, 33-36 (2018).
1185   73.   Hait, E.J. & McDonald, D.R. Impact of Gastroesophageal Reflux Disease on Mucosal
1186   Immunity and Atopic Disorders. *Clin Rev Allergy Immunol* **57**, 213-225 (2019).
1187   74.   Plessen, K.J. *et al.* Hippocampus and amygdala morphology in attention-
1188   deficit/hyperactivity disorder. *Arch Gen Psychiatry* **63**, 795-807 (2006).
1189   75.   Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for attention
1190   deficit/hyperactivity disorder. *Nat Genet* **51**, 63-75 (2019).
1191   76.   Enard, W. *et al.* A humanized version of Foxp2 affects cortico-basal ganglia circuits in
1192   mice. *Cell* **137**, 961-71 (2009).
1193   77.   Floresco, S.B., Todd, C.L. & Grace, A.A. Glutamatergic afferents from the hippocampus
1194   to the nucleus accumbens regulate activity of ventral tegmental area dopamine
1195   neurons. *J Neurosci* **21**, 4915-22 (2001).

78.   Kim, M.S. *et al.* Prefrontal Cortex and Amygdala Subregion Morphology Are Associated
With Obesity and Dietary Self-control in Children and Adolescents. *Front Hum Neurosci*
14, 563415 (2020).
79.   Elsayed, E.F. *et al.* Waist-to-hip ratio and body mass index as risk factors for
cardiovascular events in CKD. *Am J Kidney Dis* **52**, 49-57 (2008).
80.   Peterson, C.G., Skoog, V. & Venge, P. Human eosinophil cationic proteins (ECP and EPX)
and their suppressive effects on lymphocyte proliferation. *Immunobiology* **171**, 1-13
(1986).
81.   Nakagome, K. *et al.* IL-5-induced hypereosinophilia suppresses the antigen-induced
immune response via a TGF-beta-dependent mechanism. *J Immunol* **179**, 284-94 (2007).
82.   Onyema, O.O. *et al.* Eosinophils downregulate lung alloimmunity by decreasing TCR
signal transduction. *JCI Insight* **4**(2019).
83.   Wei, H.S. *et al.* Erythrocytes Are Oxygen-Sensing Regulators of the Cerebral
Microcirculation. *Neuron* **91**, 851-862 (2016).
84.   Olusi, S.O. Obesity is an independent risk factor for plasma lipid peroxidation and
depletion of erythrocyte cytoprotectic enzymes in humans. *Int J Obes Relat Metab*
Disord **26**, 1159-64 (2002).
85.   Ozata, M. *et al.* Increased oxidative stress and hypozincemia in male obesity. *Clin*
Biochem **35**, 627-31 (2002).
86.   Druml, W., Laggner, A.N., Lenz, K., Grimm, G. & Schneeweiss, B. Pancreatitis in acute
hemolysis. *Ann Hematol* **63**, 39-41 (1991).
87.   Sakai, N.S., Taylor, S.A. & Chouhan, M.D. Obesity, metabolic disease and the pancreas-
Quantitative imaging of pancreatic fat. *Br J Radiol* **91**, 20180267 (2018).
88.   Schafer, W.R. & Kenyon, C.J. A calcium-channel homologue required for adaptation to
dopamine and serotonin in Caenorhabditis elegans. *Nature* **375**, 73-8 (1995).
89.   Weinshenker, D., Garriga, G. & Thomas, J.H. Genetic and pharmacological analysis of
neurotransmitters controlling egg laying in C. elegans. *J Neurosci* **15**, 6975-85 (1995).
90.   Triarhou, L.C. Introduction. Dopamine and Parkinson's disease. *Adv Exp Med Biol* **517**, 1-
14 (2002).
91.   Zielonka, M. *et al.* Dopamine-Responsive Growth-Hormone Deficiency and Central
Hypothyroidism in Sepiapterin Reductase Deficiency. *JIMD Rep* **24**, 109-13 (2015).
92.   Comings, D.E. *et al.* The dopamine D2 receptor (DRD2) as a major gene in obesity and
height. *Biochem Med Metab Biol* **50**, 176-85 (1993).
93.   Yao, D.W., O'Connor, L.J., Price, A.L. & Gusev, A. Quantifying genetic effects on disease
mediated by assayed gene expression levels. *Nat Genet* **52**, 626-633 (2020).
94.   Leyden, G.M. *et al.* Harnessing tissue-specific genetic variation to dissect putative causal
pathways between body mass index and cardiometabolic phenotypes. *Am J Hum Genet*
109, 240-252 (2022).
95.   Thom, C.S., Wilken, M.B., Chou, S.T. & Voight, B.F. Body mass index and adipose
distribution have opposing genetic impacts on human blood traits. *Elife* **11**(2022).
96.   Umans, B.D., Battle, A. & Gilad, Y. Where Are the Disease-Associated eQTLs? *Trends*
Genet **37**, 109-124 (2021).
97.   Weissbrod, O. *et al.* Functionally informed fine-mapping and polygenic localization of
complex trait heritability. *Nat Genet* **52**, 1355-1363 (2020).

98. Nathan, A. *et al.* Single-cell eQTL models reveal dynamic T cell state dependence of
disease loci. *Nature* **606**, 120-128 (2022).
99. Perez, R.K. *et al.* Single-cell RNA-seq reveals cell type-specific molecular and genetic
associations to lupus. *Science* **376**, eabf1970 (2022).
100. Soskic, B. *et al.* Immune disease risk variants regulate gene expression dynamics during
CD4(+) T cell activation. *Nat Genet* **54**, 817-826 (2022).
101. Stegle, O., Parts, L., Durbin, R. & Winn, J. A Bayesian framework to account for complex
non-genetic factors in gene expression levels greatly increases power in eQTL studies.
PLoS Comput Biol **6**, e1000770 (2010).
102. Genomes Project, C. *et al.* An integrated map of genetic variation from 1,092 human
genomes. *Nature* **491**, 56-65 (2012).
1251