1
2
3 **Target-enriched enzymatic methyl sequencing: flexible, scalable and**
4 **inexpensive hybridization capture for quantifying DNA methylation**
5
6

7 Dustin R. Rubenstein[a,b,1] and Joseph Solomon[a]
8
9
10 [a] Department of Ecology, Evolution and Environmental Biology, Columbia University, New
11 York, NY 10027, USA
12 [b] Center for Integrative Animal Behavior, Columbia University, New York, NY 10027, USA
13
14 [1] Corresponding Author: Phone: +1-212-854-4881; Email: dr2497@columbia.edu

15
16 Running Head: Targeted capture to study DNA methylation

17   **Abstract**
18
The increasing interest in studying DNA methylation to understand how traits or diseases
develop requires new and flexible approaches for quantifying methylation states in a diversity of
organisms. Here, we develop TEEM-Seq (target-enriched enzymatic methyl sequencing), a
method that combines enzymatic methyl sequencing with a custom-designed hybridization
capture bait set that can be scaled to reactions including large numbers of samples in any species
for which a reference genome is available. TEEM-Seq is able to quantify DNA methylation
states similarly well to the more traditional approaches of whole-genome and reduced-
representation sequencing, and the downstream bioinformatic analysis is the same as for any
sequence-based approach to studying DNA methylation. By maximizing the number of samples
in the hybridization reaction and minimizing sequencing depth, TEEM-Seq is an inexpensive and
flexible sequence-based approach for quantifying DNA methylation in species where other
capture-based methods are unavailable or too expensive, particularly for non-model organisms.
31
32   **Keywords**: TEEM-Seq, epigenetics, DNA methylation, CpG site, hybridization capture, target
33   enrichment

Introduction

There is increasing recognition in fields as diverse as ecology, evolutionary biology, conservation biology, neuroscience, and biomedicine that epigenetics is critical to understanding how traits, diseases, or other phenotypes develop (Waddington 1942, Jablonka and Lamb 2006, Cavalli and Heard 2019, Rey et al. 2020). One of the most widely studied epigenetic mechanisms underlying phenotypic variation is DNA methylation (Robertson and Wolffe 2000, Bossdorf et al. 2008), the addition of a methyl group to a DNA nucleotide that impacts gene expression through transcriptional silencing (Bird 1986). In mammals and many other vertebrates, DNA methylation often occurs at CpG sites, or regions of DNA where a cytosine nucleotide is followed by a guanine nucleotide. With the increasing availability of reference genomes for non-model organisms and the continually decreasing costs of high-throughput sequencing, sequencing-based approaches are becoming the preferred method for studying DNA methylation across the diversity of life because they can provide quantitative values of CpG methylation states (Harris et al. 2010). Sequence-based approaches to studying DNA methylation range from whole- or reduced-representation sequencing of genomes to sequencing one or a few candidate genes, yet the performance and cost of different sequence-based approaches to quantifying and mapping DNA methylation varies greatly (Bock et al. 2010, Consortium 2016, Crary-Dooley et al. 2017, Gouil and Keniry 2019). Whole-genome sequencing for DNA methylation profiling remains the gold standard, but is costly because of the necessary sequencing depth (Harris et al. 2010). In contrast, reduced-representation approaches are cheaper than whole-genome sequencing because they cover only a fraction of the genome, but since such approaches require restriction enzymes to target specific genomic regions (e.g., CpG islands), they inevitably miss potentially important CpG sites in genomic regions of interest (Husby 2022). Despite the differences in cost and coverage, most whole-genome and reduced-representation sequencing approaches share the chemical conversion of unmethylated cytosines to uracils with bisulfite treatment (Gu et al. 2011), a method that enables quantification of methylation states but that can fragment DNA and complicate downstream library preparation and even bias results (Tanaka and Okamato 2007, Gouil and Keniry 2019).

As the number of studies of DNA methylation grows, it becomes increasingly important to develop new methods that can quantitatively measure CpG methylation states over large regions of the genome in an efficient yet cost-effective way. Such approaches exist for humans and some model organisms in the form of bead or probe panels, but flexible methods for studying targeted genomic regions in any organism are much rarer. Although hybridization capture using biotinylated oligonucleotide baits (probes) to hybridize specific regions of interest across the genome of any species have become standard in molecular biology (Kozarewa et al. 2015), their application in epigenetics has been more limited (Lee et al. 2011, Li et al. 2015, Masser et al. 2016). Commercial approaches such as Agilent's SureSelect[XT] Methyl-Seq Target Enrichment System and Roche's SeqCap Epi Enrichment System are neither particularly scalable nor cost-effective, and both use bisulfite conversion. Yet, with the recent development of alternative approaches to bisulfite treatment for converting DNA (Vaisvila et al. 2020), newer methods of hybridization capture may be viable for quantitatively studying DNA methylation, particularly in non-model organisms.

Here, we develop a flexible, scalable, and inexpensive method called target-enriched enzymatic
methyl sequencing (TEEM-Seq) that uses hybridization capture to quantitively measure CpG
methylation states. We achieve this goal by combining enzymatic methyl sequencing (EM-seq)
(Vaisvila et al. 2020) with a custom-designed, species-specific hybridization capture bait set that
can be scaled to reactions that include much larger numbers of samples than other currently
available methods and that only requires low amounts of input DNA to capture both 5mC and
5hmC methylation. EM-seq effectively overcomes DNA fragmentation caused by bisulfite
treatment, and a custom bait set allows for cost-effective and efficient targeting of (multiple)
specific genomic regions. In this study, we selected genomic regions encompassing selected
genes known to be associated with the hypothalamic-pituitary-adrenal axis (HPA), the
hypothalamic–pituitary–gonadal axis (HPG), and DNA methyltransferase activity. We chose
these specific HPA and HPG genes because of the critical roles that they play in the avian stress
response and in reproduction in vertebrates (Sapolsky et al. 2000). After designing baits targeting
all of the putative promoters (4 kb upstream of the transcription start site) and some of the exons
of 22 of these genes, we developed the TEEM-Seq protocol using DNA extracted from the blood
of the superb starling (*Lamprotornis superbus*), a passerine bird with a published reference
genome (Rubenstein et al. 2021). We also tested our starling bait set performance on another
passerine species, the house sparrow (*Passer domesticus*). To gauge the effectiveness and
efficiency of TEEM-Seq against more traditional methods of quantifying DNA methylation, we
compared data from some of the same individual starlings to data from whole-genome enzymatic
methyl sequencing (WGEM-Seq) and/or reduced-representation bisulfite sequencing (RRBS).
We found that necessary sequencing coverage levels were similar to RRBS, but an order of
magnitude lower than for WGEM-Seq for similarly high-quality coverage of complete target
regions. Ultimately, we demonstrate the utility of TEEM-Seq for quantifying DNA methylation
in an efficient and cost-effective manner that should be applicable to studying any organism in a
variety of tissue and sample types, including formalin-fixed, paraffin-embedded (FFPE) samples
and those from degraded or ancient DNA.
106
107   **Results**
108
TEEM-Seq metrics
Our results from a pool of 48 libraries prepared using avian DNA from whole blood in a single
capture reaction sequenced on an Illumina HiSeq at 110 Gb (2.3 Gb per sample) indicated that
target read coverage for the superb starling samples prior to deduplication ranged from a mean of
400-4300x, with an overall mean of 1700x for both DNA strands. Moreover, background, non-
target coverage was mostly scattered but evenly distributed single strand alignment, with a mean
of 22x. We determined that while duplication of total reads was 90-95% according to Bismark
deduplication reports (versus 15-17% duplicated reads for three WGEM-Seq starling samples,
which suggests that sequencing optimization for this protocol might first focus post-capture
before reducing pre-capture PCR cycles and per-sample template for the capture reaction),
TEEM-Seq target region read coverage depth after deduplication ranged from an average per
target of 20-270x, with an overall mean coverage depth for all targets of 173x (Figure 1). In
contrast, WGEM-Seq target region coverage after deduplication ranged from an average per
target of 2-41x, with an overall mean coverage of 15x (Figure 1). Figure 2 illustrates
deduplicated read coverage depth for TEEM-Seq and WGEM-Seq for three representative target
regions using the Integrative Genomics Viewer (IGV) v2.12.3 (Robinson et al. 2011). In

addition, the average mean depth of TEEM-Seq coverage after deduplication for just the seven
starling samples also included in the RRBS study ranged from 30-374x for the lowest to highest
depth promoter targets, with an overall average of 248x. For those same RRBS libraries, the
average mean coverage depth per target promoter region (non-deduplication) ranged from 0-17x,
with an overall mean of 7x. Although RRBS coverage is biased toward restriction sites (C/CGG
for MspI) and certain targets had zero coverage, on average 120 CpGs were covered above 5x
across strictly 4 kb promoter regions (Table S4). With the TEEM-Seq and WGEM-Seq libraries,
the average was also coincidentally 120 CpGs above 5x in the three samples compared across
methods, though the minimum number of 5x sites covered was never zero for either method
(Table S2). Finally, for four samples run as duplicate or triplicate libraries, there were no
significant differences in mean DNA methylation between the replicates (all paired $t_{23} < 1.63$, all
$P > 0.12$; Table S1).
137
Assessing bait performance in another avian species
Although our bait set was designed specifically for the superb starling, we included some house
sparrow samples to assess starling-derived bait performance in another passerine bird species.
For four house sparrow libraries, SAMtools reported that 19 of 23 putative promoter targets had
an average percent of covered bases of <50%, and 11 of 23 targets had <25% coverage. In
contrast, 20 of 23 targets had an average percent of covered bases >99.5% for the 44 superb
starling libraries, with *POMC* having 96% covered bases, and *NR3C1* and *VTG* having reference
gaps of 488 bp and 182 bp, respectively, but still >99% coverage elsewhere in the putative
promoters. The maximum average for the target with the highest coverage was 71% in the house
sparrow, and the mean depth of coverage after deduplication ranged from an average across
samples of 0-39.5x (versus 20-270x for the superb starling) for the lowest to highest depth
promoter targets, with an overall average of 9.3x (versus 173x for the superb starling). These
results suggest that superb starling bait sequence was not well conserved for most targets in
house sparrows, and that species-specific baits should be designed for each study organism using
its own reference genome whenever possible.
153
TEEM-Seq comparison and validation
To compare and validate TEEM-Seq against more traditional methods of analysis for DNA
methylation, we compared many of the same starling DNA samples prepared for WGEM-Seq
and RRBS. At the CpG site-level, TEEM-Seq and WGEM-Seq libraries from the same
individual were highly correlated (Figure 3). Moreover, we found no major differences in mean
DNA methylation between approaches. CpG methylation varied little at the site-level when
absolute value mean differences in DNA methylation for sites were compared across gene bodies
and putative promoter regions (Figures 4a-c), or when sites were averaged across target regions,
as methylation did not differ between WGEM-Seq and TEEM-Seq data within putative promoter
($F_{1,134} = 0.0013$, $P = 0.97$; Figure 4d and Table S2) or exonic regions ($F_{1,68} = 0.00001$, $P = 0.99$;
Table S3), between TEEM-Seq and RRBS data within putative promoter regions ($F_{1,249} = 0.049$,
$P = 0.83$; Figure 4e and Table S4), or between WGEM-Seq and RRBS in 20 random 2.5 kb
"promoter" regions ($F_{1,114} = 1.36$, $P = 0.25$; Figure 4f and Table S5). Finally, we note that
TEEM-Seq data with and without deduplication were nearly identical (Figure S1) and that
despite the high number of duplicates in the TEEM-Seq data, comparisons of WGEM-Seq versus
TEEM-Seq were qualitatively similar when using TEEM-Seq data with and without
deduplication (Tables S2 and S6). As expected, CpG site-level correlations were still higher for

TEEM-Seq deduplicated versus non-deduplicated data when using the same sites that were
highly correlated between TEEM-Seq and WGEM-Seq data (Figure S1 legend and Figure 3).
Overall, mean DNA methylation was comparable in all three methods, indicating that TEEM-
Seq is an effective approach for quantifying DNA methylation states in specific regions of the
genome.
176
177
178 **Discussion**
179
We introduce a flexible, scalable, and inexpensive method for studying DNA methylation in a
large number of candidate genes that combines enzymatic conversion of DNA (Vaisvila et al.
2020) with a custom-designed hybridization capture bait set, and uses starting input DNA as low
as 100 ng (though we expect lower amounts would also work). The TEEM-Seq approach
described here is able to quantify DNA methylation states similarly well to the more traditional
approaches of whole-genome and reduced-presentation sequencing (Figure 5), and the
downstream bioinformatic analysis is the same for all three approaches. We show that TEEM-
Seq is both reliable and repeatable, as duplicate libraries from the same samples did not vary
much. Although we demonstrate its effectiveness on DNA from birds, TEEM-Seq should work
on DNA from any organism, as well as from DNA extracted from tissues other than blood (Siller
2022). And because it employs enzymatic rather than chemical conversion of DNA (i.e., bisulfite
treatment), we also expect that TEEM-Seq will be useful for studying formalin-fixed, paraffin-
embedded (FFPE) samples and degraded or ancient DNA, particularly samples from museum
specimens that are likely to already be partially degraded and only damaged further by traditional
bisulfite conversion (Rubi et al. 2020).
195
One of the greatest strengths of TEEM-Seq is its scalability. Although we only scaled our
capture reaction to 48 libraries here, we note that in a parallel TEEM-Seq study using a similar
bait set, capture scaled consistently with a 96-library pool (Siller 2022). Moreover, New England
Biolabs currently produces 120 total unique indexes across their 24- and 96-sample EM-seq kits,
meaning that TEEM-Seq should be readily able to scale to a 120-library pool, and perhaps even
further with the development of additional unique indexes. We caution, however, that there is
likely to be a tradeoff in the number of baits used in the bait set and the number of samples used
in the hybridization capture, though assessing this was beyond the scope of the current study.
Since myBaits 80 nt baits are designed to tolerate >20% target divergence in conventional DNA-
Seq, it is likely that the eight methylation-state targets myBaits recommended for this pilot bait
design in addition to the unmodified targets were to some degree redundant. If this proves to be
the case, costs could be reduced further in bait design, though assessing the capture performance
and mismatch allowance of the unmodified baits and redundancy of methylation-state bait
permutations was beyond the scope of this study. Although we cannot directly assess bias in
whether and how specific baits hybridize and the affect on methylation levels, under the current
design parameters, no systemic bias was evident in capture products. Ultimately, TEEM-Seq is
highly scalable because the conversion of unmethylated cytosines to uracils is done prior to the
hybridization reaction. In contrast, Agilent's SureSelect[XT] Methyl-Seq Target Enrichment
System performs hybridization capture before library pooling, which limits the number of
samples to a fraction of that possible in each TEEM-Seq capture reaction (Lee et al. 2011). And
while Roche's SeqCap Epi Enrichment System (Li et al. 2015) performs hybridization capture

after bisulfite conversion, it is also currently limited to a fraction of those indexes available for
EM-seq, which similarly limits the number of samples in each capture reaction.
219
Finally, TEEM-Seq is more cost-effective and per sample data-efficient than not only other
commercial capture-based approaches, but also than whole-genome sequencing. It is comparable
in cost to reduced-representation sequencing. For an avian species with a genome size of roughly
1.07 Gb (Rubenstein et al. 2021), we sequenced 2.3 Gb per sample for TEEM-Seq, 1.3 Gb per
sample for RRBS, and 27.5 Gb per sample for WGEM-Seq. We note, however, that even with an
order of magnitude lower sequencing demand for TEEM-Seq compared to WGEM-Seq, the
specificity of the bait set meant that we routinely produced 100x coverage or more of the targets,
including control regions, and all target regions had average coverage of at least 20x after
deduplication. Sequencing depth for TEEM-Seq could be further reduced with additional
optimization and perhaps with a reduction in bait redundancy. In addition, since we were able to
scale EM-seq kit reagents down to three quarter volumes, by maximizing the number of samples
per total capture pool (up to 120 samples per reaction with current index availability), TEEM-
Seq becomes an inexpensive method for accurately quantifying DNA methylation states across
continuous regions of the genome in a large sample for a diversity of organisms. Finally, all three
methods are roughly similar in terms of library preparation time, with RRBS and WGEM-Seq
taking roughly two days and TEEM-Seq three to four days because of the added capture step.
236
Ultimately, we expect that TEEM-Seq could replace traditional approaches for studying single
candidate genes (e.g., amplicon sequencing or pyrosequencing), which are time and labor
intensive, and be effectively paired with other DNA methylation quantification methods (e.g.,
whole-genome or reduced-representation sequencing) to increase project sample sizes.
Moreover, TEEM-Seq can be paired with RNA-Seq to examine how DNA methylation in
promoters or other regulatory regions is related to the expression of individual genes or gene
networks. As enzymatic conversion of DNA and EM-seq approaches continue to expand and
replace bisulfite conversion and sequencing (e.g., Feng et al. 2020, Han et al. 2021, Liu et al.
2021, Morrison et al. 2021), we expect that TEEM-Seq can be further optimized to improve
efficiency, scale to ever larger numbers of samples, and use other bait set designs. Ultimately,
because of its flexibility, scalability, and cost-effectiveness, TEEM-Seq should prove useful to
researchers in a number of biological disciplines for rapidly quantifying DNA methylation states
in any organism of interest in order to link epigenetics to the development of phenotypes.
250
251
252   **Methods**
253
Sample collection
Superb starlings (N = 39) were captured using baited traps or mist nests at the Mpala Research
Centre, Laikipia, Kenya (0° 17′N, 37° 52′E). Whole blood was collected in Queens lysis buffer
(Seutin et al. 1991) and genomic DNA was extracted using a DNeasy Blood & Tissue Kit
(Qiagen). All research was approved by the Institutional Animal Care and Use Committee at
Columbia University (AC-AAAW6451), as well as the Kenyan National Commission for
Science, Technology and Innovation, the Kenyan National Environmental Management
Authority, the Kenya Wildlife Service, and the Mpala Research Centre.
262

Library preparation
After DNA was quantified using a Qubit 3.0 Fluorometer (Thermo Fisher Scientific), 100-150 ng
was sheared to 250-275 bp on a Covaris ME220 and inspected on an Agilent 4200 TapeStation.
The sheared DNA was then used as a template for libraries prepared with a NEBNext Enzymatic
Methyl-seq Kit (EM-seq) (New England Biolabs). We note that previous attempts to develop a
hybridization capture protocol using bisulfite-converted DNA were unsuccessful, presumably
because chemical conversion led to fragmented DNA and failed hybridization reactions.
However, the enzymatic conversion overcame challenges associated with fragmented DNA from
bisulfite conversion and allowed for successful hybridization capture. Following the EM-seq Kit
protocol, sheared DNA was end repaired at 20°C, ligated with an EM-seq adaptor at 20°C, and
then used as template in a TET2 reaction at 37°C for oxidation of 5-Methylcytosines and 5-
Hydroxymethylcytosines. The TET2 converted DNA was denatured with formamide at 85°C,
underwent APOBEC cytosine deamination at 37°C, and was then PCR amplified with index
primers and NEBNext Q5U Master Mix at an annealing temperature of 62°C. Since we intended
to perform downstream in-solution hybridization capture, we increased library yield through 12
total PCR cycles for the enzymatic conversion amplification compared with the standard
maximum of eight cycles. Although we did not vary the number of PCR cycles in this study, this
could be optimized in the future to possibly reduce PCR duplicates. Whereas the duplicate rate
increased most in the hybridization reaction and amplification, it remains unknown whether EM-
seq duplicates were compounded in the capture process through bait bias toward more highly
represented duplicate molecules. We also increased all SPRI bead volumes by 50%,
supplementing the included sample purification beads with AMPure XP beads (Beckman
Coulter) at the PCR clean-up stage. For post-ligation clean-up, we used a ratio of beads to
sample volume of 1.76:1 instead of the NEB EM-seq standard protocol's 1.18:1; for the TET2
reaction cleanup, we used a ratio of 2.65:1 instead of 1.76:1; for the APOBEC deanimation
clean-up we used 1.5:1 instead of 1:1; and for post-amplification clean-up we used 1.35:1 instead
of 0.9:1. We increased the ratio of beads to sample volume because we intended the EM-seq
libraries for a myBaits capture pool, not for direct sequencing. In addition, we varied kit reagent
volumes down to half of those recommended, ultimately settling on three quarter volumes for
our final reactions. Importantly, results with three quarter volumes of reagents were qualitatively
similar to those with full volumes (data not shown). Finally, we Qubit-quantified the libraries,
inspected them on the TapeStation, and pooled 25 ng from each library into 1.2 µg total for the
pool of 48 libraries.
296
Bait design
Target sequences for bait design were identified from the superb starling reference genome
(CU_Lasu_v1) (Rubenstein et al. 2021). Baits targeted 23 regions (*GNRHR2* had two separate
regions) of 4 kb in length upstream of the transcription start site (i.e., putative promoter regions)
and exons (excluding introns) from a subset of genes known to be associated with the HPA axis,
the HPG axis, and DNA methyltransferases (Table 1). Biotinylated RNA baits were then
prepared commercially using myBaits v4.01 Custom 1-20K 16 Reaction Kits (Daicel Arbor
Biosciences). We submitted 71 sequences (113,940 bp) for 80 nt bait design at 2x tiling density.
For each bait candidate, in addition to the unmodified original, myBaits designed eight additional
genome variants to reasonably cover conversion permutations (i.e., all methylated, a random
50% CpGs methylated, the other 50% CpGs methylated, unmethylated, and sense/antisense for
each version), resulting in 21,792 baits, all of which were quality-assessed and filtered by

myBaits Design based on masked repeats in the original sequences (repeatmasker [engine:
cross_match, organism: Aves] resulting in 5.4% of total sequences masked), GC content, and
BLAST hits on the zebra finch (*Taeniopygia guttata*), European starling (*Sturnus vulgaris*),
collared flycatcher (*Ficedula albicollis*), and great tit (*Parus major*) reference genomes for the
final myBaits kits (Tables S7 and S8). In addition to unmodified versions of each genome,
myBaits blasted bait candidates against sense strand unmethylated (i.e., all Cs converted to Ts)
and anti-sense strand unmethylated (i.e., all Gs converted to As) versions of each genome, for a
total of 12 genome variants. The top 500 blast hits (by bit score) were tabulated for each bait
candidate. The final bait set contained sequences with <25% repeat-masking, 10-80% GC
content, and ≤10 max BLAST hits across all 12 genome variants to avoid non-specific targets.
319
Hybridization capture
A pool of 48 libraries contained 39 starling samples, two house sparrow samples (see below),
and a total of seven replicate libraries from each species. Eight superb starling samples were used
in a previously unpublished DNA methylation study that used RRBS. Three of these eight were
included in duplicate libraries and one in triplicate libraries. The remaining 31 additional superb
starling samples were included to assess target capture performance when scaling up to a pool of
48 libraries. Although our bait set was designed specifically for the superb starling, we also
included two house sparrow samples run in duplicate to assess starling-derived bait performance
in more highly conserved target regions of another avian species.
329
We followed the myBaits v4.01 protocol, but lowered the default hybridization and clean up
temperature of 65°C to 63°C as myBaits recommends for fragmented DNA. We note, however,
that a parallel TEEM-Seq study worked similarly well using a temperature of 65°C (Siller 2022).
Hybridization beads bound to the pooled library-blocker mix were cleaned with three washes,
and the resuspended bead-bound DNA was then amplified with 16 PCR cycles in a KAPA HiFi
reaction using a 60°C annealing temperature and a one minute extension step at 72°C. The
amplified capture pool was subsequently cleaned up with AMPure XP beads and sequenced at
2x150 bp reads with 5% PhiX spike-in in one lane (110 Gb) of an Illumina HiSeq 4000 at the
Novogene Corporation, Sacramento, CA.
339
Control sequences
Bait library sequence alignments to NEB pUC19 methylated and lambda unmethylated controls
were sparse in background coverage because controls were not specifically targeted in our initial
bait set. Overall, coverage in control regions ranged from 0-42x without deduplication, averaging
18x for pUC19 and 25x for lambda, but methylation calls were unreliable with deduplication.
Although our three EM-seq libraries that were also used for WGEM-Seq passed control checks,
and the RRBS libraries used to compare CpG methylation percentages were spiked with lambda
controls, we confirmed that a known low methylation (average 1.5%) 57 bp region with 12 CpGs
10 bp upstream of Arginine vasopressin receptor 1A (Rubenstein and Solomon unpublished data)
also had means within 1% of this average in the seven TEEM-Seq libraries used for comparison
to other methods. Nonetheless, we then specifically targeted pUC19 and lambda control
sequences in a parallel TEEM-Seq study using a house sparrow-specific bait set to ensure that
we would have reliable EM-seq conversion estimates for all samples (Siller 2022), in the absence
of comparable libraries for these samples prepared with alternative methods. In that study, we
found that lambda unmethylated and pUC19 methylated control baits were highly specific,

targeting the first 1 kb for pUC19 and the first 2 kb for lambda included in the bait design. For a
capture pool of 96 libraries, SAMtools v1.14 (Danecek et al. 2021) reported coverage for the 1
kb pUC19 alignments as a range of 18,582-538,322x (mean = 96,689x) for non-deduplicated
BAMs and a range of 4,540-86,441x (mean = 27,929x) for deduplicated BAMs. For the 2 kb
Lambda alignments, SAMtools reported a range of 23,445-537,148x (mean = 115,380x) for non-
deduplicated BAMs and a range of 4,984-88,019x (mean = 30,943x) for deduplicated BAMs. To
ensure requisite control alignments with an untested protocol, and because we only targeted a
portion of each control genome, we followed EM-seq recommendations for a higher spike-in
with 0.3 ng of pUC19 and 6 ng of lambda controls per 100-150 ng sample. However, given the
excessive depth of capture, a spike-in at normal HiSeq levels (a 1:100 dilution, or 0.001 ng
pUC19 and 0.02 ng lambda) should be more than sufficient when control regions are targeted.
Since the bait design and hybridization reaction had no negative effects on the assessment of
control alignments, we recommend that all capture designs for TEEM-Seq libraries follow this
protocol and include baits for pUC19 and lambda controls.
369
TEEM-Seq comparison and validation
To compare and validate TEEM-Seq against more traditional methods of analysis for DNA
methylation, we compared many of the same starling DNA samples to those that underwent
WGEM-Seq and/or RRBS. For WGEM-Seq, extra material from three of the EM-seq libraries
were confirmed clear of adapter dimers on the TapeStation, pooled and sequenced at 2x150 bp
reads with 5% PhiX spike-in in 3/4[th] of a lane (82.5 Gb) on a HiSeq 4000 at Novogene with the
same parameters to provide a whole-genome sequence and partial methylome analysis
comparison to our targeted sequencing. For a previous RRBS study (Rubenstein and Solomon
unpublished data), we used NuGEN Ovation Methyl-Seq kits to prepare RRBS libraries (with
added Promega unmethylated cl857 Sam7 lambda control spike-in of 1 ng per 80-120 ng of
sample), which were sequenced in several pools of 16 samples per lane (20 Gb) at 1x100 bp
reads with 10% PhiX spike-in on a HiSeq 2500 at the Cornell Institute of Biotechnology
Genomics Core, Ithaca, NY. Libraries for RRBS were prepared according to kit protocols.
Briefly, samples were digested with MspI at 37°C, ligated with bisulfite-compatible barcoded
adaptors at 25°C, end repaired at 60°C, bisulfite converted with the Qiagen EpiTect Fast
Bisulfite Conversion Kit, PCR amplified with Nugen's standard 12 cycles and 60°C annealing,
and purified with Agencourt beads for sequencing. Eight of these RRBS samples were included
in the present study, though one failed enzymatic conversion. Data from the remaining seven
samples (from different sequencing pools) were used as a comparison to the TEEM-Seq and
WGEM-Seq derived data generated from the same individuals.
390
DNA methylation analysis
Sequencing data were trimmed with the Trim Galore v0.4.2 (Krueger et al. 2021) wrapper of
Cutadapt v1.12 (Martin 2011) with standard parameters. As per NuGEN's recommendations,
RRBS reads were not trimmed with Trim Galore's RRBS option, but instead trimmed of
adaptors with standard parameters and with NuGEN's trimRRBSdiversityAdaptCustomers.py
script. Trimmed reads were aligned to the superb starling bisulfite genome reference generated
by Bismark v0.19.0 (Krueger and Andrews 2021). WGEM-Seq and TEEM-Seq alignments were
deduplicated, and CpG coverage files with methylation percentages (100 * methylated
cytosines/total of methylated plus unmethylated cytosines in CpG context) were extracted from
alignments with Bismark (non-deduplicated methylation calls were also made for TEEM-Seq

data to compare percentages with and without deduplication). Although there does not appear to
be a consensus on whether target-enriched libraries should be deduplicated (for amplicon
sequencing it is avoided), and because (1) the EM-seq ligation adaptor is specifically designed
for the EM-seq prep and (2) EM-seq currently does not support NEB Unique Molecular
Identifiers (which would have to be methylated for compatibility with methylation sequencing
workflows), our goal for bait coverage was a minimum of 20x per 4 kb promoter target region
and 5x for smaller promoter and exonic regions after deduplication.
408
To compare DNA methylation quantification methods in superb starlings, Bismark coverage files
from TEEM-Seq and WGEM-Seq alignments were intersected with 4kb bait "promoter" and
exon ranges (for a separate exon comparison) using Bedtools v2.29.2 (Quinlan and Hall 2010).
To confirm read and CpG site coverage, BAM alignments and Bismark coverage tracks were
visually inspected with Geneious v10.0.4 using an updated chromosomal-level genome assembly
(CU_Lasu_v2; GenBank: GCA_015883425.2) that combined Hi-C data with the previously
published superb starling reference genome (Rubenstein et al. 2021). SAMtools coverage was
used to calculate mean read coverage stats in bait target ranges. To compare WGEM-Seq and
RRBS data, Bedtools was used to generate 200 random 10,000 bp tracks across the starling
genome that were then intersected with putative 2.5 kb promoter tracks, leaving us with 27
random "promoter" regions. This targeted 2.5kb range was chosen instead of a whole genome
comparison because RRBS coverage is dense at restriction sites (e.g., in CG clusters localized
just prior to the transcription start site). Visual inspection confirmed that these random regions
appeared rich in RRBS coverage. Bismark coverage files from WGEM-Seq and RRBS
alignments were likewise intersected with these random regions, and then filtered for unique
sites (in case of promoter overlap) above 5x coverage (Ziller et al. 2015). Intersected coverage
files for each sample were then matched between each method for common CpG sites. That is,
we compared matched sites on a per sample basis within random regions between WGEM-Seq
versus RRBS data (resulting in 20 regions with 5x CpG site coverage for all but one sample), or
within target regions for WGEM-Seq versus RRBS or WGEM-Seq versus TEEM-Seq data.
Thus, WGEM-Seq, TEEM-Seq, and RRBS data were compared at the CpG site-level in target
regions, and WGEM-Seq and RRBS data were compared at the CpG site-level in 20 random
"promoter" regions that were not included in the bait set. We chose this approach of an
additional random 2.5 kb "promoter" region comparison because we expected that some RRBS
regions corresponding to target ranges might have sparse coverage in comparison to the high
coverage bait alignments. Finally, Bedtools map was used to calculate the mean of Bismark
methylation percentages for these regions.
436
Statistical analysis
DNA methylation values target regions for libraries run in duplicate (or triplicate) were
compared using paired t-tests. Pearson correlations were used to compare site-level CpG data for
TEEM-Seq and WGEM-Seq libraries. Data for each pairwise comparison of methods of
quantifying DNA methylation (TEEM-Seq versus WGEM-Seq, TEEM-Seq versus RRBS, and
WGEM-Seq versus RRBS) were analyzed using generalized linearized mixed models
(GLMMS), with DNA methylation as the dependent variable, quantification method as the
independent variable, and bird ID included as a random effect to account for repeated measures
(multiple gene regions per library). All statistical tests were done in JMP Pro v16.2 or R v4.1.3
(R Core Team 2022).

Data Availability
448
449   Raw genetic data are available at the NCBI Sequence Read Archive (SRA) (BioProject:
450   PRJNA861281). Summarized data are provided in the Supplemental Information in table format.
451
452   **Computer Code**
453
454   Code is available at https://doi.org/10.6084/m9.figshare.20634759.
455
456   **Author Contributions**
457
458   D.R.R. conceived of the project. D.R.R. and J.S. designed the experiment. D.R.R. collected the
459   samples. J.S. collected and analyzed the data. D.R.R. and J.S. wrote the paper.
460
461   **Competing Interests**
462
463   The authors declare no competing interests.
464
465   **Acknowledgements**
466

We thank W. Watetu, G. Manyaas, and J. Mosiany for assistance collecting superb starlings and
the Mpala Research Centre for logistical support in Kenya. We acknowledge A. Devault and Z.
Hanf at Daicel Arbor Biosciences for help designing the bait sets. We acknowledge the Sackler
Institute for Comparative Genomics at the American Museum of Natural History and the
Department of Biological Sciences at Columbia University for access to equipment. We thank S.
Siller for helping to organize gene sequences and, along with B. Heidinger, for supplying house
sparrow DNA. A. Bendesky and J. Merritt provided constructive feedback on previous versions
of this manuscript. This work was supported by the U.S. National Science Foundation grants
IOS-1257530 and IOS-1656098.

References
Bird, A. P. CpG-rich islands and the function of DNA methylation. *Nature* **321**, 209-213 (1986).
Bock, C., Tomazou, E. M, Brinkman, A. B., Müller, F., Simmer, F., Gu, H. et al. Quantitative
comparison of genome-wide DNA methylation mapping technologies. *Nature*
Biotechnology **10**, 1106-1116 (2010).
Bossdorf, O., Richards, C. L. & Pigliucci, M. Epigenetics for ecologists. *Ecology Letters* **11**,
106-115 (2008).
Cavalli, G. & Heard, E. Advances in epigenetics link genetics to the environment and disease.
Nature **571**, 489-499 (2019).
Consortium, B. Quantitative comparison of DNA methylation assays for biomarker development
and clinical applications. *Nature Biotechnology* **34**, 726-37 (2016).
Crary-Dooley, F. K., Tam, M. E., Dunaway, K. W., Hertz-Picciotto, I., Schmidt, R. J. & LaSalle
J. M. Comparison of existing global DNA methylation assays to low-coverage whole-
genome bisulfite sequencing for epidemiological studies. *Epigenetics* **12**, 206–214
(2017).
Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O. et al. Twelve years
of SAMtools and BCFtools. *GigaScience* **10**, 1-4 (2021).
Feng, S., Zhong, Z., Wang, M. & Jacobsen, S. E. Efficient and accurate determination
of genome-wide DNA methylation patterns in *Arabidopsis thaliana* with enzymatic
methyl sequencing. *Epigenetics & Chromatin* **13**, 42 (2020).
Gouil, Q. & Keniry, A. Latest techniques to study DNA methylation. *Essays in Biochemistry* **63**,
639–48 (2019).
Gu, H., Smith, Z. D., Bock, C., Boyle, P., Gnirke, A. & Meissner, A.Preparation of reduced
representation bisulfite sequencing libraries for genome-scale DNA methylation
profiling. *Nature Protocols* **6**, 468-481 (2011).
Han, Y., Zheleznyakova, G. Y., Marincevic-Zuniga, Y., Kakhki, M. P., Raine, A., Needhamsen,
M. & Jagodic, M. 2021. Comparison of EM-seq and PBAT methylome library methods
for low-input DNA. *Epigenetics* https://doi.org/10.1080/15592294.2021.1997406 (2021).
Harris, R. A., Wang, T., Coarfa, C., Nagarajan, R. P., Hong, C., Downey, S. L. et al. Comparison
of sequencing-based methods to profile DNA methylation and identification of
monoallelic epigenetic modifications. *Nature Biotechnology* **28**, 1097-1105 (2010).
Husby, A. Wild epigenetics: insights from epigenetic studies on natural populations. *Proceedings*
of the Royal Society B **289**, 20211633 (2022).
Jablonka, E. & Lamb, M. J. The changing concept of epigenetics. *Annals of the New York*
Academy of Sciences **981**, 82-96 (2006).
Kozarewa, I., Armisen, J., Gardner, A. F., Slatko, B. E. & Hendrickson, C. L. Overview of target
enrichment strategies. *Current Protocols in Molecular Biology* **112**, 7.21.1-7.21.23
(2015).
Krueger, F., James, F., Ewels, P., Afyounian, E. & Schuster-Boeckler B. TrimGalore v0.4.2.
https://github.com/FelixKrueger/TrimGalore (2021).
Krueger, F. & Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-
Seq applications. *Bioinformatics* **27**, 1571–1572 (2011).
Lee, E.-J., Pei, L., Srivastava, G., Joshi, T., Kushwaha, G., Choi J.-H. et al. Targeted bisulfite
sequencing by solution hybrid selection and massively parallel sequencing. *Nucleic Acids*
Research **39**, e127–e127 (2011).

Li, Q., Suzuki, M., Wendt, J., Patterson, N., Eichten, S. R., Hermanson, P. J. et al. 2015. Post-
conversion targeted capture of modified cytosines in mammalian and plant genomes.
Nucleic Acids Research **43**, e81 (2015).
Liu, Y., Hu, Z., Cheng, J., Siejka-Zielinska, P., Chen, J., Inoue, M., Ahmed, A. A. & Song, C.
Subtraction-free and bisulfite-free specific sequencing of 5-methylcytosine and its
oxidized derivatives at base resolution. *Nature Communications* **12**, 618 (2021).
Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**, 10-12 (2011).
Masser, D. R., Stanford, D. R., Hadad, N., Giles, C. B., Wren, J. D., Sonntag, W. E. et al.
Bisulfite oligonucleotide-capture sequencing for targeted base- and strand-specific
absolute 5-methylcytosine quantitation. *AGE* **38**, 49 (2016).
Morrison, J., Koeman, J. M., Johnson, B. K., Foy, K. K., Beddows, I., Zhou, W. et al. Evaluation
of whole-genome DNA methylation sequencing library preparation protocols.
Epigenetics & Chromatin **14**, 28 (2021).
Quinlan, A. R. & Hall I. M. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).
R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria; https://www.R-project.org/ (2022).
Rey, O., Eizaguirre, C., Angers, B., Baltazar-Soares, M., Sagonas, K., Prunier, J. G. et al.
Linking epigenetics and biological conservation: Towards a conservation epigenetics
perspective. *Functional Ecology* **34**, 414-427 (2020).
Robertson, K. D. & Wolffe, A. P. DNA methylation in health and disease. *Nature Reviews*
Genetics **1**, 11-19 (2000).
Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G. &
Mesirov, J. P. Integrative Genomics Viewer. *Nature Biotechnology* **29**, 24–26 (2011).
Rubenstein, D. R., Corvelo, A., MacManes, M. D., Maia, R., Narzisi, G., Rousaki, A. et al.
Feather gene expression elucidates the developmental basis of iridescence in African
starlings. *Journal of Heredity* **112**, 417-429 (2021).
Rubi, T. L., Knowles, L. L. & Dantzer, B. Museum epigenomics: characterizing cytosine
methylation in historic museum specimens. *Molecular Ecology Resources* 20, 1161-1170
(2020).
Sapolsky, R. M., Romero, L. M. & Munck, A. U. How do glucocorticoids influence stress
responses? Integrating permissive, suppressive, stimulatory, and preparative actions.
Endocrine Reviews **21**, 55-89 (2000).
Seutin, G., White, B. N. & Boag, P. T. Preservation of avian blood and tissue samples for DNA
analysis. *Canadian Journal of* Zoology **69**, 82–90 (1991).
Siller, S. J. Epigenetic modification of the hypothalamic-pituitary-adrenal axis during early life
of the house sparrow (*Passer domesticus*). Ph.D. Thesis, Columbia University (2022).
Tanaka, K. & Okamato, A. Degradation of DNA by bisulfite treatment. *Bioorganic & Medicinal*
Chemistry Letters **17**, 1912-1915 (2007).
Vaisvila, R., Chaithanya Ponnaluri, V. K., Sin, Z., Langhorst, B. W., Saleh, L., Guan, S. et al.
Enzymatic methyl sequencing detects DNA methylation at single-base resolution from
picograms of DNA. *Genome Research* **31**, 1280-1289 (2021).
Waddington, C. H. The epigenotype. *Endeavor* **81**, 18–20 (1942).

Ziller, M. J., Hanson, K. D., Meissner, A. & Aryee, M. J. Coverage recommendations for
methylation analysis by whole-genome bisulfite sequencing. *Nature Methods* **12**, 230-
232 (2015).

Figure 1. Comparison of mean target region coverage (after deduplication) for target-enriched enzymatic methyl sequencing (TEEM-Seq) and whole-genome enzymatic methyl sequencing (WGEM-Seq) in (a-c) putative promoter and (d-f) exonic regions. Libraries for three of the same individual superb starlings (**a** and **d**) BB-17168, (**b** and **e**) BB-17501, and (**c** and **f**) BB-14232 were sequenced using TEEM-Seq (red bars) and WGEM-Seq (blue bars).

575
576
**Figure 2. Representative examples of target-enriched enzymatic methyl sequencing
(TEEM-Seq) versus whole-genome enzymatic methyl sequencing (WGEM-Seq) coverage**.
Integrative Genomics Viewer (IGV) logarithm-scale coverage and read alignment plots (for
deduplicated data) of superb starling sample BB-17168 multi-exonic targets for (**a**) FK506
binding protein 5 (*FKBP5*), (**b**) the glucocorticoid receptor (*NR3C1*), and (**c**) the androgen
receptor (*AR*). *FKBP5* includes the entire promoter and gene region, whereas *NR3C1* and *AR* are
focused on the end of each gene region to enable IGV's display of read alignments. Alignments
are colored in bisulfite mode by CG, with red indicating methylated sites (non-converted
cytosines) and blue indicating unmethylated sites (converted cytosines). Target tracks for the
superb starling genome assembly (CU_Lasu_v2) include GenBank sequence ID, locus tag, and
protein IDs (with exonic spans also in blue).

Figure 3. CpG site-level Pearson correlations between target-enriched enzymatic methyl sequencing (TEEM-Seq) and whole-genome enzymatic methyl sequencing (WGEM-Seq) libraries from the same individual superb starlings. (a) BB-17168 (R = 0.95, N = 2733, P < 0.0001), (b) BB-17501 (R = 0.95, N = 2770, P < 0.0001), and (c) BB-14232 (R = 0.93, N = 1588, P < 0.0001). CpG sites at 5x coverage or above were used for comparisons.

595
596
Figure 4. Pairwise comparisons of three methods for quantifying DNA methylation: target-enriched enzymatic methyl sequencing (TEEM-Seq), whole-genome enzymatic methyl sequencing (WGEM-Seq), and reduced-representation bisulfite sequencing (RRBS). (**a** and **d**) Libraries for three of the same individual superb starlings were sequenced using both TEEM-Seq and WGEM-Seq. (**b** and **e**) Libraries for seven of the same individual superb starlings were sequenced using both TEEM-Seq and RRBS. (**c** and **f**) Libraries for three of the same individual superb starlings were sequenced using both WGEM-Seq and RRBS. WGEM-Seq and RRBS data were compared in 20 random 2.5 kb "promoter" regions that were not included in the bait set, as we expected that some RRBS regions corresponding to target ranges might have sparse coverage in comparison to the high coverage bait alignments. In all cases, there were no differences in DNA methylation among quantification methods, as the absolute value mean differences in DNA methylation for promoter regions and gene bodies or random 2.5 kb "promoter" regions did not differ among pairwise comparisons (**a-c**) In each plot, grey violins show the distribution of CpG site absolute value mean differences in DNA methylation for (**a** and **b**) promoter regions and gene bodies or (**c**) random 2.5 kb "promoter" regions, black lines depict the median values, and white lines show the 1st and 3rd quartiles. Additionally, (**d** and **e**) mean DNA methylation levels for each target or (**f**) random region did not significantly differ. Target regions are indicated by different colors according to the legends on panels d-f. See Tables S2-S5 for more detailed information.

Figure 5. Diagram comparing the workflow for target-enriched enzymatic methyl
sequencing (TEEM-Seq), whole-genome enzymatic methyl sequencing (WGEM-Seq), and
reduced-representation bisulfite sequencing (RRBS).

Table 1. Targeted genomic regions for superb starling baits. Although we targeted putative
promoter regions of all of the genes (i.e., 4 kb upstream of the transcription start site where
annotated), we also targeted complete exonic regions (indicated by *) or gene body regions
(based on similarity with Ensembl genome sequences, including the collared flycatcher for
GNRH1 and zebra finch for *MC4R*, indicated by **) in a subset of the genes.
626

Target region	Gene name
*AR**	Androgen receptor
*AVPR1A**	Arginine vasopressin receptor 1A
*AVPR1B**	Arginine vasopressin receptor 1B
*CRH**	Corticotropin releasing hormone
DNMT1	DNA methyltransferase 1
DNMT3a	DNA methyltransferase 3a
DNMT3b	DNA methyltransferase 3b
*EGR1**	Early growth response 1
ESR1	Estrogen receptor 1
*FKBP5**	FK506 binding protein 5
GNIH	Gonadotropin-inhibitory hormone
*GNRH1***	Gonadotropin releasing hormone 1
*GNRHR2**	Gonadotropin-releasing hormone receptor
MC2R	Melanocortin receptor 2
*MC4R***	Melanocortin receptor 4
*NR3C1**	Glucocorticoid receptor
*NR3C2**	Mineralcorticoid receptor
*OXTR**	Oxytocin receptor
*POMC**	Proopiomelanocortin
SERPINA1	Serpin family A member 1
VT	Vasotocin
VTG1	Vitellogenin-1

627