Structures of LIG1 uncover a lack of sugar discrimination against a ribonucleotide

at the 3'-end of nick DNA

Qun Tang[1], Mitchell Gulkis[1], and Melike Çağlayan[1,*]

[1]Department of Biochemistry and Molecular Biology, University of Florida, Gainesville, FL

32610, USA

*To whom correspondence should be addressed. Tel.: +1 352-294-8383; E-mail:

caglayanm@ufl.edu

Ribonucleotides can be incorporated by DNA polymerases and the subsequent joining of

3'-OH and 5'-P ends in the phosphodiester backbone at the nick by DNA ligase during

DNA replication and repair is critical for maintaining genome stability. Although it has

been extensively studied for DNA polymerases across families, the sugar discrimination

mechanism of a human DNA ligase at atomic resolution is entirely missing. Here, for the

first time, we determine X-ray structure of DNA ligase I (LIG1) in complex with nick

DNA containing rG:C at the 3'-end and capture the ligase at the final phosphodiester

bond formation step of the ligation reaction involving an adenylate (AMP) release.

Moreover, we show mutagenic end joining of the nick DNA substrate with preinserted 3'-

rG:C by LIG1 *in vitro*. Our findings reveal an important source of ribonucleotides

embedded in genomic DNA, which could come from the failure of LIG1 to discriminate

against a ribonucleotide at the 3'-end during nick sealing step of DNA replication and

repair.

Keywords: Ribonucleotide, rNTP, DNA polymerase, DNA ligase I, DNA ligase IIIα/XRCC1

complex, mutagenic ligation, DNA repair, DNA replication, genome stability

Introduction

The concentrations of ribonucleotide triphosphates (rNTPs) are much more abundant than

those of deoxyribonucleotide triphosphates (dNTPs) within a cell, and therefore, their

misincorporation into genomic DNA by DNA polymerases occur at higher frequencies than

mismatch nucleotide insertions (1-4). It has been estimated that >10,000 or >1,000,000

ribonucleotides can be incorporated into yeast and mouse genomes, respectively, making

genomic ribonucleotides the most prevalent source of cellular DNA damage (5-13).

Ribonucleotides embedded into genomic DNA can confound structural and chemical integrity

of duplex DNA, increase its susceptibility to endogenous or exogenous damage, and lead to

several types of genome instability such as replication blockage, mutagenesis, aberrant

recombination, protein-DNA crosslinks, double-strand breaks (DSBs), and chromosome

alterations if not repaired by ribonucleotide excision repair (RER) (14-20).

In human cells, ribonucleotide incorporation can occur during almost all cellular DNA

transactions such as DNA replication by replicative polymerases (pols α, δ, and ε), non-

homologous end joining (NHEJ) of DSB repair by pol μ and base excision repair (BER) by

pols β and λ, translesion synthesis (TLS) by pol ι, and mitochondrial repair by pol γ (6-11,21-

26). Due to the high cellular rNTP/dNTP imbalance, discrimination against ribonucleotide

incorporation by these DNA polymerases is essential for the maintenance of genomic integrity

(1-4). It has been extensively shown by structure/function studies that most DNA polymerases

use a side chain named the "steric gate" (*i.e*., B-, and Y- family pols) or a protein backbone

segment in minor groove nucleotide binding pocket (*i.e*, X-family pols) to govern

ribonucleotide exclusion (6-26). Replicative DNA polymerases effectively remove mismatches

through their 3'-5' exonuclease domains, which is found to be relatively weaker at proofreading

ribonucleotides during DNA synthesis (27). The kinetic basis of inefficient rNTP incorporation

by DNA polymerases is mainly governed by weaker binding and slower rate of incorporation

compared to dNTPs (21).

DNA polymerases and DNA ligases have fundamental roles in maintaining genome stability

during the downstream steps of DNA replication and repair (35). Indeed, the efficiency of DNA

ligation by DNA ligase at the final step relies on the accuracy of DNA synthesis that requires

a Watson-Crick base pair between 5'-phosphate (P) and 3'-hydoxyl (OH) termini of the nick

after DNA polymerase incorporates a correct dNTP (36). In our previous studies, we reported

the importance of the coordination between DNA polymerase (pol) β and DNA ligase I (LIG1)

or DNA ligase IIIα/XRCC1 (LIGIIIα/XRCC1) complex at the downstream steps of BER

pathway to maintain accurate repair (37-47). Our biochemical studies have revealed that, in

certain situations, BER responses can lead to mutagenic outcomes and the DNA ligation step

of the BER pathway can be compromised in case of damaged (8-oxodGTP) or mismatched

nucleotide incorporation by polβ (35-52). Furthermore, at atomic resolution, we recently

reported that the LIG1 discriminates against the mutagenic mismatches distinctly depending

on the architecture of 3'-terminus at the nick DNA. Our structures of LIG1/nick DNA

complexes with G:T and A:C mismatches revealed the ligase strategies that favor or deter the

ligation of base substitution errors at the final step of BER pathway (53).

Although the distinct features of the sugar selection mechanism have been extensively reported

for DNA polymerases across families by X-ray crystallography studies (7-34), the mechanism

of ribonucleotide discrimination by human DNA ligases at atomic resolution is entirely

missing. Given the fact of high cellular rNTP/dNTP ratio, the mutagenic ligation of

ribonucleotides by a DNA ligase at the final step following their incorporation by a DNA

polymerase at a non-negligible rate during DNA replication or repair could constitute a major

threat to genome stability. In the present study, we aimed to elucidate the mechanism by which

human LIG1 facilitates or hinders sealing ribonucleotide at the 3'-end of nick DNA. Our

LIG1/nick DNA structure uncovers a lack of a proficient sugar discrimination and

demonstrates that the ligase active site can accommodate 3'-riboG opposite template base C.

For the first-time, we showed a phosphodiester bond formation between 3'-OH and 5'-P ends

of the nick DNA, which refers to the final step of the ligation reaction involving an adenylate

(AMP) release. Moreover, we demonstrated the mutagenic ligation of nick DNA substrate

containing 3'-preinserted rG:C by LIG1 and LIGIIIα/XRCC1 complex in the ligation assays *in*

vitro. Overall, our findings provide a novel insight into the characterization of ribonucleotide

selectivity on the downstream events of DNA repair and replication as well as a lack of sugar

discrimination by LIG1, demonstrating its distinctive ability to seal nick DNA with 3'-rG:C at

atomic resolution.

Results

Structures of LIG1/rG:C demonstrate a failure to discriminate against a ribonucleotide

at the 3'-end of nick DNA

To elucidate the sugar discrimination mechanism of LIG1 at atomic resolution, we determined

the structure of LIG1 in complex with nick DNA containing rG:C at the 3'-end (Table 1). Our

LIG1/3'-rG:C structure shows density for the 2'-hydroxyl (2'-OH) group on the ribose ring and

demonstrates the formation of a phosphodiester bond between adjacent 3'-OH and 5'-P ends of

DNA termini at the nick (Figure 1). This LIG1 structure represents the first structure of human

DNA ligase that engages with a ribonucleotide-containing nick DNA during the final step of

the ligation reaction where 3'-OH terminus of the nick attacks the 5'-P terminus downstream of

the nick to form a phosphodiester bond that is coupled to a release of adenylate (AMP).

In our previously solved LIG1/A:T structure, we reported that the active site can accommodate

a cognate A:T base pair where DNA-AMP intermediate is formed (53). The comparison of the

2Fc-Fo maps at 3'-OH and 5'-P ends of the nick DNA demonstrated continuous and

discontinuous maps in the LIG1 structures of rG:C and A:T, respectively (Figure 2A-B). This

means that both DNA ends are positioned for nick sealing in the LIG1/A:T structure, which

refers to the step 2 of the ligation reaction, and a phosphodiester bond between 3'-OH and 5'-P

ends is formed in the LIG1/3'-rG:C structure. The Fo-Fc map at 3σ (Figure 1B) and the 2Fo-

Fc map at 1σ (Figure 2B) demonstrate that the AMP is incomplete in the LIG1/rG:C and shows

less density than the LIG1/A:T structure. This observation indicates that AMP is still linked to

the 5'-P end to form DNA-AMP intermediate in the LIG1/A:T structure, while it's already

released in the LIG1/3'-rG:C structure, which refers to the step 3 of the ligation reaction. The

overlay of both LIG1 structures shows a superimposed Cα root mean square deviation of 0.398

Å (Figure 2C) and no significant conformational difference in the domain organization of the

LIG1 catalytic core (Supplementary Figure 1).

Our previously solved LIG1/mismatch structures demonstrated that the A:C is positioned at

the initial step (step 1) where the ligase active site stays adenylated (LIG1-AMP) at its active

site lysine residue (K568), and that the LIG1/G:T mismatch is captured during step 2 when an

adenylate is transferred to the 5'-end of the nick (53), while the LIG1/rG:C structure shows a

phosphodiester bond formation coupled to AMP release at step 3 of the ligation reaction

(Supplementary Figure 2). We also compared the map of LIG1/rG:C (EE/AA) with the

previously solved structure of LIG1/C:G (wild-type) (54,55). The overlay of both LIG1

structures demonstrated the 2Fc-Fo discontinuous map at 3'-OH and 5'-P ends of the nick for

LIG1/C:G and a continuous map for LIG1/rG:C (Supplementary Figure 3). We observed a

Watson-Crick conformation for rG:C and sugar pucker analysis of LIG1/3'-rG:C structure

showed that the ribose adopts a C3'-endo conformation, which is similar to that of the

LIG1/A:T structure (Supplementary Figure 4).

Mutagenic ligation of nick DNA containing a ribonucleotide at 3'-end by LIG1

To investigate the sugar discrimination mechanism of LIG1 against a ribonucleotide at the 3'-

end of the nick *in vitro*, we tested the ligation efficiency of wild-type and EE/AA mutant of

LIG1 using the nick DNA substrate with 3'-preinserted rG opposite template base C that could

be formed during DNA replication or repair when DNA polymerase incorporates a rNTP

(Supplementary Scheme 1). In the control reactions, we used the nick DNA substrate with 3'-

dG:C.

Our results demonstrated the mutagenic ligation of nick DNA with 3'-rG:C by wild-type

(Figure 3A, lines 9-14) and EE/AA mutant (Figure 3C, lines 9-14) of LIG1. This was

comparable with the ligation products of nick DNA containing a cognate 3'-dG:C (Figure 3A

and 3C, lines 2-7). We observed a similar amount of ligation products for both DNA substrates

(3'-rG:C *versus* 3'-dG:C) with wild-type (Figure 3B) and the low fidelity mutant of the ligase

(Figure 3D). Similarly, our results demonstrated no significant difference between the

mutagenic ligation efficiencies of both LIG1 proteins (Supplementary Figures 5-6).

In our structure studies, we use the EE/AA mutant of LIG1 carrying Glu(E)346 and Glu(E)592

mutations at the Mg^{2+}-dependent high-fidelity site to crystallize the ligase in the conditions

lacking Mg^{2+} (53,55). In the reaction conditions that mimic the LIG1 crystals, we also observed

efficient nick sealing of the DNA substrate with 3'-rG:C in the absence of Mg^{2+} for longer time

points of incubation (Supplementary Figure 7).

In our previous studies (38,42,45,53), we reported that LIG1 can ligate nick DNA containing

a Watson-Crick like mismatch (G:T) and oxidative damage (8-oxoG:A). We next compared

the mutagenic ligation efficiency of LIG1 for the nick DNA substrates containing a

mismatched, damaged, and ribonucleotide-containing ends (Supplementary Figure 8). Our

results showed that the ligation of 3'-8-oxoG:A and 3'-G:T yields nick sealing products along

with the accumulation of DNA-AMP intermediates (Supplementary Figure 8A, lines 2-7 and

9-14, respectively), while there was only mutagenic ligation products in the presence of the

nick DNA with 3'-rG:C (Supplementary Figure 8A, lines 16-21). These results demonstrate

more efficient nick sealing of a ribonucleotide than a mismatched or damaged base at the 3'-

end, which could explain why the ligase active site engages with those DNA ends at different

steps of the ligation reaction (Supplementary Figure 8B).

Role of LIG1 active site residues for sugar discrimination against 3'-rG:C at the nick

DNA

We previously reported that the LIG1 active site shows distinct DNA conformations depending

on the identity of the 3'-terminus (cognate *versus* mismatched) (53). The overlay of LIG1

structures in complex with nick DNA containing 3'-rG:C and 3'-A:T showed no conformational

change at the positions of the LIG1 active site residues F635, R871, and F872, which reside in

the close proximity to the 2'-OH of the ribose (Figure 4A). Furthermore, we did not observe a

shift in the position of F872 at the ligase active site engaging with rG:C as we reported

previously in the LIG1/A:C structure (Figure 4B).

To further investigate this observation, we tested the importance of the ligase active site residue

F872 for the end joining efficiency of the nick DNA substrate with 3'-rG:C by LIG1 in the

ligation assays (Supplementary Scheme 1). For this purpose, we used the LIG1 mutants

carrying a F872A mutation in the wild-type (F872A) and low fidelity mutant (EE/AA)

background. Our results demonstrated that LIG1 F872A (Figure 5A, lines 2-7) and LIG1

EE/AA/F872A (Figure 5A, lines 8-13) can also efficiently seal nick DNA with 3'-rG:C. There

were slightly less ligation products at the initial time points of the reaction (Figure 5B and

Supplementary Figure 9) in comparison with the wild-type and EE/AA mutant of LIG1 (Figure

3). These findings demonstrate that the active site residue F872 that is located upstream of the

nick and positioned close to the deoxyribose moiety of the nucleotide at the 5'-end has no role

in sugar discrimination by LIG1.

Discussion

Ribonucleotides (rNTPs) are found at a concentration several orders of magnitude higher than

dNTPs in most cells (1-3). It has been shown that the eukaryotic replicative polymerases insert

around 10,000 rNTPs per round of DNA replication and repair polymerases are also capable

of inserting rNTPs during DNA synthesis with some being better at discriminating against

rNTP insertion than others (6,14). Therefore, rNMPs are the most common type of DNA lesion

found in the genomic DNA with critical effects on overall genome stability (27). Although

DNA polymerases have evolved selectivity against rNTPs favoring dNTP insertion and have

an intrinsic exonuclease proofreading function to remove rNTPs, this is not sufficient to

prevent large amounts of rNTPs remaining in the DNA backbone (6-26). The RER pathway,

involving RNase H2 function, can efficiently remove ribonucleotides from RNA:DNA hybrids

by cleaving at the 5'-end of the ribonucleotide, which is followed by strand displacement

synthesis by pol δ and flap removal by Flap Endonuclease 1 (FEN1) and/or Dna2, and finally

ligation step by LIG1 (56-58).

In addition to its role in the RER pathway, LIG1 is responsible for the ligation of over 10

million Okazaki fragments during each round of DNA replication and nick sealing of the repair

products at the final step of all DNA repair pathways which process DNA damage that could

be formed due to a variety of endogenous and exogenous sources (59-61). The nick repair

products are indeed potentially toxic intermediates, and this may lead to DNA strand breaks

that could trigger harmful nuclease activities, cytotoxicity, double strand breaks in replicating

DNA, and ultimately genomic instability (61-63). Therefore, DNA ligases could be challenged

by various genomic insults such as oxidative damage, non-canonical mismatches, and

ribonucleotides embedded into genomic DNA due to their incorporations by DNA polymerases

on the downstream steps of DNA repair and replication (35,36). DNA ligases have evolved a

tolerance for ribonucleotides at 3'-end while heavily discriminating against a ribonucleotide at

5'-end (54,59-63). On the other hand, the mechanism of sugar discrimination during nick

sealing reaction by DNA ligase at atomic resolution remains unknown. ATP-dependent DNA

ligases catalyze the ligation reaction including three sequential steps (64-69): (i) enzyme

adenylation at the active site lysine residue (*i.e,* K568) and formation of the ligase-AMP

intermediate, (ii) transfer of AMP group to 5'-P end of nick DNA resulting in the formation of

DNA-AMP intermediate, and finally (iii) phosphodiester bond formation between adjacent 3'-

OH and 5'-P DNA termini at a nick.

Our previous and present LIG1 structures revealed the features of DNA substrate and LIG1

interaction that dictate accurate *versus* mutagenic outcomes (53). Our structures contribute to

the understanding of the mechanism by which LIG1 discriminates against non-canonical DNA

intermediates during nick sealing at each individual step of the ligation reaction as well as how

the ligase active site engages with nick DNA distinctly depending on the architecture of

cognate (step 2), mismatched (step 1 or 2), or ribonucleotide (step 3)-containing ends

(Supplementary Table 1). Overall LIG1 structures in complex with nick DNA substrates have

provided considerable mechanistic details about steps 1 and 2 of the ligation reaction while

relatively less information has been reported for step 3 (53-55,70,71). In addition to our present

study of the LIG1/rG:C structure showing the human ligase active site at step 3 in the crystal

conditions lacking Mg^{2+} using EDTA pre-treated LIG1 protein, another step 3 structure has

been shown in ATP-dependent ligase from *Prochlorococcus marinus* (Pmar-Lig) (72). In this

structure, Pmar-Lig (step 2) crystals were immersed in cryoprotectant buffer containing $MgCl_2$

and then frozen immediately to get the step 3 Pmar-Lig/DNA structure, which is poised for

phosphodiester bond formation with partially occupied AMP in the active site. Since it is well

known that Mg^{2+} is required for step 3 of the ligation reaction, it is at first puzzling why our

LIG1/rG:C without Mg^{2+} shows step 3 (73). However, this is likely because of the 3'-endo

sugar pucker that ribonucleotides adopt. In B-form DNA, the sugar moieties usually adopt the

2'-endo sugar pucker conformation, while in A-form DNA or RNA duplexes the sugar moieties

usually adopt the 3'-endo sugar pucker conformation (74). The ligase active site forces the

DNA upstream of the nick to adopt an A-form conformation which causes the sugar at the 3'-

end to assume the 3'-endo conformation, both of which are necessary for ligation (54). However,

Mg^{2+} plays an important role in this step. In previous studies, it has been shown that the

catalytic Mg^{2+} can activate the 3'-OH for nucleophilic attack on the 5'-P (73). However, this

catalytic Mg^{2+} plays a dual role where the interaction between Mg^{2+} and the 3'-OH also

stabilizes the 3'-endo conformation of the 3'-end. When we compare the Mg^{2+}-containing

structure of LIG1 wild-type/3'-ddC:G (dideoxycytidine), which lacks a 3'-OH group, as well

as the structure of Pmar-Lig/3'-dC:G with non-active Mn^{2+}, the divalent metal atom is in almost

the same position, which demonstrates that the position of the divalent metal is conserved for

the ligases. If we imagine the position of the 3'-OH in the LIG1/3'-ddC:G structure, it is also in

the same position as Pmar-Lig/3'-dC:G; additionally, the divalent metal interacts with both 3'-

OH and 5'-P (Supplementary Figure 10). Since rG at the 3'-end adopts a stable 3'-endo

conformation due to its identity as a ribonucleotide, it does not require the rigidity enforced by

the catalytic Mg^{2+} to properly align the 3'-OH for catalysis. We assume that this is the reason

LIG1/rG:C shows step 3 of ligation reaction even without Mg^{2+}. Indeed, this is further

supported biochemically by the ligation of nick DNA substrate with 3'-rG:C in the absence of

Mg^{2+} (Supplementary Figure 7).

To understand the mechanism of sugar discrimination by human DNA ligases more

comprehensively, further structural and biochemical studies are required with other repair

ligases such as DNA ligase IIIα and DSB repair ligase IV (75-81). We previously reported that

the BER DNA ligases, LIG1 and DNA ligase IIIα/XRCC1 complex, are compromised by

subtle changes in all 12 possible mismatched base pairs at the 3'-end of the nick repair

intermediate (42). In the present study, our ligation assays with DNA ligase IIIα/XRCC1

complex also demonstrated mutagenic nick sealing of DNA substrate with 3'-rG:C as efficient

as a cognate 3'-dG:C ligation (Supplementary Figure 11), and we did not observe any

significant difference between the mutagenic ligation efficiencies of LIG1 and DNA ligase

IIIα/XRCC1 complex (Supplementary Figure 6). Although these findings indicate a lack of

sugar discrimination against a ribonucleotide by LIG1 for 3'-rG:C nick DNA only, further

structure/function characterizations of LIG1 against the nick DNA intermediates containing all

possible ribonucleotides mismatches at either 3'- or 5'-end at a nick are necessary to provide a

comprehensive insight into how LIG1 bestows substrate specificity during DNA replication and repair, which is important to understand the disease mechanisms and to develop potential therapeutic targets and strategies (82-84).

Methods

Preparation of nick DNA substrates

DNA substrates with a 6-carboxyfluorescein (FAM) label were prepared as described previously (35-53). Nick DNA substrates containing 3'-preinserted dG:C, dG:T, 8oxodG:A, and rG:C were used in the ligation assays including DNA ligase I (wild-type, and F872A, EE/AA, and EE/AA/F872A mutants) or DNA ligase IIIα/XRCC1 complex (Supplementary Table 2). Nick DNA substrates for LIG1 X-ray crystallography were prepared by annealing upstream, downstream, and template primers (Supplementary Table 3).

Protein purifications

His-tag (pET-24b) human C-terminal (Δ261) wild-type and E346A/E592A (EE/AA) mutant of LIG1 proteins were purified as described previously (35-53). Briefly, they were overexpressed in Rosetta (DE3) *E. coli* cells in Terrific Broth (TB) media with kanamycin (50 μgml^{-1}) and chloramphenicol (34 μgml^{-1}) at 37 °C. The cells were induced with 0.5 mM isopropyl β-D-thiogalactoside (IPTG) when the OD_{600} was reached to 1.0, and the overexpression was continued for overnight at 28 °C. Cells were lysed in the lysis buffer [50 mM Tris-HCl (pH 7.0), 500 mM NaCl, 20 mM imidazole, 10% glycerol, 1 mM PMSF, an EDTA-free protease inhibitor cocktail tablet] by sonication at 4 °C and the cell lysate was pelleted at 31,000 x g for 1 h at 4 °C. Proteins were purified by a HisTrap HP column with an increasing imidazole concentration (20-300 mM) after being equilibrated in the binding buffer [50 mM Tris-HCl (pH 7.0), 500 mM NaCl, 20 mM imidazole, 10% glycerol] at 4 °C. The collected fractions were subsequently loaded onto a HiTrap Heparin column that was equilibrated with the binding buffer [50 mM Tris-HCl (pH 7.0), 50 mM NaCl, 1.0 mM EDTA,

and 10% glycerol], and then eluted with a linear gradient of NaCl up to 1 M. LIG1 proteins

were further purified by Superdex 200 10/300 columns in the buffer [20 mM Tris-HCl (pH

7.0), 200 mM NaCl, and 1 mM DTT]. The plasmid DNA constructs encoding human C-

terminal (△261) LIG1 genes carrying a single F872A and triple EE/AA/F872A mutations were

generated by side-directed mutagenesis using the plasmid DNA of LIG1 wild-type and

E346A/E592A (EE/AA) mutant, respectively, as previously described (35-53). His-tag LIG1

mutant proteins F872A and EE/AA/F872A were overexpressed and purified as described

above. Human his-tag DNA ligase IIIα/XRCC1 complex (pDUET1) was purified as previously

described (35-53). Briefly, the complex protein was overexpressed in BL21(DE3) *E. coli* cells

in LB media at 37 °C and the cell lysis was prepared as described above. The supernatant was

loaded onto a HisTrap HP column (GE Health Sciences) and purified with an increasing

imidazole gradient (20-300 mM) elution at 4 °C. The collected fractions were subsequently

loaded onto a HiTrap Heparin column that was equilibrated with the binding buffer [25 mM

Tris-HCl (pH 7.5), 50 mM NaCl, and 10% glycerol], and then eluted with a linear gradient of

NaCl up to 1 M. The protein complex was further purified by Superdex 200 Increase 10/300

chromatography (GE Healthcare) in the buffer [25 mM Tris-HCl (pH 7.0), 200 mM NaCl,

glycerol 5%, and 1 mM DTT]. All proteins purified in this study were dialyzed against storage

buffer including 25 mM Tris-HCl (pH 7.0), 200 mM NaCl, concentrated, frozen in liquid

nitrogen, and stored at -80 °C.

Crystallization and structure determination

LIG1 C-terminal (△261) EE/AA mutant that was pre-treated with EDTA before using in the

crystallization experiments in the absence of Mg^{2+} as described previously (53). All LIG1-nick

DNA complex crystals were grown at 20 °C using the hanging drop method. LIG1 (at 27 mgml^{-}

^{1})/DNA complex solution was prepared in 20 mM Tris-HCl (pH 7.0), 200 mM NaCl, 1 mM

DTT, 1 mM EDTA and 1 mM ATP at 1:4:1 DNA:protein molar ratio and then mixed with 1

μl reservoir solution containing 100 mM MES (pH 6.1), 100 mM lithium acetate, and 14%

(w/v) polyethylene glycol PEG3350. All crystals grew in one day. They were then washed in

the reservoir solution with 20% glycerol and flash cooled in liquid nitrogen for data collection.

The crystals were maintained at 100° K during X-ray diffraction data collection using the

beamline 7B2 at Cornell High Energy Synchrotron Source (CHESS). The diffraction images

were indexed and integrated using HKL2000. All structures were solved by the molecular

replacement method using PHASER with PDB entry 7SUM as a search model (53,85,86).

Iterative rounds of model building were performed in COOT and the final models were refined

with PHENIX or REFMAC5 (86-89). 3DNA program was used for sugar pucker analysis (90).

All structural images were drawn using PyMOL (The PyMOL Molecular Graphics System,

V0.99, Schrödinger, LLC). Detailed crystallographic statistics are provided in Table 1.

DNA ligation assays

The nick sealing efficiency of DNA ligase (DNA ligase I or DNA ligase IIIα/XRCC1 complex)

was investigated to determine the sugar discrimination mechanism against 3'-preinserted rG

opposite template base C in the ligation assays (Supplementary Scheme 1) as described

previously (35-53). Briefly, the reaction mixture containing 50 mM Tris-HCl (pH 7.5), 100

mM KCl, 10 mM $MgCl_2$, 1 mM ATP, 1 mM DTT, 100 μgml^{-1} BSA, 10% glycerol, and 500

nM nick DNA substrate was initiated by the addition of 100 nM DNA ligase, incubated at 37

°C, and stopped at the time points indicated in the figure legends. The reaction products were

then quenched with an equal amount of gel loading buffer [95% formamide, 20 mM

ethylenediaminetetraacetic acid, 0.02% bromophenol blue and 0.02% xylene cyanol]. After

incubation at 95 °C for 3 min, the reaction products were separated by electrophoresis on an

18% denaturing polyacrylamide gel. The gels were scanned with a Typhoon Phosphor Imager

(Amersham Typhoon RGB), and the data were analyzed using ImageQuant software. The

ligation assays were performed similarly with nick DNA substrates containing 3'-dG:C, 3'-

8oxodG:A, or 3'-dG:T. In addition, nick sealing of DNA with 3'-rG:C by EE/AA mutant of

LIG1 in the absence of Mg^{2+} was performed as described above in the reaction buffer lacking

10 mM $MgCl_2$.

Data availability

330 Atomic coordinates and structure factors for the reported crystal structure of LIG1 in complex

331 with nick DNA/3'-riboG:C has been deposited in the RCSB Protein Data Bank under accession

332 number 8DVT. All relevant data are available from the authors upon reasonable request.

References

1. Klein, H. L. Genome instabilities arising from ribonucleotides in DNA. *DNA Repair*

56, 26-32 (2017).

2. Potenski, C. J. & Klein, H. L. How the misincorporation of ribonucleotides into

genomic DNA can be both harmful and helpful to cells. *Nucleic Acids Res.* **42**, 10226-

10234 (2014).

3. Cerritelli, S. M. & Crouch, R. J. The balancing act of ribonucleotides in DNA. *Trends*

Biochem. Sci. **41**, 434-445 (2016).

4. Caldecott, K. W. Ribose-an internal threat to DNA. *Science* **343**, 260-261 (2014).

5. Yao, N. Y., Schroeder, J. W., Yurieva, O., Simmons, L. A. & O'Donnell, M. E. Cost

of rNTP/dNTP pool imbalance at the replication fork. *Proc. Natl. Acad. Sci.* **110**,

12942-12947 (2013).

6. Williams, J. S., Lujan, S. A. & Kunkel, T. A. Processing ribonucleotides incorporated

during eukaryotic DNA replication. *Nat. Rev. Mol. Cell Biol.* **17**, 350-363 (2016).

7. Clausen, A. R., Murray, M. S., Passer, A. R., Pedersen, L. C., Kunkel, T. A., Clausen,

A. R., Murray, M. S., Passer, A. R., Pedersen, L. C. & Kunkel, T. A. Structure-function

analysis of ribonucleotide bypass by B family DNA replicases. *Proc. Natl. Acad. Sci.*

U.S.A., **110**, 16802-16807 (2013).

8. Clausen, A. R., Zhang, S., Burgers, P. N., Lee, M. Y. & Kunkel, T. A. Ribonucleotide

incorporation, proofreading and bypass by human DNA polymerase delta. *DNA Repair*

12, 121-127 (2013).

9. Nick McElhinny, S. A., Watts, B. E., Kumar, D., Watt, D. L., Lundstrom, E. B.,

Burgers, P. M., Johansson, E., Chabes, A. & Kunkel, T. A. Abundant ribonucleotide

incorporation into DNA by yeast replicative polymerases. *Proc. Natl. Acad. Sci. U. S.*

A. **107**, 4949-4954 (2010).

10. Gosavi, R. A., Moon, A. F., Kunkel, T. A., Pedersen, L. C. & Bebenek, K. The catalytic

cycle for ribonucleotide incorporation by human DNA pol lambda. *Nucleic Acids Res.*

40, 7518-7527 (2012).

11. Goksenin, A. Y. *et al*. Human DNA polymerase epsilon is able to efficiently extend

from multiple consecutive ribonucleotides. *J. Biol. Chem.* **287**, 42675-42684 (2012).

12. Watt, D. L., Johansson, E., Burgers, P. M., Kunkel, T. A., Watt, D. L., Johansson, E.,

Burgers, P. M. & Kunkel, T. A. Replication of ribonucleotide-containing DNA

templates by yeast replicative polymerases. *DNA Repair*, **10**, 897-902 (2011).

13. Lujan, S. A., Williams, J. S., Clausen, A. R., Clark, A. B. & Kunkel, T. A.

Ribonucleotides are signals for mismatch repair of leading-strand replication errors.

Mol. Cell **50**, 437-443 (2013).

14. Nick McElhinny, S. A., Kumar, D., Clark, A. B., Watt, D. L., Watts, B. E., Lundstrom,

E. B., Johansson, E., Chabes, A. & Kunkel, T. A. Genome instability due to

ribonucleotide incorporation into DNA. *Nat. Chem. Biol*. **6**, 774-781 (2010).

15. Sassa, A., Yasui, M. & Honma, M. Current perspectives on mechanisms of

ribonucleotide incorporation and processing in mammalian DNA. *Genes Environ*. **41**,

3 (2019).

16. DeRose, E. F., Perera, L., Murray, M. S., Kunkel, T. A. & London, R. E. Solution

structure of the Dickerson DNA dodecamer containing a single ribonucleotide.

Biochemistry **51**, 2407-2416 (2012).

17. Jaishree, T. N., van der Marel, G. A., van Boom, J. H. & Wang, A. H Structural

influence of RNA incorporation in DNA: quantitative nuclear magnetic resonance

refinement of d(CG)r(CG)d(CG) and d(CG)r(C)d(TAGCG). *Biochemistry* **32**, 4903-

4911 (1993).

18. Hovatter, K. R. & Martinson, H. G. Ribonucleotide-induced helical alteration in DNA

prevents nucleosome formation. *Proc. Natl. Acad. Sci. U. S. A*. **84**, 1162-1166 (1987).

19. Conover, H. N., Lujan, S. A., Chapman, M. J., Cornelio, D. A., Sharif, R., Williams, J.

S., Clark, A. B., Camilo, F., Kunkel, T. A. & Argueso, J. L. Stimulation of

chromosomal re- arrangements by ribonucleotides. *Genetics* **201**, 951-961 (2015).

20. Huang, S. N., Williams, J. S., Arana, M. E., Kunkel, T. A. & Pommier, Y.

Topoisomerase I-mediated cleavage at unrepaired ribonucleotides generates DNA

double-strand breaks. *EMBO J*. **36**, 361-373 (2016).

21. Joyce, C. M. Choosing the right sugar: how polymerases select a nucleotide substrate.

Proc. Natl. Acad. Sci. U. S. A. **94**, 1619-1622 (1997).

22. Brown, J. A., Fiala, K. A., Fowler, J. D., Sherrer, S. M., Newmister, S. A., Duym, W.

W. & Suo, Z. A novel mechanism of sugar selection utilized by a human X-family

DNA polymerase. *J. Mol. Biol*. **395**, 282-290 (2010).

23. Cavanaugh, N. A., Beard, W. A. & Wilson, S. H. DNA polymerase β ribonucleotide

discrimination. *J. Biol. Chem*. **285**, 24457-24465 (2010).

24. Donigan, K. A., McLenigan, M. P., Yang, W., Goodman, M. F. & Woodgate, R. The

steric gate of DNA polymerase iota regulates ribonucleotide incorporation and

deoxyribonucleotide fidelity. *J. Biol. Chem*. **289**, 9136-9145 (2014).

25. Moon, A. F., Pryor, J. M., Ramsden, D. A., Kunkel, T. A., Bebenek, K. & Pedersen, L.

C. Structural accommodation of ribonucleotide incorporation by the DNA repair

enzyme polymerase μ. *Nucleic Acids Res*. **45**, 9138-9148 (2017).

26. Kasiviswanathan, R. & Copeland, W. C. Ribonucleotide discrimination and reverse

transcription by the human mitochondrial DNA polymerase. *J. Biol. Chem*. **286**, 31490-

31500 (2011).

27. Williams, J. S. & Kunkel, T. A. Ribonucleotides in DNA: Origins, repair and

consequences. *DNA Repair* **19**, 27-37 (2014).

28. Brown, J. A. & Suo, Z. Unlocking the sugar steric gate of DNA polymerases.

Biochemistry **50**, 1135-1142 (2011).

29. DeLucia, A. M., Grindley, N. D. & Joyce, C. M. An error-prone family Y DNA

polymerase (DinB homolog from Sulfolobus solfataricus) uses a 'steric gate' residue

for dis- crimination against ribonucleotides. *Nucleic Acids Res*. **31**, 4129-4137 (2003).

30. Brown, J. A. & Suo, Z. Unlocking the sugar 'steric gate' of DNA polymerases.

Biochemistry **50**, 1135-1142 (2011).

31. Ruiz, J. F., Juarez, R., Garcia-Diaz, M., Terrados, G., Picher, A. J., Gonzalez-Barrera,

S., Fernandez de Hanestrosa, A. R. & Blanco, L. Lack of sugar discrimination by

human pol μ requires a single glycine residue. *Nucleic Acids Res*. **31**, 4441-4449 (2014).

32. Reijns, M. A. M. *et al.* Enzymatic removal of ribonucleotides from DNA is essential

for mammalian genome integrity and development. *Cell* **149**, 23 (2012).

33. Vaisman, A. & Woodgate, R. Ribonucleotide discrimination by translesion synthesis

DNA polymerases. *Crit. Rev. Biochem. Mol. Biol*. **53**, 382-402 (2018).

34. Bonnin, A., Lazaro, J. M.M., Blanco, L. & Salas, M. A single tyrosine prevents

insertion of ribonucleotides in the eukaryotic-type φ29 DNA polymerase. *J. Mol. Biol*.

290, 241-251 (1999).

35. Çağlayan, M. Interplay between DNA polymerases and DNA ligases: Influence on

substrate channeling and the fidelity of DNA ligation. *J. Mol. Biol*. **431**, 2068-2081

(2019).

36. Çağlayan, M. & Wilson, S. H. Oxidant and environmental toxicant-induced effects

compromise DNA ligation during base excision DNA repair. *DNA Repair* **35**, 85-89

(2015).

37. Çağlayan, M., Batra, V. K., Sassa, A., Prasad, R. & Wilson, S. H. Role of polymerase

β in complementing aprataxin deficiency during abasic-site base excision repair. *Nat.*

Struct. Mol. Biol. **21**, 497-499 (2014).

38. Çağlayan, M. *et al.* Oxidized nucleotide insertion by pol β confounds ligation during

base excision repair. *Nat. Commun*. **8**, 14045 (2017).

39. Çağlayan, M. *et al.* Complementation of aprataxin deficiency by base excision repair

enzymes. *Nucleic Acids Res*. **43**, 2271-2281 (2015).

40. Çağlayan, M. *et al.* Complementation of aprataxin deficiency by base excision repair

enzymes in mitochondrial extracts. *Nucleic Acids Res*. **45**, 10079-10088 (2017).

41. Çağlayan, M. & Wilson, S. H. Role of DNA polymerase β oxidized nucleotide insertion

in DNA ligation failure. *J. Radiat. Res*. **58**, 603-607 (2017).

42. Çağlayan, M. The ligation of pol β mismatch insertion products governs the formation

of promutagenic base excision DNA repair intermediates. *Nucleic Acids Res*. **48**, 3708-

3721 (2020).

43. Çağlayan, M. Pol β gap filling, DNA ligation and substrate-product channeling during

base excision repair opposite oxidized 5-methylcytosine modifications. *DNA Repair*

95,102945 (2020).

44. Tang, Q., Kamble, P. & Çağlayan, M. DNA ligase I variants fail in the ligation of

mutagenic repair intermediates with mismatches and oxidative DNA damage.

Mutagenesis **35**, 391-404 (2020).

45. Kamble, P., Hall, K., Chandak, M., Tang, Q. & Çağlayan, M. DNA ligase I fidelity the

mutagenic ligation of pol β oxidized and mismatch nucleotide insertion products in base

excision repair. *J. Biol. Chem*. **296**, 100427 (2021).

46. Tang, Q. & Çağlayan, M. The scaffold protein XRCC1 stabilizes the formation of

polβ/gap DNA and ligase IIIα/nick DNA complexes in base excision repair. *J. Biol.*

Chem. **297**, 101025 (2021).

47. Horton, J. K., Stefanick, D. F., Çağlayan, M., Zhao, M. L., Gassman, N. R. & Wilson,

S.H. XRCC1 phosphorylation affects aprataxin recruitment and DNA deadenylation

activity. *DNA Repair* **64**, 26-33 (2018).

48. Prasad, R., Çağlayan, M., Da-Peng, D., Nadalutti, C. A., Gassman, N. R., Zhao, M.,

Stefanick, D. F., Horton, J. K., Krasich, R., Longley, M. J., Copeland, W. C., Griffith,

J. D. & Wilson, S. H. DNA polymerase β: The missing link of the base excision repair

machinery in mammalian mitochondria. *DNA Repair* **60**, 77-88 (2017).

49. Sassa, A., Çağlayan, M., Rodriguez, Y., Beard, W. A., Wilson, S. H., Nohmi, T.,

Honma, M. & Yasui, M. Impact of ribonucleotide backbone on translesion synthesis

and repair of 7,8-Dihydro-8-oxoguanine. *J. Biol. Chem*. **291**, 24314-24323 (2016).

50. Sassa, A., Çağlayan, M., Dyrkheeva, N. S., Beard, W. A. & Wilson, S. H. Base excision

repair of tandem modifications in a methylated CpG dinucleotide. *J. Biol. Chem*. 289,

13996-4008 (2014).

51. Çağlayan, M. & Wilson, S. H. Pol μ dGTP mismatch insertion opposite T coupled with

ligation reveals a promutagenic DNA intermediate during double strand break repair.

Nat. Commun. **9**, 4213 (2018).

52. Çağlayan, M. Pol μ ribonucleotide insertion opposite 8-oxodG facilitates the ligation of premutagenic DNA repair intermediate. *Sci. Rep*. **10**, 940 (2020).

53. Tang Q., Gulkis, M., McKenna R. & Çağlayan, M. Structures of LIG1 that engage with mutagenic mismatches inserted by polβ in base excision repair. *Nat. Commun*. **13**, 3860 (2022).

54. Pascal, J. M. *et al.* Human DNA ligase I completely encircles and partially unwinds nicked DNA. *Nature* **432**, 473-478 (2004).

55. Tumbale, P. P. *et al.* Two-tiered enforcement of high-fidelity DNA ligation. *Nat. Commun.* **10**, 5431 (2019).

56. Sparks, J. L. *et al.* RNase H2-initiated ribonucleotide excision repair. *Mol. Cell* **47**, 980-986 (2012).

57. Cerritelli, S. M. & Crouch, R. J. Ribonuclease H: the enzymes in eukaryotes. *FEBS J.* **276**, 1494-1505 (2009).

58. Hiller, B., Achleitner, M., Glage, S., Naumann, R., Behrendt, R. & Roers, A. Mammalian RNase H2 removes ribonucleotides from DNA to maintain genome integrity. *J. Exp. Med*. **209**, 1419-1426 (2012).

59. Timson, D. J., Singleton, M. R. & Wigley, D. B. DNA ligases in the repair and replication of DNA. *Mutat. Res*. **460**, 301-318 (2000).

60. Shuman, S. DNA ligases: progress and prospects. *J. Biol. Chem*. **284**, 17365-17369 (2009).

61. Tomkinson, A. E., Vijayakumar, S., Pascal, J. M. & Ellenberger, T. DNA ligases: structure, reaction mechanism, and function. *Chem. Rev*. **106**, 687-699 (2006).

62. Doherty, A. J. & Suh, S. W. Structural and mechanistic conservation in DNA ligases. *Nuc. Acids Res*. **28**, 4051-4058 (2000).

63. Ellenberger, T. & Tomkinson, A. E. Eukaryotic DNA ligases: Structural and functional
insights. *Annu. Rev. Biochem.* **77**, 313-338 (2008).

64. Unciuleac, M., Goldgur, Y. & Shuman, S. Two-metal versus one-metal mechanisms of
lysine adenylylation by ATP-dependent and NAR$^+$-dependent polynucleotide ligases.
Proc. Natl. Acad Sci. U S A **114**, 2592-2597 (2017).

65. Tomkinson, A. E., Tappe, N. J. & Friedberg, E. C. DNA Ligase I from *Saccharomyces*
cerevisiae: Physical and biochemical characterization of the CDC9 gene product.
Biochemistry **31**, 11762-11771(1992).

66. Nishida, H., Kiyonari, S., Ishino, Y. & Morikawa, K. The closed structure of an
archaeal DNA ligase from *Pyrococcus furiosus*. *J. Mol. Biol.* **360**, 956-967 (2006).

67. Chen, Y. *et al.* Structure of the error-prone DNA ligase of African swine fever virus
identifies critical active site residues. *Nat. Commun.* **10**, 387 (2019).

68. Unciuleac, M., Goldgur, Y. & Shuman, S. Structures of ATP-bound DNA ligase D in
a closed domain conformation reveal a network of amino acid and metal contacts to the
ATP phosphates. *J. Biol. Chem.* **294**, 5094-5104 (2019).

69. Odell, M., Sriskanda, V., Shuman, S. & Nikolov, D. Crystalstructure of eukaryotic
DNA ligase–adenylate illuminates the mechanism of nick sensing and strand joining.
Mol. Cell **6**, 1183-1193 (2000).

70. Williams, J. S. *et al.* High-fidelity DNA ligation enforces accurate Okazaki fragment
maturation during DNA replication. *Nat. Commun.* **12**, 482 (2021).

71. Jurkiw, T. J. *et al.* LIG1 syndrome mutations remodel a cooperative network of ligand
binding interactions to compromise ligation efficiency. *Nucleic Acids Res.* **49**, 1619-
1630 (2021).

72. Williamson, A. & Leiros, H. K. S. Structural intermediates of a DNA-ligase complex

illuminate the role of the catalytic metal ion and mechanism of phosphodiester bond

formation. *Nucleic Acids Res*. **47**, 7147-7162 (2019).

73. Unciuleac, M., Goldgur, Y. & Shuman, S. Two-metal versus one-metal mechanisms of

lysine adenylylation by ATP-dependent and NAR$^+$-dependent polynucleotide ligases.

Proc. Natl. Acad Sci. U S A **114**, 2592-2597 (2017).

74. Dejaegere, A. P. & Case, D. A. Density functional study of ribose and deoxyribose

chemical shifts. *J. Phys. Chem*. **102**, 5280-5289 (1998).

75. Cotner-Gohara, E., Kim, I., Hammel, M., Tainer, J. A., Tomkinson, A. E. &

Ellenberger, T. Human DNA ligase III recognizes DNA ends by dynamic switching

between two DNA-bound states. *Biochemistry* **49**, 6165-6176 (2010) .

76. Kukshal, V., Kim, I., Hura, G. L., Tomkinson, A. E., Trainer, J. A. & Ellenberger, T.

Human DNA ligase III bridges two DNA ends to promote specific intermolecular DNA

end joining. *Nucleic Acids Res*. **43**, 7021-7031 (2015) .

77. Cuneo, M. J., Gabel, S. A., Krahn, J. A., Ricker, M. A. & London, R. E. The structural

basis for partitioning of the XRCC1/DNA ligase IIIα BRCT-mediated dimer

complexes. *Nucleic Acids Res*. **39**, 7816-7827 (2011).

78. Conlin, M. P., Reid, D. A., Small, G. W., Chang, H. H., Watanabe, G., Lieber, M. R.,

Ramsden, D. A. & Rothenberg, E. DNA Ligase IV guides end-processing choice during

nonhomologous end joining. *Cell Rep*., **20**, 2810-2819 (2017).

79. Ochi, T., Gu, X. & Blundell, T. L. Structure of the catalytic region of DNA ligase IV

in complex with an Artemis fragment sheds light on double-strand break repair.

Structure, **21**, 672-679 (2013).

80. Ochi, T., Wu, Q., Chirgadze, D. Y., Grossmann, J. G., Bolanos-Garcia, V. M. &

Blundell, T. L. Structural insights into the role of domain flexibility in human DNA

ligase IV. *Structure*, **20**, 1212-1222 (2012).

81. Kaminski, A., Tumbale, P. P., Schellenberg, M. J., Williams, R. S., Williams, J. G.,

Kunkel, T. A., Pedersen, L. C. & Bebenek, K. Structures of DNA-bound human ligase

IV catalytic core reveal insights into substrate binding and catalysis. *Nat. Commun*., **9**,

2642 (2018).

82. Tomkinson, A. E., Howes, T. R. & Wiest, N. E. DNA ligases as therapeutic targets.

Transl. Cancer Res. **2**, 31 (2013).

83. Zhong, S. *et al.* Identification and validation of human DNA ligase inhibitors using

computer-aided drug design. *J. Med. Chem*. **51**, 4553-4562 (2008).

84. Chen, X. *et al.* Rational design of human DNA ligase inhibitors that target cellular DNA

replication and repair. *Cancer Res*. **68**, 3169-3177 (2008).

85. Otwinowski, Z. & Minor, W. Processing of X-ray diffraction data collected in

oscillation mode. *Methods Enzymol*. **276**, 307-326 (1997).

86. McCoy, A. J. *et al*. Phaser crystallographic software. *J. Appl. Crystallogr.* **40**, 658-674

(2007).

87. Emsley, P. *et al.* Features and development of Coot. *Acta Crystallogr. D. Biol.*

Crystallogr. **66**, 486-501 (2010).

88. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for

macromolecular structure solution. *Acta Crystallogr. D. Biol. Crystallogr.* **66**, 213-221

(2010).

89. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal

structures. *Acta Crystallogr. D. Biol. Crystallogr.* **67**, 355-367 (2011).

90. Li, S., Olson, W. K. & Lu, X. J. Web 3DNA 2.0 for the analysis, visualization, and

modeling of 3D nucleic acid structures. *Nucleic Acids Res*. **47**, W26-W34 (2019).

Funding

This work was supported by a grant 1R35GM147111-01 from the National Institute of General

Medical Sciences (NIGMS).

Acknowledgements

This work is based upon research conducted at the Center for High Energy X-ray Sciences

(CHEXS), which is supported by the National Science Foundation under award DMR-

1829070, and the Macromolecular Diffraction at CHESS (MacCHESS) facility, which is

supported by award 1-P30-GM124166-01A1 from the National Institute of General Medical

Sciences, National Institutes of Health, and by New York State's Empire State Development

Corporation (NYSTAR). The authors thank Dr. Robert McKenna and Jacob Combs

(University of Florida) for beamline 7B2 at CHESS, and an assistance with crystal shipment

and data collection.

Author contributions

Conceptualization M.Ç., methodology and investigation T.Q., M.G., M.Ç.; writing-original

draft T.Q., M.G., M.Ç.; writing-reviewing and editing T.Q., M.G., M.Ç.; funding acquisition

M.Ç.

Figure 1. Structure of LIG1 bound to nick DNA duplex containing 3'-rG:C. A, X-ray

crystal structure of LIG1/nick DNA duplex containing a ribonucleotide (rG) at 3'-terminus of

the nick DNA. Simulated annealing omit map (Fo-Fc) of LIG1/rG:C (cyan) is displayed for 2'-

OH of 3'-terminus contoured at 3σ. Schematic view of DNA including DNA:RNA hybrid used

in the LIG1/nick DNA duplex crystallization. **B,** Simulated annealing omit map (Fo-Fc) of

LIG1/rG:C is displayed for 3'-OH terminus and AMP contoured at 3σ. DNA and AMP are

shown as sticks and the map is depicted in green.

Figure 2. LIG1/rG:C structure demonstrates a phosphodiester bond formation and AMP release during the final ligation step at atomic resolution. A-B, The 2Fo - Fc density map (1σ) is displayed for 3'- and 5'-terminus of the nick in the LIG1/rG:C (cyan) and LIG1/A:T (magenta) structures. AMP is incomplete and shows lower density in the LIG1/rG:C (step 3) than the LIG1/A:T (step 2). DNA and AMP are shown as sticks and the map is depicted in grey. **C**, Overlay of LIG1 structures bound to the nick DNA duplexes containing rG:C (cyan) and A:T (PDB: 7SUM, magenta) ends shows the difference at the positions of 3'- and 5'-terminus of the nick.

Figure 3. Mutagenic ligation of the nick DNA with 3'-ribonucleotide by LIG1. A-C, Lanes 1 and 8 are the negative enzyme controls of the nick DNA substrates with 3'-dG:C and 3'-rG:C, respectively. Lanes 2-7 and 9-14 are the ligation products in the presence of 3'-dG:C and 3'-rG:C, respectively, by wild-type (A) and EE/AA mutant (C) of LIG1, and correspond to time points of 0.5, 1, 3, 5, 8, and 10 min. **B-D,** Graphs show time-dependent change in the amount of ligation products. The data represent the average from three independent experiments ± SD.

Figure 4. Comparison of LIG1 active site residues between the structures of LIG1/nick

DNA complexes with rG:C versus cognate and mismatched ends. Overlay of LIG1

structures bound to the nick DNA duplexes containing rG:C (cyan), A:T (PDB: 7SUM,

magenta), and A:C (PDB: 7SX5, grey) ends shows the differences in the positions of 3'- and

5'-terminus at the nick DNA and no difference in the LIG1 active site F635, R871, and F872

residues around 2'-OH of the ribose.

Figure 5. Role of LIG1 active side residue F872 for mutagenic ligation of the nick DNA

with 3'-rG:C. A, Lane 1 is the negative enzyme control of the nick DNA substrate with 3'-

rG:C. Lanes 2-7 and 8-13 are the ligation reaction products by LIG1 F872 and EE/AA/F872A

mutants, respectively, and correspond to time points of 0.5, 1, 3, 5, 8, and 10 min. **B,** Graph

shows time-dependent change in the amount of ligation products. The data represent the

average from three independent experiments ± SD.

Table 1. X-ray data collection and refinement statistics of LIG1/nick DNA duplex with a

ribonucleotide (rG:C) at the 3'-end.

	$LIG1^{EE/AA}$ Nick DNA with 3'-rG:C
PDB entry ID	
Data collection	
Space group	$P2_12_12_1$
Cell dimensions	
a, b, c (Å)	65.065 115.633 126.338
a, b, g (°)	90
Resolution (Å)	20-3.0 (3.05-3.0)
R_{sym}	0.181 (1.451)
$I / \sigma (I)$	17.7 (1.7)
$CC_{1/2}$	1.000 (0.655)
CC^*	1.000 (0.890)
Completeness (%)	96.3 (100)
Redundancy	12.5 (13.6)
Refinement	
Resolution (Å)	20-3.0
No. reflections	17321
R_{work}/R_{free}	0.172/0.206
Non-H atoms:	
Protein/DNA	5665
AMP	21
Average B-factors ($Å^2$):	60.1
R.M.S.D	
Bond lengths (Å)	0.009
Bond angles (°)	1.174