# The DUSP1 on CaMKII-positive neurons in mPFC mediates adolescent cocaine exposure-induced higher sensitivity to cocaine during adulthood

Xiaoyan Wei[1,#], Jiasong Chang[1,#], Zhen Cheng[1], Wenwen Chen[1], Hao Guo[1], Zhaoyu Liu[1], Yuning Mai[1], Tao Hu[1], Yuanyuan Zhang[1], Qinglong Cai[1], Feifei Ge[1], Yu Fan[1,*], Xiaowei Guan[1,*]

[1]Department of Human Anatomy and Histoembryology, Nanjing University of Chinese Medicine, Nanjing, China

[#]These authors are equally contribute to this work

**Running title**: Adolescent cocaine and mPFC DUSP1

*Correspondence authors: Prof. Xiaowei Guan (guanxw918@njucm.edu.cn), Dr. Yu Fan (yfan@njucm.edu.cn)

Phone number: +86-25-85811387

**Summary**

Adolescent cocaine exposure causes higher cocaine-preferred behaviors during adulthood, along with evoked mPFC activity in response to cocaine challenge. Locally overexpressing but not knocking-down the dual specificity phosphatase 1 (DUSP1) on CaMKII-positive neurons (DUSP1$^{CaMKII}$) suppresses mPFC activation, and ultimately rescues the higher sensitivity to cocaine during adulthood.

**Abstract**

Adolescent cocaine abuse increases the risk for developing addiction in later life, which remains a big public health concern, but the underlying molecular mechanism is poorly understood. Here, adolescent cocaine-exposed (ACE) male mice models were established by administrating cocaine during adolescent period. When growing to adult age, mice were subjected to conditioned place preference (CPP) to evaluate the sensitivity to cocaine, then potential molecule of dual specificity phosphatase 1 (DUSP1) were screened out by transcriptomic sequencing. Subthreshold dose of cocaine (sdC), that is insufficient to produce CPP, was used to induce CPP in adulthood. The sdC treatment effectively induced CPP in ACE mice during adulthood, accompanied with the more triggered CaMKII-positive neurons, and induced higher *Dusp1* gene, lower DUSP1 protein, lower DUSP1 activity and lower DUSP1 expression on CaMKII-positive neurons (DUSP1$^{CaMKII}$) in medial prefrontal cortex (mPFC). Overexpressing DUSP1$^{CaMKII}$ suppressed CaMKII-positive neuronal activation, and ultimately blocked sdC-induced CPP in ACE mice during adulthood. While, knocking-down DUSP1$^{CaMKII}$ activated more CaMKII-positive neurons, and aggravated sdC-preferred behavior in ACE mice during adulthood. ERK1/2 might be potential subsequent signal for DUSP1 in the process. Collectively, our findings reveal a novel molecular mechanism underlying adolescent drug abuse-induced susceptibility to addiction during adulthood, and mPFC DUSP1$^{CaMKII}$ is a promising pharmacological target to predict or treat addiction, especially caused by adolescent substance use.

**Keywords:** adolescent cocaine exposure; sensitivity to cocaine; mPFC

activation; DUSP1$^{CaMKII}$

**Introduction**

Adolescents present vulnerable to substance abuse such as cocaine (1, 2).

Adolescent experiences of drug abuse enhances the risk for developing

addiction later in life (3, 4), however, the underlying molecular mechanism

remains much to be clarified. The medial prefrontal cortex (mPFC), a critical

brain area involved in the susceptibility to drug addiction (5, 6), undergoes

rapid development during adolescent period (7, 8). A single cocaine treatment

alters mPFC dendritic spines in adolescent rather than adult animals (9, 10),

indicating a sensitive response of adolescent mPFC to drugs (11, 12).

Previously, we found that adolescent cocaine exposure (ACE) enhanced the

GABAergic transmission on pyramidal neurons of mPFC (13), and produced a

prolonged modification on mPFC synapses in adult rodents (14).

The dual specificity protein kinase-phosphatase-1 (DUSP1), also known as

mitogen-activated protein kinase-phosphatase-1 (MKP-1), is a negative

regulatory factor on the activation of mitogen activated protein kinases

(MAPKs) (15). DUSP1 are expressed widely in the brain, including PFC (16).

By regulating dephosphorization of MAPKs such as ERK1/2, p38 and the

c-Jun N-terminal kinases (JNKs), DUSP1 has been proved to achieve

neuroprotection in hearing loss (17), Huntington's disease (18), cerebral

ischemia reperfusion injury (19) and neuropathic pain (20) through its

anti-inflammatory, antioxidant, mitochondrial repair and supporting axon

survival functions. In addition, *Dusp1* gene are known to be activity-dependent

responded to stimuli signals in the brain, thus often acting as an immediate

early gene (IEGs) like *c-fos*. For example, psilocybin strongly upregulates

*Dusp1* levels in PFC (21), while synthetic cathinone (substitutes for illicit

psychostimulants including cocaine) produces increased *Dusp1* genes in

striatum (22).

In the present study, ACE male mice model was established by exposing

adolescent male mice to cocaine from postnatal day 28 (P28) to P42, a period

that represents adolescent in rodents (23). When growing up to adult age,

mice were subjected to conditioned place preference (CPP) to evaluate the

sensitivity to drug. Subthreshold dose of cocaine, which is insufficient to

produce CPP, was used to induce CPP (sdC). With method of transcriptomic

sequencing and locally regulating DUSP1 on CaMKII-positive neurons

(DUSP1$^{CaMKII}$), the role of mPFC DUSP1$^{CaMKII}$ in the sensitivity to cocaine were

explored in ACE mice during adulthood.

**Results**

**The subthreshold dose of cocaine produce CPP and trigger more mPFC**

**CaMKII-positive neurons in ACE mice during adulthood**

To examine the sensitivity to cocaine, mice were subjected to cocaine-induced

CPP training and tests in mice during adulthood (Figure 1A). Subthreshold

dose (1 mg / kg) of cocaine (sdC), which is insufficient to induce CPP in naïve

mice, were used in this study. As shown in Figure 1B-D, the sdC treatment did

not produce CPP in adolescent saline-exposed (ASE-sdC) mice (n = 18 mice, *t*

= 1.306, *p* = 0.3586), but produced CPP in ACE mice during adulthood

(ACE-sdC) (n = 24 mice, *t* = 10.91, *p* < 0.0001, Figure 1B, 1D). The ΔCPP

score was higher in ACE-sdC mice than that in ASE-sdC mice (n = 42 mice, *t* =

8.129, *p* < 0.0001, Figure 1C, 1D). These results indicate that adolescent

cocaine exposure produce a higher sensitivity to cocaine in mice during

adulthood.

To map the brain activity in respond to subthreshold dose administration of

cocaine in adult mice, c-Fos and NeuN were used as markers to evaluate

neuronal activation and to label neuron, respectively. Compared with ASE-sdC

mice during adulthood, the c-Fos-stained levels of neurons is much higher in

mPFC (n = 12 slice from 6 mice, $t$ = 2.89, $p$ = 0.0161) and its subregions of

cingulate cortex (Cg1, n = 12 slice from 6 mice, $t$ = 2.912, $p$ = 0.0155) and

prelimbic cortex (PrL, n = 12 slice from 6 mice, $t$ = 2.311, $p$ = 0.0435), but

similar in infralimbic cortex (IL, n = 12 slice from 6 mice, $t$ = 0.8092, $p$ = 0.4372,

Figure 1E), orbitofrontal cortex (OFC, n = 12 slice from 6 mice, $t$ = 2.055, $p$ =

0.0670), nucleus accumbens core (NAc core, n = 12 slice from 6 mice, $t$ =

0.3219, $p$ = 0.7542), nucleus accumbens shell (NAc shell, n = 12 slice from 6

mice, $t$ = 1.632, $p$ = 0.1337), bed nucleus of the stria terminalis (BNST, n = 12

slice from 6 mice, $t$ = 1.172, $p$ = 0.2685), triangular septal nucleus (TS, n = 12

slice from 6 mice, $t$ = 1.768, $p$ = 0.1074), basolateral amygdala (BLA, n = 12

slice from 6 mice, $t$ = 1.304, $p$ = 0.2215), central nucleus of the amygdala (CeA,

n = 12 slice from 6 mice, $t$ = 0.2851, $p$ = 0.7814), medial habenula (mHb, n =

12 slice from 6 mice, $t$ = 0.2165, $p$ = 0.8329), lateral habenula (LHb, n = 12

slice from 6 mice, $t$ = 1.151, $p$ = 0.2764), dorsal hippocampal CA1 (dCA1, n =

12 slice from 6 mice, $t$ = 0.09203, $p$ = 0.9285), dorsal hippocampal CA3 (dCA3,

n = 12 slice from 6 mice, $t$ = 0.6802, $p$ = 0.5118), dorsal dentate gyrus (dDG, n

= 12 slice from 6 mice, $t$ = 0.5304, $p$ = 0.6074), ventral hippocampal CA1

(vCA1, n = 10 slice from 6 mice, $t$ = 0.1572, $p$ = 0.8790) and ventral tegmental

area (VTA, n = 12 slice from 6 mice, $t$ = 0.2915, $p$ = 0.7767, Figure S1A) in

ACE-sdC mice during adulthood. Further, In ACE-sdC mice, the c-Fos staining

in mPFC (n = 6 mice, r = 0.9048, $p$ = 0.0132), Cg1 (n = 6 mice, r = 0.9206, $p$ =

0.0092) and PrL (n = 6 mice, r = 0.8646, $p$ = 0.0262), but not in IL (n = 6 mice, r

= 0.4181, $p$ = 0.4094) showed significant and positive correlation with the

corresponding ΔCPP score in ACE-sdC mice during adulthood (Figure 1F).

These results imply that mPFC activation, especially Cg1 and PrL, might be

involved in the adolescent cocaine exposure-induced higher sensitivity to

cocaine in mice during adulthood.

To observe the effects of adolescent cocaine exposure on baseline levels of

mPFC activity, the c-Fos levels without administration of sdC were test in mice

on P74. As shown in Figure S2A, before the sdC administration, the c-Fos

protein levels (n = 6 mice, $t$ = 5.899, $p$ = 0.0041) of mPFC was higher in ACE

than that in ASE mice on P74. These results indicate that adolescent cocaine

exposure evoke mPFC activation in mice during adulthood.

To explore the activity of mPFC different cell types in response to sdC

treatment, CaMKII, GAD67, PV and SOM were used as markers to label

glutamatergic, GABAergic, PV-positive GABAergic and SOM-positive

GABAergic neuron, respectively. As shown in Figure S2B, the c-Fos-stained

CaMKII-positive glutamatergic neurons (n = 12 slice from 6 mice, $t$ = 3.656, $p$ =

0.0044), but not GAD67-positive GABAergic neurons (n = 12 slice from 6 mice,

$t$ = 1.180, $p$ = 0.2652) and its subtypes of PV-positive (n = 12 slice from 6 mice,

$t$ = 0.5668, $p$ = 0.5833) and SOM-positive neurons (n = 12 slice from 6 mice, $t$

= 0.6119, $p$ = 0.5543), were much more in numbers at mPFC of ACE-sdC mice

than that in ASE-sdC mice during adulthood.

To *in vivo* record real-time calcium signals of mPFC CaMKII-positive

neurons, *rAAV2/9-CaMKII-GCaMp6m-WPRE-PA* virus were injected into

mPFC, specially limiting to subregions of Cg1 and PrL (Figure 1G). As shown

in Figure 1H-J, at first 4 sec-duration after mice entering into the

cocaine-paired chamber, the calcium signals of CaMKII-positive neurons were

significantly at higher levels in GCaMp6m-transfected ACE-sdC mice than that

in ASE-sdC mice (n = 6 mice, $t$ = 3.839, $p$ = 0.0185). These results indicate

that mPFC CaMKII-positive neurons might be involved in adolescent cocaine

exposure-induced higher sensitivity to cocaine in mice during adulthood.

**Subthreshold dose of cocaine increase mPFC *Dusp1* gene while**

**decrease DUPS1 protein and activity in ACE mice during adulthood**

To screen potential molecules that contribute to the higher mPFC activity in

response to sdC administration, the RNA sequencing was performed in mPFC

on P78 following CPP test. As shown in Figure S3A, following administration of

sdC, 180 genes were upregulated and 248 genes were downregulated in

mPFC of ACE-sdC mice, when compared to ASE-sdC mice. According to

Protein-Protein Interaction (PPI) network analysis of upregulated genes

(Figure S3B) and heatmaps of top 40 upregulated genes (Figure S3C), *Dusp1*

is one of the most significantly upregulated genes and a hub interactor gene

that associated with other upregulated genes including *c-Fos*. Further,

MAPK-related signals were both in the top 30 enriched GO (Figure S3D) and

the top enriched 20 KEGG pathway (Figure S3E) of upregulated genes in

mPFC of ACE-sdC mice. The transcriptomic results imply that *Dusp1*/DSUP1

and related MAPK signals might be a critical molecule that implicated in

adolescent cocaine exposure-increased sensitivity to cocaine.

The baseline level and responsive level to cocaine of mPFC *Dusp1* gene,

DUSP1 protein and activity were examined on P74 and P78, respectively.

There is similar *Dusp1* gene (n = 6 mice, $t$ = 0.3691, $p$ = 0.7307, Figure 2A),

DUSP1 protein (n = 6 mice, $t$ = 1.905, $p$ = 0.1295, Figure 2B), and DUSP1

activity (n = 6 mice, $t$ = 0.1855, $p$ = 0.8618, Figure 2C) between ACE and ASE

mice on P74, indicating that adolescent cocaine exposure did not influence the

baseline levels of *Dusp1*/DUSP1 in mPFC. However, on P78 following

administration of sdC, *Dusp1* gene (n = 11 mice, $t$ = 4.100, $p$ = 0.0027, Figure

2D) were much higher, while DUSP1 protein level (n = 6 mice, $t$ = 4.929, $p$ =

0.0079, Figure 2E), DUSP1 activity (n = 12 mice, $t$ = 3.240, $p$ = 0.0089, Figure

2F) and DUSP1-staining value on CaMKII-positive neurons (n = 6 mice, $t$ =

5.751, $p$ = 0.0045, Figure 2G) were much lower in ACE-sdC mice than that in

ASE-sdC mice. With these results, we further infer that *Dusp1*/DUSP1 on

CaMKII-positive neurons (DUSP1$^{CaMKII}$) may contribute to the sdC-evoked

mPFC CaMKII-positive neurons and higher sensitivity to cocaine in ACE mice

during adulthood.

Since DUSP1 is important regulatory molecule in the MAPK-related signal

molecules, the total ERK1/2 and its phosphorylation (p-ERK) levels of mPFC

was examined on P78 following sdC administration. As shown in Figure 2H,

the levels of ratio of p-ERK to ERK (n = 6 mice, $t$ = 3.338, $p$ = 0.0289) and

p-ERK (n = 6 mice, $t$ = 3.143, $p$ = 0.0348) were significantly higher, while total

ERK1/2 (n = 6 mice, $t$ = 1.030, $p$ = 0.3611) was similar in ACE-sdC than that in

ASE-sdC mice.

**Overexpressing DUSP1$^{CaMKII}$ suppress cocaine-induced CPP and reduce**

**mPFC activation in ACE mice during adulthood**

To explore the role of mPFC DUSP1$^{CaMKII}$ in the higher sensitivity to cocaine in

ACE mice, *AAV9-CaMKII-DUSP1-RNAOE-EGFP* (OE virus) and

*AAV9-CaMKII-EGFP* (Ctrl virus) were bilaterally infused into mPFC, especially

to Cg1 and PrL (Figure 3A-B). As shown in Figure 3C, about 80%

CaMKII-positive neurons were transfected with virus. Compared to Ctrl

virus-treated ACE-sdC mice (ACE-sdC$_{Ctrl}$), *Dusp1* gene (n = 6 mice, $t$ = 4.413,

$p$ = 0.0116, Figure 3D), DUSP1 protein (n = 6 mice, $t$ = 4.071, $p$ = 0.0152,

Figure 3E) and DUSP1 activity (n = 12 from 6 mice, $t$ = 3.336, $p$ = 0.0075,

Figure 3F) levels were much higher in OE virus-treated ACE-sdC mice

(ACE-sdC$_{OE}$), indicating the good efficiency of OE virus.

As shown in Figure 3G-I, ACE-sdC$_{Ctrl}$ mice still showed sdC-induced

preference behavior (n = 13 mice, $t$ = 3.215, $p$ = 0.0074), while ACE-sdC$_{OE}$

mice did not exhibit sdC-related CPP (n = 13 mice, $t$ = 2.051, $p$ = 0.1001),

accompanied by a lower ΔCPP score in ACE-sdC$_{OE}$ mice (n = 26 mice, $t$ =

3.723, $p$ = 0.0011). These results indicate overexpressing DUSP1$^{CaMKII}$ could

suppress adolescent cocaine exposure-induced higher sensitivity to cocaine

during adulthood.

To explore the regulatory role of DUSP1 in the activity of transfected CaMKII

neurons (EGFP-positive), c-Fos staining were performed on mPFC. As shown

in Figure 3J, the percentage of c-Fos-positive & EGFP-positive in transfected

CaMKII neurons was much lower in ACE-sdC$_{OE}$ mice than that in ACE-sdC$_{Ctrl}$

mice (n = 12 slice from 6 mice, $t$ = 3.624, $p$ = 0.0047). In parallel, the ratio of

p-ERK to ERK (n = 6 mice, $t$ = 3.503, $p$ = 0.0248) and p-ERK (n = 6 mice, $t$ =

9.063, $p$ = 0.0008) levels were significantly decreased, but ERK level (n = 6

mice, $t$ = 0.1728, $p$ = 0.8712) was not changed by OE virus, compared to Ctrl

virus (Figure 3K). These results indicate DUSP1$^{CaMKII}$ negatively regulate

mPFC CaMKII-positive neurons.

**223　Knocking-down DUSP1$^{CaMKII}$ aggravate cocaine-preferred behaviors and**

**224　further activate mPFC activity in ACE mice during adulthood**

To explore the role of ACE-increased *Dusp1* gene in further confirm the role of

mPFC DUSP1$^{CaMKII}$ in mPFC activity and sensitivity to cocaine,

*AAV9-CaMKII-DUSP1-RNAi-EGFP* (KD virus) and *AAV9-CaMKII-EGFP* (Ctrl

virus) were bilaterally infused into mPFC, especially to Cg1 and PrL (Figure

4A-B). As shown in Figure 4C, above 70% CaMKII-positive neurons were

transfected with virus. Compared to Ctrl virus-treated ACE-sdC mice

(ACE-sdC$_{Ctrl}$ mice), *Dusp1* gene (n = 8 mice, $t = 9.250$, $p < 0.0001$, Figure 4D),

DUSP1 protein (n = 6 mice, $t = 11.00$, $p = 0.0004$, Figure 4E) and DUSP1

activity (n = 11 from 6 mice, $t = 3.256$, $p = 0.0099$, Figure 4F) were lower in KD

virus-treated ACE-sdC mice (ACE-sdC$_{KD}$ mice), indicating good efficiency of

KD virus.

As shown in Figure 4G-I, both ACE-sdC$_{Ctrl}$ mice (n = 7 mice, $t = 2.839$, $p =$

0.0236) and ACE-sdC$_{KD}$ mice (n = 11 mice, $t = 7.104$, $p < 0.0001$) exhibited

cocaine-preferred behaviors. The ΔCPP score in ACE-dsC$_{KD}$ mice is higher

than that in ACE-sdC$_{Ctrl}$ mice (n = 18 mice, $t = 2.211$, $p = 0.0419$), indicating an

aggravating cocaine-preferred behavior in ACE-sdC$_{KD}$ mice.

The percentage of c-Fos-positive & EGFP-positive in transfected CaMKII

neurons was much higher in ACE-sdC$_{KD}$ mice than that in ACE-sdC$_{Ctrl}$ mice (n

= 16 slice from 8 mice, $t = 4.744$, $p = 0.0003$, Figure 4J). The ratio of p-ERK to

ERK (n = 6 mice, $t = 3.293$, $p = 0.0301$), p-ERK (n = 6 mice, $t = 3.903$, $p =$

0.0175) levels were increased, and ERK (n = 6 mice, $t$ = 0.1856, $p$ = 0.8618)

level was not changed by DUSP1 KD virus, compared to Ctrl virus (Figure 4K).

Together with the results of overexpressing DUSP1[CaMKII], these results

indicate that DUSP1[CaMKII] mediate adolescent cocaine exposure-induced

higher sensitivity to cocaine during adulthood, which is through regulating

mPFC CaMKII-positive neuronal activation.

**Discussion**

The current study found that ACE induce higher sensitivity to cocaine during

adulthood. Subthreshold dose of cocaine (sdC), triggered more

CaMKII-positive neurons in mPFC, especially subregions of Cg1 and PrL,

accompanied with lower DUSP1 protein and higher *Dusp1* gene in mPFC of

ACE mice during adulthood. To explore the roles of converse phenotypes of

DUSP1 protein and *Dusp1* gene in response to sdC treatment, both

upregulated and downregulated modulations of DUSP1[CaMKII] were performed

in mPFC of ACE mice. We found that specific overexpressing DUSP1[CaMKII]

efficiently blocked cocaine-induced CPP in ACE mice, and reduced

CaMKII-positive neuronal activation in mPFC. By contrast, knocking-down

DUSP1 aggravated cocaine-preferred behavior in ACE mice, and further

activated CaMKII-positive neuronal activation in mPFC. ERK1/2 might be

potential subsequent signal for DUSP1 in the process. (Figure 5).

Adolescence is a critical period of brain development and maturation so that

makes the adolescent brain highly susceptible to neurotoxicity of drugs.

Exposure to Δ9-tetrahydrocannabinol (psychoactive ingredient of cannabis)

enhanced sensitivity to cocaine in rats even into adulthood (24), and

adolescent methylphenidate use (psychostimulant drug to treat attention deficit

hyperactivity disorder) increased reactivity and vulnerability to cocaine in rats

of adulthood (25), suggesting that adolescent use of the psychostimulant is an

important risk factor for addiction susceptibility later in life. Previously using the

similar ACE mice models with this study, we found that there was no difference

on acquisition, extinction and reinstatement of CPP by repeated exposure to

addictive doses of cocaine (15 mg/kg) in adulthood between ACE mice and

controls (13). However, here we found that subthreshold dose administration

of cocaine (1 mg/kg) effectively induced cocaine-preferred behaviors in ACE

mice but not in control mice of their adulthood, indicating that ACE increase the

sensitivity to cocaine which may render adult mice more susceptible to cocaine.

Cass et.al (26) found that adolescent cocaine administration produced mPFC

disinhibition that endured throughout adulthood. Our previous studies found

that adolescent cocaine exposure induced prolonged modifications on

synapses (14) and triggered neuronal activation (13) in mPFC of adult rodents,

which contribute to the increased anxiety-like behaviors later in life. Here, by

mapping the activation of potential brain regions involved in the increased

sensitivity to cocaine, we found that mPFC, especially glutamatergic neurons

in Cg1 and PrL, were obviously and strongly evoked in response to cocaine

challenge in adult ACE mice. These findings suggest that ACE-induced plastic

changes in mPFC play critical roles in facilitation of susceptibility to drugs later

in life. Previously, we found that ACE suppressed mPFC activity partially

through enhancing local GABAergic transmission on mPFC pyramidal neurons

(13) and reduced dendritic spines of mPFC pyramidal neurons (14) in

adulthood. However, unlike baseline condition, here we found that cocaine

challenge in adulthood activated more glutamatergic neurons but did not

change GABAergic including its subtypes of PV-positive and SOM-positive
neurons in mPFC of ACE mice during adulthood. Based on these results, we
hypothesized that the cocaine-changed signals in glutamatergic neurons of
mPFC should be the mechanic basis for mPFC activation and susceptibility to
cocaine-induced CPP during adulthood in mice treated with cocaine during
adolescence.

In order to screen key signals, mPFC transcriptome was performed after
cocaine administration in adult mice. We found that *Dusp1* gene and genes
related to mitogen-activated protein kinase phosphatases (MAKPs) pathway
were obviously upregulated in mPFC after cocaine challenge in ACE mice.
While, level and activity of DUSP1 protein were decreased at the same time.
We believe that the inconsistency between gene and protein changes of
DUSP1 is due to their different meanings reflected in response to cocaine
challenge. Takaki et al. (27) reported that *Dusp1* mRNA levels were
significantly increased after acute or chronic methamphetamine administration
in PFC of animals. We thought that the increased *Dusp1* gene in response to
cocaine challenge, similar to immediate early genes, such as *c-fos*, *Jun*, *Ark*,
reflected rapid and sensitive response of mPFC to cocaine in ACE mice,
further indicating the important involvement of mPFC in process of
susceptibility to drugs. In contrast, we found that the decreased DUSP1 protein
level and activity was consistent with upregulated genes in MAPK-related
signals by transcriptome analysis in mPFC of ACE mice by cocaine challenge
during adulthood. Thus, DUSP1 protein might be the key regulatory molecule
to plastic changes in mPFC which underlies the increased sensitivity to drugs
in ACE mice during adulthood. Indeed, we found that specific overexpression

but not knockdown of mPFC DUSP1[CaMKII] efficiently suppressed

cocaine-activated mPFC, and ultimately reduced cocaine-preferred behaviors

in ACE mice. As a negative regulatory factor on MAPK-related signals, DUSP1

is also known as MKP-1. By dephosphorylating p38, JNKs and ERK (28),

DUSP1 regulates the brain function. Among them, ERK-related signals are

well-studied in cocaine addiction. The ERK signaling pathway is a canonical

MAPK signaling pathway that has been implicated in synaptic plasticity and

neuronal growth (29). Acute cocaine administration is sufficient to increased

phosphorylation of ERK in PFC, which contribute to the long-term plastic

changes by cocaine in the brain (30-32). While, blockade of ERK activation

during re-exposure to cocaine erased previously cocaine-preferred behaviors

(33). In the present study, we found that levels of p-ERK increased when some

changes in mPFC by cocaine, while it was decreased in

DUSP1-overexpressed mPFC with reduced activity in ACE mice, suggesting

that DUSP1 may recruit ERK-related signals to regulate mPFC activity, and

therefore influence sensitivity to cocaine challenge in ACE mice. In the future

study, the mechanistic investigation of DUSP1/MAPK signal pathway

underlying the phenotypes of higher cocaine-preferred behaviors should be

further explored in ACE mice.

Collectively, our findings reveal a novel role of mPFC DUSP1[CaMKII] in higher

sensitivity to cocaine in ACE mice during adulthood, which is through

regulating mPFC activity. The mPFC DUSP1[CaMKII] may represent a promising

therapeutic target for treatment of addiction, especially caused by adolescent

substance use.

**Materials and methods**

**Animals**

Male C57BL/6 wild type (WT) mice weighing 12-15 g were used. All animals were housed at constant humidity (40~60%) and temperature ($24 \pm 2^\circ$C) with a 12-hour light/dark cycle (lights on at 8 a.m.) and allowed free access to food and water. All mice were handled for three days before onset of experiments. All procedures were carried out in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee (IACUC) at Nanjing University of Chinese Medicine.

**Drug treatment**

From P28 to P42, male C57BL/6 WT mice were assigned to receive cocaine hydrochloride (15 mg/kg, dissolved in saline, i.p., Qinghai Pharmaceutical, China) or saline (0.2 mL, i.p.) once daily for 15 consecutive days. From P43-P74, these mice were kept at their home cage (4 mice per cage). From P75 to P78, a subset of mice were subjected to CPP training and test induced by subthreshold dose of cocaine hydrochloride (1 mg/kg).

**CPP**

The CPP test was performed in the TopScan3D CPP apparatus (CleverSys, VA, USA), which is constructed of two distinct chambers (15 × 15 × 23 cm each) separated by a removable guillotine door. The CPP procedure consisted of three phases: the pre-conditioning test (Pre-test, P75), conditioning (CPP training, P76-77) and post-conditioning test (Test, P78). Here, a subthreshold dose (1 mg/kg) of cocaine, which is believed not enough to induce CPP in naïve mouse, was used to induce CPP in mice on P78. During CPP training, mice were injected with saline (0.2 mL, i.p.) in the morning and cocaine (1

mg/kg, i.p.) in the afternoon once daily. After each injection, the mice were

confined to one chamber (non-drug-paired chamber or drug-paired chamber)

for 45 min. During the Pre-test and Test, mice freely access two chambers for

15 min. The CPP score is the time spent in drug-paired chamber minus that in

non-drug-paired chamber, and the ΔCPP score is the test CPP score minus

the pre-test CPP score.

**Immunofluorescence**

On P78, brain tissue were collected from a subset of mice within 90 min after

CPP test. Some brains of those were perfused with 4% paraformaldehyde

(PFA). The coronal brain sections (30 µm) were cut on a cryostat (Leica,

Germany). The sections were incubated in 0.3% (v/v) Triton X-100 for 0.5 h,

blocked with 5% donkey serum for 1.5 h at room temperature, and incubated

overnight at 4°C with the following primary antibodies: rabbit anti-c-Fos

(1:1500, RRID: AB_2247211, Cell Signalling Technology, USA), mouse

anti-NeuN (1:800, RRID: AB_2298772, Millipore, USA), DUSP1 (1:100,

BM4856, Boster Biotechnology, China), mouse anti-GAD67 (1:500, RRID:

AB_2278725, Millipore, USA), mouse anti-PV (1:250, BM1339, Boster

Biotechnology, China), mouse anti-SOM (1:50, RRID:AB_2271061, Santa

Cruz, USA), mouse anti-CaMKIIα (1:100, SC-13141, Santa Cruz, USA) and

mouse anti-CaMKIIα (1:200, RRID: AB_2721906, Cell Signalling Technology,

USA), followed by the corresponding fluorophore-conjugated secondary

antibodies for 1.5 h at room temperature. The following secondary antibodies

were used here: Alexa Fluor 555-labeled donkey anti-rabbit secondary

antibody (1:500, RRID: AB_2762834, Invitrogen, USA), Alexa Fluor

488-labeled donkey anti-mouse secondary antibody (1:500, RRID:

AB_141607, Invitrogen, USA). Fluorescence signals were visualized using a

Leica DM6B upright digital research microscope (Leica, Germany) or Leica

TCS SP8 (Leica, Germany).

**Fiber photometry**

On P51, the *rAAV2/9-CaMKII-GCaMp6m* (PT-0111, 3.24E+12 vg/ml, Brain

VTA, China) virus was bilaterally injected into mPFC (AP + 2.10 mm, ML ± 0.3

mm, DV − 2.15 mm), especially into PrL of mice. On P78, the GCaMp6m

signals in mPFC were recorded during CPP test. An optical fiber (200 μm outer

diameter, 0.37 numerical aperture, Inper Ltd., China) was placed 100 μm

above the viral injection site. The calcium-dependent fluorescence signals

were obtained by stimulating cells that transfected GCaMp6m virus with a 470

nm LED (35-40 μW at fiber tip), while calcium-independent signals were

obtained by stimulating these cells with a 405 nm LED (15-20 μW at fiber tip).

The LED lights of 470 nm and 405 nm were alternated at 66 fps and light

emission was recorded using sCMOS camera containing the entire fiber

bundle (2 m in length, NA = 0.37, 200 μm core, Inper Ltd.). The analog voltage

signals fluorescent was filtered at 30 Hz and digitalized at 100 Hz. The

GCaMp6m signals were recorded and analyzed by Inper Studio Multi Color

EVAL15 software (Inper Ltd., China) and Inper Data Process V0.5.9 (Inper Ltd.,

China), respectively. F0 is the baseline fluorescence signal which was

recorded for 1 sec before mice entering into the cocaine-paired chamber. F is

the real-time fluorescence signal which was recorded for 4 sec after mice

entering into the cocaine-paired chamber during CPP test. Each recording trail

was with 5 sec recording duration. The values of ΔF/F are calculated by

(F-F0)/F0. The area under curve (AUC) is the integral under recording duration

related to corresponding baseline at every trial.

**The mRNA sequencing and data processing in mPFC**

On P78, mPFC tissue (n = 6 per group) were collected from mice after CPP

test. Library preparation was performed by Novogene Bioinformatics

Technology Co., Ltd (China). RNA degradation and purity were monitored on 1%

agarose gels and using the NanoPhotometer ® spectrophotometer (IMPLEN,

CA, USA). RNA integrity was assessed using the RNA Nano 6000 Assay Kit of

the Bioanalyzer 2100 system (Agilent Technologies, CA, USA). The mRNA

was purified and fragmented from a total amount of 1 μg RNA per sample. The

cDNA was synthesized, adenylated and ligated with hairpin-structured

NEBNext Adaptor before hybridization. 250-300 bp cDNA fragments were

purified and preferentially selected for hybridization. The PCR products were

purified (AMPure XP system) and library quality was assessed on the Agilent

Bioanalyzer 2100 system. Index-coded samples were clustered on a cBot

Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS (Illumia)

and sequenced on an Illumina Novaseq platform. Differential mRNA

expression analysis was performed using the DESeq2 R package (1.16.1).

The $p$ values were adjusted using the Benjamini & Hochberg method. Cutoffs

of an absolute fold change > 1 and a correlated $p$ value < 0.05 were set for

significantly differential expression. The Bioinformatics analysis, including GO,

KEGG and PPI of upregulated genes by a subthreshold dose of cocaine in

ACE mice were performed in this study.

**RNA extraction and quantitative real-time PCR**

On P78, brain tissue were collected from a subset of mice within 90 min after

CPP test. The mPFC tissue were dissected immediately on dry ice. Total RNA

was extracted from mPFC using FastPure Cell/Tissue Total RNA Isolation Kit

(Vazyme, China). After RNA extraction, the amount of RNA was normalized

across samples, and cDNA was created using Vazyme (HiScript II Q RT

SuperMix for Qpcr (+gDNA wiper)). The β-actin was used as the internal

control. The primers of DUSP1 and β-actin were as follows: DUSP1 (forward,

GTTGTTGGATTGTCGCTCCTT; reverse, TTGGGCACGATATGCTCCAG),

β-actin (forward, GGCTGTATTCCCCTCCATCG; reverse,

CCAGTTGGTAACAATGCCATGT). DUSP1 mRNA levels were normalized to

β-actin mRNA levels. The relative mRNA level was calculated by the

comparative CT method ($2^{-\Delta\Delta Ct}$).

**Western blot**

On P74 and P78, brain tissue were collected from two subsets of mice without

CPP training (P74) or within 90 min after CPP test (P78). The mPFC tissue

were dissected immediately on dry ice. Total mPFC protein was extracted

using RIPA lysis buffer (Shanghai Beyotime, China). Protein samples (15 μg)

was separated by 10% SDS–PAGE or 12% SDS–PAGE and

electrophoretically transferred onto PVDF membranes. The transferred

membranes were blocked with 5% non-fat dry milk and 0.1% Tween 20 in 10

mM Tris–HCl (TBST buffer, pH 7.5) for 1.5 h at room temperature, then

subsequently incubated with the following primary antibodies: c-Fos (1:1500,

Rabbit, RRID: AB_2247211, Cell Signaling Technology, USA), DUSP1 (1:1000 ,

Rabbit, BM4856, Boster Biotechnology, China), $ERK_{1/2}$ (1:1000, Rabbit,

BM4326, Boster Biotechnology, China), p-$ERK_{1/2}$ (1:1000, Rabbit, BM4156,

Boster Biotechnology, China). The next day, the membranes were washed

three times in Tris-buffered saline with Tween 20 (TBS-T) and incubated with

horseradish peroxidase (HRP)-conjugated secondary antibody goat anti-rabbit

(1:5000, RRID: AB_2814709, Beijing ComWin Biotech Co., China) at room

temperature for 1 h. The blots were visualized by the ECL kit (Beijing ComWin

Biotech Co., China) and the signal was visualized by imaging system

(Tanon-5200, Shanghai, China). The blots were washed with stripping buffer

(Beyotime Institute of Biotechnology) to allow reprobing with other antibodies.

In this study, β-actin was used as control. Values for target protein levels were

calculated using Image J software (NIH, USA). The relative level of each

protein was normalized to β-actin.

**Total dual specificity phosphatase 1 (DUSP1) measurement in mPFC**

**by ELISA**

On P74 and P78, brain tissue were collected from two subsets of mice without

CPP training (P74) or within 90 min after CPP test (P78). The mPFC tissue

were dissected immediately on dry ice. The activity of mPFC DUSP1 was

measured by commercial ELISA (Mouse DUSP1 ELISA Kit instruction,

Shanghai shyx-bio.com, China). The mPFC tissues were diluted 1:10 in saline

and mashed using Tissue Grinder (Wuhan Servicebio technology CO., China),

then centrifuged for 30 min at 3000×g at 2-8 ℃ and the supernatants were

store samples at -20 ℃. The 50 µL mixture of sample and Sample Diluent (1:4)

were put in one well of plate. Then, add 100 µL of HRP-conjugate reagent to

each well, cover with an adhesive strip and incubate for 60 min at 37°C.

Washing the wells with Wash Solution (400 µL) five times. Then, added

chromogen solution A 50 µL and chromogen solution B 50 µL to each well and

incubated for 15 min at 37°C. The reaction was stopped by adding 50 µL Stop

Solution. Read the Optical Density (O.D.) at 450 nm using a standard

microplate reader (BIO-RAD iMark Microplate Reader, USA) within 15 min.

**496  Specific overexpression or knockdown of mPFC DUSP1$^{CaMKII}$**

On P51, four cohorts of mice were anesthetized with 2% isoflurane in oxygen,

and were fixed in a stereotactic frame (RWD, China). A heating pad was used

to maintain the core body temperature of the animals at 36°C. The coordinates

of mPFC were AP + 2.10, ML ± 0.3, DV − 2.15 in mm. In Experiment of

overexpression of DUSP1, a volume of 400 nL of

*CaMKIIap-MCS-EGFP-3Flag-SV40* *PloyA*

(*AAV9-CaMKIIap-DUSP1-RNAOE-EGFP,* 4.90E+13v.g/ml; Cat#:

GOSV0322969, Gene Chem, China) or *CaMKIIap-EGFP-SV40 PloyA*

(*AAV9-CaMKIIap-EGFP*, 4.01E+12v.g/ml; Cat#: AAV9/CON540, Gene Chem,

China) bilaterally into the mPFC (AP + 2.10 mm, ML ± 0.3 mm, DV − 2.15 mm)

at a rate of 80 nL/min. In Experiment of knockdown of DUSP1, a volume of 400

nL of *CaMKIIap-EGFP-MIR155(MCS)-SV40* *PloyA*

(*AAV9-CaMKIIap-DUSP1-RNAi-EGFP,* 1.10E+13 v.g/ml; Cat#: GIDV0295686,

Gene Chem, China) or *CaMKIIap-EGFP-SV40 PloyA* (*AAV9-CaMKIIap-EGFP*,

4.01E+12v.g/ml; Cat#: AAV9/CON540, Gene Chem, China) bilaterally into the

at a rate of 80 nL/min. After surgery, mice were maintained at home cage

about 3 weeks.

**514  Statistical analysis**

515  Statistical analysis was carried out using GraphPad Prism 8.0.2 software. All

516  data are presented as the Mean ± SEM. The data of CPP score between

517  ASE-C and ACE-C groups are analyzed by two-way analysis of variance

(ANOVA), while CPP score between Pre-test and Test in each mouse were analyzed by paired *t*-tests. Other data were analyzed by unpaired *t*-tests. GO and KEGG enrichment analysis of differentially expressed genes was implemented by the clusterProfiler R package, and a corrected $p < 0.05$ were considered as significantly enriched. The PPI network was drawn via the cytoscape 3.4.0 analysis software. All statistical significance was set as $p < 0.05$.

**Author contributions:** Wei X and Chang J have equal contribution to the manuscript. Wei X, Chang J, Cheng Z and Fan Y performed behavioral tests, morphological tests and virus-related experiment. Chen W, Guo H, Liu Z, Mai Y, Hu T, Zhang Y and Cai Q assist the data analysis and organized figures. Fan Y, Chang J and Ge F performed data analysis. Guan X and Fan Y wrote the manuscript. Guan X developed the overall concept.

**Acknowledgments:** This work is supported by National Natural Science Foundation of China (82271531 and 82071495), Natural Science Foundation of Jiangsu Province, China (BK20201398) and Natural Science Foundation of the Higher Education Institutions of Jiangsu Province, China (21KJB360007).

**Competing interests:** The authors declare that they have no competing interests.

**Data availability statement:** The data that support the findings of this study are available from the corresponding author upon reasonable request.

## References

1. Chambers RA, et al. Developmental neurocircuitry of motivation in adolescence: a critical period of addiction vulnerability. *The American journal of psychiatry.* 2003;160(6):1041-52.

2. Spear LP. Consequences of adolescent use of alcohol and other drugs: Studies using rodent models. *Neuroscience and biobehavioral reviews.* 2016;70(228-43.

3. Wong WC, and Marinelli M. Adolescent-onset of cocaine use is associated with heightened stress-induced reinstatement of cocaine seeking. *Addiction biology.* 2016;21(3):634-45.

4. Jordan CJ, and Andersen SL. Sensitive periods of substance abuse: Early risk for the transition to dependence. *Developmental Cognitive Neuroscience.* 2017;25(29-44.

5. Perry JL, et al. Prefrontal cortex and drug abuse vulnerability: translation to prevention and treatment interventions. *Brain Res Rev.* 2011;65(2):124-49.

6. Zhang Y, et al. Molecular changes in the medial prefrontal cortex and nucleus accumbens are associated with blocking the behavioral sensitization to cocaine. *Scientific reports.* 2015;5(1):16172.

7. Koss WA, et al. Dendritic remodeling in the adolescent medial prefrontal cortex and the basolateral amygdala of male and female rats. *Synapse.* 2014;68(2):61-72.

8. Shapiro LP, et al. Differential expression of cytoskeletal regulatory factors in the adolescent prefrontal cortex: Implications for cortical development. *J Neurosci Res.* 2017;95(5):1123-43.

9. Caffino L, et al. A single cocaine administration alters dendritic spine morphology and impairs glutamate receptor synaptic retention in the medial prefrontal cortex of adolescent rats. *Neuropharmacology.* 2018;140(209-16.

10. Caffino L, et al. A single cocaine exposure disrupts actin dynamics in the cortico-accumbal pathway of adolescent rats: modulation by a second cocaine injection. *Psychopharmacology.* 2017;234(8):1217-22.

11. Chocyk A, et al. Early-life stress affects the structural and functional plasticity of the medial prefrontal cortex in adolescent rats. *The European journal of neuroscience.* 2013;38(1):2089-107.

12. Fuhrmann D, et al. Adolescence as a Sensitive Period of Brain Development. *Trends in cognitive sciences.* 2015;19(10):558-66.

13. Shi P, et al. Adolescent cocaine exposure enhances the GABAergic transmission in the prelimbic cortex of adult mice. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology.* 2019;33(7):8614-22.

14. Zhu W, et al. Adolescent cocaine exposure induces prolonged synaptic modifications in medial prefrontal cortex of adult rats. *Brain structure & function.* 2018;223(4):1829-38.

15. Low HB, and Zhang Y. Regulatory Roles of MAPK Phosphatases in Cancer. *Immune network.* 2016;16(2):85-98.

16. Wang X, et al. DUSP1 Promotes Microglial Polarization toward M2 Phenotype in the Medial Prefrontal Cortex of Neuropathic Pain Rats via Inhibition of MAPK Pathway. *ACS Chemical Neuroscience.* 2021;12(6):966-78.

584    17.    Xia L, et al. Rosiglitazone Improves Glucocorticoid Resistance in a Sudden
585           Sensorineural Hearing Loss by Promoting MAP Kinase Phosphatase-1 Expression.
586           *Mediators of inflammation.* 2019;2019(7915730.

587    18.    Taylor DM, et al. MAP kinase phosphatase 1 (MKP-1/DUSP1) is neuroprotective in
588           Huntington's disease via additive effects of JNK and p38 inhibition. *The Journal of*
589           *neuroscience : the official journal of the Society for Neuroscience.*
590           2013;33(6):2313-25.

591    19.    Xu P, et al. DUSP1 alleviates cerebral ischaemia reperfusion injury via inactivating
592           JNK-Mff pathways and repressing mitochondrial fission. *Life Sci.* 2018;210(251-62.

593    20.    Ndong C, et al. Mitogen activated protein kinase phosphatase-1 prevents the
594           development of tactile sensitivity in a rodent model of neuropathic pain. *Molecular pain.*
595           2012;8(34.

596    21.    Jefsen OH, et al. Transcriptional regulation in the rat prefrontal cortex and
597           hippocampus after a single administration of psilocybin. *Journal of*
598           *Psychopharmacology.* 2020;35(4):483-93.

599    22.    Wojcieszak J, et al. Induction of immediate early genes expression in the mouse
600           striatum following acute administration of synthetic cathinones. *Pharmacological*
601           *Reports.* 2019;71(6):977-82.

602    23.    Spear LP. The adolescent brain and age-related behavioral manifestations.
603           *Neuroscience and biobehavioral reviews.* 2000;24(4):417-63.

604    24.    Friedman AL, et al. Effects of adolescent Δ9-tetrahydrocannabinol exposure on the
605           behavioral effects of cocaine in adult Sprague-Dawley rats. *Experimental and clinical*
606           *psychopharmacology.* 2019;27(4):326-37.

607    25.    Brandon CL, et al. Enhanced reactivity and vulnerability to cocaine following
608           methylphenidate treatment in adolescent rats. *Neuropsychopharmacology : official*
609           *publication of the American College of Neuropsychopharmacology.*
610           2001;25(5):651-61.

611    26.    Cass DK, et al. Developmental disruption of gamma-aminobutyric acid function in the
612           medial prefrontal cortex by noncontingent cocaine exposure during early adolescence.
613           *Biological psychiatry.* 2013;74(7):490-501.

614    27.    Takaki M, et al. Two kinds of mitogen-activated protein kinase phosphatases, MKP-1
615           and MKP-3, are differentially activated by acute and chronic methamphetamine
616           treatment in the rat brain. *Journal of neurochemistry.* 2001;79(3):679-88.

617    28.    Lawan A, et al. Skeletal Muscle–Specific Deletion of MKP-1 Reveals a p38
618           MAPK/JNK/Akt Signaling Node That Regulates Obesity-Induced Insulin Resistance.
619           *Diabetes.* 2018;67(4):624-35.

620    29.    Sweatt JD. The neuronal MAP kinase cascade: a biochemical signal integration
621           system subserving synaptic plasticity and memory. *Journal of neurochemistry.*
622           2001;76(1):1-10.

623    30.    Valjent E, et al. Addictive and non-addictive drugs induce distinct and specific patterns
624           of ERK activation in mouse brain. *The European journal of neuroscience.*
625           2004;19(7):1826-36.

626    31.    Radwanska K, et al. Regulation of cocaine-induced activator protein 1 transcription
627           factors by the extracellular signal-regulated kinase pathway. *Neuroscience.*

2006;137(1):253-64.

32.    Sun WL, et al. Molecular Mechanism: ERK Signaling, Drug Addiction, and Behavioral
Effects. *Progress in molecular biology and translational science.* 2016;137(1-40.

33.    Valjent E, et al. Inhibition of ERK pathway or protein synthesis during reexposure to
drugs of abuse erases previously learned place preference. *Proceedings of the*
*National Academy of Sciences of the United States of America.* 2006;103(8):2932-7.

**Figure legends**

**Figure 1. Subthreshold dose of cocaine elicit cocaine CPP and trigger**

**mPFC CaMKII-positive neurons in ACE mice during adulthood. A,**

Experimental models and timeline. **B,** CPP score. **C,** ΔCPP score. **D,** Heatmap

of spent duration by mice in CPP apparatus. **E,** C-Fos staining after CPP test

in mPFC and its subregions of Cg1, PrL, IL. The c-Fos (red) and NeuN (green)

are used as markers for evaluating cell activity and labeling neuron,

respectively. Scale bar, 400 μm/100 μm. **F,** Schematic diagram of viral injection.

**G,** CPP score. **H,** ΔCPP score. **I,** Heatmap of spent duration by mice in CPP

apparatus. **J,** Heatmap of GCaMp6m fluorescence at first 4 sec when mice

entering cocaine-paired chamber. **K,** Quantification of ΔF/F. **L,** The AUC of

ΔF/F. Cg1, cingulate cortex; PrL, prelimbic cortex; IL, infralimbic cortex;

ASE-sdC, adolescent saline-exposed mice with subthreshold dose of cocaine

treatment during adulthood; ACE-sdC, adolescent cocaine-exposed mice with

subthreshold dose of cocaine treatment during adulthood; N.S., $p > 0.05$, [#], $p <$

0.05, [##], $p < 0.01$ vs baseline (pre-test CPP score); N.S., $p > 0.05$, *, $p < 0.05$,

**, $p < 0.01$ vs ASE-sdC mice.

**Figure 2. Subthreshold dose administration of cocaine increase *Dusp1***

**gene while decrease DUPS1 protein and activity in mPFC of adult ACE**

**mice. A,** *Dusp1* mRNA levels on P74. **B,** DUSP1 protein levels and activity on

P74. **C,** DUSP1 activity on P74. **D,** *Dusp1* mRNA levels on P78. **E,** DUSP1

protein levels and activity on P78. **F,** DUSP1 activity on P78. **G,** DUSP1

staining in CaMKII-positive neurons. Scale bar, 400 μm/100 μm. **H,** Levels of

p-ERK/ERK, p-ERK and ERK proteins on P78. ASE, adolescent

saline-exposed mice during adulthood; ACE, adolescent cocaine-exposed

mice during adulthood; ASE-sdC, adolescent saline-exposed mice with

subthreshold dose of cocaine treatment during adulthood; ACE-sdC,

adolescent cocaine-exposed mice with subthreshold dose of cocaine

treatment during adulthood; N.S., $p > 0.05$, *, $p < 0.05$, **, $p < 0.01$ vs ASE

mice or ASE-sdC mice.

**Figure 3. Overexpression of DUSP1$^{CaMKII}$ reduce cocaine-induced CPP**

**and attenuate mPFC activation in ACE mice during adulthood. A,**

Experimental design and timeline. **B,** Schematic diagram of viral injection.

Scale bar, 400 µm. **C,** Percentage of viral transfected neurons in

CaMKII-positive neurons of mPFC. Scale bar, 200 µm/100 µm. **D,** *Dusp1*

mRNA levels on P78. **E,** DUSP1 protein levels on P78. **F,** DUSP1 activity on

P78. **G,** CPP score. **H,** ΔCPP score. **I,** Heatmap of spent duration by mice in

CPP apparatus. **J,** The percentage of c-Fos-positive neurons in

virus-transfected neurons. Scale bar, 200 µm/100 µm. **K,** Levels of

p-ERK/ERK, p-ERK and ERK proteins on P78. ACE-sdC$_{Ctrl}$, ACE-sdC mice

injected with Ctrl virus; ACE-sdC$_{OE}$, ACE-sdC mice with overexpression virus

of DUSP1; N.S., $p > 0.05$, #, $p < 0.05$, ##, $p < 0.01$ vs baseline (pre-test CPP

score); N.S., $p > 0.05$, *, $p < 0.05$, **, $p < 0.01$ vs ACE-sdC$_{Ctrl}$ mice.

**Figure 4. Knockdown of DUSP1$^{CaMKII}$ aggravate cocaine-induced CPP and**

**further activate mPFC activity in ACE mice during adulthood. A,**

Experimental design and timeline. **B,** Schematic diagram of viral injection.

Scale bar, 400 µm. **C,** Percentage of viral transfected neurons in

CaMKII-positive neurons of mPFC. Scale bar, 200 µm/100 µm. **D,** *Dusp1*

mRNA levels on P78. **E,** DUSP1 protein levels on P78. **F,** DUSP1 activity on

P78. **G,** CPP score. **H,** ΔCPP score. **I,** Heatmap of spent duration by mice in

CPP apparatus. **J,** The percentage of c-Fos-positive neurons in

virus-transfected neurons. Scale bar, 200 μm/100 μm. **K,** Levels of

p-ERK/ERK, p-ERK and ERK proteins on P78. ACE-sdC$_{Ctrl}$, ACE-sdC mice

injected with Ctrl virus; ACE-sdC$_{KD}$, ACE-sdC mice with knockdown virus of

DUSP1; N.S., $p > 0.05$, [#], $p < 0.05$, [##], $p < 0.01$ vs baseline (pre-test CPP

score); N.S., $p > 0.05$, *, $p < 0.05$, **, $p < 0.01$ vs ACE-sdC$_{Ctrl}$ mice.

**Figure 5. Schematic summary of the present study.** Adolescent

cocaine-exposed male mice (ACE) models were established by administrating

cocaine of 15 mg / kg once daily during adolescent period. When growing to

adult age, mice were subjected to conditioned place preference (CPP) to

evaluate the sensitivity to cocaine. Subthreshold dose of cocaine (sdC), that is

insufficient to produce CPP, was used to induce CPP in adulthood. The sdC

treatment effectively induced CPP in ACE mice, but not in adolescent

saline-exposed male mice (ASE) during adulthood, indicating higher sensitivity

to cocaine caused by ACE. The sdC treatment triggered more CaMKII-positive

neurons, and induced converse phenotypes of dual specificity phosphatase 1

(DUSP1) molecule in mPFC, as indicated by higher *Dusp1* gene, lower

DUSP1 protein, lower DUSP1 activity and lower DUSP1 expression on

CaMKII-positive neurons (DUSP1$^{CaMKII}$) in mPFC of ACE mice than those of

ASE mice during adulthood. To explore the role of converse phenotypes of

DUSP1 protein and *Dusp1* gene in response to sdC treatment, both

upregulated and downregulated modulations of DUSP1$^{CaMKII}$ were performed

in mPFC of ACE mice. Overexpressing DUSP1$^{CaMKII}$ reduced CaMKII-positive

neuronal activation, and ultimately blocked sdC-induced CPP in ACE mice

during adulthood. While, knocking-down DUSP1$^{CaMKII}$ activated more

CaMKII-positive neurons, and aggravated sdC-preferred behavior in ACE mice

during adulthood. ERK1/2 might be potential subsequent signal for DUSP1 in

the process.

# Figure1

**Figure 2**

# Figure 3

# Figure 4

# Figure 5