Title: T cell responses induced by SARS-CoV-2 mRNA vaccination are
associated with clonal replacement

Authors: Hiroyasu Aoki[1,2], Masahiro Kitabatake[3], Haruka Abe[1], Shigeyuki Shichino[1], Atsushi
Hara[3], Noriko Ouji-Sageshima[3], Toshihiro Ito[3], Kouji Matsushima[1], and Satoshi Ueha[1]*

Affiliations:

[1]Division of Molecular Regulation of Inflammatory and Immune Diseases, Research Institute for
Biomedical Sciences, Tokyo University of Science, Chiba, Japan.

[2]Department of Preventive Medicine, Graduate School of Medicine, The University of Tokyo.

[3]Department of Immunology, Nara Medical University, Nara, Japan.

* Satoshi Ueha, Ph.D., Division of Molecular Regulation of Inflammatory and Immune Diseases,
Research Institute for Biomedical Sciences, Tokyo University of Science, 2669 Yamazaki, Noda
City Chiba, 278-0022, Japan

E-mail; ueha@rs.tus.ac.jp

Abstract: mRNA vaccines against the Spike glycoprotein of severe acute respiratory syndrome
type 2 coronavirus (SARS-CoV-2) elicit strong T-cell responses. However, it is unknown
whether T cell clones induced by the first vaccination or newly generated T cell clones dominate
responses to the secondary vaccination. Here, we analyzed the kinetic profile of Spike-reactive
T-cell clones before the first dose, one week after the first and second dose, and four weeks after
the second dose of the BNT162b mRNA vaccine. Interestingly, a new set of Spike-reactive CD8[+]
T cell clones exhibited the greatest expansion following secondary vaccination and replaced the
clones that had responded to the primary vaccination. Single-cell mRNA/protein/TCR analysis
revealed that the first-responder clones exhibited a terminally differentiated phenotype, whereas
second-responder clones exhibited an actively proliferating phenotype. These results show that
Spike-reactive T cell responses induced by repetitive mRNA vaccination are augmented and
maintained by replacement with newly−generated clones with proliferative potential.

One Sentence Summary: TCR repertoire analysis following mRNA vaccination against SARS-
CoV-2 revealed the replacement of Spike-reactive T cell clones.

Main Text:

INTRODUCTION

mRNA vaccines against Spike glycoprotein of severe acute respiratory syndrome type 2
coronavirus (SARS-CoV-2), including Pfizer-BioNTech's BNT162b and Moderna's mRNA-1273,
are strikingly effective at reducing coronavirus disease 2019 (COVID-19) (*1*). These mRNA
vaccines elicit both humoral and cellular immune responses specific for the viral Spike protein (*2*–
4). Studies on the cellular immune responses following SARS-CoV-2 mRNA vaccination have
demonstrated that the numbers of Spike-reactive T cells increase in the peripheral blood after the
first dose, and are additionally boosted after the second dose (*4*–*6*). The cellular immune response
may be more protective to variants of concerns (VOCs) of SARS-CoV-2 than the humoral
response, because it can target relatively conserved regions of the Spike protein (*7*, *8*).

Previous studies demonstrated that secondary, or boost, vaccinations induce the reactivation of memory CD8$^+$ T cells established following the initial vaccination (*9–12*). However, repetitive antigen stimulation in mice alters the properties of memory T cells, including enhancement of their effector function, and reductions in the proliferation potential and ability to produce IL-2 (*10, 13*). This suggests that memory CD8$^+$ T cells induced by the first vaccination become poorly, or non-responsive in the course of repetitive antigen stimulation. In line with this hypothesis, we previously reported that older memory T cell populations were replaced by newly generated T cell populations in a serial adoptive transfer model characterized by repetitive antigen exposure (*14*). However, in repetitive SARS-CoV-2 mRNA vaccinations, it is not known whether T cell clones induced by the first vaccination or newly generated clones dominate responses to subsequent vaccination.

Pathogen-reactive T cells recognize distinct antigenic epitopes via their unique T cell receptor (TCR). To analyze changes in T cell clonal dominance during SARS-CoV-2 mRNA vaccination, we undertook a longitudinal repertoire analysis by TCR sequencing. This enabled us to comprehensively quantify the changes in the frequencies of specific T-cell clones ("kinetic profile") during primary and secondary responses to mRNA vaccination. Specifically, we performed a longitudinal TCR repertoire analysis on peripheral blood T cells at four time points before and after the BNT162b mRNA vaccination. Spike-reactive T-cell clones were identified by an activation-induced marker assay (AIM assay) using a Spike peptide pool, and their response kinetics were analyzed by tracking their TCRs in longitudinal repertoire data. Furthermore, we matched the kinetics of the clonal response with immunological phenotype using single-cell analysis of gene expression and TCR sequence. These analyses revealed that Spike-reactive T-cell responses induced by secondary mRNA vaccination resulted in the replacement of Spike-reactive T-cell clones from the primary response with new clones that exhibited different response characteristics.

RESULTS

Transient expansion of CD8$^+$ T cell clones was prominent after the second vaccination

To analyze the kinetics of T cell clones following mRNA vaccination to SARS-CoV-2, we studied a cohort of 41 healthcare workers at Nara Medical University (Japan) who had received two doses of the Pfizer-BNT162b2 vaccine (*15*). Peripheral blood mononuclear cell (PBMC) samples were collected from the participants at four points: before the first dose (point 1: P1), seven days after the first dose (P2), seven days after the second dose (P3), and 28 days after the second dose (P4) (Fig. 1A). CD4$^+$ and CD8$^+$ T cells were separated from the PBMC sample, then their TCR repertoire was analyzed ("bulk TCR repertoire analysis").

To identify the kinetic profiles of T-cell clones, we applied weighted gene correlation network analysis (WGCNA) on the longitudinal bulk TCR repertoire dataset (*16*) (Fig. 1D, method). From the kinetic profile of the top 150 clones detected at four points (which occupied about 45% of CD8$^+$ and 10% of CD4$^+$ repertoire on average, fig. S1A), we identified twelve, and ten patterns, of kinetic profile in the CD8$^+$ and CD4$^+$ repertoires (Fig. 1C and fig. S1B-D). CD8$^+$ module Green (+) and CD4$^+$ module Turquoise (+) were the modules that increased at P2 and then decreased, corresponding to a transient expansion after the first dose (hereafter "first-responders"). CD8$^+$ and CD4$^+$ module Brown (+) were the modules that transiently increased at P3 and then decreased (hereafter "second-responders"). CD8$^+$ module Turquoise (+) and CD4$^+$ module Green (+) were modules that increased at P4: four weeks after the second dose (hereafter "long-term responders").

These modules were robustly identified regardless of the number of top CD8$^+$/CD4$^+$ clones used
for WGCNA analysis (fig. S1E and F).

We next classified individual clones within the kinetic profile patterns and calculated their total
frequency at each of the four points (fig. S2A-D). Second and long-term responders seemed to be
dominant in patterns at P3, and P4, respectively. Moreover, the extent of clonal response of each
pattern during vaccination, defined by the maximum changes in the total frequency through P1 to
P4, was more significant in second-responders than first- and long-term responders (first-
responders: 2.52%, second-responders: 12.93%, long-term responders: 5.72%, median) (Fig. 1E).
These results suggest that the CD8$^+$ T cells responded more strongly after the second vaccination
than the first vaccination.

We also analyzed the CD4$^+$ T cell repertoire. First- and second-responders tended to predominate
at P2, and P3, respectively (Fig. 2A and fig. S2E-H). The clonal expansion of first-responders was
greater than that of other patterns, followed by second-responders (first-responders: 11.86%,
second-responders: 8.63%, long-term responders: 2.94%, median) (Fig. 2B). These data suggest
that the clonal response of CD4$^+$ T cells to vaccination was characterized by the greater clonal
expansion after the first dose.

Spike-reactive CD8$^+$ T cell clones were induced after vaccination

Next, we analyzed whether vaccination induced Spike-reactive CD8$^+$ T cells. We utilized the
public database of SARS-CoV-2-reactive TCRs ([17–19]) and searched for biochemically similar
TCRs to our dataset ([20]). In brief, we formed TCR meta-clonotypes for each SARS-CoV-2 epitope
(groups of TCRs sharing similar complementarity determining regions, based on TCRs reported
in the three articles). We then searched for TCRs conforming to these meta-clonotypes in our
dataset. On average, 35 clones per participant belonged to SARS-Cov2-reactive meta-clonotypes.
Furthermore, the frequency of clones belonging to each SARS-CoV-2 protein meta-clonotype was
determined (Fig. 3A). The total frequency of Spike-reactive meta-clonotypes was significantly
increased at P2 and P3 (Fig. 3B). On the other hand, the total frequency of meta-clonotypes
reactive to other SARS-CoV-2 proteins remained almost constant following vaccination (Fig. 3B).
This result demonstrated that mRNA vaccination induced a CD8$^+$ T cell response to Spike, but not
to other SARS-CoV-2 proteins.

We next investigated whether individual CD8$^+$ T cell clones with Spike-reactive meta-clonotypes
were expanded after vaccination. Of 1,421 clones in Spike-reactive meta-clonotypes in 41
participants, only 140 clones were detected at more than two points. The distribution of WGCNA
modules (Fig. 1C) among these 140 clones was assessed and compared to that among clones in
other meta-clonotypes. This comparison revealed that first- and second-responders were enriched
among clones with Spike-reactive meta-clonotypes, while long-term responders were not (Fig. 3C).
Moreover, the module Yellow (-), which peaked at P3 like second-responders, was also enriched
in Spike-reactive meta-clonotypes. These three modules occupied about 60% of clones with Spike-
reactive meta-clonotypes. This result suggested that Spike-reactive T cell clones reported in the
public database were also expanded in our cohort after the first or second vaccination.

Spike-reactive CD8$^+$ T-cell clones were replaced following secondary vaccination

To detect Spike-reactive clones that were not covered in the public database, we analyzed the
repertoire of Spike-reactive T cells in our participants. For this purpose, an activation-induced
marker (AIM) assay was performed on 15 participants ([5, 21–23]). PBMCs collected at P3 were

stimulated with peptide pools of the SARS-CoV-2 Spike protein. Then, CD137$^+$ or CD107a$^+$ non-
naïve CD8$^+$ T cells, and CD40L$^+$ CD69$^+$ non-naive CD4$^+$ T cells were sorted as AIM$^+$ Spike-
reactive cells (Fig. 4A). We sequenced the TCRs of AIM$^+$ and AIM$^-$ non-naive T cells and
identified AIM$^+$ CD8$^+$ and CD4$^+$ T cell clones (fig. S3A). Then, we excluded Spike-nonreactive
clones that became AIM+ during co-culture ("by-stander" clones) based on their enrichment in the
AIM$^+$ repertoire relative to the AIM$^-$ repertoire (Fig. 4B and C, and fig. S3B and C). Notably, the
total frequency of AIM$^+$ clones is significantly higher than the proportion of AIM$^+$ cells among
CD4$^+$ or CD8$^+$ T cells (fig. S3D). This indicated that Spike-reactive clones included T cells that
did not become AIM$^+$, which is consistent with the fact that the AIM assay underestimates the
frequency of Spike-reactive T cells (2). Next, these AIM$^+$ clones were tracked in the whole TCR
repertoire dataset from P1 to P4. We observed that the total frequency of AIM$^+$ clones increased
at P3 and then decreased at P4 (Fig. 4D and fig. S3E), which is consistent with the kinetics of
Spike-reactive T cells through vaccination reported in the previous studies (4–6).

To analyze the kinetic profile of AIM$^+$ clones, we assigned AIM$^+$ and AIM$^-$ clones into WGCNA
modules as in Fig.1 and compared the proportion of each module (Fig. 4E, left). This classification
revealed that the proportion of second-responders was significantly higher in AIM$^+$ clones than in
AIM$^-$ ones, while first- and long-term responders were not (Fig. 4E, right). In the CD8$^+$ T cell
repertoire, the expansion of second-responders was remarkable. Thus, we investigated whether
AIM$^+$ clones were enriched in second-responders. The proportion of AIM$^+$ clones among second-
responders at P3 varied between participants (0.4% - 70.8%); however, the participants where the
frequency of AIM$^+$ clones at P3 was higher than the median ("AIM-high participants") exhibited
a significantly higher proportion of AIM$^+$ clones among second-responders (fig. S3D and Fig. 4F).
On the other hand, in the AIM-low participants, the frequency of AIM$^+$ second-responder clones
was low, irrespective of the total frequency of second-responders (Fig.4G). These results suggested
that the expansion of AIM$^+$ clones as second-responders was necessary to induce greater anti-
Spike CD8$^+$ T cell response following the vaccination.

Next, we investigated whether second-responders also predominated among the AIM$^+$ clones at
other time points (Fig. 4H and fig. S4A-D). As expected, the total frequency of AIM$^+$ second-
responders was the highest in nine out of the fourteen (9/14) participants at P3 (fig. S3B). However,
at P2, the AIM$^+$ second-responders tended to be less dominant than AIM$^+$ first- and long-term
responders (fig. S3B). In addition, at P4, AIM$^+$ second-responders was nearly equivalent to AIM$^+$
long-term responders (+) (fig. S3D). These results suggest that AIM$^+$ second-responders were not
dominant before, or four weeks after, the second vaccine dose. Moreover, at the clonal level, only
51% and 67% of the top 10 most expanded AIM$^+$ clones at P3 remained top 10 at P2 and P4,
respectively. Namely, half of the top 10 AIM$^+$ clones were replaced between P2 and P3, and one-
third were replaced between P3 and P4 (Fig. 4I). In summary, these results suggest that the increase
in Spike-reactive CD8$^+$ T cells by vaccination was accompanied by the replacement of T cell clones
between the first and the second dose.

We also analyzed the kinetics of AIM$^+$ clones in the CD4$^+$ T cell repertoire. The proportion of
first- and second-responders was significantly higher in AIM$^+$ clones (Fig. 5A). Moreover, AIM-
high participants showed a higher proportion of AIM$^+$ clones among first- and second-responders
at P2 or P3, as observed in the CD8$^+$ repertoire (Fig. 5B). However, the proportion of AIM$^+$ clones
was much smaller in the CD4$^+$ repertoire than in CD8$^+$ (-10% for first-responder and -40% for
second-responder). The dominant patterns of AIM$^+$ clones were replaced between time points in
the CD4$^+$ repertoire, as observed in CD8$^+$ (Fig. 5C and fig.S4E-H). Furthermore, at the clonal level,

only 35 % and 45 % of the top 10 most expanded AIM$^+$ clones at P3 remained top 10 at P2 and
P4, respectively (Fig. 5D). These results indicate that CD4$^+$ Spike-reactive clones were also
replaced after the first and the second vaccination.

First- and second-responder clones differ in their phenotype after the second vaccination.

From TCR repertoire analysis, we have determined that the T-cell clones that expanded after the
first vaccination ("first-responder") were different from those expanded after the second
vaccination ("second-responder"). This observation suggests that the first-responder clones could
not expand as much as second-responder clones after the second vaccination. However, it remained
unclear whether the phenotype of first-responders after the second vaccination differed from that
of second-responders. To address this question, the gene-expression profiles of first- and second-
responder clones after the second dose was analyzed. We performed targeted single-cell RNA-seq
(scRNA-seq) combined with single-cell TCR sequencing (scTCR-seq) and protein expression
analysis using oligo-conjugated antibodies (AbSeq) on non-naive CD4$^+$ and CD8$^+$ T cells collected
at P3 from the two participants (ID2 and ID27) (fig. S5A). Unsupervised clustering using
integrated data of mRNA and surface protein (*24*) identified six clusters in non-naive CD8$^+$ T cells
(Fig6A-C, and fig. S5B-D). Clusters 2 and 3 (terminally differentiated: TD) were characterized by
high expression of *GZMB* and CD45RA (*25, 26*), and cluster 0 (innate-like: IL) expressed *ZBTB16*
and CD161 (*27*). Semi-invariant TCRα chains are typically present in mucosal-associated invariant
T cells (MAIT) (*28–30*), TRAV1-2/TRAJ33, TRAV1-2/TRAJ20, or TRAV1-2/TRAJ12,
composes about 67 % of TCRα-detected cells in this cluster 0 (fig. S5E). Cluster 1 (activated: Act)
contained *TYMS*$^+$ proliferating cells and HLA-DR$^+$ activated cells. Compared to cluster 4 (memory
precursor: MP), cluster 5 expressed a lower level of CD127 and CCR7 and a higher level of *GZMK*
and CD279 (PD-1), consistent with the progenitor exhausted-like (PEX) phenotype described
previously (*31*). The clonality of these CD8$^+$ clusters showed that TD and PEX clusters contained
larger clones (fig. S5E), suggesting that these clusters underwent clonal expansion due to
activation. Notably, the cluster distribution of CD8$^+$ T cells differed between the two participants:
cluster TD2 and PEX were mainly present in ID2, while cluster TD1 was predominantly present
in ID27 (Fig. 6D).

To analyze the relationship between T-cell clones' kinetic profile and their phenotype in each T
cell clone, T cells in the first-, second, and long-term responder clones were extracted from the
dimensionality reduction plot of single-cell analysis. The second-responder clones mainly
belonged to the cluster Act. (90.7 % in ID2 and 90.2 % in ID27) (Fig. 6E and F). On the other
hand, the first-responder clones were predominantly composed of cluster TD2 (79.4% in ID2), and
very few belonged to cluster Act. Furthermore, we confirmed these relationships at the clonal level.
The heatmap of cluster distribution of CD8$^+$ clones exhibited that T cells in each clone were highly
skewed to one phenotype, suggesting that the differentiation stage of CD8$^+$ T cells was highly
synchronized in clones (Fig. 6G). Moreover, all second-responder clones skewed to cluster Act,
while the two first-responder clones skewed to cluster TD2 (Fig. 6G). These results suggested that
first-responders became terminally differentiated after the second dose, which led them to be
unresponsive and outcompeted by actively proliferating second-responders.

Then, we examined the phenotype of Spike-reactive clones. AIM$^+$ and AIM$^-$ T cell clones
determined in Fig. 4 and 5 were projected onto the dimensionality reduction plot, and their
distribution to clusters was compared. CD8$^+$ T cells of AIM clones were enriched in cluster Act in
participant ID2 and IL in ID27 (Fig. 6H and I). Moreover, we examined the phenotype of expanded
AIM+ clones with more than ten cells. At the clonal level, two AIM$^+$ clones skewed to cluster Act,

two clones skewed to IL, and the remaining three clones were a mixture of Act and other clusters (Fig. 6J). Interestingly, the two Act-skewed AIM$^+$ clones exhibited the second-responder pattern, while the frequency of the two IL-skewed AIM$^+$ clones remained stable at P2 and P3 (Fig. 6K). Although the number of expanded AIM$^+$ clones analyzed was small, these results indicate that Spike-reactive second-responder clones were actively proliferating after the second dose. In contrast, Spike-reactive clones with innate-like phenotypes were not responsive to mRNA vaccination.

Single-cell TCR analysis was also performed on CD4$^+$ T cells. Four clusters were identified in non-naive CD4$^+$ T cells; CCR7$^+$ CD62L$^+$ cluster 0 (central memory: CM), CD196 (CCR6)$^+$ CD127$^+$ cluster 1 (effector memory: EM), *IFNG*$^+$HLA-DR$^+$ cluster 2 (activated: Act), and *FOXP3*$^+$ CD25$^+$ cluster 3 (regulatory T cells: Treg) (Fig. 7A-C and fig. S6A-C). The higher clonality of cluster 2 supported that this cluster underwent activation-induced clonal expansion (fig. S6D). Notably, T cells in AIM$^+$ clones were mainly composed of cells in the EM cluster and rarely contained cells in Act or Treg clusters, suggesting that the phenotype of AIM$^+$ CD4$^+$ T cells was more homogeneous than AIM$^+$ CD8$^+$ (Fig. 7D and E). We also attempted to analyze the relationship between the kinetics of CD4$^+$ T cell clones and their phenotypes. However, clear differences in cluster distribution between first-, second-, and long-term responders were not observed (Fig. 7F and G).

DISCUSSION

Through a longitudinal TCR repertoire analysis combined with an AIM assay using a Spike peptide pool, we demonstrated that the increase of Spike-reactive CD8$^+$ T cells following secondary vaccination was caused by second-responder clones, which replaced the first-responders. Moreover, single-cell analysis indicated that terminally differentiated first-responder clones were outcompeted by actively proliferating second-responders after the second vaccine dose. These results suggest that the vaccine-induced anti-Spike T-cell response was accompanied by clonal replacement of Spike-reactive T cells. These data are consistent with a dynamic turnover model of T-cell memory response under repetitive antigen challenges in humans receiving SARS-CoV-2 mRNA vaccinations (*14*).

There are several hypotheses to explain the mechanism underlying the clonal replacement of memory T cells following repetitive vaccination. One hypothesis is that first-responders are memory T cells cross-reactive to 'common cold' coronaviruses (CCCoV) that were present before the pandemic (pre-existing memory) (*32–35*). After the first dose, these pre-existing memory T cells might undergo a proliferative burst and differentiate into secondary memory with a weaker proliferation ability (*10, 13, 36*). Another possible mechanism is that the first-responders have high-affinity TCRs to Spike, compared to second-responders. Signals from the high-affinity TCRs of first-responders may induce greater expansion and effector differentiation after the first dose, resulting in reduced proliferative capacity to subsequent antigen exposure. A third possible mechanism is that second-responder clones are those that migrated into the vaccine-draining lymph node (dLN) relatively late during the primary response ("late-arriving" clones). Carton et al. demonstrated that late-arriving T cells divide less than early-arriving T cells during the first antigen challenge and their progeny proliferate more extensively during the second response (*37*). These possibilities could be addressed in future studies.

The strength of our dataset is that the sampling time point for bulk TCR repertoire analysis is uniform across the participants. As far as we know, this is the first report of longitudinal TCR

repertoire data before and after the mRNA vaccination with 41 participants, enabling the global
analysis of the kinetics of T-cell clones. Moreover, all participants reported no history of SARS-
CoV-2 infection (*15*). Thus, our data represent the exclusive effect of mRNA vaccination on T-
cell repertoire, although we can't exclude that some participants may have been asymptomatically
infected. Furthermore, we introduced the WGCNA (*16*) for the analysis of T-cell clones' kinetics,
originally used for the time-course transcriptional data to identify gene clusters associated with
biological events (*38, 39*). Compared to the earlier methods for identifying expanded/contracted
clones between two time points (*40, 41*), we believe that WGCNA identifies the kinetic patterns
of clones more flexibly.

To evaluate Spike-reactive T cell responses of the participants in our cohort, we integrated AIM
assays and TCR repertoire analyses. The AIM assay is widely used to evaluate the Spike-reactive
T cell responses (*5, 21–23*). However, this assay underestimates the frequency of Spike-reactive
T cells compared to the use of peptide-major histocompatibility complex (pMHC) multimers (*2*).
The low sensitivity of the AIM assay may be because only a tiny proportion of Spike-reactive
clones become AIM$^+$, as suggested by fig. S3F. Thus, the inclusion of TCR sequencing can
improve the sensitivity of the AIM assay. Moreover, considering that the pMHC multimer assay
requires prior knowledge of participants' HLA allele, the AIM assay integrated with TCR
sequencing we performed in this study is a more suitable method to comprehensively detect
cellular immune response induced by mRNA vaccination. Through this approach, we identified
participants who exhibited a small number of AIM$^+$ second-responders, while the total amount of
second-responders was relatively high. For these participants, the epitopes recognized by Spike-
reactive clones may not have been efficiently presented on HLA from the Spike peptide pool used
in the AIM assay. Expanding the Spike peptide pool to cover a broader range of possible epitopes
may fill the gap between the total and AIM$^+$ second-responders in these participants.

One limitation in our study is the low sensitivity of the AIM assay. The frequency of AIM$^+$ CD8$^+$
T cells is reported to be as high as 1% of the non-naive fraction during the vaccination (*5, 6*). To
collect a sufficient number (more than 100 cells) of cells for TCR sequencing, we defined AIM$^+$
CD8$^+$ T cells as expressing either CD137 or CD107a. However, this definition decreased the
specificity of AIM$^+$ CD8$^+$ clones compared to the analysis of CD4$^+$ clones. Moreover, performing
AIM assays at time points other than P3 (7 days after the second dose) is difficult because the AIM
positivity is lower than that in P3. Another limitation is the limited number of participants in whom
single-cell analysis was performed. Notably, the number of AIM$^+$ clones detected in the single-cell
analysis is too small to demonstrate the relationship between their response kinetics and
phenotypes after the second dose.

An important question is whether the clonal replacement we demonstrated in this study is specific
to the mRNA vaccination or is common to the other types of vaccination (peptide, inactivated
virus, etc.) or natural infections. We initially reported the dynamic turnover model through antigen
delivery systems using plasmid and viral vectors (*14*). In addition, in the case of yellow fever
vaccination, the responding CD8$^+$ clones are replaced between the prime and boost dose (*42*).
These observations suggest clonal replacement is a general phenomenon in establishing T-cell
memory through vaccination. It should also be noted that the current SARS-CoV-2 mRNA vaccine
strongly activates Spike-reactive CD8$^+$ T cells through the production of IFNα (*43*), which may
accelerate clonal replacement compared to other types of vaccination. Although the current study
analyzed only peripheral blood T cells, natural SARS-CoV-2 infection generates tissue-resident
memory T cells (TRM) in the lung (*44*). Considering that a subset of circulating effector T cells

have a higher capacity to form TRM (*45*), the replacement of pathogen reactive clones in blood after re-infection may be associated with the re-distribution of older clones from the blood to peripheral tissues and their differentiation to TRM. Moreover, it is unknown whether these TRM clones are also replaced under re-infection or continue to function as a defense against re-infection. Collectively, a systemic analysis of peripheral tissues and blood is necessary to understand the clonal replacement by natural infection.

As of August 2022, many countries implemented the third dose of mRNA vaccination, and several countries, including Japan, started offering fourth-dose vaccines for healthcare workers and the elderly. The third- and fourth- vaccination induces a more robust humoral and cellular immune response to SARS-CoV-2 (*46*, *47*), and increases the protection against SARS-CoV-2 infection, including the Omicron variant (*48*, *49*). However, considering the maintenance of memory T-cell populations by clonal replacement, which requires the mobilization of new clones, memory populations may eventually become unsustainable after repetitive vaccination, especially in elderly participants whose naive T-cell repertoire is decreased. Thus, the sustainability of clonal replacement under repetitive vaccination needs to be verified based on the participants' age and proportion of naive T cells in the future study. We believe that the comprehensive analysis of T cell clones like this study will provide the immunological basis for optimizing the strategy for mRNA vaccination.

In summary, we demonstrated the replacement of memory T-cell clones during SARS-CoV-2 mRNA vaccination. In contrast to humoral immunity mediated by memory B cells and long-lived plasma cells, where the memory response to the first viral variant encountered in childhood is preferentially employed even after the challenge of other variants in adults (original antigenic sin) (*50*, *51*), T cells appear to be able to replace memory population after repetitive challenges. This "re-writability" of T cell immunological memory enables them to effectively generate responses to variant strains of the virus.

MATERIALS AND METHODS

Study Design

Forty-one healthcare workers (≥20 years old) at Nara Medical University in Japan who had received the Pfizer-BNT162b2 vaccine between 10 March and 30 March 2021 were invited to participate as volunteers in the study. The Nara Medical University Ethics Committee approved the study (No. 3168). All study participants provided written informed consent. All participants had no history of suspected clinical SARS-CoV-2 infection. Peripheral blood samples were collected before receiving the first dose (P1), 1 week after receiving the first dose (P2), then 1 week (P3), 1 month (P4), 3 months, and 6 months after receiving the second dose. PBMCs were isolated from whole blood with BD vacutainer CPT tubes (BD), resuspended in CELLBANKER 1 (Takara), and stored at -80°C. The detailed study design was described in Kitabatake et al. (*15*).

Evaluation of anti-SARS-CoV-2 antibodies

All serum samples from vaccinated participants were tested for anti-RBD antibodies with a COVID-19 human IgG enzyme-linked immunosorbent assay kit per the manufacturer's instructions (RayBiotech Life, Inc. Peachtree Corners, GA, USA).

T-cell isolation from PBMCs

$CD4^+$ and $CD8^+$ T cells were magnetically isolated from PBMC. PBMC were thawed by warming frozen cryovials in a 37°C water bath. Cells were stained for 20 minutes at 4°C with 1^{st} antibody

mix; Human TruStain FcX (BioLegend), anti-CD3-Pacific Blue (clone SK7, BioLegend), anti-CD4-PE (clone SK3, BioLegend), anti-CD8-APC (clone SK1, BioLegend), and anti-CD19-APC (clone HIB19, BD Biosciences). After washing with MACS buffer [0.5% BSA (Nacalai Tesque Inc.), 2 mmol/L EDTA (Nacalai Tesque Inc.) in PBS (Nacalai Tesque Inc.)], Cells were stained for 20 minutes at 4°C with 2nd antibody mix; BD IMag anti-PE Magnetic Particles-DM (Clone E31-1459, BD) and anti-APC-Biotin (clone APC003, BioLegend). Then, the cell suspension was diluted with MACS buffer and set on a magnet plate (DynaMag96SideSkirted) (Thermo Fisher Scientific). The positive fraction was collected as $CD4^+$ T cells. The negative fraction was stained with BD IMag Streptavidin Particles Plus-DM (BD) for 20 minutes at 4°C. Then, the cell suspension was diluted with MACS buffer and set on a magnet plate. The positive fraction was collected as $CD8^+$ T cells and B cells. The counts of $CD4^+$ T cells, $CD8^+$ T cells, and B cells were determined using Flow-Count fluorospheres (Beckman) and a Cytoflex flow cytometer (Beckman).

AIM assay and sorting Spike-reactive T cells.

AIM assay was performed on P3 PBMC samples of 15 participants with varying levels of anti-RBD antibody (Table S1). PBMC were thawed by warming frozen cryovials in a 37°C water bath and resuspended in 10mL of in Culture Medium (CM) [10% fetal bovine serum (MP Bio Japan), 1% HEPES (Nakarai) and 1% penicillin/streptomycin (Nakarai) in RPMI-1640 medium (Nakarai)]. Cells were resuspended in 1mL of CM, then plated in 24-well plates and incubated overnight (about 16 hours) in an incubator at 37°C, with 5% CO2. After incubation, cells were then stimulated for 24 hours in stimulation medium [BD FastImmune CD28/CD49d Costimulatory Reagent (final x100, BD Biosciences) and anti-human CD40 (final concentration of 0.5μg/mL, clone HB14, BioLegend) in CM] with or without PepTivator SARS-CoV-2 Prot_S Complete (final concentration of 1 mg/mL, Miltenyi). 20 hours post-stimulation, an anti-CD107a-FITC antibody (final concentration of 5μg/mL, clone H4A3, BioLegend) was added to the culture. After four hours, cells were washed in PBS supplemented with 10% FBS (10% FBS/PBS). Cells were stained for 20 minutes at 4°C with antibody mix against CD4 (clone SK3, BioLegend), CD8 (clone RPA-T8, BioLegend), CD14 (clone 63D3, BioLegend), CD19 (clone HIB19, BioLegend), CD45RA (clone HI100, BioLegend), CD27 (clone O323, BioLegend), CD40L (clone 24-31, BioLegend), CD69 (clone FN50) and CD137 (clone 4B4-1, BD). Cells were washed once in 10% FBS/PBS. AIM^+ non-naïve $CD4^+$, AIM^- non-naïve $CD4^+$, AIM^+ non-naïve $CD8^+$, and AIM^- non-naive $CD8^+$ cells were sorted using FACS Aria II or Aria III (BD Biosciences). AIM^+ $CD4^+$ T cells were defined by dual-expression of CD69 and CD40L. AIM^+ $CD8^+$ T cells were defined by expression of either CD107a or CD137. Nonviable cells were excluded from the analysis based on forward and side scatter profiles and propidium iodide staining. Data were analyzed using FlowJo software (version 10.8.1; BD Biosciences).

Sample preparation for single-cell analysis

PBMC were thawed as described above, and $2x10^6$ cells per sample were used for single-cell analysis. 2x AbSeq master mix of Immune Discovery Panel (BD Biosciences) prepared as the provider's protocol. Cells were resuspended with Cell Staining Buffer (BioLegend) supplemented with Human TruStain FcX (BioLegend). After 2-3 minutes, the cell suspension was added with 2x AbSeq master mix and stained for 15 minutes on ice. Then, the antibody mix for cell sorting was added to the cell suspension and left on ice for 15 minutes for staining. Antibody mix for cell sorting contained following antibodies; anti-CD3 (clone SK7, BioLegend), CD4 (clone SK3, BioLegend), CD8 (clone SK1, BioLegend), CD14 (clone 63D3, BioLegend), CD19 (clone HIB19, BioLegend), CD45RA (clone HI100, BioLegend), and CD27 (clone O323, BioLegend). Stained

cells were washed with Cell Staining Buffer. Then non-naive CD4$^+$ and CD8$^+$ cells were sorted using FACS Aria II or Aria III (BD Biosciences). Data were analyzed using FlowJo software (version 10.8.1; BD Biosciences).

Library preparation and sequencing for TCR repertoire analysis

TCR sequencing libraries for next-generation sequencing were prepared according to the previous report (*52*). In brief, mRNA in T-cell lysate was captured by 10 µL of oligo-dT immobilized Dynabeads M270 streptavidin (Thermo Fisher Scientific). mRNA-trapped beads were suspended in 10 µL of RT mix and incubated for 60 min at 42ºC to perform reverse transcription (RT) and template-switching. The composition of the RT mix was as follows; 1× First Strand buffer (Thermo Fisher Scientific), 1 mM dNTP (Roche), 2.5 mM DTT (Thermo Fisher Scientific), 1 M betaine (Sigma-Aldrich), 9 mM MgCl2 (NIPPON GENE), 1 U/µL RNaseIn Plus RNase Inhibitor (Promega), 10 U/µL Superscript II (Thermo Fisher Scientific), and 1 µM of trP1-TSO. To amplify the TCR cDNA containing complementarity determining region 3 (CDR3), nested PCR of the TCR locus was performed, followed by the purification of PCR product using an AM Pure XP kit (Beckman Coulter) at a 0.7:1 beads-to-sample ratio. The third PCR was performed to amplify TCR libraries and add adaptor sequences for the next-generation sequencer. The third-PCR products were purified as second PCR. Final TCR libraries, whose lengths were about 600 base pairs, were sequenced using an Illumina Novaseq 6000 S4 flowcell (67 bp read 1 and 140 bp read 2) (Illumina). Only read2 contained the sequence regarding the definition of T-cell clones. Oligonucleotides used and conditions for PCR reaction and purification for library preparation were summarized in Table S2 and S3, respectively.

Library preparation and sequencing for single-cell analysis

Live cells were stained with Calcein AM (Nacalai) and counted using Flow-Count fluorospheres (Beckman) and a Cytoflex flow cytometer. About 25,000 - 30,000 labeled cells were trapped and reverse-transcribed using the BD Rhapsody™ (BD Biosciences), Targeted mRNA, and AbSeq Amplification Kit (BD Biosciences), and Immune Response Panel HS (BD Biosciences) according to the manufacturer's instructions. Targeted mRNA, TCRseq, and Abseq Index libraries were prepared according to the manufacturer's instructions with the following modifications. (1) We used the original set of primers for TCR amplification, as summarized in Table S2. (2) We performed additional PCRs to introduce a unique dual index for sequencing (3rd PCR for mRNA and TCR) or to enrich the desired product (5th PCR). (3) We used AMPure XP (Beckman-Coulter) or ProNex Size-Selection DNA purification System (Promega) for the purification of PCR products, as summarized in Table S4. Amplified libraries were quantified using a KAPA SYBR Fast qPCR Kit (KAPA Biosystems), and size distribution was analyzed by Microchip Electrophoresis System MultiNA (Shimadzu). Targeted RNAseq, TCRseq, and AbSeq libraries were sequenced on an Illumina Novaseq 6000 S4 flowcell (67 bp read 1 and 140 bp read 2) (Illumina) to a depth of approximately 20,000 reads, 7,000 reads, and 10,000 reads per cell, respectively.

Data processing of TCR sequencing reads

Data processing of TCRseq was performed as previously reported (*52*). Briefly, Adapter trimming and quality filtering of sequencing data were performed using Cutadapt-3.2 (*53*) and PRINSEQ-0.20.4 (*54*). Sequencing data were processed by MiXCR-3.0.5 (*55*). In MiXCR, filtered reads were aligned to reference mouse TCR V/D/J sequences registered in the international ImMunoGeneTics (IMGT) information system. The Variable (V) and Joining (J)

segments of TCRs were represented in IMGT gene nomenclature. T-cell clones were determined
as TCR reads with the same TCR V segment, J segment, and CDR3 nucleotide sequence.

For bulk TCRseq data, we regarded the CD4$^+$-CD8$^+$ overlapping clones in the same participant
as contamination during T-cell isolation from PBMC because CD4$^+$ and CD8$^+$ T-cell repertoire
are mutually exclusive (*56*). Thus, these overlapping clones were categorized into either CD4$^+$ or
CD8$^+$ according to their dominance in CD4$^+$ and CD8$^+$ repertoire using VDJtools-1.2.1 (*57*). The
sequencing reads of the sample were normalized to the cell count (bulk TCR sequencing data) or
16 times the cell count (AIM assay-TCRseq data) in each sample by the "DownSample"
command of VDJtools-1.2.1 (*57*). For AIM assay-TCRseq data, TCR clones with 15 or fewer
reads were regarded as sampling noise and removed from the dataset (*56*). The processed data
have been deposited at the NCBI GEO; accession GSE210227 and GSE210228.

Grouping T-cell clones based on their kinetic profile using WGCNA

To identify the pattern of clones' kinetic profile, we performed WGCNA (*16*) on the top 150 clones
detected at all four time points for each participant (Fig. 1B). First, the correlation network of
kinetics profile of these selected clones (150 clones x 41 participants) was generated, and modules
consisting of highly correlated clones were identified. We identified six and five modules by
WGCNA for CD8$^+$ or CD4$^+$ repertoire, respectively. Next, the module's eigenvector, the first
principal component of the kinetic profile of clones in the module, was calculated. Components of
the eigenvectors were the standardized frequency at each time point. Then, the kinetics patterns
were defined as the eigenvectors or inverted eigenvectors (multiplied by -1) of modules identified
(Fig. 1C and fig. S1D).

We classified the clones detected at least two time points into one of these kinetics patterns. The
criteria for assigning a clone to a particular kinetic pattern were the followings; the correlation
coefficient between the kinetic pattern and the clone's kinetic profile was (1) higher than 0.85 and
(2) higher than that between other kinetic patterns. Clones that did not meet these criteria for
kinetic patterns were regarded as "unassigned."

Searching TCRs with reported Spike-reactivity using tcrdist3

To search previously reported SARS-Co-2 reactive TCRs in our dataset, we utilized the framework
introduced in Mayer-Blackwell's study (*20*). In this framework, we first formed meta-clonotypes
of public CoV-2-reactive TCRs (groups of TCRs that were biochemically similar). Then we
searched TCRs conformant to the definition of the meta-clonotypes (meta-clonotype conformant
TCRs).

Public CoV-2-reactive TCRs were curated from the three datasets (*17–19*). Francis's and
Minervina's datasets were obtained by sorting SARS-CoV-2 antigen peptide-MHC multimer
positive cells followed by single-cell TCR sequencing. Snyder's dataset was obtained by antigen
stimulation following TCR sequencing (MIRA assay) (*58*). For Francis and Minervina's dataset,
public CoV-2-reactive TCRs were grouped by their epitopes, then meta-clonotypes were formed
for each group using tcrdist3, according to their instruction. In forming meta-clonotypes, optimal
radii were estimated at which background TCRs were detected at a frequency of $< 10^{-6}$. In addition,
for Snyder's dataset, the meta-clonotypes reported in Meyer-Blackwell's study were used.

We defined meta-clonotype conformant TCRs as TCRs that met the criteria of meta-clonotype
defined by an optimized radius and the CDR3 motif constraint. Meta-clonotype conformant TCRs
were searched using Snyder's, Francis's, and Minervina's datasets independently, then assigned

with the type of SARS-CoV-2 antigen corresponding to the meta-clonotypes ("Spike,"
"Membrane," etc.). If the assigned antigen types were inconsistent between datasets, the
assignment based on Francis's and Minervina's datasets were adopted because these datasets were
generated by sorting SARS-CoV-2 reactive T cells using peptide-MHC tetramer.

Defining Spike-reactive TCRs using TCRseq data from the AIM assay

We detected Spike-reactive clones using the dataset of AIM-assay TCRseq as in Fig. 4B. T-cell
clones detected only in the AIM^+ repertoire were regarded as Spike-reactive clones. For T-cell
clones detected in AIM+ and AIM^- repertoire, we calculated the "AIM+ enrichment ratio" for each
clone by the following equation;

$$AIM\ enrichment\ ratio\ of\ clone\ X = \frac{AIM^+\ cell\ count\ in\ clone\ X/_{AIM^-\ cell\ count\ in\ clone\ X}}{Total\ AIM^+\ cell\ count/_{Total\ AIM^-\ cell\ count}},$$

where total AIM+ and AIM^- cell count were the count of sorted cells in AIM assay, and AIM^+ and
AIM^- cell count in clone X were calculated as the value obtained by multiplying total cell count
and clone's frequency in AIM^+ or AIM^- repertoire. Then, clones with an AIM^+ enrichment ratio
over 4 were also defined as Spike-reactive clones.

Data processing of scRNAseq reads

Data processing of scRNAseq reads was performed as described in Shichino et al. (*59*). Adapter
trimming of sequencing data was performed by using cutadapt 3.4 (*53*). Filtered reads were
chunked into 16 parts for parallel processing using Seqkit 0.15.0 (*60*). Filtered cell barcode reads
were annotated by Python script provided by BD with minor modifications to be compatible with
Python3.8. Associated cDNA reads were mapped to reference RNA (build GRCh38 release-104)
using bowtie2-2.4.2 (*61*) by the following parameters: -p 2 -D 20 -R 3 -N 0 -L 8 -I S,1,0.75 –norc
–seed 656565 –reorder. Then, cell barcode information of each read was added to the bowtie2-
mapped BAM files, and read counts of each gene in each cell barcode were counted by using
mawk. Resulted count data was converted to genes x cells matrix file, and the knee-plot inflection
threshold was detected using the DropletUtils package in R version 4.1.2 (*62*). In addition, we
further estimate background beads using the emptyDrops formula in the DropletUtils package.
Cells of total read count over inflection point were considered valid cells.

Data processing of AbSeq reads

For AbSeq data, adapter trimming of sequencing data was performed using cutadapt 3.4 (*53*).
Filtered reads were chunked into 64 parts for parallel processing using Seqkit 0.15.0 (*60*). Filtered
cell barcode reads were annotated by Python script provided by BD with minor modifications to
be compatible with Python3.8. Associated AbSeq reads were mapped to known barcode fasta by
using bowtie2-2.4.2 (*61*) by the following parameters: -p 2 -D 20 -R 3 -N 0 -L 8 -I S,1,0.75 –norc
–seed 656565 –reorder–trim-to 3:39 –score-min L,-9,0 –mp 3,3 –np 3 –rdg 3,3. Then, cell barcode
information of each read was added to the bowtie2-mapped BAM files, and read counts of each
AbSeq tag in each cell barcode were counted using mawk. Resulted count data were converted to
tags x cells matrix file by using the data.table package in R version 4.1.2, and top 1M cell barcodes
were extracted.

scRNAseq/AbSeq data analysis

scRNAseq/AbSeq data analysis was performed using the R software package Seurat v4.0.1 (*24*).
Expression data of targeted panel genes and surface proteins were converted to the Seurat object.

CD4$^+$ CD8$^-$ and CD4$^-$ CD8$^+$ cells were extracted based on the expression level of CD4 and CD8
proteins determined by AbSeq, as fig. S5A. Then, expression data was log-normalized and scaled
using NormalizeData and ScaleData function, respectively. For scRNAseq data, principal
component analysis (PCA) against all genes in the target panel was performed using RunPCA
(number of calculated principal components (PCs) was 100). Then, the enrichment of each PC was
calculated using the JackStraw and ScoreJackStraw function (num.replicate = 100), and PCs that
were significantly enriched (p ≤ 0.05) were selected for further analysis. For AbSeq data, PCA
against all genes included in Immune Discovery Panel was performed using RunPCA, and the
same number of PCs as scRNAseq was selected for further analysis. Cell clustering was performed
based on the weighted-nearest neighbor graph using FindMultiModalNeighbors and FindClusters
against the significant PCs of scRNAseq and AbSeq (*24*). The dimensional reduction was
performed using RunUMAP based on the nearest neighbor graph determined by
FindMultiModalNeighbors. Marker genes of each identified cluster were identified using
FindAllMarkers function (test.use="Wilcox", min.pct=0.1, return.thresh=0.05). Clusters
constituted with contaminant cells determined by marker genes were excluded, and the above
analysis procedure was repeated.

Data processing and analysis of scTCRseq

Raw data of single-cell TCR sequencing were processed by pipeline produced by
ImmunoGeneTeqs, Inc., as described previously (*52*). Briefly, sequencing reads were separated by
cell barcode, then adapter trimming and quality filtering of sequencing data were performed using
cutadapt 3.4 (*53*). PCR and sequencing errors were corrected by lighter v1.1.2 (*63*) with the
following parameters: -newQual 25 -maxcor 4 -K 20. Error-corrected sequence reads were
processed by MiXCR (*55*) to generate the list of TCRα or TCRβ sequences for each cell barcode,
using reference mouse TCR V/D/J sequences registered in the international IMGT information
system. Quality filtering of cell barcodes was performed with the following criteria: (1) more than
32 TCR reads were detected. (2) the proportion of the most frequent TCR sequence in cell barcodes
was over 0.6. (3) the cell barcode was annotated to the T cell cluster in scRNAseq/AbSeq analysis.
Finally, the most frequent sequence of TCRα and TCRβ was adopted and paired in each cell
barcode. Identified TCRα or TCRβ sequence was imported into the meta.data matrix in Seurat
object of scRNAseq/AbSeq data based on the cellular barcode identities. T-cell clones were
determined as cells with the same TCRβ sequence defined by V segment, J segment, and CDR3
nucleotide sequence. The frequency of each clone was calculated as the proportion of cells
belonging to the clone in all T cells whose TCRβ sequence was assigned. The 1 - Pielou index was
used to evaluate the clonality of clusters, which was calculated as the following formula: $1 -$
$\sum_{i=1}^{n} p_i \log_e(p_i) / \log_e(n)$.

Quantification and statistical analysis

Statistical analyses were performed using GraphPad Prism (ver9.4.0) software (GraphPad
Software). Friedman test with Dunn's multiple comparisons test was run on the comparison
between time points or kinetic profile patterns. Friedman test with Dunn's multiple comparisons
test was run on the comparison between time points or kinetic profile patterns. Wilcoxon matched-
pairs signed rank test was run on the comparison between AIM$^+$ and AIM$^-$ clones. Mann-Whitney
test was run on the comparison between AIM-high and AIM-low participants. Spearman
correlation test was run on the correlation analysis. Asterisks to indicate significance
corresponding to the following: ***P<0.001; **P<0.005; *P<0.05; n.s. non-significant (P≥0.05).

Supplementary Materials

References and Notes

1. X. Han, P. Xu, Q. Ye, Analysis of COVID-19 vaccines: Types, thoughts, and application. *J. Clin. Lab. Anal.* **35**, 1–7 (2021).

2. U. Sahin, A. Muik, I. Vogler, E. Derhovanessian, L. M. Kranz, M. Vormehr, J. Quandt, N. Bidmon, A. Ulges, A. Baum, K. E. Pascal, D. Maurus, S. Brachtendorf, V. Lörks, J. Sikorski, P. Koch, R. Hilker, D. Becker, A. K. Eller, J. Grützner, M. Tonigold, C. Boesler, C. Rosenbaum, L. Heesen, M. C. Kühnle, A. Poran, J. Z. Dong, U. Luxemburger, A. Kemmer-Brück, D. Langer, M. Bexon, S. Bolte, T. Palanche, A. Schultz, S. Baumann, A. J. Mahiny, G. Boros, J. Reinholz, G. T. Szabó, K. Karikó, P. Y. Shi, C. Fontes-Garfias, J. L. Perez, M. Cutler, D. Cooper, C. A. Kyratsous, P. R. Dormitzer, K. U. Jansen, Ö. Türeci, BNT162b2 vaccine induces neutralizing antibodies and poly-specific T cells in humans. *Nature*. **595**, 572–577 (2021).

3. A. Pegu, S. E. O'Connell, S. D. Schmidt, S. O'Dell, C. A. Talana, L. Lai, J. Albert, E. Anderson, H. Bennett, K. S. Corbett, B. Flach, L. Jackson, B. Leav, J. E. Ledgerwood, C. J. Luke, M. Makowski, M. C. Nason, P. C. Roberts, M. Roederer, P. A. Rebolledo, C. A. Rostad, N. G. Rouphael, W. Shi, L. Wang, A. T. Widge, E. S. Yang, J. H. Beigel, B. S. Graham, J. R. Mascola, M. S. Suthar, A. B. McDermott, N. A. Doria-Rose, Durability of mRNA-1273 vaccine-induced antibodies against SARS-CoV-2 variants. *Science (80-.).* **373**, 1372–1377 (2021).

4. J. Mateus, J. M. Dan, Z. Zhang, C. R. Moderbacher, M. Lammers, B. Goodwin, A. Sette, S. Crotty, D. Weiskopf, Low-dose mRNA-1273 COVID-19 vaccine generates durable memory enhanced by cross-reactive T cells. *Science (80-.).* **374**, 1–18 (2021).

5. M. M. Painter, D. Mathew, R. R. Goel, S. A. Apostolidis, A. Pattekar, O. Kuthuru, A. E. Baxter, R. S. Herati, D. A. Oldridge, S. Gouma, P. Hicks, S. Dysinger, K. A. Lundgreen, L. Kuri-Cervantes, S. Adamski, A. Hicks, S. Korte, J. R. Giles, M. E. Weirick, C. M. McAllister, J. Dougherty, S. Long, K. D'Andrea, J. T. Hamilton, M. R. Betts, P. Bates, S. E. Hensley, A. Grifoni, D. Weiskopf, A. Sette, A. R. Greenplate, E. J. Wherry, Rapid induction of antigen-specific CD4+ T cells is associated with coordinated humoral and cellular immunity to SARS-CoV-2 mRNA vaccination. *Immunity*. **54**, 2133-2142.e3 (2021).

6. G. Guerrera, M. Picozza, S. D'Orso, R. Placido, M. Pirronello, A. Verdiani, A. Termine, C. Fabrizio, F. Giannessi, M. Sambucci, M. P. Balice, C. Caltagirone, A. Salvia, A. Rossini, L. Battistini, G. Borsellino, BNT162b2 vaccination induces durable SARS-CoV-2–specific T cells with a stem cell memory phenotype. *Sci. Immunol.* **6**, 1–18 (2021).

7. P. Moss, The T cell immune response against SARS-CoV-2. *Nat. Immunol.* **23**, 186–193 (2022).

8. D. T. Skelly, A. C. Harding, J. Gilbert-Jaramillo, M. L. Knight, S. Longet, A. Brown, S. Adele, E. Adland, H. Brown, S. Chinnakannan, T. Donnison, M. Ali, P. Rongkard, M. Pace, P. Zacharopoulou, N. Robinson, A. Csala, C. De Lara, C. L. Hutchings, H. Mehta, L. N. Lee, M. Edmans, C. P. Hackstein, P. Phalora, W. Li, E. Phillips, T. Malone, A. Ogbe, C. Jay, T. Tipoe, T. Tipton, L. Stafford, A. J. Mentzer, S. A. Johnson, A. Amini, T. Marjot, S. Dimitriadis, B. Simmons, A. Deeks, S. Kerneis, H. Abuelgasim, R. Wilson, S. R. Thomas, A. Watson, A. Alhussni, J. Cutteridge, E. Weeks, L. Denly, K. Lillie, J. Holmes, P. C. Matthews, D. O'Donnell, T. K. Tan, L. Schimanski, K. Y. A. Huang, P. Rijal, L. Turtle, T. de Silva, A. Richter, C. J. A. Duncan, R. P. Payne, S. C. Moore, J. C. Knight, M. P. Cassar, B. Raman, S. Neubauer, A. Fries, N. P. Talbot, N. Petousi, L. P. Ho, Y. Peng, T. Dong, S. Camara, S. Marinou, A. Linder, S. Adlou, M. Kasanyinga, A. Bridges-Webb, J. Hill, L. Silva-Reyes, L. Blackwell, J. Frater, P. Goulder, C. P. Conlon, K. Jeffery, C. Dold, A. J. Pollard, A. Sigal, T. de Oliveira, A. R. Townsend, P. Klenerman, S. J. Dunachie, E. Barnes, M. W. Carroll, W. S. James, Two doses of SARS-CoV-2 vaccination induce robust immune responses to emerging SARS-CoV-2 variants of concern. *Nat. Commun.* **12**, 1–12 (2021).

9. D. L. Woodland, Jump-starting the immune system: Prime-boosting comes of age. *Trends Immunol.* **25**, 98–104 (2004).

10. D. Masopust, S.-J. Ha, V. Vezys, R. Ahmed, Stimulation History Dictates Memory CD8 T Cell Phenotype: Implications for Prime-Boost Vaccination. *J. Immunol.* **177**, 831–839 (2006).

11. V. Vezys, A. Yates, K. A. Casey, G. Lanier, R. Ahmed, R. Antia, D. Masopust, Memory CD8 T-cell compartment grows in size with immunological experience. *Nature*. **457**, 196–199 (2009).

12. K. A. Fraser, J. M. Schenkel, S. C. Jameson, V. Vezys, D. Masopust, Preexisting High Frequencies of Memory CD8+ T Cells Favor Rapid Memory Differentiation and Preservation of Proliferative Potential upon Boosting. *Immunity*. **39**, 171–183 (2013).

13. T. C. Wirth, H. H. Xue, D. Rai, J. T. Sabel, T. Bair, J. T. Harty, V. P. Badovinac, Repetitive antigen stimulation induces stepwise transcriptome diversification but

preserves a core signature of memory CD8+ T cell differentiation. *Immunity*. **33**, 128–140
(2010).

14.   M. Kurachi, K. Kakimi, S. Ueha, K. Matsushima, Maintenance of memory CD8+ T cell
diversity and proliferative potential by a primary response upon re-challenge. *Int.*
Immunol. **19**, 105–115 (2007).

15.   M. Kitabatake, N. Ouji-Sageshima, S. Sonobe, R. Furukawa, M. Konda, A. Hara, H. Aoki,
Y. Suzuki, N. Imakita, A. Nakano, Y. Fujita, S. Shichino, R. Nakano, S. Ueha, K.
Kasahara, S. Muro, H. Yano, K. Matsushima, T. Ito, Transition of antibody titers after the
SARS-CoV-2 mRNA vaccine in Japanese healthcare workers. *medRxiv*. 1-9, (2021). doi:
https://doi.org/10.1101/2021.12.28.21268435.

16.   P. Langfelder, S. Horvath, WGCNA: An R package for weighted correlation network
analysis. *BMC Bioinformatics*. **9** (2008), doi:10.1186/1471-2105-9-559.

17.   T. M. Snyder, R. M. Gittelman, M. Klinger, D. H. May, E. J. Osborne, H. J. Zahid, I. M.
Kaplan, J. N. Dines, M. T. Noakes, R. Pandya, X. Chen, S. Elasady, E. Svejnoha, P. Ebert,
M. W. Pesesky, P. De, H. O. Donnell, Q. Degottardi, G. Keitany, J. Lu, A. Vong, P.
Fields, J. Greissl, L. Baldo, S. Semprini, C. Cerchione, O. M. Delmonte, K. Dobbs, G.
Carreño-tarragona, S. Barrio, M. D. Angio, P. Bonfanti, M. F. Tompkins, C. Alba, C.
Dalgard, J. Martinez-lopez, J. M. Carlson, H. S. Robins, Magnitude and Dynamics of the
T-Cell Response to SARS-CoV-2 Infection at Both Individual and Population Levels.
medRxiv. 1-33, (2020). doi: 10.1101/2020.07.31.20165647.

18.   J. M. Francis, D. Leistritz-Edwards, A. Dunn, C. Tarr, J. Lehman, C. Dempsey, A. Hamel,
V. Rayon, G. Liu, Y. Wang, M. Wille, M. Durkin, K. Hadley, A. Sheena, B. Roscoe, M.
Ng, G. Rockwell, M. Manto, E. Gienger, J. Nickerson, MGH COVID-19 Collection and
Processing Team2‡, A. Moarefi, M. Noble, T. Malia, P. D. Bardwell, W. Gordon, J.
Swain, M. Skoberne, K. Sauer, T. Harris, A. W. Goldrath, A. K. Shalek, A. J. Coyle, C.
Benoist, D. C. Pregibon, N. Jilg, J. Li, A. Rosenthal, C. Wong, G. Daley, D. Golan, H.
Heller, A. Sharpe, B. A. Abayneh, P. Allen, D. Antille, K. Armstrong, S. Boyce, J. Braley,
K. Branch, K. Broderick, J. Carney, A. Chan, S. Davidson, M. Dougan, D. Drew, A.
Elliman, K. Flaherty, J. Flannery, P. Forde, E. Gettings, A. Griffin, S. Grimmel, K.
Grinke, K. Hall, M. Healy, D. Henault, G. Holland, C. Kayitesi, V. LaValle, Y. Lu, S.
Luthern, J. M. Schneider, B. Martino, R. McNamara, C. Nambu, S. Nelson, M. Noone, C.
Ommerborn, L. C. Pacheco, N. Phan, F. A. Porto, E. Ryan, K. Selleck, S. Slaughenhaupt,
K. S. Sheppard, E. Suschana, V. Wilson, M. Carrington, M. Martin, Y. Yuki, G. Alter, A.
Balazs, J. Bals, M. Barbash, Y. Bartsch, J. Boucau, M. Carrington, J. Chevalier, F.
Chowdhury, E. DeMers, K. Einkauf, J. Fallon, L. Fedirko, K. Finn, P. Garcia-Broncano,
M. S. Ghebremichael, C. Hartana, C. Jiang, K. Judge, P. Kaplonek, M. Karpell, P. Lai, E.
C. Lam, K. Lefteri, X. Lian, M. Lichterfeld, D. Lingwood, H. Liu, J. Liu, N. Ly, Z. M.
Hill, A. Michell, I. Millstrom, N. Miranda, C. O'Callaghan, M. Osborn, S. Pillai, Y.
Rassadkina, A. Reissis, F. Ruzicka, K. Seiger, L. Sessa, C. Sharr, S. Shin, N. Singh, W.
Sun, X. Sun, H. Ticheli, A. Trocha-Piechocka, B. Walker, D. Worrall, X. G. Yu, A. Zhu,
Allelic variation in class I HLA determines CD8+ T cell repertoire shape and cross-
reactive memory responses to SARS-CoV-2. *Sci. Immunol.* **3070**, eabk3070 (2021).

19.   A. A. Minervina, M. V. Pogorelyy, A. M. Kirk, J. C. Crawford, E. K. Allen, C. H. Chou,
R. C. Mettelman, K. J. Allison, C. Y. Lin, D. C. Brice, X. Zhu, K. Vegesana, G. Wu, S.

Trivedi, P. Kottapalli, D. Darnell, S. McNeely, S. R. Olsen, S. Schultz-Cherry, J. H. Estepp, A. Gaur, J. Hoffman, M. Mori, L. Tang, E. Tuomanen, R. Webby, H. Hakim, R. T. Hayden, D. R. Hijano, R. Bajracharya, W. Awad, L. A. Van de Velde, B. L. Clark, T. L. Wilson, A. Souquette, A. Castellaw, R. H. Dallas, J. Hodges, A. Gowen, J. Russell-Bell, J. Sparks, D. E. Wittman, T. P. Fabrizio, S. Cherry, E. K. Roubidoux, V. Cortez, P. Freiden, N. Wohlgemuth, K. Whitt, M. A. McGargill, J. Wolf, P. G. Thomas, SARS-CoV-2 antigen exposure history shapes phenotypes and specificity of memory CD8+ T cells. *Nat. Immunol.* **23**, 781–790 (2022).

20. K. Mayer-Blackwell, S. Schattgen, L. Cohen-Lavi, J. C. Crawford, A. Souquette, J. A. Gaevert, T. Hertz, P. G. Thomas, P. G. Bradley, A. Fiore-Gartland, Tcr meta-clonotypes for biomarker discovery with tcrdist3 enabled identification of public, hla-restricted clusters of sars-cov-2 tcrs. *Elife.* **10**, 1–32 (2021).

21. M. R. Betts, J. M. Brenchley, D. A. Price, S. C. De Rosa, D. C. Douek, M. Roederer, R. A. Koup, Sensitive and viable identification of antigen-specific CD8+ T cells by a flow cytometric assay for degranulation. *J. Immunol. Methods.* **281**, 65–78 (2003).

22. S. Reiss, A. E. Baxter, K. M. Cirelli, J. M. Dan, A. Morou, A. Daigneault, N. Brassard, G. Silvestri, J. P. Routy, C. Havenar-Daughton, S. Crotty, D. E. Kaufmann, Comparative analysis of activation induced marker (AIM) assays for sensitive identification of antigen-specific CD4 T cells. *PLoS One.* **12**, 1–22 (2017).

23. C. J. Thieme, M. Anft, K. Paniskaki, A. Blazquez-Navarro, A. Doevelaar, F. S. Seibert, B. Hoelzer, M. J. Konik, M. M. Berger, T. Brenner, C. Tempfer, C. Watzl, T. L. Meister, S. Pfaender, E. Steinmann, S. Dolff, U. Dittmer, T. H. Westhoff, O. Witzke, U. Stervbo, T. Roch, N. Babel, Robust T Cell Response Toward Spike, Membrane, and Nucleocapsid SARS-CoV-2 Proteins Is Not Associated with Recovery in Critical COVID-19 Patients. *Cell Reports Med.* **1**, 100092 (2020).

24. Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler, M. J. Lee, A. J. Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E. P. Mimitou, J. Jain, A. Srivastava, T. Stuart, L. M. Fleming, B. Yeung, A. J. Rogers, J. M. McElrath, C. A. Blish, R. Gottardo, P. Smibert, R. Satija, Integrated analysis of multimodal single-cell data. *Cell.* **184**, 3573-3587.e29 (2021).

25. F. Sallusto, D. Lenig, R. Förster, M. Lipp, A. Lanzavecchia, Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature.* **401**, 708–712 (1999).

26. D. Hamann, P. A. Baars, M. H. G. Rep, B. Hooibrink, S. R. Kerkhof-Garde, M. R. Klein, R. A. W. Van Lier, Phenotypic and functional separation of memory and effector human CD8+ T cells. *J. Exp. Med.* **186**, 1407–1418 (1997).

27. M. Dusseaux, E. Martin, N. Serriari, I. Péguillet, V. Premel, D. Louis, M. Milder, L. L. Bourhis, C. Soudais, E. Treiner, O. Lantz, Human MAIT cells are xenobiotic-resistant, tissue-targeted, CD161hi IL-17–secreting T cells. Blood. 117(4), 1250-1259 (2011). doi: 10.1182/blood-2010-08-303339.

28. C. Zheng, L. Zheng, J. K. Yoo, H. Guo, Y. Zhang, X. Guo, B. Kang, R. Hu, J. Y. Huang, Q. Zhang, Z. Liu, M. Dong, X. Hu, W. Ouyang, J. Peng, Z. Zhang, Landscape of

Infiltrating T Cells in Liver Cancer Revealed by Single-Cell Sequencing. *Cell*. **169**, 1342-
1356.e16 (2017).

29.   A. Kurioka, L. J. Walker, P. Klenerman, C. B. Willberg, MAIT cells: new guardians of
the liver. *Clin. Transl. Immunol.* **5**, e98 (2016).

30.   M. Lepore, A. Kalinicenko, A. Colone, B. Paleja, A. Singhal, A. Tschumi, B. Lee, M.
Poidinger, F. Zolezzi, L. Quagliata, P. Sander, E. Newell, A. Bertoletti, L. Terracciano, G.
De Libero, L. Mori, Parallel T-cell cloning and deep sequencing of human MAIT cells
reveal stable oligoclonal TCRβ 2 repertoire. *Nat. Commun.* **5** (2014),
doi:10.1038/ncomms4866.

31.   G. Galletti, G. De Simone, E. M. C. Mazza, S. Puccio, C. Mezzanotte, T. M. Bi, A. N.
Davydov, M. Metsger, E. Scamardella, G. Alvisi, F. De Paoli, V. Zanon, A. Scarpa, B.
Camisa, F. S. Colombo, A. Anselmo, C. Peano, S. Polletti, D. Mavilio, L. Gattinoni, S. K.
Boi, B. A. Youngblood, R. E. Jones, D. M. Baird, E. Gostick, S. Llewellyn-Lacey, K.
Ladell, D. A. Price, D. M. Chudakov, E. W. Newell, M. Casucci, E. Lugli, Two subsets of
stem-like CD8+ memory T cell progenitors with distinct fate commitments in humans.
Nat. Immunol. **21**, 1552–1562 (2020).

32.   A. Sette, S. Crotty, Pre-existing immunity to SARS-CoV-2: the knowns and unknowns.
Nat. Rev. Immunol. **20**, 457–458 (2020).

33.   A. Grifoni, D. Weiskopf, S. I. Ramirez, J. Mateus, J. M. Dan, C. R. Moderbacher, S. A.
Rawlings, A. Sutherland, L. Premkumar, R. S. Jadi, D. Marrama, A. M. de Silva, A.
Frazier, A. F. Carlin, J. A. Greenbaum, B. Peters, F. Krammer, D. M. Smith, S. Crotty, A.
Sette, Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-
19 Disease and Unexposed Individuals. *Cell*. **181**, 1489-1501.e15 (2020).

34.   D. Weiskopf, K. S. Schmitz, M. P. Raadsen, A. Grifoni, N. M. A. Okba, H. Endeman, J.
P. C. van den Akker, R. Molenkamp, M. P. G. Koopmans, E. C. M. van Gorp, B. L.
Haagmans, R. L. de Swart, A. Sette, R. D. de Vries, Phenotype and kinetics of SARS-
CoV-2-specific T cells in COVID-19 patients with acute respiratory distress syndrome.
Sci. Immunol. **5**, 1–11 (2020).

35.   P. Bacher, E. Rosati, D. Esser, G. R. Martini, C. Saggau, E. Schiminsky, J. Dargvainiene,
I. Schröder, I. Wieters, Y. Khodamoradi, F. Eberhardt, M. J. G. T. Vehreschild, H. Neb,
M. Sonntagbauer, C. Conrad, F. Tran, P. Rosenstiel, R. Markewitz, K. P. Wandinger, M.
Augustin, J. Rybniker, M. Kochanek, F. Leypoldt, O. A. Cornely, P. Koehler, A. Franke,
A. Scheffold, Low-Avidity CD4+ T Cell Responses to SARS-CoV-2 in Unexposed
Individuals and Humans with Severe COVID-19. *Immunity*. **53**, 1258-1271.e5 (2020).

36.   H. Kay Chung, B. McDonald, S. M. Kaech, The architectural design of CD8+ T cell
responses in acute and chronic infection: Parallel structures with divergent fates. *J. Exp.*
Med. **218**, 1–15 (2021).

37.   D. M. Catron, L. K. Rusch, J. Hataye, A. A. Itano, M. K. Jenkins, CD4+ T cells that enter
the draining lymph nodes after antigen injection participate in the primary response and
become central-memory cells. *J. Exp. Med.* **203**, 1045–1054 (2006).

38.   S. Shichino, S. Ueha, S. Hashimoto, M. Otsuji, J. Abe, T. Tsukui, S. Deshimaru, T.
Nakajima, M. Kosugi-Kanaya, F. H. W. Shand, Y. Inagaki, H. Shimano, K. Matsushima,

Transcriptome network analysis identifies protective role of the LXR/SREBP-1c axis in murine pulmonary fibrosis. *JCI Insight*. **4** (2019), doi:10.1172/jci.insight.122163.

39. T. A. Doering, A. Crawford, J. M. Angelosanto, M. A. Paley, C. G. Ziegler, E. J. Wherry, Network Analysis Reveals Centrally Connected Genes and Pathways Involved in CD8+ T Cell Exhaustion versus Memory. *Immunity*. **37**, 1130–1144 (2012).

40. J. Rytlewski, S. Deng, T. Xie, C. Davis, H. Robins, E. Yusko, J. Bienkowska, Model to improve specificity for identification of clinically-relevant expanded T cells in peripheral blood. *PLoS One*. **14**, 1–10 (2019).

41. W. S. DeWitt, R. O. Emerson, P. Lindau, M. Vignali, T. M. Snyder, C. Desmarais, C. Sanders, H. Utsugi, E. H. Warren, J. McElrath, K. W. Makar, A. Wald, H. S. Robins, Dynamics of the Cytotoxic T Cell Response to a Model of Acute Viral Infection. *J. Virol.* **89**, 4517–4526 (2015).

42. A. A. Minervina, M. V. Pogorelyy, E. A. Komech, V. K. Karnaukhov, P. Bacher, E. Rosati, A. Franke, D. M. Chudakov, I. Z. Mamedov, Y. B. Lebedev, T. Mora, A. M. Walczak, Primary and secondary anti-viral response captured by the dynamics and phenotype of individual T cell clones. *Elife*. **9**, 1–21 (2020).

43. C. Li, A. Lee, L. Grigoryan, P. S. Arunachalam, M. K. D. Scott, M. Trisal, F. Wimmers, M. Sanyal, P. A. Weidenbacher, Y. Feng, J. Z. Adamska, E. Valore, Y. Wang, R. Verma, N. Reis, D. Dunham, R. O'Hara, H. Park, W. Luo, A. D. Gitlin, P. Kim, P. Khatri, K. C. Nadeau, B. Pulendran, Mechanisms of innate and adaptive immunity to the Pfizer-BioNTech BNT162b2 vaccine. *Nat. Immunol.* **23**, 543–555 (2022).

44. M. M. L. Poon, K. Rybkina, Y. Kato, M. Kubota, R. Matsumoto, N. I. Bloom, Z. Zhang, K. M. Hastie, A. Grifoni, D. Weiskopf, S. B. Wells, B. B. Ural, N. Lam, P. A. Szabo, P. Dogra, Y. S. Lee, J. I. Gray, M. C. Bradley, M. A. Brusko, T. M. Brusko, E. O. Saphire, T. J. Connors, A. Sette, S. Crotty, D. L. Farber, SARS-CoV-2 infection generates tissue-localized immunological memory in humans. *Sci. Immunol.* **6** (2021), doi:10.1126/sciimmunol.abl9105.

45. L. Kok, F. E. Dijkgraaf, J. Urbanus, K. Bresser, D. W. Vredevoogd, R. F. Cardoso, L. Perié, J. B. Beltman, T. N. Schumacher, A committed tissue-resident memory T cell precursor within the circulating CD8+ effector T cell pool. *J. Exp. Med.* **217** (2020), doi:10.1084/JEM.20191711.

46. Y. Lustig, T. Gonen, L. Meltzer, M. Gilboa, V. Indenbaum, C. Cohen, S. Amit, H. Jaber, R. Doolman, K. Asraf, C. Rubin, R. Fluss, E. Mendelson, L. Freedman, G. Regev-Yochay, Y. Kreiss, Superior immunogenicity and effectiveness of the third compared to the second BNT162b2 vaccine dose. *Nat. Immunol.* **23**, 940–946 (2022).

47. A. P. S. Munro, S. Feng, L. Janani, V. Cornelius, P. K. Aley, G. Babbage, D. Baxter, M. Bula, K. Cathie, K. Chatterjee, K. Dodd, Y. Enever, E. Qureshi, A. L. Goodman, C. A. Green, L. Harndahl, J. Haughney, A. Hicks, A. A. van der Klaauw, N. Kanji, V. Libri, M. J. Llewelyn, A. C. McGregor, M. Maallah, A. M. Minassian, P. Moore, M. Mughal, Y. F. Mujadidi, K. Holliday, O. Osanlou, R. Osanlou, D. R. Owens, M. Pacurar, A. Palfreeman, D. Pan, T. Rampling, K. Regan, S. Saich, T. Bawa, D. Saralaya, S. Sharma, R. Sheridan, E. C. Thomson, S. Todd, C. Twelves, R. C. Read, S. Charlton, B. Hallis, M. Ramsay, N. Andrews, T. Lambe, J. S. Nguyen-Van-Tam, M. D. Snape, X. Liu, S. N. Faust, A. P. S.

Munro, S. Feng, L. Janani, V. Cornelius, P. K. Aley, G. Babbage, D. Baxter, M. Bula, K. Cathie, K. Chatterjee, K. Dodd, Y. Enever, E. Qureshi, A. L. Goodman, C. A. Green, L. Harndahl, J. Haughney, A. Hicks, A. A. van der Klaauw, N. Kanji, V. Libri, M. J. Llewelyn, A. C. McGregor, A. M. Minassian, P. Moore, M. Mughal, Y. F. Mujadidi, K. Holliday, O. Osanlou, R. Osanlou, D. R. Owens, M. Pacurar, A. Palfreeman, D. Pan, T. Rampling, K. Regan, S. Saich, T. Bawa, D. Saralaya, S. Sharma, R. Sheridan, M. Maallah, E. C. Thomson, S. Todd, C. Twelves, R. C. Read, S. Charlton, B. Hallis, M. Ramsay, N. Andrews, T. Lambe, J. S. Nguyen-Van-Tam, M. D. Snape, X. Liu, S. N. Faust, A. Riordan, A. Ustianowski, C. Rogers, K. Katechia, A. Cooper, A. Freedman, R. Hughes, L. Grundy, L. Tudor Jones, E. Harrison, E. Snashall, L. Mallon, K. Burton, K. Storton, M. Munusamy, B. Tandy, A. Egbo, S. Cox, N. N. Ahmed, A. Shenoy, R. Bousfield, D. Wixted, H. Gutteridge, B. Mansfield, C. Herbert, J. Murira, J. Calderwood, D. Barker, J. Brandon, H. Tulloch, S. Colquhoun, H. Thorp, H. Radford, J. Evans, H. Baker, J. Thorpe, S. Batham, J. Hailstone, R. Phillips, D. Kumar, F. Westwell, A. Sturdy, L. Barcella, N. Soussi, M. Mpelembue, S. Raj, R. Sharma, T. Corrah, L. John, A. Whittington, S. Roche, L. Wagstaff, A. Farrier, K. Bisnauthsing, M. Abeywickrama, N. Spence, A. Packham, T. Serafimova, S. Aslam, C. McGreevy, A. Borca, P. DeLosSantosDominguez, E. Palmer, S. Broadhead, S. Farooqi, J. Piper, R. Weighell, L. Pickup, D. Shamtally, J. Domingo, E. Kourampa, C. Hale, J. Gibney, M. Stackpoole, Z. Rashid-Gardner, R. Lyon, C. McDonnell, C. Cole, A. Stewart, G. McMillan, M. Savage, H. Beckett, C. Moorbey, A. Desai, C. Brown, K. Naker, K. Gokani, C. Trinham, C. Sabine, S. Moore, S. Hurdover, E. Justice, M. Stone, E. Plested, C. Ferreira Da Silva, R. White, H. Robinson, I. Turnbull, G. Morshead, R. Drake-Brockman, C. Smith, G. Li, M. Kasanyinga, E. A. Clutterbuck, S. Bibi, M. Singh, T. Champaneri, M. Irwin, M. Khan, A. Kownacka, M. Nabunjo, C. Osuji, J. Hladkiwskyj, D. Galvin, G. Patel, J. Grierson, S. Males, K. Askoolam, J. Barry, J. Mouland, B. Longhurst, M. Moon, B. Giddins, C. Pereira Dias Alves, L. Richmond, C. Minnis, S. Baryschpolec, S. Elliott, L. Fox, V. Graham, N. Baker, K. Godwin, K. Buttigieg, C. Knight, P. Brown, P. Lall, I. Shaik, E. Chiplin, E. Brunt, S. Leung, L. Allen, S. Thomas, S. Fraser, B. Choi, J. Gouriet, J. Perkins, A. Gowland, J. Macdonald, J. P. Seenan, I. Starinskij, A. Seaton, E. Peters, S. Singh, B. Gardside, A. Bonnaud, C. Davies, E. Gordon, S. Keenan, J. Hall, S. Wilkins, S. Tasker, R. James, I. Seath, K. Littlewood, J. Newman, I. Boubriak, D. Suggitt, H. Haydock, S. Bennett, W. Woodyatt, K. Hughes, J. Bell, T. Coughlan, D. van Welsenes, M. Kamal, C. Cooper, S. Tunstall, N. Ronan, R. Cutts, T. Dare, Y. T. N. Yim, S. Whittley, S. Hamal, M. Ricamara, K. Adams, H. Baker, K. Driver, N. Turner, T. Rawlins, S. Roy, M. Merida-Morillas, Y. Sakagami, A. Andrews, L. Goncalvescordeiro, M. Stokes, W. Ambihapathy, J. Spencer, N. Parungao, L. Berry, J. Cullinane, L. Presland, A. Ross Russell, S. Warren, J. Baker, A. Oliver, A. Buadi, K. Lee, L. Haskell, R. Romani, I. Bentley, T. Whitbred, S. Fowler, J. Gavin, A. Magee, T. Watson, K. Nightingale, P. Marius, E. Summerton, E. Locke, T. Honey, A. Lingwood, A. de la Haye, R. S. Elliott, K. Underwood, M. King, S. Davies-Dear, E. Horsfall, O. Chalwin, H. Burton, C. J. Edwards, B. Welham, K. Appleby, E. Dineen, S. Garrahy, F. Hall, E. Ladikou, D. Mullan, D. Hansen, M. Campbell, F. Dos Santos, N. Lakeman, D. Branney, L. Vamplew, A. Hogan, J. Frankham, M. Wiselka, D. Vail, V. Wenn, V. Renals, K. Ellis, J. Lewis-Taylor, H. Habash-Bailey, J. Magan, A. Hardy, Safety, immunogenicity, and reactogenicity of BNT162b2 and mRNA-1273 COVID-19 vaccines given as fourth-dose boosters following two doses of ChAdOx1 nCoV-19 or BNT162b2 and a third dose

of BNT162b2 (COV-BOOST): a multicentre, blinded, phase 2, randomised t. *Lancet Infect. Dis.* **2**, 1131–1141 (2022).

48. N. Andrews, J. Stowe, F. Kirsebom, S. Toffa, T. Rickeard, E. Gallagher, C. Gower, M. Kall, N. Groves, A.-M. O'Connell, D. Simons, P. B. Blomquist, A. Zaidi, S. Nash, N. Iwani Binti Abdul Aziz, S. Thelwall, G. Dabrera, R. Myers, G. Amirthalingam, S. Gharbia, J. C. Barrett, R. Elson, S. N. Ladhani, N. Ferguson, M. Zambon, C. N. J. Campbell, K. Brown, S. Hopkins, M. Chand, M. Ramsay, J. Lopez Bernal, Covid-19 Vaccine Effectiveness against the Omicron (B.1.1.529) Variant. *N. Engl. J. Med.* **386**, 1532–1546 (2022).

49. Y. M. Bar-On, Y. Goldberg, M. Mandel, O. Bodenheimer, O. Amir, L. Freedman, S. Alroy-Preis, N. Ash, A. Huppert, R. Milo, Protection by a Fourth Dose of BNT162b2 against Omicron in Israel. *N. Engl. J. Med.* **386**, 1712–1720 (2022).

50. C. Henry, A. K. E. Palm, F. Krammer, P. C. Wilson, From Original Antigenic Sin to the Universal Influenza Virus Vaccine. *Trends Immunol.* **39**, 70–79 (2018).

51. A. Biswas, A. K. Chakrabarti, S. Dutta, Current challenges: from the path of "original antigenic sin" towards the development of universal flu vaccines: Flu vaccine efficacy encounters significant hurdles from pre-existing immunity of the host suggesting assessment of host immunity before vacci. *Int. Rev. Immunol.* **39**, 21–36 (2020).

52. H. Aoki, M. Tsunoda, H. Ogiwara, H. Shimizu, H. Abe, T. Abe, S. Shichino, K. Matsushima, S. Ueha, Clonal spreading of tumor-infiltrating T cells underlies the robust antitumor immune responses. bioRxiv, 1-59, (2022). doi: https://doi.org/10.1101/2022.04.19.488731.

53. M. Martin, Cutadapt Removes Adapter Sequences From High-Throughput Sequencing Reads. EMBnet.journal. 17, 10–12 (2013). DOI: https://doi.org/10.14806/ej.17.1.200

54. R. Schmieder, R. Edwards, Quality control and preprocessing of metagenomic datasets. *Bioinformatics*. **27**, 863–864 (2011).

55. D. A. Bolotin, S. Poslavsky, I. Mitrophanov, M. Shugay, I. Z. Mamedov, E. V. Putintseva, D. M. Chudakov, MiXCR: Software for comprehensive adaptive immunity profiling. *Nat. Methods*. **12**, 380–381 (2015).

56. H. Aoki, S. Ueha, S. Shichino, H. Ogiwara, K. Shitara, M. Shimomura, T. Suzuki, T. Nakatsura, M. Yamashita, S. Kitano, S. Kuroda, M. Wakabayashi, M. Kurachi, S. Ito, T. Doi, K. Matsushima, Transient depletion of CD4þ cells induces remodeling of the TCR repertoire in gastrointestinal cancer. *Cancer Immunol. Res.* **9**, 624–636 (2021).

57. M. Shugay, D. V. Bagaev, M. A. Turchaninova, D. A. Bolotin, O. V. Britanova, E. V. Putintseva, M. V. Pogorelyy, V. I. Nazarov, I. V. Zvyagin, V. I. Kirgizova, K. I. Kirgizov, E. V. Skorobogatova, D. M. Chudakov, VDJtools: Unifying Post-analysis of T Cell Receptor Repertoires. *PLoS Comput. Biol.* **11**, 1–16 (2015).

58. M. Klinger, F. Pepin, J. Wilkins, T. Asbury, T. Wittkop, J. Zheng, M. Moorhead, M. Faham, Multiplex identification of antigen-specific T cell receptors using a combination of immune assays and immune receptor sequencing. *PLoS One*. **10**, 1–21 (2015).

59. S. Shichino, S. Ueha, S. Hashimoto, T. Ogawa, H. Aoki, B. Wu, C.-Y. Chen, M.
Kitabatake, N. Ouji-Sageshima, S. Hontsu, N. Sawabata, T. Kawaguchi, T. Okayama, E.
Sugihara, T. Ito, Y. Iwata, T. Wada, K. Ikeo, T. Sato, K. Matsushima, TAS-Seq is a robust
and sensitive amplification method for bead-based scRNA-seq. *Commun Biol*, **5(1)**, 602
(2022). doi: 10.1038/s42003-022-03536-0.

60. W. Shen, S. Le, Y. Li, F. Hu, SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q
file manipulation. *PLoS One*. **11**, 1–10 (2016).

61. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods*. **9**,
357–359 (2012).

62. A. T. L. Lun, S. Riesenfeld, T. Andrews, T. P. Dao, T. Gomes, J. C. Marioni,
Distinguishing cells from empty droplets in droplet-based single-cell RNA sequencing
data. *bioRxiv*, 1–9 (2018).

63. L. Song, L. Florea, B. Langmead, Lighter: fast and memory-efficient sequencing error
correction without counting. *Genome Biol.* **15**, 509 (2014).

Acknowledgments: We would like to thank Y. Hara for advice in cell sorting, member of IGT.
Inc. for expert technical assistance in TCR sequencing, D. Komura and S. Ishikawa for
advice in data analysis. We also thank Dr. David L. Woodland for excellent English
language editing. Fig. 1A, B, Fig. 3A, and Fig. 4B were created with BioRender.com.

Funding:
Japan Society for the Promotion of Science, Grant-in-Aid for Scientific Research on
Innovative Areas 17929397 (KM)
Japan Society for the Promotion of Science, Grants-in-Aid for Scientific Research 20281832
(SU)
Japan Agency for Medical Research and Development grant, Research Program on Hepatitis
JP22fk0310509 (SU)
Japan Agency for Medical Research and Development grant, Project for Cancer Research
and Therapeutic Evolution JP22ama221306 (KM)
Japan Agency for Medical Research and Development grant, PRIME program
P21gm6210025 (SS)

Author contributions:

Conceptualization: HA, MK, TI, SU

Data curation: HA

Software: HA, SS

Formal analysis: HA

Methodology: HA, SS, AH, SU

Investigation: HA, MK, HA, AH, NOS, SU

Visualization: HA

Funding acquisition: TI, KM, SU

Project administration: TI, SU

Supervision: SS, KM

Writing – original draft: HA, SU

Writing – review & editing: HA, MK, HA, SS, AH, NOS, TI, KM, SU

Competing interests: H.A. reports stock for ImmunoGeneTeqs, Inc. S.S. reports advisory role for
ImmunoGeneTeqs, Inc; stock for ImmunoGeneTeqs, Inc, T.I. reports research funding
from Rohto Pharmaceutical Co., Ltd. and Biometrics Sympathies Inc. K.M. reports
consulting or advisory role for Kyowa-Hakko Kirin, ImmunoGeneTeqs, Inc; research
funding from Kyowa-Hakko Kirin, and Ono; stock for ImmunoGeneTeqs, Inc, IDAC
Theranostics, Inc. S.U. reports advisory role for ImmunoGeneTeqs, Inc; stock for
ImmunoGeneTeqs, Inc, IDAC Theranostics, Inc.

Data and materials availability: All sequencing data (Bulk TCRseq, and Single-cell RNA/TCR
sequencing) have been deposited at GEO and are publicly available as of the date of
publication (GEO: GSE210231). All original code has been deposited at Github
(https://github.com/hiro-aoki-mriid/Covid_TCRrepertoire). The list of antibodies, reagents,
and algorithms used in this study was summarized in Tables S5, 6, and 7, respectively.
960

Figures:

Figure 1

Fig. 1. **Bulk TCR repertoire analysis on blood CD8$^+$ T cells.** (**A**) Study design. (**B**) Schematic overview of grouping T-cell clones based on their kinetics profile. The top 150 clones for each participant (150 clones x 41 participants) were selected, and the correlation network of their kinetics profile was generated. Based on this network, modules consisting of highly correlated clones were identified. Then, patterns of kinetics profiles were defined based on the eigenvectors of modules. Finally, clones were classified based on their correlation with each pattern. (**C**) 12 kinetic profile patterns were identified in the CD8$^+$ T-cell repertoire. Kinetic profile patterns matching first (module Green (+)), second (module Brown (+)), and long-term responders (module Turquoise (+)) were highlighted. (**D**) Total frequency of CD8$^+$ clones classified into kinetic profile patterns at each time point. Color represents the name of module. (**E**) Comparison of CD8$^+$ clonal response between kinetic profile patterns. The extent of clonal response of each pattern was defined by the maximum changes in their total frequency through P1 to P4. Bars represent the median (n = 41). The mean rank of each pattern was compared with that of

977 pattern Brown (+) (***P<0.001; *P<0.05; n.s. non-significant, Dunn's multiple
978 comparisons test).
979

Figure 2

Fig. 2. Bulk TCR repertoire analysis on blood CD4$^+$ T cells. (**A**) Total frequency of CD4$^+$
clones classified to kinetic profile patterns at each time point. Color represents the name
of module. (**B**) Comparison of CD4$^+$ clonal response between kinetic profile patterns.
The extent of clonal response of each pattern was defined by the maximum changes in
their total frequency through P1 to P4. Bars represent the median (n = 41). The mean rank
of each pattern was compared with that of pattern Turquoise (+) (***P<0.001; n.s. non-
significant, Dunn's multiple comparisons test). First: First responders (module Turquoise
(+)). Second: Second responders (module Brown (+)). Long-term: Long-term responders
(module Green (+)).

Figure 3

A

B

C

Fig. 3. Searching Spike-reactive TCRs reported in public datasets. (**A**) Schematic overview.
(left) We identified TCRs reactive to SARS-CoV-2 epitopes from the three public
datasets. (middle) TCR meta-clonotypes were formed for each SARS-CoV-2 epitope.
Meta-clonotypes are represented as logo plots, where the sizes and colors of the letters
represent the frequency and characteristics of amino acids (red: acidic, blue: basic, green:
polar, purple: neutral, black: hydrophobic). (right) TCRs conforming to the definition of
meta-clonotypes were searched in our dataset. (**B**) Total frequency of CD8$^+$ SARS-CoV-
2 reactive meta-clonotypes at each time point. Bars represent the median (n = 41). Total
frequency of meta-clonotypes was compared between time points for each SARS-CoV-
2protein (***P<0.001;**P<0.005;*P<0.05, Dunn's multiple comparisons test). (**C**)

1002   Proportion of kinetic profile pattern among clones conformant to Spike-reactive or other
1003   SARS-CoV-2-reactive mata-clonotypes. Color represents the name of module.
1004

Figure 4

Fig. 4. **Analyzing CD8+ Spike-reactive T-cell clones using AIM assay.** (**A**) Gating strategy for sorting AIM+ and AIM- non-naive T cells. Proportion of AIM+ cells among non-naive T cells with or without Spike peptide stimulation was plotted. (**B**) Schematic overview for identifying AIM+ clones. After TCR sequencing on AIM+ and AIM- T cells, by-stander clones were excluded based on their enrichment in the AIM+ repertoire. (**C**) Scatter plot of overlapping clones between AIM+ and AIM- repertoire. Individual dots represent clones with their estimated cell count in AIM+ and AIM- repertoire. (**D**) Comparison of total frequency of CD8+ AIM+ clones between time points (n=15, ***P<0.001;**P<0.005, Dunn's multiple comparisons test). (**E**) (left) Proportion of kinetic profile pattern among CD8+ AIM+ and AIM- clones. (right) Comparison of the proportion of first-, second-, and long-term responders between CD8+ AIM+ and AIM- clones (n=14, **P<0.005; n.s. non-significant, Wilcoxon matched-pairs signed rank test).

Color represents the name of module. Participant ID34 was excluded from (E) and the following analysis, because only two clones were identified as AIM$^+$ clones. (**F**) Comparison of the proportion of CD8$^+$ AIM$^+$ clones among second-responders at P3 between AIM-high and AIM-low participants (n=7 of AIM-high and n=7 of AIM-low, ***P<0.001, Mann-Whitney test). (**G**) Scatter plot of the frequency of CD8$^+$ second-responders (x-axis) and AIM$^+$ second-responders (y-axis). (**H**) Total frequency of CD8$^+$ AIM$^+$ first-, second-, and long-term responders at each time point. Points represent the median (n = 14). (**I**) Percentage of CD8$^+$ top 10 AIM$^+$ clones at P3, which remained top 10 at P2 or P4. (D-F, H, and I) Bars represent the median.

Figure 5

Fig. 5. Analyzing CD4$^+$ Spike-reactive T-cell clones using AIM assay. (**A**) (left) Proportion of kinetic profile pattern among CD4$^+$ AIM$^+$ and AIM$^-$ clones. Color represents the name of module. (right) Comparison of the proportion of first-, second-, and long-term responders between CD4$^+$ AIM$^+$ and AIM$^-$ clones (n=15, **P<0.005; *P<0.05; n.s. non-significant, Wilcoxon matched-pairs signed rank test). (**B**) Comparison of the proportion of CD4$^+$ AIM+ clones among first-responders at P2 (left) or second-responders at P3 (right) between AIM-high and AIM-low participants (n=7 of AIM-high and n=8 of AIM-low, **P<0.005; n.s. non-significant, Mann-Whitney test). (**C**) Total frequency of CD4$^+$ AIM+ first-, second-, and long-term responders at each time point. Points represent the median (n = 15). (**D**) Percentage of CD4$^+$ top 10 AIM+ clones at P3, which remained top 10 at P2 or P4. (A and B) Bars represent the median.

Figure 6

1041

Fig. 6. Single-cell analysis on non-naive CD8⁺ T cells after the second dose. (**A**) Weighted-nearest neighbor graph-based uniform manifold approximation and projection (wnnUMAP) plot of the scRNAseq/AbSeq data of non-naive CD8⁺ T cells. (**B** and **C**)

Violin plot showing the mRNA (B) and protein (C) expression of marker genes. (**D**) wnnUmap plot split by the two participants (ID2 and ID27). (**E**) wnnUmap plot for T cells in first-, second-, and long-term responder clones. (**F**) Cluster distribution of T cells in first-, second-, and long-term responder clones. Distribution of T cells in clones with other kinetic profile patterns was also shown ("Other"). (**G**) Heat map for the cluster distribution of expanded clones ($\geq$ 10 cells per clone). Each cell represents the proportion of each cluster (row) in each clone (column). The source ("ID") and kinetic profile ("Kinetics") of clones are also shown as heatmap annotation. (**H**) wnnUmap plot for T cells in AIM$^+$ or AIM$^-$ clones. Total count of cells in each plot is also shown. (**I**) Cluster distribution of T cells in AIM$^+$ or AIM$^-$ clones. (**J**) wnnUmap plot for T cells in individual expanded AIM+ clones ($\geq$ 10 cells per clone). Clone name (source, rank in clones detected by scTCR, and dominant phenotype) was shown on the top of the plot. (**K**) Kinetic profile of AIM$^+$ expanded clones. The frequency of clones at each time point in bulk TCR repertoire data is plotted. IL: innate-like; Act: activated; TD: terminally differentiated; MP: memory precursor; PEX: progenitor exhausted-like.

Figure 7

1061

Fig. 7. Single-cell analysis of non-naive CD4⁺ T cells after the second dose. (A) wnnUMAP
plot of the scRNAseq/AbSeq data of non-naive CD4⁺ T cells. (**B** and **C**) Violin plot
showing the mRNA (B) and protein (C) expression of marker genes. (**D**) wnnUmap plot
for T cells in AIM⁺ or AIM⁻ clones. Total count of cells in each plot is also shown. (**E**)
Cluster distribution of T cells in AIM⁺ or AIM⁻ clones. (**F**) wnnUmap plot for T cells in
first-, second-, and long-term responder clones. (**G**) Cluster distribution of T cells in
first-, second-, and long-term responder clones. Distribution of T cells in clones with
other kinetic profile patterns was also shown ("Other"). CM: central memory; EM:
effector memory; Act: activated; Treg: regulatory T cells.

1071

34