Kinetics-based Inference of Environment-Dependent Microbial Interactions and Their Dynamic Variation

Hyun-Seob Song[1,2][†][*], Na-Rae Lee[3][†], Aimee K. Kessell[1], Seo-Young Park[4], Kang Zhou[5], and Dong-Yup Lee[4][*]

[1]Department of Biological Systems Engineering, University of Nebraska-Lincoln, Lincoln, NE 68583

[2]Department of Food Science and Technology, Nebraska Food for Health Center, University of Nebraska-Lincoln, Lincoln, NE 68588

[3]Department of Bioscience and Biotechnology, Konkuk University, Seoul 05029, Republic of Korea

[4]School of Chemical Engineering, Sungkyunkwan University, Suwon-si, Gyeonggi-do 16419, Republic of Korea

[5]Department of Chemical and Biomolecular Engineering, National University of Singapore, 4 Engineering Drive 4, Singapore 117585, Singapore

[†]Both authors contributed equally.

[*]Correspondence: hsong5@unl.edu (H.-S.S.); dongyuplee@skku.edu (D.-Y.L.)

ABSTRACT: Microbial communities in nature are dynamically evolving as member species

change their interactions subject to environmental variations. Accounting for such context-

dependent dynamic variations in interspecies interactions is critical for predictive ecological

modeling. However, we lack a fundamental understanding of how microbial interactions are driven

by environmental factors due to the absence of generalizable theoretical foundations, significantly

limiting our capability to predict and engineer community dynamics and function. To address this

issue, we propose a novel theoretical framework that allows us to represent interspecies

interactions as an explicit function of environmental variables (such as substrate concentrations)

by combining growth kinetics and a generalized Lotka-Volterra model. A synergistic integration

of these two complementary models leads to the prediction of alterations in interspecies

interactions as the outcome of dynamic balances between positive and negative influences of

microbial species in mixed relationships. This unique capability of our approach was

experimentally demonstrated using a synthetic consortium of two *Escherichia coli* mutants that

are metabolically dependent (due to an inability to synthesize essential amino acids), but

competitively growing on a shared substrate. The analysis of the *E. coli* binary consortium using

our model not only showed how interactions between the two amino acid auxotrophic mutants are

controlled by the dynamic shifts in limiting substrates, but also enabled quantifying previously

uncharacterizable complex aspects of microbial interactions such as asymmetry in interactions.

Our approach can be extended to other related ecological systems to model their environment-

dependent interspecies interactions from growth kinetics.

IMPORTANCE: Modeling of environment-controlled interspecies interactions through separate

identification of positive and negative influences of microbes in mixed relationships is a new

capability that can significantly improve our ability to understand, predict, and engineer complex

dynamics of microbial communities. Moreover, robust prediction of microbial interactions as a

function of environmental variables can serve as valuable benchmark data to validate modeling

and network inference tools in microbial ecology, the development of which has often been

impeded due to the lack of ground truth information on interactions. While demonstrated against

microbial data, the theory developed in this work is readily applicable to general community

ecology to predict interactions among microorganisms such as plants and animals, as well as

microorganisms.

KEYWORDS: Microbial communities; competition; cooperation; context dependence; kinetic

models; Lotka-Volterra models

INTRODUCTION

Microbial communities play pivotal roles in maintaining human and animal health, plant productivity, and ecosystem services (1-4). Increasing efforts are being dedicated towards maximizing their beneficial roles in natural systems or creating new industrial applications (5). However, control and design of microbial community dynamics and function is a challenging task, primarily due to higher-order or emergent properties that are not observable from individual species in isolation but arise through nonlinear interspecies interactions (6, 7). Therefore, rational design of microbial communities or consortia requires a fundamental knowledge of microbial interactions as a mechanistic linkage between the environment and the community compositions and function, necessitating the employment of predictive mathematical models as indispensable tools (8-14).

Development of accurate models of microbial communities that are commonly subject to environmental variations is truly complicated by the following intrinsic ecological aspects. First, microorganisms in a community build *dynamic* interactions that cannot effectively be represented by a rigid network with fixed structure (15, 16). Rather, microbial communities keep reorganizing interaction networks in response to biotic or abiotic perturbations or through adaptation to long-lasting environmental changes. Second, microorganisms often build *mixed* relationships by exerting *both* promotive and inhibitive impacts on the growth of their partners/neighbors (17, 18). Individual identification of these simultaneously acting positive and negative interactions is critical because community dynamics is mainly driven by the balances between all counteracting impacts among member species (19). The lack of capability to account for these key properties of microbial interactions limits our ability to predict and engineer microbial community dynamics and functions.

Despite rapid progress in microbiome science, we still do not know how to identify

environment-controlled dynamic variation in interspecies interactions addressed above. Three

major branches of microbial interaction modeling include (20, 21): (i) network inference, (ii)

metabolic network modeling, and (iii) kinetic modeling. Network inference is widely used for

modeling microbial interactions to identify interaction networks based on correlative relationships

among microbial populations (22-25), parameter identification through regression (26-28), or a

prescribed set of rules or hypotheses (21). The resulting networks represent interspecies

interactions as single constant metrics, therefore being unable to describe dynamic variations in

interactions nor identify the balances among counteracting individual impacts in mixed

relationships. Recently, we developed the method termed MIIA (Minimal Interspecies Interaction

Adjustment) (15, 16) to predict context-dependent interactions due to the changes in memberships,

which however has not been extended to address the environmental impacts. In contrast with

network inference, metabolic network and kinetic modeling can account for both positive and

negative interactions based on cross-feeding of small molecules (essential for growth) or

competition for shared substrates/nutrients among species; in theory, kinetic models can

additionally simulate their dynamic variations. While more mechanistic than network inference,

these methods cannot quantify the magnitude or even the sign of net interactions.

In this work, we fill these gaps by proposing a novel theoretical framework that enables a

quantifiable, mechanistic representation of the dynamic linkage between microbial interactions

and the environment. For this purpose, we synergistically integrate two complementary modeling

frameworks to overcome their own limitations: a generalized Lotka-Volterra (gLV) model (29)

and population growth kinetics. Like other network inference approaches, a typical gLV model

with a focus on pairwise interactions is constructed based on an implicit assumption of constant

interactions. We relax this assumption by representing interaction coefficients in the gLV model

as a function of environmental variables (i.e., concentrations of cross-fed metabolites and shared

substrates) described in microbial growth kinetics, which is termed here kinetics-based inference

of dynamic variation in microbial interactions (KIDI). The resulting functional representation of

interactions by KIDI enables not only quantifying their dynamic variation as environmental

conditions change, but also individually identifying negative and positive influences among

species in mixed relationships. The prediction of KIDI was demonstrated accurate through a

coordinated design of experiments using a binary consortium composed of tyrosine and tryptophan

auxotrophic mutants of *Escherichia coli* (30) so that they both compete and/or cooperate

depending on environmental conditions.

RESULTS

Formulation of a conceptual model for understanding environment-dependent interactions

For illustration of the concept of KIDI, we consider a hypothetical consortium composed of two

members where species 1 (X_1) and species 2 (X_2) cooperate by cross-feeding S_1^+ and S_2^+ each other

but compete for the shared metabolite S^- (the center circle in **Fig. 1**). Growth kinetics for the i^{th}

species (X_i) (that requires two substrates S_i^+ and S^- for growth) can be represented, e.g., using a

double Michaelis-Menten equation as follows:

$$\mu_i = \mu_i^{\max} \frac{s_i^+}{\left(K_i^+ + s_i^+\right)} \frac{s^-}{\left(K_i^- + s^-\right)}, \quad i = 1, 2 \tag{1}$$

where μ_i [1/h] is the specific growth rate of X_i, μ_i^{max} is the maximal specific growth rate, s_i^+ and

s^- [g/l] are the concentrations of S_i^+ and S^-, and K_i^+ and K_i^- [g/l] are half-saturation constants

associated with the consumption of S_i^+ and S^-, respectively. As inferable from growth kinetics in

Eq. (1), the mixed relationship (i.e., competition and cooperation) between X_1 and X_2 when both

substrates are limiting can turn into diverse forms of interactions as environmental conditions

change. When S^- is present in excess (therefore, no competition is necessary) but S_1^+ and S_2^+ are

limiting, for example, their relationship is predominantly cooperative (where $\mu_i \approx \frac{\mu_i^{\max} s_i^+}{K_i^+ + s_i^+}$). As the

opposite case, if both S_1^+ and S_2^+ are excessive in the environment (so no partners are needed to

acquire them) while S^- is limiting, their relationship is governed by competition (where $\mu_i \approx$

$\frac{\mu_i^{\max} s^-}{K_i^- + s^-}$). Likewise, one can assume many other different scenarios where their relationships turn

into competition, cooperation, amensalism, commensalism and even neutrality, as illustrated in

Fig. 1.

Representation of interaction parameters as a function of environmental variables

To model such environment-dependent microbial relationships, we derived a general form of

interaction coefficients as a function of environmental variables by integrating growth kinetics and

a gLV model. As described in detail in **Materials and Methods**, our formula (KIDI) represents

interaction coefficients of species in the mixed relationship as a sum of positive and negative parts,

i.e.,

120
$$a_{i,j}\left(s_i^+, s^-\right) = a_{i,j}^+\left(s_i^+, s^-\right) + a_{i,j}^-\left(s_i^+, s^-\right), \quad (i, j) = (1, 2) \text{ or } (2, 1) \tag{2}$$

121 where $a_{i,j}^+$ and $a_{i,j}^-$ denote the positive and negative influence of X_j on the growth rate of X_i, which

122 are defined as follows:

123
$$a_{i,j}^+\left(s_i^+, s^-\right) \equiv \frac{\partial}{\partial s_i^+}\left[\mu_i\left(s_i^+, s^-\right)\right] \cdot \frac{\partial s_i^+}{\partial x_j} \tag{3}$$

124
$$a_{i,j}^-\left(s_i^+,s^-\right) \equiv \frac{\partial}{\partial s^-}\left[\mu_i\left(s_i^+,s^-\right)\right]\cdot\frac{\partial s^-}{\partial x_j} \tag{4}$$

The positive influence of X_j on the growth rate of X_i (i.e., $a_{i,j}^+$) is represented by the two

subsequent terms on the right-hand side of Eq. (3): (i) the impact of the change in the population

size of X_j on the concentration of the cross-fed substrate S_i^+ (as denoted by $\partial s_i^+/\partial x_j$) and (2) the

subsequent impact of the change in S_i^+ on the growth rate of the i^{th} species (i.e., μ_i) (as denoted

by $\partial[\mu_i(s^-,s_i^+)]/\partial s_i^+$). The negative impact of X_j on the growth rate of X_i ($a_{i,j}^-$) in Eq. (4) can be

interpreted in a similar fashion.

The derivative terms on the right-hand side of Eqs. (3) and (4) are fully identifiable from

reaction stoichiometry and kinetics. In the case of using a double Monod kinetics, for example,

incorporation of Eq. (1) into Eqs. (3) and (4) yields $a_{i,j}^+$ and $a_{i,j}^-$ as follows:

134
$$a_{i,j}^+\left(s_i^+,s^-\right) = \left[\mu_i^{max}\frac{K_i^+}{\left(K_i^+ + s_i^+\right)^2}\frac{s^-}{\left(K_i^- + s^-\right)}\right]\cdot Y_{S_i^+/X_j} \tag{5}$$

135
$$a_{i,j}^-\left(s_i^+,s^-\right) = \left[\mu_i^{max}\frac{s_i^+}{\left(K_i^+ + s_i^+\right)}\frac{K_i^-}{\left(K_i^- + s^-\right)^2}\right]\cdot\left(-Y_{S^-/X_j}\right) \tag{6}$$

136 where $Y_{S_i^+/X_j}$ and Y_{S^-/X_j} denote the stoichiometric relationships between the changes in substrate

137 and biomass concentrations associated with X_j, i.e., $Y_{S_i^+/X_j} = |\Delta s_i^+/\Delta x_j|$ and $Y_{S^-/X_j} = |\Delta s^-/\Delta x_j|$

138 (see **Materials and Methods**).

139   To identify net interactions between two species with mixed relationships, we further

140 defined a universal interaction parameter $\gamma_{i,j}$ as follows:

141
$$\gamma_{i,j} \equiv \frac{a_{i,j}^+ + a_{i,j}^-}{a_{i,j}^+ - a_{i,j}^-}, \quad (i,j) = (1,2)\text{ or }(2,1) \tag{7}$$

The parameter $\gamma_{i,j}$ ranges from -1 to 1 to represent positive influences of species j on species i

when greater than 0 and negative impacts when less than 0, respectively, consequently allowing

us to conveniently quantify the relative dominance of inhibition vs. promotion in mixed

interactions. The parameter $\gamma_{i,j}$ complements $a_{i,j}$, rather than replaces it, in that the magnitude of

interactions cannot be determined by $\gamma_{i,j}$, but by the original interaction parameter, $a_{i,j}$. Along

with $a_{i,j}$ defined in Eqs. (2) to (4), $\gamma_{i,j}$ completely characterizes the dynamic variation of

interactions between X_1 and X_2 based on co-culture growth data as demonstrated in the following

sections.

Identification of kinetics and stoichiometry via data fit

For experimental demonstration of the mathematical formulation derived in the previous section,

we constructed a synthetic consortium composed of two *E. coli* auxotrophic mutants that

cooperatively cross-feed tryptophan and tyrosine, respectively, while competitively growing on

glucose (31). This consortium is therefore an ideal, simplest model system for studying

environment-dependent dynamic variations in microbial interactions. Due to its exact

correspondence to the hypothetical consortium in **Fig. 1**, we denote two *E. coli* mutant strains

ΔtrpC and ΔtyrA by X_1 and X_2; glucose, tryptophan, and tyrosine by S^-, S_1^+, and S_2^+, respectively.

Using these two strains, we performed growth experiments under diverse culture

conditions: two individual batch experiments using X_1 (**Fig. 2A**) and X_2 (**Fig. 2B**), respectively,

and two sets of co-culture experiments (**Figs. 2C** and **2D**). The top panels in **Figs. 2C** and **2D**

denote co-growth experiments in batch cultures, while the middle and bottom panels denote semi-

batch modes, where we added glucose feedbeads (FBs) at time 7.5 and 10 hours, respectively, to

induce dramatic changes in interspecies interactions during co-growth. Other differences in co-

culture conditions in **Figs. 2C** and **2D** include initial concentrations of S^-, S_1^+, and S_2^+, and the

number of added FBs (see **Supp Datasheet**). The optical density profiles in **Fig. 2D** denote the

combined population change of both strains, i.e., $X_1 + X_2$.

Based on the four datasets in **Fig. 2**, we constructed a dynamic co-growth model of X_1 and

X_2 to determine associated kinetics and stoichiometry, key information required for quantifying

interspecies interactions parameters ($a_{i,j}$ and $\gamma_{i,j}$) in Eqs. (2) to (7). The dynamic co-growth model

is composed of five mass balance equations for X_1, X_2, S^-, S_1^+, and S_2^+. We determined

stochiometric and kinetic parameters using three subsets of data in **Figs. 2A – 2C** and validated

the model against the remaining one (in **Fig. 2D**) that was not used for model identification. The

robust consistency between simulated and measured data in **Fig. 2D**, as well as those in **Figs. 2A**

– 2C, indicates the acceptability of using the identified model parameters in inferring interaction

coefficients. We provide the full list of model equations with parameter values in **Table 1**; the

culture conditions in **Table S1**.

Variation in microbial interactions driven by the switch in limiting substrates in batch

cultures

Based on the stochiometric and kinetic parameters determined through data fit in **Table 1**, we were

able to determine microbial interactions and their variations as a function of environmental

conditions using KIDI. We first analyzed various co-culture scenarios in batch modes (**Fig. 3**). In

Fig. 3A, we considered the growth of X_1 and X_2 on relatively high and low initial concentrations

of S^- (2 g/l) and S_1^+ and S_2^+ (i.e., 0.4 mg/l for both) as a reference condition. As shown in the top

two panels of **Fig. 3A**, S^- was depleted around 22.5 hours, after which no further microbial growth

is observed for both X_1 and X_2. In the present setting, the initial relationship between X_1 and X_2 is

expected to be cooperative (because S^- is excessive in the beginning), which however turns into

competition as the level of S^- gradually decreases. Interaction coefficients extracted by KIDI

correctly captured this transition as indicated by the dramatic changes in the values of normalized

universal interaction parameters ($\gamma_{1,2}$ and $\gamma_{2,1}$) from 1 to -1 around the time when glucose is

depleted. Actual values of interaction coefficients can be seen from $a_{i,j}$, $a_{i,j}^+$, and $a_{i,j}^-$ (three bottom

panels of **Fig. 3A**, which also showed that the level of initial cooperation was moderate or small

(as indicated by small magnitudes of $a_{i,j}^+ < 2 \times 10^{-3}$), partly due to the availability of S_1^+ and S_2^+

in the environment from the beginning. Interestingly, the values of $a_{i,j}^+$ dropped to zero after the

depletion of S^- despite that the initially added S_1^+ and S_2^+ might be completely consumed by that

time and consequently metabolic dependence between X_1 and X_2 should increase. The reason for

zero values of $a_{i,j}^+$ is due to that the exchange of S_1^+ and S_2^+ is not occurring when species cannot

grow any more due to carbon limitation after the depletion of S^-.

For comparison, we analyzed two additional conditions. (1) with a lower initial

concentration of S^- (i.e., 0.5 g/l) and (2) with lower and higher initial concentrations of S^- (0.5

g/l) and S_1^+ and S_2^+ (i.e., 4 mg/l for both), respectively. In both cases, KIDI successfully captured

the expected changes in interspecies interactions. In the case of lowering the initial concentration

of S^- (**Fig. 3B**), KIDI showed that the level of initial competition increases (due to the limited

availability of S^-) as indicated by relatively lower values of $\gamma_{1,2}$ and $\gamma_{2,1}$ compared to the case of

Fig. 3A. Notably, $\gamma_{2,1}$ showed negative value throughout the co-growth (indicating the dominance

of negative influence of X_1 on the growth of X_2). In the case of increasing the initial concentrations

of S_1^+ and S_2^+ in addition to lowering S^- (**Fig. 3C**), the relationship between the two strains became

even more negative, which was also an expected outcome because metabolic dependence between

X_1 and X_2 will accordingly reduce when they can acquire what they need from the environment,

rather than from partners.

In all three cases above (including the reference condition), the relations between the two

E. coli strains were shown asymmetric, i.e., $\gamma_{1,2} \neq \gamma_{2,1}$, $a_{1,2} \neq a_{2,1}$, $a_{1,2}^- \neq a_{2,1}^-$, and $a_{1,2}^+ \neq a_{2,1}^+$.

Interestingly, KIDI consistently predicted $\gamma_{1,2} > \gamma_{2,1}$ and $a_{1,2} > a_{2,1}$, while the reasons may vary

across conditions. In the reference condition (**Fig. 3A**) where glucose is excessive (so $a_{i,j}^-$'s are

negligible), it is mostly due to $a_{1,2}^+ > a_{2,1}^+$ (i.e., X_1 has a higher comparative advantage in

exchanging amino acids with X_2 than the other way around) that leads $\gamma_{1,2} > \gamma_{2,1}$ (as well as

$a_{1,2} > a_{2,1}$) (see **Fig. S1A** for a zoom-in view). In contrast, in the case in **Fig. 3C** where the glucose

level is low while amino acids are abundant (so $a_{i,j}^+$'s are negligible), X_1 has the comparative

growth advantage because $a_{1,2}^- > a_{2,1}^-$ (i.e., the growth of X_1 is less inhibited by X_2 than the other

way around) (**Fig. S1C**). Lastly, both $a_{i,j}^+$'s and $a_{i,j}^-$'s contribute to the outcome in the case in **Fig.**

3B where all substrates (glucose and amino acids) are limitedly available in the environment (**Fig.**

S1B).

Dynamic response of microbial interactions to environmental perturbations during growth

We extend our analysis to semi-batch cultures that are perturbed by the addition of glucose FBs

during growth and therefore are expected to show more dramatic changes in interspecies

interactions and community dynamics. In contrast with the batch cultures considered in the

previous section where no further growth is possible after the depletion of the initially added S^-,

the two strains continue to grow in semi-batch cultures due to slow but continual provision of S^-

from the added FBs. Despite a general expectation that the competition level between the two

strains will be mitigated at least at the moment of FB addition, it is uncertain: (1) to what degree

this will occur under different environmental conditions, and (2) how governing microbial

interactions will shift (between competition and cooperation), particularly in a later phase when

the growth of the two strains is be limited by both S^- and S_i^+. To answer these questions, we

applied KIDI to the following three cases. For simplicity, we set the initial conditions to be the

same as before.

First, we considered the initial concentrations of 2 g/l for S^- and 0.4 mg/l for S_1^+ and S_2^+

and added 3 FBs of S^- at around 7.5 hours (**Fig. 4A**). Due to the relatively high concentration of

S^-, the impact of adding 3 FBs of S^- was minimal, i.e., showing no appreciable qualitative

changes (in glucose concentration, population size, and interaction parameters) from the batch case

with the same initial conditions in **Fig. 3A**. By contrast, when the initial concentration of S^- was

low (i.e., 0.5 g/l) (**Fig. 4B**), KIDI identified the significant impact of adding FBs on both glucose

concentration and microbial interactions, as indicated by sudden increases in S^-, $\gamma_{1,2}$ and $\gamma_{2,1}$.

More interestingly, when additionally increasing the initial concentration of S_1^+ and S_2^+ (**Fig. 4C**)

(therefore the level of competition becomes even more significant), we found no appreciable

impacts on microbial interactions ($\gamma_{1,2}$ and $\gamma_{2,1}$), while the glucose (S^-) profile was significantly

perturbed by adding FBs. As observed in the previous section, the *net* interaction parameters ($a_{i,j}$

and $\gamma_{i,j}$) are governed by $a_{i,j}^+$ and/or $a_{i,j}^-$ depending on the environmental contexts: $a_{i,j}^+$ and $a_{i,j}^-$

are major determinants of the net interaction for the cases in **Figs. 4A** and **4C**, respectively, while

the effects of $a_{i,j}^+$ and $a_{i,j}^-$ are comparable for the case in **Fig. 4B** (see zoom-in views in **Fig. S2**).

In all three cases, the interactions between the two strains were eventually governed by competition

in the later phase as indicated by the negative values of $\gamma_{1,2}$ and $\gamma_{2,1}$ close to -1. However,

governing interactions in the early phase depended on initial conditions, i.e., cooperation and

competition were dominant in the first and third cases (**Figs. 4A** and **4C**), respectively, while the

second case (**Fig.4B**) showed a balance between them. Similar interpretations can be made for

other scenarios of perturbations where the number of added FBs and the timing of addition were

subject to variation (**Fig. S3**).

Interactions between X_1 and X_2 were also identified asymmetric. Similar to the previous

batch cases, X_1 has the comparative advantage over X_2 throughout the entire growth period by

being more helped from X_2 (than X_2 is being helped from X_1) (i.e., $a_{1,2}^+ > a_{2,1}^+$) before the

exhaustion of initially added S^-, and by being less inhibited by X_1 than X_2 is being inhibited by

X_1 (i.e., $a_{1,2}^- < a_{2,1}^-$) afterwards.

DISCUSSION

In this study, we proposed a novel computational method (KIDI) that enables quantitatively

identifying environment-dependent interspecies interactions in microbial communities. By

integrating growth kinetics into a gLV model, we derived an analytical form of interaction

coefficients as a function of environmental variables (i.e., concentrations of chemical substrates

that affect interactions), the results of which were subsequently validated through a coordinated

design of co-culture experiments.

Our theoretical development significantly extends the current scope of microbial ecological

modeling by completely relaxing the typical assumption of constant interactions among species.

The gLV model, for example, has been widely used as a basic ecological modeling template for

simulation of population dynamics and inference of interspecies interactions in microbial

communities (26-28). Due to the constant interaction assumption, however, the application of the

gLV model is often confined to a narrow range of conditions where interspecies interactions are

expected to remain largely constant. KIDI addresses this limitation by representing interaction

coefficients as an explicit function of limiting substrates. As an exception, a previous study by

Momeni et al. (32) showed that pairwise interaction (i.e., gLV) models are derivable from

mechanistic (i.e., kinetic) models through empirical manipulation of equations, which is however

limited to special forms of kinetics and therefore cannot be generalizable (32). By contrast, our

chain rule-based formulation allows us to handle any complex forms of kinetic equations with no

such constraints. Consequently, KIDI enables incorporation any forms of kinetic equations as

demonstrated using a double Michaelis-Menten kinetics as a demonstration example.

Dynamic variations in microbial interactions inferred by KIDI were experimentally

validated using a synthetic binary consortium of two metabolically engineered auxotrophic *E. coli*

mutants that cross-feed amino acids they cannot synthesize (i.e., tryptophane and tyrosine). A

coordinated design of experiments provided multiple sets of data required for determining kinetic

and stoichiometric parameters in the mechanistic model along with substrate concentrations, which

are key inputs for quantifying environment-dependent interactions. Despite diverse culture

conditions including axenic and binary growth in batch and semi-batch modes, our model with a

single set of parameters showed satisfactory fit to the three training datasets and provided

consistency with the validation dataset set aside in advance. Such a fair goodness of fit indicates

the acceptability of model parameters and therefore the subsequent inference of microbial

interactions.

Our kinetic model also shows consistency with the analysis of energetic cost of

synthesizing amino acids in the literature. Mee et al. (30) estimated the energetic cost for the

synthesis of 14 individual amino acids based the amounts of extracellularly supplemented amino

acid and the observed growth yield of *E. coli* auxotrophic mutants. From the linear relationships

between these two variables, they calculated the supplemented amounts of amino acids *per cell*,

which were 1.5×10^7 and 3.7×10^7 for the tryptophane and tyrosine auxotrophic *E. coli* mutants,

respectively. These two quantities correspond to the stoichiometric coefficients $Y_{S_i^+/X_i}(i = 1,2)$

in our kinetic model, which were determined to be 0.0845 $(= Y_{S_1^+/X_1})$ and 0.133 $(= Y_{S_2^+/X_2})$

[mg/OD] through data fit (**Table 2**). As the direct one-to-one matching between them might not

be feasible, e.g., due to different units of biomass (i.e., cell number in Mee et al. (30) vs. OD in

this work), we compared the ratios, which showed consistency between the two studies, i.e.,

$\frac{1.5 \times 10^7}{3.7 \times 10^7} \approx 0.41$ vs. $\frac{0.0845}{0.133} \approx 0.64$. Both results imply that compared to tyrosine, the synthesis of

tryptophane is more costly. In support of this, Mee et al. (30) estimated the biosynthetic cost for

tryptophane is about 43% higher than that for tyrosine.

The formulation of KIDI is valid regardless of the complexity of microbial communities.

A challenge exists, however, in accurately quantifying input parameters (i.e., kinetic and

stoichiometric parameters of the kinetic model), because it requires robust measurements of

temporal profiles of substrate concentrations. In particular, the levels of metabolites that are

exchanged among member species are often quite low and sometimes below the detection limit

due to immediate consumption after being produced. This difficulty would be relieved to a degree

by performing axenic growth experiments (as we did in this work) but extracting isolates from

natural communities may not be always possible. Under these constraints, our demonstration was

focused on relatively simple consortia that facilitate to collect experimental data (temporal profiles

of substrate concentrations and population densities) required to determine the parameters in the

mechanistic model. In this regard, KIDI can serve as a valuable modeling tool for synthetic

consortia designed for engineering applications and/or model consortia derived from natural

communities to gain a better understanding of complex ecological systems (33, 34).

Despite this challenge, we highlight that inferring environment-dependent interactions and

their dynamic variations is a critical capability uniquely associated with KIDI. Even in a simple

binary consortium considered in this work, KIDI provides new insights into interspecies

interactions such as asymmetry between the two amino acid autotrophs, which might not be

obtainable otherwise. In perturbed growth experiments with glucose FBs, for example, KIDI

identified that (1) $a_{1,2}^+ > a_{2,1}^+$ while the shared substrate (glucose) is abundant, implying that the

tryptophane auxotroph (X_1) does not support the growth of the tyrosine auxotroph (X_2) as much

as X_2 does for X_1; (2) $a_{1,2}^- < a_{2,1}^-$ after the completion of initially added glucose, implying that less

favorable supporters during cooperation become worse enemies when the relationship turned into

competition. As another critical utility, KIDI can significantly facilitate the development of

network inference techniques because the validation process of new algorithms for predicting

microbial interactions has often been hampered due to the lack of benchmark data.

Beyond microbial ecology, KIDI is applicable in broad areas of community ecology

because context dependency is not a property associated only with microorganisms, but with

macro-organisms such as plants and animals. We hope that KIDI will serve as a useful tool for

understanding and engineering context-dependent microbial interactions in a wide range of

applications.

MATERIALS AND METHODS

Mathematical definition of interaction coefficients

The dynamic change in population i in a community can be formulated in a general form as follows:

345
$$\frac{1}{x_i}\frac{dx_i}{dt} = f_i\left(x_1, x_2, \cdots, x_N\right), \quad i = 1, 2, \cdots, N \tag{8}$$

where x_i is the population density of species i, the left-hand side defines a specific growth rate of

species i (hereafter denoted by $\mu_i \equiv d(\ln x_i)/dt$), and the function $f_i(x_1, x_2, \cdots, x_N)$ represents a

nonlinear dependence of μ_i on population densities of other species that affect the growth of

species i.

Using a Taylor expansion, the right-hand side of Eq. (8) can be represented as a series of

polynomial terms, i.e.,

$$\mu_i = f_{i,0} + \sum_{j=1}^{N} \left(\frac{\partial f_i}{\partial x_j} \right)_0 x_j + H.O.T., \quad i = 1, 2, \cdots, N \tag{9}$$

353   where the subscript 0 denotes a chosen reference condition, *H.O.T.* is higher-order terms.

354   Neglecting the *H.O.T.* in Eq. (9), a gLV equation describes specific growth of species i using a

355   linear equation, i.e.,

$$\mu_i (x_1, x_2, \cdots, x_N) = \mu_{i,0} + \sum_{j=1}^{N} a_{i,j} x_j, \quad i = 1, 2, \cdots, N \tag{10}$$

357   where interaction coefficient $a_{i,j}$ denotes the effect of species population j on the specific growth

358   of species i. For a binary community, Eq. (10) reduces to

$$\mu_i (x_i, x_j) = \mu_{i,0} + a_{i,i} x_i + a_{i,j} x_j, \quad i = 1, 2 \tag{11}$$

360   where $\mu_{i,0}$ is the basal growth rate, $a_{i,i}$ is the intra-specific interaction coefficient, and $a_{i,j}$ is the

361   inter-specific interaction coefficient.

362   From Eq. (11), binary interaction coefficients in gLV are defined as follows:

$$a_{i,j} \equiv \frac{\partial \mu_i (x_i, x_j)}{\partial x_j} \tag{12}$$

The gLV model assumes that $a_{i,j}$ is constant, which however leads the gLV model to fail to capture

delicate dynamics of microbial interactions. Indeed, $a_{i,j}$ is a dynamic parameter that changes its

value in varying environmental conditions as shown in the next section.

Formulation of interaction coefficients as function of environmental variables

Specific growth rates can also be kinetically represented as a function of nutrient concentrations

in the environment. In the circumstance considered in **Fig. 1**,

371
$$\mu_i = \mu_i\left(s_i^+, s^-\right) \tag{13}$$

372   where s_i^+ is the nutrient (such as Trp or Tyr) that species i needs to get either from its partner or

373   the environment, and s^- represents the shared nutrient (i.e., glucose) that two species compete for.

374   Based on the chain rule, we formulate $a_{i,j}$ as a function of nutrient concentrations by

375   plugging Eq. (13) into (12), i.e.,

376
$$\begin{aligned} a_{i,j} &= \frac{\partial}{\partial x_j}\left[\mu_i\left(s_i^+, s^-\right)\right] \\ &= \frac{\partial}{\partial s_i^+}\left[\mu_i\left(s_i^+, s^-\right)\right]\frac{\partial s_i^+}{\partial x_j} + \frac{\partial}{\partial s^-}\left[\mu_i\left(s_i^+, s^-\right)\right]\frac{\partial s^-}{\partial x_j} \end{aligned} \tag{14}$$

377   Note that the two terms on the R.H.S. of Eq. (14) represent the positive and negative effects of

378   species j on i through environmental variables, i.e., $a_{i,j}^+$ and $a_{i,j}^-$ as defined below

379
$$a_{i,j}^+ \equiv \frac{\partial}{\partial s_i^+}\left[\mu_i\left(s_i^+, s^-\right)\right]\frac{\partial s_i^+}{\partial x_j} \tag{15}$$

380
$$a_{i,j}^- \equiv \frac{\partial}{\partial s^-}\left[\mu_i\left(s_i^+, s^-\right)\right]\frac{\partial s^-}{\partial x_j} \tag{16}$$

381   In a similar fashion, we can formulate intra-specific interaction coefficients as functions of

382   environmental variables, i.e.,

$$a_{i,i} = \frac{\partial}{\partial x_i}\left[\mu_i\left(s_i^+, s^-\right)\right]$$

$$= \frac{\partial}{\partial s_i^+}\left[\mu_i\left(s_i^+, s^-\right)\right]\frac{\partial s_i^+}{\partial x_i} + \frac{\partial}{\partial s^-}\left[\mu_i\left(s_i^+, s^-\right)\right]\frac{\partial s^-}{\partial x_i}$$

383   (17)

Final forms of $a_{i,j}(a_{i,j}^+$ and $a_{i,j}^-)$ and $a_{i,i}$ depend on specific kinetics for $\mu_i(s^-, s_i^+)$.

Microorganisms and culture conditions

Tyrsoine (Tyr) and Trptophan (Trp) auxotrophic *Escherichia coli* (*E.* coli) strains were purchased

from *E. coli* Genetic Stock Center (CGSC) at Yale University (http://cgsc2.biology.yale.edu/).

Each strain was incubated overnight at 37°C and 225 rpm in 50 ml of Falcon tube containing 5 ml

of Lysogeny broth (LB) supplemented with 33 µg/l kanamycin. Culture cells were collected and

centrifuged them at 16,000 g, 4°C for 1.5 min. The cell pellets were washed with K3 basal medium

to remove residual amino acids in the samples. The washed cells were resuspended and transferred

to 150 ml flasks carrying 25 ml of K3 defined minimal medium (5) containing glucose and 33 µg/l

kanamycin, and cultivated at 37°C and 225 rpm. An initial absorbance at 600 nm (OD 600) was

0.04 with equivalent cell ratio. For batch mode, 4.5 g/l glucose was supplied in the culture medium.

For the fed-batch mode, 0.5 g/l of an initial glucose concentration was used to shorten the lag phase

and 3 or 5 glucose FeedBeads (Kühner, Basel, Switzerland), releasing glucose at constant rate,

were added when OD 600 reached 0.2. We collected 500 ul of culture medium from each flask

and centrifuged them at 16,000 g for 1.5 min. The supernatant and pellets were stored at -20°C

until further analysis.

Analysis of glucose concentration in the culture medium

The concentration of glucose was analyzed by a high-performance liquid chromatography (HPLC)

system (Agilent, Santa Ciara, CA) equipped with a 1260 refractive index detector (RID) and an

Aminex HPX-87H column (Bio-Rad, Hercules, CA). Five microliter of filtered supernatants were

injected. Analytes were separated isocratically using 5 mM sulfuric acid at a flow rate of 0.7

ml/min.

Analysis of amino acids concentration in the culture medium

The amino acids in 10 μl of filtered supernatants were analyzed using a ultra-performance liquid

chromatography (UPLC) (Waters, Milford, MA) coupled with a micrOTOF II mass spectrometry

(TOF-MS) system (Bruker, Bremen, Germany). Analytes were measured using a tunable UV

detector at 210 and 397 nm. The amino acids were separated by an Agilent Poroshell 120 EC-C18

column at 30°C. The 1 % (v/v) of formic acid in water (mobile phase A) and 1 % (v/v) of formic

acid in acetonitrile (mobile phase B) were used, respectively. The amino acids separation was

obtained at a flow rate of 0.3 ml/min with a gradient program that allowed 100% of mobile phase

A until 2.1 min followed by increasing mobile phase B to 40% for 2 min and then equilibrated at

0% of eluent B in a total analysis time of 6 min. Analysis of the amino acids was performed using

electrospray ionization (ESI) and full-scan TOF-MS spectra (50 - 650 m/z) with 500 V end plate

voltage and 4.5 kV capillary voltage. Nebulizer gas and drying gas were supplied in 1.8 bar and 8

ml/min, respectively. The dry temperature was kept at 220°C.

Quantification of cell ratio in microbial consortium

Quantitative PCR (qPCR) was carried out in a 96-well plate by using a CFX96 Real-Time

Detection System (Bio-Rad, Hercules, CA, USA). The pellets were resuspended in ultra-pure

water to make consistent concentration (OD600 = 0.4) and then, the 200 μl solution was transferred

to a 250 μl PCR tube. The solutions were incubated at 98°C for 10 min for cell disruption using a

T100 Thermal Cycler (Bio-Rad). The lysed cells were transferred to 1.5 ml of tubes and

centrifuged at 20, 000 x g for 2 min. The supernatants were analyzed by qPCR. The qPCR mixture

was composed as follows: 3 μl of 10X Xtensa® buffer, 0.3 μl of primer mix (50 uM for each),

0.15 μl of i-Taq (i-DNA Biotechnology, Singapore), 3 μl of 25 mM $MgCl_2$, 5 μl of purified cell

lysate, and 18.55 μl of ultra-pure water. The thermal cycling was programmed as follows: 95°C

for 1 min and 30 cycles of (95°C for 20 sec, 55°C for 20 sec, 68°C for 40 sec). The primers for

qPCR analysis to quantify the different *E. coli* strains were provided in **Table S3**. The qPCR

analysis was performed in triplicate for each sample.

ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by

the Korea government (MSIT) (No. 2020R1A2C2007192), and the Korea Institute of Planning

and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (iPET) funded by the

MAFRA (32136-05-1- HD050). This work was also partially supported from Nebraska Tobacco

Settlement Biomedical Research Enhancement Funds to H-SS.

References

1.   Sorboni SG, Moghaddam HS, Jafarzadeh-Esfehani R, Soleimanpour S. 2022. A

Comprehensive Review on the Role of the Gut Microbiome in Human Neurological

Disorders. Clinical Microbiology Reviews 35.

2.   Vijay A, Valdes AM. 2022. Role of the gut microbiome in chronic diseases: a narrative

review. European Journal of Clinical Nutrition 76:489-501.

3.   Sokol NW, Slessarev E, Marschmann GL, Nicolas A, Blazewicz SJ, Brodie EL, Firestone

MK, Foley MM, Hestrin R, Hungate BA, Koch BJ, Stone BW, Sullivan MB, Zablocki O,

Pett-Ridge J, Consortium LSM. 2022. Life and death in the soil microbiome: how

ecological processes influence biogeochemistry. Nature Reviews Microbiology 20:415-

430.

4.   Trivedi P, Batista BD, Bazany KE, Singh BK. 2022. Plant-microbiome interactions under

a changing world: responses, consequences and perspectives. New Phytologist 234:1951-

1959.

5.   Albright MBN, Louca S, Winkler DE, Feeser KL, Haig SJ, Whiteson KL, Emerson JB,

Dunbar J. 2022. Solutions in microbiome engineering: prioritizing barriers to organism

establishment. Isme Journal 16:331-338.

6.   Konopka A, Lindemann S, Fredrickson J. 2015. Dynamics in microbial communities:

unraveling mechanisms to identify principles. Isme Journal 9:1488-1495.

7.   Marsh JW, Ley RE. 2022. Microbiome engineering: Taming the untractable. Cell

185:416-418.

8.   Perez-Garcia O, Lear G, Singhal N. 2016. Metabolic Network Modeling of Microbial

Interactions in Natural and Engineered Environmental Systems. Frontiers in

Microbiology 7.

9.   Allen BH, Gupta N, Edirisinghe JN, Faria JP, Henry CS. 2022. Application of the

metabolic modeling pipeline in KBase to categorize reactions, predict essential genes,

and predict pathways in an isolate genome, p 291-320, Microbial Systems Biology.

Springer.

10.   Song H-S, Nelson WC, Lee J-Y, Taylor RC, Henry CS, Beliaev AS, Ramkrishna D,

Bernstein HC. 2018. Metabolic network modeling for computer-aided design of

microbial interactions.

11.   Kessell AK, McCullough HC, Auchtung JM, Bernstein HC, Song HS. 2020. Predictive

interactome modeling for precision microbiome engineering. Current Opinion in

Chemical Engineering 30:77-85.

12.   Song HS, Lindemann SR, Lee DY. 2021. Editorial: Predictive Modeling of Human

Microbiota and Their Role in Health and Disease. Frontiers in Microbiology 12.

13.   Choi YM, Lee YQ, Song HS, Lee DY. 2020. Genome scale metabolic models and

analysis for evaluating probiotic potentials. Biochemical Society Transactions 48:1309-

1321.

14.   Song HS. 2018. Design Principles of Microbial Communities: From Understanding to

Engineering. Current Genomics 19:699-700.

15.   Lee JY, Haruta S, Kato S, Bernstein HC, Lindemann SR, Lee DY, Fredrickson JK, Song

HS. 2020. Prediction of Neighbor-Dependent Microbial Interactions From Limited

Population Data. Frontiers in Microbiology 10.

16.   Song HS, Lee JY, Haruta S, Nelson WC, Lee DY, Lindemann SR, Fredrickson JK,

Bernstein HC. 2019. Minimal Interspecies Interaction Adjustment (MIIA): Inference of

Neighbor-Dependent Interactions in Microbial Communities. Frontiers in Microbiology

10.

17.   Kato S, Haruta S, Cui ZJ, Ishii M, Igarashi Y. 2008. Network relationships of bacteria in

493   a stable mixed culture. Microbial Ecology 56:403-411.

494   18.   Yamamoto K, Haruta S, Kato S, Ishii M, Igarashi Y. 2010. Determinative Factors of

495   Competitive Advantage between Aerobic Bacteria for Niches at the Air-Liquid Interface.

496   Microbes and Environments 25:317-320.

497   19.   Klitgord N, Segre D. 2010. Environments that Induce Synthetic Microbial Ecosystems.

498   Plos Computational Biology 6.

499   20.   Song H-S, Cannon WR, Beliaev AS, Konopka A. 2014. Mathematical modeling of

500   microbial community dynamics: a methodological review. Processes 2:711-752.

501   21.   Faust K, Raes J. 2012. Microbial interactions: from networks to models. Nature Reviews

502   Microbiology 10:538-550.

503   22.   Friedman J, Alm EJ. 2012. Inferring Correlation Networks from Genomic Survey Data.

504   Plos Computational Biology 8.

505   23.   Fang HY, Huang CC, Zhao HY, Deng MH. 2015. CCLasso: correlation inference for

506   compositional data through Lasso. Bioinformatics 31:3172-3180.

507   24.   Ban YG, An LL, Jiang HM. 2015. Investigating microbial co-occurrence patterns based

508   on metagenomic compositional data. Bioinformatics 31:3322-3329.

509   25.   Joseph TA, Shenhav L, Xavier JB, Halperin E, Peer I. 2020. Compositional Lotka-

510   Volterra describes microbial dynamics in the simplex. Plos Computational Biology 16.

511   26.   Fisher CK, Mehta P. 2014. Identifying Keystone Species in the Human Gut Microbiome

512   from Metagenomic Timeseries Using Sparse Linear Regression. Plos One 9.

513   27.   Stein RR, Bucci V, Toussaint NC, Buffie CG, Ratsch G, Pamer EG, Sander C, Xavier JB.

514   2013. Ecological Modeling from Time-Series Inference: Insight into Dynamics and

515   Stability of Intestinal Microbiota. Plos Computational Biology 9.

516   28.   Li CH, Chng KR, Kwah JS, Av-Shalom TV, Tucker-Kellogg L, Nagarajan N. 2019. An

517   expectation-maximization algorithm enables accurate ecological modeling using

518   longitudinal microbiome sequencing data. Microbiome 7.

519   29.   Saeedian M, Pigani E, Maritan A, Suweis S, Azaele S. 2022. Effect of delay on the

520   emergent stability patterns in generalized Lotka-Volterra ecological dynamics.

521   Philosophical Transactions of the Royal Society a-Mathematical Physical and

522   Engineering Sciences 380.

523   30.   Mee MT, Collins JJ, Church GM, Wang HH. 2014. Syntrophic exchange in synthetic

524   microbial communities. Proceedings of the National Academy of Sciences of the United

525   States of America 111:E2149-E2156.

526   31.   Kerner A, Park J, Williams A, Lin XXNN. 2012. A Programmable Escherichia coli

527   Consortium via Tunable Symbiosis. Plos One 7.

528   32.   Momeni B, Xie L, Shou WY. 2017. Lotka-Volterra pairwise modeling fails to capture

529   diverse pairwise microbial interactions. Elife 6.

530   33.   McClure R, Naylor D, Farris Y, Davison M, Fansler SJ, Hofmockel KS, Jansson JK.

531   2020. Development and Analysis of a Stable, Reduced Complexity Model Soil

532   Microbiome. Frontiers in Microbiology 11.

533   34.   Kylilis N, Tuza ZA, Stan GB, Polizzi KM. 2018. Tools for engineering coordinated

534   system behaviour in synthetic microbial consortia. Nature Communications 9.

 Table captions

Table 1. Model equations with kinetic parameters and stoichiometric coefficients determined

through the model fit to experimental data collected under various limiting conditions. R_i is the

stoichiometric growth reaction for X_i, and Y_{S^-/X_i}, $Y_{S_i^+/X_i}$, and $Y_{S_j^+/X_i}$ denote the stoichiometric

coefficients for S^-, S_i^+, and S_j^+ associated with the growth of X_i. s^-, s_i^+, and s_j^+ respectively

denote concentrations of S^-, S_i^+, and S_j^+, x_i is the population density of X_i, μ_i is the specific

growth rate of X_i, $k_{d,i}$ is the death rate of X_i, and q_{s^-} is the substrate releasing rate from FBs in

glucose-limited semi-batch cultures (i.e., $q_{s^-} = 0$ in a batch mode). μ_i^{max} is the maximal growth

rate, and K_i^- and K_i^+ are half-saturation constants associated with the consumption of S^- and S_i^+.

 Figure captions

Fig. 1. Conceptual illustration of context-dependent microbial interactions in a binary

consortium dictated by the environmental contexts. Two species X_1 and X_2 compete for the

substrate S^- but cooperate by cross-feeding metabolite S_1^+ and S_2^+ (center panel). This mixed

relationship between X_1 and X_2 diverge into six different types of interactions by excessive

addition of specific substrates S_1^+, S_2^+, and/or S^-. Symbols next to the arrows denote the

substrate(s) excessively added to the environment. (**c**) Model microbial consortium using

tyrosine or tryptophan auxotrophic E. coli strains. Glucose is sole carbon source for both *E. coli*

strains. GLC, Tyr and Trp denote glucose, tyrosine, and tryptophan, respectively.

Fig. 2. Experimental data and model simulations for the growth of two *E. coli* mutant strains in

axenic and binary culture conditions: **A** and **B** are cultures of Trp auxotrophic and Tyr

auxotrophic *E. coli* mutants (X_1 and X_2), respectively; **C** and **D** are co-cultures with two

auxotrophs in batch and semi-batch cultures. Detailed culture conditions for the twelve panels

are provided in **Table S2**. Circles and lines denote the experimentally measured values and

simulation results, respectively. Black line denotes simulation results for glucose concentration

(S^-), and the lines in blue, red, and purple are simulated population densities of X_1, X_2, and $X_1 +$

X_2. **Figs. A, B,** and **C** show data fitting to determine model parameters, while the results in **Fig.**

D validate model predictions.

Fig. 3. Inference of dynamic variations of interaction parameters ($\gamma_{i,j}, a_{i,j}, a_{i,j}^+$, and $a_{i,j}^-$) for the

two *E. coli* mutants (X_1 and X_2) co-growing in three batch cultures: **A.** initial substrate

concentrations of 2 g/l glucose and 0.4 mg/l Trp & Tyr; **B.** initial substrate concentrations of 0.5

g/l glucose and 0.4 mg/l Trp & Tyr; and **C.** initial substrate concentrations of 0.5 g/l glucose and

4 mg/l Trp & Tyr. Black line denotes simulated glucose concentration (S^-); the lines in blue and

red indicate the variables and parameters associated with X_1 and X_2, respectively.

Fig. 4. Inference of dynamic variations of interaction parameters ($\gamma_{i,j}, a_{i,j}, a_{i,j}^+$, and $a_{i,j}^-$) for the

two *E. coli* mutants (X_1 and X_2) co-growing in three semi-batch cultures: **A.** initial substrate

concentrations of 2 g/l glucose and 0.4 mg/l Trp & Tyr, and 3 FBs added at 7.5 hours; **B.** initial

substrate concentrations of 0.5 g/l glucose and 0.4 mg/l Trp & Tyr, and 3 FBs added at 7.5 hours;

and **C.** initial substrate concentrations of 0.5 g/l glucose and 4 mg/l Trp & Tyr, and 3 FBs added

at 7.5 hours. Black line denotes simulated glucose concentration (S^-); the lines in blue and red

indicate the variables and parameters associated with X_1 and X_2, respectively.

Table 1
586

Stoichiometric equation (R_i) for growth of X_i:

$$R_i : Y_{S^-/X_i}S^- + Y_{S_i^+/X_i}S_i^+ \rightarrow X_i + Y_{S_j^+/X_i}S_j^+, \quad (i,j) = (1,2) \text{ or } (2,1) \tag{T1}$$

Dynamic mass balances:

$$\frac{dx_i}{dt} = \mu_i x_i - k_{d,i} x_i, \quad i = 1,2 \tag{T2}$$

$$\frac{ds^-}{dt} = -Y_{S^-/X_1}\mu_1 x_1 - Y_{S^-/X_2}\mu_2 x_2 (+q_{S^-}) \tag{T3}$$

$$\frac{ds_i^+}{dt} = -Y_{S_i^+/X_i}\mu_i x_i + Y_{S_i^+/X_j}\mu_j x_j, \quad (i,j) = (1,2) \text{ or } (2,1) \tag{T4}$$

Double Monod kinetics:

$$\mu_i = \mu_i^{max} \frac{s_i^+}{(K_i^+ + s_i^+)} \frac{s^-}{(K_i^- + s^-)}, \quad i = 1,2 \tag{T5}$$

Kinetic parameters and stoichiometric coefficients determined through data fit:

Parameter	Value	Parameter	Value
μ_1^{max} [1/h]	0.297	Y_{S^-/X_1} [g/OD]	1.26
μ_2^{max} [1/h]	0.173	Y_{S^-/X_2} [g/OD]	1.47
K_1^- [g/l]	3.95 x 10^{-4}	$Y_{S_1^+/X_1}$ [mg/OD]	8.45 x 10^{-2}
K_2^- [g/l]	1.16 x 10^{-3}	$Y_{S_2^+/X_1}$ [mg/OD]	9.10 x 10^{-3}
K_1^+ [mg/l]	0.165	$Y_{S_1^+/X_2}$ [mg/OD]	0.107
K_2^+ [mg/l]	5.85 x 10^{-3}	$Y_{S_2^+/X_2}$ [mg/OD]	0.133
$k_{d,1}$ [1/h]	3.06 x 10^{-3}	q_{S^-} (for 3 FBs) [g/l/h]	4.80 X 10^{-2}
$k_{d,2}$ [1/h]	9.76 x 10^{-4}	q_{S^-} (for 5 FBs) [g/l/h]	8.01 x 10^{-2}

589   Figure 1

594   Figure 2

598   Figure 3

602   Figure 4

606   **SUPPLEMENTARY TABLES AND FIGURES**

608   **Table S1.** Initial culture conditions for experimental data displayed in Fig. 2.

Exp #	Corresponding figures	Initial inoculum size (OD)	Glucose (mg/l)	Tryptophan (mg/l)	Tyrosine (mg/l)	# of FBs added	Time FBs added (hr)
1	Fig. 2A (top panel)	ΔTrp only: 0.01	5.838	10	-	-	-
2	Fig. 2A (middle panel)	ΔTrp only: 0.01	5.870	20	-	-	-
3	Fig. 2A (bottom panel)	ΔTrp only: 0.01	5.936	40	-	-	-
4	Fig. 2B (top panel)	ΔTyr only: 0.01	5.920	-	10	-	-
5	Fig. 2B (middle panel)	ΔTyr only: 0.01	5.909	-	20	-	-
6	Fig. 2B (bottom panel)	ΔTyr only: 0.01	5.863	-	40	-	-
7	Fig. 2C (top panel)	ΔTrp: 0.01 ΔTyr: 0.01	3.851	0.4	0.4	-	-
8	Fig. 2C (middle panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.626	0.4	0.4	3	7.5
9	Fig. 2C (bottom panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.620	0.4	0.4	5	7.5
10	Fig. 2D (top panel)	ΔTrp: 0.01 ΔTyr: 0.01	2.878	0.4	0.4	-	-
11	Fig. 2D (middle panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.375	0.4	0.4	3	10
12	Fig. 2D (bottom panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.377	200	200	3	10

612   **Table S2.** Sequence of the primers used in the qPCR analysis.

Primer	Sequence
Trp F	GCCGATGCCTGCTTATTA
Trp R	GCTCCTGTTCCTCTTCAT
Tyr F	CATTATGTCGTCAGAGCG
Tyr R	CCTTGCGGAAACTGTCAA

614
615

619   **Figure S1.** Zoom-in views of in interaction parameters, $a_{i,j}, a_{i,j}^{+}$, and $a_{i,j}^{-}$ in **Figs. 3A, 3B, and**

620   **3C,** respectively. The color scheme is the same as in **Fig. 3**.

Figure S2. Zoom-in views of in interaction parameters, $a_{i,j}$, $a_{i,j}^+$, and $a_{i,j}^-$ in **Figs. 4A, 4B, and 4C,** respectively. The color scheme is the same as in **Fig. 4**.

Figure S3. The predicted impacts of the numbers and timings of added FBs on the substrate
profile (S^-), the population dynamics (X_1 and X_2), and the interaction parameter ($\gamma_{i,j}$): **A.** initial
substrate concentrations of 2 g/l glucose and 0.4 mg/l Trp & Tyr with 3 FBs (left), 6 FBs
(middle) and 10 FBs (right) added at 7.5 hours; **B.** initial substrate concentrations of 0.5 g/l
glucose and 0.4 mg/l Trp & Tyr with 3 FBs (left), 6 FBs (middle) and 10 FBs (right) added at 7.5
hours; and **C.** initial substrate concentrations of 0.5 g/l glucose and 4 mg/l Trp & Tyr, with 6 FBs
added at 7.5 hours (left), 10 hours (middle) and 15 hours (right). The color scheme is the same as
Fig. 4.