Kinetics-based Inference of Environment-Dependent Microbial Interactions and Their Dynamic Variation

Hyun-Seob Song[1,2†*], Na-Rae Lee[3†], Aimee K. Kessell[1], Hugh C. McCullough[1], Seo-Young Park[4], Kang Zhou[5], and Dong-Yup Lee[4*]

[1]Department of Biological Systems Engineering, University of Nebraska-Lincoln, Lincoln, NE 68583

[2]Department of Food Science and Technology, Nebraska Food for Health Center, University of Nebraska-Lincoln, Lincoln, NE 68588

[3]Research Institute for Bioactive-Metabolome Network, Konkuk University, Seoul 05029, Republic of Korea

[4]School of Chemical Engineering, Sungkyunkwan University, Suwon-si, Gyeonggi-do 16419, Republic of Korea

[5]Department of Chemical and Biomolecular Engineering, National University of Singapore, 4 Engineering Drive 4, Singapore 117585, Singapore

[†]Both authors contributed equally.

*Correspondence: hsong5@unl.edu (H.-S.S.); dongyuplee@skku.edu (D.-Y.L.)

ABSTRACT: Microbial communities in nature are dynamically evolving as member species change their interactions subject to environmental variations. Accounting for such context-dependent dynamic variations in interspecies interactions is critical for predictive ecological modeling. In the absence of generalizable theoretical foundations, we lack a fundamental understanding of how microbial interactions are driven by environmental factors, significantly limiting our capability to predict and engineer community dynamics and function. To address this issue, we propose a novel theoretical framework that allows us to represent interspecies interactions as an explicit function of environmental variables (such as substrate concentrations) by combining growth kinetics and a generalized Lotka-Volterra model. A synergistic integration of these two complementary models leads to the prediction of alterations in interspecies interactions as the outcome of dynamic balances between positive and negative influences of microbial species in mixed relationships. This unique capability of our approach was experimentally demonstrated using a synthetic consortium of two *Escherichia coli* mutants that are metabolically dependent (due to an inability to synthesize essential amino acids), but competitively growing on a shared substrate. The analysis of the *E. coli* binary consortium using our model not only showed how interactions between the two amino acid auxotrophic mutants are controlled by the dynamic shifts in limiting substrates, but also enabled quantifying previously uncharacterizable complex aspects of microbial interactions such as asymmetry in interactions. Our approach can be extended to other ecological systems to model their environment-dependent interspecies interactions from growth kinetics.

IMPORTANCE: Modeling of environment-controlled interspecies interactions through separate identification of positive and negative influences of microbes in mixed relationships is a new

capability that can significantly improve our ability to understand, predict, and engineer complex

dynamics of microbial communities. Moreover, robust prediction of microbial interactions as a

function of environmental variables can serve as valuable benchmark data to validate modeling

and network inference tools in microbial ecology, the development of which has often been

impeded due to the lack of ground truth information on interactions. While demonstrated against

microbial data, the theory developed in this work is readily applicable to general community

ecology to predict interactions among microorganisms such as plants and animals, as well as

microorganisms.

KEYWORDS: Microbial communities; competition; cooperation; context dependence; kinetic

models; Lotka-Volterra models

INTRODUCTION

Microbial communities play pivotal roles in maintaining human and animal health, plant productivity, and ecosystem services (1-4). Increasing efforts are being dedicated towards maximizing their beneficial roles in natural systems or creating new industrial applications (5). However, control and design of microbial community dynamics and function is a challenging task, primarily due to higher-order or emergent properties that are not observable from individual species in isolation but arise through nonlinear interspecies interactions (6, 7). Therefore, rational design of microbial communities or consortia requires a fundamental knowledge of microbial interactions as a mechanistic linkage between the environment and the community compositions and function, necessitating the employment of predictive mathematical models as indispensable tools (8-14).

Development of accurate models of microbial communities that are commonly subject to environmental variations is truly complicated by the following intrinsic ecological aspects. First, microorganisms in a community build *dynamic* interactions that cannot effectively be represented by a rigid network with fixed structure (15, 16). Rather, microbial communities keep reorganizing interaction networks in response to biotic or abiotic perturbations or through adaptation to long-lasting environmental changes. Second, microorganisms often build *mixed* relationships by exerting *both* promotive and inhibitive impacts on the growth of their partners/neighbors (17, 18). Individual identification of these simultaneously acting positive and negative interactions is critical because community dynamics is mainly driven by the balances between all counteracting impacts among member species (19). The lack of capability to account for these key properties of microbial interactions limits our ability to predict and engineer microbial community dynamics and functions.

Despite rapid progress in microbiome science, we still do not know how to identify

environment-controlled dynamic variation in interspecies interactions addressed above. Three

major branches of microbial interaction modeling include (20, 21): (i) network inference, (ii)

metabolic network modeling, and (iii) kinetic modeling. Network inference is widely used for

modeling microbial interactions to identify interaction networks based on correlative relationships

among microbial populations (22-25), parameter identification through regression (26-28), or a

prescribed set of rules or hypotheses (21). The resulting networks represent interspecies

interactions as single constant metrics, therefore being unable to describe dynamic variations in

interactions nor identify the balances among counteracting individual impacts in mixed

relationships. As an exception, the approach termed MIIA (Minimal Interspecies Interaction

Adjustment) (15, 16) uniquely enables predicting context-dependent interactions due to the

changes in memberships, which however has not been extended to address the environmental

impacts. In contrast with such data-driven network inference methods, metabolic network and

kinetic modeling can account for both positive and negative interactions based on cross-feeding of

small molecules (essential for growth) or competition for shared substrates/nutrients among

species; in theory, kinetic models can additionally simulate their dynamic variations. While more

mechanistic than network inference, these methods cannot quantify the magnitude or even the sign

of net interactions.

In this work, we fill these gaps by proposing a novel theoretical framework that enables a

quantifiable, mechanistic representation of the dynamic linkage between microbial interactions

and the environment. For this purpose, we synergistically integrate two complementary modeling

frameworks to overcome their own limitations: a generalized Lotka-Volterra (gLV) model (29)

and population growth kinetics. Like other network inference approaches, a typical gLV model

with a focus on pairwise interactions is constructed based on an implicit assumption of constant

interactions. We relax this assumption by representing interaction coefficients in the gLV model

as a function of environmental variables (i.e., concentrations of cross-fed metabolites and shared

substrates) described in microbial growth kinetics, which is termed here kinetics-based inference

of dynamic variation in microbial interactions (KIDI). The resulting functional representation of

interactions by KIDI enables not only quantifying their dynamic variation as environmental

conditions change, but also individually identifying negative and positive influences among

species in mixed relationships. The prediction of KIDI was demonstrated accurate through a

coordinated design of experiments using a binary consortium composed of tyrosine and tryptophan

auxotrophic mutants of *Escherichia coli* (30) so that they both compete and/or cooperate

depending on environmental conditions.

RESULTS

Formulation of a conceptual model for understanding environment-dependent interactions

For illustration of the concept of KIDI, we consider a hypothetical consortium composed of two

members where species 1 (X_1) and species 2 (X_2) cooperate by cross-feeding S_1^+ and S_2^+ each other

but compete for the shared metabolite S^- (the center circle in **Fig. 1**). Growth kinetics for the i^{th}

species (X_i) (that requires two substrates S_i^+ and S^- for growth) can be represented, e.g., using a

double Michaelis-Menten equation as follows:

$$\mu_i = \mu_i^{\max} \frac{s_i^+}{\left(K_i^+ + s_i^+\right)} \frac{s^-}{\left(K_i^- + s^-\right)}, \quad i = 1, 2 \tag{1}$$

where μ_i [1/h] is the specific growth rate of X_i, μ_i^{max} is the maximal specific growth rate, s_i^+ and

s^- [g/l] are the concentrations of S_i^+ and S^-, and K_i^+ and K_i^- [g/l] are half-saturation constants

associated with the consumption of S_i^+ and S^-, respectively. As inferable from growth kinetics in

Eq. (1), the mixed relationship (i.e., competition and cooperation) between X_1 and X_2 when both

substrates are limiting can turn into diverse forms of interactions as environmental conditions

change. When S^- is present in excess (therefore, no competition is necessary) but S_1^+ and S_2^+ are

limiting, for example, their relationship is predominantly cooperative (where $\mu_i \approx \frac{\mu_i^{max} s_i^+}{K_i^+ + s_i^+}$). As the

opposite case, if both S_1^+ and S_2^+ are excessive in the environment (so no partners are needed to

acquire them) while S^- is limiting, their relationship is governed by competition (where $\mu_i \approx$

$\frac{\mu_i^{max} s^-}{K_i^- + s^-}$). Likewise, one can assume many other different scenarios where their relationships turn

into competition, cooperation, amensalism, commensalism and even neutrality, as illustrated in

Fig. 1.

Representation of interaction parameters as a function of environmental variables

To model such environment-dependent microbial relationships, we derived a general form of

interaction coefficients as a function of environmental variables by integrating growth kinetics and

a gLV model. As described in detail in **Materials and Methods**, our formula (KIDI) represents

interaction coefficients of species in the mixed relationship as a sum of positive and negative parts,

i.e.,

121
$$a_{i,j}\left(s_i^+, s^-\right) = a_{i,j}^+\left(s_i^+, s^-\right) + a_{i,j}^-\left(s_i^+, s^-\right), \quad (i, j) = (1, 2) \text{ or } (2,1) \tag{2}$$

122 where $a_{i,j}^+$ and $a_{i,j}^-$ denote the positive and negative influence of X_j on the growth rate of X_i, which

123 are defined as follows:

124
$$a_{i,j}^+\left(s_i^+, s^-\right) \equiv \frac{\partial}{\partial s_i^+}\left[\mu_i\left(s_i^+, s^-\right)\right] \cdot \frac{\partial s_i^+}{\partial x_j} \tag{3}$$

125
$$a_{i,j}^-\left(s_i^+,s^-\right) \equiv \frac{\partial}{\partial s^-}\left[\mu_i\left(s_i^+,s^-\right)\right]\cdot\frac{\partial s^-}{\partial x_j} \tag{4}$$

The positive influence of X_j on the growth rate of X_i (i.e., $a_{i,j}^+$) is represented by the two

subsequent terms on the right-hand side of Eq. (3): (i) the impact of the change in the population

size of X_j on the concentration of the cross-fed substrate S_i^+ (as denoted by $\partial s_i^+/\partial x_j$) and (2) the

subsequent impact of the change in S_i^+ on the growth rate of the i^{th} species (i.e., μ_i) (as denoted

by $\partial[\mu_i(s^-,s_i^+)]/\partial s_i^+$). The negative impact of X_j on the growth rate of X_i ($a_{i,j}^-$) in Eq. (4) can be

interpreted in a similar fashion.

The derivative terms on the right-hand side of Eqs. (3) and (4) are fully identifiable from

reaction stoichiometry and kinetics. In the case of using a double Monod kinetics, for example,

incorporation of Eq. (1) into Eqs. (3) and (4) yields $a_{i,j}^+$ and $a_{i,j}^-$ as follows:

135
$$a_{i,j}^+\left(s_i^+,s^-\right) = \left[\mu_i^{\max}\frac{K_i^+}{\left(K_i^+ + s_i^+\right)^2}\frac{s^-}{\left(K_i^- + s^-\right)}\right]\cdot Y_{S_i^+/X_j} \tag{5}$$

136
$$a_{i,j}^-\left(s_i^+,s^-\right) = \left[\mu_i^{\max}\frac{s_i^+}{\left(K_i^+ + s_i^+\right)}\frac{K_i^-}{\left(K_i^- + s^-\right)^2}\right]\cdot\left(-Y_{S^-/X_j}\right) \tag{6}$$

137 where $Y_{S_i^+/X_j}$ and Y_{S^-/X_j} denote the stoichiometric relationships between the changes in substrate

138 and biomass concentrations associated with X_j, i.e., $Y_{S_i^+/X_j} = |\Delta s_i^+/\Delta x_j|$ and $Y_{S^-/X_j} = |\Delta s^-/\Delta x_j|$

139 (see **Materials and Methods**).

140 To identify net interactions between two species with mixed relationships, we further

141 defined a normalized interaction parameter $\gamma_{i,j}$ as follows:

142
$$\gamma_{i,j} \equiv \frac{a_{i,j}^+ + a_{i,j}^-}{a_{i,j}^+ - a_{i,j}^-}, \quad (i,j) = (1,2) \text{ or } (2,1) \tag{7}$$

Once $a_{i,j}^+$ and $a_{i,j}^-$ are identified from Eqs. (3) and (4), the parameter $\gamma_{i,j}$ is readily calculatable by

Eq. (7). The parameter $\gamma_{i,j}$ ranges from -1 to 1 to represent positive influences of species j on

species i when greater than 0 and negative impacts when less than 0, respectively, consequently

allowing us to conveniently quantify the relative dominance of inhibition vs. promotion in mixed

interactions. The parameter $\gamma_{i,j}$ complements $a_{i,j}$, rather than replaces it, in that the magnitude of

interactions cannot be determined by $\gamma_{i,j}$, but by the original interaction parameter, $a_{i,j}$. In this

regard, $\gamma_{i,j}$ provides an additional complementary explanation of dynamic changes in interspecies

interactions. Therefore, all these parameters, including $a_{i,j}$ defined in Eqs. (2) to (4) and $\gamma_{i,j}$,

sufficiently characterize the dynamic variation of interactions between X_1 and X_2 based on co-

culture growth data as demonstrated in the following sections.

Identification of kinetics and stoichiometry via data fit

For experimental demonstration of the mathematical formulation derived in the previous section,

we constructed a synthetic consortium composed of two *E. coli* auxotrophic mutants that can

cooperatively cross-feed small molecules, while competitively growing on a common substrate

(31). Among 14 amino acid auxotrophic mutants, we chose tryptophan and tyrosine auxotrophic

mutants by considering the bioenergetic cost for the synthesis of amino acids based on a previous

study in the literature (30). This consortium is considered an ideal, simplest model system for

studying environment-dependent dynamic variations in microbial interactions. Due to its exact

correspondence to the hypothetical consortium in **Fig. 1**, we denote two *E. coli* mutant strains

ΔtrpC and ΔtyrA by X_1 and X_2; glucose, tryptophan, and tyrosine by S^-, S_1^+, and S_2^+, respectively.

Using these two strains, we performed growth experiments under diverse culture

conditions: two individual batch experiments using X_1 (**Fig. 2A**) and X_2 (**Fig. 2B**), respectively,

and two sets of co-culture experiments (**Figs. 2C** and **2D**). The top panels in **Figs. 2C** and **2D**

denote co-growth experiments in batch cultures, while the middle and bottom panels denote semi-

batch modes, where we added glucose feedbeads (FBs) at time 7.5 and 10 hours, respectively, to

induce dramatic changes in interspecies interactions during co-growth. Other differences in co-

culture conditions in **Figs. 2C** and **2D** include initial concentrations of S^-, S_1^+, and S_2^+, and the

number of added FBs (see **Supp Datasheet**). We measured optical density (OD) at an absorbance

of 600 nm as a metric of cell density and determined the OD for each strain in the consortium

through additional analysis using quantitative PCR (qPCR). The OD profiles in **Fig. 2D** denote the

combined population change of both strains, i.e., $X_1 + X_2$.

Based on the four datasets in **Fig. 2**, we constructed a dynamic co-growth model of X_1 and

X_2 to determine associated kinetics and stoichiometry, key information required for quantifying

interspecies interactions parameters ($a_{i,j}$ and $\gamma_{i,j}$) in Eqs. (2) to (7). The dynamic co-growth model

is composed of five mass balance equations for X_1, X_2, S^-, S_1^+, and S_2^+. We determined

stoichiometric and kinetic parameters using three subsets of data in **Figs. 2A – 2C** and validated

the model against the remaining one (in **Fig. 2D**) that was not used for model identification. The

robust consistency between simulated and measured data in **Fig. 2D**, as well as those in **Figs. 2A**

– 2C, indicates the acceptability of using the identified model parameters in inferring interaction

coefficients. We provide the full list of model equations with parameter values in **Table 1**; the

culture conditions in **Table S1**.

While the overall performance of the kinetic model was satisfactory, we found that our

experimental setups were not ideal for accurately determining all model parameters. For example,

Fig. 2A and **2B** did not effectively show the dependence of the growth of ΔTrp and ΔTry strains

on tryptophan and tyrosine because the range of initial concentrations of amino acids (from 10 to

40 mg/l) was too high compared to the half-saturation constants K_1^+ and K_2^+ (which were

determined to be 0.0033 mg/l for tryptophan and 0.00039 mg/l for tyrosine, respectively). This

mismatch is partly due to the difficulty in identifying the magnitudes of half-saturation constants

in advance before being determined through data fit.

Variation in microbial interactions driven by the switch in limiting substrates in batch

cultures

Based on the stochiometric and kinetic parameters determined through data fit in **Table 1**, we were

able to determine microbial interactions and their variations as a function of environmental

conditions using KIDI. We first analyzed various co-culture scenarios in batch modes (**Fig. 3**). In

Fig. 3A, we considered the growth of X_1 and X_2 on relatively high and low initial concentrations

of S^- (2 g/l) and S_1^+ and S_2^+ (i.e., 1 mg/l for both) as a reference condition. In the present setting,

the relationship between X_1 and X_2 is expected to be mostly cooperative (because S^- is excessive

in the beginning) and become competitive as the level of S^- decreases. While this overall trend

was captured well by our model, the simulation results showed more intricate dynamics than our

expectations. As depicted in the top panel of **Fig. 3A**, the concentrations of S_1^+ and S_2^+ show a

decreasing and increasing trend over time, respectively. This means that X_2 does not supply

sufficient S_1^+ for X_1, whereas X_1 provides an excess of S_2^+ for X_2. As a result, the growth pattern

of X_1 exhibits two distinct exponential phases. Notably, the second phase commences with a

slower growth rate when S_1^+ becomes depleted in the medium (the second panel from the top).

This dynamic accounts for the observed slowdown in the consumption rate of S^- (in the first panel).

Overall, these results indicate that the growth of X_1 has a greater dependency on X_2 than X_2 has

on X_1, particularly when S_1^+ becomes depleted. This aspect is correctly captured by the higher

values of $\gamma_{1,2}$ than $\gamma_{2,1}$ as shown in the third panel from the top. Actual values of interaction

coefficients can be seen from $a_{i,j}$, $a_{i,j}^{+}$, and $a_{i,j}^{-}$ (three bottom panels of **Fig. 3A**), which also

showed that $a_{1,2}^{+} > a_{2,1}^{+}$ (and consequently $a_{1,2} > a_{2,1}$) in the second growth phase of X_2.

For comparison, we analyzed two additional conditions. (1) with a lower initial

concentration of S^{-} (i.e., 0.5 g/l) (**Fig. 3B**) and (2) with lower and higher initial concentrations of

S^{-} (0.5 g/l) and S_1^{+} and S_2^{+} (i.e., 4 mg/l for both) (**Fig. 3C**), respectively. Unlike the first case in

Fig. 3A, the growth profile of X_2 does not show a biphasic growth because S_1^{+} and S^{-} are depleted

almost at the same time. In the case of lowering the initial concentration of S^{-} (**Fig. 3B**), KIDI

showed that the level of initial competition increases (due to the limited availability of S^{-}) as

indicated by relatively lower values of $\gamma_{1,2}$ and $\gamma_{2,1}$ compared to the case of **Fig. 3A**. Notably, $\gamma_{2,1}$

showed negative value throughout the co-growth (indicating the dominance of negative influence

of X_1 on the growth of X_2). In the case of increasing the initial concentrations of S_1^{+} and S_2^{+} in

addition to lowering S^{-} (**Fig. 3C**), the relationship between the two strains became even more

negative (i.e., both $\gamma_{1,2}$ and $\gamma_{2,1}$ are negative), which was also an expected outcome because

metabolic dependence between X_1 and X_2 will accordingly reduce when they can acquire what

they need from the environment, rather than from partners.

In all of these cases, the relations between the two *E. coli* strains were shown asymmetric,

i.e., $\gamma_{1,2} \neq \gamma_{2,1}$, $a_{1,2} \neq a_{2,1}$, $a_{1,2}^{-} \neq a_{2,1}^{-}$, and $a_{1,2}^{+} \neq a_{2,1}^{+}$. Asymmetric interactions in terms of

$a_{i,j}$'s can also be seen over smaller time windows in **Fig. S1**. KIDI predicted $a_{1,2} > a_{2,1}$ for the

first two cases (**Fig. S1A and S1B**) but $a_{1,2} < a_{2,1}$ for the third case (**Fig. S1C**). In the reference

condition where glucose is excessive (so $a_{i,j}^{-}$'s are relatively negligible), it is mostly due to $a_{1,2}^{+} >$

$a_{2,1}^{+}$ (i.e., X_1 has a higher comparative advantage in exchanging amino acids with X_2 than the other

way around) that leads $\gamma_{1,2} > \gamma_{2,1}$ (as well as $a_{1,2} > a_{2,1}$) (**Fig. 3A** and **Fig. S1A**). A similar trend

(i.e., $\gamma_{1,2} > \gamma_{2,1}$ and $a_{1,2} > a_{2,1}$) is observed in the second case where all substrates (glucose and

amino acids) are limitedly available in the environment and therefore both $a_{i,j}^+$'s and $a_{i,j}^-$'s make

comparable contributions to the net interaction coefficients (i.e., $a_{i,j}$'s) (**Fig. 3B** and **Fig. S1B**). In

contrast with the first two cases, the net interaction coefficients are shown to be $a_{1,2} < a_{2,1}$ for the

third case where the glucose level is low while amino acids are abundant (so $a_{i,j}^+$'s are negligible)

because the magnitudes of $a_{i,j}^-$'s are greater than $a_{i,j}^+$'s. Interestingly, KIDI predicted $\gamma_{1,2} > \gamma_{2,1}$

(**Fig. 3C**) despite $a_{1,2} < a_{2,1}$ (**Fig. S1C**), which can happen because the implications of $\gamma_{i,j}$ and

$a_{i,j}$ are not necessarily identical. The former denotes the relative dominance between promotion

vs. inhibition in the relationship of species i with species j, while the latter represents the net effect

of species j on the growth of species i.

Dynamic response of microbial interactions to environmental perturbations during growth

We extend our analysis to semi-batch cultures that are perturbed by the addition of glucose FBs

during growth and therefore are expected to show more dramatic changes in interspecies

interactions and community dynamics. In contrast with the batch cultures considered in the

previous section where no further growth is possible after the depletion of the initially added S^-,

the two strains continue to grow in semi-batch cultures due to slow but continual provision of S^-

from the added FBs. Despite a general expectation that the competition level between the two

strains will be mitigated at least at the moment of FB addition, it is uncertain: (1) to what degree

this will occur under different environmental conditions, and (2) how governing microbial

interactions will shift (between competition and cooperation), particularly in a later phase when

the growth of the two strains is be limited by both S^- and S_i^+. To answer these questions, we

applied KIDI to the following three cases. For simplicity, we set the initial conditions to be the

same as before.

First, we considered the initial concentrations of 2 g/l for S^- and 1 mg/l for S_1^+ and S_2^+ and

added 3 FBs of S^- at around 7.5 hours (**Fig. 4A**). Due to the relatively high concentration of S^-,

the impact of adding 3 FBs of S^- on interactions was minimal. The profiles of interaction

parameters (i.e., $\gamma_{i,j}, a_{i,j}, a_{i,j}^+$, and $a_{i,j}^-$ in the four bottom panels in **Fig. 3A**) as well as the growth

curves of X_1 and X_2 showed no qualitative differences from the batch case (**Fig. 3A**), while the

concentration profile of S^- showed an appreciable increase at the time of addition of 3 FBs (the

top panel in **Fig. 3A**).

By contrast, when the initial concentration of S^- was low (i.e., 0.5 g/l) (**Fig. 4B**), KIDI

identified the greater impact of adding FBs on both glucose concentration and microbial

interactions, as indicated by sudden increases in S^-, $\gamma_{1,2}$ and $\gamma_{2,1}$. Interestingly, the value of $\gamma_{1,2}$

shifted to negative (from positive) when the medium was depleted of S^-, but reverted to positive

upon the depletion of S_1^+. The latter suggest a substantial rise in the dependence of X_1 on X_2 in the

absence of S_1^+ from the medium. A similar pattern was also noted for $a_{1,2}$, $a_{1,2}^+$, and $a_{1,2}^-$.

Additionally increasing the initial concentrations of S_1^+ and S_2^+ (as shown in **Fig. 4C**), thereby

intensifying the level of competition, resulted in overall patterns similar to the previous case.

However, both $\gamma_{1,2}$ and $\gamma_{2,1}$ consistently showed negative values, attributed to the heightened

competition. Unlike the previous scenario, there was no increase in interaction parameters, as S_1^+

remained available in the medium throughout the time window up to 30 hours. Asymmetry in

interaction parameters over a shorter time frame is observable in the detailed views provided in

Fig. S2.

The simulations presented in this section demonstrate that the interactions between the two

strains are highly nonlinear, influenced by the availability of S^-, S_1^+ and S_2^+ in the environment,

as well as the growth characteristics of X_1 and X_2. As a general trend, interspecies interactions

were dominated by competition when S^- levels were low, but shifted towards cooperation in the

presence of additional limitations of S_1^+ (potentially S_2^+ as well). Similar patterns were observed

in other scenarios involving perturbations, where both the number of added FBs and the timing of

their addition varied, as shown in **Fig. S3**.

DISCUSSION

In this study, we proposed a novel computational method (KIDI) that enables quantitatively

identifying environment-dependent interspecies interactions in microbial communities. By

integrating growth kinetics into a gLV model, we derived an analytical form of interaction

coefficients as a function of environmental variables (i.e., concentrations of chemical substrates

that affect interactions), the results of which were subsequently validated through a coordinated

design of co-culture experiments.

Our theoretical development significantly extends the current scope of microbial ecological

modeling by completely relaxing the typical assumption of constant interactions among species.

The gLV model, for example, has been widely used as a basic ecological modeling template for

simulation of population dynamics and inference of interspecies interactions in microbial

communities (26-28). Due to the constant interaction assumption, however, the application of the

gLV model is often confined to a narrow range of conditions where interspecies interactions are

expected to remain largely constant. KIDI addresses this limitation by representing interaction

coefficients as an explicit function of limiting substrates. As an exception, a previous study by

Momeni et al. (32) showed that pairwise interaction (i.e., gLV) models are derivable from

mechanistic (i.e., kinetic) models through empirical manipulation of equations, which is however

limited to special forms of kinetics and therefore cannot be generalizable (32). By contrast, our

chain rule-based formulation allows us to handle any complex forms of kinetic equations with no

such constraints. Consequently, KIDI enables incorporation any forms of kinetic equations as

demonstrated using a double Michaelis-Menten kinetics as a demonstration example.

Dynamic variations in microbial interactions inferred by KIDI were experimentally

validated using a synthetic binary consortium of two metabolically engineered auxotrophic *E. coli*

mutants that cross-feed amino acids they cannot synthesize (i.e., tryptophan and tyrosine). A

coordinated design of experiments provided multiple sets of data required for determining kinetic

and stoichiometric parameters in the mechanistic model along with substrate concentrations, which

are key inputs for quantifying environment-dependent interactions. Despite diverse culture

conditions including axenic and binary growth in batch and semi-batch modes, our model with a

single set of parameters showed satisfactory fit to the three training datasets and provided

consistency with the validation dataset set aside in advance. Such a fair goodness of fit indicates

the acceptability of model parameters and therefore the subsequent inference of microbial

interactions. The model

Our kinetic model also shows consistency with the analysis of energetic cost of

synthesizing amino acids in the literature. Mee et al. (30) estimated the energetic cost for the

synthesis of 14 individual amino acids based the amounts of extracellularly supplemented amino

acid and the observed growth yield of *E. coli* auxotrophic mutants. From the linear relationships

between these two variables, they calculated the supplemented amounts of amino acids *per cell*,

which were 1.5×10^7 and 3.7×10^7 for the tryptophan and tyrosine auxotrophic *E. coli* mutants,

respectively. These two quantities correspond to the stoichiometric coefficients $Y_{S_i^+/X_i}(i = 1,2)$

in our kinetic model, which were determined to be $1.550 (= Y_{S_1^+/X_1})$ and $2.994 (= Y_{S_2^+/X_2})$ [mg/OD]

through data fit (**Table 2**). As the direct one-to-one matching between them might not be feasible,

e.g., due to different units of biomass (i.e., cell number in Mee et al. (30) vs. OD in this work), we

compared the ratios, which showed consistency between the two studies, i.e., $\frac{1.5 \times 10^7}{3.7 \times 10^7} \approx 0.41$ vs.

$\frac{1.550}{2.994} \approx 0.52$. Both results imply that compared to tyrosine, the synthesis of tryptophan is more

costly. In support of this, Mee et al. (30) estimated the biosynthetic cost for tryptophan is about

43% higher than that for tyrosine.

We highlight that inferring environment-dependent interactions and their dynamic

variations is a critical capability uniquely associated with KIDI. Even in a simple binary

consortium considered in this work, KIDI provides new insights into interspecies interactions such

as asymmetry between the two amino acid auxotrophs, which might not be obtainable otherwise.

In perturbed growth experiments with glucose FBs (as in **Fig. 4B**), for example, KIDI identified

that (1) $a_{1,2}^+ > a_{2,1}^+$ while the shared substrate (glucose) is abundant, implying that the tryptophan

auxotroph (X_1) does not support the growth of the tyrosine auxotroph (X_2) as much as X_2 does for

X_1; (2) $a_{1,2}^- < a_{2,1}^-$ after the completion of initially added glucose until tryptophan is depleted,

implying that less favorable supporters during cooperation become worse enemies when the

relationship turned into competition.

While KIDI determines pairwise interaction terms following a generalized Lotka-Volterra

framework, it is also capable of accounting for the influences of additional species, provided these

impacts are reflected in the growth kinetics. For instance, in the case study of this article, species

1 and 2 exhibit a complex relationship, competing for glucose uptake while cooperating for amino

acid exchange. However, if a third species is introduced that synthesizes and contributes amino

acids to the environment more rapidly than the existing members, the dynamic between species 1

and 2 shifts. Their reliance on each other for amino acids diminishes, transforming their mixed

relationship into pure competition due to the influence of the third species.

The chain rule formulation in KIDI successfully estimates interactions from given kinetics,

a capability that remains effective across microbial communities of varying complexities. The

primary challenge, however, is in identifying growth kinetics. This issue is especially pronounced

in complex microbial communities where prior knowledge of interspecies interactions is lacking.

Considering these limitations, we showcased KIDI's effectiveness using simpler consortia, which

simplify the experimental data collection needed for parameter determination in the mechanistic

model. The study of model microbial consortia, extracted from natural communities, has been

instrumental in enhancing our understanding of complex ecological systems (33, 34).

For KIDI to be effectively applied to simpler consortia, comprehensive measurements of

all chemical and biological species involved in interspecies interactions are essential, as precise

parameter identification is otherwise challenging. In our study, integrating complete temporal

amino acid profiles would improve parameter identification accuracy. Typically, metabolite levels

exchanged between species, such as amino acids in our case, are low and often fall below detection

limits. Additional analysis of axenic culture data would help address this issue, as implemented in

our work.

Despite several challenges mentioned above, it's important to note that these stem from the

difficulties in building robust kinetic models, rather than being a limitation of KIDI itself. KIDI's

primary function is to deduce the temporal variations in interaction coefficients based on

environmental variables. Its unique ability to handle context-dependent interactions opens up

various applications. For instance, KIDI can serve as a probing tool to investigate how assumed

growth kinetics and environment-mediated mechanisms lead to specific interactions and their

temporal evolution. This aspect is crucial for understanding the link between growth mechanisms

of particular species and their interactions. Moreover, KIDI can greatly aid in advancing network

inference techniques. The development of new algorithms for predicting microbial interactions is

often hindered by a lack of benchmark data, a gap that KIDI can help fill.

KIDI's utility goes beyond microbial ecology, encompassing a wide range of community

ecology fields. This versatility comes from the fact that context dependency is not solely a

microorganism trait but is also common in macro-organisms like plants and animals. For example,

KIDI is applicable to classic ecological models, such as MacArthur's consumer-resource model

(35). In this model, MacArthur quantifies the impact of consumer j on consumer i ($a_{i,j}$) based on

resource population densities and associated parameters, with an underlying assumption that

resource populations change more rapidly than consumer populations. KIDI is adept at deducing

the competitive relationships among consumers using the chain rule, as detailed in Eq. (4)

(**Supplementary Text**). This adaptability of the KIDI framework enables its extensive use in

analyzing context-dependent interactions within and across different biological kingdoms in a

variety of ecological systems.

MATERIALS AND METHODS

Mathematical definition of interaction coefficients

The dynamic change in population i in a community can be formulated in a general form as follows:

392
$$\frac{1}{x_i}\frac{dx_i}{dt} = f_i(x_1, x_2, \cdots, x_N), \quad i = 1, 2, \cdots, N \tag{8}$$

where x_i is the population density of species i, the left-hand side defines the specific growth rate

of species i, and the function $f_i(x_1, x_2, \cdots, x_N)$ represents a nonlinear dependence of the specific

growth rate of species i on population densities of other species.

Using a Taylor expansion, the right-hand side of Eq. (8) can be represented as a series of

polynomial terms, i.e.,

$$f_i = f_{i,0} + \sum_{j=1}^{N} \left(\frac{\partial f_i}{\partial x_j}\right)_0 x_j + H.O.T., \quad i = 1,2,\cdots,N \tag{9}$$

where the subscript 0 denotes a chosen reference condition, $H.O.T.$ is higher-order terms.

Neglecting the $H.O.T.$ in Eq. (9), a gLV equation describes specific growth of species i using a

linear equation, i.e.,

$$f_i(x_1, x_2, \cdots, x_N) = f_{i,0} + \sum_{j=1}^{N} a_{i,j} x_j, \quad i = 1,2,\cdots,N \tag{10}$$

where interaction coefficient $a_{i,j}$ denotes the effect of species population j on the specific growth

of species i. For a binary community, Eq. (10) reduces to

$$f_i(x_i, x_j) = f_{i,0} + a_{i,i} x_i + a_{i,j} x_j, \quad i = 1,2 \tag{11}$$

where $f_{i,0}$ is the basal growth rate of species i, $a_{i,i}$ is the intra-specific interaction coefficient, and

$a_{i,j}$ is the inter-specific interaction coefficient.

From Eq. (11), binary interaction coefficients in gLV are defined as follows:

$$a_{i,j} \equiv \frac{\partial f_i(x_i, x_j)}{\partial x_j} \tag{12}$$

The typical formulation assumes that $a_{i,j}$ is constant, which however leads the gLV model to fail

to capture delicate dynamics of microbial interactions. Indeed, $a_{i,j}$ is a dynamic parameter (i.e.,

$a_{i,j}(t)$) that changes its value in varying environmental conditions as shown in the next section.

Formulation of interaction coefficients as function of environmental variables

For simplicity, we assume in this section that f_i in the previous section is represented by kinetic

growth rate, μ_i, which is formulated as a function of nutrient concentrations in the environment.

In the circumstance considered in **Fig. 1**,

$$\mu_i(t) = \mu_i(s_i^+(t), s^-(t)) \tag{13}$$

where $s_i^+(t)$ is the concentration of the nutrient (such as tryptophan or tyrosine) at time t that

species i needs to get either from its partner or the environment, and $s^-(t)$ represents the

concentration of the shared nutrient (i.e., glucose) at time t that two species compete for.

Based on the chain rule, we formulate $a_{i,j}(t)$ as a function of nutrient concentrations by

plugging Eq. (13) into (12), i.e.,

$$a_{i,j}(t) = \frac{\partial}{\partial x_j}[\mu_i(s_i^+(t), s^-(t)]$$

$$= \frac{\partial}{\partial s_i^+(t)}[\mu_i(s_i^+(t), s^-(t))]\frac{\partial s_i^+(t)}{\partial x_j(t)} + \frac{\partial}{\partial s^-(t)}[\mu_i(s_i^+(t), s^-(t))]\frac{\partial s^-(t)}{\partial x_j(t)} \tag{14}$$

Note that the two terms on the R.H.S. of Eq. **(14)** represent the positive and negative effects of

species j on i through environmental variables, i.e., $a_{i,j}^+$ and $a_{i,j}^-$ as defined below

$$a_{i,j}^+(t) \equiv \frac{\partial}{\partial s_i^+(t)}[\mu_i(s_i^+(t), s^-(t))]\frac{\partial s_i^+(t)}{\partial x_j(t)} \tag{15}$$

$$a_{i,j}^-(t) \equiv \frac{\partial}{\partial s^-(t)}[\mu_i(s_i^+(t), s^-(t))]\frac{\partial s^-(t)}{\partial x_j(t)} \tag{16}$$

In a similar fashion, we can formulate intra-specific interaction coefficients as functions of

environmental variables, i.e.,

$$a_{i,i}(t) = \frac{\partial}{\partial x_i}[\mu_i(s_i^+(t), s^-(t)]$$

$$= \frac{\partial}{\partial s_i^+(t)}[\mu_i(s_i^+(t), s^-(t))]\frac{\partial s_i^+(t)}{\partial x_i(t)} + \frac{\partial}{\partial s^-(t)}[\mu_i(s_i^+(t), s^-(t))]\frac{\partial s^-(t)}{\partial x_i(t)} \tag{17}$$

Final forms of $a_{i,j}(t)$ ($a_{i,j}^+(t)$ and $a_{i,j}^-(t)$) and $a_{i,i}(t)$ depend on specific kinetics for

$\mu_i(s^-(t), s_i^+(t))$. While the symbol (t) is dropped for simplicity, all $a_{i,j}$'s in the main text are

dynamic interaction coefficients, the values of which are changing in time as formulated in this

section.

Microorganisms and culture conditions

Two auxotrophic *Escherichia coli* (*E. coli*) mutant strains, JW2581-1 and JW1254-2 originally

derived from the same strain (BW25113), were purchased from *E. coli* Genetic Stock Center

(CGSC) at Yale University (http://cgsc2.biology.yale.edu/). As experimentally validated in the

literature (36), these mutant strains, JW2581-1 (ΔtyrA) and JW2581-2 (ΔtrpC), are incapable of

growing without supplementation of Tyrosine and Tryptophan, respectively. Each strain was

incubated overnight at 37°C and 225 rpm in 50 ml of Falcon tube containing 5 ml of Lysogeny

broth (LB) supplemented with 33 μg/l kanamycin. Culture cells were collected and centrifuged

them at 16,000 g, 4°C for 1.5 min. The cell pellets were washed with K3 basal medium to remove

residual amino acids in the samples. The washed cells were resuspended and transferred to 150 ml

flasks carrying 25 ml of K3 defined minimal medium (5) containing glucose and 33 μg/l

kanamycin, and cultivated at 37°C and 225 rpm. An initial absorbance at 600 nm (OD 600) was

0.04 with equivalent cell ratio. For batch mode, 4.5 g/l glucose was supplied in the culture medium.

For the fed-batch mode, 0.5 g/l of an initial glucose concentration was used to shorten the lag phase

and 3 or 5 glucose FeedBeads (Kühner, Basel, Switzerland), releasing glucose at constant rate,

were added when OD 600 reached 0.2. We collected 500 ul of culture medium from each flask

and centrifuged them at 16,000 g for 1.5 min. The supernatant and pellets were stored at -20°C

until further analysis.

Analysis of glucose concentration in the culture medium

The concentration of glucose was analyzed by a high-performance liquid chromatography (HPLC)

system (Agilent, Santa Ciara, CA) equipped with a 1260 refractive index detector (RID) and an

Aminex HPX-87H column (Bio-Rad, Hercules, CA). Five microliter of filtered supernatants were

injected. Analytes were separated isocratically using 5 mM sulfuric acid at a flow rate of 0.7

ml/min.

Analysis of amino acids concentration in the culture medium

The amino acids in 10 μl of filtered supernatants were analyzed using an ultra-performance liquid

chromatography (UPLC) (Waters, Milford, MA) coupled with a micrOTOF II mass spectrometry

(TOF-MS) system (Bruker, Bremen, Germany). Analytes were measured using a tunable UV

detector at 210 and 397 nm. The amino acids were separated by an Agilent Poroshell 120 EC-C18

column at 30°C. The 1 % (v/v) of formic acid in water (mobile phase A) and 1 % (v/v) of formic

acid in acetonitrile (mobile phase B) were used, respectively. The amino acids separation was

obtained at a flow rate of 0.3 ml/min with a gradient program that allowed 100% of mobile phase

A until 2.1 min followed by increasing mobile phase B to 40% for 2 min and then equilibrated at

0% of eluent B in a total analysis time of 6 min. Analysis of the amino acids was performed using

electrospray ionization (ESI) and full-scan TOF-MS spectra (50 - 650 m/z) with 500 V end plate

voltage and 4.5 kV capillary voltage. Nebulizer gas and drying gas were supplied in 1.8 bar and 8

ml/min, respectively. The dry temperature was kept at 220°C.

Quantification of cell ratio in microbial consortium

Quantitative PCR (qPCR) was carried out in a 96-well plate by using a CFX96 Real-Time

Detection System (Bio-Rad, Hercules, CA, USA). The pellets were resuspended in ultra-pure

water to make consistent concentration (OD600 = 0.4) and then, the 200 µl solution was transferred

to a 250 µl PCR tube. The solutions were incubated at 98℃ for 10 min for cell disruption using a

T100 Thermal Cycler (Bio-Rad). The lysed cells were transferred to 1.5 ml of tubes and

centrifuged at 20, 000 x g for 2 min. The supernatants were analyzed by qPCR. The qPCR mixture

was composed as follows: 3 µl of 10X Xtensa® buffer, 0.3 µl of primer mix (50 uM for each),

0.15 µl of i-Taq (i-DNA Biotechnology, Singapore), 3 µl of 25 mM $MgCl_2$, 5 µl of purified cell

lysate, and 18.55 µl of ultra-pure water. The thermal cycling was programmed as follows: 95℃

for 1 min and 30 cycles of (95℃ for 20 sec, 55℃ for 20 sec, 68℃ for 40 sec). The primers for

qPCR analysis to quantify the different *E. coli* strains were provided in **Table S3**. The qPCR

analysis was performed in triplicate for each sample.

Data availability. The data and code for this study are available at

https://github.com/hyunseobsong/kidi.

ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by

the Korea government (MSIT) (No. 2020R1A2C2007192), and the Korea Institute of Planning

and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (iPET) funded by the

MAFRA (32136-05-1- HD050). This material is also based on work supported by the National

Science Foundation under Grant No. 2125155 and Nebraska Tobacco Settlement Biomedical

Research Enhancement Funds to H-SS.

References

1. Sorboni SG, Moghaddam HS, Jafarzadeh-Esfehani R, Soleimanpour S. 2022. A Comprehensive Review on the Role of the Gut Microbiome in Human Neurological Disorders. Clinical Microbiology Reviews 35.

2. Vijay A, Valdes AM. 2022. Role of the gut microbiome in chronic diseases: a narrative review. European Journal of Clinical Nutrition 76:489-501.

3. Sokol NW, Slessarev E, Marschmann GL, Nicolas A, Blazewicz SJ, Brodie EL, Firestone MK, Foley MM, Hestrin R, Hungate BA, Koch BJ, Stone BW, Sullivan MB, Zablocki O, Pett-Ridge J, Consortium LSM. 2022. Life and death in the soil microbiome: how ecological processes influence biogeochemistry. Nature Reviews Microbiology 20:415-430.

4. Trivedi P, Batista BD, Bazany KE, Singh BK. 2022. Plant-microbiome interactions under a changing world: responses, consequences and perspectives. New Phytologist 234:1951-1959.

5. Albright MBN, Louca S, Winkler DE, Feeser KL, Haig SJ, Whiteson KL, Emerson JB, Dunbar J. 2022. Solutions in microbiome engineering: prioritizing barriers to organism establishment. Isme Journal 16:331-338.

6. Konopka A, Lindemann S, Fredrickson J. 2015. Dynamics in microbial communities: unraveling mechanisms to identify principles. Isme Journal 9:1488-1495.

7. Marsh JW, Ley RE. 2022. Microbiome engineering: Taming the untractable. Cell 185:416-418.

8. Perez-Garcia O, Lear G, Singhal N. 2016. Metabolic Network Modeling of Microbial

Interactions in Natural and Engineered Environmental Systems. Frontiers in

Microbiology 7.

9. Allen BH, Gupta N, Edirisinghe JN, Faria JP, Henry CS. 2022. Application of the

metabolic modeling pipeline in KBase to categorize reactions, predict essential genes,

and predict pathways in an isolate genome, p 291-320, Microbial Systems Biology.

Springer.

10. Song H-S, Nelson WC, Lee J-Y, Taylor RC, Henry CS, Beliaev AS, Ramkrishna D,

Bernstein HC. 2018. Metabolic network modeling for computer-aided design of

microbial interactions.

11. Kessell AK, McCullough HC, Auchtung JM, Bernstein HC, Song HS. 2020. Predictive

interactome modeling for precision microbiome engineering. Current Opinion in

Chemical Engineering 30:77-85.

12. Song HS, Lindemann SR, Lee DY. 2021. Editorial: Predictive Modeling of Human

Microbiota and Their Role in Health and Disease. Frontiers in Microbiology 12.

13. Choi YM, Lee YQ, Song HS, Lee DY. 2020. Genome scale metabolic models and

analysis for evaluating probiotic potentials. Biochemical Society Transactions 48:1309-

1321.

14. Song HS. 2018. Design Principles of Microbial Communities: From Understanding to

Engineering. Current Genomics 19:699-700.

15. Lee JY, Haruta S, Kato S, Bernstein HC, Lindemann SR, Lee DY, Fredrickson JK, Song

HS. 2020. Prediction of Neighbor-Dependent Microbial Interactions From Limited

Population Data. Frontiers in Microbiology 10.

549 16. Song HS, Lee JY, Haruta S, Nelson WC, Lee DY, Lindemann SR, Fredrickson JK,

550   Bernstein HC. 2019. Minimal Interspecies Interaction Adjustment (MIIA): Inference of

551   Neighbor-Dependent Interactions in Microbial Communities. Frontiers in Microbiology

552   10.

553 17. Kato S, Haruta S, Cui ZJ, Ishii M, Igarashi Y. 2008. Network relationships of bacteria in

554   a stable mixed culture. Microbial Ecology 56:403-411.

555 18. Yamamoto K, Haruta S, Kato S, Ishii M, Igarashi Y. 2010. Determinative Factors of

556   Competitive Advantage between Aerobic Bacteria for Niches at the Air-Liquid Interface.

557   Microbes and Environments 25:317-320.

558 19. Klitgord N, Segre D. 2010. Environments that Induce Synthetic Microbial Ecosystems.

559   Plos Computational Biology 6.

560 20. Song H-S, Cannon WR, Beliaev AS, Konopka A. 2014. Mathematical modeling of

561   microbial community dynamics: a methodological review. Processes 2:711-752.

562 21. Faust K, Raes J. 2012. Microbial interactions: from networks to models. Nature Reviews

563   Microbiology 10:538-550.

564 22. Friedman J, Alm EJ. 2012. Inferring Correlation Networks from Genomic Survey Data.

565   Plos Computational Biology 8.

566 23. Fang HY, Huang CC, Zhao HY, Deng MH. 2015. CCLasso: correlation inference for

567   compositional data through Lasso. Bioinformatics 31:3172-3180.

568 24. Ban YG, An LL, Jiang HM. 2015. Investigating microbial co-occurrence patterns based

569   on metagenomic compositional data. Bioinformatics 31:3322-3329.

570 25. Joseph TA, Shenhav L, Xavier JB, Halperin E, Peer I. 2020. Compositional Lotka-

571   Volterra describes microbial dynamics in the simplex. Plos Computational Biology 16.

572   26.   Fisher CK, Mehta P. 2014. Identifying Keystone Species in the Human Gut Microbiome

573   from Metagenomic Timeseries Using Sparse Linear Regression. Plos One 9.

574   27.   Stein RR, Bucci V, Toussaint NC, Buffie CG, Ratsch G, Pamer EG, Sander C, Xavier JB.

575   2013. Ecological Modeling from Time-Series Inference: Insight into Dynamics and

576   Stability of Intestinal Microbiota. Plos Computational Biology 9.

577   28.   Li CH, Chng KR, Kwah JS, Av-Shalom TV, Tucker-Kellogg L, Nagarajan N. 2019. An

578   expectation-maximization algorithm enables accurate ecological modeling using

579   longitudinal microbiome sequencing data. Microbiome 7.

580   29.   Saeedian M, Pigani E, Maritan A, Suweis S, Azaele S. 2022. Effect of delay on the

581   emergent stability patterns in generalized Lotka-Volterra ecological dynamics.

582   Philosophical Transactions of the Royal Society a-Mathematical Physical and

583   Engineering Sciences 380.

584   30.   Mee MT, Collins JJ, Church GM, Wang HH. 2014. Syntrophic exchange in synthetic

585   microbial communities. Proceedings of the National Academy of Sciences of the United

586   States of America 111:E2149-E2156.

587   31.   Kerner A, Park J, Williams A, Lin XXNN. 2012. A Programmable Escherichia coli

588   Consortium via Tunable Symbiosis. Plos One 7.

589   32.   Momeni B, Xie L, Shou WY. 2017. Lotka-Volterra pairwise modeling fails to capture

590   diverse pairwise microbial interactions. Elife 6.

591   33.   McClure R, Naylor D, Farris Y, Davison M, Fansler SJ, Hofmockel KS, Jansson JK.

592   2020. Development and Analysis of a Stable, Reduced Complexity Model Soil

593   Microbiome. Frontiers in Microbiology 11.

594   34.   Kylilis N, Tuza ZA, Stan GB, Polizzi KM. 2018. Tools for engineering coordinated

595   system behaviour in synthetic microbial consortia. Nature Communications 9.

596   35.   Chesson P. 1990. MacArthur's consumer-resource model. Theoretical Population Biology

597   37:26-38.

598   36.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M,

599   Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame, single-gene

600   knockout mutants: the Keio collection. Mol Syst Biol 2:2006 0008.

Table captions

Table 1. Model equations with kinetic parameters and stoichiometric coefficients determined

through the model fit to experimental data collected under various limiting conditions. R_i is the

stoichiometric growth reaction for X_i, and Y_{S^-/X_i}, $Y_{S_i^+/X_i}$, and $Y_{S_j^+/X_i}$ denote the stoichiometric

coefficients for S^-, S_i^+, and S_j^+ associated with the growth of X_i. s^-, s_i^+, and s_j^+ respectively

denote concentrations of S^-, S_i^+, and S_j^+, x_i is the population density of X_i, μ_i is the specific

growth rate of X_i, $k_{d,i}$ is the death rate of X_i, and q_{s^-} is the substrate releasing rate from FBs in

glucose-limited semi-batch cultures (i.e., $q_{s^-} = 0$ in a batch mode). μ_i^{max} is the maximal growth

rate, and K_i^- and K_i^+ are half-saturation constants associated with the consumption of S^- and S_i^+.

   Figure captions

Fig. 1. Conceptual illustration of context-dependent microbial interactions in a binary

consortium dictated by the environmental contexts. Two species X_1 and X_2 compete for the

substrate S^- but cooperate by cross-feeding metabolite S_1^+ and S_2^+ (center panel). This mixed

relationship between X_1 and X_2 diverge into six different types of interactions by excessive

addition of specific substrates S_1^+, S_2^+, and/or S^-. Symbols next to the arrows denote the

substrate(s) excessively added to the environment. (**c**) Model microbial consortium using

tyrosine or tryptophan auxotrophic *E. coli* strains. Glucose is sole carbon source for both *E. coli*

strains. GLC, Tyr and Trp denote glucose, tyrosine, and tryptophan, respectively.

Fig. 2. Experimental data and model simulations for the growth of two *E. coli* mutant strains in

axenic and binary culture conditions: **A** and **B** are cultures of tryptophan auxotrophic and

tyrosine auxotrophic *E. coli* mutants (X_1 and X_2), respectively; **C** and **D** are co-cultures with two

auxotrophs in batch and semi-batch cultures. Detailed culture conditions for the twelve panels

are provided in **Table S2**. Circles and lines denote the experimentally measured values and

simulation results, respectively. Black line denotes simulation results for glucose concentration

(S^-), and the lines in blue, red, and purple are simulated population densities of X_1, X_2, and $X_1 +$

X_2. **Figs. A, B,** and **C** show data fitting to determine model parameters, while the results in **Fig.**

D validate model predictions.

Fig. 3. Inference of dynamic variations of interaction parameters ($\gamma_{i,j}, a_{i,j}, a_{i,j}^+$, and $a_{i,j}^-$) for the

two *E. coli* mutants (X_1 and X_2) co-growing in three batch cultures. Initial substrate

concentrations: **A.** 2 g/l of glucose, 1 mg/l of tryptophan, and 1 mg/l of tyrosine; **B.** 0.5 g/l of

glucose, 1 mg/l tryptophan, and 1 mg/l of tyrosine; and **C.** 0.5 g/l of glucose, 4 mg/l of

tryptophan, and 4 mg/l of tyrosine. Black line denotes simulated concentration of glucose (S^-);

the lines in blue and red indicate the variables and parameters associated with X_1 and X_2,

respectively.

Fig. 4. Inference of dynamic variations of interaction parameters ($\gamma_{i,j}, a_{i,j}, a_{i,j}^+$, and $a_{i,j}^-$) for the

two *E. coli* mutants (X_1 and X_2) co-growing in three semi-batch cultures with 3 glucose FBs

added at 7.5 hours. Initial substrate concentrations: **A.** 2 g/l of glucose, 1 mg/l of tryptophan, and

1 mg/l of tyrosine; **B.** 0.5 g/l of glucose, 1 mg/l tryptophan, and 1 mg/l of tyrosine; and **C.** 0.5 g/l

of glucose, 4 mg/l of tryptophan, and 4 mg/l of tyrosine. Black line denotes simulated

concentration of glucose (S^-); the lines in blue and red indicate the variables and parameters

associated with X_1 and X_2, respectively.

Table 1
654

Stoichiometric equation (R_i) for growth of X_i:

$$R_i : Y_{S^-/X_i}S^- + Y_{S_i^+/X_i}S_i^+ \rightarrow X_i + Y_{S_j^+/X_i}S_j^+, \quad (i,j) = (1,2) \text{ or } (2,1) \tag{T1}$$

Dynamic mass balances:

$$\frac{dx_i}{dt} = \mu_i x_i - k_{d,i} x_i, \quad i = 1,2 \tag{T2}$$

$$\frac{ds^-}{dt} = -Y_{S^-/X_1}\mu_1 x_1 - Y_{S^-/X_2}\mu_2 x_2 (+q_{S^-}) \tag{T3}$$

$$\frac{ds_i^+}{dt} = -Y_{S_i^+/X_i}\mu_i x_i + Y_{S_i^+/X_j}\mu_j x_j, \quad (i,j) = (1,2) \text{ or } (2,1) \tag{T4}$$

Double Monod kinetics:

$$\mu_i = \mu_i^{max}\frac{s_i^+}{(K_i^+ + s_i^+)}\frac{s^-}{(K_i^- + s^-)}, \quad i = 1,2 \tag{T5}$$

Kinetic parameters and stoichiometric coefficients determined through data fit:

Parameter	Value	Parameter	Value
μ_1^{max} [1/h]	2.961×10^{-1}	Y_{S^-/X_1} [g/OD]	1.372
μ_2^{max} [1/h]	1.658×10^{-1}	Y_{S^-/X_2} [g/OD]	1.773
K_1^- [g/l]	3.091×10^{-4}	$Y_{S_1^+/X_1}$ [mg/OD]	1.550
K_2^- [g/l]	3.923×10^{-4}	$Y_{S_1^+/X_2}$ [mg/OD]	2.961
K_1^+ [mg/l]	3.300×10^{-3}	$Y_{S_2^+/X_1}$ [mg/OD]	1.365
K_2^+ [mg/l]	3.881×10^{-4}	$Y_{S_2^+/X_2}$ [mg/OD]	2.994
$k_{d,1}$ [1/h]	1.206×10^{-4}	q_{S^-} (for 3 FBs) [g/l/h]	5.580×10^{-2}
$k_{d,2}$ [1/h]	4.024×10^{-7}	q_{S^-} (for 5 FBs) [g/l/h]	9.300×10^{-2}

657   Figure 1

662   Figure 2

670   Figure 4

SUPPLEMENTARY TEXT:

Connection between KIDI and MacArthur's Consumer-Resource Model

In this supplementary text, we illustrate how KIDI can be extended to traditional ecological models, such as MacArthur's consumer-resource (C-R) model, by elucidating their interconnection. MacArthur (1970) formulated the per-capita growth rate of consumer species i (X_i) and resource species j (R_j) as follows:

$$\frac{1}{X_i}\frac{dX_i}{dt} = b_i \left(\sum_{k=1}^{n} c_{ik} w_k R_k - m_i \right) \quad \text{(A1)}$$

$$\frac{1}{R_k}\frac{dR_k}{dt} = r_k \left(1 - \frac{R_k}{K_k} \right) - \sum_{i=1}^{n} c_{ik} X_i \quad \text{(A2)}$$

where
- X_i is a population density of consumer specie i
- R_k is the population density of resource species k
- w_k is the value of one unit of resource species j to the consumer
- c_{ik} is the rate at which consumer species i captures resource j per unit abundance of resource species j
- m_i is, the maintenance term (i.e., the total value of resource that must be harvested per capita for maintaining the current population density of consumer species i
- b_i is a factor converting the resource excess into the per-capita growth rate
- r_k is per-capita growth rate of resource species j
- K_k is carrying capacity for resource species j

MacArthur assumed faster dynamics of resource populations compared to consumer populations (i.e., r_k and c_{ik} are much larger than $b_i c_{ik} w_k$ and $b_i m_i$) so that Eq. (A2) is reduced to an algebraic equation (by setting $\frac{dR_j}{dt} = 0$). By substituting R_k in Eq. (A1) for X_i's, MacArthur derived a'_{ij} (i.e., the competitive relationship between the two consumers i and j, or more precisely the impact of consumer j on the growth on consumer i) in the Lotka-Volterra (LV) model as follows:

$$a'_{ij} = \sum_{k=1}^{n} \frac{c_{ik} c_{jk} w_k K_k}{r_k} \quad \text{(A3)}$$

where we use the prime notation in a'_{ij} to differentiate it from the definition of a_{ij} commonly used in a generalized LV model. These two notations are related by $-b_i a'_{ij} = a_{ij}$.

709   We now show that the above equation can be derived by KIDI. Interspecies interaction

710   coefficient a_{ij} (not a'_{ij}) for the system with multiple resources in KIDI are defined as follows:

712
$$a_{ij} = \sum_{k=1}^{n} \left(\frac{\partial \mu_i}{\partial R_k}\right)\left(\frac{\partial R_k}{\partial X_j}\right) \quad \text{(A4)}$$

714   where μ_i is the specific (or per-capita) growth rate of species i (i.e., $\frac{1}{X_i}\frac{dX_i}{dt}$). The first term $\left(\frac{\partial \mu_i}{\partial R_k}\right)$

715   is obtained simply by taking the derivative of the per-capita growth rate for X_i with respect to R_k

716   in Eq. (A1), i.e.,

718
$$\left(\frac{\partial \mu_i}{\partial R_k}\right) = b_i c_{ik} w_k \quad \text{(A5)}$$

720   Next, we take the derivative of the steady-state equation of Eq. (A2) with respect to X_j to give

721   the second term $\left(\frac{\partial R_k}{\partial X_j}\right)$ as follows:

723
$$\left(\frac{\partial R_k}{\partial X_j}\right) = -\frac{K_k c_{jk}}{r_k} \quad \text{(A6)}$$

725   With the substitutions using Eqs. (A5) and (A6), the final form of a_{ij} becomes

727
$$a_{ij} = -b_i \sum_{k=1}^{n} c_{ik} w_k \frac{K_k c_{ij}}{r_k} \quad \text{(A7)}$$

729   The above equation proves $a_{ij} = -b_i a'_{ij}$.

732   **SUPPLEMENTARY TABLES AND FIGURES**

734   **Table S1.** Initial culture conditions for experimental data displayed in Fig. 2.

Exp #	Corresponding figures	Initial inoculum size (OD)	Glucose (mg/l)	Tryptophan (mg/l)	Tyrosine (mg/l)	# of FBs added	Time FBs added (hr)
1	Fig. 2A (top panel)	ΔTrp only: 0.01	5.838	10	-	-	-
2	Fig. 2A (middle panel)	ΔTrp only: 0.01	5.870	20	-	-	-
3	Fig. 2A (bottom panel)	ΔTrp only: 0.01	5.936	40	-	-	-
4	Fig. 2B (top panel)	ΔTyr only: 0.01	5.920	-	10	-	-
5	Fig. 2B (middle panel)	ΔTyr only: 0.01	5.909	-	20	-	-
6	Fig. 2B (bottom panel)	ΔTyr only: 0.01	5.863	-	40	-	-
7	Fig. 2C (top panel)	ΔTrp: 0.01 ΔTyr: 0.01	3.851	0.4	0.4	-	-
8	Fig. 2C (middle panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.626	0.4	0.4	3	7.5
9	Fig. 2C (bottom panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.620	0.4	0.4	5	7.5
10	Fig. 2D (top panel)	ΔTrp: 0.01 ΔTyr: 0.01	2.878	0.4	0.4	-	-
11	Fig. 2D (middle panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.375	0.4	0.4	3	10
12	Fig. 2D (bottom panel)	ΔTrp: 0.01 ΔTyr: 0.01	0.377	200	200	3	10

738   **Table S2.** Sequence of the primers used in the qPCR analysis.

Primer	Sequence
Trp F	GCCGATGCCTGCTTATTA
Trp R	GCTCCTGTTCCTCTTCAT
Tyr F	CATTATGTCGTCAGAGCG
Tyr R	CCTTGCGGAAACTGTCAA

740
741

745 **Figure S1.** Zoom-in views of in interaction parameters, $a_{i,j}$, $a_{i,j}^+$, and $a_{i,j}^-$ in **Figs. 3A, 3B, and**
746 **3C,** respectively. The color scheme is the same as in **Fig. 3**.

Figure S2. Zoom-in views of in interaction parameters, $a_{i,j}, a_{i,j}^+$, and $a_{i,j}^-$ in **Figs. 4A, 4B, and 4C,** respectively. The color scheme is the same as in **Fig. 4**.

758 **Figure S3.** The predicted impacts of the number of added FBs on the substrate profile (S^-), the
759 population dynamics (X_1 and X_2), and the interaction parameter ($\gamma_{i,j}$) in fed-batch cultures with
760 initial substrate concentrations of 2 g/l of glucose, 0.4 mg/l of tryptophan, and 0.4 mg/l of
761 tyrosine. The number of glucose BFs added at 7.5 hours: **A.** 3 FBs; **B.** 6 FBs; and **C.** 10 FBs.
762

Figure S4. Simulated substrate concentrations (top), and biomass concentrations (bottom) with
and without amino acid cross-feeding in the batch culture with excessive provision of glucose.
Initial substrate concentrations of amino acids: 0.4 mg/l of tryptophan and 0.4 mg/l of tyrosine.
Solid lines denote the case with cross-feeding; dashed lines denote the case of preventing cross-
feeding in simulations.