CRISPR-Cas-based Microhomology-Mediated End Joining for Precise Gene Replacement in Plant

Tien Van Vu[1,†], Gah-Hyun Lim[2,3], Seung Hee Choi[3], Ju Yeon Moon[3], Ngan Thi Nguyen[1], Swati Das[1], Mil Thi Tran[1,3], Yeon Woo Sung[1], Jihae Kim[1], Young Jong Song[1], Suk Weon Kim[3], Jae Cheol Jeong[3], Jae-Yean Kim[1,4,†]

[1]Division of Applied Life Science (BK21 FOUR Program), Plant Molecular Biology and Biotechnology Research Center, Gyeongsang National University, Jinju 660-701, Republic of Korea.

[2]Department of Biological Sciences, Pusan National University, Busan 46241, Republic of Korea

[3]Biological Resource Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Jeongeup, 56212, Republic of Korea

[4]Nulla Bio Inc 501 Jinju-daero, Jinju 52828, Republic of Korea.

[†]Correspondence: Tien Van Vu: tienvu.agi@gmail.com (ORCID: 0000-0002-6369-7664). Jae-Yean Kim: kimjy@gnu.ac.kr (ORCID ID : 0000-0002-1180-6232).

Running title: MMEJ-mediated precise gene replacement in plants.

Abstract

Key Message

We designed and tested a method, termed PREMJ, for precise gene replacement using the CRISPR-Cas-mediated double-stranded break (DSB) formation and repairing the DSBs via microhomology-mediated end joining (MMEJ) with MMEJ donor template carrying desired base changes and microhomologies flanking the DSB ends. PREMJ showed feasibility in precise gene replacement in protoplasts of tomatoes, lettuce, and cabbage, albeit its efficacy in *Agrobacterium*-mediated plant transformation requires further optimization.

Abstract

Precise gene or allele replacement is a desirable technology, but implementing it in plants remains challenging. CRISPR-Cas-based approaches, such as gene targeting (GT) and prime editing (PE), have opened up new possibilities for precise gene replacement in plants. However, their editing size and efficiency still need improvement. Recently, strategies using canonical nonhomologous end-joining (cNHEJ) and microhomology-mediated end joining (MMEJ) have been considered promising alternatives for precise gene replacement in yeast and mammals. However, these approaches have not been extensively explored and applied to plants. Here, we proposed and tested a CRISPR-Cas-based tool, termed PREMJ (precision gene replacement via microhomology-mediated end joining), for precision gene replacement in plants. The PREMJ strategy employing 20-bp microhomology MMEJ donors (~100 bp lengths) and a canonical nonhomologous end-joining (cNHEJ) inhibitor, NU7441, produced high targeted gene replacement efficiencies, up to 1.60 ± 0.14, 4.47 ± 1.98, and 8.98 ± 4.73 % in protoplasts of tomato, lettuce, and cabbage, respectively. Our data also revealed the critical impacts of the microhomology length and NU7441 concentration on PREMJ-based precision gene editing in plants. Although obtaining edited plants remains challenging due to inefficient protoplast regenerations and *Agrobacterium*-mediated delivery, PREMJ may significantly contribute to precision gene editing in plants competent to PREMJ complex delivery and plant regeneration.

Keywords: MMEJ, Precision gene editing, Microhomology, CRISPR/Cas, DNA repair, PITCh

Introduction

Precise modification of bases/genes of interest within their genomic context is highly desirable for gene functionalization and genetic engineering applications. However, efficient precision gene editing remains challenging in plants (Schindele et al. 2020). With the advent of the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated (Cas) technology (Jinek et al. 2012), plant precision gene editing has been revolutionized with homology-directed gene targeting (HGT) (Cermak et al. 2015; Wang et al. 2017; Wolter and Puchta 2019; Zhang et al. 2022; Vu et al. 2020) and prime editing (PE) (Anzalone et al. 2020; Lin et al. 2021; Vu et al. 2024b). With HGT, we can precisely edit genes from a single base to kilobase scales, while narrower-range precise gene editing could be achieved with the PE editing tools. However, HGT still requires improvement for practical applications to a wide range of *Arabidopsis* loci (Zhang et al. 2022) and crop plant loci due to low editing efficiency and inaccessibility of the genomic contexts of the loci (Wang et al. 2017; Vu et al. 2021a; Zhang et al. 2022). In the case of prime editors, while substantial improvements have been made to achieve efficient editing of short edits (Lin et al. 2021; Zong et al. 2022; Lu et al. 2020b; Vu et al. 2024b), precise insertion of DNA sequences with more than 50-bp lengths remains challenging (Vu et al. 2024a).

Double-stranded breaks (DSBs) formed by the activity of fully functional Cas9 proteins may be repaired majorly by canonical nonhomologous end-joining (cNHEJ) and homologous recombination (HR), and the former is the dominant pathway (Puchta 2005). MMEJ, previously thought to be an alternative pathway of the cNHEJ, has been shown to play a more important role in DSB repair (Tan et al. 2020; Ata et al. 2018; Sharma et al. 2015; Vu et al. 2021b). Recently, cNHEJ and microhomology-mediated end joining (MMEJ) mechanisms were repurposed for CRISPR/Cas9-based precise gene replacement at high efficiency and specificity in yeast and animals (Sakuma et al. 2016; Yao et al. 2017; Ata et al. 2018; Hayashi and Tanaka 2019; Li et al. 2019; Matsuzaki et al. 2024). In plants, Lu and co-workers engineered a cNHEJ-mediated CRISPR/Cas9-based precise gene insertion at high efficiency by employing a single cut at genomic sites in rice (Lu et al. 2020a). However, a system exploiting cNHEJ and MMEJ for precise gene replacement has not been developed in plants.

Here, we experimentally tested our previously proposed MMEJ-mediated precise gene editing (Vu et al. 2021b) and evaluated a novel MMEJ-based system for precise gene replacement, termed

PREMJ, in plant protoplasts. Using PEG-mediated protoplast transfection and targeted deep

sequencing methods, we successfully obtained highly precise editing efficiency in tomatoes,

lettuce, and cabbage. The MMEJ-based precise gene replacement could be further improved by

optimizing the microhomology lengths, donor template concentration, and suppression of cNHEJ

by NU7441. However, PREMJ remained challenging with *in planta Agrobacterium*-mediated

transformation methods, possibly due to the inefficiency in delivering sufficient amounts of MMEJ

donors. Nevertheless, this tool may still offer an alternative way for precise gene editing in plants

that facilitates efficient PREMJ tool delivery and regeneration and thereby help to advance crop

breeding in the future.

Materials and methods

Donor DNA preparation for RNP works

Chemically modified donors (Supplemental Table 1; Supplemental Supplemental file 1) were

prepared by annealing of synthesized oligos or by PCRs. The 5' end modified cNJ.HPAT3-1 was

prepared by PCRs using 5' modified oligos (5' phosphorylated, phosphorothioate bond addition

to the phosphodiester linkage between the first and the second nucleotides) (Supplemental Table

1) synthesized by Bioneer (Korea) without a template. The MMEJ donors were prepared by PCRs

using complementary oligos (Supplemental Table 1) synthesized by Bioneer (Daejeon, Korea)

without templates. A high-fidelity DNA Taq polymerase was used for the PCR amplifications. The

PCR products were cleaned using a BIOFACT PCR cleanup kit (BIOFACT, Korea). The donor

concentrations were assessed by Nanodrop2000 spectrophotometer (Thermofisher, USA) and

directly used for transfection stored at -20°C for further uses.

Construction of plasmid for *Agrobacterium*-mediated transformation in tomato.

For stable transformation and assessment of the cNHEJ and MMEJ-mediated precision gene

replacement in tomatoes, we designed and cloned the gRNA expression cassettes (Supplemental

file 1) using the Golden-gate cloning system as described previously (Vu et al. 2020). Two gRNA

expression cassettes (gR1.HPAT3 and gR2.HPAT3 for *SlHPAT3*; gR1.HKT1;2 and gR2.HKT1;2

for *SlHKT1;2*) were used to generate two DSBs at each targeted site. The cNHEJ donors

(cNJ.HPAT3-1 for SlHPAT3; cNJ.HKT1;2 for *SlHKT1;2*) and MMEJ donor (MJ.HPAT3-1 for

SlHPAT3; MJ.HKT1;2 for *SlHKT1;2*) were designed to be flanked by two gRNAs (gDR1.HPAT3

and gDR2.HPAT3 for cNJ.HPAT3-1; gDR3.HPAT3-1 and gDR4.HPAT3 for MJ.HPAT3;

gDR1.HKT1;2 and gDR2.HKT1;2 for cNJ.HKT1;2; gDR3.HKT1;2 and gDR4.HKT1;2 for

MJ.HKT1;2) cutting sites (Supplemental file 1). The binary plasmids were constructed to test both

the cNHEJ and MMEJ approaches using a conventional T-DNA and a geminiviral replicon system

(Vu et al. 2020). The NptII selection marker expression cassette (pNOS-NptII-tOCS) is driven by

the NOS promoter and terminated by the OCS terminator (Addgene # 51144). An intron-

containing plant codon-optimized SpCas9 driven by a CaMV 35S promoter, and CaMV 35S

terminator (p35S-pcoCas9I-t35S) was used (Supplemental file 1).

Isolation of protoplasts

Tomato (Solanum lycopersicum cv. Micro-Tom), lettuce (Lactuca sativa L. cv. Cheongchima),

and cabbage (Brassica oleracea) seeds were sterilized with 70% ethanol for 3 min, 1% hypochlorite

solution for 15 min, and washed five times with distilled water. The sterilized seeds were

inoculated in a medium containing 1/2 Murashige and Skoog salts, 0.4 mg/L thiamine HCl, 100

mg/L Myo-inositol, 30 g/L sucrose, and 8 g/L gelrite, pH 5.7. The seedlings were grown in a

growth chamber under a 16 h light/8 h dark photoperiod (100–130 μmol/m2 s) at 25°C for tomato,

and 20oC for lettuce, and 23oC for cabbage.

For protoplast isolation of tomato and cabbage, the cotyledons of 4-day-old tomato seedlings and

the cotyledons of 7-day-old cabbage seedlings were immersed in cell and protoplast washing

solution (CPW) containing 0.5% cellulase (Novozymes, Basgsvaerd, Denmark), 0.5% pectinase

(Novozymes, Bagsværd, Denmark), 1% viscozyme (Novozymes, Bagsværd, Denmark), 3 mM

MES (pH 5.8) and 9% mannitol. After 15 min of vacuum infiltration, the suspension was incubated

for 2-4 hr on a rotary shaker at 50 rpm at 25°C. The suspension was filtered through an eight-layer

gauze and centrifuged for 5 min at 100 g. Protoplasts were separated on a 21% sucrose density

gradient and then collected at the interface of the W5 solution (2 mM MES pH 5.8, 154 mM NaCl,

125 mM $CaCl_2$, 5 mM KCl). The harvested protoplasts were washed three times with W5 solution

and then resuspended in MMG solution (4 mM MES pH 5.7, 0.4 M mannitol, 15 mM $MgCl_2$). The

concentration of protoplasts was determined using a hemocytometer.

For the lettuce protoplast isolation, the cotyledons of 7 d-old seedlings were digested with 10 mL

of enzyme solution (1% [w/v] Viscozyme (Novozymes, Bagsværd, Denmark), 0.5% Celluclast

(Novozymes, Bagsværd, Denmark), and 0.5% Pectinex (Novozymes, Bagsværd, Denmark), 3 mM

MES (2-[N-Morpholino] ethanesulfonic acid), pH 5.7 and 9% mannitol in CPW salts with shaking

at 40 rpm for 4–6 h at 25 °C in the dark. The protoplast mixture was then filtered through a 40 µm

nylon cell strainer and collected by centrifugation at 800 rpm for 5 min in a 14 mL round tube.

The collected protoplasts were re-suspended in W5 solution (2 mM MES [pH 5.7], 154 mM NaCl,

125 mM $CaCl_2$, and 5 mM KCl) and further centrifuged at 800 rpm for 5 min. Finally, the

protoplasts were re-suspended in W5 solution and counted under a microscope using a

hemocytometer. Protoplasts were adjusted to a 1×106/mL density in MMG solution before

transfection. The transfected protoplasts were cultured in protoplast culture medium (MS medium

containing 0.4 mg/L thiamine HCl, 100 mg/L Myo-inositol, 30 g/L sucrose, 0.2 mg/L 2,4-

dichlorophenoxyacetic acid [2,4-D], and 0.3 mg/L 6-benzylaminopurine [BAP], pH 5.7) in the

dark f at 25°C for 4 weeks.

Protoplast transfections

SpCas9 protein was purchased from ToolGen, Inc. (Seoul, South Korea), and guide RNAs were

synthesized by GeneArt Precision gRNA Synthesis Kit (Invitrogen, Massachusetts, USA)

according to the manufacturer's protocol. PEG-mediated RNP and donor transfections were

performed in the previous study (Woo et al. 2015).

For RNP and donor DNA transfections with tomato and cabbage protoplasts, 2×10^5 protoplasts

were transfected with the purified SpCas9 protein (20 µg) premixed with sgRNAs (10 µg each)

and donor templates in PBS buffer followed by incubating for 10 min at 25°C. The RNP complexes

were mixed with protoplasts and then supplemented with an equal volume of 40% PEG

transfection solution (40% PEG 4000, 0.2 M mannitol, and 0.1 M $CaCl_2$). This suspension was

mixed gently and then incubated at room temperature for 10 min. An equal volume of W5 solution

was added for washing, followed by centrifugation at 100 *g* for 5 min. The supernatant was

discarded, and the protoplasts were incubated with 1 ml of W5 solution in the dark at 25°C for 48

h. Afterward, the cells were collected for gDNA isolation and subsequent targeted deep sequencing

analysis.

For lettuce protoplast transfection, SpCas9 protein and sgRNAs were premixed in $1\times$ NEB buffer

3 for at least 10 minutes at room temperature, and 2×10^5 protoplast cells were transfected with

SpCas9 protein (20 µg) premixed with sgRNAs (10 µg each) and donor DNA. A mixture of $2 \times$

10^5 protoplast cells was re-suspended in 200 µl MMG solution and then was slowly mixed with

RNP complex, donor, and 350 µl of PEG solution (40% [w/v] PEG 4000, 0.2 M mannitol, and 0.1

M CaCl2). After incubation for 10 min, the transfected protoplast cells were gently re-suspended

in 650 μL W5 solution. After additional incubation for 10 min, 650 μL W5 solution was added

slowly again and was mixed well by inverting the tube. Protoplasts were pelleted by centrifugation

at 556 rpm for 5 min and washed gently in 1 ml W5 solution. Protoplasts were pelleted by

centrifugation at 556 rpm for 5 min and re-suspended gently in 1 ml WI solution (4 mM MES [pH

5.7], 0.5 M mannitol, and 20 mM KCl). Finally, the protoplasts were transferred into a 60×15

mm petri dish (Falcon), cultured under dark conditions at 25°C for 48 h, and then analyzed for

genome editing efficiency.

***Agrobacterium*-mediated tomato transformation.**

The *Agrobacterium*-mediated tomato transformation was conducted using a protocol published by

Vu and coworkers (Vu et al. 2020) with or without 1 μM NU7441 treatment for 5 days post-

washing. Ten days post-transformation, thirty cotyledon explants were collected per

transformation plate to isolate genomic DNAs and subsequent miniseq analysis. Regenerated

plants were selected in media containing 100 mg/L kanamycin and transferred to soil pots before

analyzing for editing performance. Genomic DNAs were extracted from the plants and analyzed

by PCR amplification of the targeted sequences and by Sanger sequencing.

Targeted deep sequencing

Genomic DNAs were isolated from the protoplasts using the CTAB method. We used the miniseq

sequencing service (MiniSeqTM System, Illumina, USA) to obtain targeted deep sequencing of

the edited genomic sites. Miniseq samples were prepared in three PCRs according to the

manufacturer's guidelines, with genomic DNAs as the first PCR template. The first and second

PCRs used primers listed in Supplemental Table 1, whereas the third PCRs were conducted with

the manufacturer's primers to assign a sample ID. A high-fidelity DNA Taq polymerase (Phusion,

NEB, USA) was used for the PCRs. The miniseq raw data FASTQ files were analyzed using the

Cas-Analyzer tool (Park et al. 2016). The indel examining window was set at 5 bases with a

comparison range that covered both the read ends. A similar analysis was conducted for the

targeted base changes of lettuce and cabbage genes.

 Statistical analysis

191 The editing data, statistical analysis, and plots were further processed by the MS Excel and

192 GraphPad Prism 9.0 programs and are explained in detail in the legends of figures and tables

193 wherever applicable. Statistical analysis was performed using a two-sided, uncorrected Fisher's

194 LSD test. A p-value of 0.05 or lower is considered statistically significant.

Results and discussion

System design and validation for precise gene replacement in plants

In a PREMJ system design, a predefined genomic site with a protospacer adjacent motif (PAM) is

determined first, thus enabling the prediction of the DSB formation site (Figure 1A and 1B). The

sequence between the two predicted cut sites is employed as a donor template so that its DNA

sequence is essentially modified, allowing it to carry targeted base changes to avoid recurrent cuts

after gene replacement (Figure 1A, 1B, and Supplemental Figure 1). In the case of MMEJ-

mediated editing, the predicted flanking DSB ends would then be used to choose microhomologies

(MH1, and MH2, Supplemental Figure 1B) at different lengths. Still, they should be optimally at

least 8 nt (up to 20 nt was reported) in size (xxxxxxx and yyyyyyy in Figure 1B), as suggested

(Nakade et al. 2014; Sakuma et al. 2016; Ata et al. 2018).

Figure 1. HR-independent precision editing approaches using CRISPR/SpCas9. (**A**) Model of
sequence design for cNHEJ-based precise gene replacement. (**B**) Model of sequence design for MMEJ-

bases precise gene replacement. The cut sites are arranged to avoid recurrent cutting. Active PAM sites

are shown in purple. Guide RNA sequences or traces of them in repaired products are underlined. The red

xxxxxxxx represents MH1, whereas the green yyyyyyyy represents MH2. The orange sequences denote

donors, and the gray indicates the genomic sequence being replaced. The red asterisks represent sequence

modifications of interest. The letters n, x, and y represent any nucleotide of A, T, G, or C, except that the

sequence-making gRNA spacers should not have more than 4 T in a row to avoid early transcription

termination. To prevent the recurrent cut, the PAM sequence corresponding to the replaced genomic

sequence should be changed. Usually, nGG/CCn is changed to nHG/CDn where H=A or C or T; D=A or

G or T. One or some bases within the core spacer sequence (i.e., ~12 base pairs upstream/downstream of

nGG/CCn) could also be changed. Such changes should not lead to changes in protein sequences of the

targeted gene. (**C**) Schematic diagrams of cNHEJ- and MMEJ-mediated gene editing of tomato targeted

locus *SlHPAT3* (with their GenBank accession number), targeted sequences, gRNA binding sites, and

donor sense sequences (cNJ.HPAT3-1, MJ.HPAT3-1, MJ.HPAT3-2, and MJ.HPAT3-3). The targeted

genes are shown from the start codon (ATG) to the stop codon (TAA), exons are drawn as blue boxes,

and the discontinuous lines in each gene represent intron parts of the gene; the gRNA binding sequences

at the genomic locus are highlighted in light blue color and arrows with PAMs (bold nucleotides). The

expected cutting sites (3-nt upstream of a PAM) are indicated with the black scissors; the intended base

changes are painted in the red font that is denoted by the discontinuous red boxes and named A1, A2, B,

C, D1, and D2 on the top of the boxes. The cNHEJ donor dsDNA is synthesized with 5'-phosphorylated-

ends, and the phosphodiester bonds between the first and the second nucleotides of 5' ends were

chemically modified with phosphorothioate linkages. The two red, discontinuous crossing lines indicate

the MMEJ-mediated replacement sites of the MJ.HPAT3-1 donor. For the donors with shorter

microhomology, the recombination lengths are reduced accordingly. The diagrams are not drawn to their

scales. (**D**) cNHEJ and MMEJ-mediated editing efficiency at the SlHPAT3 locus. The editing efficiency

is calculated by dividing the total reads containing the intended bases by the total reads without

considering the indel mutation efficiency of the gRNAs. (**E**) Editing frequency of respective repaired

products/alleles revealed from miniseq data. The edited base-changed sets/alleles revealed from miniseq

data are classified according to the presence of the base changes in the edited reads. A total of 11 base-

changed sets/alleles are recorded and plotted with their relative frequency among all the edited reads.

We selected a tomato homolog of *Arabidopsis* hydroxyproline O-arabinosyltransferase 3

(*SlHPAT3*, accession no. Solyc07g021170) for our proof-of-concept experiments (Brooks et al.

2014). We then employed PEG-mediated tomato protoplast transfection to deliver the SpCas9

proteins, two sgRNAs, namely gR1.HPAT3 and gR2.HPAT3 (Figure 1C) for cutting the genomic

sites, and donors for replacing six base pairs of SlHPAT3 (Figure 1C; Supplemental Table 1 and
Supplemental file 1). With the treatments of the SpCas9 protein and either of the gR1.HPAT3 or
gR2.HPAT3, or both of them, without a donor, no precise edited products were found (Figure 1D).
When either the NHEJ or the MMEJ donor was employed with both the gRNAs, precise
replacement occurred, as shown by Sanger sequencing (Supplemental Figure 2 and Supplemental
Table 2). Further analysis of the targeted deep-sequencing data revealed editing efficiency as high
as 2.46 % (Figure 1D and Supplemental Table 3) in the case of the MJ.HPAT3-1 donor. Adding
the cNJ.HPAT3-1 donor also resulted in the precise replacement of some bases pre-introduced in
the donor sequence at a much lower frequency of 0.09 %. Significantly, the edited products of the
cNHEJ donor did not include base changes located at its two ends (Supplemental Tables 2 and 3),
indicating that either the donor terminal ends might be damaged before use for editing or that an
unknown mechanism, possibly oligonucleotide-directed mutagenesis (ODM), might involve in the
integration of the middle SNPs into the genomic sequence.

Further analysis of the edited products revealed multiple repaired products by the MJ.HPAT3-1
donor with various frequencies (Figure 1E). The precisely edited allele containing the intended
base modification accounted for 9.72 % of total edited reads (Supplemental Table 3). However,
when we observed all the edited products containing the desirable B and C base changes and
excluded the others containing one-sided base changes (A1 and A2 or D1 and D2) that were
designed to prohibit the recurrent cleavages of the SpCas9, the proportion of these alleles was
33.02% of the total edited alleles (Figure 1E; Supplemental Table 3). The diversity of the repaired
products indicates that not only was MMEJ involved in repairing the DNA damage, but it also
participated in other repair mechanisms like HR for one-sided alleles and ODM with only the B
and C base changes. The initial data showed that both cNHEJ and MMEJ-mediated precision gene
editing are feasible in tomato protoplasts, with the latter performing far better than the former.

Higher donor amounts enhance PREMJ efficiency

The copy number of donor DNA templates available at the targeting sites during the cleavage of
CRISPR/Cas complexes was shown to be essential for enhancing HDR efficiency (Cermak et al.
2015; Vu et al. 2020; Wang et al. 2017). We sought to investigate the implications of different
donor doses on the frequency of PREMJ. The PREMJ frequencies with all base changes were
increased with higher amounts of donor DNA, from 10.42 % (50 pmol) to 19.84 % (300 pmol),

and the one-sided repair frequencies were reduced accordingly (Figure 2A and Supplemental Figure 5; Supplemental Table 4). In the case of the cNJ.HPAT3-1 donor, the editing efficiency did not vary much among the donor doses and was much lower than that of the MJ.HPAT3-1 (Figure 2A). Again, most of its products contained only B and C changes (Supplemental Figure 5). These data indicate that the simultaneity of the MMEJ donor presence and the cleavages of the genomic sites was important for precise gene replacement since higher donor doses increased the chance of diffusing them to the being-cleaved targeted sites. The higher abundance of the donor templates might also ensure sufficient amounts of undamaged donor DNA templates for editing.

Figure 2. Optimization of MMEJ-mediated precision gene editing in tomato. (**A**) Editing efficiency revealed from targeted deep sequencing data of the various donor doses. (**B**) Editing efficiencies revealed from the treatments of donors with different microhomology lengths. Targeted deep sequencing data were collected in four replicates with no donor (-donor, Mock) control. (**C**) Editing efficiency revealed from targeted deep sequencing data of the treatments at different NU7441 concentrations. Targeted deep sequencing data were collected in three biological replicates. The data in (**B**) and (**C**) were analyzed by the Cas-Analyzer tool, statistically analyzed (multiple comparisons used a two-sided, uncorrected Fisher's LSD test), and plotted by GraphPad Prism 9. The p values are shown on the top of each compared pair. All the data points are shown on the plots. Targeted editing efficiency is the sum of the

290   precise editing efficiency and the only B and C editing efficiency. Precise editing efficiency was

291   calculated by dividing the total reads containing all the base changes by the total reads. The only B and C

292   editing efficiency was calculated by dividing the total reads containing only B and C base changes by the

293   total reads. One-sided editing efficiency was calculated by dividing total reads containing only either the

294   left-sided bases (A1 and A2) or the right-sided base pairs (D1 and D2) by the total reads.

The length of microhomology affects PREMJ performance

The length of microhomology arms was an important parameter for activating the MMEJ pathway upon DSB formation and end resection (Villarreal et al. 2012). Effective microhomology lengths (8-20 bases) increased MMEJ-mediated gene insertion in mammalian cells (Nakade et al. 2014; Sakuma et al. 2016; Ata et al. 2018). We, therefore, tested the impact of three different microhomology lengths (20, 10, and 5 bp) on PREMJ efficiency. When the microhomology was shorter than 20 bp, the total editing efficiency was significantly reduced from 4.22 ± 0.47 % (MJ.HPAT3-1) to 2.33 ± 0.31 % (MJ.HPAT3-2) and 2.65 ± 0.58 % (MJ.HPAT3-3) (Supplemental Table 6). More importantly, the all-base-changed precise editing efficiency was significantly higher for 20-bp microhomology (0.28 ± 0.05 %) compared to that of the 10-bp (0.12 ± 0.03 %) and 5-bp (0.04 ± 0.01 %) microhomology lengths (Figure 2B). Furthermore, when we consider the reads, including the B and C base changes, the targeted editing efficiency was 1.16 ± 0.09, 0.66 ± 0.12, and 0.55 ± 0.06 % for the 20-bp; 10-bp, and 5-bp microhomologies, respectively (Figure 2B; Supplemental Table 6). Interestingly, the activities of the gRNAs were not significantly different among all the treatments (Supplemental Table 6), indicating that the intermolecular MMEJ-mediated editing efficiency obtained from the experiments depended on microhomology length, as shown in the cases of intramolecular MMEJ (Ata et al. 2018; Bae et al. 2014; Nakade et al. 2014). These data suggest that the 20-bp microhomology was the best among the tested lengths, confirming the highest performance of the 20-bp microhomology (Sakuma et al. 2016) for MMEJ-mediated DSB repair in plants.

NU7441 improves PREMJ

Since the PREMJ efficiency was still low to be practically applicable for crop improvement, significant improvement of the editing system is needed. NU7441, a small chemical that was shown to inhibit DNA-dependent protein kinase (DNA-PKcs), an important cNHEJ component, enhancing HR-mediated repair of DSBs (Zhao et al. 2006; Robert et al. 2015; Vu et al. 2021a),

also significantly elevated the MMEJ-mediated DSB repair products in mammalian cells (Dutta et
al. 2017). We next tested if NU7441 can facilitate MMEJ repair using various concentrations of
the chemical and the MJ.HPAT3-1donors. When the NU7441 concentration was increased from 0
to 1 µM, the editing efficiency was significantly elevated (Table 1 and Figure 2C). The read
numbers of the precise products containing all the intended base changes, or only B and C base
changes, were enhanced with the targeted editing efficiency reached up to 1.60 ± 0.14 % at 1 µM
NU7441 compared to 1.03 ± 0.32 % without the addition of NU7441 (Table 1). The total editing
efficiency was significantly higher with 1 µM NU7441 treatment, reaching up to 5.96 ± 0.34 %, a
1.65-fold increase compared to mock control and majorly contributed by one-side editing (Table
1), indicating that NU7441 treatment effectively favored the choice of MMEJ pathway for DSB
repair. When NU7441 was added up to 2 µM, the repaired outcomes were reversely correlated
with the NU7441 concentrations (Figure 2C). This result suggests that over-suppression of the
cNHEJ components might lead to cellular toxicity, as shown in mammalian cells (Robert et al.
2015), and thus negatively impact the DSB repairs (Table 1 and Figure 2C).

Taken together, MMEJ-mediated precision editing is feasible in tomatoes and might be improved
by using sufficient microhomology and NU7441 treatment.

Efficient PREMJ-based gene editing in lettuce and cabbage protoplasts

To extend PREMJ-mediated precision gene editing to other plant species, we conducted PREMJ-
mediated precise gene replacement in lettuce and cabbage protoplasts using the ribonucleoprotein
(RNP) transfection method. The *THERMO-TOLERANCE 1* (*TT1*) (Li et al. 2015), *ORANGE* (*Or*)
(Yuan et al. 2015; Yazdani et al. 2019), and *ACETOLACTATE SYNTHASE 1* (*ALS1*) (Mazur et al.
1987) genes were selected as targets for both lettuce and cabbage (Supplemental Figure 6). All the
edited alleles of *TT1*, *Or*, and *ALS1* contained a single amino acid substitution (Supplemental
Figure 6) for obtaining the expected traits (heat tolerance, carotenoid overaccumulation, and
herbicide tolerance, respectively) (Li et al. 2015; Yuan et al. 2015; Yazdani et al. 2019; Mazur et
al. 1987). Sequence alignment using tomato protein sequences was conducted using NCBI Blastp
and putative *TT1*, *Or*, and *ALS1* genes were subsequently identified in lettuce (Supplemental
Tables 7) and cabbage (Supplemental Tables 8). We designed and employed two gRNAs for
cutting genomic loci and MMEJ donors containing 20-base microhomologies at two ends (Figure
3A and 3B). In lettuce, the highest precise PREMJ efficiency for all the intended base changes was

1.81 ± 0.75 % for the *LsALS1* locus, and the lowest efficiency was zero for the *LsTT1* locus. The

LsOr locus showed only 0.13 ± 0.10 % for exchanging all the intended bases (Figure 3C;

Supplemental Table 9). As mentioned earlier, the simultaneous activity of the two gRNAs is

important for PREMJ-mediated DSB repair. The data here suggest more similar cleavages between

the two gRNAs of *LsOr* and *LsALS1* compared to that of the *LsTT1* (Supplemental Table 9), which

may partially explain the inefficiency of PREMJ-mediated repair at that locus. Considering all the

repaired products that contain the targeted base changes for the desirable a.a., the efficiency

reached 4.47 ± 1.98 %, 0.78 ± 0.30 %, and 0.42 ± 0.28 % for *LsALS1*, *LsTT1*, and *LsOr* locus,

respectively (Fig 3C; Supplemental Table 9).

C

No.	Locus	Accession	Total reads	Targeted editing efficiency (%)	Precise editing efficiency (%)	gRNA1 indel rate (%)	gRNA2 indel rate (%)
1	LsTT1	XP_023737048	96871	0.78 ± 0.30	0.000	0.87 ± 0.71	0.35 ± 0.27
2	LsOr	XP_023743959	90601	0.42 ± 0.27	0.13 ± 0.10	1.07 ± 0.71	0.94 ± 0.39
3	LsALS1	XP_023747230	89394	4.47 ± 1.98	1.81 ± 0.75	2.69 ± 1.19	2.15 ± 0.99

D

No.	Locus	Accession	Total reads	Targeted editing efficiency (%)	Precise editing efficiency (%)	gRNA1 indel rate (%)	gRNA2 indel rate (%)
1	BoTT1	Bo2g095110	16905	7.27 ± 4.46	1.71 ± 0.48	4.83 ± 1.27	2.39 ± 1.20
2	BoOr	Bo9g014710	68028	3.94 ± 0.14	1.56 ± 0.65	24.37 ± 1.99	16.76 ± 6.12
3	BoALS1	Bo1g076910	90366	2.51 ± 1.19	0.73 ± 0.35	1.13 ± 0.40	10.61 ± 5.11

Figure 3. MMEJ-mediated precision gene editing in lettuce and cabbage. (A-B) Schematic diagrams

of targeted loci (*TT1*, *Or*, and *ALS1* with their GenBank accession number), targeted sequences, gRNA

binding sites, and MMEJ donors in lettuce (**A**) and cabbage (**B**). The targeted genes are shown from the

start codon (ATG) to stop codons. Exons are drawn as colored boxes (purple boxes with lettuce's loci and

pink boxes for cabbage), and the discontinuous line in each gene represents intron parts of the gene; the

gRNA binding sequences are highlighted in light blue color with PAMs (bold nucleotides). The expected

cutting sites (3-nt upstream of a PAM) are indicated with the black scissors; the intended base changes are

painted in red font; the targeted amino acid changes and corresponding triplexes are denoted by the

discontinuous red boxes, and the texts placed on their top. The diagrams are not drawn to their scales. **(C)**

MMEJ-mediated editing efficiency in lettuce at the *TT1*, *Or*, and *ALS1* loci. (**D**) MMEJ-mediated editing

efficiency in cabbage at the *TT1*, *Or*, and *ALS1* loci. Targeted deep sequencing data were collected in

three replicates and were analyzed by the Cas-Analyzer tool.

In cabbage protoplasts, the PREMJ worked more efficiently than for the tomato and lettuce

protoplasts. The highest precise editing efficiency (7.27 ± 4.46 %) was obtained at the *BoTT1* locus

and lower at the *BoOr* gene (1.68 ± 0.24 %). The *ALS1* locus for cabbage resulted in the least

precise editing efficiency, at only 0.73 ± 0.43 % (Figure 3D; Supplemental Table 10). Moreover,

the targeted editing efficiency at the *BoTT1* locus was the highest among tested loci in tomato,

lettuce, and cabbage, reaching up to 8.98 ± 4.73 % (Figure 3D). In summary, the data from lettuce

and cabbage indicate that PREMJ-mediated precision editing could be successfully extended to

other plant species.

Agrobacterium-**mediated delivery of PREMJ in tomatoes**

The plant regeneration system for tomato protoplast was shown to be of very low efficiency and

time-consuming. To overcome this challenge, we attempted to deliver the PREMJ tools into

tomato cotyledon explants by the *Agrobacterium*-mediated method (Vu et al. 2020). *SlHPAT3* and

SlHKT1;2 loci were selected as the targets for allele replacement (Supplemental Figure 7a,b and

Supplemental file 1). Unexpectedly, at the callus stage, the PREMJ efficiency was extremely low

at both the loci (Supplemental Table 11) and the editing reads were mostly with one-side editing

(Supplemental Figure 8a) or one-sided insertion of the MMEJ donor (Supplemental Figure 8b).

However, subsequent screening of transformants revealed plants carried precisely edited alleles,

albeit the editing frequency was relatively low (up to 4% for the cNHEJ event cNJ1 and 3% for

the PREMJ event #MJ5) (Supplemental Figure 9). The low efficiency of PREMJ might be

explained by the requirement of simultaneous cleavages of four sites located in the donors and

genomic sequences. In addition, the donor template amounts might also not be sufficient to be

distributed to the targeted sites immediately after DSB formations. Those hurdles are not subjected

to the protoplast transfection method.

Conclusions

We have developed and validated the PREMJ method in plant protoplasts. The PREMJ approach

has no limitation concerning the position and the number of base changes within the donor

sequence. Such gene replacement is helpful in short sequence replacement for engineering
regulatory elements, *in vivo* protein tagging, and short a.a. modification of a protein. Multiple-site
editing is also possible with the RNP-protoplast transfection, but further testing is required.
However, *in planta* transformation and regeneration efficiency using PREMJ tools should be
sufficiently improved for practical applications in plant breeding.

Data availability

All data generated or analyzed during this study are included in this published article and its
supplementary information files or it will be provided upon a reasonable request.

References

Anzalone AV, Koblan LW, Liu DR (2020) Genome editing with CRISPR-Cas nucleases, base
    editors, transposases and prime editors. Nat Biotechnol 38 (7):824-844.
    doi:10.1038/s41587-020-0561-9

Ata H, Ekstrom TL, Martinez-Galvez G, Mann CM, Dvornikov AV, Schaefbauer KJ, Ma AC,
    Dobbs D, Clark KJ, Ekker SC (2018) Robust activation of microhomology-mediated end
    joining for precision gene editing applications. PLoS Genet 14 (9):e1007652.
    doi:10.1371/journal.pgen.1007652

Bae S, Kweon J, Kim HS, Kim JS (2014) Microhomology-based choice of Cas9 nuclease target
    sites. Nat Methods 11 (7):705-706. doi:10.1038/nmeth.3015

Brooks C, Nekrasov V, Lippman ZB, Van Eck J (2014) Efficient gene editing in tomato in the first
    generation using the clustered regularly interspaced short palindromic repeats/CRISPR-
    associated9 system. Plant Physiol 166 (3):1292-1297. doi:10.1104/pp.114.247577

Cermak T, Baltes NJ, Cegan R, Zhang Y, Voytas DF (2015) High-frequency, precise modification
    of the tomato genome. Genome Biol 16:232. doi:10.1186/s13059-015-0796-9

Dutta A, Eckelmann B, Adhikari S, Ahmed KM, Sengupta S, Pandey A, Hegde PM, Tsai MS,
    Tainer JA, Weinfeld M, Hegde ML, Mitra S (2017) Microhomology-mediated end joining
    is activated in irradiated human cells due to phosphorylation-dependent formation of the
    XRCC1 repair complex. Nucleic Acids Res 45 (5):2585-2599. doi:10.1093/nar/gkw1262

Hayashi A, Tanaka K (2019) Short-Homology-Mediated CRISPR/Cas9-Based Method for
    Genome Editing in Fission Yeast. G3 (Bethesda) 9 (4):1153-1163.
    doi:10.1534/g3.118.200976

Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. Science 337 (6096):816-821. doi:10.1126/science.1225829

Li P, Zhang L, Li Z, Xu C, Du X, Wu S (2019) Cas12a mediates efficient and precise endogenous gene tagging via MITI: microhomology-dependent targeted integrations. Cell Mol Life Sci. doi:10.1007/s00018-019-03396-8

Li XM, Chao DY, Wu Y, Huang X, Chen K, Cui LG, Su L, Ye WW, Chen H, Chen HC, Dong NQ, Guo T, Shi M, Feng Q, Zhang P, Han B, Shan JX, Gao JP, Lin HX (2015) Natural alleles of a proteasome alpha2 subunit gene contribute to thermotolerance and adaptation of African rice. Nat Genet 47 (7):827-833. doi:10.1038/ng.3305

Lin Q, Jin S, Zong Y, Yu H, Zhu Z, Liu G, Kou L, Wang Y, Qiu J-L, Li J, Gao C (2021) High-efficiency prime editing with optimized, paired pegRNAs in plants. Nature Biotechnology 39 (8):923-927. doi:10.1038/s41587-021-00868-w

Lu Y, Tian Y, Shen R, Yao Q, Wang M, Chen M, Dong J, Zhang T, Li F, Lei M, Zhu JK (2020a) Targeted, efficient sequence insertion and replacement in rice. Nat Biotechnol 38 (12):1402-1407. doi:10.1038/s41587-020-0581-5

Lu Y, Tian Y, Shen R, Yao Q, Zhong D, Zhang X, Zhu JK (2020b) Precise genome modification in tomato using an improved prime editing system. Plant Biotechnol J. doi:10.1111/pbi.13497

Matsuzaki S, Sakuma T, Yamamoto T (2024) REMOVER-PITCh: microhomology-assisted long-range gene replacement with highly multiplexed CRISPR-Cas9. In Vitro Cellular & Developmental Biology - Animal 60 (7):697-707. doi:10.1007/s11626-024-00850-1

Mazur BJ, Chui CF, Smith JK (1987) Isolation and characterization of plant genes coding for acetolactate synthase, the target enzyme for two classes of herbicides. Plant Physiol 85 (4):1110-1117. doi:10.1104/pp.85.4.1110

Nakade S, Tsubota T, Sakane Y, Kume S, Sakamoto N, Obara M, Daimon T, Sezutsu H, Yamamoto T, Sakuma T, Suzuki KT (2014) Microhomology-mediated end-joining-dependent integration of donor DNA in cells and animals using TALENs and CRISPR/Cas9. Nat Commun 5:5560. doi:10.1038/ncomms6560

457   Park J, Lim K, Kim J-S, Bae S (2016) Cas-analyzer: an online tool for assessing genome editing
458   results using NGS data. Bioinformatics 33 (2):286-288.
459   doi:10.1093/bioinformatics/btw561 %J Bioinformatics

460   Puchta H (2005) The repair of double-strand breaks in plants: mechanisms and consequences for
461   genome evolution. J Exp Bot 56 (409):1-14. doi:10.1093/jxb/eri025

462   Robert F, Barbeau M, Ethier S, Dostie J, Pelletier J (2015) Pharmacological inhibition of DNA-
463   PK stimulates Cas9-mediated genome editing. Genome Med 7:93. doi:10.1186/s13073-
464   015-0215-6

465   Sakuma T, Nakade S, Sakane Y, Suzuki KT, Yamamoto T (2016) MMEJ-assisted gene knock-in
466   using TALENs and CRISPR-Cas9 with the PITCh systems. Nat Protoc 11 (1):118-133.
467   doi:10.1038/nprot.2015.140

468   Schindele A, Dorn A, Puchta H (2020) CRISPR/Cas brings plant biology and breeding into the
469   fast lane. Curr Opin Biotechnol 61:7-14. doi:10.1016/j.copbio.2019.08.006

470   Sharma S, Javadekar SM, Pandey M, Srivastava M, Kumari R, Raghavan SC (2015) Homology
471   and enzymatic requirements of microhomology-dependent alternative end joining. Cell
472   Death Dis 6:e1697. doi:10.1038/cddis.2015.58

473   Tan J, Zhao Y, Wang B, Hao Y, Wang Y, Li Y, Luo W, Zong W, Li G, Chen S, Ma K, Xie X,
474   Chen L, Liu YG, Zhu Q (2020) Efficient CRISPR/Cas9-based plant genomic fragment
475   deletions by microhomology-mediated end joining. Plant Biotechnol J.
476   doi:10.1111/pbi.13390

477   Villarreal DD, Lee K, Deem A, Shim EY, Malkova A, Lee SE (2012) Microhomology directs
478   diverse DNA break repair pathways and chromosomal translocations. PLoS Genet 8
479   (11):e1003026. doi:10.1371/journal.pgen.1003026

480   Vu TV, Doan DTH, Tran MT, Sung YW, Song YJ, Kim J-Y (2021a) Improvement of the
481   LbCas12a-crRNA System for Efficient Gene Targeting in Tomato. 12.
482   doi:10.3389/fpls.2021.722552

483   Vu TV, Nguyen NT, Kim J, Hong JC, Kim JY (2024a) Prime editing: Mechanism insight and
484   recent applications in plants. Plant Biotechnol J 22 (1):19-36. doi:10.1111/pbi.14188

485   Vu TV, Nguyen NT, Kim J, Song YJ, Nguyen TH, Kim JY (2024b) Optimized dicot prime editing
486   enables heritable desired edits in tomato and Arabidopsis. Nat Plants. doi:10.1038/s41477-
487   024-01786-w

Vu TV, Sivankalyani V, Kim E-J, Doan DTH, Tran MT, Kim J, Sung YW, Park M, Kang YJ, Kim J-Y (2020) Highly efficient homology-directed repair using CRISPR/Cpf1-geminiviral replicon in tomato. Plant Biotechnology Journal 18 (10):2133-2143. doi:https://doi.org/10.1111/pbi.13373

Vu TV, Thi Hai Doan D, Kim J, Sung YW, Thi Tran M, Song YJ, Das S, Kim J-Y (2021b) CRISPR/Cas-based precision genome editing via microhomology-mediated end joining. Plant Biotechnology Journal 19 (2):230-239. doi:https://doi.org/10.1111/pbi.13490

Wang M, Lu Y, Botella JR, Mao Y, Hua K, Zhu JK (2017) Gene Targeting by Homology-Directed Repair in Rice Using a Geminivirus-Based CRISPR/Cas9 System. Mol Plant 10 (7):1007-1010. doi:10.1016/j.molp.2017.03.002

Wolter F, Puchta H (2019) In planta gene targeting can be enhanced by the use of CRISPR/Cas12a. The Plant Journal 100 (5):1083-1094. doi:https://doi.org/10.1111/tpj.14488

Woo JW, Kim J, Kwon SI, Corvalan C, Cho SW, Kim H, Kim SG, Kim ST, Choe S, Kim JS (2015) DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. Nat Biotechnol 33 (11):1162-1164. doi:10.1038/nbt.3389

Yao X, Wang X, Liu J, Hu X, Shi L, Shen X, Ying W, Sun X, Wang X, Huang P, Yang H (2017) CRISPR/Cas9 - Mediated Precise Targeted Integration In Vivo Using a Double Cut Donor with Short Homology Arms. EBioMedicine 20:19-26. doi:10.1016/j.ebiom.2017.05.015

Yazdani M, Sun Z, Yuan H, Zeng S, Thannhauser TW, Vrebalov J, Ma Q, Xu Y, Fei Z, Van Eck J, Tian S, Tadmor Y, Giovannoni JJ, Li L (2019) Ectopic expression of ORANGE promotes carotenoid accumulation and fruit development in tomato. Plant Biotechnol J 17 (1):33-49. doi:10.1111/pbi.12945

Yuan H, Owsiany K, Sheeja TE, Zhou X, Rodriguez C, Li Y, Welsch R, Chayut N, Yang Y, Thannhauser TW, Parthasarathy MV, Xu Q, Deng X, Fei Z, Schaffer A, Katzir N, Burger J, Tadmor Y, Li L (2015) A Single Amino Acid Substitution in an ORANGE Protein Promotes Carotenoid Overaccumulation in Arabidopsis. Plant Physiol 169 (1):421-431. doi:10.1104/pp.15.00971

Zhang Z, Zeng W, Zhang W, Li J, Kong D, Zhang L, Wang R, Peng F, Kong Z, Ke Y, Zhang H, Kim C, Zhang H, Botella JR, Zhu J-K, Miki D (2022) Insights into the molecular mechanisms of CRISPR/Cas9-mediated gene targeting at multiple loci in Arabidopsis. Plant Physiology:kiac431. doi:10.1093/plphys/kiac431

Zhao Y, Thomas HD, Batey MA, Cowell IG, Richardson CJ, Griffin RJ, Calvert AH, Newell DR,
Smith GC, Curtin NJ (2006) Preclinical evaluation of a potent novel DNA-dependent
protein kinase inhibitor NU7441. Cancer Res 66 (10):5354-5362. doi:10.1158/0008-
5472.CAN-05-4275
Zong Y, Liu Y, Xue C, Li B, Li X, Wang Y, Li J, Liu G, Huang X, Cao X, Gao C (2022) An
engineered prime editor with enhanced editing efficiency in plants. Nature Biotechnology
40 (9):1394-1402. doi:10.1038/s41587-022-01254-w

Acknowledgments

527 Not applicable.

Funding

529 This work was supported by the National Research Foundation of Korea (the Bio & Medical
530 Technology Development Program 2020M3A9I4038352, 2020R1A6A1A03044344,
531 2021R1A5A8029490, 2022R1A2C3010331).

Author information

Authors and Affiliations

Division of Applied Life Science (BK21 FOUR Program), Plant Molecular Biology and
Biotechnology Research Center, Gyeongsang National University, Jinju 660-701, Republic of
Korea

Tien Van Vu, Ngan Thi Nguyen, Swati Das, Mil Thi Tran, Yeon Woo Sung, Jihae Kim, Young
Jong Song, Jae-Yean Kim

Department of Biological Sciences, Pusan National University, Busan 46241, Republic of Korea

Gah-Hyun Lim

Biological Resource Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB),
Jeongeup, 56212, Republic of Korea

Gah-Hyun Lim, Seung Hee Choi, Ju Yeon Moon, Suk Weon Kim, Mil Thi Tran, Jae Cheol Jeong

Nulla Bio Inc 501 Jinju-daero, Jinju 52828, Republic of Korea

Jae-Yean Kim

Contributions

Conceptualization, T.V.V and J.Y.K.; Methodology, T.V. V and J.Y.K.; Conducted experiments, T.V.V., G.H.L, S.H.C, J.Y.M., S.W.K., J.C.J, N.T.N, S.D., M.T.T., Y.W.S, J.K, Y.J.S; Data analysis, T.V.V. and J.Y.K.; Writing – Original Draft, T.V.V.; Writing – Review & Editing, T.V.V. and J.Y.K.; Funding Acquisition, T.V.V., and J.Y.K.; and Supervision, T.V.V and J.Y.K.

Corresponding authors

Correspondence to Tien Van Vu (tienvu.agi@gmail.com, ORCID: 0000-0002-6369-7664) and Jae-Yean Kim (kimjy@gnu.ac.kr, ORCID ID : 0000-0002-1180-6232).

Ethics declarations

Consent for publication

Not applicable.

Ethical approval

Not applicable.

Competing interests

Based on the results reported in this paper, the authors have applied for a Korean patent (10-2021-0089814) and a PCT patent (PCT-KR2021-008727). J.Y.K. is a founder and the CEO of Nulla Bio Inc.

SHORT LEGENDS FOR SUPPORTING INFORMATION

Supplemental figure legends

Supplemental Figure 1. HR-independent strategies for precision gene/allele replacement using CRISPR/SpCas9

Supplemental Figure 2. Sanger sequencing data revealed precise replacement of DNA by MMEJ donor

Supplemental Figure 3. Indel alleles revealed from transfection of SpCas9, gR1.HPAT1 and gR2.HPAT3 with cNJ.HPAT3-1 donor

Supplemental Figure 4. Indel alleles revealed from transfection of SpCas9, gR1.HPAT1 and
gR2.HPAT3 with MJ.HPAT3-1 donor

Supplemental Figure 5. The frequency of the MMEJ-mediated repaired products at different donor
doses

Supplemental Figure 6. Identification of *TT1*, *Or*, and *ALS1* genes in lettuce and cabbage for
MMEJ-mediated gene targeting

Supplemental Figure 7. Agrobacterium-mediated system for cNHEJ and MMEJ-mediated gene
editing in tomato.

Supplementary Figure 8. Representative repaired products obtained by the pMJ1 and pMJ2 vectors.

Supplemental Figure 9. cNHEJ and MMEJ-mediated precision editing events revealed from
Agrobacterium-mediated transformation.

Supplemental Tables

Supplemental Table 1. gRNA, donor, and primer sequences employed in the study.

Supplemental Table 2. cNHEJ- and MMEJ-mediated editing rates at *SlHPAT3* locus revealed by
Sanger sequencing.

Supplemental Table 3. cNHEJ- and MMEJ-mediated editing rates of various repaired products at
SlHPAT3 locus.

Supplemental Table 4. Editing frequency of various repaired products revealed from treatments of
different donor types and doses.

Supplemental Table 5. MMEJ-mediated gene editing using various MJ.HPAT3-1 donor amounts
(replicate).

Supplemental Table 6. MMEJ-mediated editing efficiency revealed from DSBs repair using
donors with different microhomology lengths.

Supplemental Table 7. Sequence alignment for selection of targeted genes in lettuce.

Supplemental Table 8. Sequence alignment for selection of targeted genes in cabbage.

Supplemental Table 9. Data revealed from the targeted deep-sequencing analysis of MMEJ-
mediated editing in lettuce.

Supplemental Table 10. Data revealed from the targeted deep-sequencing analysis of MMEJ-

mediated editing in cabbage.

Supplementary Table 11. PREMJ-mediated gene editing in tomato using the Agrobacterium-

mediated delivery of the editing tools.

Supplemental files

Supplemental file 1: Sequences used in the study

604 **Table 1. The detailed targeted deep sequencing data revealed from the various NU7441 concentrations**

Treatment	Replicate	NU7441 (uM)	Total reads	Total editing efficiency (%)	Precise editing efficiency (%)[a]	Targeted editing efficiency (%)[b]	Only B-C editing efficiency (%)[c]	One-sided editing efficiency (%)[d]	gRNA1 indel rate (%)	gRNA2 indel rate (%)
NU0	1	0.0	22605	3.67	0.23	0.86	0.63	2.81	3.36	4.20
	2	0.0	29443	2.18	0.09	0.58	0.49	1.59	3.05	2.97
	3	0.0	88974	4.97	0.32	1.65	1.33	3.32	0.58	0.69
Average		-		3.60	0.21	1.03	0.82	2.57	2.33	2.62
SEM		-		0.81	0.07	0.32	0.26	0.51	0.88	1.03
NU0.5	1	0.5	23931	5.39	0.24	1.35	1.11	4.04	4.76	4.71
	2	0.5	22605	4.14	0.23	1.13	0.91	3.00	3.36	3.40
	3	0.5	78738	5.70	0.42	1.86	1.44	3.84	0.53	0.55
Average		-		5.08	0.29	1.45	1.15	3.63	2.88	2.89
SEM		-		0.48	0.06	0.22	0.16	0.32	1.24	1.23
p-value (compared to NU0)				0.0740	0.0387	0.2280	0.2157	0.0544	0.7459	0.8778
NU1	1	1.0	28953	6.55	0.32	1.69	1.36	4.86	3.94	4.02
	2	1.0	30859	5.38	0.30	1.33	1.02	4.06	4.25	4.21
	3	1.0	79749	5.94	0.37	1.79	1.42	4.15	0.82	0.82
Average		-		5.96	0.33	1.60	1.27	4.35	3.00	3.02
SEM		-		0.34	0.02	0.14	0.13	0.25	1.10	1.10
p-value (compared to NU0)				0.0111	0.1666	0.1102	0.1101	0.0051	0.6938	0.8192
NU2	1	2.0	25789	3.95	0.13	0.90	0.76	3.05	3.63	3.70
	2	2.0	29652	4.44	0.24	1.14	0.90	3.30	6.03	5.91
	3	2.0	80326	4.53	0.32	1.54	1.22	3.00	1.23	1.18
Average		-		4.31	0.23	1.19	0.96	3.12	3.63	3.59
SEM		-		0.18	0.05	0.19	0.13	0.09	1.39	1.36
p-value (compared to NU0)				0.3565	0.8333	0.6226	0.5828	0.2789	0.4533	0.5769

605 a, Precise editing efficiency was calculated by dividing the total reads containing all the base changes by the total reads. b, Targeted editing efficiency is the sum of precise editing efficiency and only

606 B and C editing efficiency. c, The only B and C editing efficiency was calculated by dividing the total reads containing only B and C base changes by the total reads. d, One-sided editing efficiency

607 was calculated by dividing total reads containing only either the left-sided bases (A1 and A2) or the right-sided base pairs (D1 and D2) by the total reads. The total editing efficiency is the sum of the

608 MMEJ-mediated editing efficiency (a+b+c+d).

 FIGURE LEGEND

Figure 1. HR-independent precision editing approaches using CRISPR/SpCas9. (**A**) Model

of sequence design for cNHEJ-based precise gene replacement. (**B**) Model of sequence design for

MMEJ-bases precise gene replacement. The cut sites are arranged to avoid recurrent cutting.

Active PAM sites are shown in purple. Guide RNA sequences or traces of them in repaired

products are underlined. The red xxxxxxxx represents MH1, whereas the green yyyyyyy

represents MH2. The orange sequences denote donors, and the gray indicates the genomic

sequence being replaced. The red asterisks represent sequence modifications of interest. The letters

n, x, and y represent any nucleotide of A, T, G, or C, except that the sequence-making gRNA

spacers should not have more than 4 T in a row to avoid early transcription termination. To prevent

the recurrent cut, the PAM sequence corresponding to the replaced genomic sequence should be

changed. Usually, nGG/CCn is changed to nHG/CDn where H=A or C or T; D=A or G or T. One

or some bases within the core spacer sequence (i.e., ~12 base pairs upstream/downstream of

nGG/CCn) could also be changed. Such changes should not lead to changes in protein sequences

of the targeted gene. (**C**) Schematic diagrams of cNHEJ- and MMEJ-mediated gene editing of

tomato targeted locus *SlHPAT3* (with their GenBank accession number), targeted sequences,

gRNA binding sites, and donor sense sequences (cNJ.HPAT3-1, MJ.HPAT3-1, MJ.HPAT3-2, and

MJ.HPAT3-3). The targeted genes are shown from the start codon (ATG) to the stop codon (TAA),

exons are drawn as blue boxes, and the discontinuous lines in each gene represent intron parts of

the gene; the gRNA binding sequences at the genomic locus are highlighted in light blue color and

arrows with PAMs (bold nucleotides). The expected cutting sites (3-nt upstream of a PAM) are

indicated with the black scissors; the intended base changes are painted in the red font that is

denoted by the discontinuous red boxes and named A1, A2, B, C, D1, and D2 on the top of the

boxes. The cNHEJ donor dsDNA is synthesized with 5'-phosphorylated-ends, and the

phosphodiester bonds between the first and the second nucleotides of 5' ends were chemically

modified with phosphorothioate linkages. The two red, discontinuous crossing lines indicate the

MMEJ-mediated replacement sites of the MJ.HPAT3-1 donor. For the donors with shorter

microhomology, the recombination lengths are reduced accordingly. The diagrams are not drawn

to their scales. (**D**) cNHEJ and MMEJ-mediated editing efficiency at the SlHPAT3 locus. The

editing efficiency is calculated by dividing the total reads containing the intended bases by the

total reads without considering the indel mutation efficiency of the gRNAs. (**E**) Editing frequency of respective repaired products/alleles revealed from miniseq data. The edited base-changed sets/alleles revealed from miniseq data are classified according to the presence of the base changes in the edited reads. A total of 11 base-changed sets/alleles are recorded and plotted with their relative frequency among all the edited reads.

Figure 2. Optimization of MMEJ-mediated precision gene editing in tomato. (**A**) Editing efficiency revealed from targeted deep sequencing data of the various donor doses. (**B**) Editing efficiencies revealed from the treatments of donors with different microhomology lengths. Targeted deep sequencing data were collected in four replicates with no donor (-donor, Mock) control. (**C**) Editing efficiency revealed from targeted deep sequencing data of the treatments at different NU7441 concentrations. Targeted deep sequencing data were collected in three biological replicates. The data in (**B**) and (**C**) were analyzed by the Cas-Analyzer tool, statistically analyzed (multiple comparisons used a two-sided, uncorrected Fisher's LSD test), and plotted by GraphPad Prism 9. The p values are shown on the top of each compared pair. All the data points are shown on the plots. Targeted editing efficiency is the sum of the precise editing efficiency and the only B and C editing efficiency. Precise editing efficiency was calculated by dividing the total reads containing all the base changes by the total reads. The only B and C editing efficiency was calculated by dividing the total reads containing only B and C base changes by the total reads. One-sided editing efficiency was calculated by dividing total reads containing only either the left-sided bases (A1 and A2) or the right-sided base pairs (D1 and D2) by the total reads.

Figure 3. MMEJ-mediated precision gene editing in lettuce and cabbage. (**A-B**) Schematic diagrams of targeted loci (*TT1*, *Or*, and *ALS1* with their GenBank accession number), targeted sequences, gRNA binding sites, and MMEJ donors in lettuce (**A**) and cabbage (**B**). The targeted genes are shown from the start codon (ATG) to stop codons. Exons are drawn as colored boxes (purple boxes with lettuce's loci and pink boxes for cabbage), and the discontinuous line in each gene represents intron parts of the gene; the gRNA binding sequences are highlighted in light blue color with PAMs (bold nucleotides). The expected cutting sites (3-nt upstream of a PAM) are indicated with the black scissors; the intended base changes are painted in red font; the targeted amino acid changes and corresponding triplexes are denoted by the discontinuous red boxes, and the texts placed on their top. The diagrams are not drawn to their scales. (**C**) MMEJ-mediated

669   editing efficiency in lettuce at the *TT1*, *Or*, and *ALS1* loci. (**D**) MMEJ-mediated editing efficiency

670   in cabbage at the *TT1*, *Or*, and *ALS1* loci. Targeted deep sequencing data were collected in three

671   replicates and were analyzed by the Cas-Analyzer tool.

Figure 1

Figure 1. HR-independent precision editing approaches using CRISPR/SpCas9. (**A**) Model of sequence design for cNHEJ-based precise gene replacement. (**B**) Model of sequence design for MMEJ-bases precise gene replacement. The cut sites are arranged to avoid recurrent cutting. Active PAM sites are shown in purple. Guide RNA sequences or traces of them in repaired products are underlined. The red xxxxxxxx represents MH1, whereas the green yyyyyyyy represents MH2. The orange sequences denote donors, and the gray indicates the genomic sequence being replaced. The red asterisks represent sequence modifications of interest. The letters n, x, and y represent any nucleotide of A, T, G, or C, except that the sequence-making gRNA spacers should not have more than 4 T in a row to avoid early transcription termination. To prevent the recurrent cut, the PAM sequence corresponding to the replaced genomic sequence should be changed. Usually, nGG/CCn is changed to nHG/CDn where H=A or C or T; D=A or G or T. One or some bases within the core spacer sequence (i.e., ~12 base pairs upstream/downstream of nGG/CCn) could also be changed. Such changes should not lead to changes in protein sequences of the targeted gene. (**C**) Schematic diagrams of cNHEJ- and MMEJ-mediated gene editing of tomato targeted locus *SlHPAT3* (with their GenBank accession number), targeted sequences, gRNA binding sites, and donor sense sequences (cNJ.HPAT3-1, MJ.HPAT3-1, MJ.HPAT3-2, and MJ.HPAT3-3). The targeted genes are shown from the start codon (ATG) to the stop codon (TAA), exons are drawn as blue boxes, and the discontinuous lines in each gene represent intron parts of the gene; the gRNA binding sequences at the genomic locus are highlighted in light blue color and arrows with PAMs (bold nucleotides). The expected cutting sites (3-nt upstream of a PAM) are indicated with the black scissors; the intended base changes are painted in the red font that is denoted by the discontinuous red boxes and named A1, A2, B, C, D1, and D2 on the top of the boxes. The cNHEJ donor dsDNA is synthesized with 5'-phosphorylated-ends, and the phosphodiester bonds between the first and the second nucleotides of 5' ends were chemically modified with phosphorothioate linkages. The two red, discontinuous crossing lines indicate the MMEJ-mediated replacement sites of the MJ.HPAT3-1 donor. For the donors with shorter microhomology, the recombination lengths are reduced accordingly. The diagrams are not drawn to their scales. (**D**) cNHEJ and MMEJ-mediated editing efficiency at the SlHPAT3 locus. The editing efficiency is calculated by dividing the total reads containing the intended bases by the total reads without considering the indel mutation efficiency of the gRNAs. (**E**) Editing frequency of respective repaired products/alleles revealed from miniseq data. The edited base-changed sets/alleles revealed from miniseq data are classified according to the presence of the base changes in the edited reads. A total of 11 base-changed sets/alleles are recorded and plotted with their relative frequency among all the edited reads.

Figure 2. Optimization of MMEJ-mediated precision gene editing in tomato. (**A**) Editing efficiency revealed from targeted deep sequencing data of the various donor doses. (**B**) Editing efficiencies revealed from the treatments of donors with different microhomology lengths. Targeted deep sequencing data were collected in four replicates with no donor (-donor, Mock) control. (**C**) Editing efficiency revealed from targeted deep sequencing data of the treatments at different NU7441 concentrations. Targeted deep sequencing data were collected in three biological replicates. The data in (**B**) and (**C**) were analyzed by the Cas-Analyzer tool, statistically analyzed (multiple comparisons used a two-sided, uncorrected Fisher's LSD test), and plotted by GraphPad Prism 9. The p values are shown on the top of each compared pair. All the data points are shown on the plots. Targeted editing efficiency is the sum of the precise editing efficiency and the only B and C editing efficiency. Precise editing efficiency was calculated by dividing the total reads containing all the base changes by the total reads. The only B and C editing efficiency was calculated by dividing the total reads containing only B and C base changes by the total reads. One-sided editing efficiency was calculated by dividing total reads containing only either the left-sided bases (A1 and A2) or the right-sided base pairs (D1 and D2) by the total reads.

Figure 3

C

No.	Locus	Accession	Total reads	Targeted editing efficiency (%)	Precise editing efficiency (%)	gRNA1 indel rate (%)	gRNA2 indel rate (%)
1	LsTT1	XP_023737048	96871	0.78 ± 0.30	0.00	0.86 ± 0.71	0.35 ± 0.27
2	LsOr	XP_023743959	90601	0.42 ± 0.28	0.13 ± 0.10	1.07 ± 0.71	0.94 ± 0.39
3	LsALS1	XP_023747230	89394	4.47 ± 1.98	1.81 ± 0.75	2.69 ± 1.19	2.15 ± 0.99

D

No.	Locus	Accession	Total reads	Targeted editing efficiency (%)	Precise editing efficiency (%)	gRNA1 indel rate (%)	gRNA2 indel rate (%)
1	BoTT1	Bo2g095110	16905	8.98 ± 4.73	7.27 ± 4.46	4.83 ± 1.27	2.39 ± 1.20
2	BoOr	Bo9g014710	68028	2.95 ± 1.10	1.68 ± 0.24	24.37 ± 1.99	16.76 ± 6.12
3	BoALS1	Bo1g076910	90366	3.24 ± 1.54	0.73 ± 0.43	1.13 ± 0.49	10.61 ± 6.25

Figure 3. MMEJ-mediated precision gene editing in lettuce and cabbage. (A-B) Schematic diagrams of targeted loci (*TT1*, *Or*, and *ALS1* with their GenBank accession number), targeted sequences, gRNA binding sites, and MMEJ donors in lettuce (**A**) and cabbage (**B**). The targeted genes are shown from the start codon (ATG) to stop codons. Exons are drawn as colored boxes (purple boxes with lettuce's loci and pink boxes for cabbage), and the discontinuous line in each gene represents intron parts of the gene; the gRNA binding sequences are highlighted in light blue color with PAMs (bold nucleotides). The expected cutting sites (3-nt upstream of a PAM) are indicated with the black scissors; the intended base changes are painted in red font; the targeted amino acid changes and corresponding triplexes are denoted by the discontinuous red boxes, and the texts placed on their top. The diagrams are not drawn to their scales. **(C)** MMEJ-mediated editing efficiency in lettuce at the *TT1*, *Or*, and *ALS1* loci. **(D)** MMEJ-mediated editing efficiency in cabbage at the *TT1*, *Or*, and *ALS1* loci. Targeted deep sequencing data were collected in three replicates and were analyzed by the Cas-Analyzer tool.

Supplemental information

CRISPR-Cas-based Microhomology-Mediated End Joining for Precise Gene Replacement in Plants

Tien Van Vu[1,†], Gah-Hyun Lim[2,3], Seung Hee Choi[3], Ju Yeon Moon[3], Ngan Thi Nguyen[1], Swati Das[1], Mil Thi Tran[1,3], Yeon Woo Sung[1], Jihae Kim[1], Young Jong Song[1], Suk Weon Kim[3], Jae Cheol Jeong[3], Jae-Yean Kim[1,4,†]

[1]Division of Applied Life Science (BK21 FOUR Program), Plant Molecular Biology and Biotechnology Research Center, Gyeongsang National University, Jinju 660-701, Republic of Korea.

[2]Department of Biological Sciences, Pusan National University, Busan 46241, Republic of Korea

[3]Biological Resource Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Jeongeup, 56212, Republic of Korea

[4]Nulla Bio Inc 501 Jinju-daero, Jinju 52828, Republic of Korea.

[†]Correspondence: Tien Van Vu: tienvu.agi@gmail.com (ORCID: 0000-0002-6369-7664). Jae-Yean Kim: kimjy@gnu.ac.kr (ORCID ID : 0000-0002-1180-6232).

Running title: MMEJ-mediated precise gene replacement in plants.

Supplemental Figure 1. HR-independent strategies for precision gene/allele replacement using CRISPR/SpCas9. (**a**) cNHEJ-mediated DNA replacement model. (**b**) MMEJ-mediated DNA replacement model. MH1 and MH2 are predefined from genomic loci and are subsequently introduced into the two ends of the donor sequence. CRISPR/Cas complexes are designed with canonical PAM binding sites (nGG/CCn) to produce DSBs at genomic loci with MH1 and MH2 flanking ends. MMEJ-mediated repair using the donor sequence generates genomic loci with precise modifications. The cut sites are arranged to avoid recurrent cutting. The orange sequences denote donors, and the gray indicates the genomic sequence being replaced. The letters n represent any nucleotide of A, T, G, or C except that the sequence-making gRNA spacers should not have more than 4 T in a row to avoid early transcription termination.

Supplemental Figure 2. Sanger sequencing data revealed precise replacement of DNA by MMEJ donor. PCR amplification of the targeted site was conducted using primers flanking the site and PCR products were resolved on gel and purified for Sanger sequencing. Sanger sequencing data were decomposed using the ICE Synthego web application using gR1.HPAT3 (upper panel) and gR2.HPAT3 (lower panel) for locating the cutting site. The edited allele and its percentage among the total sequence were revealed and highlighted in the discontinuous red boxes. The indented base changes of the left and right sides of the targeted sequence are highlighted in the orange background of the up and down panels, respectively.

Miniseq read	Product length	Read number	Edited type
gRNA1 ✂ **gRNA2** ✂			
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	30461	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGAT---AGTGTGGTATGTAATATGTATACATC	64	23884	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGAT--GTGTGGTATGTAATATGTATACATC	63	16471	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCG--AGTGTGGTATGTAATATGTATACATC	57	2854	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGAT--AGTGTGGTATGTAATATGTATACATC	59	2619	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTG---AGTGTGGTATGTAATATGTATACATC	62	2249	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATT--GTGTGGTATGTAATATGTATACATC	59	1361	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTT---AGTGTGGTATGTAATATGTATACATC	61	1280	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGAT---GTGTGGTATGTAATATGTATACATC	58	1279	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATT---AGTGTGGTATGTAATATGTATACATC	60	1229	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCG---AAGTGTGGTATGTAATATGTATACATC	58	876	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTG--AAGTGTGGTATGTAATATGTATACATC	63	850	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTG--GTGTGGTATGTAATATGTATACATC	55	632	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGC--AGTGTGGTATGTAATATGTATACATC	56	487	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	131	364	Ins
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGA-----GATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	125	354	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGA---TGTGTGGTATGTAATATGTATACATC	56	270	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGAT----TAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	126	211	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTG--GTGTGGTATGTAATATGTATACATC	61	200	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATTAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	131	200	Ins
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTT---GAAAGTGTGGTATGTAATATGTATACATC	64	148	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCC----AGTGTGGTATGTAATATGTATACATC	126	129	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGA---TGTGTGGTATGTAATATGTATACATC	61	129	del
AGCTTGCTTACGGAAAAATTGGAGAG--AGTGTGGTATGTAATATGTATACATC	52	117	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGAT---TGTGTGGTATGTAATATGTATACATC	64	114	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATT--TGTGTGGTATGTAATATGTATACATC	58	113	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATT---TAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	127	107	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGAT---TGTGTGGTATGTAATATGTATACATC	62	105	del
AGCTTGCTTACGGAAAAATTGGAG---AGTGTGGTATGTAATATGTATACATC	50	96	del
AGCTTGCTTACGGAAAAATTGG--AGTGTGGTATGTAATATGTATACATC	48	92	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTG--AGGTGTGGTATGTAATATGTATACATC	63	88	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	88	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	84	WT
GGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	81	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAG-AAGTGTGGTATGTAATATGTATACATC	129	75	del

Supplemental Figure 3. Indel alleles revealed from transfection of SpCas9, gR1.HPAT1 and gR2.HPAT3 with cNJ.HPAT3-1 donor. The gRNA binding and cutting sites are shown on the top panel with the scissor icons on the WT sequence.

Miniseq read	Product length	Read number	Edited type
gRNA1 ✂ ... gRNA2 ✂			
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	44318	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAG--AGTGTGGTATGTAATATGTATACATC	64	17053	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGAT---GTGTGGTATGTAATATGTATACATC	63	11634	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGAT--AGTGTGGTATGTAATATGTATACATC	59	2190	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCG--AGTGTGGTATGTAATATGTATACATC	57	2159	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTG---AGTGTGGTATGTAATATGTATACATC	62	2060	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGAT---GTGTGGTATGTAATATGTATACATC	58	1317	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATT---AGTGTGGTATGTAATATGTATACATC	60	1039	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTT--AGTGTGGTATGTAATATGTATACATC	61	909	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCG---AAGTGTGGTATGTAATATGTATACATC	58	877	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATT---GTGTGGTATGTAATATGTATACATC	59	813	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAAGATCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	722	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTT--GTGTGGTATGTAATATGTATACATC	60	586	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	541	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGC---AGTGTGGTATGTAATATGTATACATC	56	505	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCTGAGAGTGTGGTATGTAATATGTATACATC	130	429	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTG--AGTGTGGTATGTAATATGTATACATC	63	414	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCTGAGAGTGTGGTATGTAATATGTATACATC	130	405	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	131	396	Ins
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	365	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGG---GTGTGGTATGTAATATGTATACATC	56	348	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGC---GTGTGGTATGTAATATGTATACATC	55	335	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCTGAGAGTGTGGTATGTAATATGTATACATC	130	277	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGC-----GATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	125	245	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCTGAGAGTGTGGTATGTAATATGTATACATC	130	220	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	197	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATTAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	131	185	Ins
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGAT--TGTGTGGTATGTAATATGTATACATC	62	184	del
AGCTTGCTTACGG--AAGTGTGGTATGTAATATGTATACATC	40	181	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	130	179	WT
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGAT----TAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATC	126	178	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTT--GAAAGTGTGGTATGTAATATGTATACATC	64	170	del
AGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCC----AGTGTGGTATGTAATATGTATACATC	50	135	del
AGCTTGCTTACGGAAAAATTGGAGAGTGG--AGTGTGGTATGTAATATGTATACATC	126	128	del
AGCTTGCTTACGGAAAAATTGGAGAGTGG--AGTGTGGTATGTAATATGTATACATC	55	123	del

Supplemental Figure 4. Indel alleles revealed from transfection of SpCas9, gR1.HPAT1 and gR2.HPAT3 with MJ.HPAT3-1 donor. The gRNA binding and cutting sites are shown on the top panel with the scissor icons on the WT sequence.

Supplemental Figure 5. The frequency of the MMEJ-mediated repaired products at different donor doses

Supplemental Figure 6. Identification of *TT1*, *Or*, and *ALS1* genes in lettuce and cabbage for MMEJ-mediated gene targeting. **a**-**c**, Protein sequence alignment among tomato, lettuce, and cabbage for identification of targeted a.a changes in *TT1* (**a**), *Or* (**b**), and *ALS1* (**b**) genes. The targeted a.a is highlighted by the blue discontinuous boxes.

Supplemental Figure 7. Agrobacterium-mediated system for cNHEJ and MMEJ-mediated gene editing in tomato. a-b, Schematic diagrams of cNHEJ- and MMEJ-mediated gene editing of tomato targeted loci (*SlHPAT3* (**a**) and *SlHKT1;2* (**b**) with their GenBank accession number), targeted sequences, gRNA binding sites, and donors. The targeted genes are shown from the start codon (ATG) to stop codons. Exons are drawn as blue boxes and the discontinuous lines in each gene represent intron parts of the gene; the gRNA binding sequences are highlighted in light blue color at the genomic sites and purple color at the donor vectors with PAMs (bold nucleotides). The expected cutting sites (3-nt upstream of a PAM) are indicated with the black scissors; the intended base changes are painted in the red font; the targeted amino acid changes and corresponding triplexes are denoted by the discontinuous red boxes and the texts placed above them. Two to three silent base changes corresponding to either the PAM or seed sequences (within 9 bases upstream of the 5'-NGG-3' PAMs) were introduced into the donor to prevent the cleavage of the donor DNAs after they were successfully integrated into the genomic sites. The diagrams are not drawn to scale. **c-d,** Binary vectors used for tomato transformation to edit the *SlHPAT3* gene (**c**) and *SlHKT1;2* (**d**). A plant codon-optimized SpCas9 was employed to make the DSBs at the genomic sites and to release the donors. Also, a neomycin phosphotransferase II expression cassette was used for plant selection. The *SlHPAT3* gene was tested with control vectors (pNJC1 and pMJC1) without the donor cutting gRNAs. The pNJ1 and pMJ1 vectors are conventional T-DNAs and the other entire vectors are designed with a geminiviral replicon-based system that contains a Replicative protein expression cassette (Rep) driven by a bidirectional promoter (red bar, named as LIR) and terminated by a bidirectional terminator (blue bar, named as SIR). The replicon requires two LIRs to be autonomously replicated within the nuclei of plant cells.

a

Sequence	Length	Count	Type	HDR
TGTTCAGAGTGAGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACCTCCACTTAAAAATTTATCCC |||||||||||||||||||||||||||||||||||||||||||||||||·||·||||||||||||||||||||||||||||||||||||||| TGTTCAGAGTGAGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATAAAAGATCCTATATCAAAGGACCTCCACTTAAAAATTTATCCC TGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATCTCTCCATAAATCTTCAATAAGCATGGT ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||| TGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATCTCTCCATAAATCTTCAATAAGCATGGT	168	1	WT or Sub	O

b

Sequence	Length	Count	Type	HDR
TGTTCAGAGTGAGCTTGCTTACGGAAAAA--------------------TTGGAGAGTGGCGATTTGATAAGAGGTCCTATATCAAAGGACC ||||||||||||||||||||||||||| |||||||||||||||||||||||·||·|||||||||||||||| TGTTCAGAGTGAGCTTGCTTACGGAAAAATTGGAGAGTGGCGATTTGATTTGGAGAGTGGCGATTTGATAAAAGATCCTATATCAAAGGACC TCCACTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATCTCTCCATAAATCTTCAATAAGCA |||·|||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||| TCCTCTTAAAAATTTATCCCTGCCTCCACCTAACGTCCCAGAAAGTGTGGTATGTAATATGTATACATCTCTCCATAAATCTTCAATAAGCA TGGT |||| TGGT	188	1	Ins	O

Supplementary Fig. 8. Representative repaired products obtained by the pMJ1 and pMJ2 vector. a, A one-sided precise replacement of the A1 and A2 bases by the pMJ1. **b**, A repaired product of the pMJ2 containing one-sided precise replacement of the A1 and A2 bases and insertion of the 20-bp 5'-microhomology arm.

a

Event	Indel mutation efficiency (%)	Precise editing efficiency (%)
cNJ1	12	4
cNJ2	0	0
cNJ3	0	0
MJ1	20	0
MJ2	79	0
MJ3	0	0
MJ4	31	1
MJ5	39	3
MJ6	42	0
MJ7	23	0
MJ9	64	0
MJ10	80	0

Supplemental Figure 9. cNHEJ and MMEJ-mediated precision editing events revealed from *Agrobacterium*-mediated transformation. **a**. Table showing precisely edited allele frequency from regenerated plants transformed with the cNHEJ and MMEJ editing tools for editing the *SlHPAT3* locus. **b-d**. Edited alleles reported by ICE Synthego analysis with the cNHEJ event cNJ1 (**b**); the MMEJ events MJ4 (**c**) and MJ5 (**d**). The precisely edited alleles and efficiencies are denoted by the framed orange boxes named HDR that also highlighted modified bases in orange. The WT allele is marked with the plus sign.

SUPPLEMENTAL TABLES

Supplemental Table 1. gRNA, donor, and primer sequences employed in the study

No.	Name	Sequence (5'-3')	Note
\multicolumn	**gRNA and donor sequences**		
1	gR1.HPAT3	GAGAGTGGCGATTTGATAAG	gRNA1
2	gR2.HPAT3	ACCTAACGTCCCAGAAAGTG	gRNA2
3	gDR1.HPAT3	TTGATATAGGATCTTTTGCG	Donor gRNA1
4	gDR2.HPAT3	ACCTAACGTCCCTGAGAGCG	Donor gRNA2
5	gDR3.HPAT3	AAATCGCCACTCTCCAAGCG	Donor gRNA3
6	gDR4.HPAT3	TGGTATGTAATATGTATGCG	Donor gRNA4
7	cNJ.HPAT3-1	AAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCTGAGA	cNJ.HPAT3-1 donor
8	MJ.HPAT3-1	TTGGAGAGTGGCGATTTGATAAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCTGAGAGTGTGGTATGTAATATGTAT	MJ.HPAT3-1 donor
9	MJ.HPAT3-2	GCGATTTGATAAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCTGAGAGTGTGGTATG	MJ.HPAT3-2 donor
10	MJ.HPAT3-3	TTGATAAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCTTCCTCCACCTAACGTCCCTGAGAGTGTG	MJ.HPAT3-3 donor
11	gR1.LsTT1	GAGTTCTAGTCAGGAAGAGC	gRNA1
12	gR2.LsTT1	ACAGCCCTAAAAGTGTGAAT	gRNA2
13	MJ.LsTT1	CTAGAGTTCTAGTCAGGAAGAGTAGAAAGCAAGCAGAACAATACCATCATCTATACAAAGTACGCTTATTGTCCAACTCACACTTTTAGGGCTGTTTCT	MJ.LsTT1 donor
14	gR1.LsOr	AAGATAATATTAGAAGTCGT	gRNA1
15	gR2.LsOr	GAGGAGGTGAGCAAGTTTTT	gRNA2
16	MJ.LsOr	TTCAAGATAATATTAGAAGTCATAGAAATAAAATATTTTTACACATGGAGGAGGTGAGCAAATCTTTCGGAATATTCATCTGTA	MJ.LsOr donor
17	gR1.LsALS1	CCATCACCGGCCAAGTCCCC	gRNA1
18	gR2.LsALS1	ACGCGTTACCTCCACAATCG	gRNA2
19	MJ.LsALS1	TTGCCATCACCGGCCAAGTCTCTCGTAGAATGATCGGAACCGATGCGTTTCAAGAAACTCCAATTGTGGAGGTAACGCGTTCC	MJ.LsALS1 donor
20	gR1.BoTT1	GAGTTCTTGTGCGGAAGAGT	gRNA1
21	gR2.BoTT1	TTAAGATTACTTTGTACAAG	gRNA2
22	MJ.BoTT1	TCCGAGTTCTTGTGCGGAAGAGCAGAAAGCAGGCTGAGCAATACCTCCATCTGTACAAAGTAATCTTAAATA	MJ.BoTT1 donor
23	gR1.BoOr	AGAAGGACCTGAAACAGTAC	gRNA1
24	gR2.BoOr	GCAAGAAGATCTTGTTTCGA	gRNA2
25	MJ.BoOr	CATAGAAGGACCTGAAACAGTTCAAGACTTTGCCAAAATGCAATTACAAGAGATTCAAGACAACATTAGAAGTCACCGAAACAAGATCTTCTTGCACA	MJ.BoOr donor
26	gR1.BoALS1	TCTTGTCGCCATTACAGGAC	gRNA1
27	gR2.BoALS1	AACGATTGGTGTCTCTTGGA	gRNA2

28	MJ.BoALS1	TCCTCTTGTCGCCATTACAGGTCAAGTCTCTCGCCGGATGATCGG TACTGACGCTTTTCAAGAGACACCAATCGTTGAG	MJ.BoALS1 donor
Primers for donor preparation			
29	cNJ.HPAT-F1	AAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTATCCCT	Primers for preparing cNJ.HPAT3.1 donor. The phosphodiester linkages between the red nucleotides were modified with phosphorothioate bond
30	cNJ.HPAT-R1	TCTCAGGGACGTTAGGTGGAGGAAGGGATAAATTTTTAAGAGG AGG	
31	MJ.HPAT-F1	TTGGAGAGTGGCGATTTGATAAAAGATCCTATATCAAAGGACCT CCTCTTAAAAATTTATCCCT	Primers for preparing MJ.HPAT3.1 donor
32	MJ.HPAT-R1	ATACATATTACATACCACACTCTCAGGGACGTTAGGTGGAGGAA GGGATAAATTTTTAAGAGGAGG	
33	MJ.HPAT-F2	GCGATTTGATAAAAGATCCTATATCAAAGGACCTCCTCTTAAAA ATTTATCCCT	Primers for preparing MJ.HPAT3.2 donor
34	MJ.HPAT-R2	CATACCACACTCTCAGGGACGTTAGGTGGAGGAAGGGATAAAT TTTTAAGAGGAGG	
35	MJ.HPAT-F3	TTGATAAAAGATCCTATATCAAAGGACCTCCTCTTAAAAATTTAT CCCT	Primers for preparing MJ.HPAT3.3 donor
36	MJ.HPAT-R3	CACACTCTCAGGGACGTTAGGTGGAGGAAGGGATAAATTTTTA AGAGGAGG	
37	MJ.BoTT1-F1	TCCGAGTTCTTGTGCGGAAGAGCAGAAAGCAGGCTGAGCAATA CCTCC	Primers for preparing MJ.BoTT1 donor
38	MJ.BoTT1-R1	TATTTAAGATTACTTTGTACAGATGGAGGTATTGCTCAGCCTGCT TTC	
39	MJ.LsTT1-F1	CTAGAGTTCTAGTCAGGAAGAGTAGAAAGCAAGCAGAACAATA CCATCATCTATACAAAGTACGC	Primers for preparing MJ.LsTT1 donor
40	MJ.LsTT1-R1	AGAAACAGCCCTAAAAGTGTGAGTTGGACAATAAGCGTACTTT GTATAGATGATGGTATTGT	
41	MJ.BoOr-F1	CATAGAAGGACCTGAAACAGTTCAAGACTTTGCCAAAATGCAAT TACAAGAGATTCAAGACA	Primers for preparing MJ.BoOr donor
42	MJ.BoOr-R1	TGTGCAAGAAGATCTTGTTTCGGTGACTTCTAATGTTGTCTTGAA TCTCTTGTAATTGCATT	
43	MJ.LsOr-F1	TTCAAGATAATATTAGAAGTCATAGAAATAAAATATTTTTACACA TGGAGGAGGTG	Primers for preparing MJ.LsOr donor
44	MJ.LsOr-R1	TACAGATGAATATTCCGAAAGATTTGCTCACCTCCTCCATGTGTA AAAATATTT	
45	MJ.BoALS1-F1	TCCTCTTGTCGCCATTACAGGTCAAGTCTCTCGCCGGATGATCGG TACTGAC	Primers for preparing MJ.BoALS1 donor
46	MJ.BoALS1-R1	CTCAACGATTGGTGTCTCTTGAAAAGCGTCAGTACCGATCATCC GGCGAGAG	
47	MJ.LsALS1-F1	TTGCCATCACCGGCCAAGTCTCTCGTAGAATGATCGGAACCGAT GCGTTTCAAG	Primers for preparing MJ.LsALS1 donor
48	MJ.LsALS1-R1	GGAACGCGTTACCTCCACAATTGGAGTTTCTTGAAACGCATCGG TTCCGATCATT	
Primers for miniseq sample preparation			
49	BoTT1-F1	GCTCCTTTGCTCACTTAGATCA	First PCR

50	BoTT1-R1	GCAAATTGTGTATTGAAGTGAATTATGATC	
51	BoTT1-F2	ACACTCTTTCCCTACACGACGCTCTTCCGATCTGCTTCTCTGTAGTGGTATGGGCCC	Second PCR
52	BoTT1-R2	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTACAATACTCCAAGAAAAGTAGATGAG	
53	BoOr-F1	GTTCGTAATAGAGTGTGGTGGT	First PCR
54	BoOr-R1	AGAGGATAATCCCAGCAATGAG	
55	BoOr-F2	ACACTCTTTCCCTACACGACGCTCTTCCGATCTTGATTATTTTTGGTGCAGCTTTTGTATC	Second PCR
56	BoOr-R2	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTCCACTTTAAGAAAGTAAACCTCTTCC	
57	BoALS1-F1	CCCTCTACCCATTTCCAGATTC	First PCR
58	BoALS1-R1	CTATGCTCCACAGCGTAGTTAG	
59	BoALS1-F2	ACACTCTTTCCCTACACGACGCTCTTCCGATCTGTTAGCAGACGCGATGCTTGACA	Second PCR
60	BoALS1-R2	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGGTATGTCATCAACATCCATCACCAA	
61	LsTT1-F1	TCAGATTGAACATGCCCTCAC	First PCR
62	LsTT1-R1	GCCTTACACCACTACACAAAGA	
63	LsTT1-F2	ACACTCTTTCCCTACACGACGCTCTTCCGATCTTCATTTCCAGTGGCATGGGGCC	Second PCR
64	LsTT1-R2	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGTCCAAATTAAGTGCTTAACCCATTAAG	
65	LsOr-F1	AAGGAGGGCGACTGATAATTG	First PCR
66	LsOr-R1	CTTACTGTAGGTGCAAGGAGAC	
67	LsOr-F2	ACACTCTTTCCCTACACGACGCTCTTCCGATCTAACAGTCCAGGACTTTGATAAAATGG	Second PCR
68	LsOr-R2	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGCATAACGCAATTAGATAACTCACGC	
69	LsALS1-F1	CCTTCTATCACCCACCCATTTC	First PCR
70	LsALS1-R1	TCCGAGCATTTGAAGGGATAAA	
71	LsALS1-F2	ACACTCTTTCCCTACACGACGCTCTTCCGATCTACGCGCTGCTCGACAGTATCCC	Second PCR
72	LsALS1-R2	GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTACATCCAAAACAAGGTAATTGTGTTTAG	
Primers for Sanger sequencing			
73	HPAT3-sF1	CACCTTGGGATTTGGACAGTAG	SlHPAT3

Supplemental Table 2. cNHEJ- and MMEJ-mediated editing rates at *SlHPAT3* locus revealed by Sanger sequencing

Treatment	SpCas9	gR1.HPAT3	gR2.HPAT3	Donor	Precise editing efficiency* (%)	Targeted editing efficiency** (%)
1	-	-	-	-	0.0	0.0
2	+	+	-	-	0.0	0.0
3	+	-	+	-	0.0	0.0
4	+	+	+	-	0.0	0.0
5	+	+	+	cNHEJ.HPAT3-1	0.0	1.0
6	+	+	+	MMEJ.HPAT3-1	2.0	3.0

Precise editing efficiency was calculated by dividing total reads containing all the base changes by the total reads.
**Targeted-editing efficiency is the portion of the precisely edited products containing all the base changes and those containing only B and C base changes among all the edited products.

Supplemental Table 3. cNHEJ- and MMEJ-mediated editing rates of various repaired products at *SlHPAT3* locus

Edited SNPs	Donor			
	cNJ.HPAT3-1		MJ.HPAT3-1	
	Number of edited reads	Relative editing frequency (%)	Number of edited reads	Relative editing frequency (%)
A1-A2-B-C-D1-D2	0	0.00	237	9.72
A1-A2	0	0.00	872	35.77
A1-A2-B-C	0	0.00	150	6.15
B-C	72	94.74	169	6.93
B-C-D1-D2	4	5.26	249	10.21
D1-D2	0	0.00	761	31.21
Number of edited reads	76	100.000	2438	100.00
Total reads	93054	-	99885	-

Supplemental Table 4. Editing frequency of various repaired products revealed from treatments of different donor types and doses

Edited alleles	Mock		cNJ.HPAT3.1 donor amount (pmol)									MJ.HPAT3.1 donor amount (pmol)							
	0		50		100		200		300		50		100		200		300			
	Number of reads	Frequency (%)	Number of reads	Frequency (%)	Number of reads	Frequency (%)	Number of reads	Frequency (%)	Number of reads	Frequency (%)	Number of reads	Frequency (%)	Number of reads	Frequency (%)	Number of reads	Frequency (%)	Number of reads	Frequency (%)
A1-A2	0	0.00	0	0.00	0	0.00	0	0.00	2	0.42	719	39.66	699	37.85	522	33.42	461	32.56
D1-D2	0	0.00	1	0.17	8	1.48	8	1.43	5	1.05	525	28.96	534	28.91	358	22.92	278	19.63
A1-A2-B-C-D1-D2	0	0.00	0	0.00	0	0.00	1	0.18	0	0.00	189	10.42	185	10.02	238	15.24	281	19.84
A1-A2-B-C	0	0.00	0	0.00	0	0.00	1	0.18	1	0.21	65	3.59	68	3.68	65	4.16	75	5.30
B-C	0	0.00	601	99.67	533	98.34	541	96.43	459	96.63	122	6.73	94	5.09	107	6.85	72	5.08
B-C-D1-D2	0	0.00	1	0.17	1	0.18	10	1.78	8	1.68	193	10.65	267	14.46	272	17.41	249	17.58
Total reads with edited bases	0	0.000	603	100.00	542	100.00	561	100.00	475	100.00	1813	100.00	1847	100.00	1562	100.00	1416	100.00
Total read	27391	-	31995	-	36917	-	33600	-	27420	-	29937	-	27998	-	28611	-	28296	-
Total editing efficiency (%)*	0.00		1.88		1.47		1.67		1.73		6.06		6.60		5.46		5.00	
Targeted editing ratio (%)**	0.00		99.83		98.52		98.57		98.53		31.38		33.24		43.66		47.81	

Total editing efficiency is a sum of all the MMEJ-mediated editing efficiency ((Total reads with edited bases/Total read) x 100).

*** Targeted-editing ratio is the portion of the precisely edited products containing all the base changes and those containing only B and C base changes among all the edited products.*

Supplemental Table 5. MMEJ-mediated gene editing using various MJ.HPAT3-1 donor amounts (replicate)

Treatment	Donor amount (pmol)	Total reads	Precise editing efficiency* (%)	Only B and C editing efficiency (%)**	gRNA1 indel rate (%)	gRNA2 indel rate (%)
1	300	23848	0.105	0.587	3.329	3.350
2	200	29643	0.078	0.523	3.077	3.080
3	100	26579	0.090	0.764	3.525	3.424
4	50	25108	0.008	0.119	2.880	2.824

* Precise editing efficiency was calculated by dividing total reads containing all the base changes by the total reads.

 ** The only B and C editing efficiency were calculated by dividing total reads containing the only B and C base changes by the total reads

Supplemental Table 6. MMEJ-mediated editing efficiency revealed from DSBs repair using donors with different microhomology lengths.

Donor	Replicate	Total reads	Total editing efficiency (%)**	Targeted editing efficiency (%)	Precise editing efficiency (%)	Only B and C editing efficiency (%)	One-sided editing efficiency (%)	gRNA1 indel rate (%)	gRNA2 indel rate (%)
-	1	76598	0.00	0.00	0.00	0.00	0.00	0.62	0.62
-	2	76002	0.00	0.00	0.00	0.00	0.00	1.39	1.36
Average		-	0.00	0.00	0.00	0.00	0.00	1.01	0.99
SEM		-	0.00	0.00	0.00	0.00	0.00	0.38	0.37
MJ.HPAT3-1	1	74776	4.62	1.12	0.32	0.80	3.50	0.73	0.73
MJ.HPAT3-1	2	72689	5.32	1.40	0.40	1.00	3.92	0.69	0.68
MJ.HPAT3-1	3	28491	3.80	1.12	0.21	0.92	2.67	2.23	2.18
MJ.HPAT3-1	4	20879	3.16	1.01	0.19	0.81	2.16	4.26	4.26
Average		-	4.22	1.16	0.28	0.88	3.06	1.98	1.96
SEM		-	0.47	0.09	0.05	0.05	0.40	0.84	0.84
p (MJ.HPAT3-1 vs MJ.HPAT3-2)*		-	0.0064	0.0007	0.0025	0.0024	0.0016	0.9310	0.9419
p (MJ.HPAT3-1 vs MJ.HPAT3-3)		-	0.0179	0.0002	<0.0001	0.0015	<0.0001	0.5456	0.5619
MJ.HPAT3-2	1	75083	1.76	0.42	0.10	0.32	1.34	0.18	0.19
MJ.HPAT3-2	2	67743	3.01	0.76	0.19	0.57	2.25	0.34	0.40
MJ.HPAT3-2	3	25992	2.72	0.94	0.15	0.79	1.92	4.21	4.19
MJ.HPAT3-2	4	28214	1.85	0.52	0.06	0.46	1.33	2.71	2.68
Average		-	2.33	0.66	0.12	0.54	1.71	1.86	1.86
SEM		-	0.31	0.12	0.03	0.10	0.23	0.97	0.96
p (MJ.HPAT3-2 vs MJ.HPAT3-3)		-	0.5903	0.3657	0.0731	0.7876	0.0654	0.6539	0.6544
MJ.HPAT3-3	1	65392	1.91	0.62	0.06	0.56	1.29	0.39	0.38
MJ.HPAT3-3	2	68109	1.44	0.41	0.06	0.35	1.03	0.32	0.35
MJ.HPAT3-3	3	28571	3.37	0.49	0.02	0.47	0.72	2.50	2.47
MJ.HPAT3-3	4	26584	3.89	0.70	0.04	0.66	1.09	2.09	2.14
Average		-	2.65	0.55	0.04	0.51	1.03	1.32	1.33
SEM		-	0.58	0.06	0.01	0.07	0.12	0.57	0.57

*p-value returned from Uncorrected Fisher's LSD test. ** Total editing efficiency is the sum of all the MMEJ-mediated editing efficiency without considering the indel mutation efficiency of the gRNAs. Targeted editing efficiency is the sum of the precise editing efficiency and the only B and C editing efficiency. Precise editing efficiency was calculated by dividing total reads containing all the base changes by the total reads. The only B and C editing efficiency were calculated by dividing total reads containing the only B and C base changes by the total reads. One-sided editing efficiency was calculated by dividing total reads containing either the left-sided bases (A1 and A2) or the right-sided base pairs (D1 and D2) by the total reads

Supplemental Table 7. Sequence alignment for selection of targeted genes in lettuce

Gene ID	Identity (%)	Alignment length (a.a)	E-value	Bitscore
TT1				
XP_023737048.1	93.617	235	3.60E-164	453
XP_023733423.1	38.767	227	1.51E-50	165
XP_023758433.1	35.526	228	5.80E-47	156
XP_023734413.1	36.325	234	2.22E-46	154
XP_023770770.1	35.043	234	2.55E-45	152
Or				
XP_023743959.1	85.657	462	3.03E-165	453
PLY66076.1	85.657	462	3.22E-163	165
XP_023734734.1	68.775	362	1.56E-125	156
ALS1				
XP_023747230.1	83.043	659	0	1068
PLY79534.1	61.836	249	1.14E-82	262
PLY79513.1	65.574	652	5.42E-52	176
XP_023754650.1	25.664	630	3.49E-40	155
XP_023750264.1	23.897	578	1.03E-15	81.3

Supplemental Table 8. Sequence alignment for selection of targeted genes in cabbage

Gene ID	Identity (%)	Alignment Length (a.a)	E-value	Bitscore
TT1				
Bo2g095110	94.1	236	7.4e-120	433.0
Bo6g081130	93.6	236	1.6e-119	431.8
Bo6g123660	93.2	236	1.6e-119	431.8
Or				
Bo9g014710	74.4	312	2.0e-104	382.1
Bo2g007140	68.3	249	3.3e-75	285.0
ALS1				
Bo1g076910	74.4	666	6.0e-244	846.7
Bo7g082530	68.3	638	4.6e-143	511.5

Supplemental Table 9. Data revealed from the targeted deep-sequencing analysis of MMEJ-mediated editing in lettuce

No.	Targeted gene	Replicate	Total reads	Total editing efficiency (%)*	Targeted editing efficiency (%)**	Precise editing efficiency (%)***	Desirable base change efficiency (%)****	One-sided editing efficiency (%)*****	gRNA1 indel rate (%)	gRNA2 indel rate (%)
1	*LsTT1*	1	31767	2.62	0.81	0.00	0.81	1.81	2.26	0.88
2	*LsTT1*	2	33468	4.44	1.29	0.00	1.29	3.15	0.31	0.13
3	*LsTT1*	3	31636	0.82	0.26	0.00	0.26	0.56	0.02	0.03
Average			96871	2.63	0.78	0.00	0.78	1.84	0.86	0.35
SEM******			590	1.05	0.30	0.00	0.30	0.75	0.71	0.27
4	*LsOr*	1	29362	0.48	0.09	0.02	0.07	0.39	2.48	1.73
5	*LsOr*	2	29362	0.82	0.20	0.04	0.15	0.62	0.32	0.50
6	*LsOr*	3	31877	3.94	0.97	0.33	0.64	2.97	0.40	0.60
Average			90601	1.74	0.42	0.13	0.29	1.33	1.07	0.94
SEM			838.33	1.10	0.28	0.10	0.18	0.82	0.71	0.39
7	*LsALS1*	1	25553	2.74	1.53	0.44	1.09	1.22	0.97	0.41
8	*LsALS1*	2	26629	13.62	8.23	3.03	5.19	5.39	2.13	2.20
9	*LsALS1*	3	37212	5.37	3.65	1.96	1.68	1.72	4.96	3.83
Average			89394	7.24	4.47	1.81	2.66	2.78	2.68	2.15
SEM			3720	3.28	1.98	0.75	1.28	1.31	1.19	0.99

*Total editing efficiency is a sum of all the MMEJ-mediated editing efficiency without considering the indel mutation efficiency of the gRNAs. ** Targeted editing efficiency is the sum of the precise editing efficiency and the desirable base change efficiency. ***Precise editing efficiency was calculated by dividing total reads containing all the base changes by the total reads. ****The desirable base change efficiency was calculated by dividing total reads containing only the targeted a.a changes by the total reads. *****One-sided editing efficiency was calculated by dividing total edited reads not containing targeted a.a change by the total reads. ******SEM: Standard error of the mean.*

Supplemental Table 10. Data revealed from the targeted deep-sequencing analysis of MMEJ-mediated editing in cabbage

No.	Targeted gene	Replicate	Total reads	Total editing efficiency (%)*	Targeted editing efficiency (%)**	Precise editing efficiency (%)***	Desirable base change efficiency (%)****	One-sided editing efficiency (%)*****	gRNA1 indel rate (%)	gRNA2 indel rate (%)
1	*BoTT1*	1	6647	1.64	1.37	0.57	0.8	0.27	2.72	0.56
2	*BoTT1*	2	3719	20.25	17.64	15.73	1.91	2.61	7.13	4.65
3	*BoTT1*	3	6539	10.25	7.94	5.51	2.43	2.31	4.65	1.97
Average			16905	10.71	8.98	7.27	1.71	1.73	4.83	2.39
SEM******			-	5.38	4.73	4.46	0.48	0.73	1.27	1.2
4	*BoOr*	1	17556	2.49	1.95	1.38	0.57	0.54	21.94	9.27
5	*BoOr*	2	28622	5.69	3.95	1.98	1.97	1.75	26.8	24.26
6	*BoOr*	3	21850	0.28	0.14	0.01	0.13	0.14	11.73	6.58
Average			68028	4.09	2.95	1.68	1.27	1.15	24.37	16.76
SEM			-	1.31	1.10	0.24	0.57	0.49	1.99	6.12
7	*BoALS1*	1	30301	0.38	0.22	0.04	0.18	0.16	0.43	6.7
8	*BoALS1*	2	38778	7.93	5.32	1.19	4.13	2.61	1.16	4.39
9	*BoALS1*	3	21287	6.2	4.17	0.96	3.21	2.02	1.81	20.74
Average			90366	4.84	3.24	0.73	2.51	1.6	1.13	10.61
SEM			-	2.8	1.54	0.43	1.46	0.9	0.49	6.25

*Total editing efficiency is a sum of all the MMEJ-mediated editing efficiency without considering the indel mutation efficiency of the gRNAs. ** Targeted editing efficiency is the sum of the precise editing efficiency and the desirable base change efficiency. ***Precise editing efficiency was calculated by dividing total reads containing all the base changes by the total reads. ****The desirable base change efficiency was calculated by dividing total reads containing only the targeted a.a changes by the total reads. *****One-sided editing efficiency was calculated by dividing total edited reads not containing targeted a.a change by the total reads. ******SEM: Standard error of the mean.*

Supplementary Table 11. PREMJ-mediated gene editing in tomato using the *Agrobacterium*-mediated delivery of the editing tools

Vector	Donor	NU7441 (µM)	Rep 1			Rep 2		
			Precise editing rate (%)	gRNA1 indel rate (%),	gRNA3 indel rate (%),	Precise editing rate (%)	gRNA1 indel rate (%),	gRNA3 indel rate (%),
pNJC1	cNJ.HPAT3-1	0	0.000	14.727	14.579	0.000	12.296	12.006
pMJC1	MJ.HPAT3-1	0	0.000	16.150	16.008	0.000	8.772	8.439
pMJC1	MJ.HPAT3-1	1	0.000	36.014	35.346	0.000	16.115	15.502
pNJ1	cNJ.HPAT3-1	0	0.000	7.137	7.122	0.003	5.024	5.005
pNJ2	cNJ.HPAT3-1	0	0.000	5.279	5.209	0.003	5.798	5.814
pMJ1	MJ.HPAT3-1	0	0.000	5.100	5.252	0.000	3.819	3.946
pMJ1	MJ.HPAT3-1	1	0.004	5.943	5.790	0.000	4.654	4.647
pMJ2	MJ.HPAT3-1	0	0.000	7.807	7.714	0.000	9.917	9.832
pMJ2	MJ.HPAT3-1	1	0.000	8.384	8.169	0.003	7.778	7.775
pNJ3	cNJ.HKT1;2	0	0.024	0.013	0.238	0.000	N/A*	N/A
pMJ3	MJ.HKT1;2	0	0.000	0.018	0.399	0.000	N/A	N/A
pMJ3	MJ.HKT1;2	1	0.009	0.015	0.425	0.000	N/A	N/A

N/A: not available

SEQUENCES USED IN THE STUDY

❖ **HPAT3 cNHEJ donor (cNJ.HPAT3-1) for RNP works**

5'-AAaAGaTCCTATATCAAAGGACCTCCtCTTAAAAATTTATCCCTtCCTCCACCTAACGTCCCtGAgA-3'

Blue font: introduced bases. From 5'-3': the bases were name as A1; A2; B; C; D1; D2, respectively.

❖ **HPAT3 MMEJ donor (MJ.HPAT3-1) for RNP works**

5'-
ATTGGAGAGTGGCGATTTGATAAaAGaTCCTATATCAAAGGACCTCCtCTTAAAAATTTATCCCTtCCTCCACCT
AACGTCCCtGAgAGTGTGGTATGTAATATGTAT-3'

black font: replaced sequence; Blue font: introduced bases. From 5'-3': the bases were name as A1; A2; B; C; D1; D2, respectively; red font: MH1; green font: MH2

❖ **HPAT3 MMEJ donor (MJ.HPAT3-2) for RNP works**

5'-
GGCGATTTGATAAaAGaTCCTATATCAAAGGACCTCCtCTTAAAAATTTATCCCTtCCTCCACCTAACGTCCCtGA
gAGTGTGGTATG-3'

black font: replaced sequence; Blue font: introduced bases. From 5'-3': the bases were name as A1; A2; B; C; D1; D2, respectively; red font: MH1; green font: MH2

❖ **HPAT3 MMEJ donor (MJ.HPAT3-3) for RNP works**

5'-
TTGATAAaAGaTCCTATATCAAAGGACCTCCtCTTAAAAATTTATCCCTtCCTCCACCTAACGTCCCtGAgAGTGT
G-3'

black font: replaced sequence; Blue font: introduced bases. From 5'-3': the bases were name as A1; A2; B; C; D1; D2, respectively; red font: MH1; green font: MH2

❖ **HPAT3 cNHEJ donor (cNJ.HPAT3-1) for *Agrobacterium*-mediated transformation works**

CCgcgcAAaAGaTCCTATATCAAAGGACCTCCtCTTAAAAATTTATCCCTtCCTCCACCTAACGTCCCtGAgAgcgtG
G

Blue font: introduced bases. From 5'-3': the bases were name as A1; A2; B; C; D1; D2, respectively; yellow highlighted font: gDR1.HPAT3 binding sequence; gray highlighted font: gDR2.HPAT3 binding sequence.

❖ **HPAT3 MMEJ donor (MJ.HPAT3-1) for *Agrobacterium*-mediated transformation works**

CCgcgcTTGGAGAGTGGCGATTTGATAAaAGaTCCTATATCAAAGGACCTCCtCTTAAAAATTTATCCCTtCCTCC
ACCTAACGTCCCtGAgAGTGTGGTATGTAATATGTATgcgtGG

Blue font: introduced bases. From 5'-3': the bases were name as A1; A2; B; C; D1; D2, respectively; yellow highlighted font: gDR3.HPAT3 binding sequence; gray highlighted font: gDR4.HPAT3 binding sequence. red font: MH1; green font: MH2

❖ **sgR1.HPAT3**

GAGAGTGGCGATTTGATAAGgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtg
gcaccgagtcggtgc

Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

- **sgR2.HPAT3**

  ACCTAACGTCCCAGAAAGTGgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtgg
  caccgagtcggtgc

  Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

- **gDR1.HPAT3**

  TTGATATAGGAtCTtTTgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagtc
  ggtgcTTTTTTT

  Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

- **gDR2.HPAT3**

  ACCTAACGTCCCtGAgAgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagt

  cggtgcTTTTTTT

  Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

- **gDR3.HPAT3**

  AAATCGCCACTCTCCAAgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagt

  cggtgcTTTTTTT

  Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

- **gDR4.HPAT3**

  TGGTATGTAATATGTATgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgag
  tcggtgcTTTTTTT

  Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

- **HKT1;2 cNHEJ donor (cNJ.HKT12) for *Agrobacterium*-mediated transformation works**

  CCgcgcAcGTt<mark>GTTTTTGGATAT</mark>ATTCTTGTTGTTATTCTTCTTGGTTCATCGTTAGTTTCTCTCTATATAATAATCA
  TCCCTAGCGCCAAACAAATCCTTGACCAAAAAGGCCTTAATTTACATACTTTTTCACTATTCACCACAGTATCAA
  CTTTTGCAgATTGTGGTTTTTTACCTACAAATGAAAACATGATGATTTTCAAGAAAAATTCAGGaCTTCTTCTCA
  TTCTT<mark style="background:#c0c0c0">ATCCCTCAAGTC</mark>tTgCTgcgtGG

  Blue font: introduced bases; <mark>yellow highlighted font: gDR1.HKT12 binding sequence;</mark> <mark style="background:#c0c0c0">gray
  highlighted font: gDR2.HKT12 binding sequence</mark>.

- **HKT1;2 MMEJ donor (MJ.HKT12) for *Agrobacterium*-mediated transformation works**

  CCgcgc<mark>ATCGATAAAATTGTTGAGTA</mark>AcGTtGTTTTTGGATATATTCTTGTTGTTATTCTTCTTGGTTCATCGTTAG
  TTTCTCTCTATATAATAATCATCCCTAGCGCCAAACAAATCCTTGACCAAAAAGGCCTTAATTTACATACTTTTT
  CACTATTCACCACAGTATCAACTTTTGCAgATTGTGGTTTTTTACCTACAAATGAAAACATGATGATTTTCAAGA
  AAAATTCAGGaCTTCTTCTCATTCTTATCCCTCAAGTCtTgCTAGG<mark style="background:#c0c0c0">GAACACTTTGTTTGCTC</mark>gcgtGG
  Blue font: introduced bases. <mark>yellow highlighted font: gDR3.HKT12 binding sequence;</mark> <mark style="background:#c0c0c0">gray
  highlighted font: gDR4.HKT12 binding sequence</mark>. red font: MH1; green font: MH2

- **sgR1.HKT12**

GATAAAATTGTTGAGTAATGgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccga
gtcggtgcTTTTTTT

Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

❖ **sgR2.HKT12**

CAAACAAAGTGTTCCCTAGAgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccga
gtcggtgcTTTTTTT

Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

❖ **gDR1.HKT12**

ATATCCAAAAACaACgTgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagt
cggtgcTTTTTTT

Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

❖ **gDR2.HKT12**

ATCCCTCAAGTCtTgCTgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagtc
ggtgcTTTTTTT

Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

❖ **gDR3.HKT12**

TCAACAATTTTATCGATgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagt
cggtgcTTTTTTT

Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

❖ **gDR4.<u>H</u>KT12**

GAACACTTTGTTTGCTCgcggttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagt
cggtgcTTTTT

Black font: gRNA sequence; green font: CRISPR gRNA scaffold.

❖ p35S-pcoCas9I-t35S sequence

GGAGGAATTCCAATCCCACAAAAATCTGAGCTTAACAGCACAGTTGCTCCTCTCAGAGCAGAATCGGGTATTCAAC
ACCCTCATATCAACTACTACGTTGTGTATAACGGTCCACATGCCGGTATATACGATGACTGGGGTTGTACAAAGGC
GGCAACAAACGGCGTTCCCGGAGTTGCACACAAGAAATTTGCCACTATTACAGAGGCAAGAGCAGCAGCTGACGC
GTACACAACAAGTCAGCAAACAGACAGGTTGAACTTCATCCCCAAAGGAGAAGCTCAACTCAAGCCCAAGAGCTT
TGCTAAGGCCCTAACAAGCCCACCAAAGCAAAAAGCCCACTGGCTCACGCTAGGAACCAAAAGGCCCAGCAGTGA
TCCAGCCCCAAAAGAGATCTCCTTTGCCCCGGAGATTACAATGGACGATTTCCTCTATCTTTACGATCTAGGAAGG
AAGTTCGAAGGTGAAGGTGACGACACTATGTTCACCACTGATAATGAGAAGGTTAGCCTCTTCAATTTCAGAAAG
AATGCTGACCCACAGATGGTTAGAGAGGCCTACGCAGCAAGTCTCATCAAGACGATCTACCCGAGTAACAATCTCC
AGGAGATCAAATACCTTCCCAAGAAGGTTAAAGATGCAGTCAAAAGATTCAGGACTAATTGCATCAAGAACACAG
AGAAAGACATATTTCTCAAGATCAGAAGTACTATTCCAGTATGGACGATTCAAGGCTTGCTTCATAAACCAAGGCA

AGTAATAGAGATTGGAGTCTCTAAAAAGGTAGTTCCTACTGAATCTAAGGCCATGCATGGAGTCTAAGATTCAAAT
CGAGGATCTAACAGAACTCGCCGTCAAGACTGGCGAACAGTTCATACAGAGTCTTTTACGACTCAATGACAAGAA
GAAAATCTTCGTCAACATGGTGGAGCACGACACTCTGGTCTACTCCAAAAATGTCAAAGATACAGTCTCAGAAGAT
CAAAGGGCTATTGAGACTTTTCAACAAAGGATAATTTCGGGAAACCTCCTCGGATTCCATTGCCCAGCTATCTGTC
ACTTCATCGAAAGGACAGTAGAAAAGGAAGGTGGCTCCTACAAATGCCATCATTGCGATAAAGGAAAGGCTATCA
TTCAAGATCTCTCTGCCGACAGTGGTCCCAAAGATGGACCCCCACCCACGAGGAGCATCGTGGAAAAGAAGAGG
TTCCAACCACGTCTACAAAGCAAGTGGATTGATGTGACATCTCCACTGACGTAAGGGATGACGCACAATCCCACTA
TCCTTCGCAAGACCCTTCCTCTATATAAGGAAGTTCATTTCATTTGGAGAGGACACGCTCGAGTATAAGGTAAATTT
CTGTGTTCCTTATTCTCTCAAAATCTTCGATTTTGTTTTCGTTCGATCCCAATTTCGTATATGTTCTTTGGTTTAGATTC
TGTTAATCTTAGATCGAAGATGATTTTCTGGGTTTGATCGTTAGATATCATCTTAATTCTCGATTAGGGTTTCATAG
ATATCATCCGATTTGTTCAAATAATTTGAGTTTTGTCGAATAATTACTCTTCGATTTGTGATTTCTATCTAGATCTGG
TGTTAGTTTCTAGTTTGTGCGATCGAATTTGTCGATTAATCTGAGTTTTTCTGATTAACAGGAGCTCATTTTTACAAC
AATTACCAACAACAACAAACAACAAACAACATTACAATTACATTTACAATTATCGATACAATGacc
ATGAAACGGACAGCCGACGGAAGCGAGTTCGAGGCCCCAAAGAAGAAGCGGAAGGTCGATAAGAAGTACTCTAT
CGGACTCGATATCGGAACTAACTCTGTGGGATGGGCTGTGATCACCGATGAGTACAAGGtaaagcctcgattttggggttt
aggtgtctgcttattagagtaaaaacacatcctttgaaattgtttgtggtcatttgattgtgctcttgatccattgaattgctGCAGGTGCCATCTAA
GAAGTTCAAGGTTCTCGGAAACACCGATAGGCACTCTATCAAGAAAAACCTTATCGGTGCTCTCCTCTTCGATTCTG
GTGAAACTGCTGAGGCTACCAGACTCAAGAGAACCGCTAGAAGAAGGTACACCAGAAGAAAGAACAGGATCTGC
TACCTCCAAGAGATCTTCTCTAACGAGATGGCTAAAGTGGATGATTCATTCTTCCACAGGCTCGAAGAGTCATTCCT
CGTGGAAGAAGATAAGAAGCACGAGAGGCACCCTATCTTCGGAAACATCGTTGATGAGGTGGCATACCACGAGA
AGTACCCTACTATCTACCACCTCAGAAAGAAGCTCGTTGATTCTACTGATAAGGCTGATCTCAGGCTCATCTACCTC
GCTCTCGCTCACATGATCAAGTTCAGAGGACACTTCCTCATCGAGGGTGATCTCAACCCTGATAACTCTGATGTGG
ATAAGTTGTTCATCCAGCTCGTGCAGACCTACAACCAGCTTTTCGAAGAGAACCCTATCAACGCTTCAGGTGTGGA
TGCTAAGGCTATCCTCTCTGCTAGGCTCTCTAAGTCAAGAAGGCTTGAGAACCTCATTGCTCAGCTCCCTGGTGAG
AAGAAGAACGGACTTTTCGGAAACTTGATCGCTCTCTCTCTCGGACTCACCCCTAACTTCAAGTCTAACTTCGATCT
CGCTGAGGATGCAAAGCTCCAGCTCTCAAAGGATACCTACGATGATGATCTCGATAACCTCCTCGCTCAGATCGGA
GATCAGTACGCTGATTTGTTCCTCGCTGCTAAGAACCTCTCTGATGCTATCCTCCTCAGTGATATCCTCAGAGTGAA
CACCGAGATCACCAAGGCTCCACTCTCAGCTTCTATGATCAAGAGATACGATGAGCACCACCAGGATCTCACACTT
CTCAAGGCTCTTGTTAGACAGCAGCTCCCAGAGAAGTACAAAGAGATTTTCTTCGATCAGTCTAAGAACGGATACG
CTGGTTACATCGATGGTGGTGCATCTCAAGAAGAGTTCTACAAGTTCATCAAGCCTATCCTCGAGAAGATGGATGG
AACCGAGGAACTCCTCGTGAAGCTCAATAGAGAGGATCTTCTCAGAAAGCAGAGGACCTTCGATAACGGATCTAT
CCCTCATCAGATCCACCTCGGAGAGTTGCACGCTATCCTTAGAAGGCAAGAGGATTTCTACCCATTCCTCAAGGAT
AACAGGGAAAAGATTGAGAAGATTCTCACCTTCAGAATCCCTTACTACGTGGGACCTCTCGCTAGAGGAAACTCAA
GATTCGCTTGGATGACCAGAAAGTCTGAGGAAACCATCACCCCTTGGAACTTCGAAGAGGTGGTGGATAAGGGT
GCTAGTGCTCAGTCTTTCATCGAGAGGATGACCAACTTCGATAAGAACCTTCCAAACGAGAAGGTGCTCCCTAAGC
ACTCTTTGCTCTACGAGTACTTCACCGTGTACAACGAGTTGACCAAGGTTAAGTACGTGACCGAGGGAATGAGGA
AGCCTGCTTTTTTGTCAGGTGAGCAAAAGAAGGCTATCGTTGATCTCTTGTTCAAGACCAACAGAAAGGTGACCGT
GAAGCAGCTCAAAGAGGATTACTTCAAGAAATCGAGTGCTTCGATTCAGTTGAGATTTCTGGTGTTGAGGATAG
GTTCAACGCATCTCTCGGAACCTACCACGATCTCCTCAAGATCATTAAGGATAAGGATTTCTTGGATAACGAGGAA
AACGAGGATATCTTGGAGGATATCGTTCTTACCCTCACCCTCTTTGAAGATAGAGAGATGATTGAAGAAAGGCTCA
AGACCTACGCTCATCTCTTCGATGATAAGGTGATGAAGCAGTTGAAGAGAAGAAGATACACTGGTTGGGGGAAGG
CTCTCAAGAAAGCTCATTAACGGAATCAGGGATAAGCAGTCTGGAAAGACAATCCTTGATTTCCTCAAGTCTGATG

GATTCGCTAACAGAAACTTCATGCAGCTCATCCACGATGATTCTCTCACCTTTAAAGAGGATATCCAGAAGGCTCA
GGTTTCAGGACAGGGTGATAGTCTCCATGAGCATATCGCTAACCTCGCTGGATCTCCTGCAATCAAGAAGGGAAT
CCTCCAGACTGTGAAGGTTGTGGATGAGTTGGTGAAGGTGATGGGAAGGCATAAGCCTGAGAACATCGTGATCG
AAATGGCTAGAGAGAACCAGACCACTCAGAAGGGACAGAAGAACTCTAGGGAAAGGATGAAGAGGATCGAGGA
AGGTATCAAAGAGCTTGGATCTCAGATCCTCAAAGAGCACCCTGTTGAGAACACTCAGCTCCAGAATGAGAAGCT
CTACCTCTACTACCTCCAGAACGGAAGGGATATGTATGTGGATCAAGAGTTGGATATCAACAGGCTCTCTGATTAC
GATGTTGATCATATCGTGCCACAGTCATTCTTGAAGGATGATTCTATCGATAACAAGGTGCTCACCAGGTCTGATA
AGAACAGGGGTAAGAGTGATAACGTGCCAAGTGAAGAGGTTGTGAAGAAAATGAAGAACTATTGGAGGCAGCT
CCTCAACGCTAAGCTCATCACTCAGAGAAAGTTCGATAACTTGACTAAGGCTGAGAGGGGAGGACTCTCTGAATT
GGATAAGGCAGGATTCATCAAGAGGCAGCTTGTGGAAACCAGGCAGATCACTAAGCACGTTGCACAGATCCTCGA
TTCTAGGATGAACACCAAGTACGATGAGAACGATAAGTTGATCAGGGAAGTGAAGGTTATCACCCTCAAGTCAAA
GCTCGTGTCTGATTTCAGAAAGGATTTCCAATTCTACAAGGTGAGGGAAATCAACAACTACCACCACGCTCACGAT
GCTTACCTTAACGCTGTTGTTGGAACCGCTCTCATCAAGAAGTATCCTAAGCTCGAGTCAGAGTTCGTGTACGGTG
ATTACAAGGTGTACGATGTGAGGAAGATGATCGCTAAGTCTGAGCAAGAGATCGGAAAGGCTACCGCTAAGTATT
TCTTCTACTCTAACATCATGAATTTCTTCAAGACCGAGATTACCCTCGCTAACGGTGAGATCAGAAAGAGGCCACTC
ATCGAGACAAACGGTGAAACAGGTGAGATCGTGTGGGATAAGGGAAGGGATTTCGCTACCGTTAGAAAGGTGCT
CTCTATGCCACAGGTGAACATCGTTAAGAAACCGAGGTGCAGACCGGTGGATTCTCTAAAGAGTCTATCCTCCCT
AAGAGGAACTCTGATAAGCTCATTGCTAGGAAGAAGGATTGGGACCCTAAGAAATACGGTGGTTTCGATTCTCCT
ACCGTGGCTTACTCTGTTCTCGTTGTGGCTAAGGTTGAGAAGGGAAAGAGTAAGAAGCTCAAGTCTGTTAAGGAA
CTTCTCGGAATCACTATCATGGAAAGGTCATCTTTCGAGAAGAACCCAATCGATTTCCTCGAGGCTAAGGGATACA
AAGAGGTTAAGAAGGATCTCATCATCAAGCTCCCAAAGTACTCACTCTTCGAACTCGAGAACGGTAGAAAGAGGA
TGCTCGCTTCTGCTGGTGAGCTTCAAAAGGGAAACGAGCTTGCTCTCCCATCTAAGTACGTTAACTTTCTTTACCTC
GCTTCTCACTACGAGAAGTTGAAGGGATCTCCAGAAGATAACGAGCAGAAGCAACTTTTCGTTGAGCAGCACAAG
CACTACTTGGATGAGATCATCGAGCAGATCTCTGAGTTCTCTAAAAGGGTGATCCTCGCTGATGCAAACCTCGATA
AGGTGTTGTCTGCTTACAACAAGCACAGAGATAAGCCTATCAGGGAACAGGCAGAGAACATCATCCATCTCTTCAC
CCTTACCAACCTCGGTGCTCCTGCTGCTTTCAAGTACTTCGATACAACCATCGATAGGAAGAGATACACCTCTACCA
AAGAAGTGCTCGATGCTACCCTCATCCATCAGTCTATCACTGGACTCTACGAGACTAGGATCGATCTCTCACAGCTC
GGTGGTGATTCAAGGGCTGATCCTAAGAAGAAGAGGAAGGTTggttcgTACCCATACGATGTTCCTGACTATGCGG
GCTATCCCTATGACGTCCCGGACTATGCAGGATTGTATCCATATGACGTTCCAGATTACgccactagagctgcttaaGCTT
CTCTAGCTAGAGTCGATCGACAAGCTCGAGTTTCTCCATAATAATGTGTGAGTAGTTCCCAGATAAGGGAATTAGG
GTTCCTATAGGGTTTCGCTCATGTGTTGAGCATATAAGAAACCCTTAGTATGTATTTGTATTTGTAAAATACTTCTAT
CAATAAAATTTCTAATTCCTAAAACCAAAATCCAGTACTAAAATCCAGAT

Red font: CaMV 35S promoter with UBQ10, intron 1 (p35S); green font: bpNLS; purple font: SV40 NLS; black font: plant codon-optimized SpCas9 (pcoCas9) containing; dark blue font: Arabidopsis thaliana phosphoribosylanthranilate transferase (PAT1, Genbank no. M96073.1) intron 1 (Trp1 intron 1); orange font: 3xHA tag; blue font: CaMV 35S terminator (t35S).

❖ **Plant selection marker**: pNOS-NptII-tOCS cloned from pICSL11024 (pICH47732::NOSp-NPTII-OCST) (Addgene Plasmid #51144).