1 **Title: Prioritizing landscapes to reconcile biodiversity conservation, ecosystem services,**
2 **and human well-being in India**
3

4 **Authors:** Arjun Srivathsa[1]*, Divya Vasudev[2]*, Tanaya Nair[1,3]*, Stotra Chakrabarti[4], Pranav
5 Chanchani[5], Ruth DeFries[6], Arpit Deomurari[5,7], Sutirtha Dutta[8], Dipankar Ghose[5], Varun R.
6 Goswami[2], Rajat Nayak[9], Amrita Neelakantan[10], Prachi Thatte[5], Srinivas Vaidyanathan[9], Madhu
7 Verma[11], Jagdish Krishnaswamy[12,13,14]*, Mahesh Sankaran[1,14], Uma Ramakrishnan[1,14]*
8
9 *Authors contributed equally to this work
10
11
12
Author affiliations:
1 National Centre for Biological Science, TIFR, Bengaluru, India
2 Conservation Initiatives, Guwahati, India
3 Division of Biosciences, University College London, London, UK
4 Department of Biology, Macalester College, Minnesota, USA
5 World Wildlife Fund, Delhi, India
6 Department of Ecology, Evolution, and Environmental Biology, Columbia University, New
York, USA
7 RoundGlass H2O Pvt. Ltd, Mohali, India
8 Wildlife Institute of India, Dehradun, India
9 Foundation for Ecological Research, Advocacy and Learning, Bengaluru, India
10 Network for Conserving Central India, India
11 World Resources Institute, New Delhi, India
12 Ashoka Trust for Research in Ecology and the Environment, Bengaluru, India
13 School of Environment and Sustainability, Indian Institute for Human Settlements,
Bengaluru, India
14 Biodiversity Collaborative, India
30
31 **Corresponding authors:**
32 Arjun Srivathsa: arjuns@ncbs.res.in
33 Uma Ramakrishnan: uramakri@ncbs.res.in

Abstract

Biodiversity conservation and human well-being are tightly interlinked; yet mismatches in the scale at which both priorities are planned have led to biodiversity loss, erosion of ecosystem services, and declining human quality of life. In India, <5% of the land area is effectively protected for conservation, while the rest is impacted by the second largest human population on the planet. This warrants landscape-level conservation planning through a judicious mix of *land-sharing* and *land-sparing* approaches, and co-production of ecosystem services. Through a multi-faceted assessment, we prioritize spatial extents of land parcels that, in the face of anthropogenic threats, can safeguard conservation landscapes across India's biogeographic zones. We find that only a fraction (~15%) of such priority areas identified here are encompassed under India's extant PA network, and several landscapes of high importance were omitted in all previous global-scale assessments. We then examined the spatial congruence of priority areas with administrative units earmarked for economic development by the Indian government, and propose management-zoning through State-driven and participatory approaches. Our spatially explicit insights can help meet the twin goals of biodiversity conservation and sustainable development in a strategically important country for Asia.

Keywords: *conservation; ecosystem services; global targets; landscape conservation; spatial prioritization; sustainable development*

Main

Concomitant impacts of biodiversity decline, climate change, unsustainable land use, and inequitable extraction of natural resources, have degraded the quality of human life (Levett 2002; Harrison 2010; Otero et al. 2020). Biodiversity underpins several provisioning, regulatory, and cultural or aesthetic ecosystem services; ecosystem services linked to biodiversity, in turn, are crucial for ensuring long-term human well-being (Seppelt et al. 2013). However, the relationship between these two entities can be complex (Mace et al. 2010). Recent discourse on nature-based solutions acknowledges nature's contributions as humanity's 'safety net' (IPBES 2019; Dinerstein et al. 2020), and that ecological processes are shaped by a complex interplay between ecological *and* social systems (Ostrom 2009; Summers et al. 2012; Bennett et al. 2015). A sustainable future can only be ensured by adopting an ecosystems approach to conservation that emphasizes links between human and natural systems to address global, regional and biome-scale threats to their functioning (Haines-Young & Potschin 2010). Siloed management approaches to meet conservation targets that ignore this interplay have failed to yield equitable and sustainable progress; meanwhile, there has been an increase in the frequency and magnitude of calamities arising from crossing the planet's ecological thresholds (Tallis & Kareiva 2006; Steffen et al. 2018). A renewed global resolve will need to consider nature in the Anthropocene, and explicitly reconcile land-use planning for economic development with ecosystem functioning, biodiversity conservation and long-term human well-being.

For nearly two decades, several priority-setting tools have been proposed the world over to identify and rank species, habitats and locations based on their relative importance, vulnerability, or ease of management interventions (Wilson et al. 2007). These approaches generally adopt systematic spatial conservation planning—a set of decision-making tools that help determine the spatial locations where resources and actions need to be directed to optimize conservation gains (Moilanen et al. 2005, 2011; Game et al. 2013). Earlier iterations of these tools largely involved an inclusion-exclusion approach, which helped identify locations for declaring Protected Areas (PAs), determine optimal reserve design, or demarcate conservation corridors (Kremen et al. 2008). But human interests like economic aspirations, the need for infrastructure development, and agricultural expansion to ensure food security must be integrated synergistically with conservation goals to enhance traction with policy makers (Pressey et al. 2007). Incorporating these attributes necessitates adopting *landscape approaches* to conservation (see Sayer et al. 2013), a key goal of which is to provide pragmatic, results-oriented solutions in the Anthropocene. These considerations have led to the idea of 'zoning' landscapes, whereby assessors not only identify priority locations, but also stratify them into various zones and determine appropriate management interventions as part of the planning exercise (Watts et al. 2009; Arkema et al. 2015). This combination of prioritization–zoning therefore holds promise to guide better management of landscapes in an increasingly human-modified world.

Since its conception, the idea of prioritizing areas for conservation at the global level has seen
wide application in scientific literature. While these investigations can offer important insights at
macro-scales, their real-world applications have been limited (Wyborn & Evans 2021). One
reason for this is perhaps the pronounced lack of alignment between scientific 'boundaries'
(sampling units) and administrative jurisdictions; yet, it is within these jurisdictions that
implementation typically occurs (Seppelt et al. 2013). Aligning ecological findings with
administrative boundaries, or existing policy, may thus increase the utility of prioritization
exercises. Second, making assessments at global, cross-national scales may compromise the
spatial resolution of analyses, hampered by the availability and comparability of data at scale, or
miss out on local socio-political nuances (Brooks et al. 2006; Turner et al. 2007; Jenkins et al.
2013; Brum et al. 2017). Third, macro-scale analyses could entail intrinsic biases in
representation of key features or attributes. For example, certain historically overlooked biomes
or under-studied regions may continue to remain ignored (Hoeskstra et al. 2005; Silviera et al.
2021); habitats conventionally presumed to be 'unproductive' may not be prioritized (e.g.,
grasslands, see Bond & Parr 2010); indices such as species richness, which ignore community
composition, may discount rarity or endemism (Veach et al. 2017); PAs or 'intact' wilderness
areas may take precedence over heterogeneous multi-use conservation landscapes (Venter et al.
2014; Potapov et al. 2017); and important dimensions such as human populations may
altogether be excluded (Jung et al. 2021).
127
Achieving conservation goals, sustaining ecosystem services, and ensuring human well-being
while balancing economic development, are universal as well as site-specific issues that present
formidable challenges in implementation (Otero et al. 2020). India exemplifies this premise and
associated challenges; (i) it is a large, diverse country with ten distinct biogeographic zones and
four biodiversity hotspots; (ii) its Protected Area (PA) network, conservation landscapes and
riverscapes support several ecologically important and evolutionarily distinct species
assemblages; (iii) it has the second largest human population in the world, with a large
proportion of the people directly dependent on natural resources drawn from forests, grasslands,
rivers and other ecosystems; and (iv) with rapid infrastructure development and liberal
investment policies, India is currently among the fastest-growing economies in Asia. In this
study, we: (1) identify spatial scale(s), resolutions, and thematic dimensions for a priority-setting
exercise across the country (Fig. 1); (2) use a systematic spatial prioritization approach to
optimize landscapes for biodiversity conservation and ecosystem service gains, while penalizing
locations facing negative anthropogenic impacts; and (3) align our results with global
biodiversity targets (Post-2020 Global Biodiversity Framework, CBD 2020) and the Indian
government's development initiative to identify 'aspirational' economic districts. We provide
insights that can guide future environmental planning and policy to help meet the twin goals of
biodiversity conservation and sustainable development in a strategically important country for
Asia. More broadly, our framework can be adapted and applied for conservation planning in
other eco-regions of developing countries.

Figure 1. Schematic of the study framework, biogeographic zones of India, and themes and input layers used for spatial prioritization. Theme-wise input layers include: (1) Habitats: 28 vegetation classes depicted as 5 composite panels; (2) Ecosystem Services: blue water flux, above- and below-ground carbon, and green water flux; (3) Biodiversity: Protected Areas, Key Biodiversity Areas, and a diversity index of threatened mammals, birds, reptiles and amphibians; (4) Threats: human population density, livestock population density, urbanization, linear infrastructure, mines, river fragmentation, agricultural expansion, vegetation greening, vegetation browning, future climate warming and future rainfall anomaly.

Results and Discussion

a. Overlap in priority sites across themes

Among the most important considerations in conservation prioritization is the extent of spatial congruence among the attributes or dimensions. Priority maps generated independently for habitats, ecosystem services and biodiversity (Fig. 2a) represented sites of high value in isolation, i.e., sites that were: (a) representative of important and rare natural habitats, (b) responsible for the provision of key ecosystem services, viz. water and carbon, and (c) important solely from the perspective of threatened species diversity and turnover. We found moderate overlap amongst these three layers, which varied across biogeographic zones (hereafter, 'zones'; Fig. 2). The overlap of priority sites for biodiversity (defined as the top 30% rankings) and habitats was 37% across the country, ranging from 30% in deserts, to 48% in the Western Ghats (Fig. S1). There was significant addition of sites when ecosystem services was added as a criterion, in addition to biodiversity and habitats. Around 38% of priority sites for ecosystem services were not covered by either biodiversity or habitat priority sites; these locations

172 collectively contain a population of 2.3 million people (based on spatial overlap alone) who are
173 directly dependent on these watersheds (Fig. S1).
174
Several conservationists have highlighted a mismatch in priority sites when different criteria are
used to assign a 'conservation value' (Westgate et al. 2014; Cadotte & Tucker 2018). Our
findings support this concern (Fig. 2) and point to the value of incorporating multiple
contributors to conservation value. Of significance, our multi-criterion approach selected for
India's Open Natural Ecosystems (Madhusudan & Vanak 2021) — prioritized based on habitats,
but not based on biodiversity — encompassing open grasslands, savannas, hot and cold deserts,
ravines, rocky boulders and escarpments. These systems are extremely fragile, with unique
endemic flora and fauna, but inappropriately classified as "wastelands" as per India's land-use
and conservation policy (Wastelands Atlas of India 2019). Similarly, areas ranked high for
habitat and biodiversity, but low in terms of ecosystem services represented some locations in
the arid/semi-arid dry zones of western India, the cold deserts of trans-Himalayas, and parts of
the terai grasslands along the India-Nepal border (Fig. 2a). Conversely, areas ranked high for
ecosystem services, critical for water and carbon storage, did not rank high for biodiversity or
habitats in some places (e.g., parts of Northeast India). This, perhaps, is due to low coverage of
traditional PAs, or the lack of sufficient data on biodiversity in the region.
190
b. Synergies and trade-offs among themes
Alongside the spatial congruence discussed above, we acknowledge that there could be trade-
offs between the thematic dimensions considered here. Carbon sequestration and hydrologic
services may not necessarily work in synergy (Chisholm 2010). Similarly, biodiversity
conservation could constrain provisioning services such as non-timber forest products or
livestock-grazing in certain habitats. Such trade-offs could also be spatially asynchronous. For
instance, up-stream water regulation and abstraction in rivers, and disruption of sediment
transport by trapping behind dams could bear deleterious impacts on aquatic biodiversity and
ecosystem services downstream. Regulatory services, like hydrologic or water services, can be
generated only at certain size of catchment depending on factors such as climate, soils, geology
and vegetation type. And while there could be synergy between carbon and water services to
people at a local or regional scale (Clark et al. 2021), the transpiration from a large patch of
forest can increase rainfall in other regions and benefit agriculture and communities elsewhere
(Paul et al. 2018). We also caution that we used only a subset of ecosystem services deemed
important for human well-being; services like pollination, forest produce extraction, rangeland
services, or freshwater dependence, could not be included due to unavailability of spatially
explicit data on these attributes. Incorporating these features, together with other factors such as
the presence and population size of endemic species, the conservation value of community-
managed forests, or rare habitats that we might have missed, may yield a more comprehensive
assessment of priority locations and landscapes.

212
Figure 2. (a) Theme-wise results from prioritization analyses of Habitats (top left), Ecosystem services (centre left), Biodiversity (bottom left), and Composite of the three themes (right); (b) Theme-specific results from prioritization analysis of Threats.

c. Appraisal of anthropogenic pressures

Reconciling biodiversity conservation concerns and the demands of economic development is among the greatest challenges of the 21st century. We explicitly incorporated *Threats* (Fig. 2b) both, as factors that make conservation more costly, and also such that the final priority ranks represent such a reconciliation (Fig. 3). Across zones, a consistent pattern we observed was the compounding effects of agricultural expansion and urbanization, coupled with vegetation greening (indicative of increased year-round irrigation in agriculture areas) and linear infrastructure (Fig. 2b). Thus, the agricultural belts of (i) the northern semi-arid zone, (ii) the lowland plains of Northeast India, (iii) western and southern parts of the Deccan Peninsula, and urban hotspots representing major cities were ranked the highest in terms of anthropogenic pressures (Fig. 2b; Fig. S1). Of these, the northern semi-arid zone and the western parts of Deccan peninsula also had pronounced signatures of vegetation browning and greening, respectively. Vegetation browning could arise from forest degradation, deforestation, or forest fires; it could also be due to natural shifts in woody vegetation to grass dominated systems, or loss of green foliar biomass due to climate change impacts. Vegetation greening could be due to ecological restoration or natural regeneration of native species; but the more ubiquitous source is the proliferation of a single invasive alien species (e.g., *Prosopis*, *Lantana*, etc.; Krishnaswamy et al. 2014).

Our results pertaining to threats, when viewed in the context of projected human population growth and the demands of meeting future food security concerns, are representative of areas that either have or will soon surpass thresholds beyond which interventions for sustainable land-use practices may not be feasible (Martin et al. 2022). On the other hand, our analyses could not

fully elicit the broader deleterious impacts of localized threats imposed by mines and river dams.
Further, while we did incorporate projected temperature increase and future rainfall anomalies to
account for climate change impacts, zones that are likely to be the most vulnerable to these
threats, i.e., coastal areas and island systems, were not part of our assessment. Nevertheless,
climate change is still projected to impact various ecosystems considered in our analysis in
significant ways. For example, predicted increases in the magnitude and frequency of major
floods can shape the distribution of biodiversity on the floodplains of Northeast India (see
Goswami et al. 2021). Such influences underscore the importance of spatial conservation
planning based on prioritization efforts of the kind we undertake here. We also note that climatic
influences on terrestrial systems are extremely dynamic and are pivotally linked to national and
global policy changes in the future.
251
d. Landscape-level approach to conservation
Recent conservation literature has highlighted the importance of viewing shared habitats as
coupled human–natural systems that are encompassed within 'conservation landscapes' (Das et
al. 2006; Rodrigues et al. 2021; Vasudev et al. 2021). Across zones, we found that most
designated PAs, which constitute ~5% of India's land area and span an average size ~300 km^2
(Ghosh-Harihar et al. 2019), were included as priority sites. But these were embedded within
larger landscapes that also included high priority non-PA locations; 85% of the top 30% priority
sites were outside PAs (Fig. 3). This was reinforced by the specific inclusion of ecosystems
(such as the Open Natural Ecosystems referred to above) that are not part of India's PA network,
and sites critical for the supply of ecosystem services. Interestingly, locations that constituted the
top 30% priority ranks in our study were largely connected (Fig. 3), even when we did not
explicitly impose connectivity parameters via the prioritization model. While there was spatial
clumping at the regional scale, there remains a degree of fragmentation in the final priority
locations, especially in north-western and south-eastern India (Fig. 3b). Although our zone-wise
analyses allowed for better geographic representation of sites, it still indicated lack of
connectivity across zone boundaries. These issues together emphasize that functional
connectivity is an important consideration while implementing landscape-scale conservation
interventions within our priority sites (see Brennan et al. 2022).
270
Traditional PAs, which typify a *land-sparing* approach to conservation, are mostly focused on
forested habitats in the country. A substantially large proportion of biodiversity continues to
inhabit unprotected, human-use landscapes, warranting a *land-sharing* approach. Realizing
conservation and human well-being goals in the landscapes that encompass the 30% priority
areas will therefore necessitate invoking Other Effective area-based Conservation Measures
(OECMs). The core tenets of these approaches hinge fundamentally on effective and *equitable*
models of governance that are fully cognizant of the complexities of socio–ecological systems,
all of which are only beginning to be recognized in environmental and conservation policy
(Alves-Pinto et al. 2021; Jonas et al. 2021). In India, this may be achieved through

implementation of existing frameworks, for instance, in locations where communities are granted Community Forest Rights, and declaring areas as Critical Wildlife Habitats under the Forest Rights Act — provisions which, at present, remain extremely under-utilized. Our spatially explicit landscape approach that considers linked biodiversity–ecosystem service dimensions can only succeed if the beneficiaries of ecological restoration, and those who may be otherwise impacted (typically the marginalized sections of society) are addressed in policies and implementation.

Figure 3. (a) Conservation priority ranks for the country, computed zone-wise and combined; (b) Top 30% priority areas from each zone demarcated as a set of three 10% blocks. Designated Protected Areas are overlaid to show the extent of overlap and spatial congruence with high priority locations.

e. Linking landscape prioritization and administration

Overlaying administrative (district) boundaries, we highlight 338 districts that play a key role in maintaining India's biodiversity and ecosystem services (Fig. 4a and 4b); of these, 169 are 'high priority' districts, where the management focus needs to be on retention of habitats, ecosystem services and biodiversity through both, State-driven and participatory approaches. The next 169 are 'potential priority' districts (Fig. 4, centre panel), where the management focus, besides retention of important sites, should also aim for proactive rewilding and ecological restoration efforts. In conjunction, we also refer to India's aspirational districts, identified by the Government of India's NITI Aayog for economic development (see Methods for details). Locations where aspirational districts overlap with 'high priority' districts (Fig. 4c) require deprioritizing mega-infrastructure projects while promoting sustainable models of nature protection besides demarcation of PAs. Such approaches may entail community stewardship for biodiversity protection, co-management of habitats outside PAs, and nature-friendly livelihood

306 development, within larger conservation landscapes. In locations where aspirational districts
307 overlap with 'potential priority' districts, targeted actions are essential to mitigate negative
308 impacts of infrastructure development through economically incentivized instruments like
309 paying for ecosystem services, promoting agroforestry, and where appropriate, demarcating
310 conservation and/or community reserves.
311
The relationship between ecosystem services and poverty alleviation (especially in the
aspirational and poorer districts of India as defined by the NITI Aayog discussed above) is also
driven by political economy of negotiations between stakeholders and those who manage or
regulate ecosystem services (Vira et al. 2012). A detailed understanding of the associated
vulnerabilities is important to enable ecosystem services to benefit the poor. If not internalized,
an ecosystem services approach to conservation planning may fail because of resistance from
those who are excluded or those who stand to lose most from such undertakings. The actions
discussed above will also be critical in other countries of the Global South, where a large
proportion of the people are directly dependent on forests and other natural ecosystems for their
livelihoods. The nuances and location-based differences in prescriptive actions we discuss here
reiterate the importance of (a) spatially explicit prioritization assessments incorporating regional
expertise, and (b) mainstreaming biodiversity and ecosystem service goals into district and state
management plans for socio-economic development. In this context, we posit that India's
ambitious endeavours on climate change mitigation through reversal of land degradation (26
million ha), the Green India Mission (Ravindranath & Murthy 2010), as well as renewable
energy projects initiated under the Intended Nationally Determined Contribution (NDC) will
require a degree of reorientation.
329

Figure 4. Results from spatial prioritization analyses processed at the scale of administrative units in the
country. Left panel: Proportion of each district covered by areas identified as top 30% priority locations.
Centre panel: Districts in the top quantile (25%) identified as 169 'high priority' districts and the next 25%
quantile identified as 169 'potential priority' districts. Darker lines denote state boundaries. Right panel:
NITI Aayog aspirational districts (112) earmarked by the Government of India, and their overlap with 338
priority districts identified in this study (72 districts overlap). Darker lines in the centre and right panels
denote state boundaries.
338
339

Conclusion
Repercussions of the COVID-19 pandemic, and a clear recognition of links between ecosystem
integrity and human health, have led to broad and ambitious goals under global targets such as
the Post-2020 Global Biodiversity Framework. However, there exists a wide geographical
imbalance in the capacities, resources, and vulnerabilities of different countries to achieve such
goals. The Global South, in particular, aims to conserve biodiversity and meet climate change
goals under scenarios of high dependency of people on natural resources (Fedele et al. 2021),
coupled with aspirations of economic progress, and better standards of human life. To address
these expectations, national scale prioritization exercises such as ours, which combine
conservation priorities, human well-being indices, economic development and infrastructure
considerations can guide countries and governments towards meeting international targets set
forth for the next decade (e.g., Strassburg et al. 2020); linkages between prioritization exercises
and existing government schemes and administrative boundaries are crucial in this regard (see
Srivathsa et al. 2020; Belote et al. 2021). In the context of elevated competition for scarce land
and water, exacerbated due to future climate change, the importance of our framework lies in its
ability to objectively and effectively address trade-offs at the intersection of sustainable
development goals, conservation of biodiversity and ecosystem services.
357
358

Methods

a. Framework for prioritization

The main goal of our study was to prioritize representative landscapes for biodiversity conservation, and securing of ecosystem services and human well-being targets, while balancing these with the economic aspirations of 1.4 billion people. Our prioritization exercise is part of a larger program aimed at mainstreaming biodiversity conservation into the discourse on development and human well-being — the Government of India's National Mission for Biodiversity and Human Well-Being (NMBHWB; Bawa et al. 2021). Recognizing the importance of delineating, characterizing, and evaluating functional landscapes, the NMBHWB set up a working group in 2020, which was tasked with deliberating on and determining landscape-level approaches for biodiversity conservation, keeping nature's benefit to human well-being as the central theme. Through consultations with 42 experts from across the country, the working group first outlined the need for, salient features of, and the prerequisites for a priority-setting exercise. Subsequently, the working group engaged with 18 field and domain experts (authors of the present study) to undertake the prioritization exercise in India.

b. Prioritization approach

We used a two-step prioritization approach (Fig. 1) to ensure representation of geographies and eco-regions across India. We first chose biogeographic zones ('zones') as described by Rodgers and Panwar (1988) as an appropriate level of primary spatial classification. This level of classification splits the country into zones that have some similarity of biogeography, while across zones, they capture diversity of species, ecosystems, and human–nature relationships. The country has 10 zones: (i) trans-Himalayas, (ii) Himalayas, (iii) deserts, (iv) semi-arid, (v) Western Ghats, (vi) Deccan plateau, (vii) Gangetic plains, (viii) Northeast India, (ix) coasts and (x) islands. We excluded coasts and islands from our analyses as we deemed them to be unique and requiring a separate treatment of biodiversity, ecosystem services and threats. The eight selected zones are shown in Fig. 1 (additional descriptions are in Table S1). We then identified three broad focus themes, each of which encompassed a set of criteria for prioritization.

i)   We considered ecosystems or natural habitats that require inclusion in priority landscapes. These layers, with zone-specific sets of priority ecosystems, represented our *Habitats* theme.

ii) Recognizing the increasing emphasis on human–nature links and joint well-being of biodiversity and people, we considered *Ecosystem Services* as the second theme.

iii) Species diversity, biodiversity hotspots and locations of species population sources collectively formed the *Biodiversity* theme.

Lastly, we included a *Threats* theme to recalibrate conservation priorities based on spatial variation in anthropogenic pressures and impacts on natural ecosystems and biodiverse areas that provision essential ecosystem services. The full list of input layers is referenced in Fig. 1 and detailed in Table S2.

c. Spatial prioritization
We used software Zonation 4.0 (Moilanen et al. 2014) to determine priority areas in each zone.
The program's algorithm follows an iterative process to rank cells (in our case, 1-km^2 pixels)
within the extent of interest, by excluding cells one at a time and measuring the collective
conservation value of retained cells. In other words, cells with the least value are removed first
and those with the highest values are retained till the end. The final output thus generated is an
optimized map whose pixels are ranked based on relative priority values. For additional details
see Moilanen et al. (2014). For each zone, we implemented a set of six prioritization analyses
(four theme-wise runs and two composite runs) as explained below.
407
Habitats. We used data from Roy et al. (2015) with 156 land-cover classes across the country.
We first reduced the data into 28 habitat classes by grouping similar land cover classes, broadly
following the forest type classification of Champion and Seth (1968). For instance, land cover
types labelled "Grassland", "Manmade grassland", "*Lasiurus-Panicum* grassland", "*Cenchrus-*
Dactyloctenium grassland", "*Aristidia-oropetium* grassland" and "*Sehima-Dichanthium*
grassland" were reclassified collectively as "Grasslands". Data from 25-m^2 pixels of the original
dataset were resampled at a 1-km^2 resolution; the value assigned to each 1-km^2 pixel for each of
the 28 habitat classes reflected the proportion of 25-m^2 pixels of the corresponding class. For
each zone, we further sub-selected habitat classes to retain (a) the dominant vegetation classes
(covering >1% of the total zone area), and (b) select rare, vulnerable habitats, determined based
on our experience and knowledge of the landscape(s). For example, we included wet grasslands
in the Northeast zone due to its ecological and livelihood importance, even though this habitat
class covered only 0.25% of the zone. The final number of habitat types chosen for each zone
thus varied between three (Desert zone) and 10 (Western Ghats zone; Table S2). All habitats
were assigned equal weights for prioritization. We chose the 'Core Area Zonation' cell removal
rule as it prioritizes pixels with the most important yet rare features; this way, pixels that
contained high proportions of rare habitats were retained even if the other zone-wide dominant
habitats within these pixels were low.
426
Ecosystem Services. We identified and used three layers that collectively reflected key ecosystem
services, for which we were able to obtain or generate spatially explicit metrics across the
country. These included (i) blue water flux, or the amount of water available as surface water or
as ground water to meet utilitarian needs and for ecological flows in rivers, (ii) green water flux,
or the amount of water lost via surface evaporation or through transpiration, which contributes to
multiple ecosystem services, from flood-control in very wet regions to recycling back as rainfall
in other regions, and (iii) carbon stock, a harmonized metric of aboveground and belowground
biomass carbon density calculated using woody plant, grassland, and cropland biomass (Table
S2). While these do not represent a comprehensive set of ecosystem services, we believe they
capture two critical policy mandates for India: water security and climate change mitigation. We
assigned equal weights to all three layers and chose the Additive Benefit Function 'ABF' cell

removal rule in Zonation, to ensure that locations with relatively high values of all three features
are prioritized higher than areas where individual features or subsets of the three features had
high values.
441
Biodiversity. To represent important locations for biodiversity, we first considered designated
Protected Areas (PAs), reasonably assuming that these locations currently harbor source
populations of many species. In addition, we also used Key Biodiversity Areas (KBAs; BirdLife
International 2021), as these include locations outside designated PAs with high diversity of
species that are of ecological or conservation importance. Next, we collated distribution ranges
of imperilled mammals, birds, amphibians, and reptiles—species categorized as Critically
Endangered, Endangered, Vulnerable or Threatened as per the IUCN Red List (IUCN 2020). For
each zone, we stacked the species maps and generated a beta-diversity index, calculated as the
Bray-Curtis distance between a focal cell and a hypothetical reference cell which had all
threatened species of the corresponding zone. This gave us an index of zone-wise species
turnover. We assigned equal weights to all three layers and chose the ABF cell removal rule in
Zonation to prioritize areas based on this theme.
454
Threats. We identified 11 attributes that negatively impact biodiversity, ecosystem services and
habitats to varying degrees, and are likely to reduce the ease of implementing conservation
actions. These included human population density, livestock population density, urbanization,
linear infrastructure, mines, river fragmentation, agricultural expansion, vegetation greening and
vegetation browning (both of which can have positive or negative impacts on ecosystem services
and biodiversity), future climate warming and future rainfall anomaly (see Table S2 for details).
We assigned zone-specific weights to these layers following a consensus-approach, based on the
collective field knowledge of the assessors. The weights assigned to the threats varied across
zones, reflecting the relative severity of impacts in the corresponding zone (see Table S3 for
zone-wise threat ranks). We chose the ABF cell removal rule again, so that areas retained
towards the end were representative of locations with high values of multiple threats combined.
466
Composite. In addition to the four sets of theme-wise analyses described above, we also
undertook two composite analyses. First, we used the outputs from the theme-wise analyses of
Habitats, *Ecosystem Services* and *Biodiversity* features. All input layers were assigned equal
weights and the ABF cell removal rule was specified. Second, we used the aforementioned three
layers along with the rank output from the *Threats* analysis. In this case, we assigned equal
positive weights to *Habitats*, *Ecosystem Services* and *Biodiversity*, and an equal but negative
weight to *Threats* (i.e., 1, 1, 1, and -1). The first analysis thus gave us an assessment of
conservation importance, while the second incorporated feasibility.

d. Alignment with global and national policy

We collated results across zones post-prioritization to demarcate the top 30% priority areas in the country, so as to align our results with the Post-2020 Global Biodiversity Framework targets. The areas were selected based on pixel ranks in each zone; these covered 30% of the area in every zone, and therefore, collectively included 30% of the country (sans the coasts and islands). This approach ensured representation of unique features from each zone, encompassing a more diverse set of habitats, ecosystem services, biodiversity, and associated threats. We then overlaid administrative units (districts) and calculated the proportion of each district covered by pixels with the top 30% ranks. We chose two sets of districts: (1) those in the top 25% quartile, which we term 'high priority' districts, and (2) those in the subsequent 25% quartile, which we term 'potential priority' districts. We juxtapose these results with districts earmarked by the Government of India's NITI Aayog under its flagship Aspirational Districts Programme for economic development (www.niti.gov.in/aspirational-districts-programme). Our maps were generated with the goal of directly offering prescriptive management actions for the governments of the corresponding administrative units.

Acknowledgements

This work was catalysed and supported by the Office of the Principal Scientific Adviser to the Government of India as part of the National Mission for Biodiversity and Human Well-Being. A.S. was supported by the Department of Science and Technology–Government of India's Innovation in Science Pursuit for Inspired Research (DST INSPIRE) Faculty Award. We are grateful to J. Zacharias, V. Athreya, P. Bindra, A. Harihar, U. Ganguly, A. Kumar, M. Manuel, S. Babu, A. Kshettry, S. Nijhawan, M. Muralidharan, N. Namboothiri, Y. Jhala, K.S. Subin, S. Datta, M. Sen, S. Madhulkar, T. Thekaekara, V. Vasudevan, I. Anwardeen, S. Sahu, S. Reddy, V. Varadhan, K.T. Subramaniam, C. Madegowda, C. Meenakshi, K. Karanth, P. G. Krishnan, T. Dash, A. Chanchani and A. Bijoor for partaking in discussions on landscape-scale conservation in India. We thank K. Bawa and R. Chellam for their guidance, R.G. Rodrigues, A. Samrat and N. Srinivas for assistance with data processing and compilation, and the National Centre for Biological Sciences-TIFR and Ashoka Trust for Research in Ecology and the Environment (part of the Biodiversity Collaborative) for facilitating this study.

References

Alves-Pinto, H., Geldmann, J., Jonas, H., Maioli, V., Balmford, A., Latawiec, A. E., ... & Strassburg, B. (2021). Opportunities and challenges of other effective area-based conservation measures (OECMs) for biodiversity conservation. *Perspectives in Ecology and Conservation*, *19*(2), 115–120.

Arkema, K. K., Verutes, G. M., Wood, S. A., Clarke-Samuels, C., Rosado, S., Canto, M., ... & Guerry, A. D. (2015). Embedding ecosystem services in coastal planning leads to better

515   outcomes for people and nature. *Proceedings of the National Academy of Sciences*, *112*(24),
516   7390–7395.
517
518   Bawa, K. S., Sengupta, A., Chavan, V., Chellam, R., Ganesan, R., Krishnaswamy, J., ... &
519   Vanak, A. T. (2021). Securing biodiversity, securing our future: A national mission on
520   biodiversity and human well-being for India. *Biological Conservation*, *253*, 108867.
521
522   Belote, R. T., Aplet, G. H., Carlson, A. A., Dietz, M. S., May, A., McKinley, P. S., ... &
523   Garncarz, J. (2021). Beyond priority pixels: Delineating and evaluating landscapes for
524   conservation in the contiguous United States. *Landscape and Urban Planning*, *209*, 104059.
525
526   Bennett, E. M., Cramer, W., Begossi, A., Cundill, G., Díaz, S., Egoh, B. N., ... & Woodward, G.
527   (2015). Linking biodiversity, ecosystem services, and human well-being: three challenges for
528   designing research for sustainability. *Current Opinion in Environmental Sustainability*, *14*,
529   76–85.
530
BirdLife International (2021). *World Database of Key Biodiversity Areas*. Developed by
the KBA Partnership: BirdLife International, International Union for the Conservation of
Nature, American Bird Conservancy, Amphibian Survival Alliance, Conservation
International, Critical Ecosystem Partnership Fund, Global Environment Facility, Global
Wildlife Conservation, NatureServe, Rainforest Trust, Royal Society for the Protection of
Birds, Wildlife Conservation Society and World Wildlife Fund. March 2021 version.
Accessed form: www.keybiodiversityareas.org/kba-data/request
538
539   Bond, W. J., & Parr, C. L. (2010). Beyond the forest edge: Ecology, diversity and conservation
540   of the grassy biomes. *Biological Conservation*, *143*(10), 2395–2404.
541
542   Brennan, A., Naidoo, R., Greenstreet, L., Mehrabi, Z., Ramankutty, N., & Kremen, C. (2022).
543   Functional connectivity of the world's protected areas. *Science*, *376*(6597), 1101–1104.
544
545   Brooks, T. M., Mittermeier, R. A., Da Fonseca, G. A., Gerlach, J., Hoffmann, M., Lamoreux, J.
546   F., ... & Rodrigues, A. S. (2006). Global biodiversity conservation
547   priorities. *Science*, *313*(5783), 58–61.
548
549   Brum, F. T., Graham, C. H., Costa, G. C., Hedges, S. B., Penone, C., Radeloff, V. C., ... &
550   Davidson, A. D. (2017). Global priorities for conservation across multiple dimensions of
551   mammalian diversity. *Proceedings of the National Academy of Sciences*, *114*(29), 7641–
552   7646.
553

554 Cadotte, M. W., & Tucker, C. M. (2018). Difficult decisions: Strategies for conservation
555   prioritization when taxonomic, phylogenetic and functional diversity are not spatially
556   congruent. *Biological Conservation*, *225*, 128–133.
557

558 CBD (2020). First draft of the Post-2020 Global Biodiversity Framework; released in June 2021.
559   Accessed from: www.cbd.int/conferences/post2020
560

561 Champion, H. G., & Seth, S. K. (1968). *A revised survey of the forest types of India*. Manager of
562   Publications, Government of India.
563

564 Chisholm, R.A. (2010). Trade-offs between ecosystem services: water and carbon in a
565   biodiversity hotspot. *Ecological Economics*, *69*(10), 1973–1987.
566

567 Clark, B., DeFries, R., & Krishnaswamy, J. (2021). India's commitments to increase tree and
568   forest cover: consequences for water supply and agriculture production within the central
569   Indian Highlands. *Water*, *13*(7), 959.
570

571 Das, A., Krishnaswamy, J., Bawa, K. S., Kiran, M. C., Srinivas, V., Kumar, N. S., & Karanth, K.
572   U. (2006). Prioritisation of conservation areas in the Western Ghats, India. *Biological
573   Conservation*, *133*(1), 16–1.
574

575 Dinerstein, E., Joshi, A. R., Vynne, C., Lee, A. T. L., Pharand-Deschênes, F., França, M., ... &
576   Olson, D. (2020). A "Global Safety Net" to reverse biodiversity loss and stabilize Earth's
577   climate. *Science Advances*, *6*(36), eabb2824.
578

579 Fedele, G., Donatti, C. I., Bornacelly, I., & Hole, D. G. (2021). Nature-dependent people:
580   Mapping human direct use of nature for basic needs across the tropics. *Global Environmental
581   Change*, *71*, 102368.
582

583 Game, E. T., Kareiva, P., & Possingham, H. P. (2013). Six common mistakes in conservation
584   priority setting. *Conservation Biology*, *27*(3), 480–485.
585

586 Ghosh-Harihar, M., An, R., Athreya, R., Borthakur, U., Chanchani, P., Chetry, D., ... & Price, T.
587   D. (2019). Protected areas and biodiversity conservation in India. *Biological Conservation*,
588   *237*, 114–124.
589

590 Goswami, V. R., Vasudev, D., Joshi, B., Hait, P., & Sharma, P. (2021). Coupled effects of
591   climatic forcing and the human footprint on wildlife movement and space use in a dynamic
592   floodplain landscape. *Science of The Total Environment*, *758*, 144000.
593

594 Haines-Young, R., & Potschin, M. (2010). The links between biodiversity, ecosystem services
595   and human well-being. *Ecosystem Ecology: A new synthesis*, *1*, 110–139.
596
597 Harrison, P. A. (2010). Ecosystem services and biodiversity conservation: an introduction to the
598   RUBICODE project. *Biodiversity and Conservation*, *19*(10), 2767–2772.
599
600 Hoekstra, J. M., Boucher, T. M., Ricketts, T. H., & Roberts, C. (2005). Confronting a biome
601   crisis: Global disparities of habitat loss and protection. *Ecology Letters*, *8*(1), 23–29.
602
603 IPBES (2019). Summary for policymakers of the global assessment report on biodiversity and
604   ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and
605   Ecosystem Services. Accessed from: https://ipbes.net/document-library-categories
606
607 Jenkins, C. N., Pimm, S. L., & Joppa, L. N. (2013). Global patterns of terrestrial vertebrate
608   diversity and conservation. *Proceedings of the National Academy of Sciences*, *110*(28),
609   e2602–e2610.
610
611 Jonas, H. D., Ahmadia, G. N., Bingham, H. C., Briggs, J., Butchart, D. H. M., Cariño, J., ... &
612   von Weizsäcker, C. (2021). Equitable and effective area□based conservation: towards the
613   conserved areas paradigm. *PARKS: The International Journal of Protected Areas and
614   Conservation*, *27*, 74–81.
615
616 Jung, M., Arnell, A., De Lamo, X., García-Rangel, S., Lewis, M., Mark, J., ... & Visconti, P.
617   (2021). Areas of global importance for conserving terrestrial biodiversity, carbon and
618   water. *Nature Ecology & Evolution*, *5*(11), 1499–1509.
619
620 Kremen, C., Cameron, A., Moilanen, A., Phillips, S. J., Thomas, C. D., Beentje, H., ... & Zjhra,
621   M. L. (2008). Aligning conservation priorities across taxa in Madagascar with high-
622   resolution planning tools. *Science*, *320*(5873), 222–226.
623
624 Krishnaswamy, J., John, R., & Joseph, S. (2014). Consistent response of vegetation dynamics to
625   recent climate change in tropical mountain regions. *Global Change Biology*, *20*(1), 203–215.
626
627 Levett, R. (1998). Sustainability indicators—integrating quality of life and environmental
628   protection. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, *161*(3),
629   291–302.
630
631 Mace, G.M., Norris, K., & Fitter, A.H. (2012). Biodiversity and ecosystem services: A multi-
632   layered relationship. *Trends in Ecology & Evolution*, *27*(1), 19–26.
633

Madhusudhan, M.D., & Vanak, A.T. (2021). Mapping the distribution and extent of India's
semi-arid open natural ecosystems. Earth and Space Science Open Archive. Accessed from
www.doi.org/10.1002/essoar.10507612.1

Martin, D. A., Andrianisaina, F., Fulgence, T. R., Osen, K., Rakotomalala, A. A. N. A.,
Raveloaritiana, E., ... & Kreft, H. (2022). Land-use trajectories for sustainable land system
transformations: Identifying leverage points in a global biodiversity hotspot. *Proceedings of*
the National Academy of Sciences, *119*(7), e2107747119.

Moilanen, A., Franco, A. M., Early, R. I., Fox, R., Wintle, B., & Thomas, C. D. (2005).
Prioritizing multiple-use landscapes for conservation: Methods for large multi-species
planning problems. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1575),
1885–1891.

Moilanen, A., Anderson, B. J., Eigenbrod, F., Heinemeyer, A., Roy, D. B., Gillings, S., ... &
Thomas, C. D. (2011). Balancing alternative land uses in conservation prioritization.
Ecological Applications, *21*(5), 1419–1426.

Moilanen, A., Pouzols, F.M., Meller, L., Veach, V., Arponen, A., Leppänen, J., & Kujala. H.
(2014). Zonation spatial conservation planning methods and software. *Version 4. User*
Manual. Helsinki, Finland: C-BIG Conservation Biology Informatics Group, Department of
Biosciences, University of Helsinki. Accessed from:
www.cbig.it.helsinki.fi/software/zonation

Ostrom, E. (2009). A general framework for analyzing sustainability of social-ecological
systems. S*cience*, *325*(5939), 419-422.

Otero, I., Farrell, K. N., Pueyo, S., Kallis, G., Kehoe, L., Haberl, H., ... & Pe'Er, G. (2020).
Biodiversity policy beyond economic growth. *Conservation Letters*, *13*(4), e12713.

Paul, S., Ghosh, S., Rajendran, K., & Murtugudde, R. (2018). Moisture supply from the Western
Ghats forests to water deficit east coast of India. *Geophysical Research Letters*, *45*(9), 4337–
4344.

Potapov, P., Hansen, M. C., Laestadius, L., Turubanova, S., Yaroshenko, A., Thies, C., ... &
Esipova, E. (2017). The last frontiers of wilderness: Tracking loss of intact forest landscapes
from 2000 to 2013. *Science Advances*, *3*(1), e1600821.

Pressey, R. L., Cabeza, M., Watts, M. E., Cowling, R. M., & Wilson, K. A. (2007). Conservation
planning in a changing world. *Trends in Ecology & Evolution*, *22*(11), 583–592.

Ravindranath, N.H., & Murthy, I.K. (2010). Greening India Mission. *Current Science*, *99(4)*,
444–449.

Rodgers, W.A., & Panwar, H.S. (1988). Planning a wildlife Protected Area network in India,
Vol. 1. A Report. Wildlife Institute of India, Dehradun, India.

Rodrigues, R. G., Srivathsa, A., & Vasudev, D. (2022). Dog in the matrix: envisioning
countrywide connectivity conservation for an endangered carnivore. *Journal of Applied*
Ecology, *59*(1), 223–237.

Roy, P. S., Roy, A., Joshi, P. K., Kale, M. P., Srivastava, V. K., Srivastava, S. K., ... &
Kushwaha, D. (2015). Development of decadal (1985–1995–2005) land use and land cover
database for India. *Remote Sensing*, *7*(3), 2401–2430.

Sayer, J., Sunderland, T., Ghazoul, J., Pfund, J. L., Sheil, D., Meijaard, E., ... & Buck, L. E.
(2013). Ten principles for a landscape approach to reconciling agriculture, conservation, and
other competing land uses. *Proceedings of the National Academy of Sciences*, *110*(21),
8349–8356.

Seppelt, R., Lautenbach, S., & Volk, M. (2013). Identifying trade-offs between ecosystem
services, land use, and biodiversity: a plea for combining scenario analysis and optimization
on different spatial scales. *Current Opinion in Environmental Sustainability*, *5*(5), 458–463.

Silveira, F. A., Ordóñez Parra, C. A., Moura, L. C., Schmidt, I. B., Andersen, A. N., Bond, W.,
... & Pennington, R. T. (2021). Biome Awareness Disparity is BAD for tropical ecosystem
conservation and restoration. *Journal of Applied Ecology*. Doi: 10.1111/1365-2664.14060.

Srivathsa, A., Majgaonkar, I., Sharma, S., Singh, P., Punjabi, G. A., Chawla, M. M., & Banerjee,
A. (2020). Opportunities for prioritizing and expanding conservation enterprise in India using
a guild of carnivores as flagships. *Environmental Research Letters*, *15*(6), 064009.

Steffen, W., Rockström, J., Richardson, K., Lenton, T. M., Folke, C., Liverman, D., ... &
Schellnhuber, H. J. (2018). Trajectories of the Earth System in the Anthropocene.
Proceedings of the National Academy of Sciences, *115*(33), 8252–8259.

Strassburg, B. B., Iribarrem, A., Beyer, H. L., Cordeiro, C. L., Crouzeilles, R., Jakovac, C. C., ...
& Visconti, P. (2020). Global priority areas for ecosystem restoration. *Nature*, *586*(7831),
724–729.

Summers, J. K., Smith, L. M., Case, J. L., & Linthurst, R. A. (2012). A review of the elements of
human well-being with an emphasis on the contribution of ecosystem services. *Ambio*, *41*(4),
327–340.
717

718 Tallis, H. M., & Kareiva, P. (2006). Shaping global environmental decisions using socio-
719   ecological models. *Trends in Ecology & Evolution*, *21*(10), 562–568.
720

721 Turner, W. R., Brandon, K., Brooks, T. M., Costanza, R., Da Fonseca, G. A., & Portela, R.
722   (2007). Global conservation of biodiversity and ecosystem services. *BioScience*, *57*(10),
723   868–873.
724

725 Vasudev, D., Goswami, V. R., Srinivas, N., Syiem, B. L. N., & Sarma, A. (2021). Identifying
726   important connectivity areas for the wide□ranging Asian elephant across conservation
727   landscapes of Northeast India. *Diversity and Distributions*, *27*(12), 2510–2526.
728

729 Veach, V., Di Minin, E., Pouzols, F. M., & Moilanen, A. (2017). Species richness as criterion for
730   global conservation area placement leads to large losses in coverage of biodiversity.
731   *Diversity and Distributions*, *23*(7), 715–726.
732

733 Venter, O., Fuller, R. A., Segan, D. B., Carwardine, J., Brooks, T., Butchart, S. H., ... & Watson,
734   J. E. (2014). Targeting global protected area expansion for imperiled biodiversity. *PLoS*
735   *Biology*, *12*(6), e1001891.
736

737 Vira, B., Adams, B., Agarwal, C., Badiger, S., Hope, R.A., Krishnaswamy, J., & Kumar, C.,
738   (2012). Negotiating trade-offs: choices about ecosystem services for poverty
739   alleviation. *Economic and Political Weekly*, 67–75.
740

741 Wastelands Atlas of India (2019). Department of Land Resources (Ministry of Rural
742   Development); National Remote Sensing Centre, Indian Space Research Organisation
743   (Department of Space). Government of India. Accessed from:
744   www.dolr.gov.in/documents/wasteland-atlas-of-india
745

746 Watts, M. E., Ball, I. R., Stewart, R. S., Klein, C. J., Wilson, K., Steinback, C., ... & Possingham,
747   H. P. (2009). Marxan with Zones: Software for optimal conservation based land-and sea-use
748   zoning. *Environmental Modelling & Software*, *24*(12), 1513-1521.
749

750 Westgate, M. J., Barton, P. S., Lane, P. W., & Lindenmayer, D. B. (2014). Global meta-analysis
751   reveals low consistency of biodiversity congruence relationships. *Nature Communications*,
752   *5*(1), 1–8.
753

Wilson, K. A., Underwood, E. C., Morrison, S. A., Klausmeyer, K. R., Murdoch, W. W., Reyers,
B., ... & Possingham, H. P. (2007). Conserving biodiversity efficiently: What to do, where,
and when. *PLOS Biology*, *5*(9), e223.

Wyborn, C., & Evans, M. C. (2021). Conservation needs to break free from global priority
mapping. *Nature Ecology & Evolution*, *5*(10), 1322–1324.

Figure captions

Figure 1. Schematic of the study framework, biogeographic zones of India, and themes and
input layers used for spatial prioritization. Theme-wise input layers include: (1) Habitats: 28
vegetation classes depicted as five composite panels; (2) Ecosystem Services: blue water flux,
above- and below-ground carbon, and green water flux; (3) Biodiversity: Protected Areas, Key
Biodiversity Areas, and beta-diversity index of threatened mammals, birds, reptiles and
amphibians; (4) Threats: human population density, livestock population density, urbanization,
linear infrastructure, mines, river fragmentation, agricultural expansion, vegetation greening,
vegetation browning, future climate warming and future rainfall anomaly.

Figure 2. (a) Theme-wise results from prioritization analyses of Habitats (top left), Ecosystem
services (centre left), Biodiversity (bottom left), and Composite of the three themes (right); (b)
Theme-specific results from prioritization analysis of Threats.

Figure 3. (a) Conservation priority ranks for the country, computed zone-wise and combined; (b)
Top 30% priority areas from each zone demarcated as a set of three 10% blocks. Designated
Protected Areas are overlaid to show the extent of overlap and spatial congruence with high
priority locations.

Figure 4. Figure 4. Results from spatial prioritization analyses processed at the scale of
administrative units in the country. Left panel: Proportion of each district covered by areas
identified as top 30% priority locations. Centre panel: Districts in the top quantile (25%)
identified as 169 'high priority' districts and the next 25% quantile identified as 169 'potential
priority' districts. Darker lines denote state boundaries. Right panel: NITI Aayog aspirational
districts (112) earmarked by the Government of India, and their overlap with 338 priority
districts identified in this study (72 districts overlap). Darker lines in the centre and right panels
denote state boundaries.

Supplementary Information

Table S1. Eco-climatic characteristics of the 8 biome-zones considered in this
study. Classification of the biomes are as described by Rodgers and Panwar (1988) This level of

classification splits the country into zones that have some similarity of biogeography, while
across zones, they capture diversity of species, ecosystems, and human–nature relationships.
796
797 **Table S2.** Details and description of data used as input attributes for zone-wise prioritization
798 analyses in India. The table also includes information on the data source, spatial resolution and
799 the corresponding year from which the data were sourced.
800
801 **Table S3.** Ranks assigned to individual *Threat* attributes to reflect their relative severity in terms
802 of impact on habitats, ecosystems services and biodiversity in each biogeographic zone. Ranks
803 were assigned based on a consensus-approach. The assessors (authors of the current study) with
804 relevant expertise for each zone/region together ranked the individual threats for the
805 corresponding zone/region. All assessors reviewed and gave final approval for all threat ranks.
806

proportion of districts with top 30% priority areas

0%   100%

high priority districts   NITI aspirational districts

potential priority districts   overlap districts