Gli2 and *Gli3* Regulate Horizontal Basal Cell-Mediated Regeneration of the Olfactory Epithelium

Anna Shirazyan[1], Haeyoung Park[1], Ariell M. Joiner[2], Justine Ra[1], Melissa S. Kim[3], Archana Kumari[4], Nicole E. Franks[1], Tien Peng[5], Robert K. Bradley[6], Charlotte M. Mistretta[6], Andrzej A. Dlugosz[7,8], and Benjamin L. Allen[1,8]*

[1]Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, MI, [2]Cerenovus, Irvine, CA, [3]Developmental, Stem Cell and Regenerative Biology Program, University of Pennsylvania, Philadelphia, PA, [4]Cell Biology and Neuroscience, Rowan University School of Osteopathic Medicine, Glassboro, NJ, [5]Department of Medicine and Cardiovascular Research Institute, University of California, San Francisco, CA, [6]Department of Biologic and Materials Sciences & Prosthodontics, University of Michigan School of Dentistry, Ann Arbor, MI, [7]Department of Dermatology, University of Michigan, Ann Arbor, MI. [8]Rogel Cancer Center, University of Michigan, Ann Arbor, MI

*Correspondence: benallen@umich.edu (B.L.A.)

Summary

The olfactory epithelium (OE) is a specialized neuroepithelium that is replenished by two stem cell

populations: globose basal cells (GBCs) and horizontal basal cells (HBCs). Previous work indicated that HBCs

contain primary cilia, organelles that mediate Hedgehog (HH) pathway activity. However, a role for HH

signaling in HBCs has not been investigated. We find that GLI2 and GLI3, transcriptional effectors of the HH

pathway, are expressed in HBCs in the adult OE and that their expression expands following injury. Further,

Gli2-expressing descendants contribute to all major OE cell types during OE regeneration. HBC-specific

expression of constitutively active GLI2 drives inappropriate HBC proliferation, alters HBC identity, and

culminates in a failure of HBCs to differentiate into olfactory sensory neurons (OSNs) following injury. HBC-

specific deletion of endogenous *Gli2* and *Gli3* results in decreased HBCs and OSNs following OE injury. These

data identify GLI2 and GLI3 as key regulators of HBC-mediated OE regeneration.

Introduction

Introduction

Olfactory dysfunction affects approximately 13.3 million older Americans (Hoffman et al., 2016). This

can be due to factors that affect regeneration of the olfactory epithelium (OE), including physical injuries or

pathogenic infections. The recent COVID-19 pandemic has resulted in loss of smell (anosmia) in patients (Aziz

et al., 2021). To more effectively treat olfactory disorders, we need to better understand the signals that govern

OE regeneration. The OE is a highly specialized neuroepithelium that contains olfactory sensory neurons

(OSNs), which relay smell information (Graziadei and Graziadei, 1979). OSNs are vulnerable to the

environmental toxins and pathogens present in the air, making the OE one of the few adult cites of neurogenesis

(Graziadei and Graziadei, 1979; Morrison and Costanzo, 1989). Fortunately, two presumed stem cell

populations can replenish OSNs: rapidly dividing globose basal cells (GBCs), and relatively quiescent

horizontal basal cells (HBCs) (Schwob et al., 2017). HBCs lie in a monolayer along the OE basement

membrane, with GBCs situated just above the HBCs. Both HBCs and GBCs can generate neuronal and non-

neuronal progenitors, which give rise to immature OSNs (iOSNs), supporting glial-like Sustentacular cells

(Sus), Bowman's gland, and Microvillar cells (MVCs; (Schwob *et al.*, 2017); Figure 1A). Although HBCs and

GBCs mediate OE regeneration, the signals governing their function in homeostasis and injury-mediated repair

are largely unexplored.

Hedgehog (HH) signaling is necessary for adult stem cell maintenance and function in many tissues,

including multiple different epithelia (Petrova and Joyner, 2014). Notably, HH signaling is necessary for taste

bud maintenance and renewal in the tongue, another chemosensory organ (Mistretta and Kumari, 2019).

Targeted deletion of the HH transcription factor *Gli2* in the lingual epithelium results in taste bud loss and

formation of atypical taste organs (Ermilov et al., 2016). Similar experiments with the HH pathway inhibitor

LDE225 demonstrated a loss of taste buds and taste sensation in mice (Kumari et al., 2015; Kumari et al.,

2017). Given the importance of HH signaling in lingual epithelial regeneration, it is possible that HH signaling

is playing a similar role in another chemosensory epithelium – the olfactory epithelium.

HH signaling is mediated by the primary cilium, a microtubule-based signaling center (Huangfu et al.,

2003). In the absence of HH ligand, the canonical HH receptor Patched 1 (PTCH1) inhibits the activity of the

GPCR-like protein Smoothened (SMO) resulting in phosphorylation and processing/degradation of GLI proteins, the transcriptional effectors of the HH pathway (Briscoe and Therond, 2013). In the presence of HH ligand, HH binding to PTCH1 results in de-repression of SMO, allowing SMO to accumulate in the cilium, resulting in GLIs accumulating at tips of primary cilia, and processing of GLIs into transcriptional activators that induce HH target gene expression (e.g. *Ptch1* and *Gli1*; (Briscoe and Therond, 2013)). GLI1 functions exclusively as a transcriptional activator and is also a target of HH signaling; GLI2 is typically the major transcriptional activator of the HH pathway; conversely GLI3 acts largely as a transcriptional repressor (Briscoe and Therond, 2013). GLI2 and GLI3 contain an N-terminal repressor domain and a C-terminal activator domain, and can be post-translationally processed into either their full length activator form or truncated repressor form (Sasaki et al., 1997; Sasaki et al., 1999). Though GLI2 and GLI3 are typically thought to have opposing roles, they can also play redundant roles depending on the context of the tissue (Chang et al., 2016; McDermott et al., 2005).

Previous work demonstrated that HBCs possess primary cilia and that HBC-specific ablation of primary cilia result in defective OE regeneration, indicating a functional role for primary cilia in HBCs (Joiner et al., 2015). Given that 1) HH signaling is a key regulator of tissue renewal across multiple epithelia, 2) HH signaling also is a key regulator of adult neurogenesis, and that 3) HH/GLI signaling rely on the primary cilium, a structure that is essential for HBC-mediated OE regeneration, we wondered whether HH signaling might play a role in stem cell-mediated adult OE regeneration.

Here, we show that the HH transcription factors *Gli2* and *Gli3* are expressed in HBCs and a subset of Sus cells in the OE. Additionally, we show that *Gli2+* HBCs can give rise to GBCs, OSNs, and Sus cells following injury. Further, we show that expression of constitutively active *Gli2* (GLI2A) in HBCs results in abnormal HBC proliferation and altered cell identity. Further, GLI2A-expressing HBCs fail to differentiate into OSNs following injury. Conditional *Gli2* and *Gli3* deletion in HBCs results in loss of HBCs and OSNs following injury. Together, these data suggest novel roles for GLI2 and GLI3 transcription factors in adult OE regeneration.

<u>Results</u>

Gli2 and Gli3 are expressed in HBCs and expand following OE injury

To investigate the expression of GLI transcription factors in the adult OE we utilized mice carrying *lacZ*

reporter alleles for $Gli1^{lacZ/+}$ (Bai et al., 2002), $Gli2^{lacZ/+}$ (Bai and Joyner, 2001), and $Gli3^{lacZ/+}$ (Garcia et al.,

2010). Specifically, we performed X-GAL staining on coronal sections, focusing on two different regions, the

dorsal septum, and a ventral turbinate (Figure 1A). Like *Gli1*, *Gli2* is also expressed in the stroma underlying

the OE; however, *Gli2* is more broadly expressed in both the proximal and distal stroma of the OE compared to

Gli1 (Figure 1F-F'). In contrast to *Gli1*, *Gli2* is also expressed in basal epithelial cells throughout the OE

(Figure 1F-F', H-H'). Further, *Gli2* is expressed in a subset of apical epithelial cells in the turbinates (Figure

1H-H'; black arrowheads). *Gli3* is also expressed broadly in the underlying stroma, and in basal cells

throughout the OE (Figure 1G-G', I-I'). Distinct from *Gli2*, *Gli3* is expressed in apical epithelial cells both at

the septum (Figure 1G-G'; black arrowheads) and at turbinates (Figure 1I-I';black arrowheads).

To define the cell types that expressed *Gli2* and *Gli3* in the OE, we used immunofluorescent antibody

detection to mark *Gli*-expressing cells (BGAL$^+$) and horizontal basal cells (HBCs; p63$^+$) (Packard et al., 2011).

While no BGAL signal was detected in WT animals (Figure 1K inset), in $Gli2^{lacZ/+}$ mice, BGAL co-localizes

with NP63 (Figure 1K-K'''; yellow arrow). Likewise, in $Gli3^{lacZ/+}$ mice, BGAL also co-localizes with p63

(Figure 1L-L'''; yellow arrow), demonstrating that both *Gli2* and *Gli3* are expressed in HBCs of the adult OE.

Co-staining with BGAL and a sustentacular (Sus) cell marker, SOX2 (Guo et al., 2010), revealed that BGAL

co-localizes with apical SOX2$^+$ cells in the ventral turbinates in both $Gli2^{lacZ/+}$ (Figure S1B-B''') and $Gli3^{lacZ/+}$

mice (Figure S1C-C'''), indicating that *Gli2* and *Gli3* are expressed in distinct subsets of Sus cells in the OE.

Together, these data demonstrate that *Gli2* and *Gli3*, but not *Gli1*, are expressed in the adult OE, displaying

similar, but not identical expression patterns.

To further explore *Gli2* and *Gli3* expression in HBCs, a cell type that is quiescent during homeostasis

(Fletcher et al., 2011; Packard *et al.*, 2011), we sought to activate HBCs with an injury to the OE. Upon delivery

of methimazole to the OE, Sus cells in the OE metabolize methimazole into a toxicant that results in the

destruction of most of the OE (Genter et al., 1995). Notably, HBCs will be spared– they will proliferate and

give rise to both neuronal and non-neuronal lineages, and at 8 weeks will fully regenerate the OE (Leung et al.,

2007). We delivered 75mg/kg of methimazole to $Gli2^{lacZ/+}$ and $Gli3^{lacZ/+}$ via intraperitoneal (IP) injection,

followed by analysis at 4 days of recovery post-injury a time point when HBCs are most active during

regeneration (Packard *et al.*, 2011). We performed *in situ* hybridization detection of *lacZ* transcripts, followed

by immunofluorescent antibody detection of NP63 to mark HBC nuclei. Importantly, no *lacZ* expression is

detected in wild-type uninjured and injured OE, demonstrating specificity of the *lacZ* probe (Figure 2A-C, D-F).

Sparse *lacZ*+ puncta are detected in NP63+ HBCs and stromal cells in uninjured $Gli2^{lacZ/+}$ (Figure 2G-I) and

Gli3$^{lacZ/+}$ (Figure 2M-O) animals. At four days post-injury, however, *Gli2* expression is significantly increased

throughout the entire OE (Figure 2J-L, S) as well as specifically in NP63$^+$ HBCs (Figure 2U). Similarly, *Gli3*

expression is also significantly increased in the OE following injury (Figure 2P-R, T) and in HBCs (Figure 2V).

We confirmed these findings by performing X-GAL staining on $Gli2^{lacZ/+}$ and $Gli3^{lacZ/+}$ mice at 4 days and 8

weeks post methimazole injury (Figure 2S2 A-L'). Notably, X-GAL staining was not detected in WT animals at

either timepoint (Figure S2 A-B', G-H'). Similar to the *in situ* data, increased X-GAL staining was detected in

$Gli2^{lacZ/+}$ mice at 4 days following injury, especially at ventral turbinates compared to the dorsal septum region

(Figure S2C-D'). Since *Gli1* is a transcriptional readout for active HH signaling, we also used an endogenous

Gli1 in situ probe to visualize *Gli1* expression prior to and following injury (Figure S1D-K). *Gli1* expression

was not detected in the OE of uninjured mice, similar to X-GAL staining of $Gli1^{lacZ/+}$ animals (Figure S1E). No

Gli1 expression was detected in the OE 4 days following injury (Figure S1I), although stromal *Gli1* levels are

significantly upregulated (Figure S1J-K). Taken together, these data suggest *Gli2* and *Gli3,* but not *Gli1,* are

upregulated in the OE during early injury recovery.

Gli2 descendants can give rise to all major OE cell types

Since *Gli2* expression in HBCs expands following injury, we employed a genetic lineage tracing

approach to investigate the ability of *Gli2*-expressing cells and their descendants to contribute to different OE

cell types. To achieve this, we bred *Gli2Cre^{ER}* mice with a *ROSA26-lox-STOP-lox-tdtomato* reporter allele

(Madisen et al., 2010; Wang et al., 2018). Coronal sections from the heads of mice treated with vehicle (corn

oil) or tamoxifen (Figure 3A) were immunolabelled with antibodies directed against NP63 (HBCs) and SEC8

(GBCs). After either vehicle or tamoxifen treatment mice were rested for 72 hours then injected with

methimazole to injure the OE (Figure 3A-G''''). We analyzed *Gli2*-expressing descendants at 8 weeks (full

recovery) after injury by immunolabelling for different OE cell types- NP63/CD54 (HBCs), SEC8 (GBCs),

SOX2 (Sus cells), and CBX8 (OSNs). Notably, no tdTomato+ progeny were visible in vehicle treated

Gli2Cre^{ER}; tdTomato/+ mice (Figure 3B-B'''', D-D'''', F-F''''). We detected tdTomato+ labelled HBCs

(Figure 3C'', C''''; white arrowhead) and GBCs (Figure 3C''', C''''; yellow arrowhead) in fully recovered

mice. Further, we detected tdTomato+ Sus cells (Figure 3E-E''''; blue arrowheads) and OSNs (Figure 3G-

G''''; green arrowheads). These findings suggest that *Gli2*+ HBCs can give rise to all major cell types of adult

OE, namely GBCs, Sus cells, and OSNs.

GLI2 activator drives HBC proliferation and alters HBC identity

Given that increased *Gli2* expression correlates with HBC activation and that *Gli2* descendants

contribute to all major OE cell types, we wondered whether the primarily transcriptional activator function

ascribed to GLI2 in other systems (Bai *et al.*, 2002) might drive HBC activation in the OE. To test this notion,

we utilized a doxycycline inducible system to express a constitutively active form of GLI2 lacking the N-

terminal repressor domain (GLI2ΔN) in HBCs. We crossed mice that contain a reverse tetracycline

transactivator driven by a Keratin 5 promoter (Krt5rtTA) (Diamond et al., 2000) to mice containing a tet operon

inducible MYC-tagged *GLI2ΔN* transgene (Grachtchouk et al., 2011) (Figure 4). Mice were fed a doxycycline

diet to induce *GLI2ΔN* expression in HBCs and analyzed at 1 day, 3 days, 5 days, and 8 days following

induction (Figure 4B-G'''', Figure S3A-D). Due to the aggressive nature of driving overactive HH signaling

these mice do not survive past 7-8 days (Grachtchouk *et al.*, 2011). Coronal sections of heads from control and

experimental mice were stained with antibodies directed against MYC (GLI2ΔN), CD54 (HBCs), and Ki67

(actively cycling cells). While no MYC expression was detected in vehicle-treated *Krt5rtTA;GLI2ΔN* mice

(Figure 4B'), MYC immunostaining was detected in HBCs of *Krt5rtTA;GLI2ΔN* mice treated with doxycycline

for 1 day (Figure 4C'). Immunostaining for Ki67 (Figure 4B'', C''), a marker for actively cycling cells,

identifies GBCs lying apical to CD54+ HBCs (Figure 4B''', C'''; white arrowheads) but not HBCs themselves,

indicating that at 1 day post doxycycline induction HBCs are not actively proliferating. In contrast, at 3 days

post doxycycline induction MYC$^+$ HBCs (Figure 4E', E'''; yellow arrowheads) express Ki67 (Figure 4E'';

yellow arrowheads), while control HBCs do not (Figure 4D-D''''). At 5 days post doxycycline induction MYC$^+$

HBCs (Figure 4G', G'''; yellow arrowheads) continue expressing Ki67 (Figure 4G''; yellow arrowheads),

while control HBCs do not (Figure G-G''''). Notably, GLI2ΔN protein is detected in HBC-associated primary

cilia, an important organelle for HH signal transduction, at 8 days following doxycycline administration (Figure

S4 F-J; white arrowhead). HBCs reach peak proliferation at 3 days post doxycycline induction (Figure S3 C-D),

resulting in significantly increased HBC number at 3, 5, and 8 days post doxycycline induction (Figure S3 A,

D). These data suggest that GLI activator can induce HBC proliferation.

To further characterize the consequences of HBC-specific GLI2A expression on the OE, we examined

markers for other cell types, starting with GBCs. Strikingly, and in contrast to control mice (Figure 5A-E), at 8

days post doxycycline induction HBCs co-express SEC8, a pan GBC marker (Joiner *et al.*, 2015) (Figure 5 F-

M). The total number of HBCs (Figure 5K) is significantly increased while the total number of GBCs remains

unchanged (Figure 5 L). Further, we performed immunostaining for SOX2, which labels a subset of HBCs and

GBCs basally and Sus cells apically in control mice (Figure 5N-Q). GLI2A-expressing HBCs also co-express

SOX2 (Figure 5R-V). Immunolabelling for SOX9 (normally a marker for Bowman's gland cells) also revealed

extensive co-labelling with HBCs in *Krt5rtTA;GLI2ΔN* mice (Figure S4O-R), compared to control mice (Figure

S4K-N, S). Interestingly, *Krt5rtTA;GLI2ΔN* HBCs do not co-express the neuronal marker CBX8 (Figure S4T),

and there was no significant change in neurons in *Krt5rtTA;GLI2ΔN* OE (Figure S4U). In contrast, there is a

subtle but significant increase in Sus cell number in mice expressing GLI2A (Figure 5W). Taken together, these

data suggest that constitutive GLI activator function drives altered HBC identity and increased Sus cells.

GLI2 activator promotes HBC differentiation to Sus Cell, but not OSN lineages following OE injury

Next, we assessed the effect of GLI2A on HBC function. Since HBCs are primarily quiescent in

homeostatic conditions (Fletcher *et al.*, 2011; Packard *et al.*, 2011), we sought to activate them by injuring the

OE. After inducing control and *Krt5rtTA;GLI2ΔN* mice with doxycycline for 24 hours, we injured the OE with

an IP injection of methimazole at 75mg/kg (Figure 6, Figure S5). Mice were maintained on a doxycycline diet

following injury and allowed to recover for 7 days (Figure 6A). We first analyzed the basal cells in control and

GLIA mice following methimazole injury (Figure S5A-N). Coronal sections of mice collected at 7 days

following injury were stained with antibodies for different OE markers in addition to Ki67 to mark actively

cycling cells. While only a small fraction of HBCs in control OE are actively cycling (less than 5%)

proliferative (Figure S5A-E, K), GLI2A-expressing HBCs are significantly more proliferative (Figure S5F-K).

As a consequence there is a significant increase in HBC number in injured *Krt5rtTA;GLI2ΔN* mice (Figure

S5L). In contrast, GBC number is significantly downregulated (Figure S5M). As with uninjured animals, we

also observed an upregulation in SOX9+ cells in injured GLI2A-expressing animals (Figure S5N).

To investigate the consequences of HBC-specific GLI2A expression on OE differentiation, including the

formation of neurons and Sus cells, we first immunostained coronal sections of control and *Krt5rtTA;GLI2ΔN*

mice with SOX2 to mark Sus cell progenitors. We discovered that GLI2A-expressing HBCs give rise to a

significantly higher amount of total SOX2$^+$ progenitors, in comparison to control HBCs (Figure 6J). The OE in

control HBCs also appeared more organized with a clear layer of early apical Sus cells separate from basal cells

(Figure 6B-E), whereas the OE in *Krt5rtTA;GLI2ΔN* mice appeared much more disorganized with expanded

SOX2 staining (Figure 6F-I). Additionally, we immunolabelled for neurons using TUJ1 (Figure 6K-S).

Surprisingly, there was a significant decrease in TUJ1$^+$ progenitors in GLI2A OE compared to control OE

(Figure 6S). GLI2A-expressing HBCs give rise to very few neurons (Figure 6O-R) compared to control HBCs

which generate a well-defined TUJ1$^+$ layer of cells (Figure 6K-N). In short, while wild-type HBCs can give rise

to both neurons and Sus cells following injury, HBCs with overactive HH signaling primarily give rise to Sus

cells and not neurons (Figure 6T). This suggests that GLI activator not only drives HBC proliferation, but also

drives HBCs preferentially toward a Sus cell lineage.

GLI2 and GLI3 are necessary for proper OE regeneration following injury

212  To investigate the function of endogenous GLIs in HBCs we utilized a conditional deletion mouse

model to selectively delete *Gli2* and *Gli3* in HBCs. We crossed in an HBC-specific, tamoxifen-inducible *Krt5-*

Cre^{ER} (Diamond *et al.*, 2000) to *Gli2$^{fl/fl}$* (Corrales et al., 2006) and *Gli3$^{fl/fl}$* (Blaess et al., 2008) alleles

surrounded by loxP sites. A *ROSA26-lox-STOP-LOX-tdTomato* (Madisen *et al.*, 2010) allele was also crossed in

to track HBC lineages following methimazole injury. Mice were treated with 100mg/kg of tamoxifen for 5

consecutive days, thus generating *Gli2^{CKO}*, *Gli3^{CKO}*, and *GLI2/3^{CKO}* mice lacking either *Gli2* or *Gli3*

individually, or combination in HBCs. We then performed methimazole injury at 75mg/kg to stimulate HBC

proliferation and differentiation. We collected and analyzed HBC progeny in *Gli2^{CKO}* (Figure S6A-E), *Gli3^{CKO}*

(Figure S6F-J), and *GLI2/3^{CKO}* (Figure 7) mice 8 weeks following methimazole injury (full recovery). We used

the tdTomato reporter to track HBC-derived cells and performed immunolabeling for different OE markers to

determine the effect of HBC-specific conditional *Gli* deletion on OE recovery. *Gli2^{CKO}* OE are comparable to

control OE across all cell types. We observed no significant difference in the numbers of HBCs (Figure S6B),

GBCs (Figure S6C), OSNs (Figure S6D), or Sus cells (Figure S6E). In contrast, *Gli3^{CKO}* animals display subtle

but significant increases in HBCs (Figure S6G), GBCs (Figure S6H), and Sus cells (Figure S6J). The number of

OSNs are slightly but not significantly decreased (Figure S6I). Given that GLI3 largely functions as a

transcriptional repressor in other tissues (Petrova et al., 2013; Sasaki *et al.*, 1997; Sasaki *et al.*, 1999; Wang et

al., 2000), the increase in basal and Sus cells in *Gli3^{CKO}* OE complement our findings with HBC-specific

constitutive GLI2 activator expression (cf. Figure 6).

230  Since *Gli2* and *Gli3* can have redundant functions (McDermott *et al.*, 2005), we sought to delete *Gli2*

and *Gli3* simultaneously in HBCs (*GLI2/3^{CKO}*) (Figure 7). We first confirmed efficient deletion of *Gli2* and *Gli3*

in our genetic mouse model using *in situ* hybridization prior to methimazole injury (Figure S7A-F). *Gli2*

expression was 86% downregulated in *GLI2/3^{CKO}* OE (Figure S7C, E) compared to control mice (Figure S7A,

E). *Gli3* was 74% downregulated in *GLI2/3^{CKO}* OE (Figure S7D, F) compared to control mice (Figure S7B, F).

GLI2/3^{CKO} animals display a range of OE phenotypes following injury- while some regions are grossly

disturbed and thin (Figure S7G, I) others looked normal (Figure S7H, J). Quantitation of basal cell numbers in

GLI2/3^{CKO} OE revealed that HBCs are significantly decreased (Figure 7E-H), compared to control OE (Figure

7B-D, H). GBC numbers are also subtly decreased, although not significantly (Figure 7E-G, Figure S7R). Next,

we co-immunolabelled for TUJ1 and CBX8 to mark OSNs (Figure 7I-O). We discovered that OSNs are

significantly decreased in *GLI2/3^{CKO}* OE (Figure 7L-N, O), compared to control OE (Figure 7I-K, O). Notably,

we observed no change in Sus cell number (Figure S7K-Q), even in more significantly perturbed areas (Figure

S7N inset). These findings suggest that combined HBC-specific *Gli2* and *Gli3* deletion results in reduced HBC

numbers and a subsequent failure to properly regenerate the OE following injury, particularly with regard to

OSN differentiation. Taken together, these data illustrate novel individual and combined roles for GLI

transcription factors in HBC-mediated OE regeneration by regulating the proliferation and differentiation of

HBCs following injury (Figure 7S).

<u>Discussion</u>

Here, we explored the contribution of the HH pathway to OE regeneration. Specifically, we found that

Gli2 and *Gli3*, encoding two key transcriptional effectors of the HH pathway, are expressed in HBCs and

distinct subset of Sus cells in the OE. Further, we discovered that *Gli2* and *Gli3* expression expands following

OE injury. Constitutive GLI2A expression in HBCs results in HBC hyperproliferation and defective

differentiation following OE injury. These data suggest that transient GLI activation following OE injury allows

for HBC activation and proliferation, but that abrogation of GLI activator function is necessary for subsequent

HBC differentiation. HBC-specific conditional *Gli2* deletion does not result in any detectable phenotype, while

HBC-specific conditional *Gli3* deletion results in increased HBC, GBC and Sus cell numbers. Simultaneous

conditional *Gli2* and *Gli3* deletion in HBCs results in improper OE regeneration and significantly decreased

HBC, GBC, and OSN numbers. Together, these data illustrate novel roles for GLI signaling in HBC function

and demonstrate that GLI2 and GLI3 are necessary for OE regeneration post-injury.

Diversity of HH response across different types of epithelia.

HH signaling plays important but diverse roles in many adult regenerative epithelia. Two such examples

are the lingual and respiratory epithelium. HH signaling is essential for proper maintenance and function of

taste organs in the lingual epithelium (Mistretta and Kumari, 2019). SHH ligand is present in the stem cells and

nerves that innervate the taste bud (Ermilov *et al.*, 2016; Liu et al., 2013). As in the OE, *Gli2* is also expressed

in basal cells of the lingual epithelium as well as in the underlying stroma (Liu *et al.*, 2013). *Gli1* is also present

in the lingual stroma and epithelium, which differs from the OE where it is expressed exclusively in the stroma

(Ermilov *et al.*, 2016). GLI2A expression in Keratin5+, *Gli2*+ cells in the tongue results in aberrant

proliferation of lingual epithelial cells (Liu *et al.*, 2013). These findings are remarkably like the OE, where

GLI2A-expressing HBCs become hyperproliferative. Further, the lingual epithelium is dependent on active HH

signaling for maintenance. Chemical HH pathway inhibition using a SMO antagonist LDE225 results in

disrupted taste organs and taste sensation (Kumari *et al.*, 2015). Further studies in the OE with chemical HH

pathway blockade can elucidate if canonical HH signaling is necessary for OE maintenance and regeneration.

HH signaling is also a key component of adult lung regeneration and maintenance (Wang et al., 2019). During

lung homeostasis, SHH ligand in the epithelium signals to surrounding GLI1+ mesenchymal cells (Peng et al.,

2015). Genetic ablation of *Shh* results in mesenchymal cell proliferation and an increase in epithelial cells in the

airway (Peng *et al.*, 2015). Additionally, genetic activation of HH signaling in the lung mesenchyme (through

expression of oncogenic *Smo*) results in an emphysema-like phenotype (Wang *et al.*, 2018). Thus, HH

signaling appears to have a restrictive role and maintains quiescence in the lung. Conversely, HH signaling is

necessary for HBC proliferation and can drive proliferation in the olfactory epithelium. The adult lingual,

respiratory, and olfactory epithelia are all regenerative, diverse, and unique structures. The differential

deployment of HH/GLI signaling in each of these epithelial appears to be one mechanism by which a single

pathway can contribute to the maintenance and regeneration of related, but distinct tissues.

Regulation of GLI expression and function in the olfactory epithelium.

Our data indicate an increase in *Gli2* and *Gli3* expression following injury in the OE. In contrast, there is

no detectable expression of the HH target gene *Gli1* in the OE with *Gli1* expression remains solely stromal even

following OE injury. These data suggest potentially non-canonical roles for GLI2 and GLI3 in the OE. While a

definitive source of HH-producing cells in the OE remains to be elucidated, one potential source includes SHH

in nerves that innervate the OE (Kumari et al., 2017). SHH ligand is also present in the olfactory bulb (OB),

particularly in the glomeruli (Gong et al., 2009). Further, OSNs are able to respond to SHH ligand *in vitro*

(Gong et al., 2009). Since OSNs directly connect the OE with the OB, it is possible that SHH is delivered to

HBCs and the underlying stroma through OSN axonal projections. Another promising source of ligand is IHH,

which is expressed in bone and regulates proliferation and differentiation of chondrocytes (Vortkamp et al.,

1996). The OE and underlying stroma are directly in contact with the bone in the turbinates, making the

turbinates a potential source of IHH. Further careful investigation into *Shh* and *Ihh* expression in the OE and

surrounding tissues is necessary to elucidate if canonical HH signaling is regulates *Gli2* and *Gli3* in the OE.

Since HH target gene expression is not detected in the OE, it is possible that GLI2 and GLI3 function

independently of HH ligands. A possible avenue of GLI regulation in the OE is through the NP63 transcription

factor. NP63 is an important regulator of stemness in HBCs as illustrated in (Packard *et al.*, 2011). NP63

maintains HBC quiescence and transiently turns off following MeBr gas injury to the OE (Packard *et al.*, 2011).

Our data illustrate an upregulation of *Gli2* and *Gli3* immediately following injury to the OE. It is possible that

NP63 could negatively regulate *Gli2* and *Gli3* during OE homeostasis, resulting in upregulation of *Gli2* and

Gli3 when NP63 levels decrease following injury. Previous studies in mammary cancer stem cells illustrated

that NP63 can directly regulate HH pathway genes such as SHH and GLI2 by binding to upstream regulatory

regions (Memmi et al., 2015). Additionally, our data indicates that driving GLI2A results in significant

upregulation of NP63. This suggests a possible feedback loop between GLIs and NP63 expression. Careful

investigation of whether NP63 can bind at *Gli2* and *Gli3* loci in HBCs can further our understanding of GLI

regulation in HBCs.

Crosstalk between Hedgehog and Notch signaling.

While the primary focus of this study is on HH signaling and GLI transcription factors, other signaling

pathways have been described in the adult OE. Specifically, Notch signaling has been implicated in OE

regeneration (Schwob *et al.*, 2017). Importantly, Notch signaling pathway components are present in HBCs,

along with Sus cells and underlying stroma (Herrick et al., 2018; Herrick et al., 2017). Previously work

demonstrated that modulation of various Notch signaling pathway components altered HBC cell fate following

OE injury (Herrick *et al.*, 2018). These findings echo our observations in the OE following GLI2A

overexpression in HBCs. Similarly, in our HH overactivation model HBC cell fate is altered to Sus/non-

neuronal following injury. It is possible that HH and Notch signaling coordinate to dictate HBC cell fates in

response to injury. Along these lines, previous studies have demonstrated that Notch and HH signaling can

work in conjunction in the context of development. For example, Notch signaling directs HH morphogen

activity in the developing ventral spinal cord (Kong et al., 2015). Additionally, the HH co-receptor GAS1

directly binds to NOTCH1 in the developing forebrain, and coordinates HH and Notch signaling in the

neuroepithelium (Marczenke et al., 2021). Future studies examining Notch signaling pathway components in

our genetic models can elucidate cooperation between HH and Notch signaling in HBCs.

Limitations of Study

This study used multiple gain-of-function and loss-of-function mouse genetic approaches to investigate

the roles of GLI2 and GLI3 in HBCs during OE regeneration. Nevertheless, we could broaden our study by

including human tissue in our analysis. Previous studies have successfully isolated single cell transcripts from

mOSNs of human patients (Durante et al., 2020). A similar approach could be utilized by sorting HBCs and

examining HH pathway transcripts in human patients. Understanding the role of HH signaling in human OE can

also open therapeutic avenues for patients with loss of smell.

Though our work focused primarily on HBCs and regeneration, we also discovered that *Gli2* and *Gli3*

are expressed in other cell types in the OE as well. For example, GLI2 and GLI3 are regionally expressed in Sus

cells in the OE, opening a potential avenue of study of HH signaling in Sus cells. There is a precedent for key

developmental pathways playing a role in Sus cell maintenance – for example genetic ablation of Notch2 results

in Sus cell death in post-natal OE (Rodriguez et al., 2008). It is possible that genetic ablation of *Gli2* and *Gli3*

specifically in Sus cells could affect Sus cell function. Additionally, it is possible that Sus cells and HBCs

directly communicate through HH signaling. Similar studies have shown expression of both Notch ligand and

receptors in HBCs and Sus cells, suggesting possible cell-cell communication (Herrick *et al.*, 2017). Using a

similar approach, we could study the expression of additional HH pathway components in Sus cells to assess the

potential for juxtracrine signaling between HBCs and Sus cells. Overall, a closer study of HH signaling in Sus

cells would broaden our understanding of Sus cell function in the adult OE.

The functional role of the underlying stroma, also known as the lamina propria, during OE injury

remains largely unexplored. The stroma is a highly vascularized tissue where fibroblasts, dendritic bundles from

OSNs, and blood vessels are all present. During infection, the stroma is an important source of inflammatory

cytokines and defense against invading parasites (Imamura and Hasegawa-Ishii, 2016). Studies have examined

the stroma in chronic inflammatory rhinosinusitis models of mice and demonstrated that the immune system can

directly affect HBCs and the OE (Chen et al., 2019). Thus, the immune response could be playing a role in OE

regeneration following methimazole-induced injury. Notably, recent work from our lab indicate that GLIs

regulate the immune response in the context of pancreatic cancer (Scales et al., 2022). In this study we

determined that all three *Glis* are expressed in the stroma underlying the OE. These data raise the question of

GLI-dependent stromal contributions to maintenance and regeneration of the OE.

Acknowledgements

We thank members of the Allen lab for their feedback and technical support throughout this study. We thank Dr. Bradley Goldstein (Duke University) for technical advice and valuable feedback on the OE injury experiments. We acknowledge the Department of Cell and Developmental Biology at the University of Michigan and the O'Shea laboratory for access to research equipment. We acknowledge the Biomedical Research Core Facilities Microscopy Core for access to and maintenance the microscopes used for this study. This work was supported by a Ruth L. Kirschstein Predoctoral Individual National Research Service Award (F31 DC018206 to A.S.), Bradley M. Patten Fellowship (to A.S.), Rackham Merit Fellowship (to A.S.), and the Hearing, Balance, and Chemical Senses Training Program (T32 DC00011, University of Michigan to A.S.). This work was also supported by the National Institutes of Health (R01 DC014428 to B.L.A., R.K.B., A.A.D., C.M.M.).

Author Contributions

Conceptualization, A.S., A.M.J., and B.L.A.; Methodology, A.S., A.M.J., and N.E.F., and B.L.A.; Investigation, A.S., H.P., A.M.J., J.R., M.S.K., and A.K.; Formal Analysis, A.S.; Writing – Original Draft, A.S.; Writing – Review & Editing, A.S. and B.L.A.; Funding Acquisition, A.S., R. B., A. A. D., C.M.M., and B.L.A.; Resources, T.P., A.A.D., and B.L.A.; Supervision, B.L.A.

Declaration of Interests

Authors declare no competing interests.

 Figure Legends

Figure 1. *Gli2* and *Gli3* are expressed in horizontal basal cells of the adult olfactory epithelium. (A)

Cartoon of a sagittal view of the adult mouse head (left). Coronal view of a section through the adult mouse

olfactory epithelium (middle). Schematic of OE cell types (right), including horizontal basal cells (HBCs; blue),

globose basal cells (GBCs; orange), immature olfactory sensory neurons (iOSNs; green), mature olfactory

sensory neurons (mOSNs; red), sustentacular cells (Sus; purple), and microvillar cells (MVCs; magenta). (B-I)

X-gal staining of coronal sections from adult wildtype (B, B', D, D'), *Gli1*$^{lacZ/+}$ (C, C', E, E'), *Gli2*$^{lacZ/+}$ (F, F',

H, H'), and *Gli3*$^{lacZ/+}$ (G, G', I, I') reporter mice. Images are taken from both the dorsal septum (B-C', F-G')

and a ventral turbinate (D-E', H-I'). Black arrowheads indicate BGAL$^+$ apical cells in *Gli2*$^{lacZ/+}$ (H') and

Gli3$^{lacZ/+}$ (G'. I') mice. (J-M) Antibody detection of β-galactosidase (BGAL, green; J', K', L', M') and the HBC

nuclear marker NP63 (magenta; J", K", L). DAPI denotes nuclei (gray; J, K, L, M). NP63 (magenta) and BGAL

co-localization (green) in *Gli2*$^{lacZ/+}$ animals (K'''; yellow arrowhead) and *Gli3*$^{lacZ/+}$ (L'''; yellow arrowhead).

Note that *Gli3*, but not *Gli2*, is expressed in apical cells at the septum of the adult OE (L'; L''''; white

arrowhead). Inset in K denotes the lack of BGAL detection in wildtype animals. Scale bars, B, B', J = 50μm.

Figure 2. Selective upregulation of *Gli2* and *Gli3* expression in mice following methimazole-induced OE

injury. (A-V) Adult WT, *Gli2*$^{lacZ/+}$ and *Gli3*$^{lacZ/+}$ reporter mice were either collected at 6-8 weeks of age or

injured with a 75mg/kg IP injection of methimazole. Coronal sections were collected from the heads of either

uninjured (A-C, G-I, M-O) or 4 days post methimazole injury mice (D-F, J-L, P-R). *In situ* hybridization

detection of *lacZ* transcripts in sections from WT, *Gli2*$^{lacZ/+}$ and *Gli3*$^{lacZ/+}$ reporter mice prior to and following

injury (yellow; B, H, N, E, K, Q). Inverted images of *lacZ* transcripts (black; C, I, O, F, L, R). Co-

immunostaining with NP63 (blue; B, H, E, K, Q) demarcates HBCs. DAPI denotes nuclei (A, D, G, J, M, P),

scale bar 50μm (A) and 25μm (B). Quantitation of *lacZ* puncta in the OE in *Gli2*$^{lacZ/+}$ (S) and *Gli3*$^{lacZ/+}$ (T)

reporter mice prior and following methimazole injury. Quantitation of *lacZ*+ puncta in HBCs from *Gli2*$^{lacZ/+}$ (U)

and *Gli3*$^{lacZ/+}$ (V) reporter mice prior and following methimazole injury. N=2 uninjured *Gli2*$^{lacZ/+}$, n=3 injured

Gli2$^{lacZ/+}$, n=3 uninjured *Gli3*$^{lacZ/+}$, n=3 injured *Gli3*$^{lacZ/+}$. Data are mean ± standard deviation. P-values were

determined by a two-tailed unpaired t-test; n.s.= not significant (p>0.05). Scale Bar=50μm (A), 25μm (B).

Figure 3. *Gli2*-descendants give rise to GBCs, Sus cells, and neurons following injury. (A) Cartoon of *Gli2*

lineage tracing model using a tamoxifen inducible *Gli2Cre*^{ER} mouse to mark *Gli2*-expressing cells. Mice

carrying *Gli2Cre*^{ER}*; tdTomato* alleles were given an IP injection with either vehicle (corn oil) or tamoxifen for 5

consecutive days at 100mg/kg. Upon tamoxifen administration, Cre recombinase excised the stop codon

preceding the tdTomato gene, labelling *Gli2*-expressing cells and their descendants red. Mice were then injured

with 75mg/kg methimazole and collected 8 weeks following injury (B-G''''). Coronal sections of vehicle- and

tamoxifen-treated mice were stained with antibodies directed against NP63 label HBCs (magenta; B', C'; white

arrowheads) and SEC8 mark GBCs (cyan; B''', C'''; yellow arrowheads). Antibodies directed against SOX2

label Sus cells apically (magenta; D', E'; blue arrowheads) while antibodies directed against CD54 label HBCs

(cyan; D''', E''', F''', G'''). Antibodies directed against CBX8 denote neurons (magenta; F', G'; green

arrowheads). Cells originating from a *Gli2* lineage are labelled with tdTomato (red; B'', C''. D'', E'', F'', G'').

Merged images of NP63, tdTomato and SEC8 (B'''', C''''), SOX2, tdTomato, CD54 (D'''', E''''), and CBX8,

tdTomato, and CD54 (F'''', G''''). DAPI denotes nuclei (gray; B, C, D, E, F, G), scale bar (B, D, F) = 50μm.

Figure 4. Constitutively active GLI2 drives HBC proliferation. (A) Cartoon of *GLI2ΔN* transgene activation

using a doxycycline-inducible mouse model. Upon doxycycline (DOX, green) administration, mice carrying a

Keratin 5 reverse tetracycline transactivator (*Krt5rtTA*, blue) transgene and a tet-regulated MYC-tagged

GLI2ΔN transgene (pink) express constitutively active *GLI2ΔN* specifically in HBCs of the OE by the rtTA

(brown) protein binding at the tet operon (oragnge). (B-G'''') Antibody detection of MYC (magenta, B'-G'),

Ki67 (red, B''-G''), and CD54 (cyan, B'''-G''') at 1 Day Vehicle treatment (B-B''''), 1 Day DOX treatment

(C-C''''), 3 days DOX treatment (D-D'''', E-E''''), and 5 days DOX treatment (F-F'''', G-G''''). DAPI

denotes nuclei (gray; B-G). Merged MYC, Ki67, and CD54 images (B''''-G''''). MYC::GLI2ΔN protein is

detectable at 1 day following DOX treatment (magenta, C', C''''; white arrowheads). Following 3 days of DOX

administration HBCs express Ki67 (red, E'', E''''; yellow arrowheads). Ki67 expression in HBCs persists

following 5 days of DOX administration (red, G'', G''''; yellow arrowheads). Scale bar (B) = 50μm.

Figure 5. Constitutively active GLI2 alters HBC identity. (A-J) Coronal sections from adult mice treated

with DOX for 8 days were analyzed with immunostaining using different OE markers. Antibodies directed

against SEC8 (red; B, G) identify GBCs, while antibody detection of CD54 (cyan; C,H) and NP63 (magenta;

D,I) denotes HBCs. DAPI indicates nuclei (gray; A, F). Merged SEC8, CD54, and NP63 images (E, J).

Quantitation of HBCs (NP63$^+$, SEC8$^-$; K), GBCs (NP63$^-$,SEC8$^+$; L), and mixed identity HBCs (NP63$^+$, SEC8$^+$;

M). n=8 control and n=8 *Krt5rtTA*;*GLI2ΔN* animals analyzed (K-M). (N-U) Coronal sections from adult mice

treated with DOX for 8 days were analyzed with immunostaining using different OE markers. Antibody

detection of CD54 (cyan; O,S) and apical SOX2 (Sus cell marker, magenta; P,T). DAPI indicates nuclei (N, R).

Merged CD54 and SOX2 images (Q,U). Quantitation of SOX2$^+$ HBCs (SOX2$^+$, CD54$^+$; V) and SOX2$^+$ Sus

cells (Apical SOX2$^+$ cells; W). n=6 control and n=7 *Krt5rtTA*; *GLI2ΔN* animals analyzed (V-W). Scale bars (A,

N) = 50μm. Control refers to mice containing either the *Krt5rtTA* or *GLI2ΔN* transgene. Data are mean ±

standard deviation. P-values were determined by a two-tailed unpaired t-test; n.s.= not significant (p>0.05).

Figure 6. GLI2A-expressing HBCs fail to contribute to neuronal lineages following injury. (A) Cartoon of

GLI2ΔN transgene activation using a doxycycline-inducible mouse model. Upon doxycycline (DOX, green)

administration, mice carrying a Keratin 5 reverse tetracycline transactivator (*Krt5-rtTA*, blue) transgene and a

tet-regulated MYC-tagged *GLI2ΔN* transgene (pink) express constitutively active *GLI2ΔN* specifically in HBCs

of the OE. 1 day following doxycycline induction, mice are injured with a 75mg/kg intraperitoneal (IP)

injection of methimazole. Following methimazole injury, mice are kept on doxycycline and collected 7 days

following injury. (B-S) Coronal sections from adult mice treated with DOX for 8 days, 7 days following

methimazole injury, were analyzed with immunostaining using different OE markers. Antibodies directed

against SOX2 (magenta; C, G) denote apical lying Sus cells and a subset of basal HBCs and GBCs, while

antibody detection of CD54 (cyan; D, H, M, Q) exclusively identifies HBCs. Merged SOX2 and CD54 images

(E, I). DAPI indicates nuclei (gray; B, F, K, O). Quantitation of the percentage of SOX2$^+$ cells in the OE (J).

Antibodies directed against TUJ1 (red; L, P) denote neurons. Merged TUJ1 and CD54 images (N, R).

Quantitation of the percentage of TUJ1$^+$ cells in the OE (S). Scale bar (B, K), 50μm. n=3 control and n=5

Krt5rtTA; *GLI2ΔN* animals analyzed. Control refers to mice containing either the *Krt5rtTA* or *GLI2ΔN*

transgene. Data are mean ± standard deviation. P-values were determined by a two-tailed unpaired t-test; n.s.=

not significant (p>0.05). (T) Summary of HBC-mediated OE reconstitution at 7 days following injury in wild

type (left) and *Krt5rtTA*;*GLI2ΔN* mice (right).

Figure 7. Simultaneous HBC-specific *Gli2* and *Gli3* deletion results in defective OE regeneration. (A)

Cartoon of tamoxifen-inducible HBC-specific *Gli2* and *Gli3* deletion. Mice carrying a *Krt5-Cre^{ER}* allele were

crossed with mice containing *Gli2$^{fl/fl}$* and *Gli3$^{fl/fl}$* alleles as well as a *tdTomato* reporter allele. Tamoxifen

administration induces Keratin 5 (*Krt5*)-expressing cells to drive *Gli2* and *Gli3* deletion and *tdTomato*

expression. After tamoxifen administration (I.P., 100mg/kg for 5 consecutive days), mice were rested for 72

hours, then injured with methimazole (I.P., 75mg/kg) and allowed to recover for 8 weeks. (B-O) Coronal

sections from adult mice collected 8 weeks following injury were analyzed with immunostaining for various OE

markers. Antibodies directed against NP63 (magenta; D, G) mark HBCs, while antibodies directed against

SEC8 (cyan; D, G) denote GBCs. Antibodies directed against Tuj1 (magenta; K, N) and CBX8 (cyan; K, N)

mark OSNs. DAPI denotes nuclei (gray; B, E, I, L), scale bar (B, I), 50μm. Quantitation of HBC (H) and OSN

(O) number per mm of OE. n=4 control and n=5 *Krt5-Cre^{ER}*; *tdTomato/+*; *Gli2$^{fl/fl}$*; *Gli3$^{fl/fl}$* animals analyzed.

Endogenous tdTomato reporter immunofluorescence (red; C, F, J, M) demarcates HBCs and their lineages.

Control animals refers to *Krt5-Cre^{ER}*;*tdTomato/+*;*Gli2$^{fl/+}$*;*Gli3$^{fl/+}$* mice or mice lacking the *Krt5-Cre^{ER}* allele (H,

O). Data are mean ± standard deviation. P-values were determined by a two-tailed unpaired t-test; n.s.= not

significant (p>0.05).

Materials and Methods

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-p63-alpha (D2K8X) (1:500)	Cell Signaling	13109T
Goat anti-CD54 (ICAM1) (1:100)	R&D Systems	AF796
Mouse anti-SEC8 (1:100)	BD Biosciences	610659
Mouse anti-Tuj1 (1:2500)	Promega	G7121
Mouse anti-MYC Epitope (1:500)	DSHB	9E 10
Rabbit anti-Ki67 (1:500)	Abcam	ab15580
Rabbit anti-CBX8 (1:500)	Cell Signaling	14696
Chicken anti-Beta-Galactosidase (1:2500)	Abcam	ab134435
Rabbit anti-SOX2 (1:2000)	Seven Hills Bio	WRAB-1236
Rabbit anti-SOX9 (1:500)	Millipore	AB5535
Rabbit anti-ARL13B (1:2000)	Proteintech	17711-1-AP
Biotin-Donkey anti-Rabbit (1:500)	Jackson Immuno	711-065-152
Biotin-Donkey anti-Mouse (1:500)	Jackson Immuno	715-065-150
Alexa488-Donkey anti-Chicken (1:500)	Jackson Immuno	703-545-155
Alexa488-Donkeyanti-Goat (1:500)	Invitrogen	A11055
Alexa488-Donkeyanti-Mouse (1:500)	Invitrogen	A21202
Alexa555-Donkey anti-Goat (1:500)	Invitrogen	A21432
Alexa555-Donkey anti-Mouse (1:500)	Invitrogen	A31570
Alexa647-Donkey anti-Rabbit (1:500)	Invitrogen	A31573
Alexa647-Donkey anti-Goat (1:500)	Invitrogen	A11055
Chemicals, peptides, and recombinant proteins		
Tamoxifen	Sigma	T5648
2-mercapto-1-methylimdazole (Methimazole)	Sigma	301507
Normal Donkey Serum	Jackson Immuno	NC9624464
Doxycycline	Thermo Fisher	BP26535
Doxycycline Chow (1g/kg)	Bio-Serv	F3949
EDTA	EMD/VWR	4005
Sucrose	EMD/VWR	8550
Paraformaldehyde	Thermo Fisher	50980489
DAPI	Thermo Fisher	D1306
BSA	Sigma	A7906
L-lysine	Sigma	W384720
Sodium Periodate	Sigma	311448
Sodium Monobasic phosphate (Na_2HPO_4)	Sigma	SX0709
Sodium Dibasic phosphate (Na_2HPO_4)	Sigma	8210-OP
Igepal (NP-40)	Sigma	I8896
$K_3Fe(CN)_6$	Sigma	PX1455
$K_4Fe(CN)_6$	Sigma	P9387
$MgCl_2$	VWR	0288
NaCl	Sigma	SX0420-3
Na deoxycholate	VWR	SX0480-2
Permount	Thermo Fisher	SP15100
Triton X-100	VWR	9410
Tween-20	VWR	9480

X-gal	Goldbio	X4281C
Xylenes	VWR	XX00555
EtOH	Thermo Fisher	04355222
OCT	Thermo Fisher	23730571
Antigen Unmasking Solution, Tris-Based	Vector	H-3301-250
Critical commercial assays		
Alexa-488 Tyramide SuperBoost Kit, streptavidin	Thermo Fisher	NC1439037
RNAscope Multiplex Fluorescent Reagent Kit v2	ACD	323100
Experimental models: Organisms/strains		
Mouse: $Keratin5^{CreERT2}$	(Indra et al., 1999)	JAX: 018394
Mouse: $Gli2Cre^{CreERT2}$	(Wang *et al.*, 2018)	
Mouse: $Gli1^{lacZ/+}$	(Bai *et al.*, 2002)	JAX: 008211
Mouse: $Gli2^{lacZ/+}$	(Bai and Joyner, 2001)	JAX: 007922
Mouse: $Gli3^{lacZ/+}$	(Garcia *et al.*, 2010)	MGI: 4838603
Mouse: $Gli2^{fl}$	(Corrales *et al.*, 2006)	JAX: 007926
Mouse: $Gli3^{fl}$	(Blaess *et al.*, 2008)	JAX: 008873
Mouse: Smo^{fl}	(Long et al., 2001)	JAX: 004526
Mouse: $ROSA26fl$-$STOP$-fl-$tdTomato$	(Madisen *et al.*, 2010)	JAX: 007908
Mouse: $Keratin5rtTA$	(Diamond *et al.*, 2000)	MGI: 4867436
Mouse: $tetOGli2\Delta N$	(Grachtchouk *et al.*, 2011)	
Software and algorithms		
Prism 9	Graph Pad	https://www.graphpad.com/scientific-software/prism/
FIJI	FIJI	https://imagej.net/software/fiji/
In Situ Hybridization Probes		
RNAscope Ecoli-lacZ probe	ACD	313451
RNAscope Probe Ms-Gli1	ACD	311001
RNAscope Probe Ms-Gli2E7E8	ACD	412101
RNAscope Probe Ms-Gli3	ACD	445511

Animals and breeding

All mice were maintained on a mixed BL/6, 129, and CD1 genetic background. $Gli1^{lacZ/+}$ {Bai, 2002 #3},

$Gli2^{lacZ/+}$ (Bai and Joyner, 2001), and $Gli3^{lacZ/+}$ (Garcia *et al.*, 2010) mice have all been described previously.

Keratin5rtTA;tetOGli2ΔN (Grachtchouk et al., 2000) mice were provided by Dr. Andrzej Dlugosz (University

of Michigan, Ann Arbor MI). $Gli2Cre^{ERT2}$ mice were provided by Dr. Tien Peng (University of California, San

Francisco). HBC-specific *Gli2* and *Gli3* deletion was accomplished by breeding animals carrying a

$Keratin5Cre^{ERT2}$ allele with mice carrying *Gli2* and *Gli3* alleles flanked by loxP sites. To lineage trace progeny

from HBCs that underwent *Cre*-mediated recombination, we crossed in a *ROSA26fl-STOP-fl-tdTomato* allele

(JAX: 007908) into all conditional deletion mouse lines (*Gli2^{fl}* and *Gli3^{fl}*). For all experiments male and female

adult mice (6-8 weeks of age) were used. All animal procedures were reviewed and approved by the

Institutional Animal Care and Use Committee (IACUC) at the University of Michigan.

Tamoxifen preparation and administration

Tamoxifen was dissolved in sterile corn oil at 55°C for approximately 2-3 h at 40mg/mL with occasional

agitation until in solution. For all experiments with conditional deletion mice (*ROSA26fl-STOP-fl-tdTomato*,

Gli2^{fl}, and *Gli3^{fl}*) mice were injected 100mg/kg of tamoxifen intraperitoneally (I.P.) for 5 consecutive days.

Mice were allowed to rest for 72 h prior to methimazole lesion (described below).

Doxycycline preparation and administration

Doxycycline powder was dissolved at 200μg/ml in 5% sucrose in autoclaved water and administered to mice

through drinking water. *Keratin5rtTA;tetOGli2ΔN* mice were given doxycycline water during first three days of

doxycycline chow treatment. Doxycycline chow (1g/kg, Bio-Serv #F3949) was administered to

Keratin5rtTA;tetOGli2ΔN mice at 6-8 weeks until date of euthanasia.

Methimazole lesion

Methimazole (2-mercapto-1-methylimdazole) was dissolved in sterile 1X PBS and administered to control and

experimental mice through an intraperitoneal (IP) injection at 75mg/kg following either tamoxifen or

doxycycline treatment.

X-gal staining

Mice were anesthetized with 30% isoflurane, transcardially perfused with 2% PLP solution (2%

paraformaldehyde, 0.01M sodium periodate, 0.01M monobasic and dibasic phosphates, and 90mM L-lysine as

described in (Packard *et al.*, 2011), and decapitated. Heads were post-fixed in 2% PLP for 1hr at 4 °C. Tissue

was then decalcified in 0.5 M EDTA overnight at 4°C; cryoprotected in 10% (1h), 20% (1h), and 30% sucrose

overnight at 4°C; and frozen in OCT compound. Coronal sections of the olfactory epithelium and olfactory bulb

(OB) were cut at 12μm thickness on a Epredia™ Microm HM525 NX Cryostat. Sections were stored at -80 °C

until the day of staining. Beta-galactosidase activity was detected with X-gal staining solution (5mM

$K_3Fe(CN)_6$, 5mM $K_4Fe(CN)_6$, 2mM $MgCl_2$, 0.01% Na deoxycholate, 0.02% NP-40, 1mg/mL X-gal). Sections

were incubated with X-gal staining solution overnight in 37 °C. After staining, sections were washed 3 x 5 min

with 1x PBS, pH 7.4, counterstained with nuclear fast red for 5 min and dehydrated in an ethanol series (70%

ethanol, 95% ethanol, 100% ethanol and 100% Xylenes) followed by application of coverslips with permount

mounting media. Sections were visualized on a Nikon E-800 Upright Widefield Microscope.

Immunohistochemistry

Mice were anesthetized with 30% Fluriso (isoflurane, VetOne), transcardially perfused with 4%

paraformaldehyde (PFA), and decapitated. Heads were post-fixed in 4% PFA for 20–24 h at 4°C. Tissue was

then decalcified in 0.5 M EDTA overnight at 4°C; cryoprotected in 10% (1h), 20% (1h), and 30% sucrose

overnight at 4°C; and frozen in OCT compound. 12⌈m thick coronal sections of the olfactory epithelium and

olfactory bulb (OB) were collected on a Epredia™ Microm HM525 NX Cryostat. Sections were stored at -80°C

until day of staining. On the day of staining, sections were baked at 70°C for 10 min, washed 3 x 5 min with 1X

PBST (0.01% Triton X), pH 7.4, then immediately placed in TRIS antigen retrieval solution for 15 minutes in a

92°C water bath. Sections were then blocked with 10% donkey serum for 1h at RT then incubated with primary

antibody overnight at 4°C. The next day sections were washed with 3 x 5 min with 1X PBST (0.01% Triton X),

pH 7.4, then incubated with secondary antibody for 1h at RT, For some antibodies (SEC8, CBX8) TSA

amplification was used to detect signal. Nuclei were labeled with DAPI for 10 min at room temperature (RT)

and slides were mounted with coverslips using Immu-mount aqueous mounting medium. Sections were

visualized on a Leica upright SP5X confocal microscope.

In situ hybridization

Mice were anesthetized with 30% Fluriso (isoflurane, VetOne), transcardially perfused with 10% neutral

buffered formalin (10% NBF), and decapitated. Heads were post-fixed in 10% NBF for 24 h at RT. Tissue was

then decalcified in 0.5 M EDTA overnight at 4°C; cryoprotected in 10% (1h), 20% (1h), and 30% sucrose

overnight at 4°C; and frozen in OCT compound. Coronal sections of the olfactory epithelium and olfactory bulb

(OB) were cut at 12μm thickness on a Epredia™ Microm HM525 NX Cryostat. Sections were stored at -80 °C

until the day of staining when sections were baked at 70°C for 10 min then washed 5 min in 1X PBST (0.01%

Triton X), pH 7.4. Fluorescent *in situ* hybridization was performed using ACD RNAscope Multiplex

Fluorescent Reagent Kit v2, according to manufacturer's instructions (ACD 323100-USM). Pretreatment

conditions were optimized for the olfactory epithelium: antigen retrieval was performed for 15 min, and

protease treatment was performed for 1 min. Following the RNAscope assay, slides were incubated in primary

antibody and followed the immunohistochemistry staining protocol described above. Sections were visualized

on a Leica upright SP5X confocal microscope.

Quantitation and Statistical analysis

All the data are represented as mean ± standard deviation. All statistical analyses were performed using

GraphPad statistic calculator (GraphPad Software, La Jolla California USA, www.graphpad.com). Statistical

significance was determined using two-tailed Student's t-test. Significance was defined according GraphPad

Prism style: non-significant ($p > 0.05$) and significant ($p \leq 0.05$). For all the experimental analyses a minimum of

3 mice of each genotype were examined; each n represents a mouse. To quantify cell counts all images were

blinded and four separate areas of the OE (dorsal septum, dorsal turbinate, ventral septum, and ventral

turbinate) were averaged per n. All the statistical details (statistical tests used, statistical significance and exact

value of each n) for each experiment are specified in the figure legends.

 <u>Supplemental Figure Legends</u>

Figure S1. *Gli2* and *Gli3* are expressed in turbinate-associated sustentacular cells, while *Gli1* expression is stromally restricted, even following OE injury. (A-B) Coronal sections of adult WT, *Gli2$^{lacZ/+}$*, and *Gli3$^{lacZ/+}$* mice were stained with antibodies against β-galactosidase (green; A'-C') and a marker for Sus cells, SOX2 (red; A''-C''). DAPI denotes nuclei (gray; A-C). Scale bar= 50μm. (D-K) Adult mice were either collected at 6-8 weeks of age or injured with a 75mg/kg IP injection of methimazole. Coronal sections of either uninjured OE or OE 4 days following injury were processed for *in situ* hybridization with a probe to detect endogenous mouse *Gli1* (yellow; E, G). Antibodies directed against NP63 demarcate HBCs (blue; E, G). Inverted images of the *mGli1* probe in OE (black; H, J) and stroma (black; I, J). DAPI denotes nuclei (D, F), scale bars, 50μm (A, D), 25μm (H). Quantitation of *mGli1* puncta in the stroma underlying the OE, n=4 uninjured, n=4 injured (K). Data are mean ± standard deviation. P-values were determined by a two-tailed unpaired t-test; n.s.= not significant (p>0.05).

Figure S2. *Gli2* expression is up-regulated in ventral turbinate-associated OE at 4 days following methimazole injury. Adult wild type, *Gli2$^{lacZ/+}$*, and *Gli3$^{lacZ/+}$*mice were injured with 75mg/kg methimazole and allowed to recover for either 4 days (A-F') or 8 weeks for full recovery (G-L'). X-GAL staining of coronal sections from adult wildtype (A-B', G-H'), *Gli2$^{lacZ/+}$* (C-D', I-J'), and *Gli3$^{lacZ/+}$* (E-F', K-L') reporter mice. Images are taken from both the dorsal septum (A, A', C, C', E, E', G, G', I, I', K, K') and a ventral turbinate (B, B', D, D', F, F', H, H', J, J', L, L').

Figure S3. Quantitation of GLI2A-mediated HBC proliferation. (A) Quantitation of HBC number per millimeter (mm) of olfactory epithelium at 1 Day, 3 Days, 5 Days, and 8 Days following doxycycline induction. (B) Quantitation of the percentage of Ki67$^+$ HBCs of total HBCs in the olfactory epithelium at 1 Day, 3 Days, 5 Days, and 8 Days following doxycycline induction. n=3 control, 3 *Krt5rtTA*; *GLI2ΔN* animals analyzed for 1 Day DOX. (C) Quantitation of Ki67$^+$ HBCs per mm of olfactory epithelium at 1 Day, 3 Days, 5 Days, and 8 Days following doxycycline induction. n=4 control, n=5 *Krt5rtTA*; *GLI2ΔN* animals analyzed for 3 Days DOX.

n=3 control, 4 *Krt5rtTA*; *GLI2ΔN* animals analyzed for 5 Days DOX. n=6 control, 7 *Krt5rtTA*; *GLI2ΔN* animals

analyzed for 8 Days DOX. Control refers to mice containing either the *Krt5rtTA* transgene or *GLI2ΔN*

transgene. Data are represented as the mean ± standard deviation. P-values were determined by a two-tailed

unpaired t-test; n.s.= not significant. (D) Summary graph of HBC dynamics in *Krt5rtTA*; *GLI2ΔN* animals, red

indicates the number of HBCs per mm OE, green indicates the number of Ki67$^+$ HBCs per mm OE, and blue

indicates the percentage of Ki67$^+$ HBCs from total HBCs in the OE.

Figure S4. Constitutively Active GLI2 localizes to primary cilia in HBCs and increases SOX9 expression.

(A-U) Mice containing the *Krt5rtTA* and *GLI2ΔN* transgenes were treated with doxycycline for 8 days. Coronal

sections from adult mice were treated with antibodies directed against MYC (magenta; B, G; white arrowhead)

which detects GLI2ΔN, Arl13b (red; C, H; white arrowhead) which labels primary cilia, and CD54 (cyan; D, I)

which denotes HBCs. Merged MYC, Arl13b, CD54 images (E, J). DAPI denotes nuclei (A, F), scale bar (A)

50μm. (K-R) Coronal sections from adult mice treated with DOX for 8 days were analyzed with

immunostaining using different OE markers. Antibodies directed against CD54 (cyan; L, P) denote HBCs,

while antibody detection of SOX9 (magenta; M, Q) identifies Bowman's gland cells. DAPI indicates nuclei

(gray; K, O). Scale bar (A), 50μm. Merged CD54 and SOX9 images (N, R). Quantitation of the percentage

SOX9$^+$ HBCs, n=3 control and n=3 *Krt5rtTA*; *GLI2ΔN* animals analyzed (S). Quantitation of the percentage of

CBX8$^+$ HBCs, n=5 control and n=5 *Krt5rtTA*; *GLI2ΔN* animals analyzed (T). Quantitation of mOSNs, n=5

control and n=5 *Krt5rtTA*; *GLI2ΔN* animals analyzed (U). Control refers to mice containing either the *Krt5rtTA*

or *GLI2ΔN* transgene. Data are mean ± standard deviation. P-values were determined by a two-tailed unpaired t-

test; n.s.= not significant (p>0.05).

Figure S5. Constitutively Active GLI2 drives HBC proliferation following injury. (A-J) Coronal sections

from adult mice treated with DOX for 8 days, 7 days following methimazole injury, were analyzed with

immunostaining using different OE markers. Antibodies directed against MYC (magenta; B, G) denote MYC

tagged GLI2ΔN, while antibodies directed against CD54 (cyan; C, H) stain HBCs. Actively cycling cells are

stained with Ki67 (red; D, I), DAPI denotes nuclei (gray; A, F), MYC, CD54, Ki67 merged images (E, J). Scale

bar (A), 50μm. Quantitation of the percentage of Ki67$^+$ HBCs (K). Quantitation of the number of NP63+ HBCs

(L), SEC8+ GBCs (M), and SOX9+ cells (N) per mm of OE. n=3 control and n=5 *Krt5rtTA*; *GLI2ΔN* animals

analyzed. Control refers to mice containing either the *Krt5rtTA* or *GLI2ΔN* transgene. Data are mean ± standard

deviation. P-values were determined by a two-tailed unpaired t-test; n.s.= not significant (p>0.05).

Figure S6. HBC-specific individual *Gli3*, but not *Gli2* deletion results in improper OE regeneration. (A, F)

Cartoon of HBC-specific individual deletion of either *Gli2* or *Gli3*. Mice containing a *Krt5-Cre^{ER}* Cre allele

were crossed with mice carrying either *Gli2$^{fl/fl}$* (A) or *Gli3$^{fl/fl}$* (F), alleles in addition to a *tdTomato* reporter

allele. Upon tamoxifen administration (I.P., 100mg/kg for 5 consecutive days), mice were rested for 72 hours,

then injured with methimazole (I.P., 75mg/kg) and allowed to recover for 8 weeks. Quantitation of HBCs (B),

GBCs (C), OSNs (D), and Sus Cells (E) per mm of OE. n=5 control and n=6 *Krt5-Cre^{ER}*; *tdTomato/+*; *Gli2$^{fl/fl}$*

animals analyzed. Control refers to *Krt5-Cre^{ER}*; *tdTomato/+*; *Gli2$^{fl/+}$* mice or mice lacking the *Krt5-Cre^{ER}* allele.

Quantitation of HBCs (G), GBCs (H), OSNs (I), and Sus Cells (J) per mm of OE. n=7 control and n=6 *Krt5-*

Cre^{ER}; *tdTomato/+*; *Gli3$^{fl/fl}$* animals analyzed. Control refers to *Krt5-Cre^{ER}*; *tdTomato/+*; *Gli3$^{fl/+}$* mice or mice

lacking the *Krt5-Cre^{ER}* allele. Data are mean ± standard deviation. P-values were determined by a two-tailed

unpaired t-test; n.s.= not significant (p>0.05).

Figure S7. Simultaneous HBC-specific *Gli2* and *Gli3* deletion results in defective OE regeneration. (A-E)

Coronal sections from adult *Krt5-Cre^{ER}*; *tdTomato/+*; *Gli2$^{fl/fl}$*; *Gli3$^{fl/fl}$* mice collected after 5 days of consecutive

tamoxifen injections (I.P., 100mg/kg). Sections were treated with mRNA probes directed against endogenous

Gli2 (yellow; A, C) and *Gli3* (yellow; B, D). Antibodies directed against NP63 demarcate HBCs (blue; A-D).

Quantitation of *Gli2* (E) puncta and *Gli3* (F) puncta adjacent to NP63+ nuclei. Scale bar (A), 25μm. Data are

mean ± standard deviation. P-values were determined by a two-tailed unpaired t-test; n.s.= not significant

(p>0.05). (G-P) Coronal sections from adult *Krt5-Cre^{ER}*; *tdTomato/+*; *Gli2$^{fl/fl}$*; *Gli3$^{fl/fl}$* mice collected 8 weeks

following injury were analyzed with immunostaining for various OE markers. Mice lacking both *Gli2* and *Gli3*

display a range of morphological phenotypes (G-J). Antibodies directed against SOX2 (magenta; M, P, inset in

N) denote Sus cells apically, whereas antibodies directed against CD54 (cyan, M, P, inset in N) demarcate

HBCs. DAPI denotes nuclei (gray; G-J, K, N), tdTomato marks HBCs and their progeny following injury (red;

G-J, L, O), scale bar (G, K), 50μm. Quantitation of Sus cells (Q) and GBCs (R) per mm of OE. n=4 control and

n=5 *Krt5-Cre^{ER}*; *tdTomato/+*; *Gli2$^{fl/fl}$*; *Gli3$^{fl/fl}$* animals analyzed. Control refers to *Krt5-Cre^{ER}*; *tdTomato/+*;

Gli2$^{fl/+}$; *Gli3$^{fl/+}$* mice or mice lacking the *Krt5-Cre^{ER}* allele. Data are mean ± standard deviation. P-values were

determined by a two-tailed unpaired t-test; n.s.= not significant (p>0.05). (S) Table summarizing OE cell

composition in *Gli2^{CKO}*, *Gli3^{CKO}*, and *Gli2/3^{CKO}* mice compared to control mice. Green indicates increase in cell

number, red indicates decrease, light gray indicates no change.

References

Aziz, M., Goyal, H., Haghbin, H., Lee-Smith, W.M., Gajendran, M., and Perisetti, A. (2021). The Association
of "Loss of Smell" to COVID-19: A Systematic Review and Meta-Analysis. Am J Med Sci *361*, 216-225.
10.1016/j.amjms.2020.09.017.

Bai, C.B., Auerbach, W., Lee, J.S., Stephen, D., and Joyner, A.L. (2002). Gli2, but not Gli1, is required for
initial Shh signaling and ectopic activation of the Shh pathway. Development *129*, 4753-4761.

Bai, C.B., and Joyner, A.L. (2001). Gli1 can rescue the in vivo function of Gli2. Development *128*, 5161-5172.

Blaess, S., Stephen, D., and Joyner, A.L. (2008). Gli3 coordinates three-dimensional patterning and growth of
the tectum and cerebellum by integrating Shh and Fgf8 signaling. Development *135*, 2093-2103.
10.1242/dev.015990.

Briscoe, J., and Therond, P.P. (2013). The mechanisms of Hedgehog signalling and its roles in development and
disease. Nat Rev Mol Cell Biol *14*, 416-429. 10.1038/nrm3598.

Chang, C.F., Chang, Y.T., Millington, G., and Brugmann, S.A. (2016). Craniofacial Ciliopathies Reveal
Specific Requirements for GLI Proteins during Development of the Facial Midline. PLoS Genet *12*, e1006351.
10.1371/journal.pgen.1006351.

Chen, M., Reed, R.R., and Lane, A.P. (2019). Chronic Inflammation Directs an Olfactory Stem Cell Functional
Switch from Neuroregeneration to Immune Defense. Cell Stem Cell *25*, 501-513 e505.
10.1016/j.stem.2019.08.011.

Corrales, J.D., Blaess, S., Mahoney, E.M., and Joyner, A.L. (2006). The level of sonic hedgehog signaling
regulates the complexity of cerebellar foliation. Development *133*, 1811-1821. 10.1242/dev.02351.

Diamond, I., Owolabi, T., Marco, M., Lam, C., and Glick, A. (2000). Conditional gene expression in the
epidermis of transgenic mice using the tetracycline-regulated transactivators tTA and rTA linked to the keratin
5 promoter. J Invest Dermatol *115*, 788-794. 10.1046/j.1523-1747.2000.00144.x.

Durante, M.A., Kurtenbach, S., Sargi, Z.B., Harbour, J.W., Choi, R., Kurtenbach, S., Goss, G.M., Matsunami,
H., and Goldstein, B.J. (2020). Single-cell analysis of olfactory neurogenesis and differentiation in adult
humans. Nat Neurosci *23*, 323-326. 10.1038/s41593-020-0587-9.

Ermilov, A.N., Kumari, A., Li, L., Joiner, A.M., Grachtchouk, M.A., Allen, B.L., Dlugosz, A.A., and Mistretta,
C.M. (2016). Maintenance of Taste Organs Is Strictly Dependent on Epithelial Hedgehog/GLI Signaling. PLoS
Genet *12*, e1006442. 10.1371/journal.pgen.1006442.

Fletcher, R.B., Prasol, M.S., Estrada, J., Baudhuin, A., Vranizan, K., Choi, Y.G., and Ngai, J. (2011). p63
regulates olfactory stem cell self-renewal and differentiation. Neuron *72*, 748-759.
10.1016/j.neuron.2011.09.009.

Garcia, A.D., Petrova, R., Eng, L., and Joyner, A.L. (2010). Sonic hedgehog regulates discrete populations of
astrocytes in the adult mouse forebrain. J Neurosci *30*, 13597-13608. 10.1523/JNEUROSCI.0830-10.2010.

Genter, M.B., Deamer, N.J., Blake, B.L., Wesley, D.S., and Levi, P.E. (1995). Olfactory toxicity of
methimazole: dose-response and structure-activity studies and characterization of flavin-containing
monooxygenase activity in the Long-Evans rat olfactory mucosa. Toxicol Pathol *23*, 477-486.
10.1177/019262339502300404.

Gong, Q., Chen, H., and Farbman, A.I. (2009). Olfactory sensory axon growth and branching is influenced by sonic hedgehog. Dev Dyn *238*, 1768-1776. 10.1002/dvdy.22005.

Grachtchouk, M., Mo, R., Yu, S., Zhang, X., Sasaki, H., Hui, C.C., and Dlugosz, A.A. (2000). Basal cell carcinomas in mice overexpressing Gli2 in skin. Nat Genet *24*, 216-217. 10.1038/73417.

Grachtchouk, M., Pero, J., Yang, S.H., Ermilov, A.N., Michael, L.E., Wang, A., Wilbert, D., Patel, R.M., Ferris, J., Diener, J., et al. (2011). Basal cell carcinomas in mice arise from hair follicle stem cells and multiple epithelial progenitor populations. J Clin Invest *121*, 1768-1781. 10.1172/JCI46307.

Graziadei, P.P., and Graziadei, G.A. (1979). Neurogenesis and neuron regeneration in the olfactory system of mammals. I. Morphological aspects of differentiation and structural organization of the olfactory sensory neurons. J Neurocytol *8*, 1-18. 10.1007/BF01206454.

Guo, Z., Packard, A., Krolewski, R.C., Harris, M.T., Manglapus, G.L., and Schwob, J.E. (2010). Expression of pax6 and sox2 in adult olfactory epithelium. J Comp Neurol *518*, 4395-4418. 10.1002/cne.22463.

Herrick, D.B., Guo, Z., Jang, W., Schnittke, N., and Schwob, J.E. (2018). Canonical Notch Signaling Directs the Fate of Differentiating Neurocompetent Progenitors in the Mammalian Olfactory Epithelium. J Neurosci *38*, 5022-5037. 10.1523/JNEUROSCI.0484-17.2018.

Herrick, D.B., Lin, B., Peterson, J., Schnittke, N., and Schwob, J.E. (2017). Notch1 maintains dormancy of olfactory horizontal basal cells, a reserve neural stem cell. Proc Natl Acad Sci U S A *114*, E5589-E5598. 10.1073/pnas.1701333114.

Hoffman, H.J., Rawal, S., Li, C.M., and Duffy, V.B. (2016). New chemosensory component in the U.S. National Health and Nutrition Examination Survey (NHANES): first-year results for measured olfactory dysfunction. Rev Endocr Metab Disord *17*, 221-240. 10.1007/s11154-016-9364-1.

Huangfu, D., Liu, A., Rakeman, A.S., Murcia, N.S., Niswander, L., and Anderson, K.V. (2003). Hedgehog signalling in the mouse requires intraflagellar transport proteins. Nature *426*, 83-87. 10.1038/nature02061.

Imamura, F., and Hasegawa-Ishii, S. (2016). Environmental Toxicants-Induced Immune Responses in the Olfactory Mucosa. Front Immunol *7*, 475. 10.3389/fimmu.2016.00475.

Indra, A.K., Warot, X., Brocard, J., Bornert, J.M., Xiao, J.H., Chambon, P., and Metzger, D. (1999). Temporally-controlled site-specific mutagenesis in the basal layer of the epidermis: comparison of the recombinase activity of the tamoxifen-inducible Cre-ER(T) and Cre-ER(T2) recombinases. Nucleic Acids Res *27*, 4324-4327. 10.1093/nar/27.22.4324.

Joiner, A.M., Green, W.W., McIntyre, J.C., Allen, B.L., Schwob, J.E., and Martens, J.R. (2015). Primary Cilia on Horizontal Basal Cells Regulate Regeneration of the Olfactory Epithelium. J Neurosci *35*, 13761-13772. 10.1523/JNEUROSCI.1708-15.2015.

Kong, J.H., Yang, L., Dessaud, E., Chuang, K., Moore, D.M., Rohatgi, R., Briscoe, J., and Novitch, B.G. (2015). Notch activity modulates the responsiveness of neural progenitors to sonic hedgehog signaling. Dev Cell *33*, 373-387. 10.1016/j.devcel.2015.03.005.

Kumari, A., Ermilov, A.N., Allen, B.L., Bradley, R.M., Dlugosz, A.A., and Mistretta, C.M. (2015). Hedgehog pathway blockade with the cancer drug LDE225 disrupts taste organs and taste sensation. J Neurophysiol *113*, 1034-1040. 10.1152/jn.00822.2014.

Kumari, A., Ermilov, A.N., Grachtchouk, M., Dlugosz, A.A., Allen, B.L., Bradley, R.M., and Mistretta, C.M. (2017). Recovery of taste organs and sensory function after severe loss from Hedgehog/Smoothened inhibition with cancer drug sonidegib. Proc Natl Acad Sci U S A *114*, E10369-E10378. 10.1073/pnas.1712881114.

Leung, C.T., Coulombe, P.A., and Reed, R.R. (2007). Contribution of olfactory neural stem cells to tissue maintenance and regeneration. Nat Neurosci *10*, 720-726. 10.1038/nn1882.

Liu, H.X., Ermilov, A., Grachtchouk, M., Li, L., Gumucio, D.L., Dlugosz, A.A., and Mistretta, C.M. (2013). Multiple Shh signaling centers participate in fungiform papilla and taste bud formation and maintenance. Dev Biol *382*, 82-97. 10.1016/j.ydbio.2013.07.022.

Long, F., Zhang, X.M., Karp, S., Yang, Y., and McMahon, A.P. (2001). Genetic manipulation of hedgehog signaling in the endochondral skeleton reveals a direct role in the regulation of chondrocyte proliferation. Development *128*, 5099-5108.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat Neurosci *13*, 133-140. 10.1038/nn.2467.

Marczenke, M., Sunaga-Franze, D.Y., Popp, O., Althaus, I.W., Sauer, S., Mertins, P., Christ, A., Allen, B.L., and Willnow, T.E. (2021). GAS1 is required for NOTCH-dependent facilitation of SHH signaling in the ventral forebrain neuroepithelium. Development *148*. 10.1242/dev.200080.

McDermott, A., Gustafsson, M., Elsam, T., Hui, C.C., Emerson, C.P., Jr., and Borycki, A.G. (2005). Gli2 and Gli3 have redundant and context-dependent function in skeletal muscle formation. Development *132*, 345-357. 10.1242/dev.01537.

Memmi, E.M., Sanarico, A.G., Giacobbe, A., Peschiaroli, A., Frezza, V., Cicalese, A., Pisati, F., Tosoni, D., Zhou, H., Tonon, G., et al. (2015). p63 Sustains self-renewal of mammary cancer stem cells through regulation of Sonic Hedgehog signaling. Proc Natl Acad Sci U S A *112*, 3499-3504. 10.1073/pnas.1500762112.

Mistretta, C.M., and Kumari, A. (2019). Hedgehog Signaling Regulates Taste Organs and Oral Sensation: Distinctive Roles in the Epithelium, Stroma, and Innervation. Int J Mol Sci *20*. 10.3390/ijms20061341.

Morrison, E.E., and Costanzo, R.M. (1989). Scanning electron microscopic study of degeneration and regeneration in the olfactory epithelium after axotomy. J Neurocytol *18*, 393-405. 10.1007/BF01190842.

Packard, A., Schnittke, N., Romano, R.A., Sinha, S., and Schwob, J.E. (2011). DeltaNp63 regulates stem cell dynamics in the mammalian olfactory epithelium. J Neurosci *31*, 8748-8759. 10.1523/JNEUROSCI.0681-11.2011.

Peng, T., Frank, D.B., Kadzik, R.S., Morley, M.P., Rathi, K.S., Wang, T., Zhou, S., Cheng, L., Lu, M.M., and Morrisey, E.E. (2015). Hedgehog actively maintains adult lung quiescence and regulates repair and regeneration. Nature *526*, 578-582. 10.1038/nature14984.

Petrova, R., Garcia, A.D., and Joyner, A.L. (2013). Titration of GLI3 repressor activity by sonic hedgehog signaling is critical for maintaining multiple adult neural stem cell and astrocyte functions. J Neurosci *33*, 17490-17505. 10.1523/JNEUROSCI.2042-13.2013.

Petrova, R., and Joyner, A.L. (2014). Roles for Hedgehog signaling in adult organ homeostasis and repair. Development *141*, 3445-3457. 10.1242/dev.083691.

Rodriguez, S., Sickles, H.M., Deleonardis, C., Alcaraz, A., Gridley, T., and Lin, D.M. (2008). Notch2 is
required for maintaining sustentacular cell function in the adult mouse main olfactory epithelium. Dev Biol *314*,
40-58. 10.1016/j.ydbio.2007.10.056.

Sasaki, H., Hui, C., Nakafuku, M., and Kondoh, H. (1997). A binding site for Gli proteins is essential for HNF-
3beta floor plate enhancer activity in transgenics and can respond to Shh in vitro. Development *124*, 1313-1322.
10.1242/dev.124.7.1313.

Sasaki, H., Nishizaki, Y., Hui, C., Nakafuku, M., and Kondoh, H. (1999). Regulation of Gli2 and Gli3 activities
by an amino-terminal repression domain: implication of Gli2 and Gli3 as primary mediators of Shh signaling.
Development *126*, 3915-3924. 10.1242/dev.126.17.3915.

Scales, M.K., Velez-Delgado, A., Steele, N.G., Schrader, H.E., Stabnick, A.M., Yan, W., Mercado Soto, N.M.,
Nwosu, Z.C., Johnson, C., Zhang, Y., et al. (2022). Combinatorial Gli activity directs immune infiltration and
tumor growth in pancreatic cancer. PLoS Genet *18*, e1010315. 10.1371/journal.pgen.1010315.

Schwob, J.E., Jang, W., Holbrook, E.H., Lin, B., Herrick, D.B., Peterson, J.N., and Hewitt Coleman, J. (2017).
Stem and progenitor cells of the mammalian olfactory epithelium: Taking poietic license. J Comp Neurol *525*,
1034-1054. 10.1002/cne.24105.

Vortkamp, A., Lee, K., Lanske, B., Segre, G.V., Kronenberg, H.M., and Tabin, C.J. (1996). Regulation of rate
of cartilage differentiation by Indian hedgehog and PTH-related protein. Science *273*, 613-622.
10.1126/science.273.5275.613.

Wang, B., Fallon, J.F., and Beachy, P.A. (2000). Hedgehog-regulated processing of Gli3 produces an
anterior/posterior repressor gradient in the developing vertebrate limb. Cell *100*, 423-434. 10.1016/s0092-
8674(00)80678-9.

Wang, C., Cassandras, M., and Peng, T. (2019). The Role of Hedgehog Signaling in Adult Lung Regeneration
and Maintenance. J Dev Biol *7*. 10.3390/jdb7030014.

Wang, C., de Mochel, N.S.R., Christenson, S.A., Cassandras, M., Moon, R., Brumwell, A.N., Byrnes, L.E., Li,
A., Yokosaki, Y., Shan, P., et al. (2018). Expansion of hedgehog disrupts mesenchymal identity and induces
emphysema phenotype. J Clin Invest *128*, 4343-4358. 10.1172/JCI99435.
796

Figure 1. Shirazyan et al., 2022

Figure 2. Shirazyan et. al., 2022

Figure 3. Shirazyan et. al., 2022

A HBCs reconsititute OE in response to injury → Analyze

Gli2-cre/ERT2 · tdTomato

8 Weeks Post Methimazole Injury (Full Recovery)

| DAPI | NP63 | tdTomato | SEC8 | NP63/tdTomato/SEC8 |

Gli2Cre$^{ER/+}$; tdTomato/+

Vehicle (B–B'''') · Tamoxifen (C–C'''')

8 Weeks Post Methimazole Injury (Full Recovery)

| DAPI | SOX2 | tdTomato | CD54 | SOX2/tdTomato/CD54 |

Gli2Cre$^{ER/+}$; tdTomato/+

Vehicle (D–D'''') · Tamoxifen (E–E'''')

8 Weeks Post Methimazole Injury (Full Recovery)

| DAPI | CBX8 | tdTomato | CD54 | CBX8/tdTomato/CD54 |

Gli2Cre$^{ER/+}$; tdTomato/+

Vehicle (F–F'''') · Tamoxifen (G–G'''')

Figure 4. Shirazyan et. al., 2022

Figure 5. Shirazyan et. al., 2022

Figure 6. Shirazyan et. al., 2022

Figure 7. Shirazyan et. al., 2022

Figure S1. Shirazyan et. al., 2022

Figure S2. Shirazyan et. al., 2022

Figure S4. Shirazyan et. al., 2022

Figure S7. Shirazyan et. al., 2022

5 Tamoxifen I.P. Injections

NP63/*mGli2* — A, C (*Krt5-Cre^{ER}; tdTomato/+*, *Krt5-Cre^{ER}; Gli2^{fl/fl}; Gli3^{fl/fl}; tdTomato/+*)

NP63/*mGli3* — B, D

E *Gli2* mRNA in HBCs — p=0.0084, *mGli2*+ puncta (per mm OE)

F *Gli3* mRNA in HBCs — p=0.0109, *mGli3*+ puncta (per mm OE)

■ *Krt5-Cre^{ER}; tdTomato/+*
□ *Krt5-Cre^{ER}; Gli2^{fl/fl}; Gli3^{fl/fl}; tdTomato/+*

8 Weeks Post Methimazole Injury (Full Recovery) — DAPI/tdTomato (G, H, I, J)

8 Weeks Post Methimazole Injury (Full Recovery) — DAPI (K, N); tdTomato (L, O); SOX2/CD54 (M, P)

Q Sus Cells — n.s.(p=0.6608), Apical SOX2+ (per mm OE), Full Recovery

R GBCs — n.s.(p=0.1492), SEC8+ (per mm OE), Full Recovery

■ Control
□ *Krt5-Cre^{ER}; Gli2^{fl/fl}; Gli3^{fl/fl}; tdTomato/+*

S	OE Cell Composition (Compared to Control)			
Genotype	HBCs	GBCs	OSNs	Sus
Gli2^{CKO}	No Change	No Change	No Change	No Change
Gli3^{CKO}	↑	↑	No Change	↑
Gli2/3^{CKO}	↓	↓	↓	No Change