Revisiting the impact of synthetic ORF sequences on engineered LINE-1 retrotransposition

Dorothy Chan[1,5], Stephanie Workman[1,5], Patricia Gerdes[1], Jeffrey S Han[2], Jef D Boeke[3], Geoffrey J

Faulkner[1,4,*], Sandra R Richardson[1,*]

1. Mater Research Institute - University of Queensland, TRI Building, Woolloongabba QLD 4102,
Australia.

2. Department of Biochemistry and Molecular Biology, Tulane University School of Medicine, New
Orleans, LA, 70112, USA.

3. Institute for Systems Genetics and Department of Biochemistry and Molecular Pharmacology, NYU
Langone Health, New York, NY, 10016, USA; Department of Biomedical Engineering, NYU Tandon
School of Engineering, Brooklyn NY 11201

4. Queensland Brain Institute, University of Queensland, Brisbane QLD 4072, Australia.
12
5. These authors contributed equally.

*Correspondence: S.R.R. (sandra.richardson@mater.uq.edu.au), G.J.F. (faulknergj@gmail.com)

Abstract: The retrotransposon Long Interspersed Element 1 (L1) contains adenosine rich ORFs, a

characteristic that limits its expression in mammalian cells. A synthetic mouse L1 (smL1) with ORF

adenosine content decreased from 40% to 26% showed increased mRNA expression and

retrotransposed far more efficiently than the native parental element, L1spa (1). Here, we observe

two nonsynonymous substitutions between the L1spa and smL1 ORF1 sequences, and note that the

smL1 3'UTR lacks a conserved guanosine-rich region (GRR) which could potentially take on a G-

quadruplex secondary structure. We find that the combined effect of a single amino acid change and

the GRR 3'UTR deletion, rather than synthetic ORF sequences, accounts for the increase in smL1

retrotransposition efficiency over L1spa. Furthermore, we demonstrate that the position of the GRR

within the L1 reporter construct impacts retrotransposition efficiency. Our results prompt a

reevaluation of synthetic L1 activity and suggest native mouse L1 mobility has in some cases been

underestimated in engineered retrotransposition assays.

Author Summary: L1 retrotransposons are mobile DNA elements or "jumping genes" that can copy-

and-paste their sequences to new locations in the host genome. The jumping ability, or

retrotransposition efficiency, of individual L1 elements can be evaluated using a cultured cell assay in

which the L1 is tagged in its 3' untranslated region (3'UTR) with a reporter gene that becomes

expressed upon successful retrotransposition. In a previous study, authors Han and Boeke reported

that the retrotransposition efficiency of a mouse L1 element could be enhanced dramatically by

synthetically increasing the GC content of the L1 ORFs without changing their amino acid sequence.

Curiously, a similarly constructed synthetic human L1 achieved only a modest increase in

retrotransposition efficiency over the native element. Here, we find that two coding changes and

partial deletion of the mouse L1 3'UTR sequence which occurred during construction of the synthetic

mouse L1 reporter actually are responsible for the increased jumping of this construct. We also find

that changing the placement as well as the presence of this deleted 3'UTR region within the reporter

construct determines its impact on engineered retrotransposition efficiency. Together, our study

reconciles the disparate impacts of synthetic sequences upon human and mouse L1 retrotransposition

efficiency, prompts a reconsideration of numerous studies using synthetic L1 constructs, and will

inform the ongoing use of synthetic and natural mouse L1 reporter constructs in vivo and in vitro.

Introduction: L1 retrotransposons are an ongoing source of mutagenesis in mammalian genomes (2,

3). Mice contain ~3000 retrotransposition-competent L1s (RC-L1s) representing three subfamilies

(L1_T_F, L1_G_F, L1_A), distinguished by their monomeric promoter-harboring 5'UTRs (4-9). RC-L1s

encode two proteins required for their mobility in cis: ORF1p, a nucleic acid binding and chaperone

protein, and ORF2p, which has endonuclease and reverse transcriptase activities critical for the

generation of new L1 insertions by target-primed reverse transcription (TPRT) (10-18). The L1 3'UTR

contains a weak polyadenylation (poly-A) signal and a poly-purine tract which is found in L1s across

species but which varies in length and nucleotide sequence (19-23). Hallmarks of TPRT-mediated L1

integrants include insertion at the degenerate consensus sequence 5'-TTTT/AA-3', terminal poly-A

tracts, and flanking target site duplications (TSDs) (24-28). Approximately 1 in 8 mice and 1 in 62.5

humans harbor a de novo L1 insertion (29, 30). Moreover, >100 cases of human genetic disease and

numerous spontaneous mouse mutants have been linked to L1-mediated retrotransposition events,

highlighting the impact of L1 mutagenesis on mammalian genomes (31, 32).

The retrotransposition efficiency of individual L1 copies can be evaluated in cultured cells and

in vivo (33-35). In these assays, a retrotransposition indicator cassette (36), consisting of a reporter

gene in the opposite transcriptional orientation to the L1 and equipped with its own promoter and

polyadenylation (poly-A) signal, is placed into the L1 3'UTR (Figure 1A). The reporter gene is

interrupted by a backwards intron, with expression achieved only upon splicing and reverse

transcription of the L1 mRNA to deliver an intact copy of the reporter cassette into genomic DNA.

Expression and polyadenylation of the full-length L1 mRNA can be augmented by a strong

heterologous promoter and poly-A signal flanking the L1 element and reporter cassette (33, 37). In

selection-based cultured cell assays, quantification of antibiotic-resistant foci provides a readout of

retrotransposition efficiency (33).

The adenosine richness of L1 ORFs limits L1 expression in mammalian cells and influences

expression of genes containing intronic sense oriented L1 insertions (38-40). A synthetic mouse L1

sequence (smL1, or ORFeus_Mm) was previously developed by Boeke and Han to increase the

retrotransposition efficiency of engineered L1 reporter constructs, improving their utility for random

mutagenesis screens of mammalian genomes (1). smL1 was derived from the $L1\_T_F$ subfamily element

L1spa (5, 41) whereby the adenosine content of the L1spa ORFs was reduced from 40% to 26% via

synonymous substitutions. The synthetic ORF sequences of smL1 yielded markedly increased mRNA

expression levels, and smL1 retrotransposed ~200-fold more efficiently in HeLa cells than the

retrotransposition indicator construct pTN201/L1spa (1, 5). Notably, similar constructs containing a

synthetic human L1 showed at most a 3-fold increase in retrotransposition efficiency over the native

element (42). Here, we note both coding and non-coding differences between smL1 and

pTN201/L1spa, and we systematically evaluate the impact of these differences, and of synthetic ORF

Figure 1

Figure 1. L1spa and smL1 differ at two ORF1p amino acid positions.

A. The cultured cell retrotransposition assay. A full-length L1 containing intact open reading frames (ORF1 and ORF2) is driven by its native 5'UTR promoter or a strong heterologous promoter (grey), and equipped with a polyadenylation signal (native and/ or engineered, open lollipop). The L1 is tagged with a retrotransposition indicator cassette (green) inserted in its 3'UTR. The reporter consists of a backwards intron-containing neomycin phosphotransferase (NEO) reporter gene in opposite transcriptional orientation to the L1 and equipped with a separate promoter and polyadenylation signal (upside-down black arrow and black lollipop, respectively). Expression of the reporter gene from genomic DNA is achieved upon retrotransposition.

B. L1spa/pTN201 and smL1 plasmids. Features are depicted as follows: CMV promoter (grey), L1spa 5'UTR (white triangles, representing monomers). NEO reporter cassette (green) with SV40 early promoter (black arrow) and HSV-Tk polyadenylation signal (filled black lollipop). Native L1spa nucleotide sequence (white); synthetic sequence (pink). The amino acids at positions 53 and 159 in ORF1p are indicated. Engineered Kozak consensus sequence (star). L1spa 3'UTR (dark blue); G-rich region (red). Mutated smL1 3'UTR (light blue), deleted GRR (Δ symbol). Human L1 polyadenylation signal (open grey lollipop) and the SV40 polyadenylation signal (open black lollipop).

C. Effects of ORF1p amino acid changes on retrotransposition. From left to right: a schematic of each construct (as described in B), a representative well from the retrotransposition assay, and its retrotransposition efficiency relative to smL1. The histogram displays the mean of three biological replicate assays, each comprising three technical replicates per construct. Triangle, square, and diamond shapes indicate the individual values for each biological replicate, and the error bars represent standard deviation among biological replicates.

sequences, on mouse L1 retrotransposition efficiency. We determine that the presence and position

of a conserved poly purine tract in the mouse L1 3'UTR profoundly impacts the retrotransposition

efficiency of engineered mouse L1 reporter constructs.

Results: We capillary sequenced the pCEP4-based smL1 and pTN201/L1spa plasmids (Supplemental

Table 1), and observed that the synthetic sequences of the smL1 ORFs largely comprised synonymous

nucleotide substitutions relative to L1spa. However, smL1 ORF1 contained two nonsynonymous

substitutions (G53D and H159D) and bore an engineered Kozak consensus at its initiation codon that

was absent from pTN201/L1spa. Furthermore, the terminal 159 bp of 3'UTR sequence present in

pTN201/L1spa construct was absent from smL1 (Figure 1B; Supplemental Figure 1). We refer to this

159 bp sequence as the guanine-rich region (GRR), as it encompasses the conserved poly-purine tract.

We traced this deletion to the inadvertent loss of a NdeI restriction fragment during subcloning of the

smL1 construct (Supplemental Figure 2). The remaining 3'UTR sequence of smL1 contained 19

nucleotide substitutions and two single nucleotide deletions relative to the L1spa 3'UTR, which arose

during synthesis of the smL1 construct (Supplemental Figure 1).

The $L1\_T_F$ subfamily consensus, and previously described active mouse $L1\_T_F$ elements,

contain aspartic acid at ORF1p positions 53 and 159 (5, 6, 29, 43). Indeed, the substitution H159D

improves L1spa ORF1p nucleic acid chaperone activity and retrotransposition efficiency (44). To

quantify the contribution of G53D and H159D to the increased retrotransposition efficiency of smL1

over L1spa, we generated L1spa_G53D, L1spa_H159D, and a construct with both changes, L1spa_Dbl

(Figure 1C). In a HeLa cell transient transfection-based retrotransposition assay (34), L1spa mobilized

at ~6% the efficiency of smL1, and, consistent with previous results(44), the substitution G53D had

little effect on retrotransposition. However, L1spa_H159D and L1spa_Dbl mobilized at ~19% the

efficiency of smL1. The reciprocal changes (D53G and D159H) reduced smL1 retrotransposition

efficiency by ~40% (smL1_Dbl; Figure 1C). Thus, the H159D amino acid substitution partially explains

the increased retrotransposition efficiency of smL1 over L1spa.

To determine the contribution of non-coding changes between smL1 and L1spa_Dbl to their

differing retrotransposition efficiencies, we equipped L1spa_Dbl ORF1 with a Kozak consensus and

replaced the L1spa 3'UTR with the deletion-containing smL1 3'UTR to generate L1spa_Dbl_smBB.

Despite entirely lacking synthetic ORF sequences, L1spa_Dbl_smBB retrotransposed with equal

efficiency to smL1 (Figure 2). When tested independently, removing the Kozak consensus from this

construct (L1spa_Dbl_smBB_delete_Kozak) had no impact on retrotransposition efficiency. However,

restoring the GRR-containing L1spa 3'UTR (L1spa_Dbl_smBB_restore_3'UTR) reduced

retrotransposition to similar levels as L1spa_Dbl (Figure 2). We therefore concluded that the deletion-

containing smL1 3'UTR largely accounts for the increased retrotransposition efficiency of smL1 over

L1spa.

We speculated that the GRR might diminish observed retrotransposition by preventing

transcription through the NEO cassette. Indeed, G-quadruplex forming sequences have been

demonstrated to impede RNA polymerase procession (45-48) and implicated in the use of alternative

polyadenylation sites (49). To approach these possibilities, we first asked whether placement of the

GRR relative to the NEO cassette impacts retrotransposition efficiency. We replaced the L1spa GRR

into L1spa_Dbl_SMBB, inserting it downstream of NEO but upstream of the pCEP4 vector SV40

polyadenylation signal, to generate L1spa_Dbl_smBB_GRRAN. Unexpectedly, this construct

retrotransposed with equal efficiency to smL1 and L1spa_Dbl_smBB (Figure 2). We therefore

investigated whether formation of a G-quadruplex secondary structure within the GRR diminished

retrotransposition via a transcriptional block when placed upstream of the NEO cassette. To infer the

structure of L1spa_Dbl_smBB_GRRAN mRNA molecules from which successful retrotransposition

events arose, we characterized 10 L1spa_Dbl_smBB_GRRAN integrants by inverse PCR. Like smL1

insertions previously characterized in HeLa cells (1), all 10 L1spa_Dbl_smBB_GRRAN insertions bore

the hallmarks of bona fide retrotransposition events and incorporated L1 sequence, the spliced NEO

cassette, and the GRR. Importantly, all insertions terminated in poly-A tracts directed by the SV40

poly-A signal (Figure 3A, Supplemental Figure 3, Supplemental Table 1). We concluded that these

Figure 2

A.

Figure 2. The presence and position of the L1spa 3'UTR G-rich region impact retrotransposition.

A. From left to right: a schematic of each construct, a representative well from the retrotransposition assay, and its retrotransposition efficiency relative to smL1. L1spa_Dbl_smBB contains the ORF1 and ORF2 sequences of L1spa_Double with a Kozak consensus at the ORF1p initiation codon (as in smL1), and the deletion 3'UTR of the smL1 construct. Restore_3'UTR is equivalent to L1spa_Dbl_smBB but with the full L1spa 3'UTR restored upstream of the NEO cassette. GRR_after_NEO is equivalent to L1spa_Dbl_smBB but with the G-rich region of the L1spa 3'UTR placed downstream of the NEO cassette, and the human L1 polyadenylation signal removed. The histogram displays the mean of three biological replicate assays, each comprising three technical replicates per construct. Triangle, square, and diamond shapes indicate the individual values for each biological replicate, and the error bars represent standard deviation among biological replicates.

insertions occurred following complete transcription of the L1spa_Dbl_smBB_GRRAN construct,

including RNA polymerase procession through the GRR and utilization of the SV40 poly-A signal,

arguing against hinderance of RNA polymerase by the GRR.

We next asked whether the GRR can promote premature polyadenylation. In the L1spa_Dbl

construct, where the GRR is upstream of the NEO cassette, polyadenylation in the vicinity of the GRR

would result in an L1 mRNA molecule lacking the NEO cassette, the retrotransposition of which would

be undetectable in a selection-based assay. To determine whether positioning the GRR upstream of

the NEO cassette results in premature polyadenylation, we performed 3'RACE assays on RNA

extracted from cells transfected with L1spa_Dbl constructs with the GRR upstream of the NEO cassette

(L1spa_Dbl and Restore_3'UTR), downstream of the NEO cassette (L1spa_Dbl_smBB_GRRAN), or

deleted entirely (L1spa_Dbl_SMBB). We used L1-specific primers located in the SV40 promoter of the

antisense NEO cassette to detect mRNAs correctly polyadenylated at the SV40 polyadenylation signal,

and in the L1spa 3'UTR to detect mRNAs potentially prematurely polyadenylated proximal to the GRR.

We also used a primer annealing to the 3' end of the hygromycin resistance gene in the pCEP4

backbone as an internal control (Figure 3B). For all constructs, we detected the expected ~520bp band

corresponding to polyadenylated transcripts from the hygromycin resistance gene. Using a primer to

the antisense SV40 promoter responsible for NEO expression, we detected the expected ~350 bp band

corresponding to L1 reporter transcripts polyadenylating at the SV40 polyadenylation signal for

L1spa_Dbl, L1spa_Dbl_SMBB, and Restore_3'UTR, and a ~520 bp band from

L1spa_Dbl_SMBB_GRR_after_NEO, the increased size of which is accounted for by the inclusion of the

GRR sequence between the antisense SV40 promoter and the SV40 polyadenylation signal. Using a

primer annealing to the L1spa 3'UTR, we did not detect bands in the range of 200-400 bp indicative of

polyadenylation within or proximal to the GRR in either the construct L1spa_Dbl or Restore_3'UTR.

Thus, we conclude that premature polyadenylation of sense engineered L1 transcripts is unlikely to be

responsible for the observed reduction in retrotransposition efficiency when the GRR is situated

upstream of the NEO cassette.

Figure 3

A.

L1spa_Dbl_smBB_GRRAN (spliced)

B.

Figure 3: The GRR does not promote premature polyadenylation

A. Structural hallmarks of ten insertions generated with the GRR_after_NEO construct are shown, including the 5' TSD, 5' truncation position relative to the L1spaDbl_smBB_GRRAN, polyadenylation position, poly-A tail length and 3'TSD. Below, the structure of GRR after NEO, with the spliced NEO cassette in green, the GRR in red, and the SV40 polyadenylation signal in black. Schematic is not to scale. Delta symbol indicates a target-site deletion. Insertion I contained a 5' inversion for which the break-points are indicated. Insertions I and J arose from a transcript initiating upstream of the L1 5'UTR, at the transcription start site of the CMV promoter within the pCEP4 vector(67) denoted as position -43. For two insertions bearing >100 bp TSDs, the TSD length, rather than TSD sequence, is shown.

B. 3'RACE assay to identify sites of polyadenylation. Above, schematics representing the constructs 1) L1spa_Dbl, 2) L1spa_Dbl_SMBB, 3) Restore_3'UTR and 4) GRR_after_NEO. The positions of gene-specific primers annealing to the L1spa 3'UTR (blue), the antisense SV40 promoter driving NEO expression (grey), and the hygromycin resistance gene in the pCEP4 vector backbone (orange) are denoted by arrows. Schematics are not drawn to scale. Below, agarose gel showing 3'RACE products detected using RNA isolated from cells transfected with the four above constructs, using each of the primers shown.

Finally, we investigated the sequence characteristics of the L1spa GRR responsible for

diminished engineered retrotransposition efficiency when placed upstream of the NEO cassette. An

L1spa_Dbl construct containing the L1spa GRR and lacking the remaining 3'UTR sequence

retrotransposed comparably to L1spa_Dbl, confirming that the L1spa GRR alone is sufficient to impede

engineered retrotransposition (L1spa_GRR, Figure 4A). To determine whether this effect is common

to the GRR sequences from ancient (V, F) and presently active (G_F, A) L1s in addition to the T_F subfamily

(represented by L1spa), we replaced the L1spa GRR with the consensus GRR sequence from each of

these subfamilies, which contain 9 (G_F, A) or 10 (V, F) runs of 3 or more guanine residues (6). In all

cases, retrotransposition was reduced to <20% of L1spa_Dbl_smBB (Figure 4A). To examine in greater

detail the role of the L1spa GRR sequence in retrotransposition inhibition, we generated a series of

deletion constructs. The L1spa GRR contains 9 runs of 3 or more guanine residues, separated by

variable stretches of other bases (Figure 4B). Relative to L1spa_Dbl_SMBB (no GRR sequence), the

sequential restoration of guanine runs in a 5'-to-3' direction resulted in a diminution of

retrotransposition efficiency, with the largest difference found between the constructs containing 5

and 6 guanine runs (Figure 4C). Thus, the position-dependent inhibitory effect of the GRR on

engineered L1 retrotransposition is shared by both ancient and presently active mouse L1s, and the

magnitude of inhibition is positively correlated with the length of the GRR.

Discussion: The highly active smL1 construct and its derivatives have facilitated numerous insights into

the mechanism, regulation, and consequences of L1 retrotransposition in vivo and in vitro (50-59).

Here we demonstrate that synthetic reduction in mouse L1 ORF adenosine content has little impact

on retrotransposition efficiency, at least in HeLa cells. We find instead that the elevated activity of

smL1 over pTN201/L1spa arises partly from the nonsynonymous ORF1 substitution H159D (44), and

to a larger extent from the deletion of the L1spa GRR from the smL1 3'UTR. Our results resolve a long-

standing incongruity regarding the impact of synthetic ORF sequences on L1 retrotransposition

efficiency, as the similarly re-coded synthetic human L1 ORFeus_Hs retrotransposes no more than 3-

fold more efficiently than the native parental element L1$_{RP}$ (42, 60). Thus, while both human and

A.

B. L1spa GRR nucleotide sequence

5'-TCAAAAGATGGCCTAGTCGGCCATCACTGGAAAGAGAGGCCCATTGGACACGCAGA

CTTTGTGTGCCCCGGTACA**GGGG**AACGCCA**GGG**CCAAA**GGGGGGGGG**AGT**GGGT**GGG
                        G1   G2   G3   G4   G5

TA**GGGG**AGT**GGGGG**TAGGT**GGG**TAA**GGGGG**ACTTTTGGTATAGCATT-3'
   G6   G7   G8   G9

C.

Percent Retrotransposition relative to smBB

Figure 4: Variations in G-rich region sequence content differentially impact retrotransposition.

A. From left to right: a schematic of each construct, a representative well from the retrotransposition assay, and its retrotransposition efficiency relative to smL1. L1spa_GRR contains the GRR portion only of the L1spa 3'UTR. V_GRR, F_GRR, GF_GRR and A_GRR contain the GRR portion of the 3'UTR from the L1Md_V$_I$, L1Md_F$_I$, L1Md_G$_{FI}$ and L1Md_A$_I$ consensus sequences. The histogram displays the mean of three biological replicate assays, each comprising three technical replicates per construct. Triangle, square, and diamond shapes indicate the individual values for each biological replicate, and the error bars represent standard deviation among biological replicates.

B. Nucleotide sequence of the 159bp region comprising the G-rich region (GRR) of the L1spa 3'UTR. Runs of 3 or more consecutive G nucleotides are denoted as G1 through G9 and highlighted in red.

C. From left to right: a schematic of each construct, a representative well from the retrotransposition assay, and its retrotransposition efficiency relative to L1spa_Dbl_SMBB. Runs of guanine residues have been progressively restored to the GRR of constructs G1 through G8, with the full GRR present in construct G9. The histogram displays the mean of three biological replicate assays, each comprising three technical replicates per construct. Triangle, square, and diamond shapes indicate the individual values for each biological replicate, and the error bars represent standard deviation among biological replicates.

mouse synthetic L1 ORFs yield massively increased mRNA levels relative to native ORF sequences, this

elevated expression does not effect a proportionate increase in retrotransposition (1, 42). This result

is consistent with the modest increase in retrotransposition achieved by augmenting L1 expression

with a strong heterologous promoter (33), and suggests that L1 mRNA levels are not the rate-limiting

determinant of retrotransposition efficiency.

The molecular mechanism by which deleting or repositioning the L1spa GRR impacts

retrotransposition efficiency remains unclear, and does not appear to be caused by direct impediment

of RNA polymerase processing or by premature polyadenylation. Importantly, the impaired

retrotransposition efficiency observed when the L1spa GRR is placed upstream of the NEO cassette is

detected using GRR sequences from active and ancient mouse L1 elements, but based on previous

reports does not extend to human L1 reporter constructs. In contrast, deletions encompassing the

human poly-purine tract from NEO- and EGFP-based L1 constructs, in which this motif is also

positioned upstream of the reporter cassette, reduced retrotransposition by ~10% and ~30%,

respectively (33, 61). Furthermore, experimental evidence suggests that the human poly-purine tract

can form a G-quadruplex (G4) secondary structure (19, 62) and that stabilization of this structure very

modestly stimulates engineered retrotransposition (61). The decreased retrotransposition efficiency

of constructs incorporating the L1spa GRR upstream of the NEO cassette may represent an artifact

specific to the L1spa GRR sequence composition and its ability to form secondary structures in the

context of the reporter construct (19), rather than evidence that this motif intrinsically inhibits

retrotransposition. Indeed, the possibility that disruption of the L1 3'UTR by an indicator cassette

could impact retrotransposition has long been acknowledged (63) and, to speculate, the conservation

of a self-attenuating motif is difficult to reconcile with the evolutionary imperative of L1 as a selfish

genetic element (19, 20).

We observed no difference in retrotransposition efficiency when the L1spa GRR was deleted

entirely, compared to when it is repositioned downstream of the NEO cassette. This result suggests

that a potential function for the GRR in endogenous mouse L1 retrotransposition is likely to be subtle,

perhaps influenced by the stabilization of RNA or DNA G4 secondary structures and dependent on the

cellular milieu. The extent to which the deletion of the GRR from the smL1 construct and its derivatives

impacts the conclusions drawn from previous in vitro and in vivo studies using these reporters remains

to be evaluated (50-59); at least some transgenic mouse models generated using smL1 based

constructs lack the GRR (50, 53, 57, 64). Engineered mouse L1 constructs reconfigured with the GRR

downstream of the reporter cassette represent useful tools for investigating the function of this

conserved motif, and for studies of mouse L1 biology in general, as high efficiency retrotransposition

and structurally normal L1 integrants can be achieved using a construct containing the entire native

mouse L1 sequence.

Materials and Methods:

Constructs used in this study:

pTN201/L1spa: Has been described previously (5). It consists of the pCEP4 backbone with the element

L1spa truncated at position 7418 in the 3'UTR (removing the native mouse L1 polyadenylation signal

but leaving the G-rich region (GRR) of the 3'UTR in place). The L1spa 3'UTR is followed by the mneoI

retrotransposition indicator cassette (33, 36). The mneoI cassette is followed by the human L1

polyadenylation signal which is directly upstream of the SV40 polyadenylation signal in the pCEP4

backbone.

L1spa_G53D: Identical to pTN201/L1spa but with glycine 53 of ORF1 replaced by aspartic acid.

L1spa_H159D: Identical to pTN201/L1spa but with histidine 159 of ORF1 replaced by aspartic acid.

L1spa_dbl: Identical to pTN201/L1spa but with glycine 53 of ORF1 replaced by aspartic acid and

histidine 159 of ORF1 replaced by aspartic acid.

smL1: Has been described previously (1). It consists of the pCEP4 backbone with a synthetic mouse L1

element based on L1spa. The initiation codon of ORF1 has been placed into an engineered Kozak

consensus. The amino acid sequence of smL1 is identical to that of L1spa except for two residues in

ORF1 (position 53 and position 159). The 3'UTR of smL1 contains several single nucleotide

substitutions relative to L1spa, as well as two single-nucleotide deletions, and is truncated at position

7259 of L1spa. This truncation deletes the GRR of the L1spa 3'UTR. The smL1 3'UTR is followed by the

mneoI retrotransposition indicator cassette. The mneoI cassette is followed by the human L1

polyadenylation signal which is directly upstream of the SV40 poly-A signal in the pCEP4 backbone.

smL1_mut2: Has been described previously (1). It is identical to smL1 but with inactivating mutations

in the endonuclease and reverse transcriptase active sites of ORF2.

smL1_Dbl: Identical to smL1 but with aspartic acid at position 53 of ORF1 replaced by glycine and

aspartic acid at position 159 of ORF1 replaced by histidine.

L1spa_dbl_smBB: "L1spa double; smL1 backbone". It consists of the L1spa_dbl sequence, with a Kozak

consensus at the initiation codon of ORF1 and the mutated 3'UTR of smL1.

L1spa_dbl_smBB _Del_Kozak: Identical to L1spa_dbl_smBB but with the Kozak consensus removed.

L1spa_dbl_smBB_Restore_3'UTR: Identical to L1spa_dbl_smBB but with the mutated 3'UTR of L1SM

replaced by the native L1spa 3'UTR as found in pTN201/L1spa.

L1spa_Dbl_smBB_GRRAN: "L1spa double; smL1 backbone; G-rich region after NEO". Identical to

L1spa_dbl_smBB but with the GRR of L1spa (nucleotides 7260-7416) and an engineered AflII

restriction site placed after the mneoI cassette and before the SV40 polyadenylation signal of the

pCEP4 backbone, and the human L1 polyadenylation signal deleted.

L1spa_Dbl_SMBB_L1spaGRR: Identical to L1spa_Dbl_SMBB, but the GRR sequence from the L1spa

3'UTR placed between the end of ORF2 and the NEO cassette, using an existing AgeI site and

introducing an engineered SbfI site.

L1spa_Dbl_SMBB_V_GRR,   L1spa_Dbl_SMBB_F_GRR,   L1spa_Dbl_SMBB_G_F_GRR,   and

L1spa_Dbl_SMBB_A_GRR: Identical to L1spa_Dbl_SMBB_L1spaGRR, but with the GRR sequence from

the L1Md_VI, L1Md_FI, L1Md_GFI and L1Md_AI consensus sequences (6) placed between the end of

ORF2 and the NEO cassette, using AgeI and SbfI sites described above.

L1spa_Dbl_SMBB_G1 through L1spa_Dbl_SMBB_G9: Identical to L1spa_Dbl_SMBB, but with

increasing portions of the GRR sequence from the L1spa 3'UTR, each restoring an additional run of 3

or more guanine residues, introduced between the end of ORF2 and the NEO cassette, using the AgeI

and SbfI sites as described for L1spa_Dbl_SMBB_L1spaGRR.

pCEP4_GFP: Consists of the pCEP4 backbone with a humanized renilla GFP reporter gene.

Construct generation: all cloning was performed using restriction enzymes from New England Biolabs

(NEB) according to manufacturer's instructions. PCR fragments were generated using oligonucleotides

ordered from Integrated DNA Technologies (IDT) and Q5 High Fidelity DNA Polymerase 2X master mix

(NEB), according to manufacturer's instructions. Constructs were thoroughly capillary sequenced to

confirm the absence of random PCR-induced mutations.

Cell culture: HeLa-JVM cells (33) were maintained in Dulbecco's Modified Eagle's Medium (DMEM)

high glucose supplemented with 1% L-glutamine, 1% pen-strep and 10% fetal bovine serum. Cells

were passaged every 3-5 days at 70-80% confluence using 0.25% Trypsin-EDTA (Life Technologies).

Cells were grown at 37°C in 5% CO_2.

Retrotransposition assays: For transient retrotransposition assays performed in 6-well plates as

previously described(34), $5x10^3$ HeLa-JVM cells were plated per well. 24 hours later, each well was

transfected with 1 μg of plasmid DNA using 4 ul FuGene and 96 uL opti-MEM. Each construct was

transfected in triplicate. The media was replaced 24 hours post-transfection. At 72 hours post-

transfection, selection was initiated with 400 μg/ml G418 (Geneticin, Life Technologies). G418 media

was replaced every other day. At 12 days post-transfection, cells were fixed with 2%

formaldehyde/0.4% glutaraldehyde solution and stained with 0.1% crystal violet solution.

Assays to measure transfection efficiency for each L1 reporter plasmid were carried out as follows:

4×10^4 HeLa cells were plated per well of a 6-well dish. 24 hours later, each well was transfected with

1 μg total plasmid DNA, consisting of 0.5 μg L1 reporter plasmid and 0.5 μg pCEP4-GFP, using 4 ul of

FuGene and 96 ul opti-MEM. Each L1 reporter+GFP plasmid was transfected in duplicate. The media

was replaced 24 hours post-transfection. At 72 hours post-transfection, cells were trypsinized,

resuspended in PBS (Life Technologies) and assessed for GFP expression by flow cytometry using a

Cytoflex flow cytometer (Translational Research Institute flow cytometry core). The average percent

GFP-positive cells was determined for each L1 reporter construct, and used to normalize the results

of the retrotransposition assay.

Clonal cell lines for inverse PCR characterization of L1spa_Dbl_smBB_GRRAN insertions were

generated by plating 5×10^3 HeLa cells per 10 cm dish, and 18 hours later transfecting with 1 ug plasmid

DNA using 4 ul FuGene and 96 ul opti-MEM. Media was replaced 24 hours after transfection. Selection

with 400 ug/ml Geneticin was initiated at 72 hours post-transfection and continued for 12 days to

generate isolated colonies harboring retrotransposition events. Individual colonies were manually

picked and expanded to generate clonal cell lines. Approximately 5×10^6 cells were harvested per clonal

cell line and subjected to phenol-chloroform DNA extraction. DNA concentration was measured via

NanoDrop. Inverse PCR was carried out as described previously (65, 66). Briefly, 4 ug from each cell

line was digested with 25 units EcoRI (New England Biolabs) in 100 ul total volume. Reactions were

heat inactivated and digested DNA was ligated under dilute conditions to promote intramolecular

ligation (1 ml total volume) using 3200 units T4 DNA ligase (NEB) for 24 hours. Ligations were purified

by chloroform extraction and DNA was resuspended in 40 ul water. For each ligation, 4 ul of DNA was

used as a template in the first-round inverse PCR reaction using the Roche Expand long-template PCR

system. Each reaction consisted of 10 pmol primer NEO210as (5' GACCGCTTCCTCGTGCTTTACG 3), 10

pmol primer NEO1720s (5' TGCGCTGACAGCCGGAACACG 3'), 20 nM each dNTP, and 2.5 units of

enzyme. Cycling conditions were 95°C for 2 minutes, followed by 30 cycles of 94°C for 15 seconds,

64°C for 30s, and 68°C for 15 min, followed by a 30 min extension at 68°C. For the second-round

inverse PCR reaction, 4 ul of the product from the first-round inverse PCR was used directly as

template, with 10 pmol primer NEO173as (5' CATCGCCTTCTATCGCCTTCTTG 3') 10 pmol primer

NEO1808s (5' GCGTGCAATCCATCTTGTTCAATG 3'), 20 nM each dNTP, and 2.5 units of enzyme. Cycling

conditions were identical to first-round inverse PCR. Products were run on a 1% agarose gel and bands

were excised and purified using the Qiagen Min-Elute gel extraction kit. Bands <2kb were cloned using

the pGEM-T Easy kit (Promega) and bands >2kb were cloned using the Topo-XL2 PCR kit (Life

Technologies) following manufacturer's instructions. Cloned PCR products were sequenced using M13

forward and reverse primers Australian Genome Research Facility, Brisbane). Validation primers

(Integrated DNA Technologies) were designed based on the putative genomic location of each

insertion detected by inverse PCR, and insertions were validated as empty/filled and/or by 5' and 3'

junction PCRs using MyTaq HS DNA polymerase (Bioline). Validation products were run on 1% agarose

gels, purified using the Qiagen Min-Elute gel extraction kit, and capillary sequenced (Australian

Genome Research Facility, Brisbane) to verify the genomic location and structural hallmarks of each

insertion.

3'RACE: For each construct, $1x10^5$ HeLa-JVM cells were plated per well of a 6-well dish and transfected

~18 hours later with 1ug of plasmid DNA using 3ul of FugeneHD and 97ul of Opti-MEM as described

above. Cells were selected for 7 days using 200ug/ml of Hygromycin B, and total RNA was extracted

using the Qiagen RNeasy Mini kit (cat no. 74104). 3'RACE was performed using the 3'RACE module of

the Invitrogen First Choice RLM-RACE kit (cat no. AM1700) according to manufacturer's instructions.

Briefly, 1ug of total RNA was used in a reverse transcription reaction with kit provided M-MLV reverse

transcriptase,   RNase   inhibitor,   and   3'RACE   adapter   (5'-

GCGAGCACAGAATTAATACGACTCACTATAGGT12VN-3'). RNA was reverse transcribed at 42°C for 1

hour. PCR amplification of 3'RACE products was performed using gene-specific primers to the 5' region

of the L1spa 3'UTR (3'RACE_3'UTR-3 CGAAAGGACCCAGATGTAGC), the antisense SV40 early promoter

within the NEO retrotransposition indicator cassette (asSV40prom_F CGGGACTATGGTTGCTGACT),

and the 3' region of the hygromycin resistance gene in the pCEP4 plasmid backbone (Hygro_F3

ATTTCGGCTCCAACAATGTC) in combination with the adaptor specific primer from the RLM-RACE kit

(3'RACE Outer Primer 5'-GCGAGCACAGAATTAATACGACT-3'). PCR cycling conditions were as follows:

3 minutes at 94°C, 35 cycles of 94°C 30 seconds, 55°C 30 seconds, 72°C 30 seconds followed by 7

minutes at 72°C. PCR products were run on a 2% agarose gel, and bands were excised and purified

using the Qiagen MinElute gel extraction kit (cat no. 28604). Purified products were cloned using TA

Cloning Kit (Invitrogen, K204001.) and sequenced using primers T7 TAATACGACTCACTATAGGG and

Neosplice_F2 AGTGACAACGTCGAGCACAG by AGRF Brisbane.

Acknowledgements: HeLa-JVM cells were a gift from John V. Moran. The pTN201/L1spa construct was

a gift from Haig H. Kazazian. We thank members of the Richardson and Faulkner laboratories for

helpful discussion and critical reading of the manuscript. J.D.B. is a Founder and Director of CDI Labs,

Inc., a Founder of Neochromosome, Inc, a Founder and SAB member of ReOpen Diagnostics, and

serves or served on the Scientific Advisory Board of the following: Sangamo, Inc., Modern Meadow,

Inc., Sample6, Inc. and the Wyss Institute.

References

1.   Han JS, Boeke JD. A highly active synthetic mammalian retrotransposon. Nature.
2004;429(6989):314-8.
2.   Richardson SR, Doucet AJ, Kopera HC, Moldovan JB, Garcia-Perez JL, Moran JV. The influence
of LINE-1 and SINE retrotransposons on mammalian genomes. Microbiology Spectrum. 2015;3(2).
3.   Kazazian HH, Jr., Moran JV. Mobile DNA in Health and Disease. N Engl J Med. 2017;377(4):361-
70.
4.   Goodier JL, Ostertag EM, Du K, Kazazian HH, Jr. A novel active L1 retrotransposon subfamily in
the mouse. Genome Res. 2001;11(10):1677-85.
5.   Naas TP, DeBerardinis RJ, Moran JV, Ostertag EM, Kingsmore SF, Seldin MF, et al. An actively
retrotransposing, novel subfamily of mouse L1 elements. EMBO J. 1998;17(2):590-7.

6.   Sookdeo A, Hepp CM, McClure MA, Boissinot S. Revisiting the evolution of mouse LINE-1 in
the genomic era. Mobile DNA. 2013;4(1):3.

7.   DeBerardinis RJ, Goodier JL, Ostertag EM, Kazazian HH, Jr. Rapid amplification of a
retrotransposon subfamily is evolving the mouse genome. Nat Genet. 1998;20(3):288-90.

8.   Penzkofer T, Jager M, Figlerowicz M, Badge R, Mundlos S, Robinson PN, et al. L1Base 2: more
retrotransposition-active LINE-1s, more mammalian genomes. Nucleic Acids Res. 2017;45(D1):D68-
D73.

9.   Zhou M, Smith AD. Subtype classification and functional annotation of L1Md retrotransposon
promoters. Mobile DNA. 2019;10:14.

10.   Luan DD, Korman MH, Jakubczak JL, Eickbush TH. Reverse transcription of R2Bm RNA is primed
by a nick at the chromosomal target site: a mechanism for non-LTR retrotransposition. Cell.
1993;72(4):595-605.

11.   Feng Q, Moran JV, Kazazian HH, Jr., Boeke JD. Human L1 retrotransposon encodes a conserved
endonuclease required for retrotransposition. Cell. 1996;87(5):905-16.

12.   Mathias SL, Scott AF, Kazazian HH, Jr., Boeke JD, Gabriel A. Reverse transcriptase encoded by
a human transposable element. Science. 1991;254(5039):1808-10.

13.   Kolosha VO, Martin SL. In vitro properties of the first ORF protein from mouse LINE-1 support
its role in ribonucleoprotein particle formation during retrotransposition. Proc Natl Acad Sci U S A.
1997;94(19):10155-60.

14.   Kolosha VO, Martin SL. High-affinity, non-sequence-specific RNA binding by the open reading
frame 1 (ORF1) protein from long interspersed nuclear element 1 (LINE-1). J Biol Chem.
2003;278(10):8112-7.

15.   Khazina E, Weichenrieder O. Non-LTR retrotransposons encode noncanonical RRM domains
in their first open reading frame. Proc Natl Acad Sci U S A. 2009;106(3):731-6.

16.   Khazina E, Truffault V, Buttner R, Schmidt S, Coles M, Weichenrieder O. Trimeric structure and
flexibility of the L1ORF1 protein in human L1 retrotransposition. Nat Struct Mol Biol. 2011;18(9):1006-
14.

17.   Baldwin ET, van Eeuwen T, Hoyos D, Zalevsky A, Tchesnokov EP, Sanchez R, et al. Structures,
functions and adaptations of the human LINE-1 ORF2 protein. Nature. 2024;626(7997):194-206.

18.   Thawani A, Ariza AJF, Nogales E, Collins K. Template and target-site recognition by human
LINE-1 in retrotransposition. Nature. 2024;626(7997):186-93.

19.   Howell R, Usdin K. The ability to form intrastrand tetraplexes is an evolutionarily conserved
feature of the 3' end of L1 retrotransposons. Mol Biol Evol. 1997;14(2):144-55.

20.   Boissinot S, Sookdeo A. The Evolution of LINE-1 in Vertebrates. Genome Biol Evol.
2016;8(12):3485-507.

21.   Goodier JL, Ostertag EM, Kazazian HH, Jr. Transduction of 3'-flanking sequences is common in
L1 retrotransposition. Hum Mol Genet. 2000;9(4):653-7.

22.   Moran JV, DeBerardinis RJ, Kazazian HH, Jr. Exon shuffling by L1 retrotransposition. Science.
1999;283(5407):1530-4.

23.   Pickeral OK, Makalowski W, Boguski MS, Boeke JD. Frequent human genomic DNA
transduction driven by LINE-1 retrotransposition. Genome Res. 2000;10(4):411-5.

24.   Jurka J. Sequence patterns indicate an enzymatic involvement in integration of mammalian
retroposons. Proc Natl Acad Sci U S A. 1997;94(5):1872-7.

25.   Cost GJ, Boeke JD. Targeting of human retrotransposon integration is directed by the
specificity of the L1 endonuclease for regions of unusual DNA structure. Biochemistry.
1998;37(51):18081-93.

26.   Gilbert N, Lutz S, Morrish TA, Moran JV. Multiple fates of L1 retrotransposition intermediates
in cultured human cells. Mol Cell Biol. 2005;25(17):7780-95.

27.   Gilbert N, Lutz-Prigge S, Moran JV. Genomic deletions created upon LINE-1 retrotransposition.
Cell. 2002;110(3):315-25.

28.   Sultana T, van Essen D, Siol O, Bailly-Bechet M, Philippe C, Zine El Aabidine A, et al. The
Landscape of L1 Retrotransposons in the Human Genome Is Shaped by Pre-insertion Sequence Biases
and Post-insertion Selection. Mol Cell. 2019;74(3):555-70 e7.

29.   Richardson SR, Gerdes P, Gerhardt DJ, Sanchez-Luque FJ, Bodea GO, Munoz-Lopez M, et al.
Heritable L1 retrotransposition in the mouse primordial germline and early embryo. Genome Res.
2017;27(8):1395-405.

30.   Feusier J, Watkins WS, Thomas J, Farrell A, Witherspoon DJ, Baird L, et al. Pedigree-based
estimation of human mobile element retrotransposition rates. Genome Res. 2019;29(10):1567-77.

31.   Hancks DC, Kazazian HH, Jr. Roles for retrotransposon insertions in human disease. Mobile
DNA. 2016;7:9.

32.   Gagnier L, Belancio VP, Mager DL. Mouse germ line mutations due to retrotransposon
insertions. Mobile DNA. 2019;10:15.

33.   Moran JV, Holmes SE, Naas TP, DeBerardinis RJ, Boeke JD, Kazazian HH, Jr. High frequency
retrotransposition in cultured mammalian cells. Cell. 1996;87(5):917-27.

34.   Wei W, Morrish TA, Alisch RS, Moran JV. A transient assay reveals that cultured human cells
can accommodate multiple LINE-1 retrotransposition events. Anal Biochem. 2000;284(2):435-8.

35.   Ostertag EM, DeBerardinis RJ, Goodier JL, Zhang Y, Yang N, Gerton GL, et al. A mouse model
of human L1 retrotransposition. Nat Genet. 2002;32(4):655-60.

36.   Freeman JD, Goodchild NL, Mager DL. A modified indicator gene for selection of
retrotransposition events in mammalian cells. Biotechniques. 1994;17(1):46, 8-9, 52.

37.   Rangwala SH, Kazazian HH, Jr. The L1 retrotransposition assay: a retrospective and toolkit.
Methods. 2009;49(3):219-26.

38.   Han JS, Szak ST, Boeke JD. Transcriptional disruption by the L1 retrotransposon and
implications for mammalian transcriptomes. Nature. 2004;429(6989):268-74.

39.   Ilik IA, Glazar P, Tse K, Brandl B, Meierhofer D, Muller FJ, et al. Autonomous transposons tune
their sequences to ensure somatic suppression. Nature. 2024;626(8001):1116-24.

40.   Seczynska M, Bloor S, Cuesta SM, Lehner PJ. Genome surveillance by HUSH-mediated silencing
of intronless mobile elements. Nature. 2022;601(7893):440-5.

41.   Kingsmore SF, Giros B, Suh D, Bieniarz M, Caron MG, Seldin MF. Glycine receptor beta-subunit
gene mutation in spastic mouse associated with LINE-1 element insertion. Nat Genet. 1994;7(2):136-
41.

42.   An W, Dai L, Niewiadomska AM, Yetil A, O'Donnell KA, Han JS, et al. Characterization of a
synthetic human LINE-1 retrotransposon ORFeus-Hs. Mobile DNA. 2011;2(1):2.

43.   Schauer SN, Carreira PE, Shukla R, Gerhardt DJ, Gerdes P, Sanchez-Luque FJ, et al. L1
retrotransposition is a common feature of mammalian hepatocarcinogenesis. Genome Research.
2018;28(5).

44.   Martin SL, Bushman D, Wang F, Li PW, Walker A, Cummiskey J, et al. A single amino acid
substitution in ORF1 dramatically decreases L1 retrotransposition and provides insight into nucleic
acid chaperone activity. Nucleic Acids Res. 2008;36(18):5845-54.

45.   Belotserkovskii BP, Liu R, Tornaletti S, Krasilnikova MM, Mirkin SM, Hanawalt PC. Mechanisms
and implications of transcription blockage by guanine-rich DNA sequences. Proc Natl Acad Sci U S A.
2010;107(29):12816-21.

46.   Belotserkovskii BP, Neil AJ, Saleh SS, Shin JH, Mirkin SM, Hanawalt PC. Transcription blockage
by homopurine DNA sequences: role of sequence composition and single-strand breaks. Nucleic Acids
Res. 2013;41(3):1817-28.

47.   Broxson C, Beckett J, Tornaletti S. Transcription arrest by a G quadruplex forming-trinucleotide
repeat sequence from the human c-myb gene. Biochemistry. 2011;50(19):4162-72.

48.   Belotserkovskii BP, Soo Shin JH, Hanawalt PC. Strong transcription blockage mediated by R-
loop formation within a G-rich homopurine-homopyrimidine sequence localized in the vicinity of the
promoter. Nucleic Acids Res. 2017;45(11):6589-99.

49.   Beaudoin JD, Perreault JP. Exploring mRNA 3'-UTR G-quadruplexes: evidence of roles in both
alternative polyadenylation and mRNA shortening. Nucleic Acids Res. 2013;41(11):5898-911.
50.   Newkirk SJ, Lee S, Grandi FC, Gaysinskaya V, Rosser JM, Vanden Berg N, et al. Intact piRNA
pathway prevents L1 mobilization in male meiosis. Proc Natl Acad Sci U S A. 2017;114(28):E5635-E44.
51.   Kannan M, Li J, Fritz SE, Husarek KE, Sanford JC, Sullivan TL, et al. Dynamic silencing of somatic
L1 retrotransposon insertions reflects the developmental and cellular contexts of their genomic
integration. Mobile DNA. 2017;8:8.
52.   Richardson SR, Narvaiza I, Planegger RA, Weitzman MD, Moran JV. APOBEC3A deaminates
transiently exposed single-strand DNA during LINE-1 retrotransposition. eLife. 2014;2014(3).
53.   O'Donnell KA, An W, Schrum CT, Wheelan SJ, Boeke JD. Controlled insertional mutagenesis
using a LINE-1 (ORFeus) gene-trap mouse model. Proc Natl Acad Sci U S A. 2013;110(29):E2706-13.
54.   Monot C, Kuciak M, Viollet S, Mir AA, Gabus C, Darlix JL, et al. The specificity and flexibility of
l1 reverse transcription priming at imperfect T-tracts. PLoS Genet. 2013;9(5):e1003499.
55.   Xie Y, Mates L, Ivics Z, Izsvak Z, Martin SL, An W. Cell division promotes efficient
retrotransposition in a stable L1 reporter cell line. Mobile DNA. 2013;4(1):10.
56.   Grandi FC, Rosser JM, An W. LINE-1-derived poly(A) microsatellites undergo rapid shortening
and create somatic and germline mosaicism in mice. Mol Biol Evol. 2013;30(3):503-12.
57.   An W, Han JS, Schrum CM, Maitra A, Koentgen F, Boeke JD. Conditional activation of a single-
copy L1 transgene in mice by Cre. Genesis. 2008;46(7):373-83.
58.   Alisch RS, Garcia-Perez JL, Muotri AR, Gage FH, Moran JV. Unconventional translation of
mammalian LINE-1 retrotransposons. Genes Dev. 2006;20(2):210-24.
59.   Wagstaff BJ, Barnerssoi M, Roy-Engel AM. Evolutionary conservation of the functional
modularity of primate and murine LINE-1 elements. PLoS One. 2011;6(5):e19672.
60.   Kimberland ML, Divoky V, Prchal J, Schwahn U, Berger W, Kazazian HH, Jr. Full-length human
L1 insertions retain the capacity for high frequency retrotransposition in cultured cells. Hum Mol
Genet. 1999;8(8):1557-60.
61.   Sahakyan AB, Murat P, Mayer C, Balasubramanian S. G-quadruplex structures within the 3'
UTR of LINE-1 elements stimulate retrotransposition. Nat Struct Mol Biol. 2017;24(3):243-7.
62.   Lexa M, Steflova P, Martinek T, Vorlickova M, Vyskot B, Kejnovsky E. Guanine quadruplexes
are formed by specific regions of human transposable elements. BMC Genomics. 2014;15:1032.
63.   Moran JV. Human L1 retrotransposition: insights and peculiarities learned from a cultured cell
retrotransposition assay. Genetica. 1999;107(1-3):39-51.
64.   An W, Han JS, Wheelan SJ, Davis ES, Coombes CE, Ye P, et al. Active retrotransposition by a
synthetic L1 element in mice. Proc Natl Acad Sci U S A. 2006;103(49):18662-7.
65.   Morrish TA, Gilbert N, Myers JS, Vincent BJ, Stamato TD, Taccioli GE, et al. DNA repair
mediated by endonuclease-independent LINE-1 retrotransposition. Nat Genet. 2002;31(2):159-65.
66.   Li J, Shen H, Himmel KL, Dupuy AJ, Largaespada DA, Nakamura T, et al. Leukaemia disease
genes: large-scale cloning and pathway predictions. Nat Genet. 1999;23(3):348-53.
67.   Isomura H, Stinski MF, Kudoh A, Nakayama S, Murata T, Sato Y, et al. A cis element between
the TATA Box and the transcription start site of the major immediate-early promoter of human
cytomegalovirus determines efficiency of viral replication. J Virol. 2008;82(2):849-58.

Supplemental Figure 1: Alignment of the 3'UTRs and flanking sequence of pTN201/L1spa and smL1.
Alignment depicts the 3' end of ORF2 (L1spa ORF2, dark red; smL1 ORF2, bright red), followed by the
3'UTR (L1spa 3'UTR dark blue; smL1 3'UTR, light blue). The smL1 3'UTR deletion is highlighted in
yellow, and the runs of guanine bases in the L1spa 3'UTR are highlighted in red. The positions of
engineered restriction sites in each construct are shown in pink, and the NdeI sites resulting in the
GRR deletion are shown in green. The HSV Tk polyadenylation signal belonging to the NEO cassette
(purple) and the second exon of the NEO cassette (orange) are identical between pTN201/L1spa and
smL1.

Supplemental Figure 2: Deletion of the GRR during cloning of the smL1 reporter construct. A. The
pTN201/L1spa construct(5) and an intermediate pBluescript clone containing the smL1 sequence, with
features annotated as in Figure 1. The positions of known NdeI and BamHI sites are shown in black.
Our map of pTN201 did not include 15 base pairs of human L1 sequence present in pTN201
immediately following the L1spa 3'UTR, which was carried over from subcloning described in(5). This
15 bp of human L1 sequence contained an NdeI site (shown in red).

B) Deletion of the GRR-containing NdeI fragment during cloning. Left: we intended to move an
NdeI/BamHI fragment containing the GRR of the L1spa element and the NEO cassette downstream of
the smL1 ORFs and 5' portion of the 3'UTR in pBluescript. Right: due to the unanticipated NdeI site, a
180 bp NdeI fragment containing the terminal GRR-containing 159 bp of the L1spa 3'UTR and an
engineered PacI site (not shown) was deleted from the resulting construct. Owing to its small size, the
deletion was not detected when screening the constructs with diagnostic restriction digests.

Supplemental Figure 3: Detailed structures of L1spa_Dbl_smBB_GRRAN integrants characterized via
inverse PCR. The structure of each insertion is presented as follows: L1spa 5'UTR (triangles), ORFs
(white rectangles), first part of the smL1 3'UTR (light blue), spliced NEO cassette (green), sv40
promoter (grey), GRR (red) sv40 polyadenylation signal (purple). The 5' truncation point for each
insertion is shown. Insertions I and J arose from a transcript initiating upstream of the L1 5'UTR, at the
transcription start site of the CMV promoter within the pCEP4 vector(67) denoted as position -43.
Insertion J had a 5' inversion with breakpoint at position 1493 shown. Flanking genomic DNA is shown
to the right and left, with target-site duplications in red. The empty site is shown below, with TSD
sequence in red, the first-strand cleavage site indicated by a black arrow and the second-strand
cleavage site indicated by a grey arrow. Insertion A had a 2 bp target-site deletion, depicted in blue.
The chromosomal coordinates of the TSD sequence and the endonuclease cleavage motif for each
insertion is shown.

Supplemental Table 1: The first tab contains the sequences of L1spa (pTN201) and smL1 as
determined by capillary sequencing. The second tab contains the sequences (iPCR_L1_rev and
iPCR_L1_fwd) of inverse PCR products characterizing of L1spa_Dbl_smBB_GRRAN integrants and the
corresponding validation primers. The third tab contains retrotransposition assay data (average
colony counts from three technical replicate wells, normalized to transfection efficiency) and
mean/standard deviation values for three biological replicates used to generate the graphs shown in
Figure 1C, Figure 2, and Figure 4 A and C.

Supplemental Figure 1

```
                     L1Spa Orf2
L1spa  6742  TCCTGAGTGAGGTAACACAATCACAAAGGAACTCACACAATATGTACTCACTGATAAGTG  6801
             ||||||| ||||| || ||    || | |||  ||||| ||||||  ||||| || |
L1SM   6777  TCCTGAGCGAGGTGACCCAGAGCCAGCGCAACAGCCACAACATGTACAGCCTGATCAGCG  6836
             L1SM Orf2
                      SacI   L1spa 3'UTR
L1spa  6802  GATACTAGCGAGCTC-CAAAACCTAGGATACCCACGATATAAGATACAATTTCCTAAACA  6860
             | |||||||      |         |||||||||||||||||||||| ||||||||
L1SM   6837  GCTACTAGCCTAAAACCGGT------GATACCCACGATATAAGATACAATTGCCTAAACA  6890
                      AgeI         L1SM 3'UTR

L1spa  6861  CATGAAACTCAAGAAAAATGAAGACTGAAGTGTGGACACTATGCCCCTCCTTAGAAGTGG  6920
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
L1SM   6891  CATGAAACTCAAGAAAAATGAAGACTGAAGTGTGGACACTATGCCCCTCCTTAGAAGTGG  6950

L1spa  6921  GAACAAAACACCCATGGAAGGAGTTACAGAAACAAAGTTTGGAGCTGAGATGAAAGGAGG  6980
             ||||||||| |||| ||| ||||||||||||||||||||||||||||||||||||||||
L1SM   6951  GAACAAAACAGCCATAGAAAGAGTTACAGAAACAAAGTTTGGAGCTGAGATGAAAGGAGA  7010

L1spa  6981  GACCATGTAGAGACTGCCATATCCAGGGATCCACCCCATAATCAGCATCCAAACGC--TG  7038
             |||||||||| ||||||||||||||||| |||||||||||||||||||||||||  || ||
L1SM   7011  GACCATGTAGAAACTGCCATATCCAGGGTTCCACCCCATAATCAGCATCCAA--GCTTTG  7068

L1spa  7039  ACACCATTGCATATACTAGCAAGATTTTATCGAAAGGACCCAGATGTAGCTGTCTCTTGT  7098
             |||||||||||||||||||| |||||||||||||||||||||||||||||||||||||||
L1SM   7069  ACACCATTGCATATACTAGGAAGATTTTATCGAAAGGACCCAGATGTAGCTGTCTCTTGT  7128

L1spa  7099  GAGACTATGCCGGGGCCTAGCAAACACAGAAGTGGATGCTCACAGTCAGCTAATGGATGG  7158
             ||||||||||||||||||||||| ||||||||||||||||||||||||||| |||||||
L1SM   7129  GAGACTATGCCGGGGCCTAGCAA-CACAGAAGTGGATGCTCACAGTCAGCTA-TGGATGG  7186

L1spa  7159  ATCACAGGGCTCCCAATGGAGGAGCTAGAGAAAGTACCCAAGGAGCTAAAGGGATCTGCA  7218
             ||||||||| |||||||||||||||||||||||||||||||||||||||||||||||||
L1SM   7187  ATCACAGGGCTTTCAATGGAGGAGCTAGAGAAAGTACCCAAGGAGCTAAAGGGATCTGCA  7246

L1spa  7219  ACCCTATAGGTGGAACAACATTATGAAC-TAACCAGTACCCCTGAGCTCTTGACTCTAGC  7277
             || |||||||||||||||||||||||||  |||||||||||| |||||||||| | |||||
L1SM   7247  ACTCTATAGGTGGAACAACATTATGA-GTTAACCAGTACCCTTGAGCTCTTGTCGCTAGC  7305

L1spa  7278  TGCATATGTATCAAAAGATGGCCTAGTCGGCCATCACTGGAAAGAGAGGCCCATTGGACA  7337
             ||||||||||
L1SM   7306  TGCATATGTA--------------------------------------------------  7315

L1spa  7338  CGCAGACTTTGTGTGCCCCGGTACAGGGGAACGCCAGGGCCAAAGGGGGGGGGGAGTGGGTG  7397

L1SM   7316  ------------------------------------------------------------  7315

                                                                   PacI
L1spa  7398  GGTAGGGGAGTGGGGGTAGGTGGGTAAGGGGGACTTTTGGTATAGCATTAATTAACTCGA  7457

L1SM   7316  ------------------------------------------------------------  7315

                                   HSV Tk pA (antisense)
L1spa  7458  TACATATGTAACTAACCCGATCCGAACAAACGACCCAACACCCGTGCGTTTTATTCTGTC  7517
                          ||||||||||||||||||||||||||||||||||||||||||||||
L1SM   7316  ---------ACTAACCCGATCCGAACAAACGACCCAACACCCGTGCGTTTTATTCTGTC   7365

                    NEO (antisense)
L1spa  7518  TTTTTATTGCCGATCCCCTCAGAAGAACTCGTCAAGAAGGCGATAGAAGGCGATGCGCTG  7577
             ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
L1SM   7366  TTTTTATTGCCGATCCCCTCAGAAGAACTCGTCAAGAAGGCGATAGAAGGCGATGCGCTG  7425
```

Supplemental Figure 2

A.

B.

Supplemental Figure 3

Insertion A

5'-TTTATCCATAAAGATGGAAATCAAACTCTT **NEO** A~49 AAAATGTTCTTATATTTTTCACTAAAAG-3'
3'-AAATAGGTATTTCTACCTTTAGTTTGAGAA TTTTACAAGAATATAAAAGTGATTTTC-5'

5'-TTTATCCATAAAGATGGAAATCAAACTCTTTTAAAATGTTCTTATATTTTTCACTAAAAG-3'
3'-AAATAGGTATTTCTACCTTTAGTTTGAGAAAATTTTACAAGAATATAAAAGTGATTTTC-5'

Target-site deletion: Chr3:130997869-130997870

EN cleavage: 5'-TTTT/AA-3'

Insertion B

5'-AGCAATATGATATTAATGCAATAAAA **NEO** A~86 AAAAGAAATTCCCCCAAAGAAT-3'
3'-TCGTTATACTATAATTACGTTATTTT TTTTCTTTAAGGGGGTTTCTTA-5'

5'-TTCTAGCAATATGATATTAATGCAATAAAAGAAATTCCCCCAAAGAATACTTCCAG-3'
3'-AAGATCGTTATACTATAATTACGTTATTTTCTTTAAGGGGGTTTCTTATGAAGGTC-5'

TSD: Chr3:148918423-148918426

EN cleavage: 5'-TTTT/AT-3'

Insertion C

5'-AAACATGGATGTATAAAAAATAAAGAAT **NEO** A~104 AAAAAATAAAGAATACCTAAGAGAGTTAAA-3'
3'-TTTGTACCTACATATTTTTTATTTCTTA TTTTTTATTTCTTATGGATTCTCTCAATTT-5'

5'-GTCTTAATAAAAGAAAAAAACATGGATGTATAAAAAATAAAGAATACCTAAGAGAGTTAAATTCCCAATAAGTAC-3'
3'-CAGAATTATTTTCTTTTTTTTGTACCTACATATTTTTTATTTCTTATGGATTCTCTCAATTTAAGGGTTATTCATG-5'

TSD: Chr5: 21539952-21539965

EN cleavage: 5'-TTTT/AT-3'

Supplemental Figure 3

Insertion D

```
5'-ATAATTTTTGTGTCATGGTTTATGTTTTT
3'-TATTAAAAACACAGTACCAAATACAAAAA
```

```
AAAAAAACTTACCTTTTAACATGCTAAAAAG
TTTTTTTGAATGGAAAATTGTACGATTTTTTC
```

```
ACAGTACTTCTTGAATGGTAGAATAAAGACCA
TGTCATGAAGAACTTACCATCTTATTTCTGGT
```

```
TCAAAAATCTGCTCATCCATAAAAGCAATGAG
AGTTTTTAGACGAGTAGGTATTTTCGTTACTC
```

```
ATCAGTGGCAAAAATTGTCAAAATTAACTTTT
TAGTCACCGTTTTTAACAGTTTTAATTGAAAA
```

```
TCAGAACTCTGGAAATTAACTAAAAGATTGC
AGTCTTGAGACCTTTAATTGATTTTCTAACG
```

6327 | □ | NEO | ▨ ▮ ▮ | A~66

```
AAAAAAACTTACCTTTTAACATGCTAAAAAG
TTTTTTTGAATGGAAAATTGTACGATTTTTTC
```

```
ACAGTACTTCTTGAATGGTAGAATAAAGACCA
TGTCATGAAGAACTTACCATCTTATTTCTGGT
```

```
TCAAAAATCTGCTCATCCATAAAAGCAATGAG
AGTTTTTAGACGAGTAGGTATTTTCGTTACTC
```

```
ATCAGTGGCAAAAATTGTCAAAATTAACTTTT
TAGTCACCGTTTTTAACAGTTTTAATTGAAAA
```

```
TCAGAACTCTGGAAATTAACTAAAAGATTGC
AGTCTTGAGACCTTTAATTGATTTTCTAACG
```

TSD: Chr21: 29204205-29204047

EN cleavage: 5'-TTTT/AA-3'

```
AAAATGTCAAGGACTGTTTAGTCAGGAAAA-3'
TTTTACAGTTCCTGACAAATCAGTCCTTTT-5'
```

```
5'-TGTCATGGTTTATGTTTTTTAAAAAAACTTACCTTTTAACATGCTAAAAAAGACAGTACTTCTTGAATGGTAGAATAAAG
3'-ACAGTACCAAATACAAAAATTTTTTTGAATGGAAAATTGTACGATTTTTTCTGTCATGAAGAACTTACCATCTTATTTC
```
                              ↑

```
ACCATCAAAAATCTGCTCATCCATAAAAGCAATGAGATCAGTGGCAAAAATTGTCAAAATTAACTTTTTTCAGAACTCTGG
TGGTAGTTTTTAGACGAGTAGGTATTTTCGTTACTCTAGTCACCGTTTTTAACAGTTTTAATTGAAAAAGTCTTGAGACC
```
                ↓

```
AAATTAACTAAAAGATTGCAAAATGTCAAGGACTGTTT-3'
TTTAATTGATTTTCTAACGTTTTACAGTTCCTGACAAA-5'
```

Insertion E

4772 | □ | NEO | ▨ ▮ ▮ | A~55

```
5'-TTTATATAAACAATATATAAACAA   ACCTCAGTTAGAATATTTAATTAA-3'
3'-AAATATATTTGTTATATATTTGTT   TGGAGTCAATCTTATAAATTAATT-5'
```

```
                    ↓
5'-TTTATATAAACAATATATAAACAACCTCAGTTAGAATATTTAATTAA-3'
3'-AAATATATTTGTTATATATTTGTTGGAGTCAATCTTATAAATTAATT-5'
                    ↑
```

TSD: chr7: 153197463

EN cleavage: 5'-AGGT/TG-3'

Supplemental Figure 3

Insertion F

```
5'-AATTCTTTGGAAAAAAGACAAAGTTGAAATAATGTCAGAGAGAATACATCAGGAAAAAGGTTGAAAATGTAAGTTCTA
3'-TTAAGAAACCTTTTTTCTGTTTCAACTTTATTACAGTCTCTCTTATGTAGTCCTTTTTCCAACTTTTACATTCAAGAT
```

```
TTAAACACAACATACTTAGAGGGGGAGAAAATCACAGAAGAGCCAGTTATGGCAGCTGAAAGAGGGGCAATGCTAAAGAC
AATTTGTGTTGTATGAATCTCCCCCTCTTTTAGTGTCTTCTCGGTCAATACCGTCGACTTTCTCCCCGTTACGATTTCTG
```

```
TCACATTTTTAATTTAAAAATCAGTGTTCCCTGGGCAGCAAGAAATTTTTGCCTATACTATATACAAAAAATTCATTCAA
AGTGTAAAAATTAAATTTTTAGTCACAAGGGACCCGTCGTTCTTTAAAAACGGATATGATATATGTTTTTTAAGTAAGTT
```

```
AGAGAATAAAAGACCTAAACCTGAGAGCTAAAACTCTAAAACCTTTAGAAGAAAACATAAGGGAAATGCTTCATGACATT
TCTCTTATTTTCTGGATTTGGACTCTCGATTTTGAGATTTTGGAAATCTTCTTTTGTATTCCCTTTACGAAGTACTGTAA
```

```
GAACTTTACAGTGGTGTCTTGGATATGATACCAAAAGCATGGACAATGAAAGAAAAAAAACACATAGATTTCATCGAAG
CTTGAAATGTCACCACAGAACCTATACTATGGTTTTCGTACCTGTTACTTTCTTTTTTTTTGTGTATCTAAAGTAGCTTC
```

```
TAAAAACTTCTGTGCATTAAGTAACATTC          GAAAATGTAAGTTCTATTAAACACAACAT
ATTTTTGAAGACACGTAATTCATTGTAAG          CTTTTACATTCAAGATAATTTGTGTTGTA
```

```
ACTTAGAGGGGGAGAAAATCACAGAAGAGCCAGTTATGGCAGCTGAAAGAGGGGCAATGCTAAAGACTCACATTTTTAAT
TGAATCTCCCCCTCTTTTAGTGTCTTCTCGGTCAATACCGTCGACTTTCTCCCCGTTACGATTTCTGAGTGTAAAAATTA
```

```
TTAAAAATCAGTGTTCCCTGGGCAGCAAGAAATTTTTGCCTATACTATATACAAAAAATTCATTCAAAGAGAATAAAAGA
AATTTTTAGTCACAAGGGACCCGTCGTTCTTTAAAAACGGATATGATATATGTTTTTTAAGTAAGTTTCTCTTATTTTCT
```

```
CCTAAACCTGAGAGCTAAAACTCTAAAACCTTTAGAAGAAAACATAAGGGAAATGCTTCATGACATTGAACTTTACAGTG
GGATTTGGACTCTCGATTTTGAGATTTTGGAAATCTTCTTTTGTATTCCCTTTACGAAGTACTGTAACTTGAAATGTCAC
```

```
GTGTCTTGGATATGATACCAAAAGCATGGACAATGAAAGAAAAAAAACACATAGATTTCATCGAAGTAAAAACTTCTGT
CACAGAACCTATACTATGGTTTTCGTACCTGTTACTTTCTTTTTTTTTGTGTATCTAAAGTAGCTTCATTTTTGAAGACA
```

```
GCATTAAGTAACATTCTCAACAGAGTGAAAAGTCACAGAGGAAATATTTGCAAATCATATATCTGATAAGGGATTAATA-3'
CGTAATTCATTGTAAGAGTTGTCTCACTTTTCAGTGTCTCCTTTATAAACGTTTAGTATATAGACTATTCCCTAATTAT-5'
```

```
5'-AATTCTTTGGAAAAAAGACAAAGTTGAAATAATGTCAGAGAGAATACATCAGGAAAAAGGTTGAAAATGTAAGTTCTA
3'-TTAAGAAACCTTTTTTCTGTTTCAACTTTATTACAGTCTCTCTTATGTAGTCCTTTTTCCAACTTTTACATTCAAGAT
                                                                        ↑
```

```
TTAAACACAACATACTTAGAGGGGGAGAAAATCACAGAAGAGCCAGTTATGGCAGCTGAAAGAGGGGCAATGCTAAAGAC
AATTTGTGTTGTATGAATCTCCCCCTCTTTTAGTGTCTTCTCGGTCAATACCGTCGACTTTCTCCCCGTTACGATTTCTG
```

```
TCACATTTTTAATTTAAAAATCAGTGTTCCCTGGGCAGCAAGAAATTTTTGCCTATACTATATACAAAAAATTCATTCAA
AGTGTAAAAATTAAATTTTTAGTCACAAGGGACCCGTCGTTCTTTAAAAACGGATATGATATATGTTTTTTAAGTAAGTT
```

```
AGAGAATAAAAGACCTAAACCTGAGAGCTAAAACTCTAAAACCTTTAGAAGAAAACATAAGGGAAATGCTTCATGACATT
TCTCTTATTTTCTGGATTTGGACTCTCGATTTTGAGATTTTGGAAATCTTCTTTTGTATTCCCTTTACGAAGTACTGTAA
```

```
GAACTTTACAGTGGTGTCTTGGATATGATACCAAAAGCATGGACAATGAAAGAAAAAAAACACATAGATTTCATCGAAG
CTTGAAATGTCACCACAGAACCTATACTATGGTTTTCGTACCTGTTACTTTCTTTTTTTTTGTGTATCTAAAGTAGCTTC
```

```
                                                   ↓
TAAAAACTTCTGTGCATTAAGTAACATTCTCAACAGAGTGAAAAGTCACAGAGGAAATATTTGCAAATCATATATCTG-3'
ATTTTTGAAGACACGTAATTCATTGTAAGAGTTGTCTCACTTTTCAGTGTCTCCTTTATAAACGTTTAGTATATAGAC-5'
```

TSD: Chr12:77126920-77127284

EN cleavage: 5'-TTTC/AA3'