1

Germline piRNAs counteract endogenous retrovirus invasion from somatic cells.

Marianne Yoth[1], Nathalie Gueguen[1], Silke Jensen[1]*, Emilie Brasset[1]*

[1] iGReD, Université Clermont Auvergne, CNRS, INSERM, Faculté de Médecine, 63000 Clermont-Ferrand, France.

* Correspondence: emilie.brasset@uca.fr, silke.jensen@uca.fr

Abstract

In metazoan gonads, transposable elements (TEs) mobilization is limited by PIWI-interacting RNAs (piRNAs). These small RNAs originate from specific source loci, the piRNA clusters. piRNAs are known to silence TEs in the cells where they are produced. Endogenous retroviruses (ERVs), a subclass of TEs, pose a particular threat because they are capable of transiting from cell to cell. In this study, we reveal that piRNAs produced locally in germ cells counteract invasion by ERVs arriving from adjacent somatic cells. We reactivated the Drosophila ERV *ZAM* in somatic gonadal cells by deleting, using CRISPR-Cas9 genome editing, its single copy in the somatic *flamenco* piRNA cluster, while keeping the piRNA pathway fully functional. Upon reactivation, *ZAM* invaded the oocytes, resulting in transposition and severe fertility defects. We show that once *ZAM*-piRNAs are produced in germ cells they counter the invasion. Our study sheds new light on the mechanisms of recognition and regulation of invasive genetic elements, which is essential for the maintenance of genome integrity.

Introduction

The gonad is a major organ in animals because of its role in producing germ cells, the only cells that can transfer genetic information to the next generation. Germ cells require a set of supporting somatic cells, called gonadal somatic cells. Studies in model organisms, such as *Caenorhabditis elegans*, mice and Drosophila, revealed the essential role of somatic cells in regulating germ cell development [1–5].

Transposable elements (TEs), discovered by Barbara McClintock in the 1940s [6], are mobile genetic entities that are present in nearly all organisms. Genome integrity relies on transcriptional and post-transcriptional repression of TE activity. In animal gonads, the genome defense against TEs is based essentially on the PIWI-interacting RNA (piRNA) pathway. piRNAs, the central players in this regulatory pathway, are small (23 to 29 nucleotides) RNAs that interact with proteins of the PIWI family [7–9]. They originate from transposon-rich genomic loci called piRNA clusters [10–12]. In Drosophila ovaries, there are significant differences of TE control between germline and somatic cells. Transposons have been classified as "germline", "somatic" or "intermediate" TEs, according to the origin of the piRNAs that target them [13]. Indeed, there are germline- and soma-expressed piRNA clusters. In germ cells, most piRNA clusters are expressed in a bidirectional manner and produce piRNAs from both genomic strands. Other clusters are specifically expressed in the somatic cells that surround the ovary in a unidirectional manner and produce piRNAs that are predominantly antisense to transposons [10]. Nonetheless, the relationship between piRNA clusters and TE control is incompletely understood. Surprisingly, while depletion of piRNA pathway components in the germline or somatic lineage leads to TE derepression and sterility, the deletion of the three largest germline piRNA clusters (*42AB*, *38C*, and *20A*) does not have any impact on transposon regulation or on fertility [14–16]. Conversely, *flamenco (flam)*, the major somatic piRNA cluster, is directly

involved in the regulation of at least three TEs: *ZAM*, *Idefix,* and *Gypsy* [17–19] and female fertility

is affected when *flam* is mutated [20–22].

In the RevI-H2 line, in which the proximal part of *flam* is deleted (i.e., the region containing its

only *ZAM* insertions), *ZAM* is derepressed in the somatic cells of the ovary. We previously

showed that *ZAM* somatic piRNA production is abolished, whereas germline sense and

antisense *ZAM*-derived piRNAs are newly produced [23,24]. As *Gypsy, ZAM* is an endogenous

retrovirus (ERV), a class of TEs capable of transiting from cell to cell. However, whether and

how germline piRNAs contribute to protect the germline against invasion of TEs coming from

the surrounding somatic cells had not been investigated yet.

Here we report, using *ZAM* as a model, that when an ERV is derepressed in the somatic cells

surrounding germ cells, it invades the germline. Furthermore, no initial small RNA response

takes place after invasion of the germ cells. Many years after the *flam* partial deletion appeared

in the RevI-H2 line, *ZAM* is still expressed in somatic cells that surround germ cells and no new

ZAM insertion in the *flam* piRNA cluster has been detected. However, once the germline

produces piRNAs against this new invader, either from newly inserted dispersed copies or

because the invading TE has jumped into a piRNA cluster expressed in the germline, invasion

is arrested while the retrotransposon is still activated in the surrounding tissue. Our findings

reveal the critical role of the germline piRNA pathway as an effective defense mechanism to

counteract invasion by TEs from an adjacent tissue. Our work provides a unique opportunity to

better understand the role of each piRNA cluster type in TE regulation at the cellular and tissue

scales.

Results

In an ancient *flam* mutant line, *ZAM* is still expressed and no new *ZAM* copy is inserted in the *flam* locus.

In the RevI-H2 line, in which a deletion of the proximal part of *flam* eliminats the region in which *ZAM* was inserted, *ZAM* piRNA production by *flam* is completely lost in follicle cells, which are somatic cells of the ovary [23]. RevI-H2i2, RevI-H2i3 and RevI-H2i4 are three isogenic lines derived from the RevI-H2 line that has more than 25 years of laboratory history [25]. *ZAM* actively transposed in the initial RevI-H2 line, but it was not known whether *ZAM* is still transcriptionally active in the RevI-H2-derived lines. We found that these three isogenic RevI-H2 lines have a very similar *ZAM* expression pattern (**Fig 1a, Supplementary Fig. S1a**). Indeed, single-molecule RNA fluorescence *in situ* hybridization (smRNA FISH) experiments revealed that *ZAM* RNA was produced very early during oogenesis in the somatic cells of the germarium. Then, after stage 8, *ZAM* transcripts accumulated in a patch of follicle cells located at the posterior side of the follicles (**Fig. 1a**). We did not detect any *ZAM* staining in germ cells (nurse cells or oocytes). RNase A treatment led to complete loss of *ZAM* staining in the follicle cells (**Supplementary Fig. S1b**). We did not detect any *ZAM* RNA in the Iso1A, w[1118] and w[IR6] control ovaries (**Supplementary Fig. S1c**). Then, we used RT-qPCR to confirm that *ZAM* was derepressed in the RevI-H2i2 ovaries. Conversely, other soma-specific TEs that are not included in the deletion, such as *Gypsy*, and germline-specific TEs, such as *Burdock*, were not upregulated (**Fig. 1b**). As *ZAM* is an endogenous retrovirus that encodes Gag, Pol and Env proteins, we showed by immunostaining that Gag and Env proteins accumulated in the posterior follicle cells in late follicles (> stage 9) (**Fig. 1c**). All these results showed that *ZAM* is still expressed in the RevI-H2i2 line. The long reads obtained by Oxford Nanopore Technology (ONT) sequencing of the RevI-H2i2 genome allowed identifying *ZAM* insertions, even in

repeated sequences, but no *ZAM* insertion was present in the *flam* piRNA cluster (**Fig. 2a**).

Thus, no *ZAM* piRNA is produced from *flam*, the major somatic piRNA source locus,

explaining why *ZAM* was still expressed in follicle cells in the RevI-H2i2 line. Although *ZAM*

was expressed in follicle cells, the RevIH2i2 line is fertile. Indeed, the ovary morphology and

the number of eggs laid were comparable in a RevIH2i2 and a control female (**Fig. 1d, e**).

Altogether, these results suggest that *ZAM* expression in follicle cells has no major deleterious

effect in the RevI-H2i2 line.

RevI-H2i2 is a stable line although *ZAM* is actively expressed in follicle cells.

The ONT long-read sequencing data allowed identifying 18 *ZAM* copies in the RevI-H2i2

genome (**Fig. 2a**). Only one of these *ZAM* copies was also present in the reference genome

(http://flybase.org). The other 17 new insertions were very similar to the *ZAM* reference element

identified in the initial RevI-H2 line [25]. Specifically, nine were full-length *ZAM* elements, one

harbored a deleted 5'-UTR, three had large internal deletions in the coding regions, and five

were what we called "*ZAM*-variants", i.e. full-length insertions with a small deletion at the C-

terminal end of the *pol* gene. This small deletion corresponds to positions 5500 to 6126 in the

ZAM internal sequence [25] (Repbase *ZAM*_I, https://www.girinst.org/repbase/ [26]). This deleted

part of the *pol* gene does not correspond to any known protein domain and the fact that the five

ZAM-variants were at different locations in the RevI-H2i2 line genome indicates that this *ZAM*

copy is competent for transposition. The full-length copies and ZAM-variants all potentially

encode Gag, Pol and Env proteins. We could clearly localize 14 of the 17 new *ZAM* insertions

(**Fig. 2a** and Supplementary **Table S1** for coordinates). In the initial RevI-H2 line, a *ZAM*

insertion was found in the germline piRNA cluster *9*, which is close to the X chromosome

centromere [12,23]. Here, using ONT sequencing data, we identified three *ZAM* insertions in

piRNA cluster *9* (one full-length and two *ZAM*-variants), and also a full-length *ZAM* insertion,

which is localized in either piRNA cluster *13* or cluster *56* (uncertainty is due to flanking

repeats, piRNA clusters as in [12]).

Comparison of the Illumina sequencing data for the initial RevI-H2 line, performed ten years

ago [23] , and the ONT sequencing data showed that at least 14 of the new *ZAM* insertions in the

RevI-H2i2 line, including all *ZAM* insertions in piRNA clusters, were already present ten years

ago. This result indicates that the genomic *ZAM* profile is the same as ten years ago and suggests

that no new *ZAM* insertion occurred since then in the germline. Therefore, the RevI-H2 line is

stable.

Germline *ZAM* piRNAs counteract *ZAM* invasion from adjacent somatic cells.

The next step was to determine the mechanisms of the RevI-H2i2 line stability despite *ZAM*

expression in follicle cells. It was previously reported [27] that the *ZAM* TE actively transposed

in the RevI-H2 line. These transposition events occurred in the germline, as attested by the

transmission of the *ZAM* insertions to the offspring despite the fact that *ZAM* is specifically and

exclusively expressed in the somatic follicle cells. This indicates that *ZAM* invades the RevI-

H2 germline from follicle cells. We showed previously that germ cells of RevI-H2 flies produce

high amounts of *ZAM*-derived piRNAs with a ping-pong signature (i.e. an enrichment of 10-

nucleotide overlap between sense and antisense *ZAM*-derived piRNAs). One possible source of

these germline *ZAM*-targeting piRNAs is a *ZAM* copy inserted in the germline piRNA cluster

9 (piRNA clusters as referenced in [12]).

First, we checked whether the RevI-H2i2 line harbors the same characteristics. We found that

the RevI-H2i2 line produced sense and antisense *ZAM*-derived-piRNAs with a ping-pong

signature and silenced a *ZAM* sensor transgene in germ cells (**Fig. 2b, c, d**). We obtained similar

results in the RevI-H2i3 and RevI-H2i4 lines (**Supplementary Fig. S2a, b, c**). Moreover, in all

three RevI-H2 isogenic lines the *ZAM* copy was deleted in the *flam* piRNA cluster and *ZAM*

was inserted in the germline piRNA cluster 9 (**Supplementary Fig. S2d**).

Then, to test whether piRNAs produced in germ cells can thwart *ZAM* invasion from somatic

cells to the germline, we abolished the piRNA pathway in the germ cells by germline-specific

knock-down (GLKD), in the RevI-H2i2 genetic background, and then followed *ZAM* RNA

subcellular localization in ovaries. We first confirmed the loss of *ZAM*-targeting piRNAs. We

observed a decrease in the level of antisense piRNAs that map to *ZAM,* on the same extend as

intermediate and germ-cell-specific TEs, in ovaries in which *argonaute3* (*ago3*), *zucchini* (*zuc*),

or *piwi* (three genes encoding piRNA pathway components) was knocked down in the RevI-

H2i2 genetic background (**Fig. 3a**). No decrease was observed for *Gypsy10*, a somatic TE.

Furthermore, in *ago3*-GLKD ovaries, the ping-pong signature for *ZAM* and for many other

germline-specific TEs was abolished, while it was maintained in control *white*-GLKD and also

in *zuc-* and *piwi*-GLKD ovaries (**Fig. 3b, c**).

We then analyzed the subcellular localization of *ZAM* RNAs by smRNA FISH in the *white*-

GLKD RevI-H2i2 control line in which *ZAM*-derived piRNAs are produced in the germline,

but not in follicle cells. *ZAM* RNA expression was restricted to follicle cells and almost no *ZAM*

RNA signal was detectable in the oocyte (**Fig. 3d**). Conversely, all RevI-H2i2 lines with GLKD

of piRNA pathway actors (*vreteno (vret)*, *zuc, ago3*, *armitage (armi)*) clearly showed *ZAM*

expression in posterior follicle cells and *ZAM* RNA invasion in oocytes. *ZAM* RNA spread

throughout the ooplasm, but was clearly enriched at the posterior pole, adjacent to *ZAM*-

expressing follicle cells. Actually, 30-70% of stage 10 follicles showed a strong *ZAM* RNA

signal in oocytes (**Fig. 3d, e**). The strongest invasion phenotype was observed for the RevI-

H2i2 *ago3*-GLKD condition where we found a drastic decrease in the production of *ZAM*-

derived piRNAs (**Fig. 3a**). In this condition, *ZAM* RNA accumulation in the ooplasm was

correlated with a strong increase of *ZAM* RNAs in total ovaries (RT-qPCR data) (**Fig. 3f**). These

results showed that when *ZAM*-derived piRNAs are absent or downregulated in the germline,

ZAM RNAs transit from the somatic follicle cells, where they are produced, to the oocyte.

Altogether, our data strongly suggest that *ZAM* piRNAs produced by the germline piRNA

pathway trigger post-transcriptional silencing of *ZAM* RNAs arriving from follicle cells. In line

with this conclusion no *ZAM* RNA was detected in stage 10 ooplasm and *ZAM* expression was

not affected when Piwi, which acts thought transcriptional gene silencing, was knocked down

in the RevI-H2i2 germline (**Fig. 3e, Supplementary Fig. S3a, b**).

We confirmed this finding using a line where no *ZAM*-derived piRNA is produced in the

germline (w[1118] genetic background) and in which we knocked down the piRNA pathway in

follicle cells by somatic knock-down (sKD) of *armi*, *vret*, *piwi* or *yb*. sKD of a piRNA pathway

actor, such as *vret* or *yb,* led to a strong decrease in *ZAM* piRNA production (**Fig. 4a,**

Supplementary Fig. S4a). The amount of antisense Piwi-bound piRNAs that map to soma-

specific TEs was strongly decreased (**Fig. 4b, Supplementary Fig. S4b**). RT-qPCR analysis

confirmed that *ZAM* and other soma-specific TEs, such as *Gypsy*, were derepressed (**Fig. 4c**).

Moreover, smRNA FISH results revealed that *ZAM* RNAs were not restricted to the posterior

follicle cells. Indeed, like in ovaries harboring GLKD of the piRNA pathway in RevI-H2i2

background, *ZAM* RNAs were present in oocytes. Actually, we detected *ZAM* RNAs in the

ooplasm in 90% of stage 10 follicles (**Fig. 4d, e, Supplementary Fig. S4c**). This result

confirmed that when no *ZAM* piRNA is produced in the germline, *ZAM* RNAs expressed in the

somatic follicle cells transit to the oocyte.

232 ***ZAM* RNAs transcribed in follicle cells transit to the oocyte and are deposited in early**

233 **embryos.**

234

235 Although *ZAM* should be transcribed only in follicle cells [28,29], we wanted to rule out the

236 possibility that *ZAM* RNAs originated from germinal nurse cells. *ZAM* RNAs strongly

237 accumulated at the posterior pole of oocytes, suggesting that these RNAs arrive from the

238 adjacent follicle cells (**Fig. 3e, 4d**). Moreover, we detected *ZAM* RNA staining only in the

239 cytoplasm of the oocyte at late stages of oogenesis (stage 10) upon *ago3*-GLKD in RevI-H2i2

240 ovaries. By contrast to TEs expressed in the germline (eg, Burdock), we never observed *ZAM*

241 RNA staining in the cytoplasm of other germ cells, the nurse cells, at any stage. (**Fig. 5a**). Our

242 experiments indicated that *ZAM* RNAs detected in oocytes originate from the somatic follicle

243 cells and not from nurse cells.

244 Throughout the animal kingdom, the first embryonic development stages are controlled by

245 transcripts and proteins deposited by the mother during oogenesis [30]. In Drosophila, most of the

246 maternal mRNAs are dumped from nurse cells into the oocyte during oogenesis. Although *ZAM*

247 transcripts originate from the somatic follicle cells, when we investigated *ZAM* RNA

248 transmission to the oocyte, we found such transcripts also in early embryos. Indeed, smRNA

249 FISH experiments revealed a strong accumulation of *ZAM* RNAs in early embryos (before the

250 zygotic transition, 0-2 hours after egg laying) from a RevI-H2i2 mother with GLKD of the

251 piRNA pathway, and also from a mother without germinal *ZAM* piRNAs with sKD of the

252 piRNA pathway in follicle cells (**Fig. 5b, Supplementary Fig. S5a**). Furthermore, *ZAM* RNAs

253 accumulated at the posterior pole of early embryos where future germ cells cellularize (**Fig. 5c,**

254 **Supplementary Fig. S5b**). We detected *ZAM* RNA in high quantity until stage 5 and few *ZAM*

255 RNA signal persisting until the cellularization of the blastoderm (~stage 8-9 of embryogenesis)

256 (**Fig. 5d**). In the progeny of the RevI-H2i2 line, we did not detect any *ZAM* RNA, although

ZAM RNAs were produced in follicle cells of RevI-H2i2 ovaries (**Fig 5b**). This confirmed that

ZAM invasion of the germline is not allowed in this line. These data revealed an efficient post-

transcriptional silencing of *ZAM* RNAs arriving from follicle cells into the oocyte by piRNAs

produced in the germline and dumped into the oocyte. This mechanism should limit the

transposition of the somatic *ZAM* retrotransposon into the germline genome.

***De novo ZAM* reactivation leads to subfertility.**

In addition to leading to the desilencing of many TEs, the loss of the piRNA pathway in the

germline or in the somatic follicle cells results in severe developmental defects during

oogenesis [31–34]. We sought to exclude the possibility that these defects could underlie the *ZAM*

invasion into the oocyte observed when the piRNA pathway is knocked down. To this end,

using CRISPR-Cas9, we *de novo* deleted the *ZAM* copy in the *flam* piRNA cluster in a line

carrying the X chromosome of the IsoA1 reference line and in which the piRNA pathway is

fully functional. As in this line, the *flam ZAM* copy is at the genomic position

X:21,778,810…21,783,994, we used two guides to create a deletion spanning over

X:21,777,135…21,784,062 (6926 pb). We named the resulting line *flamΔZAM*. Mapping of

unique piRNAs to the *flam* locus highlighted the complete loss of piRNA production at the

deleted region compared with the control IsoA1 line (**Fig. 6a**). Conversely, the global

production of unique piRNAs mapping upstream and downstream of *ZAM* was not affected.

We confirmed that the deletion induced a strong decrease of all piRNAs mapping to internal

regions of *ZAM*. However, antisense piRNAs targeting the *ZAM* LTR were still produced (**Fig.**

6b). In line with these results, PCR amplification and DNA sequencing of the CRISPR-Cas9

target locus showed that only the internal regions of *ZAM* were deleted from the *flam* cluster,

resulting in the retention of a solo-LTR at the initial *ZAM* insertion site (**Supplementary Fig.**

S6).

In addition, only the production of *ZAM*-derived antisense piRNAs was strongly altered,

whereas the production of antisense piRNAs mapping to other TEs was not affected (**Fig. 6c**).

This shows that *ZAM* deletion from *flam* impairs *ZAM*-derived piRNA production, but does not

affect the global piRNA production in ovaries. Thus, in the *flamΔZAM* line, the piRNA pathway

was functional and no *ZAM* piRNA targeting the internal *ZAM* sequences was produced. We

then analyzed *ZAM* RNA expression by smRNA FISH in *flamΔZAM* ovaries. Surprisingly, we

did not detect any *ZAM* expression (**Fig. 6d- panel 1**). We hypothesized that there was no

functional *ZAM* copy in the *flamΔZAM* line. To check this, we exchanged chromosome II or

III of the *flamΔZAM* line with chromosome II or III of the RevI-H2i2 line that each contains

four or three recently transposed potentially functional *ZAM* copies, respectively. smRNA FISH

revealed that *ZAM* RNAs were strongly expressed when these RevI-H2i2 *ZAM* copies were

added to the genome of the *flamΔZAM* line (**Fig. 6d – panels 2 and 3**). We also observed a

strong invasion of oocytes: approximately 70% of stage 10 follicles presented *ZAM* RNAs in

the ooplasm. We obtained similar results when we added the two *ZAM* copies located close to

the X chromosome telomeres of the RevI-H2i2 line to the *flamΔZAM* genome (**Fig. 6d – panel**

4). Overall, our findings show that in the absence of *ZAM*-derived piRNAs in the somatic

follicle cells and in the germline, transcripts from functional *ZAM* copies massively invade the

oocytes. Our results also demonstrated that the presence of a solo-LTR of *ZAM* (454 bp) in the

flam piRNA cluster of the *flamΔZAM* line was not sufficient to silence *ZAM* expression in

follicle cells (**Supplementary Fig. S6**).

Moreover, the *flamΔZAM* flies that contain active *ZAM* copies displayed atrophied ovaries and

reduced fertility compared with the control *flamΔZAM* flies that lack functional *ZAM* copies or

the RevI-H2i2 line (**Fig 6. e, f**). Specifically, 90% of *flamΔZAM* females with active *ZAM*

copies did not lay eggs. These findings indicate that the reactivation of one single

retrotransposon in a patch of follicle cells is sufficient to induce severe fertility defects. These

results suggest that when *ZAM* piRNAs are produced in the germline, they contribute to ensure

genome integrity and fertility by protecting oocytes against *ZAM* invasion from adjacent cells.

No initial small RNA response takes place after *ZAM* retrotransposon reactivation.

The reactivation of an endogenous retrovirus, such as *ZAM,* in a genetic context where the

piRNA pathway is functional provides a unique opportunity to directly examine the genomic

immune response to ERV reactivation and germline invasion. piRNAs and siRNAs are the two

main classes of small RNAs produced to control the expression of transposable elements and

counteract viral infection, respectively. Moreover, invading viral RNAs can be processed to

small RNAs to initiate a primary response [35,36]. To characterize the small RNA response upon

ERV reactivation, we sequenced small RNAs that are complexed with Argonaute proteins

(named here "regulatory" piRNAs) from *flamΔZAM* ovaries containing four or three functional

ZAM copies (introduced with the RevI-H2i2 chromosome II or III, respectively). Regardless of

the *flamΔZAM* line analyzed, *ZAM* piRNAs (23 to 29 nt) were not *de novo* produced upon *ZAM*

reactivation and despite germline invasion. Indeed, only few sense and antisense *ZAM* piRNAs

were produced in the ovaries of the *flamΔZAM* line that contained functional *ZAM* copies (**Fig.**

7a – Supplementary Fig. S7a). Moreover, this lack of piRNA production concerned only *ZAM*

and the level of regulatory piRNAs targeting other TEs was not reduced (**Fig. 7b-**

Supplementary Fig. S7b,).

We sequenced total small RNAs in *vret-* or *yb*-sKD ovaries where *ZAM* is also reactivated.

Surprisingly we observed sense *ZAM*-mapping small RNAs of various sizes, including 23-29

nt small RNAs that are not produced in a *white*-sKD control condition (**Supplementary Fig.**

S7c). However, the sense 23-29 nt *ZAM*-derived small RNAs detected in *yb-* and *vret*-sKD

ovaries did not display uridine bias at the 5' end, a feature of mature primary piRNAs

(**Supplementary Fig. S7d**). Furthermore, these sense *ZAM*-mapping small RNAs were not

immunoprecipitated with Piwi proteins. We detected almost no Piwi-bound sense or antisense

ZAM-mapping piRNAs in *vret-* or *yb*-sKD ovaries (**Supplementary Fig. S7e**). In addition, no

sense regulatory *ZAM*-piRNAs complexed with Argonaute proteins were detected (Fig. 4a).

These results suggest that the sense *ZAM*-mapping small RNAs are most likely degradation

products of excess *ZAM* sense transcripts and not piRNAs. Thus, in these different conditions,

we did not detect any primary piRNA response either in the somatic cells where *ZAM* is

expressed or in the oocyte that it invades.

As *ZAM* is an endogenous retrovirus, we also asked whether a siRNA response was triggered

upon *ZAM* reactivation. For this, we analyzed the production of 21 nt small RNAs complexed

with Argonaute proteins ("regulatory" siRNAs). Similar levels of *ZAM* 21 nt small RNAs were

produced in the *flamΔZAM* lines with functional *ZAM* copies (**Fig. 7c**). Globally, the production

of 21 nt small RNAs was comparable for all TEs upon *ZAM* reactivation (**Fig. 7d**). Their

production was only slightly increased in the *vret*-sKD condition, compared to the *white*-sKD

control line without *ZAM* expression (**Supplementary Fig. S7f**). Overall, our results suggest

that no or very few siRNAs that could protect ovaries from *ZAM* activity are produced upon

activation of this endogenous retrovirus in follicle cells.

Therefore, we propose that no innate small RNA response is triggered in the lines where *ZAM*

is reactivated, suggesting that *ZAM* will not be brought under control until the establishment of

specific adaptive immunity and immune memory.

Discussion

The discovery that piRNAs produced in germ cells not only restrict TE expression in the germline itself but also counteract the invasion of endogenous retroviruses from adjacent cells reveals a novel role of piRNAs produced locally as an effective defense mechanism at the tissue scale. When piRNAs against *ZAM* are produced in the germline, *ZAM* transcripts are limited to a patch of follicle cells and flies are fertile. Conversely, in the absence of *ZAM*-derived piRNAs in the germline, transcripts from functional *ZAM* copies massively invade the oocytes and are even transferred to the embryos. At the molecular level, our study revealed that piRNAs produced in nurse cells and dumped into the oocyte can target RNAs produced by follicle cells and delivered to the oocyte. It is tempting to speculate that the *ZAM* RNAs arriving from somatic cells are targeted by complementary antisense piRNAs and degraded through the ping-pong cycle. This would result in the production of *ZAM* sense piRNAs, thus participating in the amplification of the pool of piRNAs against *ZAM* and strengthening the defense mechanism against the invasion.

Transposition in the germ cell genome is crucial for TE propagation in a population because it allows the vertical transmission of new insertions. At the time of invasion (e.g. in the *flamΔZAM* line), no *ZAM*-derived piRNA was produced in somatic or germ cells. Therefore, this condition could be compared to what happens when a TE first enters a new species by horizontal transfer. In this case, according to several studies, an initial transposition burst occurs that leads to TE accumulation in the genome before the induction of an adaptive response by the host to control transposition [37–40]. We found at least 17 new *ZAM* insertions in the RevI-H2i2 genome, testifying that *ZAM* actively transposed. Surprisingly, we identified three *ZAM* insertions in the same germline piRNA cluster on the X chromosome. It has been proposed that two different mechanisms can explain the *de novo* piRNA production required to specifically silence a novel

invading TE. Indeed, piRNAs can be produced from a new TE insertion into a pre-existing

piRNA cluster, and also by euchromatic TE insertions converted into piRNA-producing loci

[14,41–43]. Interestingly, when active ZAM copies are added into a *flamΔZAM* genetic background,

these copies do not produce regulatory *ZAM*-derived piRNAs and are not converted into piRNA

producing regions. Actually, *ZAM* should be transcriptionally inactive in the germline because

ZAM expression is regulated by the somatic transcription factor Pointed2 [28]. Therefore, although

the contribution of individual copies needs to be investigated, our data strongly suggest that

germline *ZAM* piRNAs originate at least in part from piRNA clusters. In line with our findings,

computer simulations of the TE invasion dynamics suggested that several insertions in piRNA

clusters are likely to be required to stop the invasion in the germline [44]. Conversely, we found

that a single insertion in a somatic piRNA cluster, such as *flamenco*, is sufficient for TE

silencing in the somatic follicle cells, as previously suggested [24]. Indeed, we showed that in

follicle cells, the *flamenco* piRNA cluster acts alone as the principal regulator of transposon

activity. A precise deletion of the internal part of the endogenous retrovirus *ZAM*, inserted in

flamenco, leads to a complete derepression of functional *ZAM* copies. Surprisingly, reactivation

of this endogenous retrovirus in only a patch of somatic cells leads to subfertility. However, in

the germline the deletion of the three most highly expressed germline piRNA clusters (42AB,

38C, and 20A) affected neither fertility nor TE silencing [14]. This could be explained by the fact

that in the germline, but not in somatic cells, redundancy in piRNA production among different

piRNA clusters is observed, with probably multiple piRNA clusters involved in the silencing

of the same TE. Moreover, many TE copies have acquired mutations and are non-functional for

transcription or for transposition. In the *flamΔZAM* line, although no *ZAM* piRNA was

produced, we did not detect any *ZAM* expression. However, when we introduced functional

copies of *ZAM*, *ZAM* was expressed demonstrating that depending on the genetic background,

functional copies can be absent or present. In general, the TE content varies considerably among

populations [42,45]. Interestingly, we identified *ZAM* copies with a deletion that affects the *pol*

gene. The copy numbers of this *ZAM*-variant and of classical *ZAM* elements that have

transposed in the RevI-H2 line are similar (5 and 8 copies), suggesting that this *ZAM*-variant

can transpose very efficiently. Studies in different organisms have shown that TEs rapidly

diversify. For instance, the mariner DNA transposon is present in 68 different versions in grass

genomes [46]. Such rapid diversification certainly promotes speciation of new families of active

TEs.

One interesting question is how *ZAM* RNAs transit into the oocyte and at which stage during

Drosophila oogenesis *ZAM* transposes in germ cells. We noticed that *ZAM* RNAs, although

transcribed in follicle cells, were deposited and accumulated at the posterior pole of early

embryos, where future germ cells cellularize. *ZAM* might have evolved to preferentially

mobilize in the dividing primordial germ cells of the offspring instead of the developing oocyte,

which is not in a phase favorable to transposition because of its prolonged arrest in prophase I

associated with highly compacted chromatin [47].

Previous experiments demonstrated that *ZAM* RNAs accumulate in follicle cells when

vitellogenesis is defective, suggesting that *ZAM* transmission to the oocytes requires functional

vitellogenin trafficking [48]. Moreover, endogenous retroviruses, such as *ZAM* and *gypsy*, can

form pseudo-viral particles in follicle cells when they are expressed [29,48–50]. *ZAM* Gag end Env

proteins are expressed in RevI-H2i2 follicle cells. Retroviral envelope glycoproteins undergo

proteolytic processing by cellular serine endoproteases (furin and proprotein convertases) in

order to produce the two functional subunits, a glycosylated hydrophilic polypeptide (SU) and

a transmembrane domain (TM). This step is necessary to achieve protein competence to

promote membrane fusion and virus infectivity [51,52] . The *ZAM env* gene is similar to the

retroviral *env* gene, which is responsible for the infectious properties [25]. Indeed, *ZAM* Env is

composed of a SU, a TM and an Arg–X–Lys–Arg conserved domain, which is considered to

be a furin consensus proteolytic cleavage site [25,53,54]. A recent study demonstrated that

intracellular retrotransposition of the *gypsy* endogenous retrovirus can occur in the absence of

the Env protein, while intercellular transmission requires a functional Env [50]. Currently, whether

ZAM can transpose intracellularly, in follicle cells, and the exact mechanism(s) (Env protein-

dependent or independent) used for *ZAM* transmission to the oocyte are unknown.

Once a virus enters into the target cells, three models for retrovirus uncoating have been

proposed: the capsid (1) is disassembled soon after virus entry, (2) remains intact until it comes

into contact with the nuclear pore complex, or (3) is partially disassembled during the

cytoplasmic traffic to the nucleus (for review [55]). Capsid disassembly of endogenous

retroviruses has not been investigated yet. In our study, we observed the loss of *ZAM* RNA in

the oocyte when germline *ZAM*-derived piRNAs are produced. We could hypothesize that *ZAM*

capsid is dissociated soon after entry in the oocyte and thus piRNAs can easily target *ZAM*

RNAs. Alternatively, *ZAM* RNAs are still encapsidated in the oocyte but the capsid does not

protect the *ZAM* RNA from piRNAs and their associated endonuclease. Previous studies on

virus-like particles of the yeast retrotransposon Ty suggest that these particles form open

structures that leave the RNA at the interior of the capsid accessible to RNase [56,57].

It has been proposed that in different species, an "innate" response at the piRNA or siRNA level

could be initiated in the gonads upon retrovirus invasion (reviewed in [58], [36,59–63]). We found that

no clear innate small RNA response is triggered upon *ZAM* reactivation in Drosophila ovaries.

Total small-RNA sequencing after *ZAM* reactivation revealed the presence of sense *ZAM*-

mapping small RNAs, but they lack characteristics of siRNAs or piRNAs. We hypothesize that

these sense small RNAs are degradation products generated when there is an excessive amount

of *ZAM* RNAs. Altogether, these results show that no initial small RNA response is mounted

upon *ZAM* reactivation, leaving follicle cells, where *ZAM* is expressed, and also oocytes

unprotected against *ZAM* transposition. Our findings suggest that the piRNA response is a

robust response that, in the case of *ZAM*, must be established either in the follicle cells or in the

germ cells to efficiently control the invading TE. However, the time required for developing a

robust piRNA response to sustainably control TE transposition remains unknown. Lastly, it is

thought that the piRNA pathway has no antiviral role in Drosophila [64]. However, it would be

interesting to determine whether this is true also if parts of the viral genome are integrated into

a germline piRNA cluster during infection.

Online Methods

Fly stocks, transgenic lines, and crosses

Flies were maintained at 25°C under a light/dark cycle and 60% humidity. Between 3 and 6

days after hatching, flies were used for experiments. The RNAi lines against components of the

piRNA pathway used in this study are listed in **Supplementary Table S2**. Isogenic lines from

the RevI-H2 stock were generated by crossing a single female with a single male for five

generations. Germline and somatic knock-down have been performed using the nanos-gal4 and

the traffic-jam-gal4 driver lines respectively. The Drosophila line carrying the *ZAM* sensor is

described in [23]. The sensor expresses the GFP reporter gene under the control of an inducible

Upstream Activation Sequence promoter (UASp) and harbors a *ZAM* fragment in its 3′UTR

(pGFP-*ZAM*).

Single-molecule RNA fluorescence *in situ* hybridization (smRNA FISH) in ovaries and

embryos.

ZAM smRNA FISH was performed using 48 probes that target *ZAM* transcripts in a region that

is absent in the *ZAM* inserted in the *flamenco* piRNA cluster to detect only transcripts of active

ZAM copies (sequences in **Supplementary Table S3**). Ovaries from 3 to 6-day-old flies were

dissected in Schneider's Drosophila Medium and fixed in Fixing Buffer (4% formaldehyde,

0.3% Triton X-100, 1x PBS) for 20 min at room temperature, rinsed three times in 0.3% Triton

X-100, once in PBS, and permeabilized in 70% ethanol at 4°C overnight. Permeabilized ovaries

were rehydrated in smRNA FISH wash buffer (10% formamide in 2x SSC) for 10 min. Ovaries

were resuspended in $50\mu L$ hybridization buffer (10% dextran sulfate, 10% formamide in 2x

SSC) supplemented with $1\mu L$ of smRNA FISH probes. Hybridization was performed with

rotation at 37°C overnight. Ovaries were then washed twice with smRNA FISH wash buffer at

37°C for 30 min and twice with 2xSSC solution. Then, DNA was stained with DAPI (1/500

dilution in 2x SSC) at room temperature for 20 min. Ovaries were mounted in 30 μL

Vectashield mounting medium and imaged on a Zeiss LSM-980 or Zeiss LSM-800 confocal

microscope. The resulting images were processed using FIJI/ImageJ. The RNA signal

specificity was confirmed by adding 1 mg/mL of RNase A in 2x SSC for 2h before the

hybridization step.

For embryo staining, flies were caged and fed yeast paste. Embryos (0-2h) were collected,

dechorionated in 50% bleach for 4 min and rinsed in water. Eggs were fixed in 4%

paraformaldehyde/heptane for 20 min, devitellinized by vigorous shaking in 100% methanol

and stored in methanol at -20°C. Embryos were rehydrated with 1:1 methanol in 1xPBS, 0.1%

Tween-20 for 5 min and twice in 1xPBS, 0.1% Tween-20. Embryos were resuspended in

smRNA FISH wash buffer (10% formamide in 2x SSC) for 10 min and then processed for

smRNA FISH as described for ovaries. Immunostaining combined with smRNA FISH was

performed by adding primary antibodies to the smRNA FISH probes in the hybridization buffer

and incubating at 37°C overnight. Embryos were washed twice with smRNA FISH wash buffer

and incubated with the secondary antibody (1/1000 diluted in 2x SSC) for 90 min. After two

washes in 2x SSC, embryos were mounted in 30 μL Vectashield mounting medium.

Immunofluorescence analysis of ovaries and embryos.

Ovaries from 3-5-day-old flies were dissected in supplemented Schneider's medium, ovarioles were separated, and the muscle sheath was removed before fixation to obtain undistorted follicles. Then, ovaries were fixed in 4% formaldehyde/1x PBS/2% Tween-20 for 15 min, rinsed three times with 1x PBS/2% Tween-20, and incubated 2 hours in PBT (1x PBS, 0.1% Triton X-100, 1% BSA) for blocking. Ovaries were incubated with primary antibodies in PBT at 4°C overnight (antibodies are described in **Supplementary Table S4)**. After three washes in PBT, ovaries were incubated with the corresponding secondary antibodies (1/1000 diluted in PBT), coupled to Alexa-488 or Cy3 for 90 min. After two washes in 1x PBS, DNA was stained with DAPI (1/500 dilution in 1x PBS) for 20 min. Ovaries were mounted in 30 μL Vectashield mounting medium and imaged on a Zeiss LSM-980 or Zeiss LSM-800 confocal microscope. The resulting images were processed using FIJI/ImageJ.

RT-qPCR analysis of transposon expression

10-20 pairs of dissected ovaries were homogenized in TRIzol reagent (Ambion). Following DNase I treatment, cDNA was prepared from 1 μg RNA by random priming of total RNA using Superscript IV Reverse Transcriptase (ThermoFisher Scientific). Quantitative PCR was performed with Roche FastStart SYBR Green Master and the Lightcycler® 480 Instrument. cDNA was used as template for RT-qPCR quantification of transposon mRNA levels (primer sequences in **Supplementary Table S5**). All experiments were done with three biological replicates and with technical triplicates. Steady-state RNA levels were calculated from the threshold cycle for amplification with the $2^{-\Delta\Delta CT}$ method; *rp49* was used for normalization.

Immunoprecipitation of Piwi-ribonucleoprotein complexes

For each genotype, 50 pairs of ovaries from 3-5-day-old flies were dissected, lysed in 1 ml lysis

buffer (20 mM HEPES-NaOH at pH 7.0, 150 mM NaCl, 2.5 mM $MgCl_2$, 250 mM sucrose,

0.05% NP40, 0.5% Triton X-100, 1x Roche-Complete). Samples were cleared by centrifugation

at 10000 × g at 4°C for 10 min. Extracts were incubated with rotation with rabbit polyclonal

anti-Piwi antibodies (4 μg per sample) at 4°C for 4h followed by overnight incubation with

Dynabeads Protein A (50 μl, Invitrogen, 10002D) at 4°C with rotation. Before incubation,

beads were equilibrated in NT2 buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1 mM $MgCl_2$,

0.05% NP40). Beads were washed twice with ice-cold NT2 and twice with NT2 in which NaCl

concentration was adjusted to 300 mM. Nucleic acids that co-immunoprecipitated with Piwi

were isolated by treating beads with 0.7 mg/ml proteinase K in 0.3ml proteinase K buffer (0.5%

SDS, 10mM Tris-HCl pH 7.4, 50mM NaCl, 5mM EDTA), followed by phenol/chloroform

extraction (phenol at neutral pH) and ethanol precipitation.

DNA Isolation, Oxford Nanopore Technology sequencing and genome analysis

DNA was extracted from 200 RevI-H2i2 females using the Qiagen Blood & Cell Culture DNA

Midi kit. The genomic DNA quality and quantity were evaluated using a Femto Pulse (Agilent)

and a Qbit 3.0 (Invitrogen) respectively. Oxford Nanopore Technology (ONT) sequencing was

performed by I2BC (Gif-sur-Yvette, France) using five micrograms of genomic DNA. Adapter-

trimmed ONT reads were analyzed using NCBI Blastn

(https://blast.ncbi.nlm.nih.gov/Blast.cgi) or command-line Blastn (Blastn 2.8.1). The RevI-

H2i2 ONT reads containing *ZAM* insertions were detected by Blastn with the reference *ZAM*

[65], Repbase *ZAM*_I and *ZAM*_LTR sequences (https://www.girinst.org/repbase/, [26]). The *ZAM*

containing reads were then recovered with bedtools getfasta

(https://bedtools.readthedocs.io/en/latest/content/tools/getfasta.html, [66]). The *ZAM* insertion

sites where determined using Blastn of the *ZAM* containing reads to the *D. melanogaster*

Release 6 genome (http://flybase.org). The absence of empty sites, without a given *ZAM*

insertion, was verified by Blastn of the respective empty insertion sites, recovered from the *D.*

melanogaster Release 6 genome, to all ONT reads.

Small RNA sequencing

Regulatory small RNA extraction was performed as described in [67]. Briefly, 50 pairs of ovaries

from 3-5-day-old flies were lysed and Argonaute-small RNA complexes were isolated using

TraPR ion exchange spin columns. Small RNAs were subsequently purified using TRIzol

(Ambion). Total RNA also was extracted for some samples (80-100 ovaries for each sample)

using the classical TRIzol extraction, followed by 2S depletion and size selection before

sequencing. Sequencing was performed by Fasteris SA (Geneva, Switzerland) on an Illumina

NextSeq550 instrument (13-15 million reads per sample).

Illumina small RNA sequencing reads were loaded in the small RNA pipeline sRNAPipe [68] for

mapping to the various genomic sequence categories of the *D. melanogaster* genome (Release

6). For the analysis, 23-29nt genome mappers were selected as piRNAs, and 21nt genome

mappers were selected as siRNAs. "Unique" piRNAs were defined as 23-29 reads that mapped

uniquely across the reference genome. piRNAs or unique piRNAs were mapped to TEs (0-3

mismatches) or to piRNA clusters, as defined in [10] (no mismatch allowed). The window size

was of 91nt for *ZAM* and 140nt for the *flamenco* region at X:21,769,891…21,789,891 to

establish the piRNA density profile. For comparison between samples, all read counts were

normalized by the number of reads used as input (genome-mapped piRNAs, unique genome-

mapped piRNAs, or genome-mapped siRNAs). The ping-pong signature was assessed by

calculating the percentage of sense piRNAs with 10nt 5'-overlap with antisense piRNAs, based

on piRNA mapping to *ZAM* (0 to 3 mismatches). The percentages of 1 to 23nt-long overlaps

were determined, and the percentage of 10nt overlaps was defined as the ping-pong signature.

The Z-score for 10nt overlaps was determined using the percentages of 1 to 23nt-long overlaps

as a background and was considered significant when it was >1.96. Scatter plots were done

with RStudio with antisense piRNAs or siRNAs mapping to individual internal sequence of

TEs (0-3 mismatches for piRNAs and 0-1 mismatches for siRNAs). All small RNA sequencing

done for this study is available in **Supplementary Table S6.**

Sterility tests

For the sterility tests, RevI-H2i2 homozygous females were compared to w[1118] females, and

homozygous *flamΔZAM* females that carry two functional *ZAM* copies from the RevI-H2i2 X

chromosome were compared to *flamΔZAM* females without functional *ZAM* copies in their

genome. Thirty 3-5-day-old virgin females of each genotype were individually mated with three

w[1118] males, and eggs were collected at 25°C for 24 hours. The number of eggs laid by each

female was determined. Then, eggs were kept at 25°C for another 24 hours before determining

the egg hatching rate. The experiment was done twice for a total of approximatively 60

replicates for each condition.

Generation of the *flamΔZAM* line using the CRISPR-Cas9 technology

The *ZAM* copy in the *flamenco* piRNA cluster was deleted using a pCFD6 plasmid that

expresses two single guide RNAs (sgRNAs) under the Gal4/UAS system. The sgRNAs were

designed to target genome-unique sequences located upstream and downstream of the *flamenco*

ZAM copy: TTGTAGCGCTCTTCTTCTCT (sgRNA *flamΔZAM_1*) and

AGCGCAACCACGTACAGCGA (sgRNA *flamΔZAM_2*). The pCFD6 plasmid harboring

these sgRNAs was injected by the Bestgene company into embryos from the BL#9736 stock

(Bloomington stock center) for integration of the plasmid into the genomic site 53B2. The

obtained flies were crossed with nanos-gal4; UAS-Cas9 flies (BL#54593, Bloomington stock center). Prior to crossing, the X chromosomes of both Drosophila lines were replaced by the X chromosome of the IsoA1 Drosophila line. After establishment of 100 lines, homozygote females were screened by PCR using primers specific for the flamenco *ZAM* copy. Lines without any amplification of these *flamenco ZAM* sequences were selected for PCR amplification of the target locus with primers that framed the *flamenco ZAM* copy and the target sequences of the sgRNAs. The obtained amplicons were sequenced.

ACKNOWLEDGEMENTS

We thank the *Bloomington Stock Center* for providing stocks. We thank Julius Brennecke for providing antibodies. We thank flybase.org for providing databases. We thank Séverine Chambeyron and Kirsten-André Senti for helpful discussions and the members of the Brasset laboratory for comments on the manuscript. We thank Yoan Renaud and Pierre Pouchin for helpful advices and development of the GReD bioinfomatic platform. We thank Gentyane Facility for the Quality Control of gDNA.

FUNDING

This work was supported by grants from the Agence Nationale pour la Recherche (ANR-CHApiTRE (AAPG2020 – CE12), ANR-BiopiC (AAPG2021 – PRC – CE12), and ANR-EpiTET projects (17-CE12-0030-03)), the Association de Recherche Cancer (PJA20171206129). M.Y. was supported by the Ministère de l'Enseignement Supérieur et de la Recherche (MESR) and the Fondation pour la Recherche Medical (FRM) (FDT202106012950). This research is supported by the French government Emergence-IDEX-ISITE initiative 16-IDEX-0001 (CAP20-25). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

COMPETING INTERESTS

The authors declare that no competing interest exists.

REFERENCES

1.   Byrd, D. T. & Kimble, J. Scratching the niche that controls Caenorhabditis elegans

germline stem cells. *Semin. Cell Dev. Biol.* **20**, 1107–1113 (2009).

2.   Kidder, G. M. & Vanderhyden, B. C. Bidirectional communication between oocytes

and follicle cells: ensuring oocyte developmental competence. *Can. J. Physiol.*

Pharmacol. **88**, 399–413 (2010).

3.   Saunders, P. T. K. Germ cell-somatic cell interactions during spermatogenesis. *Reprod.*

(Cambridge, England). Suppl. **61**, 91–101 (2003).

4.   Jemc, J. C. Somatic gonadal cells: the supporting cast for the germline. *Genesis* **49**,

753–775 (2011).

5.   Hall, D. H. *et al.* Ultrastructural features of the adult hermaphrodite gonad of

Caenorhabditis elegans: relations between the germ line and soma. *Dev. Biol.* **212**,

101–123 (1999).

6.   McClintock, B. The origin and behavior of mutable loci in maize. *Proc. Natl. Acad.*

Sci. U. S. A. **36**, 344–355 (1950).

7.   Aravin, A. *et al.* A novel class of small RNAs bind to MILI protein in mouse testes.

442, 203–207 (2006).

8.   Grivna, S. T., Beyret, E., Wang, Z. & Lin, H. A novel class of small RNAs in mouse

spermatogenic cells. *Genes Dev.* **20**, 1709–1714 (2006).

9.   Lau, N. C. *et al.* Characterization of the piRNA complex from rat testes. *Science (80-.*

). **313**, 363–367 (2006).

10.   Brennecke, J. *et al.* Discrete Small RNA-Generating Loci as Master Regulators of

Transposon Activity in Drosophila. *Cell* **128**, 1089–1103 (2007).

11.   Aravin, A. a, Hannon, G. J. & Brennecke, J. The Piwi-piRNA Pathway Provides an

Adaptive Defense in the Transposon Arms Race. *Science (80-.).* **318**, 761–764 (2007).

12.   George, P. *et al.* Increased production of piRNAs from euchromatic clusters and genes

in Anopheles gambiae compared with Drosophila melanogaster. *Epigenetics and*

Chromatin **8**, (2015).

13.   Malone, C. D. *et al.* Specialized piRNA Pathways Act in Germline and Somatic

Tissues of the Drosophila Ovary. *Cell* **137**, 522–535 (2009).

14.   Gebert, D. *et al.* Large Drosophila germline piRNA clusters are evolutionarily labile

and dispensable for transposon regulation. *Mol. Cell* 1–14 (2021)

doi:10.1016/j.molcel.2021.07.011.

15.   Handler, D. *et al.* The genetic makeup of the drosophila piRNA pathway. *Mol. Cell* **50**,

762–777 (2013).

16.   Olivieri, D., Sykora, M. M., Sachidanandam, R., Mechtler, K. & Brennecke, J. An in

vivo RNAi assay identifies major genetic and cellular requirements for primary piRNA

biogenesis in Drosophila. *EMBO J.* **29**, 3301–3317 (2010).

17.   Desset, S., Meignin, C., Dastugue, B. & Vaury, C. COM, a heterochromatic locus

governing the control of independent endogenous retroviruses from Drosophila

melanogaster. *Genetics* **164**, 501–509 (2003).

18.   Pélisson, A. *et al.* Gypsy transposition correlates with the production of a retroviral

envelope-like protein under the tissue-specific control of the Drosophila flamenco

gene. *EMBO J.* **13**, 4401–4411 (1994).

19.   Bucheton, A. The relationship between the flamenco gene and gypsy in Drosophila :

how to tame a retrovirus. **11**, 349–353 (1995).

20.   Robert, V., Prud'homme, N., Kim, A., Bucheton, A. & Pélisson, A. Characterization of

the flamenco region of the Drosophila melanogaster genome. *Genetics* **158**, 701–713

(2001).

21.   Mével-Ninio, M., Pelisson, A., Kinder, J., Campos, A. R. & Bucheton, A. The

flamenco Locus Controls the gypsy and ZAM Retroviruses and Is Required for

Drosophila Oogenesis ´. **1624**, 1615–1624 (2007).

22.   Lécher, P., Bucheton, A. & Pélisson, A. Expression of the Drosophila retrovirus gypsy

as ultrastructurally detectable particles in the ovaries of flies carrying a permissive

flamenco allele. *J. Gen. Virol.* **78**, 2379–2388 (1997).

23.   Duc, C. *et al.* Trapping a somatic endogenous retrovirus into a germline piRNA cluster

immunizes the germline against further invasion. *Genome Biol.* **20**, 1–14 (2019).

24.   Zanni, V. *et al.* Distribution, evolution, and diversity of retrotransposons at the

flamenco locus reflect the regulatory properties of piRNA clusters. *Proc. Natl. Acad.*

Sci. **110**, 19842–19847 (2013).

25.   Leblanc, P., Desset, S., Dastugue, B. & Vaury, C. Invertebrate retroviruses: ZAM a

new candidate in D. melanogaster. *EMBO J.* **16**, 7521–7531 (1997).

26.   Jurka, J. *et al.* Repbase Update, a database of eukaryotic repetitive elements.

Cytogenet. Genome Res. **110**, 462–467 (2005).

27.   Desset, S. *et al.* Mobilization of two retroelements, ZAM and Idefix, in a novel

unstable line of Drosophila melanogaster. *Mol. Biol. Evol.* **16**, 54–66 (1999).

28.   Meignin, C., Dastugue, B. & Vaury, C. Intercellular communication between germ line

and somatic line is utilized to control the transcription of ZAM, an endogenous

retrovirus from Drosophila melanogaster. *Nucleic Acids Res.* **32**, 3799–3806 (2004).

29.   Leblanc, P. *et al.* Life Cycle of an Endogenous Retrovirus, ZAM, in Drosophila

melanogaster. *J. Virol.* **74**, 10658–10669 (2000).

30.   Atallah, J. & Lott, S. E. Evolution of maternal and zygotic mRNA complements in the

early Drosophila embryo. *PLoS Genet.* **14**, e1007838 (2018).

31.   Aravin, A. A. & Hannon, G. J. Small RNA silencing pathways in germ and stem cells.

Cold Spring Harb. Symp. Quant. Biol. **73**, 283–290 (2008).

32.   Khurana, J. S. & Theurkauf, W. piRNAs, transposon silencing, and Drosophila

germline development. *J. Cell Biol*. **191**, 905–913 (2010).

33.   Saito, K. & Siomi, M. C. Small RNA-mediated quiescence of transposable elements in

animals. *Dev. Cell* **19**, 687–697 (2010).

34.   Klattenhoff, C. *et al*. Drosophila rasiRNA pathway mutations disrupt embryonic axis

specification through activation of an ATR/Chk2 DNA damage response. *Dev. Cell*

12, 45–55 (2007).

35.   Parameswaran, P. *et al*. Six RNA viruses and forty-one hosts: viral small RNAs and

modulation of small RNA repertoires in vertebrate and invertebrate systems. *PLoS*

Pathog. **6**, e1000764 (2010).

36.   Yu, T. *et al*. The PiRNA Response to Retroviral Invasion of the Koala Genome. *SSRN*

Electron. J. 1–12 (2019) doi:10.2139/ssrn.3391942.

37.   Le Rouzic, A. & Capy, P. The first steps of transposable elements invasion: Parasitic

strategy vs. genetic drift. *Genetics* **169**, 1033–1043 (2005).

38.   Kofler, R., Senti, K. A., Nolte, V., Tobler, R. & Schlötterer, C. Molecular dissection of

a natural transposable element invasion. *Genome Res*. **28**, 824–835 (2018).

39.   Robillard, É., Le Rouzic, A., Zhang, Z., Capy, P. & Hua-Van, A. Experimental

evolution reveals hyperparasitic interactions among transposable elements. *Proc. Natl.*

Acad. Sci. U. S. A. **113**, 14763–14768 (2016).

40.   Schaack, S., Gilbert, C. & Feschotte, C. Promiscuous DNA: Horizontal transfer of

transposable elements and why it matters for eukaryotic evolution. *Trends Ecol. Evol*.

25, 537–546 (2010).

41.   Mohn, F., Sienski, G., Handler, D. & Brennecke, J. The Rhino-Deadlock-Cutoff

complex licenses noncanonical transcription of dual-strand piRNA clusters in

Drosophila. *Cell* **157**, 1364–1379 (2014).

42.   Luo, S. *et al*. The evolutionary arms race between transposable elements and piRNAs

738   in Drosophila melanogaster. *BMC Evol. Biol.* **20**, 1–18 (2020).

739   43.   Shpiz, S., Ryazansky, S., Olovnikov, I., Abramov, Y. & Kalmykova, A. Euchromatic

740   Transposon Insertions Trigger Production of Novel Pi- and Endo-siRNAs at the Target

741   Sites in the Drosophila Germline. *PLoS Genet.* **10**, (2014).

742   44.   Kofler, R. Dynamics of transposable element invasions with piRNA clusters. *Mol. Biol.*

743   *Evol.* **36**, 1457–1472 (2019).

744   45.   Lerat, E. *et al*. Population-specific dynamics and selection patterns of transposable

745   element insertions in European natural populations. *Mol. Ecol.* **28**, 1506–1522 (2019).

746   46.   Feschotte, C. & Wessler, S. R. Mariner-like transposases are widespread and diverse in

747   flowering plants. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 280–285 (2002).

748   47.   Navarro-Costa, P. *et al*. Early programming of the oocyte epigenome temporally

749   controls late prophase I transcription and chromatin remodelling. *Nat. Commun.* **7**,

750   12331 (2016).

751   48.   Brasset, E. *et al*. Viral particles of the endogenous retrovirus ZAM from Drosophila

752   melanogaster use a pre-existing endosome/exosome pathway for transfer to the oocyte.

753   *Retrovirology* **3**, 1–9 (2006).

754   49.   Chalvet, F. *et al*. Proviral amplification of the Gypsy endogenous retrovirus of

755   Drosophila melanogaster involves env-independent invasion of the female germline.

756   *EMBO J.* **18**, 2659–2669 (1999).

757   50.   Keegan, R. M., Talbot, L. R., Chang, Y. H., Metzger, M. J. & Dubnau, J. Intercellular

758   viral spread and intracellular transposition of Drosophila gypsy. *PLoS Genet.* **17**, 1–22

759   (2021).

760   51.   Seidah, N. G. & Prat, A. Precursor convertases in the secretory pathway, cytosol and

761   extracellular milieu. *Essays Biochem.* **38**, 79–94 (2002).

52.   Bergeron, F., Leduc, R. & Day, R. Subtilase-like pro-protein convertases: from

molecular specificity to therapeutic applications. *J. Mol. Endocrinol.* **24**, 1–22 (2000).

53.   Klenk, H. D. & Garten, W. Host cell proteases controlling virus pathogenicity. *Trends*

Microbiol. **2**, 39–43 (1994).

54.   Hosaka, M. *et al.* Arg-X-Lys/Arg-Arg motif as a signal for precursor cleavage

catalyzed by furin within the constitutive secretory pathway. *J. Biol. Chem.* **266**,

12127–12130 (1991).

55.   Yamashita, M. & Engelman, A. N. Capsid-Dependent Host Factors in HIV-1 Infection.

Trends Microbiol. **25**, 741–755 (2017).

56.   Palmer, K. J. *et al.* Cryo-electron microscopy structure of yeast Ty retrotransposon

virus-like particles. *J. Virol.* **71**, 6863–6868 (1997).

57.   Burns, N. R. *et al.* Symmetry, flexibility and permeability in the structure of yeast

retrotransposon virus-like particles. *EMBO J.* **11**, 1155–1164 (1992).

58.   Kolliopoulou, A. *et al.* PIWI pathway against viruses in insects. *Wiley Interdiscip. Rev.*

RNA **10**, e1555 (2019).

59.   Rozhkov, N. V. *et al.* Small RNA-based silencing strategies for transposons in the

process of invading Drosophila species. *Rna* **16**, 1634–1645 (2010).

60.   Rozhkov, N. V. *et al.* Evolution and dynamics of small RNA response to a

retroelement invasion in drosophila. *Mol. Biol. Evol.* **30**, 397–408 (2013).

61.   Kawamura, Y. *et al.* Drosophila endogenous small RNAs bind to Argonaute 2 in

somatic cells. *Nature* **453**, 793–797 (2008).

62.   Rehwinkel, J. *et al.* Genome-Wide Analysis of mRNAs Regulated by Drosha and

Argonaute Proteins in Drosophila melanogaster . *Mol. Cell. Biol.* **26**, 2965–2975

(2006).

63.   Katsuma, S. *et al.* Transcriptome profiling reveals infection strategy of an insect

maculavirus. *DNA Res. an Int. J. rapid Publ. reports genes genomes* **25**, 277–286

(2018).

64. Petit, M. *et al.* piRNA pathway is not required for antiviral defense in Drosophila

melanogaster . *Proc. Natl. Acad. Sci.* **113**, E4218–E4227 (2016).

65. Leblanc, P., Desset, S., Dastugue, B. & Vaury, C. Invertebrate retroviruses: ZAM a

new candidate in D. melanogaster. *EMBO J.* **16**, 7521–7531 (1997).

66. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**, 841–842 (2010).

67. Grentzinger, T. *et al*. A universal method for the rapid isolation of all known classes of

functional silencing small RNAs. *Nucleic Acids Res.* **48**, (2020).

68. Pogorelcnik, R., Vaury, C., Pouchin, P., Jensen, S. & Brasset, E. SRNAPipe: A

Galaxy-based pipeline for bioinformatic in-depth exploration of small RNAseq data.

Mob. DNA **9**, 4–9 (2018).

Figure 1: *ZAM* is expressed in follicle cells in the RevI-H2i2 isogenic line.
(a) Color-inverted projection of confocal images of ovarioles, germarium, and stage 10 follicles showing *ZAM* expression by smRNA FISH (in black) in ovaries from the RevI-H2i2 line (Z-projection of 20 stacks). (b) Fold-change in the steady-state *ZAM*, *Gypsy* and *Burdock* RNA levels for RevI-H2i2 ovaries compared with w[1118] ovaries (control), quantified by RT-qPCR (primer sequences in Supplementary Table S5). At least three biological replicates and three technical replicates were used. **p-value <0.01; ns, not-significant (p-value >0.05) (Mann-Whitney test). Error bars indicate the standard deviation (SD). (c) Confocal sections of stage 10 RevI-H2i2 egg chambers showing *ZAM*-encoded Gag (green) and Env (red) proteins. DNA was stained with DAPI (blue). (d) Morphology of control and RevI-H2i2 adult ovaries. (e) Box plot displaying the number of eggs laid per fly per day by control and RevI-H2i2 females. Each dot represents an individual female. The midline indicates the median value, and the box shows the first and third quartile. Error bars indicate the SD. ns, not-significant (p-value >0.05) (Mann-Whitney test).

828
Figure 2: *ZAM* elements in the RevI-H2i2 genome.
(**a**) Each triangle represents a *ZAM* insertion. Blue: full-length *ZAM* elements; orange: *ZAM*-
variants, in which positions 5500-6126 of the *ZAM* internal sequences (*ZAM*_I in RepBase) are
deleted; violet: *ZAM* full-length with deleted 5'-UTR; green: *ZAM* with internal deletions other
than the *ZAM*-variant; gray: *ZAM* copy also present in the reference Release 6 genome
(http://flybase.org). Four of the X chromosome *ZAM* copies are in X pericentromeric
heterochromatin (in black), all the others are in euchromatic regions. No *ZAM* insertion was
found in chromosome 4 (1.35 Mb, not shown). For coordinates and details see Supplementary
Table S1. (**b**) Density plot of *ZAM*-mapping regulatory piRNAs along the *ZAM* sequence in
RevI-H2i2 ovaries (up to 3 mismatches). (**c**) Histogram showing the percentage of 5'-overlaps
between sense and antisense *ZAM*-derived piRNAs (23–29 nt) in RevI-H2i2 ovaries. The
proportion of 10 nt 5'-overlap pairs is in red and the corresponding Z-score is indicated.
(**d**) Confocal images of ovarioles with GFP (green) and DNA (blue) staining. Ovarioles of the
progeny of a cross between RevI-H2i2 females and males carrying the pGFP-*ZAM* sensor
transgene driven by actin-Gal4.
844
845

846
Figure 3: Germline *ZAM* piRNAs produced in RevI-H2i2 germ cells counteract *ZAM*
invasion.
(**a**)Antisense piRNAs mapping to TE internal sequences (0-3 mismatches) in the RevI-H2i2
line upon *white-* (control), *ago3-*, *zuc-*, or *piwi*-GLKD. Normalized per million of genome-
mapping piRNAs. (**b**) Box plot displaying the Z-score distribution of 10 nt 5'-overlaps for TE-
mapping piRNAs in the *white*-GLKD and *ago3*-GLKD RevI-H2i2 lines. Each dot represents a
TE with a Z-score >1.96 (dotted line) in the control condition (*white*-GLKD). *ZAM* is
highlighted in red. The midline indicates the median value, and the box shows the first and third
quartile. Error bars indicate the SD. ***p-value* <0.001 (Mann-Whitney test). (**c**) Histogram
showing the percentage of 5'-overlaps between sense and antisense *ZAM*-mapping piRNAs (up
to 3 mismatches) in *white-*, *ago3-*, *zuc-* or *piwi*-GLKD RevI-H2i2 ovaries. The percentage of
10 nt overlapping pairs is in red. (**d**) Color-inverted confocal images of stage 10 egg chambers
showing *ZAM* smRNA FISH signal in ovaries of the indicated genotypes. (**e**) Bar plot showing
the percentage of stage 10 follicles with *ZAM* RNA detected in the ooplasm by smRNA FISH
of RevI-H2i2 ovaries with the indicated GLKD. This experiment was done three times and
approximately 50 follicles per conditions were observed for each experiment. Error bars
indicate the SD. (**f**) Bar plot showing the steady-state *ZAM* RNA levels in total RNA from
ovaries with the indicated genotypes (normalized to *rp49*). At least three
biological replicates and two technical replicates were used. ***p-value* <0.01 (Mann-Whitney
tst); error bars indicate the SD.

**Figure 4: *ZAM* invades the oocyte when no *ZAM*-derived piRNA is produced in the germ
cells.**

(**a**) Density plot along the *ZAM* sequence of *ZAM*-mapping regulatory piRNAs produced in
white- and *vret*-sKD ovaries (up to 3 mismatches). (**b**) Scatter plot showing the log2 of the
normalized counts [log2(RPM)] of antisense Piwi-bound piRNAs mapping to individual
internal TEs sequences in control ovaries (*white*-sKD) and *vret*-sKD ovaries. Antisense piRNA
counts, mapped allowing up to 3 mismatches, were normalized per million of genome-mapping
piRNAs (RPM). *TEs in red have a *vret*-sKD/*white*-sKD ratio <0.3. (**c**) Fold-change in the
steady-state *ZAM*, *Gypsy* and *Burdock* RNA level for the indicated sKD ovaries, compared with
white-sKD ovaries, measured by RT-qPCR (primer sequences in Supplementary Table S5). At
least three biological replicates and two technical replicates were used. **p-value* <0.01 (Mann-
Whitney test); error bars indicate the SD. (**d**) Color-inverted confocal images of ovarioles or
stage 10 egg chambers showing *ZAM* smRNA FISH signal in *white-* and *vret*-sKD ovaries.
ZAM RNA is observed in follicle cells and invaded the ooplasm in *vret*-sKD ovaries (zoomed
inset). (**e**) Bar plot showing the percentage of stage 10 follicles with *ZAM* RNA invasion of the
ooplasm, assessed by *ZAM* smRNA FISH, of *white-* (control) and *vret*-sKD ovaries. This
experiment was done twice and ~50 follicles per condition were observed for each experiment.
Error bars indicate the SD

926
Figure 5: *ZAM* RNA transcribed in the follicle cells transits to the oocyte and is deposited in early embryos.

(a) Confocal images of ovarioles (upper panels) and of an isolate stage 8 egg chamber (lower panels) in the *ago3*-GLKD RevI-H2i2 line. *Burdock* (green) and *ZAM* mRNA (red) were detected by smRNA FISH. DNA was stained with DAPI (blue). (b) Color-inverted confocal images showing *ZAM* smRNA FISH signal in early embryos collected 0-2 hours after egg laying (AEL). Mothers were from the indicated genotype. (c) Zoom on the posterior part of a *vret*-sKD early embryo collected at 0-1 hours AEL showing *ZAM* RNA accumulation at the posterior pole. (d) Color-inverted confocal images showing *ZAM* smRNA FISH signal in embryos laid by *vret*-sKD females at the indicated developmental stages.

Figure 6: *ZAM* deletion from the *flamenco* piRNA cluster leads to *ZAM* reactivation and subfertility.

(a) Density plots showing genome-unique piRNAs mapping over 10 kb of the *flamenco* piRNA cluster (without mismatch) where the *ZAM* insertion is located (Release 6: X:21,769,891..21,789,891) in the control (Iso1A) and *flamΔZAM* lines. The position of the *flamenco ZAM* copy *(ZAM-flam)* is indicated. The positions of the sgRNAs used for *ZAM-flam* deletion by CRISPR-Cas9 are shown in red. **(b)** Density plot of *ZAM*-mapping regulatory piRNAs along the *ZAM* sequence in control and *flamΔZAM* ovaries (up to 3 mismatches). **(c)** Scatter plot showing the log2 of normalized counts [log2(RPM)] of antisense regulatory piRNA mapping to individual internal TE sequences in control ovaries (Iso1A) versus *flamΔZAM* ovaries. Antisense piRNA counts, mapped allowing up to 3 mismatches, were normalized per million of genome-mapping piRNAs (RPM). *TEs in red have a *flamΔZAM/Iso1A* ratio <0.3. **(d)** Color-inverted confocal images of ovarioles (upper panels) and stage 10 egg chambers (lower panels) from the indicated genotypes showing *ZAM* smRNA FISH signal. **(e)** Ovaries of the *flamΔZAM* line and of the *flamΔZAM* line with two active *ZAM* copies introduced by genetic crossing with the RevI-H2i2 line (the two euchromatic copies of the X chromosome). **(f)** Box plot displaying the number of eggs laid per fly per day by RevI-H2i2, *flamΔZAM and flamΔZAM* with 2 active *ZAM* copies females. Each dot represents an individual female. The midline indicates the median value, and the box shows the first and third quartile. Error bars indicate the SD. ****p-value* <0.001; ns, not-significant (*p-value* >0.05) (Mann-Whitney test).

Figure 7: No initial piRNA and siRNA response after *ZAM* retrotransposon reactivation
(**a**) Density plot of regulatory piRNAs along the *ZAM* internal sequence in ovaries from a control line (*white*-sKD) and from the *flamΔZAM*-RevI-H2i2 chr III line (up to 3 mismatches). (**b**) Scatter plots showing the log2 of normalized counts [log2(RPM)] of regulatory piRNAs mapping to individual internal TE sequences in control ovaries (*white-sKD*) versus *flamΔZAM*-RevIH2 chr III ovaries. piRNA counts, mapped allowing up to 3 mismatches, were normalized per million of genome-mapping piRNAs (RPM). (**c**) Density plot of *ZAM*-mapping regulatory siRNAs (21nt small-RNAs complexed with Argonaute proteins) along the *ZAM* internal sequence produced in control line (*white*-sKD) and *flamΔZAM*-RevI-H2i2 chr III ovaries (0-1 mismatch). (**d**) Scatter plot showing the log2 of normalized counts [log2(RPM)] of antisense regulatory siRNAs mapping to individual internal TE sequences in control ovaries (*white-sKD*) versus *flamΔZAM*-RevIH2 chr III ovaries. Antisense siRNA counts, mapped allowing up to 1 mismatch, were normalized per million of genome-mapping siRNAs (RPM).

972
973
Figure S1 (related to Figure 1)
(a) Color-inverted confocal images of ovarioles showing *ZAM* RNA signal in ovaries (smRNA
FISH) from the indicated RevI-H2 isogenic lines. **(b)** Color-inverted confocal images of
ovarioles from RevI-H2i2 ovaries showing *ZAM* RNA signal without (left panel) and after
(right panel) RNase A treatment. **(c)** Color-inverted confocal images of ovarioles showing that
ZAM RNA is completely absent in Iso1A, w[1118], and w[IR6] ovaries.

Figure S2 (related to Figure 2)

(a) Density plot of regulatory piRNAs along the *ZAM* sequence in ovaries of the indicated RevI-H2 isogenic lines (up to 3 mismatches). **(b)** Histogram showing the percentage of 5'-overlaps between sense and antisense *ZAM*-derived piRNAs in ovaries of the indicated RevI-H2 isogenic lines. The percentage of 10-nt 5'-overlaps is in red. **(c)** Confocal images of ovarioles with the GFP sensor transgene (GFP in green) and DNA (blue) staining. Ovarioles were from the progeny of a cross between RevI-H2i3 or RevI-H2i4 females and males carrying the pGFP-*ZAM* sensor transgene driven by actin-Gal4. **(d)** Genomic DNA from the RevI-H2 isogenic lines (RevIH-2i2, RevI-H2i3 and RevI-H2i4) was PCR amplified using primers in order to detect the presence of the *ZAM* insertion in *flamenco* or a *ZAM* insertion in piRNA cluster *9* (specific of the RevI-H2 line). Amplification can only occur if *ZAM* is present at the target locus (primer sequences in Table S5). In all tested RevI-H2 isogenic lines, no amplification was observed for *ZAM* in *flamenco,* while a PCR product of ~1000 pb was observed for the *ZAM* insertion in piRNA cluster *9*.

a

RevI-H2i2 *piwi*-GLKD

50μm

b

ns

Figure S3 (related to Figure 3)

(a) Color-inverted confocal images of stage 10 egg chambers from RevI-H2i2 *piwi*-GLKD ovaries showing *ZAM* smRNA FISH signal. (b) Bar plot showing the fold-changes in steady-state *ZAM* RNA levels in total ovarian RNA from the indicated genotypes (normalized to *rp49*). At least three biological replicates and two technical replicates were used. ns: *p*-value >0.05 (Mann-Whitney test). Error bars indicate the SD.

Table Analyzed	Col: Entering replicate data
Column B	
Column A	vs.
vs.	RevIH2i2white-GLKD
Column A	
Mann-Whitney test	
P value	0.8201
Exact or approximate P value?	Exact
P value summary	ns

1017
1018
1019
Figure S4 (related to Figure 4)
(a) Amount of antisense piRNAs mapping to *ZAM* (up to 3 mismatches) in *white-*, *vret-*, and
yb-sKD ovaries (normalized per millions of genome-mapping piRNAs). **(b)** Scatter plot
showing the log2 of normalized counts [log2(RPM)] of antisense Piwi-bound piRNAs mapping
to individual internal TEs sequence in control ovaries (*white*-sKD) versus *yb*-sKD ovaries.
Antisense piRNA counts, mapped allowing up to 3 mismatches, were normalized per million
of genome-mapping piRNAs (RPM). *TEs in blue have a *yb*-sKD/*white*-sKD ratio >3; TEs in
red have a ratio <0.3. **(c)** Color-inverted confocal images of ovarioles (upper panels) and stage
10 egg chambers (lower panels) from the indicated sKD lines showing *ZAM* RNA expression
in follicle cells and in the ooplasm in all conditions.
1030
1031
1032
1033

Early embyos (< 2 h AEL)

Figure S5 (related to Figure 5):
(a) Color-inverted confocal images of early embryos collected 0-2 hours after egg laying
(AEL). Mothers harbor the indicated sKD of the piRNA pathway. *ZAM* RNA signal was
detected by smRNA FISH. **(b)** Projection of confocal images of stage 1 embryos (0-1 h AEL)
laid by *vret*-sKD females showing *ZAM* RNA (smRNA FISH) and Aub protein
immunostaining. Both accumulated at the posterior pole of the embryo where future germ cells
cellularize (Z-projection on 25 stacks).

1051
1052
Figure S6 (related to Figure 6):
Sequence alignment of the RevI-H2 consensus *ZAM* LTR and the *ZAM* solo-LTR, which
remains after CRISPR-Cas9 deletion of the internal *ZAM* sequence in *flamenco* in the
flamΔZAM line. The image shows the RevI-H2 consensus *ZAM* LTR (upper sequence) and the
ZAM solo-LTR (lower sequence) with their TATA-box and polyadenylation signal (pA-signal)
at position 312 of the *ZAM* consensus (upper sequence), the transcription start site (TSS, at
position 327), the start of the poly-A tail at position 341 (CAAGCAGC-(A)n), according to [25].

(a) Density plot of *ZAM*-mapping regulatory piRNAs (23-29 nt small-RNAs complexed with
Argonaute proteins) along the *ZAM* internal sequence in *flamΔZAM*- RevIH2 chr II ovaries (up
to 3 mismatches). **(b)** Scatter plots showing the log2 of normalized counts [log2(RPM)] of
regulatory piRNAs mapping to individual internal TE sequences in control ovaries (*white*-sKD)
versus *flamΔZAM*-RevIH2 chr II ovaries. Antisense piRNA counts, mapped allowing up to 3
mismatches, were normalized per million of genome-mapping piRNAs (RPM). **(c)** Density plot
of total small RNAs of 23 to 29 nt along the *ZAM* sequence in *white*-sKD, *vret*-sKD and *yb*-
sKD ovaries (up to 3 mismatches). **(d)** Sequence logo for the first ten positions in ZAM-
mapping sense small RNAs of 23 to 29nt (from total ovarian small RNA) in *vret*-sKD ovaries.
The nucleotide height represents its relative frequency at that position. **(e)** Density plot of *ZAM*-
mapping Piwi-bound piRNAs along the *ZAM* internal sequence in *white*-sKD, *vret*-sKD and
yb-sKD ovaries (up to 3 mismatches). **(f)** Density plot of *ZAM*-mapping regulatory siRNAs
(21nt small-RNAs complexed with Argonaute proteins) along the *ZAM* internal sequence
produced in *white*-sKD and *vret*-sKD ovaries (0-1 mismatch).

Supplementary Table S1 – ZAM insertions in the RevI-H2i2 genome

number	chromosome	insertion site (Release 6 genome)	orientation	type	locus	comment
1	X	X:1602129-1602133	minus	ZAM-variant	intron of the overlapping Mur-2B and broad genes	ZAM deleted for internal positions 5500-6126 (Repbase ZAM_I)
2	X	X:1980714-1980717	minus	ZAM full-length	intron of gene Hr4	
3	X	X:19249375-19249375	minus	ZAM full-length	intron of gene Hs3st-B	inserted in a Jockey TE which is absent in the reference genome
4	X	X:22257753 or X:22373661	minus	ZAM full-length	piRNA cluster 56 or 13 (George, Jensen et al., 2015)	
5	X	X:23212829	plus	ZAM full-length	piRNA cluster 9 (X:23150001-23545000, George, Jensen et al., 2015)	
6	X	X:23368838-23368841	minus	ZAM-variant	piRNA cluster 9 (X:23150001-23545000, George, Jensen et al., 2015)	ZAM deleted for internal positions 5500-6126 (Repbase ZAM_I)
7	X	X:23513106-23513113	minus	ZAM-variant	piRNA cluster 9 (X:23150001-23545000, George, Jensen et al., 2015)	ZAM deleted for internal positions 5500-6126 (Repbase ZAM_I)
8	2L	2L:8158963-8158966	plus	ZAM full-length	intron of gene PAPLA1	inserted in a NOMAD TE which is absent in the reference genome
9	2L	2L:15773891-15773894	plus	ZAM full-length	intergenic region	
10	2R	2R:1808663-1817084	plus	ZAM full-length	intergenic region	ZAM copy also present in the reference genome Release 6
11	2R	2R:6793620-6793623	plus	ZAM with an internal deletion	intron of gene Epac	ZAM deleted for internal positions 2332-5754 (Repbase ZAM_I)
12	2R	2R:11601808-11601811	minus	ZAM full-length	intron of gene toutatis	
13	3L	3L:4299131-4299137	minus	ZAM full-length	intron of gene dyl	ZAM deleted within the 5'-UTR, internal positions 60-1218 (Repbase ZAM_I)
14	3L	3L:19980882-19980883	minus	ZAM-variant with additional internal deletion	intergenic region	ZAM deleted for internal positions 13-2993 and 5500-6126 (Repbase ZAM_I)
15	3R	3R:12079947-12079950	plus	ZAM-variant	intron of gene ATP8B	ZAM deleted for internal positions 5500-6126 (Repbase ZAM_I)
16	unknown	NW_007931121.1:45136-45142	plus	ZAM full-length	rDNA	
17	unknown	possibly close to X:23239427	(plus)	ZAM with internal deletions	possibly piRNA cluster 9 (X:23150001-23545000, George, Jensen et al., 2015)	ZAM deleted for internal positions 2332-5754 and 6116-7265 (Repbase ZAM_I)
18	unknown	NW_001845412.1:1075-1078	minus	ZAM with an internal deletion	unknown	ZAM deleted for internal positions 2929-5615 (Repbase ZAM_I)

1075

Supplementary Table S2 – Drosophila lines

Category	Gene name	CG_Number	Stock number	Genotype	Source	Notes
RNAi line	white	CG2759	BDSC_35573	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GL00094}attP2	Bloomington	
RNAi line	armitage	CG11513	BDSC_34789	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS00098}attP2	Bloomington	
RNAi line	yb	CG6272	VDRC_110056	P{KK105851}VIE-260B	VDRC	
RNAi line	Piwi	CG6122	BDSC_37483	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GL00626}attP40	Bloomington	
RNAi line	vreteno	CG4771	VDRC_34897	w1118; P{GD11424}v34897	VDRC	
RNAi line	zucchini	CG12314	BDSC_35227	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GL00111}attP2	Bloomington	
RNAi line	Argonaute 3	CG40300	BDSC_35232	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GL00117}attP2	Bloomington	
RNAi line	Aubergine	CG6137	BDSC_35201	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.GL00076}attP2	Bloomington	
Wildtype	white (w1118)		BDSC_3605	w1118;;;	Bloomington	
Nos-Gal4 driver			BDSC_4937	w[1118]; P{w[+mC]=GAL4::VP16-nos.UTR}CG6325[MVD1]	Bloomington	
Traffic jam gal4 driver				w, Tjgal4/cyO; ;	Unknown	
RevI-H2i1					Produced in the lab	Isogenic line derived from the RevI-H2 stock
RevI-H2i2					Produced in the lab	Isogenic line derived from the RevI-H2 stock
RevI-H2i3					Produced in the lab	Isogenic line derived from the RevI-H2 stock
RevI-H2i4					Produced in the lab	Isogenic line derived from the RevI-H2 stock

1076

Supplementary Table S3 – smRNA FISH probes

Oligo number	ZAM (this paper) 5'- 3'	Burdock (from Senti and al,.2015) 5' - 3'
1	GAAAAAGAAATGTTGGCAAT	AGGTTATCACTGTCACTCAT
2	AATATGGGCAGTAAAATACT	GCTTGACACTAGGTTGTCTA
3	AGACCATATATTTATGGCGT	TGGTGCGCTGACCATTTATT
4	TACAGATCACAAGCCACTAA	TTTTTTCTGCACTGTTTTGC
5	TGAATTTCAAAGAACCCAAC	CCACCAATTATCTGATGACT
6	AAAAATAATTCGTTGGAGAC	CGCTAGTTGTCTTTGTTTTT
7	TCCAACTCATGGAATACAAT	AACTGAGCTTCAAATTCCAT
8	TCACAAGAAAGGTTCACAAA	AACAGCACTCTCTACGATCG
9	TGTAATTGCAGACGCCTTAA	TGATTGTTTTTGGACTGCGA
10	GTAGAGCGGACCCAAATTTA	TCATTCATCTCCTGCAATTG
11	AACTACAACGAAACACTGAC	GGTGTTCACTGTTAATTTCC
12	TGTTAAGCCTTGCCCCACAT	CTACATAAGTTTCCACCTCT
13	CCGAAAAACCTATTAACGAA	TATATCTAGAGGCTCGCTAC
14	AACACGCAACTCATACTAGA	GCCATCAAAATCTGGCAGAG
15	TCTTACCAAACTACAACACC	CTTCCCTCGTAATCTTTGAA
16	AGGAAAAAATATTCACAGCC	CCAAGAGCTTGGTAAAATGT
17	AAAATCTTGAAAGGAACCCT	AAGCCAATACTGTATTCGCT
18	AAAACCTAACAGGATTTGCG	GCTTTGAAATGTAACGGAGT
19	CATTCTTGGCGGACGATAAT	TGTGAAATCAAGACGGCTGA
20	TCCGCATTAATCGAAAAAGC	TAGATAGATCGGCCTTTTGT
21	TTCTCCACGAAATCGTAGGA	GTCGCAAAGTTGATAGCTCT
22	ACCCCGAACAAAACAAATTC	TCAGTAAGAGTCATGTCTCC
23	GGAATATCACACTAACAGCA	TCTTGTTGGTAAGTAGGGTC
24	GAGGCATAGACGAAACATTC	GCCAAGGCACTATCAAATGT
25	TTCACCTCAAACGAGAAACC	GTCCTGTACTTTTCATTCAG
26	ACTTCCCCAATATGAAAAAC	CTCCGGTTACAAATACACGT
27	AATTAGGAATTGCGAAACCT	ATGTCGCTCAACGATTTCTT
28	GTCTAAAACTCAAATACGAC	ATCTTTTGGACCTTTTGCAA
29	AGGCAACCACAAAATATAGT	TGGTTCGACTCGACCTCTTG
30	ATTTGAAACCCCAGAAACCC	AAGGGCGAATTGGTAACGCT
31	GAAACCCCTCGACATAATAC	CCCTGTCTCCAATATTTTTA
32	ACCTGACAATCATAGACAAA	TGCCTTTGTTCGATTTTCTG
33	TCTCAAAATTCGCAGCTGTC	TGAGTTTCTATCCTTATCGC
34	TACCCCATCCCAAATAGAAA	TTTTTAGTTTGCATGGGCAT
35	CGGCATCAATTGCATCAAAG	CACCTGACGCTTGCTAAAAT
36	TTCAGTCAATTCGGACTACC	GGATTTGTTAACTGGACTGG
37	CAAAAAACTAATACACGACC	TTCTTCTAGATCGCATGGAT
38	AGGAGTAGAATTTTGCAACG	GGAGTTTGTCCAAAATTTCC
39	CCATGTCACATCCTTCCAGC	AATATTTCCCTGAGTCGGAA
40	AGTAATTCTCCAGTAGAACG	GGTTGCGATGCGAATTGATT
41	ACACTCCTCTTTGACAGAGA	CTGACTTGTGTTTTGCTGTT
42	AATACTAGACACACGGAAAA	TTGCGCATACCCAAAATGTC
43	ACACAAATTACCTACAGACC	CACTCGTTGGTCTTTTTGAT
44	ACGAAGAAATAATGTCAGAA	TTTTGGCCCTGTAAACCTTG
45	ATAACAACGCAATCCACTCC	GGTAAGTAGTTGATCCTCTG
46	CCACCAAACACACCCCTTTT	CTCACATTGACCTTTCTCAT
47	TGGTAGGACCCATTTATTCG	CCTTTTGATAACCGTCTGTA
48	ATACCCAATAATGAGCATGA	ATAATCATCAACCTCCGCTT

1077

Antibody	Source	Dilution	Fluorochrome	Species	Reference
Anticorps primaires					
Goat anti-GFP	abcam	1/1000			ab5450
Rabbit anti-Aubergine	Homemade (Eurogentec) - polyclonal antibody	1/1000			
Rabbit anti-Gag-ZAM	Home-made	1/100			
Rabbit anti Env-ZAM	Home-made	1/1000			
Secondary antibody					
Alexa Fluor 488 donkey anti-goat IgG (H+L)	Invitrogen (Fisher)	1/1000	Alexa 488	goat	A11055
Donkey anti-rabbit Alexa 488 IgG (H+L) 2mg/ml	life tech	1/1000	Alexa 488	rabbit	A21206
Goat anti-rat Cy3	Jackson Immunoresearch	1/1000	Cy3	rat	112-165-143
Fluorescent dyes					
DAPI	Sigma	1/500			D8417-1MG

1078

Supplementary Table S5 - Primers used

	Name	Sequence	Used for
Rp49	rp49_R	AAACGCGGTTCTGCATGAG	RT-qPCR
	rp49_F	GACGCTTCAAGGGACAGTATCTG	
Burdock	Burdock_F	GGACAAACTCCATTTCCGACTC	RT-qPCR
	Burdock_R	TCCCTGAGCCTGACTTGTGT	
Gypsy	Gypsy-F	CGCTCGAAGGTTACCAGGTAGGTTC	RT-qPCR
	Gypsy_R	CTTCACGTTCTGCGAGCGGTCT	
ZAM	Zam_env_F	CTCCCAATCATCTCCTCCAA	RT-qPCR
	Zam_env_R	AGAAGTCGGGGTGCTCTTCT	
ZAM-flam	ZAM-FLAM_F	TCGCTTAGGAGGGTAGCTTG	CRISPR SCREEN
	ZAM-FLAM_R	CTTAACACCTACTGTCAGTGC	
ZAM insertion in *flamenco*	flam_R	ctagtgtctttggtaccatgg	Screen
	flam_F	tatcaactatcacgactatcact	
ZAM insertion in *cluster 9*	ZAM insertion_R	CTCACCATTTCCTCCTTGAC	Screen
	ZAM insertion_F	CTCCCAATCATCTCCTCCAA	

1079

Supplementary Table S6 – small-RNA seq

Small RNA seq
- sRNA from Revl-H2i1
- sRNA from Revl-H2i2
- sRNA from Revl-H2i3
- sRNA from Revl-H2i4
- sRNA- from Revl-H2i2- nos-Gal4- expressed dsRNA-white
- sRNA- from Revl-H2i2- nos-Gal4- expressed dsRNA Ago3
- sRNA- from Revl-H2i2- nos-Gal4- expressed dsRNA Piwi
- sRNA- from Revl-H2i2- nos-Gal4- expressed dsRNA Zuc
- sRNA from tj-Gal4-expressed dsRNA-Vret
- sRNA from tj-Gal4-expressed dsRNA-white
- sRNA from CRISPR flamΔZAM
- Total RNA from tj-Gal4-expressed dsRNA-Vret
- Total RNA from tj-Gal4-expressed dsRNA-white
- Total RNA from tj-Gal4-expressed dsRNA-yb

1080