Boolean modeling of mechanosensitive Epithelial to Mesenchymal Transition and its reversal

Emmalee Sullivan[1,2], Marlayna Harris[1,3], Arnav Bhatnagar[1,4], Eric Guberman[1,5], Ian Zonfa[1,6], Erzsébet Ravasz Regan[1]

[1] Biochemistry and Molecular Biology, The College of Wooster, Wooster, OH
[2] Ohio University Heritage College of Osteopathic Medicine, Cleveland, OH
[3] Stanford University School of Medicine, Stanford, CA
[4] Molecular, Cellular, and Developmental Biology, UC Santa Barbara, Santa Barbara, CA
[5] Clinical Trial Operations Development, AbbVie, Chicago, IL
[6] Case Western Reserve University, Cleveland, OH

ABSTRACT

The significance of biophysical modulators of the Epithelial to Mesenchymal Transition is demonstrated by experiments that show EMT on nano-patterned ECM without biochemical signals, or stiffness-induced / shape-dependent induction of EMT transcription factors. Yet, current models focus on biochemical triggers without addressing the mechanosensitive drivers of EMT. Here we built a Boolean model of EMT triggered by crosstalk between mechanosensing and mitogens. Our model reproduces epithelial, hybrid E/M and mesenchymal phenotypes, the role of autocrine *TGFβ* signaling in maintaining mesenchymal cells in the absence of biomechanical drivers, inhibition of proliferation by *TGFβ,* and its apoptotic effects on soft ECM. We offer testable predictions on the density-dependence and molecular drivers of partial EMT and the tug of war between mitosis and the stability hybrid E/M state. Our model serves as a stepping-stone towards modeling the effects of the biomechanical environment on cancer cell stemness linked to the hybrid E/M state, as well as the mutually inhibitory crosstalk between EMT and senescence.

INTRODUCTION

Carcinomas emerge from healthy epithelial tissue due to oncogenic mutations, their growth enabled by the loss of contact inhibition of proliferation [1,2]. The weakening of cell junctions is critical for tissue invasion and migration leading to metastasis. This transformation from epithelial to an invasive, migratory mesenchymal phenotype is called the epithelial to mesenchymal transition (EMT) [3,4]. A rate-limiting step in the development of metastatic cancer, EMT remodels the cytoskeleton to enhance migration, disrupts the ECM with proteolytic enzymes to promote tissue invasion, and enhances resistance to apoptosis as well as senescence [5,6]. The reverse mesenchymal-epithelial transition (MET) often aids the settling of these migratory cells into distant metastases [7]. As metastatic cancers account for 90% of cancer deaths worldwide [8], development of effective therapeutic approaches requires predictive models of the

complex interaction between contact-mediated signaling, engagement of the EMT transcriptional program, and its reversal during MET.

A closer look at cancer cells within solid tumors revealed a striking cell-to-cell heterogeneity in the expression of epithelial and mesenchymal markers [9–12]. The initial picture of an all-or-nothing transition from an epithelial to a mesenchymal state gave way to the insight that at least one but perhaps several stable cell phenotypes between these two extremes exist *in vitro* and *in vivo* [7]. These hybrid E/M phenotypes show characteristics of both epithelial and mesenchymal cells, including an ability to maintain adherens junctions while polarizing horizontally to migrate in a mesenchymal fashion [13]. As a result, hybrid E/M cells migrate and invade as clusters [14], which gives them an advantage in seeding colonies [15,15–17], traversing narrow capillary vessels by forming single-cell trains [18]. Indeed, hybrid E/M cells were shown to have up to 50-fold greater metastatic potential than single circulating cancer cells [15]. In healthy epithelial tissues such as blood vessels, lung, or liver epithelia, balancing birth and death in response to injury is complemented by partial EMT and migration at monolayer wounds [3]. Accompanied by proliferation, this helps wounded epithelial sheets migrate as collectives, close wounds, and use MET to reestablish their junctional barriers. Yet, the precise cellular contexts that generate hybrid E/M rather than mesenchymal cells are not well understood [7].

EMT is triggered by a series of transforming signals such as *TGF-β, Wnt, Notch, Hedgehog* and *IL6*, aided by growth factors, hypoxia and stiff/fibrous ECM [19–21]. These influences converge on a core transcriptional circuit rich in positive feedback that can robustly maintain both epithelial and mesenchymal states [22]. The *Snail* (*Snail*) transcription factor is a key convergence point; its double-negative feedback with the epithelial micro-RNA *miR-34* is the first switch in the transcriptional cascade of EMT [23]. Once fully active, *Snail* lowers the barrier for toggling a second switch generated by mutual inhibition between *miR-200* microRNAs and the transcription factor *Zeb1* by lowering *miR-200* expression while promoting *Zeb1* [23]. High *Zeb1* and *Snail* activity fully repress *E-cadherin* transcription, leading to complete disassembly of adherens junctions [24]. Mesenchymal cells then secrete *TGFβ* to engage an autocrine positive feedback loop that stabilizes their phenotype in the absence of external stimuli [25,26]. This core regulatory architecture was shown to generate a three-state cascading switch that can maintain a distinct epithelial, hybrid E/M and mesenchymal phenotypes [27].

This four-node *Snail / miR-34 / Zeb1 / miR-200* circuit is far from capturing the richness of the regulatory links and feedback that modulate EMT. Yet its central role in generating the nonlinearity of the transition allowed computational models detailing their dynamics to reproduce and predict a wide range of experimentally observed behaviors [22]. For example, the existence of a stable hybrid E/M phenotype was predicted by an ODE-based computational model of this core network [26,27] before it was confirmed *in vitro* [23,28–30] and *in vivo* [31–34]. A series of continuous models followed to explore the modulating effect of additional regulators, such as phenotypic stability factors that increase the robustness or number of hybrid E/M states [35,36], the

relationship between the hybrid E/M state and de-differentiation to a stem-like state [30], the effect of *Notch*-mediated spatial patterning in tumors [37], or biological noise generating population heterogeneity along the EMT spectrum [38]. Integrating the crosstalk between signaling pathways controlling EMT and the core transcriptional network was accomplished by a large Boolean network model which linked *TGF-β, Wnt, Notch, Hedgehog* and growth signaling to well-established EMT TFs [39]. They identified generalized positive feedback motifs (stable motifs) that lock in the mesenchymal state, involving feedback between the core EMT network, major growth pathways, and autocrine *TGF-β/Wnt/Hedgehog* signaling. They went on to characterize perturbations that stabilize hybrid E/M, providing testable interventions to control the network [40,41]. A subsequent study using roughly the same network but simpler rules that sum positive and negative influences showed that the system can generate hundreds of metastable hybrid states along a spectrum between epithelial and mesenchymal cells, mimicking the heterogeneity of bulk and single-cell gene expression data [42].

As Boolean models characterize regulatory signals as expressed/active or absent/inactive [43], they rarely deal with processes that depend on the uneven distribution of regulatory molecules in different parts of a single cell. Yet, several key aspects of EMT controlled by the cell's biophysical environment involve such asymmetry. The transition from apical/basal polarity in an intact monolayer to horizontal polarity and directed migration at lower density requires a different set of regulatory molecules to be ON at the cell's leading vs. trailing edge [44]. In a recent study modeling the crosstalk between epithelial contact inhibition of proliferation, migration, cell cycle progression, and anoikis we focused on the mechanosensitive regulation of cell behaviors by ECM stiffness and a cell's neighbors [45]. Rather than building a space-aware 3D model, we captured key features of the biology in a Boolean framework by modeling the logic of pathways that push cells past discrete thresholds: anoikis as cells loose ECM anchorage, establishment of apical/basal polarity and contact inhibition in dense monolayers, and the coupling between contact inhibition of proliferation and migration. Our model predicted a metastable migratory state in non-dividing cells at low density but stopped short of integrating these mechanosensing mechanisms with the regulation of EMT-driving transcription factors.

The importance of the biophysical microenvironment in controlling EMT is demonstrated by a series of experiments published EMT model cannot yet reproduce. These include the stiffness-dependent effects of *TGF-β,* leading to apoptosis on soft ECM vs. EMT on stiff substrates [46], the demonstration of full EMT in the absence of any biochemical transforming signal on stiff nano-patterned ECMs [47], mechanical stress-mediated EMT transcription factor induction [48–50], or the effect of cytoskeletal and nuclear morphology on the transcriptional regulation of EMT [51]. Yet, we lack predictive mechanistic model that can reproduce and predict the effects of a single cell's biomechanical environment on its behavior in different signaling contexts, such as varying growth stimuli or transforming signals. Moreover, current computational models only address the process of MET in response to either the withdrawal of input signals that activate *Snail* [26,27,30], or to drugs / mutations [40,41].

To address these needs, we have built on our published mechanosensitive model of epithelial contact inhibition, proliferation, and apoptosis [45] to model the process of EMT triggered by biomechanical and growth signaling crosstalk. Our goal was to examine the stability of hybrid E/M and M states in the absence vs. presence of autocrine *TGFβ* signaling. In addition to EMT regulation (**Fig. 1A-B**), our 150-node Boolean model explicitly accounts for the regulatory networks driving growth signaling, cell cycle commitment and progression, apoptosis, adhesion to the ECM and to neighboring cells, contact inhibition, and key regulators of migration (**Fig. 1C**). We reproduce a range of cell behaviors linked to EMT, including 1) the ability of the core EMT transcriptional network to maintain distinct epithelial, hybrid E/M and mesenchymal states, 2) EMT driven by strong mitogens such as EGF on stiff ECMs [52,53], 3) increased mesenchymal transcription factor activity and EMT in response to the loss of *E-cadherin* [54,55]; 4) TGFβ-induced EMT and the need for strong autocrine TGFβ signaling to maintain the mesenchymal state in the absence of mitogens, on softer matrices, or at high density [25], 5) TGFβ-induced inhibition of proliferation [56]; 6) TGFβ-induced apoptosis on soft ECM [46]; 7) anoikis resistance in mesenchymal cells [57] driven by TGFβ [58], and 8) lack of TGFβ-induced EMT in very dense monolayers [59]. In addition, we offer experimentally testable predictions on the effect of neighbors on partial vs. full EMT, the tug of war between mitosis and the maintenance of migratory hybrid E/M states, the biomechanical triggers of MET and its effect on autocrine TGFβ signaling, and finally, cell cycle defects in heterogeneous populations of epithelial, hybrid E/M and mesenchymal cells.

METHODS

Boolean modeling methods. Boolean network models focus on the complex combinatorial logic by which molecular species and pathways intersect, sacrificing the precise concentration and time-dependence of interactions in favor for scalability [43,60]. The ON (expressed and active) vs. OFF (not expresses or inactive) response of each molecule to every combination of its regulators dictate the model's temporal dynamics (**Table S1**). Ideally, Boolean regulatory rules should be directly inferred from experimental data. Yet, detailed computer-accessible time-series data on the dose-dependent response of single cells to their biochemical environment, along with molecular activity responsible for these responses are not currently available. Instead, we compiled qualitative experimental data from 479 papers into a 150-node Boolean model in which each of the 630 links is justified by experiments (**Table S1**; *blue:* changes from [45]; *green*: further additions in version with *TGFβ*). Logic gates were inferred from literature and/or chosen based on reported effects of molecular input combinations. Each iteration of the model described in *Results* is available in BooleanNet format (**Files S1-3**: EMT switch, model without / with *TGFβ* module) [61] and in SBML format on the BioModels database (MODEL2208050001, MODEL2208050002; *https://www. ebi.ac.uk/biomodels*) [62]. Editable graphic layouts from **Figs. 1C** and **S3** are available as glm files (**Files S4-5**).

Our analysis methods are unchanged from [45,63]. Briefly, we used synchronous Boolean modeling to find every stable phenotype and/or oscillation (attractor) of the model [45,63,64]. The update-robustness of our cell cycle / apoptosis model was examined in [63]. To test the effect of non-saturating, intermediate environmental inputs we simulated the model while overriding one or more input node in each time-step with a stochastically generated ON/OFF value of a set probability, thus tuning the cell's average "signal exposure" between 0 (none) to 1 (saturating). Similarly, we model partial or complete knockdown/hyperactivation of a node by forcing it OFF/ON with a set probability in each time-step, otherwise leaving it to obey the Boolean rule that normally governs its behaviors [45,63]. Finally, we use an automated process to assign specific biological phenotypes to any state of the model and monitor phenotypic transitions; namely healthy vs. erroneous cell cycle progression, apoptosis, as well as contact inhibited, migratory (signatures detailed in [45]), epithelial, mesenchymal and hybrid E/M states, defined as:

Epithelial   = (miR_34=1, miR_200=1, $Ecadherin\_mRNA$=1, $ZEB1\_H$=0)
Mesenchymal = (miR_34=0, miR_200=0, $Ecadherin\_mRNA$=0, $ZEB1\_H$=1, $N\_bcatenin\_H$=1)
Hybrid_EM   = (miR_34=0, miR_200=1, $Ecadherin\_mRNA$=1, $ZEB1\_H$=0, $ZEB1$=1, $SNAI1$=1)

Previously published network modules are briefly described in *Supplementary Methods* (details in [45,63,64]), followed by in-depth description of new pathways (*NF-κB*, *TGFβ* signaling, EMT).

RESULTS

Modeling the epithelial to mesenchymal toggle switch

To model the effect of mechanosensitive signals on EMT, we added an EMT switch to our contact inhibition model [45] (**Fig. 1A**). We built this switch around feedback loops common to published continuous and Boolean models [26,27,39]. These include direct double-negative feedback between *Snai1* and the epithelial microRNA *miR-34* [65] as well as between *Zeb1* and *miR-200* [66]. They are reinforced by positive feedback between *Snai1*, *Zeb1*, nuclear *β-catenin* and the two miRs [67–70]. This switch controls *E-cadherin* mRNA expression, repressed cooperatively by mesenchymal transcription factors [71–73]. As *Zeb1* is not expressed in epithelial cells, expressed at a moderate level in hybrid E/M states, and is only high in mesenchymal cells [74,75], we used two Boolean nodes to distinguish between these three levels of *Zeb1*. Similarly, we modeled the effects of adherens junctions on *β-catenin* nuclear localization with two nodes, distinguishing between a) cells entirely surrounded by neighbors in a dense monolayer (all *β-catenin* is localized to adherens junctions; both nodes OFF), b) cells firmly attached to the edge of a monolayer (some but not all *β-catenin* is tied to adherens junctions; medium nuclear localization, *N_bcatenin*=ON but *N_bcatenin_H*=OFF), and c) cells that do not touch their neighbors (all *β-catenin* is freed from junctions even if *E-cadherin* is expressed; nuclear localization is maximal as long as *GSK3β* is blocked by growth signals; both nodes ON). **Fig. 1A** highlights key signals that converge on the EMT module from cell junctions (*blue*), growth signaling (*green*), and the migration prompting *Pak1* kinase (*pink*).

Modeling the EMT switch in isolation reveals three stable phenotypes (**Fig. 1B**; automated isolation of a module described in [46]). As expected, its epithelial stable state is characterized by *E-cadherin*, *miR-34* and *miR-200* expression, and no transcriptional activity in EMT driver genes (**Fig. 1B**, *top*). The opposite pattern of expression and activity characterizes the mesenchymal state, a complete flip of the switch (**Fig. 1B**, *bottom*). In between the two, the network maintains a stable hybrid E/M state characterized by fully suppressed *miR-34* but not *miR-200* [27], along with *Snai1*, *Twist*, *Snai2* and moderate activation of both *Zeb1* and nuclear *β-catenin* (**Fig. 1B**, *middle*). The regulatory environment outside the switch can bias or dictate the state(s) it can maintain in different environments. To model this we linked the EMT switch into our contact inhibition model [45], where it is influenced primarily by *i)* mechanosensing and junction-mediated contact inhibition via *β-catenin* localization, *ii)* migration via *Pak1,* and *iii)* growth signaling via *Akt1*-mediated *NF-κB* activation and *GSK3-β* suppression (**Fig. 1A,C**).

Figure 1. The transcriptional circuit driving EMT is a tree-way switch influenced by cell junctions, migration, and growth signaling. A) Regulatory switch controlling EMT (*yellow*) with relevant inputs from regulators of cell cycle (*purple, lavender*), adhesion (*light & dark blue*), growth (*greens*), migration (*pink*) and apoptosis (*dark red*). **B)** Three stable attractors of the isolated EMT regulatory switch. *Orange/blue node borders:* ON/OFF. **C)** Modular network representation of our extended Boolean model. *Gray:* inputs representing environmental factors; *dark blue:* Adhesion signals; *red: TGFβ* signaling; *green*: Growth Signaling (*lime green*: basal AKT & MAPK, *bright green:* PI3K/AKT oscillations, *mustard*: mTORC1, *dark green: NF-κB, GSK3, FoxO1*); *dark red:* Apoptotic Switch*; light blue:* Contact Inhibition; *pink/light orange:* Migration; *light brown:* Origin of Replication Licensing; *lilac:* Restriction Switch; *purple:* Phase Switch; *dark orange:* cell cycle processes; *yellow:* EMT switch; → : activation; ⊣ : inhibition.

Growth factors cooperate with matrix stiffness and loss of cell-cell contacts to drive reversible mechanosensitive EMT

The key difference between the EMT model on **Fig. 1C** and previous models is that it does not include the traditional exogenous drivers of EMT such as *TGF-β, Wnt*, *Notch*, *Sonic Hedgehog*, or *IL-6* [39,76]. Instead, we focus on signals that can trigger EMT in the *absence* of the above signals – namely the cell's biomechanical environment. Evidence from different cell lines indicates that epithelial cells undergo an epithelial to mesenchymal transition when plated on stiff substrates [47–50]. To model this, we started by scanning all stable phenotypes of our model on a soft ECM at different cell densities and growth factor exposures. Due to an inability to form stress fibers and activate *YAP* on soft ECM, the only stable model phenotype is epithelial – even at very low densities that preclude adherens junctions (**Fig. 2A,** *initial state*).

Figure 2. Strong growth signals on a stiff matrix can drive the epithelial to mesenchymal transition, leading to mesenchymal cells at very low density vs. a hybrid E/M state at monolayer edges. A-D) Dynamics of regulatory molecule expression/activity during exposure of (A) isolated, growth-stimulated epithelial cells to stiff ECM, (B) isolated epithelial cells in low growth conditions on a stiff ECM to strong mitogens, (C) isolated, growth-stimulated epithelial cells at a monolayer edge to stiff ECM, and (D) isolated epithelial cells in low growth conditions on a stiff ECM to a brief pulse of strong mitogens. *X-axis:* time-steps; *y-axis:* nodes organized in regulatory modules; *orange/blue:* ON/OFF; *black/white labels & arrows:* molecular changes that drive EMT.

Plating our model epithelial cells on a stiff matrix triggered stress fiber formation, *YAP* activation and *Rac1/Pak1*-driven migration (**Fig. 2A**, transition in *Migration* module at the start of *Stiff ECM* exposure). In addition, *YAP* led to the induction of high *PI3K 110* (*p110)* protein expression [77]. In growth-promoting environments (serum, EGF, HGF), high expression of *p110* increases the potency of *PI3K / AKT* signaling, which activates *NF-κB* and turns off *GSK3β* [78]. *NF-κB*, in turn, lead to *Snail* transcription, stabilized by low *GSK3β* activity [79,80]. *Snail* nuclear localization is subsequently aided by *Pak1*-mediated phosphorylation, a key early event leading to EMT [81]. Thus, *YAP*-mediated sensing of a stiff ECM works cooperatively with growth signaling to promote *Snail* activation and subsequent *miR-34* repression by *Snail*. Without *miR-34* to lower *β-catenin* and neighbors to trap it at adherens junctions, nuclear *β-catenin* levels rise sufficiently to support high *Zeb1* activation, *miR-200* repression, and full EMT (**Fig. 2A**, *final state*). This effect is followed by cell cycle entry, which does not require EMT but is also supported by growth signals and a stiff ECM (**Fig. 2A**, *cell cycle oscillations*). In contrast, plating cells on a stiff ECM in the *absence* of strong growth signaling does not result in high *AKT* signaling, migration, *Snail* activation, or EMT (**Fig. S1A**).

An alternate route to the same mesenchymal phenotype is to start with isolated cells plated on a stiff ECM in the absence of strong growth signaling. Exposing these cells to growth factors also triggers EMT via the same cascade of signals (**Fig. 2B**). Modeling intermediate levels of stiffness and growth signal exposure by stochastically toggling the *Stiff_ECM* and/or *GF_High* nodes ON for a fraction of the time also shows that cells on stiffer matrices spend increasing amounts of time in a mesenchymal state (**Fig. S1B,** *top row, orange*). The response to growth signals is similar, except that we predict a high prevalence of hybrid E/M cells at intermediate mitogen exposure (**Fig. S1B,** *bottom row, orange*).

Modeling the emergence of hybrid E/M states: cells at the edge of a monolayer undergo partial EMT in response to stiff ECM exposure

Repeating the above simulations with cells placed in cell densities akin to the edge of a confluent monolayer leads to a different outcome. As **Fig. 2C** indicates, exposing cells that maintain adherens junctions with some neighbors but not enough to establish apical/basal polarity respond to stiff ECM exposure by undergoing partial EMT. In this case, *Snail* activation and *miR-34* inhibition do not result in high nuclear *β-catenin.* Instead, cells stabilize in a state that matches the observed transcription factor expression and phenotype of hybrid E/M. Namely, *Snail*, *Twist1* and *Snai2* are expressed, *Zeb1* activity is moderate, and *miR-34* is repressed but *miR-200* is still active (**Fig. 2C**, EMT module). These cells are migratory and have reduced *E-cadherin* expression compared to dense monolayers. Yet, they maintain adherens junctions with neighbors and migrate as a group (**Fig. 2C**, *Migration* and *CIP* modules). Modeling intermediate levels of stiffness also shows that cells with adherens junctions spend an increasing amount of time in a hybrid E/M on stiffer substrates, but do not undergo full EMT (**Fig. S1C**).

Interestingly, we predict that while a sufficiently stiff ECM to sustain stress fiber formation is mandatory for entry into and maintenance of a hybrid E/M phenotype, sustained *growth* signaling is not necessary. Rather, entry into hybrid E/M only requires a short-lived boost of growth signaling (**Fig. 2D**). Once established, positive feedback linking the *EMT* and *Migration* modules can sustain hybrid E/M. Consequently, even weak / transient growth factor exposure has a large effect on the amount of time these cells spend in a hybrid E/M state (**Fig. S1C,** *bottom right*). Saturating growth signals actually *decrease* the stability of the hybrid E/M state.

To further probe the importance of adherens junctions in partial vs. full EMT, we ran a large sample of cells stochastically exposed to a matrix with relatively high stiffness (*Stiff ECM* = ON in 95% of time-steps) and near-saturating growth factors (95% GF_{High} = ON). In this environment, a randomly sampled cell population will be 40% epithelial (**Fig. S1D,** *leftmost blue bar*s), ~55% in hybrid E/M (*leftmost orange bar*), and 0% mesenchymal (the remaining 5% are in transition). Increasingly potent inhibition of junctional *E-cadherin* leads to a moderate decrease in epithelial states, but a drastic shift from hybrid E/M to full mesenchymal cells (**Fig. S1D**). This simulation, akin to either experimental inhibition of *E-cadherin* or gradual decrease in density, underscores the importance of cell contacts in stabilizing the hybrid E/M state even in the absence of *Notch*-mediated juxtracrine signaling [37].

Modeling the Mesenchymal to Epithelial Transition: in the absence of strong autocrine signaling, loss of YAP or ERK activity reverses EMT

As our model lacks autocrine signals that stabilize the mesenchymal or hybrid E/M state, their stability relies on ongoing feedback between the EMT switch, growth signaling, and migration. To investigate the consequences of breaking this feedback, we simulated the response of isolated mesenchymal cells proliferating on a stiff matrix to two types of environmental change. First, we simulated their re-plating onto very soft ECM, where the maintenance of stress fibers, *YAP* activity, horizontal cell polarity and migration is no longer possible (**Fig. 3A**). This led to a loss of *Pak1* signaling, nuclear exclusion of *Snai1*, and MET. Indeed, even partial matrix softening was sufficient to lock cells into an epithelial state, as mesenchymal entry below 50% *Stiff_ECM* is rare (**Fig. S1B,** *top row*).

Placing mesenchymal cells into a high cell density environment has a similar effect to soft ECM, as it also triggers MET (**Fig. 3B**). This, however, only occurs at densities that forbid cell spreading and stress fiber maintenance. Placing a mesenchymal cell into a neighborhood that resembles the edge of a monolayer does not alter its mesenchymal state (**Fig. S2A**). The presence of neighbors alone fails to trigger partial MET to a hybrid E/M state because fully mesenchymal cells lack *E-cadherin* expression and do not form adherens junctions. Thus, MET only occurs in our model when feedback between the EMT switch and migration is interrupted, and *Snai1 /*

Zeb1 activity is lost. Only then do cells re-express *E-cadherin* and establish adherens junctions. To highlight the key role of *YAP* and *Pak1* in the maintenance of the mesenchymal state, we modeled partial *YAP* or *Pak1* knockdown in a 95% *Stiff_ECM,* 95% *GF$_{High}$* environment. As **Fig. 3C** indicates, partial downregulation of both *YAP* or *Pak1* can trigger MET, despite the lack of neighbors and a stiff ECM. In contrast, forced activation of *Pak1* can partially counteract MET at relatively high density by stabilizing hybrid E/M, while hyperactive YAP alone cannot (**Fig. 3D**).

Figure 3. The mesenchymal to epithelial transition (MET) is triggered by both soft matrix exposure and high cell density in the absence of autocrine or external EMT-promoting signals. A-B) Dynamics of regulatory molecule expression/activity during exposure of an isolated, growth-stimulated and proliferating mesenchymal cell to (A) soft ECM, and (B) very high cell density. *X-axis:* time-steps; *y-axis:* nodes organized in regulatory modules; *orange/blue:* ON/OFF; *black/white labels & arrows:* molecular changes that drive MET. **C)** *Top/bottom:* fraction of time spent in a mesenchymal (*orange*) vs. epithelial (*blue*) state as a function of increasing *YAP1* (*top*) or *PAK1* (*bottom*) inhibition in isolated cells on 95% *Stiff ECM* and exposed to 95% saturating growth stimuli. **D)** *Top/bottom:* fraction of time spent in *i*) mesenchymal (*orange*) vs. epithelial (*blue*) state

The fact that even transient interruption of migration can block *Snai1* and flip the EMT switch leads to an unexpected model prediction for proliferating mesenchymal cells. Such temporary loss of *Pak1* is triggered by *Cdk1/Cyclin B* during mitosis, when cells loosen their contact with the ECM, loose their stress fibers and round up to assemble the mitotic spindle (**Fig. 2** *Migration* module in mesenchymal cells) [83]. The resulting *Pak1* inactivation does not impact the EMT switch directly, as both *miR-34* and *miR-200* remain repressed by *Zeb1* and *Snai1*. That said, in cells that complete mitosis in the *absence* of strong growth signals, loss of *NF-κB* and reactivation of *GSK3β* compounds the loss of *Pak1,* all tipping the scales towards the loss of active *Snai1*. Thus, our model predicts that the loss of mitogenic stimuli leading to cell cycle exit is accompanied by MET (**Fig. S2B**). A similar fate awaits hybrid E/M cells exiting the cell cycle (**Fig. S2C**). In contrast, cells that undergo partial EMT with the aid of a *short-lived* growth signal which does not push them into the cell cycle remain migratory and maintain a hybrid E/M state in low growth factor environments (**Fig. 2D**). This behavior explains the biphasic response to increasing growth factors; namely that moderate growth signals increase, but saturating signals *decrease* the stability of the hybrid E/M state (**Fig. S1C**).

The full phenotype repertoire of our model showcases the mechanosensitive routes to EMT and MET

To summarize the model's ability to reproduce the cell phenotypes we expect to see in different microenvironments, we visualized our model's phenotypes (attractors) across all input-combinations (**Fig. 3E**). To this end we created a coordinate system in which each axis represents an independent environmental factor [45,63,64]. Omitting all *Trail* = ON and *ECM* = OFF conditions where all cells eventually die allowed us to reduce the dimensionality of the environment space to three independent inputs: growth signals (*x* axis), cell density (*y* axis) and ECM stiffness (*z* axis). Distinct positions (coordinates) along the resulting 3D grid represent unique environment-combinations. At each unique coordinate we placed a small picture to represent a stable cell phenotype. For example, apoptosis (*blue apoptotic cell*) is the only stable state in environments with no growth factors – the leftmost position along the *x* axis representing

growth signals. In environments that also support non-apoptotic attractors, there is more than one phenotype-symbol (e.g., apoptotic cells and green epithelial cells in growth factors with soft ECM; **Fig. 3E** *bottom plane*). As **Fig. 3E** indicates, a mesenchymal phenotype in the absence of external or autocrine drivers of EMT is only stable on stiff ECM, at medium to low cell density, in the presence of strong mitogens (*red stars*). Weaker, non-proliferative growth stimuli still permit the maintenance of hybrid E/M, provided they can stretch on stiff ECM (moderate / low density, *orange triangles*). If they become isolated when exposed to strong growth signals, these hybrid E/M cells undergo full EMT to a mesenchymal state (*lone red star*). In contrast, both soft ECM and high cell density force cells into an epithelial state (*green boxes*), while loss of all growth signals leads to apoptosis (*light blue images*).

Our model makes intriguing predictions about the bi-stable mix of epithelial and hybrid E/M cells that coexist under low growth signaling. Namely, we predict that short spikes of growth signaling that fail to trigger cell cycle entry can push epithelial cells into hybrid E/M at a monolayer's edge. This hybrid E/M state is maintained, along with a migratory state, until interrupted by mitosis, soft ECM exposure, or high cell density. In contrast, transitions triggered by *sustained* growth signals trigger cell cycle entry and lead to reversible EMT (**Fig. 3E,** EMT: *red/orange arrows;* MET: *mustard arrows*).

Autocrine TGF-β signaling locks in EMT triggered by growth signaling on stiff ECM and blocks mesenchymal cells proliferation

Next, we set out to model the crosstalk between the biomechanical environment and autocrine *TGFβ* signaling required to sustain the mesenchymal state *in vitro* [25,26]. To this end we added a *TGFβ* signaling cascade, as well as a secreted *TGFβ* node to the EMT switch (**Figs. 4A, S3**). In this new module, externally applied or secreted *TGFβ* can bind to *TGFβ* receptors I and II, which in turn activate *Smad 2/3/4* [84]. *Smads* drive EMT by promoting nuclear translocation of *β-catenin,* and inducing *Snai2*, *Lef1*, and *HMGA2* expression. *HMGA2* further induces *Snai1, Snai2*, and *Twist*. In addition, we included *TGFβ*'s paradoxical effect on the MAPK cascade along with PI3K/AKT activation, anti-proliferative *p15* induction, and pro-apoptotic *BIM/BIK* induction and *BCL-X$_L$* repression (**Fig. 4A**; see **Table S1-2** for details). To create autocrine positive feedback, we included *TGFβ* secretion downstream of nuclear *β-catenin/TCF4* (repressed by *miR-200*).

Adding an autocrine loop to a Boolean regulatory model makes the often-unexamined assumption that a single cell is capable of secreting sufficient signal (in our case *TGF-β*) to drive a *saturating* response (i.e., pathway is ON rather than OFF). As a result, the effects of autocrine loops can be easily overestimated in Boolean models, indicating irreversible lock-in of EMT in situations where cells *in vitro* show metastable EMT due to partial autocrine activation [85]. To mitigate this, we included an additional input named *Self_Loop*. When set to 0, *TGFβ* secreted by an isolated cell is too weak to trigger signaling, whereas *Self_Loop=ON* represents saturating

levels of autocrine *TGFβ* activation. Tuning this input to intermediate values allows us to model a range of autocrine signal strengths in sparse cells. To further account for the possibility that cells with mesenchymal *TGFβ*-secreting neighbors have stronger autocrine signaling [86], our model allows secreted *TGFβ* to saturate signaling at cell densities of a monolayer's edge or above.

To test the effects of a potent autocrine loop, we re-ran the mechanosensitive experiments shown on **Fig. 2** with the extended model. Namely, we exposed isolated epithelial cells to stiff ECM and saturating growth factors (**Figs. 4B,C**). Cells exposed to growth factors on soft ECM remain epithelial (**Fig. 4B**, *left*) but undergo EMT when plated on a stiff matrix for a sufficient interval (*middle*) and sustain their mesenchymal state when replated on soft ECM (*rigth*). As they fully engage autocrine *TGFβ* signaling, they also cease to proliferate (**Fig. 4B**). Indeed, autocrine *TGFβ* is responsible for non-monotone proliferation response with increasing matrix stiffness, where maximum proliferation is predicted at the stiffness boundary at which EMT is triggered (**Fig. 4D,** *top 2 panels; ~55% Stiff_ECM*). Below this, soft ECM limit proliferation; above the threshold it is blocked by *TGF-β.* A similarly irreversible EMT is observed in cells plated on stiff matrix first, then exposed to high growth factors (**Fig. 4C**), except for the appearance of hybrid E/M cells in response to weak growth stimuli too weak to trigger proliferation (**Fig. 4D,** *bottom 2 panels*). This hybrid E/M is triggered by short bursts of growth factors that do not persist long enough to trigger cell cycle or lock in the autocrine signaling loop (**Fig. S4A,B**). In contrast, short-lived exposure to a stiff ECM either fails to trigger sustained EMT (**Fig. S4C**) or leads to *TGF-β*-mediated apoptosis (**Fig. S4D**) (see next section). Next, we weakened it from 1 to 0 in isolated cells exposed to an environment that favors but cannot fully force mechanosensitive EMT; namely 85% growth signals and 85% Stiff ECM. As **Fig. 4E** indicates, even a slight weakening of the autocrine loop decreases the time cells spend in a mesenchymal state, restoring proliferation instead. Its full knockout leads to a mix with majority epithelial vs. mesenchymal cells (*bottom panel*). In summary, our model shows that growth signals and stiff ECM work together to drive EMT in sparsely plated cells, a transition that is subsequently maintained by strong autocrine *TGFβ* signaling.

Figure 4. At very low density, stiff ECM and growth signaling trigger stable commitment to a mesenchymal state, sustained by strong autocrine *TGFβ* signaling. A) *TGFβ* signaling module (*red*) linked to the previous model's EMT switch (*yellow;* augmented with a node for secreted *TGFβ*), growth signaling (*light green:* MAPK; *dark green:* AKT) supported by *Src* activation (*dark blue*), cell cycle entry (*lilac*) and apoptosis (*dark red*). **B, C)** Dynamics of regulatory molecule expression/activity during exposure of (B) an isolated, growth-stimulated epithelial cell to stiff ECM, and (C) an isolated epithelial cell on stiff ECM to strong mitogens. *X-axis:* time-steps; *y-axis:* nodes organized in regulatory modules; *orange/blue:* ON/OFF; *black/white labels & arrows:* molecular changes that drive EMT. **D)** *Top two vs. bottom 2 panels:* response of isolated cells to *i)* increasing *Stiff ECM* exposure in the presence of 95% saturating growth stimuli (*top two*) vs. ii) increasing growth factor exposure on 95% stiff ECM (*bottom two*). *Proliferation label:* rate of normal cell cycle completion (*blue*) vs. G2 → G1 reset (*orange, not observed*), aberrant mitosis (*green*, not observed), or failed cytokinesis (*red*, not observed) followed by genome duplication, relative to the minimum cell cycle length (21 time-steps), shown as stacked bar charts. *EMT label:* fraction of time spent in a mesenchymal (*orange*) vs. epithelial (*blue*) state. **E)** Response of isolated cells on 85% Stiff ECM exposed to 85% saturating growth stimuli to increasing inhibition of autocrine signaling (*Self-Loop knockdown*). *Top:* rate of normal cell cycle completion (*blue*) vs. G2 → G1 reset (*orange*), aberrant mitosis (*green*, not observed), or failed cytokinesis (*red*, not observed) followed by genome duplication. *Bottom:* fraction of time spent in a mesenchymal (*orange*) vs. epithelial (*blue*) state. *Total sampled live cell time:* 500,000 steps; synchronous update; *Initial state for sampling runs:* isolated epithelial cell in low mitogens on a soft ECM.

In contrast to full EMT in sparse cells, in our model partial EMT at the edge of a monolayer does not result in *TGFβ* secretion. As a result, the model's behavior is the same as that of the previous one (**Figs. 2C-D** vs. **S5-S6**). Namely, matrix stiffness and growth signaling induce reversible hybrid E/M and proliferation in cells at the edge of a monolayer (**Figs. S5A,B**), but sustained migratory hybrid E/M in cells exposed to sub-proliferative bursts of growth signaling (**Fig. S5C**). We predict that moderate levels of growth factor exposure on stiff ECMs leads to the most time spent in a hybrid E/M state (**Fig. S6B**, *bottom*), as cells reset from a hybrid E/M to an epithelial state every time they round up for mitosis. While short bursts are necessary for partial EMT, sustained exposure triggers a conflict between proliferation and migration (hybrid E/M) and requires partial EMT after each division. Knockdown of *E-cadherin* breaks the tug of war by favoring full EMT at the expense of proliferation (**Fig. S6C**).

Next, we set out to text whether altering the biomechanical environment can trigger MET in the presence of strong autocrine *TGFβ* signaling. After verifying that mesenchymal cells remained stable at high cell density (**Fig. 5A**, *left*), we exposed them to soft ECM. Loss of both space and stiff ECM triggered MET by activating *Sprouty2*, which in turn repressed both *Smad* activity and MAPK signaling. Mesenchymal cells plated on soft ECM and then crowded to high density underwent a similar MET (**Fig. 5B**). At intermediate densities that only trigger intermittent *Sprouty2* activity, our model predicts a mix of epithelial, mesenchymal and hybrid E/M states (**Fig. 5C** *at ~10% High Cell Density*). As the density increases, mesenchymal cells shift to an epithelial - hybrid E/M mix generated by crowded cells undergoing full MET, followed by partial EMT whenever space opens (**Fig. S7**) – an event that disappears in confluent epithelium.

Figure 5. Soft ECM and high cell density work together to break the autocrine *TGFβ* loop and trigger MET via contact-induced *Sprouty-2* (*Spry2*) activation. A-B) Dynamics of regulatory molecule expression/activity during exposure of (A) a growth-stimulated mesenchymal cell on a stiff matrix but at very high density to a soft ECM, and (B) an isolated mesenchymal cell on soft ECM to high density. *X-axis:* time-steps; *y-axis:* nodes organized in regulatory modules; *orange/blue:* ON/OFF; *black/white labels & arrows:* molecular changes that drive MET. **C)** Response of mesenchymal cells at the edge of a monolayer to increasing cell density in the presence of 95% saturating growth stimuli on 95% stiff ECM. *Top:* rate of normal cell cycle completion (*blue*) vs. G2 → G1 reset (*orange*, not observed), aberrant mitosis (*green*, not observed), or failed cytokinesis (*red*, not observed) followed by genome duplication, relative to the minimum cell cycle length (21 time-steps), shown as stacked bar charts. *Middle:* fraction of time spent in a mesenchymal (*orange*) vs. epithelial (*blue*) state. *Bottom:* fraction of time spent in a hybrid E/M (*orange*) vs. epithelial (*blue*) state. *Total sampled live cell time:* 100,000 steps; synchronous update; *Initial state for sampling runs:* isolated epithelial cell in low mitogens on a soft ECM.

Modeling Anoikis Resistance: mesenchymal cells lose anchorage dependence

A concerning consequence of EMT in the context of metastatic cancer is loss of anchorage dependence, or resistance to anoikis in mesenchymal cells [57]. To test whether our model can

reproduce this, we simulated loss of anchorage with an epithelial cell in low growth factor conditions (**Fig. S8A**) vs. a mesenchymal cell sustained by autocrine *TGFβ* signaling in the complete absence of growth factors (**Fig. S8B**). As expected, the epithelial cell underwent apoptosis while the mesenchymal one did not (**Fig. S8A-C**). Overexpression of either *AKT* or *ERK* rescued epithelial cells from anoikis (**Fig. S8D**, *left*). In mesenchymal cells, however, even complete knockout of *AKT* was not sufficient to induce anoikis, as *TGF-β*-mediated ERK activity could maintain survival. In contrast, *ERK* knockdown could re-sensitize these cells to anoikis (**Fig. S8C**, *right*).

External TGFβ triggers EMT and cell cycle inhibition on stiff ECM, but apoptosis on soft substrates

TGFβ was shown promote EMT on stiff matrixes, but cause apoptosis on soft ECMs in the same epithelial culture [46]. To test whether our model can reproduce both outcomes, we simulated cells at the edge of a monolayer (medium density) on soft vs. stiff ECM and exposed them to external *TGFβ*. Indeed, our model epithelial cells underwent apoptosis on soft ECM, an event driven by a combination of *BIM* overexpression, $BCL\text{-}X_L$ downregulation, and critically, lack of both *ERK* and high *AKT* activity (**Fig. 6A**). The latter is caused by weak activation of both mitogenic pathways in the absence of strong focal adhesions and stress fibers [87,88]. In contrast, on stiff ECM *TGFβ* signaling activates both pathways, counteracts its own apoptotic effects, and triggers EMT instead (**Fig. 6B/C,** *low / high growth factor env.*). As cell density increases, the ability of *TGFβ* to trigger EMT weakens [59]. Indeed, our model predicts that fully confluent monolayers dense enough to forbid stress fiber formation and at least some *YAP* activity stop responding to *TGFβ,* even if its receptors are accessible to external *TGFβ* (**Fig. 6D**). In a microenvironment with saturating growth signals and 50% external *TGFβ,* most cells on soft ECM have an epithelial phenotype and are prone to apoptosis, but also to occasional cell cycle entry (**Fig. 6E**, proliferation rate of ~20% of saturating levels seen on stiff ECM with no *TGFβ*). As matrix stiffness increases, the rate of both proliferation and apoptosis drops, giving way to EMT.

Figure 6. Exposure to external *TGFβ* leads to apoptosis on soft ECM, but EMT and cell cycle arrest on stiff ECM. A-D) Dynamics of regulatory molecule expression/activity during exposure to *TGFβ* of (A) growth-stimulated epithelial cells on soft ECM (medium density), (B) epithelial cells in low growth conditions on stiff ECM (medium density), (C) cycling hybrid E/M cells on stiff ECM (medium density), and (D) growth-stimulated but fully confluent (dense) epithelium on stiff ECM. *X-axis:* time-steps; *y-axis:* nodes organized in regulatory modules; *orange/blue:* ON/OFF; *black/white labels & arrows:* molecular changes that drive apoptosis (A), EMT (B-C), or lack of response (D). **E)** Response growth-stimulated epithelial cells at the edge of a monolayer to 50% external *TGFβ* at increasing ECM stiffness. *Top:* rate of normal cell cycle completion (*blue*) vs. G2 → G1 reset (*orange,* rare but observed between 50-70% ECM stiffness), aberrant mitosis (*green*, not observed), or failed cytokinesis (*red*, not observed) followed by genome duplication, relative to the minimum cell cycle length (21 time-steps), shown as stacked bar charts. *Second from top:* rate of apoptosis (per minimum cell cycle length). *Third from top:* fraction of time spent in a mesenchymal (*orange*) vs. epithelial (*blue*) state. *Bottom:* fraction of time spent in a hybrid E/M (*orange*) vs. epithelial (*blue*) state. *Total sampled live cell time:* 100,000 steps; synchronous update; *Initial state for sampling runs:* dividing hybrid E/M at saturating mitogen exposure, at the edge of a monolayer on stiff ECM.

To test the effects of weakening autocrine *TGFβ* signaling, we started with cells exposed to near-saturating growth signals and scanned their responses as a function of increasing ECM stiffness (**Fig. S9A,** *y axis*) at densities increasing from isolated to that of a monolayer's edge (**Fig. S9A,** *x axis*). As **Fig. S9A** illustrates, even a 10% loss of autocrine signal strength in isolated mesenchymal cells (modeled as *Self_Loop* = OFF for a random 10% of time steps), restored proliferation on stiffer ECMs at *all* densities, rather than only in hybrid E/M cells capable of maintaining adherens junctions at a monolayer's edge (**Fig. S9A,** *top*). Intriguingly, our model predicts a small uptick of cell cycle errors such as slippage from G2 back to G1 and failure to complete cytokinesis following telophase, both more common among sparce cells (~ 2-3% of all divisions). An increase of apoptosis with the weakening of autocrine signaling is also predicted on stiff ECMs at very low density, at a slightly higher rate than cell cycle errors. Meanwhile, weakening the autocrine loop severely destabilized the mesenchymal state at very low density, triggering MET on all but the stiffest ECM (**Fig. S9A,** *bottom 3 panels*). Next, we repeated the above scan with cells on 95% stiff ECM, scanning their responses as a function of increasing mitogen exposure (**Fig. S9B,** *y axis*) at increasing densities (**Fig. S9B,** *x axis*). In this case a weaker loop increased the prevalence of cell cycle errors even more (**Fig. S9B,** *top panels*), but more importantly reduced the stability of mesenchymal states in favor of hybrid E/M.

To summarize our results, we recreated the map cell phenotypes our model creates in different microenvironment-combinations, accounting for the availability of external *TGFβ* as a 4th dimension (**Fig. 7**). In contrast to the model with no autocrine *TGFβ* signaling (**Fig. 3E**), the quiescent mesenchymal phenotype is stable in the absence of external *TGFβ* in all environments except on very soft ECMs at very high density – assuming a strong autocrine signaling loop (**Fig. 7,** *left panel, red stars*). In contrast, ongoing and saturating external *TGFβ* eliminates the hybrid E/M phenotype characteristic of monolayer edges, while the epithelial phenotype is only stable at very high density (**Fig. 7,** *right panel)*. Interestingly, our model predicts that cells at this high density do not undergo apoptosis in response to *TGFβ*; rather they use basal AKT activation by *TGFβ* as a survival signal even in the absence of all other mitogens, or in the absence of anchorage – though they cannot survive the absence of both (**Fig. 7,** *right panel, green boxes*).

Figure 7. Strong autocrine *TGFβ* signaling stabilizes the mesenchymal state and external *TGFβ* forces full EMT in all but dense monolayers on soft ECM. Summary of cell phenotypes predicted for every combination of no/low/high growth-factor (*x axis*), no/low/high cell density environments (*y axis*), and no/very soft/very stiff ECM (*z axis*), in the absence (*left*) and presence (*right*) of external *TGFβ*. *Blue fragmented cell:* apoptotic state; *green box*: quiescent epithelial cell; *orange triangle:* hybrid E/M cell, *red star:* mesenchymal cell; *symbol with blue nucleus:* non-dividing cell; *symbol with mitotic spindle:* proliferating cell. Image credits: apoptotic cell [82]; mitotic spindle: *https://en.wikipedia.org/wiki/ Cell_division#/media/File:Kinetochore.jpg*.

DISCUSSION

Modeling the signaling networks responsible for EMT has been an ongoing focus for the last decade [22,26,27,30,35–42,76,89,90]. Models range from small ODE-based circuits that examine the core feedback loops responsible for locking in EMT to large Boolean networks that focus on the extensive crosstalk between EMT-promoting biochemical signals. Population-level models, while often considering the 2-3D biophysical environment of individual cells, do not integrate the effects of mechanosensitive molecular pathways such as *Hippo*, *YAP*, *Rac1/PAK1* and *NF-κB* [37,91]. Thus the biomechanical influences that control partial vs. full EMT, with or without transforming signals, are not adequately linked to molecular mechanisms responsible for each outcome. To address this, we first built a 136-node Boolean regulatory network model of partial as well as full EMT, driven by a cell's biomechanical environment in the absence of transforming factors such as *TGF-β, Wnt, Notch, Hedgehog* or *IL6* (**Fig. 1C**). Instead, EMT in this model is triggered by mitogenic signals on stiff ECM, together with a lack of contact inhibition leading to migration. Namely, *Rac1/PAK1* activation and *PI3K/AKT*-mediated *NF-κB* signaling kickstart EMT by increasing the expression, localization, and stability of *SNAI1*. In line with most published models, our network can maintain distinct epithelial, hybrid E/M and mesenchymal states (**Figs. 1B, 3E**). Unlike other models, we also track the coordination between

EMT, migration, cell cycle progression and apoptosis. With the aid of mechanosensitive inputs such as low density, our model can reproduce EMT driven by strong mitogens such as EGF on stiff ECMs (**Figs. 2,4, S1B-C**) [52,53], along with the observations that loss of *E-cadherin* can promote EMT (**Fig. S1D, S6C**) [54,55].

Our extended 150-node model endowed with a TGFβ signaling module (**Fig. 4A, S3**) reproduces the need for a potent autocrine TGFβ loop to maintain the mesenchymal state on soft ECM (**Fig. 4B**), in the absence of strong mitogens (**Fig. 4C**), or at high density (**Fig. 5A,** *left*). While this autocrine loop is a common feature to most published EMT models, here we also reproduce TGFβ-induced apoptosis on soft ECMs vs. EMT on stiff ones (**Fig. 6A-B**) [46], TGFβ-induced inhibition of proliferation in mesenchymal cells (**Fig. 6C**) [56], the lack of TGFβ-induced EMT in very dense monolayers (**Fig. 6D**) [59], and mesenchymal anoikis resistance sustained by autocrine TGFβ (**Fig. S8**) [57,58]. This board repertoire of behaviors is a direct result of a modular modeling strategy by which we built the EMT transcriptional network as an extension of a model series that started with the detailed control of cell cycle progression [64], its coordination with apoptosis [63], then examined the effects of contact inhibition and loss of anchorage on both phenotypes [45].

In addition to reproducing known cell behaviors, our model offers several mechanistic, experimentally testable predictions. *1)* In our model, the density of adherens junctions around a single cell is a biomechanical signal that biases EMT towards hybrid E/M vs. a mesenchymal outcome. Thus, we predict that relatively isolated cells in sparsely plated epithelia exposed to *EGF* on a stiff matrix are more likely to undergo full EMT, whereas cells at the edge of a monolayer or with a similar contact-surface stop at hybrid E/M instead (**Fig. 2**). *2)* We further predict that short, intermittent mitogen bursts that lack the strength to induce cell cycle entry can drive sparsely plated epithelial cells into a metastable migratory hybrid E/M state, which is maintained by uninterrupted migration (**Figs. 2D, S4A, S5C**). As a corollary, we predict a biphasic response to growth signal exposure where moderate signaling results in the largest fraction of hybrid E/M cells (**Fig. S6B**), while saturating mitogens interrupt migration and reset cells at each mitotic rounding (**Figs. S4B, S5B**). *3)* We predict that the biomechanical environment alone can trigger MET by blocking stress fiber formation and muting *ERK/AKT* signaling, while *Pak1* hyperactivation can re-establish hybrid E/M (**Fig. 3**). Furthermore, soft ECM and high cell density cooperatively break the autocrine *TGFβ* loop and trigger MET via contact induced *Sprouty 2* activation (**Fig. 5**). *4)* In line with previous model predictions and observations about the asymmetry of EMT vs. MET [27,92], our model does not undergo a direct transitions from mesenchymal to hybrid E/M state. Rather, model environments that favor hybrid E/M over the mesenchymal state induce full MET followed by partial EMT (**Fig. S7**). *5)* While TGFβ-induced EMT requires stiff ECM, we predict that the mesenchymal state can be maintained on soft ECMs by autocrine TGFβ (**Fig. 4B**). *6)* Finally, we predict that at low-to-medium cell density and medium-strength growth signals, autocrine TGFβ signaling that does not saturate the TGFβ pathway (does not reliably guarantee its downstream effects) will generate a dynamic mix of mesenchymal and hybrid E/M cells, the latter of which proliferate and thus

continually reset to undergo EMT anew (**Figs. S9**). This is accompanied by a small uptick in apoptosis as well as cell cycle errors that generate polyploidy (specifically, slippage from G2 and/or telophase to G1) in the absence of any mutations related to checkpoint control (**Figs. S9**).

Rather than our ability to reproduce known cell behaviors, we believe that the above predictions represent a more important contribution to characterizing the roles of partial vs. full EMT in non-oncogenic processes such as wound healing. Testing them could boost the credibility of our model in predicting the effects of targeted interventions. Alternately, failed predictions would point to unexpected gaps, helping us further probe the behavior of cells in different biomechanical environments. Most of our predictions can be tested by single-cell imaging experiments that carefully control ECM stiffness, cell density, and mitogen exposure while monitoring the activity of core EMT regulators such as *Snai1*, *Zeb1, E-cadherin*, or the two microRNAs. Ultimately, correct integration of mechanosensing and transforming signals will be required for building predictive cell population models that follow the 2D or 3D processes of wound healing, the initiation of cancer invasion, or the establishment and growth of metastatic lesions.

Our models have several limitations that can be addressed by future refinement; some related to the way our previous contact inhibition model's environment was built, some specific to EMT and TGFβ signaling. As detailed in [45], we use a crude approximation of the spatial asymmetry of *Rac1/Pak1* activity at the leading edge and a minimal migration module. Yet, feedback between cell contractility and ECM stiffness – itself modulated by the migrating cell – was shown to render the behavior of invading cells nonlinear, such that intermediate stiffness is optimal for high-speed migration [93]. Moreover, hybrid E/M cells use their lower contractility to move faster and invade more effectively than mesenchymal cells [94]. The migration module of our model needs significant refinement to address these phenomena before it is suited to model collective behavior such as wound healing in 2D. Other areas of improvement inherited from [45] hinge on the limits of how well Boolean models capture gradual changes in ECM stiffness and cell density (detailed in Supplementary Methods and [45]).

A potential limitation related to our integration of the EMT switch with the cell cycle is that our model's mesenchymal cells do not show anchorage-independent growth [7,95]. While we reproduce anoikis resistance, our cell cycle control relies entirely on spreading on a stiff matrix. A deeper examination of how cancer cells acquire anchorage-independent growth, and whether there are pre-requisite mutations that could drive such growth in our current model, requires further work. A related potential problem is that our mesenchymal cells with strong autocrine TGFβ loop do not divide at all, owing to a TGFβ-mediated cell cycle block. Instead, in environments that do not force full EMT (sub-optimal matrix stiffness, non-saturating autocrine TGFβ signaling, moderate density, non-saturating growth signals) our model predicts a fluid coexistence of mesenchymal and hybrid E/M cells, the latter of which divide with high frequency but then briefly reset to an epithelial state before undergoing EMT once more. It is likely that our model's

TGFβ-mediated cell cycle block is too strong, in that fully mesenchymal cells with sufficiently potent autocrine signals to keep them mesenchymal may nevertheless divide. Yet there is substantial experimental evidence for the tug of war between proliferative potential and the migratory mesenchymal state [7]. For example, oval cells capable of regenerating the liver when hepatocyte proliferation is compromised where shown to have a hybrid E/M phenotype [96]. Moreover, cancer stem cells in squamous cell carcinoma and ovarian cancer exist as a heterogeneous mix of proliferative hybrid E/M and mesenchymal states, and they dynamically regenerate this mix from either subpopulation [97,98]. Indeed, *in vivo* metastatic tumor growth in ovarian and breast cancer is primarily due to hybrid E/M cells [101,102]. We can model this division/migration conflict in environments that can induce and maintain EMT biomechanically, and thus do not rely on uninterrupted autocrine TGFβ signaling (**Fig. S9**). What our model currently does not account for, however, is the connection between the hybrid E/M state and the regulatory network responsible for a stem cell-like state, observed in all forms of EMT (i.e., developmental, regenerative, and pathological) [30,99]. Integrating a stemness module, along with an explicit accounting for the likely context-dependent role of 'phenotypic stability factors' such as *OVOL* and *GRHL2* known to stabilize the hybrid E/M state and promote stemness, would be an important next step in delineating the similarities and differences between EMT in different tissue and disease contexts [30,100].

In addition to the limitations described above, it is important to delineate the boundaries of our model; cell behaviors linked to EMT that we explicitly do not address. Two of these stand out as key to our ability to integrate our regulatory network into multiscale models of tissue behavior. *First,* our model does not address the patterning effects of *Notch* signaling in a population of cells undergoing EMT. *Notch* not only promotes EMT by inducing *Snai1* [39], but also generates distinct patterns between neighboring cells depending on the ligands expressed on neighbors [37]. Cells expressing *Delta* ligands promote lateral inhibition patterns that alternate $Delta^{High}$/$Notch^{Low}$ cells with the opposite phenotype. This is central to the endothelial to mesenchymal transition, as it generates alternating patterns of tip cells and stalk cells required for sprouting angiogenesis [76,101]. In contrast, the *Jagged* ligand promotes lateral induction, leading to a strong clustering of similar cells [102]. Previous computational models of *Notch* signaling showed that lateral induction via *Notch-Jagged* can generate intermingled clusters of epithelial, mesenchymal, and hybrid E/M cells akin to the patterning seen in tumors [37]. This pattern formation is heavily influenced by the tissue microenvironment such as inflammatory signals (*IL-6*, *TNFα*) [103,104], *Wnt* and *Hedgehog* signaling [105,106], as well as hypoxia [107], most of which promote EMT and induce *Jagged*.

The second direction that merits our attention is the relationship between EMT and senescence. The process of senescence makes short-term use of damaged cells by shutting down their ability to divide, but using them to boost survival, proliferation and EMT in their surroundings [108]. While senescent cells trigger their own immune clearance, their slow accumulation is accelerated by chronic damage such as inflammation or oxidative stress and leads to tissue aging [109]. There is

increasing evidence that senescence and the mesenchymal state are mutually exclusive in individual cells [110,111]. Mesenchymal cells are largely protected from senescence, while senescent cells do not respond to signals that trigger EMT [112]. During localized healing of an epithelium, this creates a useful distribution of labor: damaged cells go senescent but induce division and EMT in neighbors. These mesenchymal cells are then protected from senescence until they reestablish an intact epithelium. As neither senescent nor mesenchymal cells contribute to barrier formation, this can accelerate dysfunction in aging tissues [113]. Worse, in epithelial tumors this crosstalk protects invasive but damaged cells from senescence, aiding metastasis [114]. Yet, the two phenomena are usually studied in isolation, and the relationship between hybrid E/M cell states and senescence remains unexamined. Building a predictive molecular model of the mechanisms that forbid their coexistence and probe the transition from healing to the pathological roles of EMT and senescence is a key next step that benefits from our current integration of EMT with cell cycle control, apoptosis, and contact inhibition. Ultimately, we envision these single-cell network models as the engines that drive 3D tissue simulations, able to predict effective interventions that can nudge diseased tissues back to health.

ACKGNOWLEDGEMENTS

All funding for this study was provided by the College of Wooster. E.R. would like to thank the Henry Luce III Fund for Distinguished Scholarship. E.G and I.Z thank the Henry J. Copeland Independent Study Fund for research support. A.B. thanks APEX (Advising, Planning, & Experiential Learning) at the College of Wooster for support to present their work.

AUTHOR CONTRIBUTIONS

E.S. and E.G. designed the first version of our Boolean EMT switch and started its integration with the model in [45], leading to the first mechanosensitive version. M.H., A.B. and I.Z. tested and refined the model to converge in the version included here. E.R.R. supervised the research, analyzed/refined the final models, ran final simulations, and wrote the paper with edits from E.S.

DECLARATION OF INTERESTS

All authors were employees (E.R.R.) or undergraduate students (E.S., M.H., E.G., A.B., I.Z.) at the College of Wooster at the time of this study.

REFERENCES

1. Abercrombie, M. Contact inhibition and malignancy. *Nature* **281**, 259–262 (1979).
2. Eagle, H. & Levine, E. M. Growth regulatory effects of cellular interaction. *Nature* **213**, 1102–1106 (1967).

3. Kim, D. H. *et al.* Epithelial Mesenchymal Transition in Embryonic Development, Tissue Repair and Cancer: A Comprehensive Overview. *J Clin Med* **7**, (2017).
4. Lamouille, S., Xu, J. & Derynck, R. Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol* **15**, 178–196 (2014).
5. Dongre, A. & Weinberg, R. A. New insights into the mechanisms of epithelial-mesenchymal transition and implications for cancer. *Nat. Rev. Mol. Cell Biol.* **20**, 69–84 (2019).
6. Thiery, J. P., Acloque, H., Huang, R. Y. J. & Nieto, M. A. Epithelial-mesenchymal transitions in development and disease. *Cell* **139**, 871–890 (2009).
7. Jolly, M. K., Ware, K. E., Gilja, S., Somarelli, J. A. & Levine, H. EMT and MET: necessary or permissive for metastasis? *Mol Oncol* **11**, 755–769 (2017).
8. Dillekås, H., Rogers, M. S. & Straume, O. Are 90% of deaths from cancer caused by metastases? *Cancer Med* **8**, 5574–5576 (2019).
9. Hong, D. *et al.* Epithelial-to-mesenchymal transition and cancer stem cells contribute to breast cancer heterogeneity. *J Cell Physiol* **233**, 9136–9144 (2018).
10. Kim, J. E., Leung, E., Baguley, B. C. & Finlay, G. J. Heterogeneity of expression of epithelial-mesenchymal transition markers in melanocytes and melanoma cell lines. *Front Genet* **4**, 97 (2013).
11. Pereira, L., Mariadason, J. M., Hannan, R. D. & Dhillon, A. S. Implications of epithelial-mesenchymal plasticity for heterogeneity in colorectal cancer. *Front Oncol* **5**, 13 (2015).
12. Stylianou, N. *et al.* A molecular portrait of epithelial-mesenchymal plasticity in prostate cancer associated with clinical outcome. *Oncogene* **38**, 913–934 (2019).
13. Campbell, K. & Casanova, J. A common framework for EMT and collective cell migration. *Development* **143**, 4291–4300 (2016).
14. Aiello, N. M. *et al.* EMT Subtype Influences Epithelial Plasticity and Mode of Cell Migration. *Developmental Cell* **45**, 681-695.e4 (2018).
15. Aceto, N. *et al.* Circulating tumor cell clusters are oligoclonal precursors of breast cancer metastasis. *Cell* **158**, 1110–1122 (2014).
16. Fidler, I. J. The relationship of embolic homogeneity, number, size and viability to the incidence of experimental metastasis. *Eur J Cancer* **9**, 223–227 (1973).
17. Liotta, L. A., Saidel, M. G. & Kleinerman, J. The significance of hematogenous tumor cell clumps in the metastatic process. *Cancer Res* **36**, 889–894 (1976).
18. Au, S. H. *et al.* Clusters of circulating tumor cells traverse capillary-sized vessels. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 4947–4952 (2016).
19. Chang, C.-J. *et al.* p53 regulates epithelial-mesenchymal transition and stem cell properties through modulating miRNAs. *Nat Cell Biol* **13**, 317–323 (2011).
20. Liu, X. *et al.* Roles of Signaling Pathways in the Epithelial-Mesenchymal Transition in Cancer. *Asian Pac. J. Cancer Prev.* **16**, 6201–6206 (2015).
21. Nieszporek, A., Skrzypek, K., Adamek, G. & Majka, M. Molecular mechanisms of epithelial to mesenchymal transition in tumor metastasis. *Acta Biochim Pol* (2019) doi:10.18388/abp.2019_2899.
22. Tripathi, S., Levine, H. & Jolly, M. K. The Physics of Cellular Decision Making During Epithelial–Mesenchymal Transition. *Annu. Rev. Biophys.* **49**, 1–18 (2020).
23. Zhang, J. *et al.* TGF-β-induced epithelial-to-mesenchymal transition proceeds through stepwise activation of multiple feedback loops. *Sci Signal* **7**, ra91 (2014).
24. Peinado, H., Portillo, F. & Cano, A. Transcriptional regulation of cadherins during development and carcinogenesis. *Int. J. Dev. Biol.* **48**, 365–375 (2004).

25. Gregory, P. A. *et al.* An autocrine TGF-beta/ZEB/miR-200 signaling network regulates establishment and maintenance of epithelial-mesenchymal transition. *Mol Biol Cell* **22**, 1686–1698 (2011).

26. Tian, X.-J., Zhang, H. & Xing, J. Coupled reversible and irreversible bistable switches underlying TGFβ-induced epithelial to mesenchymal transition. *Biophys. J.* **105**, 1079–1089 (2013).

27. Lu, M., Jolly, M. K., Levine, H., Onuchic, J. N. & Ben-Jacob, E. MicroRNA-based regulation of epithelial-hybrid-mesenchymal fate determination. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 18144–18149 (2013).

28. Grosse-Wilde, A. *et al.* Stemness of the hybrid Epithelial/Mesenchymal State in Breast Cancer and Its Association with Poor Survival. *PLoS One* **10**, e0126522 (2015).

29. Huang, R. Y.-J. *et al.* An EMT spectrum defines an anoikis-resistant and spheroidogenic intermediate mesenchymal state that is sensitive to e-cadherin restoration by a src-kinase inhibitor, saracatinib (AZD0530). *Cell Death Dis* **4**, e915 (2013).

30. Jolly, M. K. *et al.* Stability of the hybrid epithelial/mesenchymal phenotype. *Oncotarget* **7**, 27067–27084 (2016).

31. Dong, J. *et al.* Single-cell RNA-seq analysis unveils a prevalent epithelial/mesenchymal hybrid state during mouse organogenesis. *Genome Biol* **19**, 31 (2018).

32. Gonzalez, V. D. *et al.* Commonly Occurring Cell Subsets in High-Grade Serous Ovarian Tumors Identified by Single-Cell Mass Cytometry. *Cell Rep* **22**, 1875–1888 (2018).

33. Karacosta, L. G. *et al.* Mapping lung cancer epithelial-mesenchymal transition states and trajectories with single-cell resolution. *Nat Commun* **10**, 5587 (2019).

34. Puram, S. V. *et al.* Single-Cell Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and Neck Cancer. *Cell* **171**, 1611-1624.e24 (2017).

35. Biswas, K., Jolly, M. K. & Ghosh, A. Stability and mean residence times for hybrid epithelial/mesenchymal phenotype. *Phys Biol* **16**, 025003 (2019).

36. Hong, T. *et al.* An Ovol2-Zeb1 Mutual Inhibitory Circuit Governs Bidirectional and Multi-step Transition between Epithelial and Mesenchymal States. *PLoS Comput Biol* **11**, e1004569 (2015).

37. Bocci, F. *et al.* Toward understanding cancer stem cell heterogeneity in the tumor microenvironment. *Proc Natl Acad Sci U S A* **116**, 148–157 (2019).

38. Tripathi, S., Chakraborty, P., Levine, H. & Jolly, M. K. A mechanism for epithelial-mesenchymal heterogeneity in a population of cancer cells. *PLoS Comput Biol* **16**, e1007619 (2020).

39. Steinway, S. N. *et al.* Network modeling of TGFβ signaling in hepatocellular carcinoma epithelial-to-mesenchymal transition reveals joint sonic hedgehog and Wnt pathway activation. *Cancer Res.* **74**, 5963–5977 (2014).

40. Gómez Tejeda Zañudo, J. *et al.* Towards control of cellular decision-making networks in the epithelial-to-mesenchymal transition. *Phys Biol* **16**, 031002 (2019).

41. Steinway, S. N. *et al.* Combinatorial interventions inhibit TGFβ-driven epithelial-to-mesenchymal transition and support hybrid cellular phenotypes. *NPJ Syst Biol Appl* **1**, 15014 (2015).

42. Font-Clos, F., Zapperi, S. & La Porta, C. A. M. Topography of epithelial-mesenchymal plasticity. *Proc Natl Acad Sci U S A* **115**, 5902–5907 (2018).

43. Thomas, R. Boolean formalization of genetic control circuits. *J Theor Biol* **42**, 563–585 (1973).

44. De Pascalis, C. & Etienne-Manneville, S. Single and collective cell migration: the mechanics of adhesions. *MBoC* **28**, 1833–1846 (2017).
45. Guberman, E., Sherief, H. & Regan, E. R. Boolean model of anchorage dependence and contact inhibition points to coordinated inhibition but semi-independent induction of proliferation and migration. *Comput Struct Biotechnol J* **18**, 2145–2165 (2020).
46. Tilghman, R. W. *et al.* Matrix rigidity regulates cancer cell growth and cellular phenotype. *PLoS ONE* **5**, e12905 (2010).
47. Park, J. *et al.* Switch-like enhancement of epithelial-mesenchymal transition by YAP through feedback regulation of WT1 and Rho-family GTPases. *Nat Commun* **10**, 2797 (2019).
48. Matte, B. F. *et al.* Matrix stiffness mechanically conditions EMT and migratory behavior of oral squamous cell carcinoma. *J. Cell. Sci.* **132**, (2019).
49. Rice, A. J. *et al.* Matrix stiffness induces epithelial-mesenchymal transition and promotes chemoresistance in pancreatic cancer cells. *Oncogenesis* **6**, e352 (2017).
50. Wei, S. C. *et al.* Matrix stiffness drives epithelial-mesenchymal transition and tumour metastasis through a TWIST1-G3BP2 mechanotransduction pathway. *Nat Cell Biol* **17**, 678–688 (2015).
51. Garcia, M. A. *et al.* *Epithelial-Mesenchymal Plasticity is regulated by inflammatory signalling networks coupled to cell morphology.* http://biorxiv.org/lookup/doi/10.1101/689737 (2019) doi:10.1101/689737.
52. Grasset, E. M. *et al.* Matrix Stiffening and EGFR Cooperate to Promote the Collective Invasion of Cancer Cells. *Cancer Res* **78**, 5229–5242 (2018).
53. Kim, J., Kong, J., Chang, H., Kim, H. & Kim, A. EGF induces epithelial-mesenchymal transition through phospho-Smad2/3-Snail signaling pathway in breast cancer cells. *Oncotarget* **7**, 85021–85032 (2016).
54. Aban, C. E. *et al.* Downregulation of E-cadherin in pluripotent stem cells triggers partial EMT. *Sci Rep* **11**, 2048 (2021).
55. Onder, T. T. *et al.* Loss of E-Cadherin Promotes Metastasis via Multiple Downstream Transcriptional Pathways. *Cancer Research* **68**, 3645–3654 (2008).
56. Siegel, P. M. & Massagué, J. Cytostatic and apoptotic actions of TGF-beta in homeostasis and cancer. *Nat Rev Cancer* **3**, 807–821 (2003).
57. Paoli, P., Giannoni, E. & Chiarugi, P. Anoikis molecular pathways and its role in cancer progression. *Biochim. Biophys. Acta* **1833**, 3481–3498 (2013).
58. Muñoz, N. M., Baek, J. Y. & Grady, W. M. TGF-β has paradoxical and context dependent effects on proliferation and anoikis in human colorectal cancer cell lines. *Growth Factors* **26**, 254–262 (2008).
59. Nallet-Staub, F. *et al.* Cell density sensing alters TGF-β signaling in a cell-type-specific manner, independent from Hippo pathway activation. *Dev Cell* **32**, 640–651 (2015).
60. Wang, R.-S., Saadatpour, A. & Albert, R. Boolean modeling in systems biology: an overview of methodology and applications. *Physical biology* **9**, 055001 (2012).
61. Albert, I., Thakar, J., Li, S., Zhang, R. & Albert, R. Boolean network simulations for life scientists. *Source Code Biol Med* **3**, 16 (2008).
62. Bergmann, F. T., Keating, S. M., Gauges, R., Sahle, S. & Wengler, K. SBML Level 3 package: Render, Version 1, Release 1. *J Integr Bioinform* **15**, (2018).
63. Sizek, H. Modeling the Commitment to Senescence Through Mitochondrial Fusion and Chromosomal Restructuring. *Senior Independent Study Theses* (2017).

64. Deritei, D., Aird, W. C., Ercsey-Ravasz, M. & Regan, E. R. Principles of dynamical modularity in biological regulatory networks. *Sci Rep* **6**, 21957 (2016).

65. Siemens, H. *et al.* miR-34 and SNAIL form a double-negative feedback loop to regulate epithelial-mesenchymal transitions. *Cell Cycle* **10**, 4256–4271 (2011).

66. Title, A. C. *et al.* Genetic dissection of the miR-200-Zeb1 axis reveals its importance in tumor differentiation and invasion. *Nat Commun* **9**, 4671 (2018).

67. Zhang, P. *et al.* ATM-mediated stabilization of ZEB1 promotes DNA damage response and radioresistance through CHK1. *Nat Cell Biol* **16**, 864–875 (2014).

68. Kim, N. H. *et al.* Competing Endogenous RNA of Snail and Zeb1 UTR in Therapeutic Resistance of Colorectal Cancer. *Int J Mol Sci* **22**, 9589 (2021).

69. Sánchez-Tilló, E. *et al.* β-catenin/TCF4 complex induces the epithelial-to-mesenchymal transition (EMT)-activator ZEB1 to regulate tumor invasiveness. *Proc Natl Acad Sci U S A* **108**, 19204–19209 (2011).

70. Kang, C. MicroRNA-200a suppresses the Wnt/β-catenin signaling pathway by interacting with β-catenin. *Int J Oncol* (2011) doi:10.3892/ijo.2011.1322.

71. Shirakihara, T., Saitoh, M. & Miyazono, K. Differential regulation of epithelial and mesenchymal markers by deltaEF1 proteins in epithelial mesenchymal transition induced by TGF-beta. *Mol Biol Cell* **18**, 3533–3544 (2007).

72. Vesuna, F., van Diest, P., Chen, J. H. & Raman, V. Twist is a transcriptional repressor of E-cadherin gene expression in breast cancer. *Biochem Biophys Res Commun* **367**, 235–241 (2008).

73. Bolós, V. *et al.* The transcription factor Slug represses E-cadherin expression and induces epithelial to mesenchymal transitions: a comparison with Snail and E47 repressors. *J Cell Sci* **116**, 499–511 (2003).

74. Jia, D. *et al.* Distinguishing mechanisms underlying EMT tristability. *Cancer Converg* **1**, 2 (2017).

75. Kröger, C. *et al.* Acquisition of a hybrid E/M state is essential for tumorigenicity of basal breast cancer cells. *Proc Natl Acad Sci U S A* **116**, 7353–7362 (2019).

76. Weinstein, N., Mendoza, L. & Álvarez-Buylla, E. R. A Computational Model of the Endothelial to Mesenchymal Transition. *Front Genet* **11**, 40 (2020).

77. Lin, Z. *et al.* Pi3kcb links Hippo-YAP and PI3K-AKT signaling pathways to promote cardiomyocyte proliferation and survival. *Circ. Res.* **116**, 35–45 (2015).

78. Manning, B. D. & Toker, A. AKT/PKB Signaling: Navigating the Network. *Cell* **169**, 381–405 (2017).

79. Zheng, H. *et al.* Glycogen synthase kinase-3 beta regulates Snail and β-catenin expression during Fas-induced epithelial-mesenchymal transition in gastrointestinal cancer. *Eur J Cancer* **49**, 2734–2746 (2013).

80. Min, C., Eddy, S. F., Sherr, D. H. & Sonenshein, G. E. NF-κB and epithelial to mesenchymal transition of cancer. *Journal of Cellular Biochemistry* **104**, 733–744 (2008).

81. Yang, Z. *et al.* Pak1 phosphorylation of snail, a master regulator of epithelial-to-mesenchyme transition, modulates snail's subcellular localization and functions. *Cancer Res.* **65**, 3179–3184 (2005).

82. Wells, R. G. The role of matrix stiffness in regulating cell behavior. *Hepatology* **47**, 1394–1400 (2008).

83. Jones, M. C., Askari, J. A., Humphries, J. D. & Humphries, M. J. Cell adhesion is regulated by CDK1 during the cell cycle. *Journal of Cell Biology* **217**, 3203–3218 (2018).

84. Seoane, J. & Gomis, R. R. TGF-β Family Signaling in Tumor Suppression and Cancer Progression. *Cold Spring Harb Perspect Biol* **9**, a022277 (2017).

85. Celià-Terrassa, T. *et al.* Hysteresis control of epithelial-mesenchymal transition dynamics conveys a distinct program with enhanced metastatic ability. *Nat Commun* **9**, 5005 (2018).

86. Doğaner, B. A., Yan, L. K. Q. & Youk, H. Autocrine Signaling and Quorum Sensing: Extreme Ends of a Common Spectrum. *Trends in Cell Biology* **26**, 262–271 (2016).

87. Yin, G., Zheng, Q., Yan, C. & Berk, B. C. GIT1 is a scaffold for ERK1/2 activation in focal adhesions. *J Biol Chem* **280**, 27705–27712 (2005).

88. Schrader, J. *et al.* Matrix stiffness modulates proliferation, chemotherapeutic response, and dormancy in hepatocellular carcinoma cells. *Hepatology* **53**, 1192–1205 (2011).

89. Shin, S.-Y. *et al.* Functional roles of multiple feedback loops in extracellular signal-regulated kinase and Wnt signaling pathways that regulate epithelial-mesenchymal transition. *Cancer Res* **70**, 6715–6724 (2010).

90. Gould, R., Bassen, D. M., Chakrabarti, A., Varner, J. D. & Butcher, J. Population Heterogeneity in the Epithelial to Mesenchymal Transition Is Controlled by NFAT and Phosphorylated Sp1. *PLoS Comput Biol* **12**, e1005251 (2016).

91. Neagu, A. *et al.* Computational modeling of epithelial-mesenchymal transformations. *Biosystems* **100**, 23–30 (2010).

92. Samavarchi-Tehrani, P. *et al.* Functional genomics reveals a BMP-driven mesenchymal-to-epithelial transition in the initiation of somatic cell reprogramming. *Cell Stem Cell* **7**, 64–77 (2010).

93. Ahmadzadeh, H. *et al.* Modeling the two-way feedback between contractility and matrix realignment reveals a nonlinear mode of cancer cell invasion. *Proc Natl Acad Sci U S A* **114**, E1617–E1626 (2017).

94. Margaron, Y. *et al. Biophysical properties of intermediate states of EMT outperform both epithelial and mesenchymal states.* http://biorxiv.org/lookup/doi/10.1101/797654 (2019) doi:10.1101/797654.

95. Mori, S. *et al.* Anchorage-independent cell growth signature identifies tumors with metastatic potential. *Oncogene* **28**, 2796–2805 (2009).

96. Yovchev, M. I. *et al.* Identification of adult hepatic progenitor cells capable of repopulating injured rat liver. *Hepatology* **47**, 636–647 (2008).

97. Biddle, A. *et al.* Cancer Stem Cells in Squamous Cell Carcinoma Switch between Two Distinct Phenotypes That Are Preferentially Migratory or Proliferative. *Cancer Research* **71**, 5317–5326 (2011).

98. Strauss, R. *et al.* Analysis of epithelial and mesenchymal markers in ovarian cancer reveals phenotypic heterogeneity and plasticity. *PLoS One* **6**, e16186 (2011).

99. Carmona-Gutierrez, D. *et al.* Guidelines and recommendations on yeast cell death nomenclature. *Microb Cell* **5**, 4–31.

100. Jolly, M. K. *et al.* Coupling the modules of EMT and stemness: A tunable 'stemness window' model. *Oncotarget* **6**, 25161–25174 (2015).

101. Bischoff, J. Endothelial-to-Mesenchymal Transition. *Circ Res* **124**, 1163–1165 (2019).

102. Bocci, F., Onuchic, J. N. & Jolly, M. K. Understanding the Principles of Pattern Formation Driven by Notch Signaling by Integrating Experiments and Theoretical Models. *Front. Physiol.* **11**, 929 (2020).

103. Abaurrea, A., Araujo, A. M. & Caffarel, M. M. The Role of the IL-6 Cytokine Family in Epithelial-Mesenchymal Plasticity in Cancer Progression. *Int J Mol Sci* **22**, 8334 (2021).

104.   Laha, D., Grant, R., Mishra, P. & Nilubol, N. The Role of Tumor Necrosis Factor in Manipulating the Immunological Response of Tumor Microenvironment. *Front. Immunol.* **12**, 656908 (2021).

105.   Basu, S., Cheriyamundath, S. & Ben-Ze'ev, A. Cell-cell adhesion: linking Wnt/β-catenin signaling with partial EMT and stemness traits in tumorigenesis. *F1000Res* **7**, F1000 Faculty Rev-1488 (2018).

106.   Katoh, Y. & Katoh, M. Hedgehog signaling, epithelial-to-mesenchymal transition and miRNA (review). *Int J Mol Med* **22**, 271–275 (2008).

107.   Saxena & Jolly. Acute vs. Chronic vs. Cyclic Hypoxia: Their Differential Dynamics, Molecular Mechanisms, and Effects on Tumor Progression. *Biomolecules* **9**, 339 (2019).

108.   Wilkinson, H. N. & Hardman, M. J. Senescence in Wound Repair: Emerging Strategies to Target Chronic Healing Wounds. *Front Cell Dev Biol* **8**, 773 (2020).

109.   Di Micco, R., Krizhanovsky, V., Baker, D. & d'Adda di Fagagna, F. Cellular senescence in ageing: from mechanisms to therapeutic opportunities. *Nat Rev Mol Cell Biol* **22**, 75–95 (2021).

110.   Smit, M. A. & Peeper, D. S. Epithelial-mesenchymal transition and senescence: two cancer-related processes are crossing paths. *Aging (Albany NY)* **2**, 735–741 (2010).

111.   Liu, Y., El-Naggar, S., Darling, D. S., Higashi, Y. & Dean, D. C. Zeb1 links epithelial-mesenchymal transition and cellular senescence. *Development* **135**, 579–588 (2008).

112.   Burk, U. *et al.* A reciprocal repression between ZEB1 and members of the miR-200 family promotes EMT and invasion in cancer cells. *EMBO Rep* **9**, 582–589 (2008).

113.   Krouwer, V. J. D., Hekking, L. H. P., Langelaar-Makkinje, M., Regan-Klapisz, E. & Post, J. A. Endothelial cell senescence is associated with disrupted cell-cell junctions and increased monolayer permeability. *Vasc Cell* **4**, 12 (2012).

114.   Schosserer, M., Grillari, J. & Breitenbach, M. The Dual Role of Cellular Senescence in Developing Tumors and Their Response to Cancer Therapy. *Front Oncol* **7**, 278 (2017).