Bacterial Microbiome of Foreskin Tissue, Impact of Short-course PrEP, and Relation to Host Inflammation

Brandon S. Maust[a,b], Stefan Petkov[c], Carolina Herrera[d], Colin Feng[a], Bryan P. Brown[a,b],

Limakatso Lebina[e], Daniel Opoka[f,g], Andrew Ssemata[f,g], Natasha Pillay[e], Jennifer Serwanga[f,g],

Portia Seatlholo[e], Patricia Namubiru[f,g], Geoffrey Odoch[f,g], Susan Mugaba[f,g], Thabiso Seiphetlo[h],

Clive M. Gray[h,i], Pontiano Kaleebu[f,g], Emily L. Webb[j], Neil Martinson[e], Francesca Chiodi[c*], Julie

Fox[k*], Heather B. Jaspan[a,b,h*#,] on behalf of the CHAPS team

[a] Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, WA, USA

[b] Division of Infectious Disease, Department of Pediatrics, University of Washington School of Medicine, Seattle, WA, USA

[c] Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden

[d] Department of Infectious Disease, Faculty of Medicine, Imperial College London, London, U.K.

[e] Perinatal HIV Research Unit, University of the Witwatersrand, Johannesburg, South Africa

[f] Medical Research Council/Uganda Virus Research Institute, Entebbe, Uganda

[g] London School of Hygiene & Tropical Medicine, Uganda Research Unit, Entebbe, Uganda

[h] Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa

[i] Division of Molecular Biology and Human Genetics, Department of Biomedical Sciences, Stellenbosch University, Cape Town, South Africa

[j] Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, U.K.

[k] Faculty of Life Sciences & Medicine, School of Immunology & Microbial Sciences, Kings College, London, U.K.

* Authors contributed equally (ordered alphabetically)

24   Abstract word count: 249

25   Text word count: 3692

Abstract

HIV pre-exposure prophylaxis (PrEP) is effective at preventing HIV acquisition, but its impact on the microbiome of the penis has not been described. As part of the CHAPS randomized clinical trial, we sequenced a segment of the bacterial 16S rRNA gene from foreskin tissue collected during surgical penile circumcision of 144 adolescents in South Africa and Uganda after randomization to placebo, emtricitabine with tenofovir disoproxil fumarate, or emtricitabine with tenofovir alafenamide. We found a large proportion of *Corynebacterium* in addition to other anaerobic species. *Cutibacterium acnes* was more abundant among participants from South Africa than Uganda, though this made no difference in surgical recovery. We did not find a difference in bacterial populations by treatment received nor bacterial taxa that were differentially abundant between participants who received placebo versus active drug. Using RNAseq libraries from the same participants, we found generally negative correlations between the relative abundance of bacterial taxa and the expression of genes downstream of the innate response to bacteria and regulation of the inflammatory response. When participants were divided into two clusters based on bacterial community composition, these clusters were also distinguished by high and low bacterial diversity. Random forest classification showed higher expression of *NFATC3* and *SELENOS* and lower expression of *STAP1* and *NLRP6* in the higher diversity group compared to the lower. Our results show no difference in the tissue microbiome of the foreskin in the different PrEP arms versus placebo. Bacterial taxa were largely inversely correlated with gene expression, consistent with non-inflammatory colonization.

Importance

We found that a short course of oral medicine to prevent HIV was not associated with a difference in the bacterial community of the foreskin of the penis. Previous studies found increased inflammation with certain anaerobic bacteria from swabs taken under the foreskin, but we found that higher relative abundances of the bacteria were correlated with lower

51   expression of inflammatory genes. Understanding the relationship between bacteria and

52   inflammation in the penis will help us to understand how interventions like penile circumcision

53   reduce the risk of acquiring sexually transmitted infections such as HIV.

Introduction

HIV remains a significant global health challenge despite substantial clinical, public health, and basic science research efforts. While condom use (1, 2), pre-exposure prophylaxis (PrEP) (3), and medical penile circumcision (4) are effective at reducing HIV incidence in men, the contribution of the penile microbiome to these mechanisms has not been fully explored. Previous work has shown a predominance of anaerobic species in the microbiota of swabs taken from the coronal sulcus or urethra (5-15) and have reported associations between species such as *Prevotella* and increased mucosal inflammation, HIV target cell density and risk of HIV acquisition (8, 16). Following circumcision, the surface microbiota shifts to be dominated by more aerobic species as found on other skin surfaces (10, 17, 18). Thus far, no data exist as to the microbiota of the foreskin itself, and its relation to tissue inflammation.

Antiretrovirals (ARVs) are used for both treatment and prevention of HIV, and limited data have shown a complex relationship with bacteria. When applied as topical vaginal pre-exposure prophylaxis, *L. crispatus* was shown to endocytose Tenofovir (TFV) then either actively metabolize or release it back into the environment (19). Similarly, *Gardnerella vaginalis* and other anaerobes have been shown to metabolize TFV (20) or block its entry into cells by secretion of adenine (19). Antiretrovirals may also theoretically alter bacteriophage populations which can dramatically reshape the bacterial component of the microbiome which they infect (21). At the rectal mucosa, small studies have investigated the effects of oral emtricitabine (FTC) with tenofovir disoproxil fumarate (TDF) on the bacterial microbiome in men who have sex with men (MSM) and transwomen (22-24) but with varying results.

In a randomized, open-label trial of 1-2 doses of FTC with TDF versus with tenofovir alafenamide (TAF) prior to circumcision, we characterized the bacterial microbiome and inflammatory gene expression of foreskin tissue removed during circumcision, hypothesizing that short courses of PrEP, as utilized in a dose-finding trial, would not result in significant

80 changes to the bacterial microbiome of the foreskin, but that there would be a relationship

81 between the microbiota and inflammatory gene expression.

Materials and Methods

Cohort and Specimen Collection

The Combined HIV Adolescent PrEP and Prevention Study (CHAPS) was a randomized

controlled trial that enrolled 144 men living without HIV aged 13-24 years between 2019 and

2021 from the Chris Hani Baragwanath Academic Hospital in Soweto, South Africa (n=72) and

the Entebbe Regional Referral Hospital in Entebbe, Uganda (n=72). Inclusion criteria were male

sex at birth, hemoglobin > 9 g/dL, weight > 35 kg, two successive negative rapid HIV antibody

tests, and clinical eligibility for surgical circumcision. Exclusion criteria were conditions

precluding circumcision or receipt of the study medications. Participants were randomized to

placebo versus FTC with either TDF or TAF for 1-2 days prior to surgical penile circumcision to

investigate ARV dosing for on-demand PrEP (25).

All participants underwent a physical exam at study entry and completed survey instruments

including sexual history at the randomization visit. At the circumcision visit, they provided

midstream urine for *Chlamydia trachomatis* (CT) and *Neisseria gonorrhea* (GC) testing via nucleic

acid amplification testing (NAAT) prior to surgery. If an asymptomatic sexually transmitted

infection was diagnosed, antibiotic treatment was prescribed at the post-operative visit.

Penile circumcision was performed using the dorsal slit method and the removed prepuce was

placed immediately in cold Dulbecco's Modified Eagle Medium and shipped on ice within 1

hour (median 30 minutes) to the local laboratories in Uganda (Medical Research

Council/Uganda Virus Research Institute) and South Africa (Perinatal HIV Research Unit in

Johannesburg) for processing. Smaller, 5–7 mm^2-sized sections were stored dry at -80° C until

the samples were transported on dry ice to the Seattle Children's Research Institute, U.S.A for

microbiome studies and to the Karolinska Institutet, Sweden for transcriptome analyses.

Ethics and Human Subjects

Ethical clearance to conduct the CHAPS trial was obtained from the South African Health Products Regulatory Authority (20181004), the Uganda Virus Research Institute research and ethics committee (GC/127/18/12/680), Uganda National Council of Science and Technology (HS 2534), Uganda National Drug Authority (618/NDA/DPS/09/2019), and the London School of Hygiene and Tropical Medicine research ethics committee (Ref:17403). Informed written consent was collected from all participants. The Swedish Ethics Review Authority approved the transcriptome studies of the collected specimens at the Karolinska Institutet (2020-00941). The ethics approval for the microbiome analysis was granted by the Seattle Children's Institutional Review Board (STUDY00003430).

Specimen Processing

16S rRNA Analysis

At the time of analysis, vials were thawed and approximately 25 mg of tissue was dissected and processed by a customized PowerSoil Pro protocol (Supplement S1) for extraction of DNA using a QIAcube instrument. A negative extraction control consisting of solution CD1 without specimen was also included. Specimens from each collection site were extracted on separate plates. The resulting total DNA was diluted 1:4 to reduce PCR inhibition.

The *16S* rRNA gene V3-V4 region was amplified using 319F/806R universal primers for 20 cycles of PCR as previously described (26) for each specimen along with a negative PCR control reaction consisting of mastermix without DNA template for each replicate and evenly and staggered genomic DNA from mock bacterial libraries (BEI Resources) as positive sequencing controls. The amplified products were purified using Agencourt AMPure XP beads (Beckman Coulter) and submitted to an additional 10 rounds of PCR with indexing primers (Illumina). The resulting libraries were pooled by volume with specimens at 100x the positive controls. The resulting library was purified using a MinElute PCR purification column (Qiagen), followed by

the QiaQuick gel extraction kit (Qiagen). The cleaned library was quantitated using qPCR

(NEBNext Library Quant Kit for Illumina), then pooled with PhiX, denatured, and loaded onto

a MiSeq instrument (Illumina) with a v3 2x300 flow cell following the manufacturer's protocol.

Sequences were de-multiplexed using Illumina's BaseSpace workflow. Primers and adapters

were removed by cutadapt 2.7 (27). Sequences were further trimmed for quality, then filtered

and merged using dada2 1.22.0 (28) to generate amplicon sequence variants (ASVs). Taxa were

annotated using the Silva 138.1 database (29) with additional genital-associated species (30)

using a 100% identity threshold. The phyloseq 1.40.0 (31) and vegan 2.6-2 (32) R packages were

used to manipulate ASV tables and calculate diversity measures. ASV sequences were aligned

using ssu-align 0.1.1 (33), and a maximum likelihood phylogeny was generated using PhyML

3.3.20220408 (34) with a GTR substitution model. Contaminating sequences were identified by

their presence in negative controls for the extraction and PCR amplification or mock community

using decontam 1.16.0 (35) and microfiltR (36). After decontamination, specimens with fewer

than 25-fold as many reads than extraction and PCR controls were excluded. For differential

abundance analysis, decontaminated ASVs were filtered with prevalence >= 10% and relative

abundance threshold of 1×10^{-4} before combining counts for all ASVs classified as the same

species. ALDEx2 1.28.1 (37), ANCOM-BC 1.6.2 (38), and DESeq2 1.36.0 (39) (using the poscounts

factors estimation) were used for differential abundance testing to overcome the documented

limited power and accuracy of these tools when used individually on 16S data sets which

contain a high proportion of zero counts (40, 41).

RNAseq of Foreskin Tissue

Foreskin samples were disrupted and homogenized using a Tissuelyzer (Qiagen) and total RNA

isolated using the RNeasy Kit (Qiagen) according to manufacturer's instructions. RNA was

subjected to quality control with Agilent Bioanalyzer (Agilent). To construct libraries suitable

for Illumina sequencing, the Illumina stranded mRNA prep ligation sample preparation

protocol was used with starting concentration of 200 ng total RNA. The protocol includes

mRNA isolation, cDNA synthesis, ligation of adapters and amplification of indexed libraries.

The yield and quality of the amplified libraries were analysed using Qubit by (Thermo Fisher)

and the Agilent Tapestation (Agilent). The indexed cDNA libraries were normalized and

combined, and the pools were sequenced on the Illumina Novaseq 6000 S4 flowcell to generate

150 bp paired-end reads.

Sample demultiplexing was performed using bcl2fastq 2.20.0 (Illumina), and quality and

adapter trimming of reads was performed using Cutadapt 2.8 (27). Sample quality was assessed

using FastQC 0.11.8 (Babraham Bioinformatics) and MultiQC 1.7 (42). Reads were aligned to the

Ensembl GRCh38 reference genome using STAR 2.6.1d (43). Counts for each gene were

obtained using featureCounts 1.5.1 (44).

The Gene Ontology (GO) term "inflammatory response" (GO:0006954) selected 860 putative

inflammatory genes which were filtered to only those with at least two copies detected in at

least 90% of specimens. RNA read counts were normalized then transformed by centered log

ratio (CLR). The 16S ASVs were filtered and combined as described for the differential

abundance analysis and also CLR-transformed. We calculated the correlation between the gene

counts and bacterial relative abundances, then filtered for r >0.4 and Benjamani-Hochberg-

adjusted p-value < 0.05. The resulting genes were manually inspected for their most relevant

GO annotation and grouped according to their immunological function and pro- or anti-

inflammatory nature (Supplemental table S1).

Normalized gene counts were used to perform random forest feature selection as implemented

in the Boruta R package 7.0.0 (45). Only the importance measures of statistically significant

(p<0.01) features were reported.

Statistical Analyses

All statistical analyses were performed in R version 4. Alpha diversity comparisons were

evaluated using the Wilcoxon rank sum test. Beta diversity was compared using Permutational

181 Multivariate of Variance (PERMANOVA) using the adonis2 function of the vegan R package.

182 The relationship between treatment arm and CST was assessed using multinomial logistic

183 regression. RNAseq and 16S taxa correlations were calculated using Pearson coefficient. A

184 significance threshold of $\alpha = 0.05$ was used for the differential abundance hypothesis testing.

185 ## Data Availability

186 The 16S raw reads and RNASeq libraries will be deposited in the National Center for

187 Biotechnology Information Short Read Archive and European Bioinformatics Institute

188 (respectively) upon publication. R code to reproduce the analysis is available at

189 https://github.com/bmaust/CHAPS/. The completed STORMS (Strengthening The

190 Organizing and Reporting of Microbiome Studies) checklist (46) for this project is located at

191 https://doi.org/10.5281/zenodo.7080176.

192 # Results

193 *Clinical STI testing*

194 No participants reported STI symptoms, and no physical exams revealed urethral discharge or

195 other genital abnormality. None of the participants had clinical balanoposthitis or evidence of

196 macroscopic inflammation. No GC infections were diagnosed, but NAAT for seven participants

197 was positive for CT: five from Uganda and two from South Africa.

198 *Microbiome 16S sequencing*

199 After filtering and contamination removal, 137 specimens from the 144 enrolled participants

200 had sufficient bacterial DNA reads to proceed with analysis. The identified bacterial taxa

201 include a variety of skin-associated Gram-positive and genital-associated anaerobic species in

202 addition to Gram-negative enterics (Figure 1). *Corynebacterium* was the most prevalent and

203 abundant genus, appearing in 132 (97%) of specimens at median relative abundance of 34%

204 (range: 0.14% to 98%). Anaerobic species were extremely common, including bacteria that are

commonly found in bacterial vaginosis, an inflammatory dysbiosis of the vagina, including

Prevotella, *Anaerococcus*, *Finegoldia* and *Porphyromonas*. There were no ASVs in the *Chlamydiaceae*

family which includes *C. trachomatis*.

Unsupervised partition around medioids clustering separated the unweighted Unifrac

distances into two community structure types (CST) with distinct community structure

(PERMANOVA R^2 = 0.25 with p < 0.001) and alpha diversity (p=2.39x10^{-19}) (Figure 2). Two

clusters maximized the silhouette score with acceptable within sum of squares and gap

statistics. CST1 was highly diverse with a median Shannon index of 3.05. CST2 was dominated

by *Corynebacterium tuberculostearicum* and *Finegoldia magna* and, with median relative

abundances of 21% and 9.6%, respectively. CST2 also had a significantly lower median Shannon

index of 1.68 (Wilcoxon unpaired exact, p = 2.39x10^{-19}). Though *F. magna* and *C.*

tuberculostearicum were also highly abundant in CST1 (median relative abundances of 4.6% and

2.5%), they shared high relative abundance with *Anaerococcus*, *Campylobacter*, *Fenollaria*,

Finegoldia, *Ezakiella*, *Mobiluncus*, and *Peptinophilis* species without a clear dominant taxon.

Microbiome differences by study site

The 137 participants with 16S data included 69 (50.3%) individuals from South Africa and 68

(49.7%) from Uganda. We compared microbiota between the two study sites and found no

differences in within-participant alpha diversity (Shannon entropy) or between-participant beta

diversity (unweighted Unifrac distance) (Figure 3, A and B). Differential abundance testing

identified *Cutibacterium acnes* as significantly higher in participants from South Africa compared

to those from Uganda by ALDEx2 with a CLR difference of 3.3 between sites (Wilcoxon rank

test with Benjamani-Hochberg correction p=0.0184). ANCOM-BC identified the same ASV with

a significant q-value (0.0079), but the log$_2$fold change of 1.87 failed to meet the effect size

threshold. DESeq2 did not identify any taxa as significantly differentially abundant. (Figure 3,

C-E).

Microbiome differences by treatment arm

We performed similar analyses for the bacterial populations in participants who received active

drug versus placebo. The participants with 16S data were equally distributed among treatment

arms with FTC-TAF and FTC-TDF (n=59 and 62, respectively, χ^2 p=0.437). All 16 participants

who received placebo had sufficient sequences for analysis. We found no differences in alpha or

beta diversity (Figure 4, A and B) between placebo and FTC-TAF or FTC-TDF regimens.

Combining both treatment groups and comparing to placebo, no species were significantly

differentially abundant by any of the three tools (Figure 4, C-D). An un-annotated *Dialister*

species was identified as statistically significant by ANCOM-BC but did not meet the effect size

threshold. The treatment arm was not a significant predictor of the CSTs identified above

(p=0.32 for placebo, p=0.99 for FTC/TAF, p=0.88 for FTC/TDF).

Inflammatory genes and bacterial taxa

Forty inflammatory genes showed significant correlation with 31 bacterial species (Table 1). Six

genes had insufficient evidence for inflammatory function and were excluded. The remaining

34 genes were primarily pro-inflammatory with negative correlation to bacterial species (Figure

5) without difference by environmental niche. IL-15 was the most frequently correlated gene,

with significant negative correlations to seven bacterial taxa not typically associated with

invasive infection: *Brevibacterium luteolum*, *Corynebacterium urealyticum*, *Dietzia timorensis*, and

unannotated species in the *Cutibacterium, Corynebacterium, Dietzia,* and *Nosocomiicoccus* genera.

The majority of the bacterial taxa significantly correlated with other genes were gram-positive

organisms which frequently colonize the skin. The CLR-transformed relative abundance of

Corynebacterium massilliense, in particular, was associated with significantly lower expression of

genes involved in regulation of inflammatory responses and neutrophil chemotaxis and

activation. Anaerobes also found in the oral cavity such as *Parvimonas* and *Porphyromonas* were

also correlated with primarily lower expression of inflammatory genes. However, the oral

anaerobe *Rothia amarae* was associated with higher expression of the regulatory factor *GHSR*

and an unclassified species also in the *Rothia* genus was associated with higher expression of the

pro-inflammatory gene *REG3G*. Species in three genera canonically associated with BV,

Atopobium, Prevotella, and *Sneathia,* were correlated with lower expression of inflammatory

genes, but one unannotated *Prevotella* ASV was negatively correlated with *ZFP36*, an immune

regulatory gene.

We performed a similar analysis, grouping bacterial ASVs at the genus level (Supplemental

table S2). Eight genes showed correlation with nine bacterial genera. As expected, all the

correlated genes and bacterial genera were also identified in the species-level analysis.

We conducted a separate query of associations between inflammatory genes and the CSTs

described in Figure 2 A distinguished by high and low diversity bacterial populations. In the

random forest classification, four features achieved statistical significance. Nuclear Factor of

Activated T cells 3 (*NFATC3*) and Selenoprotein S (*SELENOS*) showed higher expression in the

highly diverse CST1 relative to CST2, while Signal Transducing Adapter Family Member 1

(*STAP1*) and Nod-like Receptor Pyrin domain-containing 6 (*NLRP6*) showed lower expression

(Figure 6).

Discussion

Our study is the first of which we are aware to analyze the tissue-level microbiome of the

foreskin. Consistent with previous reports using penile swabs (5-18, 47), the bacteria we

identified are predominated by taxa commonly colonizing the skin (chiefly *Corynebacteria* spp)

in addition to anaerobic bacteria such as *Prevotella, Dialister, Murdochiella, Peptoniphilus,* and

Negativicoccus. These species have been associated with increased inflammation and HIV

acquisition in uncircumcised men (8) and bacterial vaginosis in women (48). In the foreskin, we

describe two major CSTs, one significantly more diverse than the other. While we found no

effect of one or two doses of antiretroviral exposure on the foreskin bacterial microbiome,

further studies of more prolonged ARV use may better elucidate whether there is an effect on

bacterial or viral communities of the penis.

The overall composition of the bacterial community did not appear to differ by study site;

however, we did find *Cutibacterium acnes* to be significantly more abundant in South African

than Ugandan participants. *C. acnes* is typically resident in the deep dermis in association with

sebaceous glands and hair follicles (49), which our study sampled by digesting full-thickness

specimens rather than resuspending skin swabs. While its contribution to its namesake acne

vulgaris is debated, it is otherwise non-pathogenic in immunocompetent hosts without artificial

material (49). It is frequently identified in surgical cultures, with unclear significance to sterility,

likely due to its resistance to surgical sterilization techniques and transection of deep dermal

structures during surgery (50). The differential abundance we observed may have been caused

by different surgical preparation at the two sites or by an actual difference in bacterial

populations. As there were no surgical complications observed during the study, the difference

does not appear to have clinical significance.

We did not identify bacterial species that were differentially abundant between participants

receiving placebo compared to emtricitabine with either of the two forms of tenofovir. While

vaginal-associated species such as *Lactobacillus* and *Gardnerella* take up tenofovir from their

environment (19, 20), a longer exposure than our study may be required to see a potential effect

on bacterial growth or to alter bacteriophage activity.

Lack of concordance between the clinical STI testing with NAAT and our 16S results is not

surprising, as *C. trachomatis* is an obligate intracellular pathogen targeting the urethral

epithelium and upper genital tract including the prostate and seminal vesicles but not the glans

or foreskin of the penis (51). Our lack of detection in foreskin tissue is consistent with the

previously described cellular tropisms and emphasizes the need for screening via the

appropriate site to evaluate infection.

Our inflammatory gene analysis primarily identified an inverse relationship between

expression of genes associated with response to bacteria and skin commensals such as

Corynebacterium and *Cutibacterium*, consistent with non-inflammatory colonization. IL-15, the

most commonly correlated gene, is a pleotropic cytokine secreted by a narrow range of cell

types, in foreskin tissue including epithelial cells, fibroblasts, Langerhans cells, and monocytes

(52). It has broad immunostimulatory function, promoting NK cell differentiation and survival

(53), inflammatory cytokine production by macrophages (54) and dendritic cells (55), neutrophil

activation, survival, and phagocytosis (56), germinal center B cell proliferation (57), CD8$^+$ T cell

survival (58), and it is required for development of skin-resident memory CD8$^+$ T cells (59). Its

lower expression correlating with higher relative abundances of non-pathogenic bacteria is

consistent with reduced inflammatory signaling corresponding to increased bacterial growth.

When examining gene expression between the participant groups with higher and lower

diversity bacterial communities, we did not confirm previous findings in the penis (8, 10, 16)

and vagina (60) that high bacterial diversity, particularly with anaerobic taxa, is associated with

increased inflammation. While we frequently identified anaerobic taxa, individual species

showed lower prevalence than in studies using surface swabs. This may have reduced our

power to detect an inflammatory association. Another explanation could be in the structure of

our experiment: rather than measuring secreted cytokines, we processed the entire tissue

specimen for bulk RNAseq, which likely included many cells not directly interacting with

bacteria or immune cells. Alternatively, the pre-procedure sterilization or the surgery itself may

have preferentially removed inflammatory species. The sterilization or collection could also

have altered host-bacterial interactions, though the rapid timeframe makes this explanation

seem less likely. Lastly, the bacterial strains in our cohort may employ different metabolic or

virulence strategies that render them less inflammatory. Unfortunately, our experiment was not

structured to investigate this possibility.

In the vagina, high diversity communities are highly inflammatory (60) and associated with

higher risk of adverse outcomes including HIV acquisition (61, 62) and preterm birth (63). The

foreskin microbiome had two obvious CSTs, one highly diverse CST1 and a less diverse CST2.

We hypothesized that the more diverse CST1 would be associated with higher levels of genes

related to inflammation, as in the vagina. However, when comparing gene expression between

CST1 versus CST2 specimens, random forest feature selection identified four genes, not

including canonical acute innate antibacterial response pathways. While named for its key

activity in T-Cell Receptor (TCR) signaling, NFAT family members play a broad role in cell

differentiation in the immune system and beyond (64), including in B cells (65), Toll-like

receptor (TLR) signaling in monocytes (66), and proliferation in perivascular tissue (67) and

keratinocytes (68). Higher expression of NFATC3 in CST1 specimens may represent increased

signaling from any of these cell types. SELENOS is up-regulated by cytokines such as IL-1β and

TNF-α via NF-κB and in turns acts to suppress cytokine secretion in macrophages (69). Its

higher expression in CST1 samples is consistent with increased suppression of chronic

inflammation. STAP1 has the best evidence for signaling downstream of the B-Cell Receptor

(BCR) (70), and lower expression could represent a response to sustained signaling (although

this was not supported by the expected changes in other genes). NLRP6 is both an inducer and

a component of the inflammasome, binds directly to the lipopolysaccharide of gram negative or

lipoteichoic acid of gram positive bacterial cell membranes, and plays both a pro- and anti-

inflammatory role in tissues such as liver, kidney, and intestine (71). Given these contrasting

roles, the effects of its lower expression are difficult to predict with available data. None of the

identified genes plays a first-line role in regulating bacterial sensing or inflammatory response.

In summary, we report on the first tissue-level examination of the bacterial microbiome of the

foreskin, without apparent effect of brief antiretroviral drug exposure. Correlating bacterial

species with RNAseq data revealed largely negative correlations with genes involved in the

inflammatory response, consistent with maintenance of immune tolerance.

Acknowledgements

This study is part of the EDCTP2 programme supported by the European Union (grant number
RIA2016MC-1616). BSM was supported by NIH training grant T32HD007233. The study
received support by a grant from the Swedish Research Council (Francesca Chiodi;
Vetenskapsrådet 2019-04596). The funders had no role is study design, data collection, and
interpretation, or the decision to submit the work for publication. The authors declare that no
conflict of interest exists.

References

1. Weller S, Davis K. 2002. Condom effectiveness in reducing heterosexual HIV transmission. Cochrane Database Syst Rev doi:10.1002/14651858.CD003255:CD003255.
2. Silverman BG, Gross TP. 1997. Use and effectiveness of condoms during anal intercourse. A review. Sex Transm Dis 24:11-7.
3. Spinner CD, Boesecke C, Zink A, Jessen H, Stellbrink HJ, Rockstroh JK, Esser S. 2016. HIV pre-exposure prophylaxis (PrEP): a review of current knowledge of oral systemic HIV PrEP in humans. Infection 44:151-8.
4. Friedman B, Khoury J, Petersiel N, Yahalomi T, Paul M, Neuberger A. 2016. Pros and cons of circumcision: an evidence-based overview. Clin Microbiol Infect 22:768-774.
5. Nelson DE, Dong Q, Van der Pol B, Toh E, Fan B, Katz BP, Mi D, Rong R, Weinstock GM, Sodergren E, Fortenberry JD. 2012. Bacterial communities of the coronal sulcus and distal urethra of adolescent males. PLoS One 7:e36298.
6. Carda-Diéguez M, Cárdenas N, Aparicio M, Beltrán D, Rodríguez JM, Mira A. 2019. Variations in Vaginal, Penile, and Oral Microbiota After Sexual Intercourse: A Case Report. Front Med (Lausanne) 6:178.
7. Onywera H, Williamson AL, Ponomarenko J, Meiring TL. 2020. The Penile Microbiota in Uncircumcised and Circumcised Men: Relationships With HIV and Human Papillomavirus Infections and Cervicovaginal Microbiota. Front Med (Lausanne) 7:383.
8. Prodger JL, Abraham AG, Tobian AA, Park DE, Aziz M, Roach K, Gray RH, Buchanan L, Kigozi G, Galiwango RM, Ssekasanvu J, Nnamutete J, Kagaayi J, Kaul R, Liu CM. 2021. Penile bacteria associated with HIV seroconversion, inflammation, and immune cells. JCI Insight 6.
9. Liu CM, Hungate BA, Tobian AA, Ravel J, Prodger JL, Serwadda D, Kigozi G, Galiwango RM, Nalugoda F, Keim P, Wawer MJ, Price LB, Gray RH. 2015. Penile Microbiota and Female Partner Bacterial Vaginosis in Rakai, Uganda. mBio 6:e00589.
10. Galiwango RM, Park DE, Huibner S, Onos A, Aziz M, Roach K, Anok A, Nnamutete J, Isabirye Y, Wasswa JB, Male D, Kigozi G, Tobian AAR, Prodger JL, Liu CM, Kaul R. 2022. Immune milieu and microbiome of the distal urethra in Ugandan men: impact of penile circumcision and implications for HIV susceptibility. Microbiome 10:7.
11. Li M, Mao JX, Jiang HH, Huang CM, Gao XH, Zhang L. 2021. Microbiome Profile in Patients with Adult Balanoposthitis: Relationship with Redundant Prepuce, Genital Mucosa Physical Barrier Status and Inflammation. Acta Derm Venereol 101:adv00466.
12. Mehta SD, Zhao D, Green SJ, Agingu W, Otieno F, Bhaumik R, Bhaumik D, Bailey RC. 2020. The Microbiome Composition of a Man's Penis Predicts Incident Bacterial Vaginosis in His Female Sex Partner With High Accuracy. Front Cell Infect Microbiol 10:433.

13. Mehta SD, Nandi D, Agingu W, Green SJ, Bhaumik DK, Bailey RC, Otieno F. 2020. Vaginal and
Penile Microbiome Associations with HSV-2 in Women and their Male Sex Partners. J Infect Dis
doi:10.1093/infdis/jiaa529.
14. Onywera H, Williamson AL, Cozzuto L, Bonnin S, Mbulawa ZZA, Coetzee D, Ponomarenko J,
Meiring TL. 2020. The penile microbiota of Black South African men: relationship with human
papillomavirus and HIV infection. BMC Microbiol 20:78.
15. Plummer EL, Vodstrcil LA, Danielewski JA, Murray GL, Fairley CK, Garland SM, Hocking JS,
Tabrizi SN, Bradshaw CS. 2018. Combined oral and topical antimicrobial therapy for male
partners of women with bacterial vaginosis: Acceptability, tolerability and impact on the genital
microbiota of couples - A pilot study. PLoS One 13:e0190199.
16. Liu CM, Prodger JL, Tobian AAR, Abraham AG, Kigozi G, Hungate BA, Aziz M, Nalugoda F,
Sariya S, Serwadda D, Kaul R, Gray RH, Price LB. 2017. Penile Anaerobic Dysbiosis as a Risk
Factor for HIV Infection. mBio 8.
17. Liu CM, Hungate BA, Tobian AA, Serwadda D, Ravel J, Lester R, Kigozi G, Aziz M, Galiwango
RM, Nalugoda F, Contente-Cuomo TL, Wawer MJ, Keim P, Gray RH, Price LB. 2013. Male
circumcision significantly reduces prevalence and load of genital anaerobic bacteria. mBio
4:e00076.
18. Price LB, Liu CM, Johnson KE, Aziz M, Lau MK, Bowers J, Ravel J, Keim PS, Serwadda D, Wawer
MJ, Gray RH. 2010. The effects of circumcision on the penis microbiome. PLoS One 5:e8422.
19. Taneva E, Sinclair S, Mesquita PM, Weinrick B, Cameron SA, Cheshenko N, Reagle K, Frank B,
Srinivasan S, Fredricks D, Keller MJ, Herold BC. 2018. Vaginal microbiome modulates topical
antiretroviral drug pharmacokinetics. JCI Insight 3.
20. Klatt NR, Cheu R, Birse K, Zevin AS, Perner M, Noël-Romas L, Grobler A, Westmacott G, Xie IY,
Butler J, Mansoor L, McKinnon LR, Passmore JS, Abdool Karim Q, Abdool Karim SS, Burgener
AD. 2017. Vaginal bacteria modify HIV tenofovir microbicide efficacy in African women. Science
356:938-945.
21. Sutton TDS, Hill C. 2019. Gut Bacteriophage: Current Understanding and Challenges. Front
Endocrinol (Lausanne) 10:784.
22. Fulcher JA, Li F, Cook RR, Zabih S, Louie A, Okochi H, Tobin NH, Gandhi M, Shoptaw S,
Gorbach PM, Aldrovandi GM. 2019. Rectal Microbiome Alterations Associated With Oral Human
Immunodeficiency Virus Pre-Exposure Prophylaxis. Open Forum Infect Dis 6:ofz463.
23. Dubé MP, Park SY, Ross H, Love TMT, Morris SR, Lee HY. 2018. Daily HIV pre-exposure
prophylaxis (PrEP) with tenofovir disoproxil fumarate-emtricitabine reduced Streptococcus and
increased Erysipelotrichaceae in rectal microbiota. Sci Rep 8:15212.
24. Perler BK, Reinhart EM, Montgomery M, Maynard M, Shapiro JM, Belenky P, Chan PA. 2021.
Evaluation of the Microbiome in Men Taking Pre-exposure Prophylaxis for HIV Prevention.
AIDS Behav doi:10.1007/s10461-020-03130-7.
25. Nash S, Dietrich J, Ssemata AS, Herrera C, O'Hagan K, Else L, Chiodi F, Kelly C, Shattock R,
Chirenje M, Lebina L, Khoo S, Bekker LG, Weiss HA, Gray C, Stranix-Chibanda L, Kaleebu P,
Seeley J, Martinson N, Fox J, team C. 2020. Combined HIV Adolescent Prevention Study
(CHAPS): comparison of HIV pre-exposure prophylaxis regimens for adolescents in sub-Saharan
Africa-study protocol for a mixed-methods study including a randomised controlled trial. Trials
21:900.
26. Dabee S, Mugo N, Mudhune V, McLellan-Lemal E, Peacock S, O'Connor S, Njoroge B, Nyagol B,
Thurman AR, Ouma E, Ridzon R, Wiener J, Haugen HS, Gasper M, Feng C, Allen SA, Doncel GF,
Jaspan HB, Heffron R, Kisumu Combined Ring Study T. 2022. Genital microbiota of women
using a 90 day tenofovir or tenofovir and levonorgestrel intravaginal ring in a placebo controlled
randomized safety trial in Kenya. Sci Rep 12:12040.
27. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnetjournal 17:10.
28. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods 13:581-3.
29. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner F. 2013. The
SILVA ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic acids research 41.
30. Brown BP. 2021. Updated 16S databases for marker gene taxonomic assignment.
31. McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. PLoS One 8:e61217.

32. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB,
Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2020. vegan: Community Ecology
Package, https://CRAN.R-project.org/package=vegan.
33. Nawrocki EP. 2009. Structural RNA Homology Search and Alignment Using Covariance Models.
PhD. Washington University in St. Louis, St. Louis, MO.
34. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML
3.0. Syst Biol 59:307-21.
35. Davis N, Proctor D, Holmes S, Relman D, Callahan B. 2018. Simple statistical identification and
removal of contaminant sequences in marker-gene and metagenomics data. Microbiome 6.
36. Brown BB. microfiltR, v1.2. https://github.com/itsmisterbrown/microfiltR.
37. Fernandes AD, Reid JN, Macklaim JM, McMurrough TA, Edgell DR, Gloor GB. 2014. Unifying
the analysis of high-throughput sequencing datasets: characterizing RNA-seq, 16S rRNA gene
sequencing and selective growth experiments by compositional data analysis. Microbiome 2:15.
38. Lin H, Peddada SD. 2020. Analysis of compositions of microbiomes with bias correction. Nat
Commun 11:3514.
39. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 15:550.
40. Nearing JT, Douglas GM, Hayes MG, MacDonald J, Desai DK, Allward N, Jones CMA, Wright
RJ, Dhanani AS, Comeau AM, Langille MGI. 2022. Microbiome differential abundance methods
produce different results across 38 datasets. Nat Commun 13:342.
41. Lin H, Peddada SD. 2020. Analysis of microbial compositions: a review of normalization and
differential abundance analysis. NPJ Biofilms Microbiomes 6:60.
42. Ewels P, Magnusson M, Lundin S, Kaller M. 2016. MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics 32:3047-8.
43. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15-21.
44. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30:923-30.
45. Kursa MB, Rudnicki WR. 2010. Feature Selection with the Boruta Package. Journal of Statistical
Software 36:1-13.
46. Mirzayi C, Renson A, Genomic Standards C, Massive A, Quality Control S, Zohra F, Elsafoury S,
Geistlinger L, Kasselman LJ, Eckenrode K, van de Wijgert J, Loughman A, Marques FZ,
MacIntyre DA, Arumugam M, Azhar R, Beghini F, Bergstrom K, Bhatt A, Bisanz JE, Braun J,
Bravo HC, Buck GA, Bushman F, Casero D, Clarke G, Collado MC, Cotter PD, Cryan JF, Demmer
RT, Devkota S, Elinav E, Escobar JS, Fettweis J, Finn RD, Fodor AA, Forslund S, Franke A,
Furlanello C, Gilbert J, Grice E, Haibe-Kains B, Handley S, Herd P, Holmes S, Jacobs JP, Karstens
L, Knight R, Knights D, Koren O, et al. 2021. Reporting guidelines for human microbiome
research: the STORMS checklist. Nat Med 27:1885-1892.
47. Srinivasan S, Chambers LC, Tapia KA, Hoffman NG, Munch MM, Morgan JL, Domogala D,
Sylvan Lowens M, Proll S, Huang ML, Soge OO, Jerome KR, Golden MR, Hughes JP, Fredricks
DN, Manhart LE. 2021. Urethral Microbiota in Men: Association of Haemophilus influenzae and
Mycoplasma penetrans With Nongonococcal Urethritis. Clin Infect Dis 73:e1684-e1693.
48. Dabee S, Passmore JS, Heffron R, Jaspan HB. 2021. The Complex Link between the Female
Genital Microbiota, Genital Infections, and Inflammation. Infect Immun 89.
49. Achermann Y, Goldstein EJ, Coenye T, Shirtliff ME. 2014. Propionibacterium acnes: from
commensal to opportunistic biofilm-associated implant pathogen. Clin Microbiol Rev 27:419-40.
50. Lee MJ, Pottinger PS, Butler-Wu S, Bumgarner RE, Russ SM, Matsen FA, 3rd. 2014.
Propionibacterium persists in the skin despite standard surgical preparation. J Bone Joint Surg
Am 96:1447-50.
51. Mackern-Oberti JP, Motrich RD, Breser ML, Sanchez LR, Cuffini C, Rivero VE. 2013. Chlamydia
trachomatis infection of the male genital tract: an update. J Reprod Immunol 100:37-53.
52. Perera PY, Lichy JH, Waldmann TA, Perera LP. 2012. The role of interleukin-15 in inflammation
and immune responses to infection: implications for its therapeutic use. Microbes Infect 14:247-
61.
53. Waldmann TA, Tagaya Y. 1999. The multifaceted regulation of interleukin-15 expression and the
role of this cytokine in NK cell differentiation and host response to intracellular pathogens. Annu
Rev Immunol 17:19-49.

54.   Alleva DG, Kaser SB, Monroy MA, Fenton MJ, Beller DI. 1997. IL-15 functions as a potent
515   autocrine regulator of macrophage proinflammatory cytokine production: evidence for
516   differential receptor subunit utilization associated with stimulation or inhibition. J Immunol
517   159:2941-51.
518   55.   Mattei F, Schiavoni G, Belardelli F, Tough DF. 2001. IL-15 is expressed by dendritic cells in
519   response to type I IFN, double-stranded RNA, or lipopolysaccharide and promotes dendritic cell
520   activation. J Immunol 167:1179-87.
521   56.   Girard D, Paquet ME, Paquin R, Beaulieu AD. 1996. Differential effects of interleukin-15 (IL-15)
522   and IL-2 on human neutrophils: modulation of phagocytosis, cytoskeleton rearrangement, gene
523   expression, and apoptosis by IL-15. Blood 88:3176-84.
524   57.   Park CS, Yoon SO, Armitage RJ, Choi YS. 2004. Follicular dendritic cells produce IL-15 that
525   enhances germinal center B cell proliferation in membrane-bound form. J Immunol 173:6676-83.
526   58.   Wu TS, Lee JM, Lai YG, Hsu JC, Tsai CY, Lee YH, Liao NS. 2002. Reduced expression of Bcl-2 in
527   CD8+ T cells deficient in the IL-15 receptor alpha-chain. J Immunol 168:705-12.
528   59.   Mackay LK, Rahimpour A, Ma JZ, Collins N, Stock AT, Hafon ML, Vega-Ramos J, Lauzurica P,
529   Mueller SN, Stefanovic T, Tscharke DC, Heath WR, Inouye M, Carbone FR, Gebhardt T. 2013. The
530   developmental pathway for CD103(+)CD8+ tissue-resident memory T cells of skin. Nat Immunol
531   14:1294-301.
532   60.   Anahtar MN, Byrne EH, Doherty KE, Bowman BA, Yamamoto HS, Soumillon M, Padavattan N,
533   Ismail N, Moodley A, Sabatini ME, Ghebremichael MS, Nusbaum C, Huttenhower C, Virgin HW,
534   Ndung'u T, Dong KL, Walker BD, Fichorova RN, Kwon DS. 2015. Cervicovaginal bacteria are a
535   major modulator of host inflammatory responses in the female genital tract. Immunity 42:965-76.
536   61.   McClelland RS, Lingappa JR, Srinivasan S, Kinuthia J, John-Stewart GC, Jaoko W, Richardson BA,
537   Yuhas K, Fiedler TL, Mandaliya KN, Munch MM, Mugo NR, Cohen CR, Baeten JM, Celum C,
538   Overbaugh J, Fredricks DN. 2018. Evaluation of the association between the concentrations of key
539   vaginal bacteria and the increased risk of HIV acquisition in African women from five cohorts: a
540   nested case-control study. Lancet Infect Dis 18:554-564.
541   62.   Masson L, Passmore JA, Liebenberg LJ, Werner L, Baxter C, Arnold KB, Williamson C, Little F,
542   Mansoor LE, Naranbhai V, Lauffenburger DA, Ronacher K, Walzl G, Garrett NJ, Williams BL,
543   Couto-Rodriguez M, Hornig M, Lipkin WI, Grobler A, Abdool Karim Q, Abdool Karim SS. 2015.
544   Genital inflammation and the risk of HIV acquisition in women. Clin Infect Dis 61:260-9.
545   63.   Wylie KM, Wylie TN, Cahill AG, Macones GA, Tuuli MG, Stout MJ. 2018. The vaginal eukaryotic
546   DNA virome and preterm birth. Am J Obstet Gynecol 219:189.e1-189.e12.
547   64.   Horsley V, Pavlath GK. 2002. NFAT: ubiquitous regulator of cell differentiation and adaptation. J
548   Cell Biol 156:771-4.
549   65.   Venkataraman L, Francis DA, Wang Z, Liu J, Rothstein TL, Sen R. 1994. Cyclosporin-A sensitive
550   induction of NF-AT in murine B cells. Immunity 1:189-96.
551   66.   Minematsu H, Shin MJ, Celil Aydemir AB, Kim KO, Nizami SA, Chung GJ, Lee FY. 2011. Nuclear
552   presence of nuclear factor of activated T cells (NFAT) c3 and c4 is required for Toll-like receptor-
553   activated innate inflammatory response of monocytes/macrophages. Cell Signal 23:1785-93.
554   67.   Graef IA, Chen F, Chen L, Kuo A, Crabtree GR. 2001. Signals transduced by Ca(2+)/calcineurin
555   and NFATc3/c4 pattern the developing vasculature. Cell 105:863-75.
556   68.   Santini MP, Talora C, Seki T, Bolgan L, Dotto GP. 2001. Cross talk among calcineurin, Sp1/Sp3,
557   and NFAT in control of p21(WAF1/CIP1) expression in keratinocyte differentiation. Proc Natl
558   Acad Sci U S A 98:9575-80.
559   69.   Curran JE, Jowett JB, Elliott KS, Gao Y, Gluschenko K, Wang J, Abel Azim DM, Cai G, Mahaney
560   MC, Comuzzie AG, Dyer TD, Walder KR, Zimmet P, MacCluer JW, Collier GR, Kissebah AH,
561   Blangero J. 2005. Genetic variation in selenoprotein S influences inflammatory response. Nat
562   Genet 37:1234-41.
563   70.   Ohya K, Kajigaya S, Kitanaka A, Yoshida K, Miyazato A, Yamashita Y, Yamanaka T, Ikeda U,
564   Shimada K, Ozawa K, Mano H. 1999. Molecular cloning of a docking protein, BRDG1, that acts
565   downstream of the Tec tyrosine kinase. Proc Natl Acad Sci U S A 96:11976-81.
566   71.   Angosto-Bazarra D, Molina-Lopez C, Pelegrin P. 2022. Physiological and pathophysiological
567   functions of NLRP6: pro- and anti-inflammatory roles. Commun Biol 5:524.

Tables

Taxon (Genus species)	Gene Symbol	Pearson corr	adj p
(Uknown *Muribaculaceae* family)	ADAMTS12	-0.404	5.90E-04
(Uknown *Muribaculaceae* family)	TUSC2	-0.497	2.63E-06
Aliicoccus (unclassified sp)	IL15	-0.557	2.41E-08
Anaerococcus lactolyticus	ODAM	0.479	8.41E-06
Arcanobacterium (unclassified sp)	CXCL9	-0.415	3.31E-04
Atopobium (unclassified sp)	JAM3	-0.420	2.64E-04
Brevibacillus (unclassified sp)	IL1RN	-0.401	6.89E-04
Brevibacterium luteolum	IL15	-0.563	1.48E-08
Campylobacter (unclassified sp)	CCL11	0.492	3.68E-06
Campylobacter (unclassified sp)	ITGB2	-0.465	2.03E-05
Campylobacter (unclassified sp)	RIPK2	-0.461	2.55E-05
Corynebacterium confusum	FPR3	-0.402	6.46E-04
Corynebacterium confusum	MS4A2	-0.439	9.72E-05
Corynebacterium confusum	SMPDL3B	-0.543	7.86E-08
Corynebacterium coyleae	APOA2	0.542	8.40E-08
Corynebacterium genitalium	BDKRB2	-0.474	1.14E-05
Corynebacterium genitalium	POLB	-0.434	1.23E-04
Corynebacterium genitalium	PPBP	0.414	3.52E-04
Corynebacterium massiliense	CAMK1D	-0.405	5.56E-04
Corynebacterium massiliense	DAB2IP	-0.595	7.04E-10
Corynebacterium massiliense	KDM6B	-0.482	6.95E-06
Corynebacterium massiliense	LILRB4	-0.419	2.83E-04
Corynebacterium massiliense	PLCG2	-0.412	4.01E-04
Corynebacterium massiliense	PRCP	-0.469	1.51E-05
Corynebacterium massiliense	RELA	-0.454	3.88E-05
Corynebacterium massiliense	SPATA2	-0.556	2.43E-08
Corynebacterium riegelii	APOA2	0.516	6.69E-07
Corynebacterium urealyticum	IL15	-0.607	2.22E-10
Cutibacterium (unclassified sp)	IL15	-0.650	1.75E-12
Cutibacterium (unclassified sp)	S1PR3	-0.407	5.03E-04
Cutibacterium (unclassified sp)	TLR1	-0.402	6.55E-04
Cutibacterium acnes	GPRC5B	-0.405	5.46E-04
Dermabacter vaginalis	FOS	-0.401	6.84E-04
Dietzia (unclassified sp)	IL15	-0.583	2.48E-09
Dietzia timorensis	IL15	-0.470	1.49E-05
Enhydrobacter (unclassified sp)	HDAC4	-0.423	2.21E-04
Lactobacillus iners	BDKRB2	-0.440	8.68E-05
Lactobacillus iners	POLB	-0.428	1.72E-04
Lactobacillus iners	PSTPIP1	-0.423	2.21E-04
Mycoplasma spermatophilum	IL9	0.432	1.36E-04
Nosocomiicoccus (unclassified sp)	IL15	-0.424	2.11E-04
Nosocomiicoccus (unclassified sp)	SMPDL3B	-0.489	4.44E-06
Parvimonas (unclassified sp)	RIPK2	-0.430	1.44E-04
Parvimonas micra	CCL18	-0.409	4.56E-04
Porphyromonas somerae	C4A	-0.409	4.52E-04
Prevotella bergensis	CXCR6	-0.430	4.52E-04
Prevotella_7 (unclassified sp)	ZFP36	-0.464	1.51E-04
Rothia (unclassified sp)	REG3G	0.534	2.18E-05
Rothia amarae	EGFR	-0.468	1.62E-07
Rothia amarae	GHSR	0.480	1.66E-05
Sneathia amnii	XCR1	-0.419	7.99E-06

569   Table 1 Bacterial taxa and human gene correlations

Figures

572   Figure 1 Top 25 most abundant bacterial genera across participants by study treatment received.

573   (anaerobic taxa in earth tones, aerobes in purples, uncultured or unidentified species in grays)

575   Figure 2 (A) Heatmap of top 25 bacterial species by CST, (B) PCoA ordination of bacterial beta

576   diversity by unweighted Unifrac distance, colored by CST

Figure 3 Comparisons by study site: alpha (A) and beta (B) diversity; differential abundance

testing with ALDEx2 (C), ANCOM-BC (D), and DESeq2 (E)

Figure 4 Analysis by treatment arm: alpha (A) and beta (B) diversity by drug received;

differential abundance testing with ALDEx2 (C), ANCOM-BC (D), and DESeq2 (E)

586   Figure 5 Bacterial taxa and inflammatory gene associations by immune function

CST1 vs CST2 feature importance (normalized counts)

Red represents upregulation and blue downregulation of genes in CST1

588 Figure 6 Random Forest Feature Importance comparing RNAseq results between CST 1 and

589 CST 2

Supplemental items

591 Supplemental Table S1: Gene categorization

592 Supplemental Table S2: Bacterial genera and human gene correlations

593 Supplement S1: Foreskin Tissue DNA Extraction Workflow