Evolution of an extreme hemoglobin phenotype contributed to the sub-Arctic specialization of extinct Steller's sea cows.

Anthony V. Signore[1†], Phillip R. Morrison[2‡], Colin J. Brauner[2], Angela Fago[3], Roy E. Weber[3] and Kevin L. Campbell[1]*

[1]Department of Biological Sciences, University of Manitoba, Winnipeg, R3T 3N2, Canada

[2]Department of Zoology, University of British Columbia, Vancouver, V6T 1Z4, Canada

[3]Department of Biology, Zoophysiology, Aarhus University, DK-8000 Aarhus, Denmark.

*For correspondence: Kevin.Campbell@umanitoba.ca

Present address: [†]National Centre for Foreign Animal Disease, Winnipeg, R3E 3M4, Canada; [‡]Department of Resource Management and Protection, and Biology Department, Vancouver Island University, Nanaimo, V9R 5S5, Canada

Author contributions: Anthony V Signore, Conceptualization, Project administration, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review and editing; Phillip R Morrison, Data curation, Investigation, Methodology, Writing – review and editing; Colin J Brauner, Project administration, Funding acquisition, Resources, Supervision, Writing – review and editing; Angela Fago, Project administration, Funding acquisition, Resources, Supervision, Writing – review and editing; Roy E Weber, Project administration, Funding acquisition, Resources, Supervision, Writing – review and editing; Kevin L Campbell, Conceptualization, Project administration, Funding acquisition, Resources, Formal analysis, Investigation, Methodology, Visualization, Supervision, Writing – original draft, Writing – review and editing.

Competing interests: Authors declare that they have no competing interests.

Keywords: hemoglobin, thermal adaptation, protein biochemistry, paleophysiology, Sirenia

This PDF file includes:

Main Text
Figures 1 to 4

Abstract

The extinct Steller's sea cow (*Hydrodamalis gigas*; †1768) was a whale-sized marine mammal that manifested profound morphological specializations to exploit the harsh coastal climate of the North Pacific. Yet despite first-hand accounts of their biology, little is known regarding the physiological adjustments underlying their evolution to this environment. Here, the adult-expressed hemoglobin (Hb; $\alpha_2\beta/\delta_2$) of this sirenian is shown to harbor a fixed amino acid replacement at an otherwise invariant position (β/δ82Lys→Asn) that alters multiple aspects of Hb function. First, our functional characterization of recombinant sirenian Hb proteins demonstrate that the Hb–O_2 affinity of this sub-Arctic species was less affected by temperature than those of living (sub)tropical sea cows. This phenotype presumably safeguarded O_2 delivery to cool peripheral tissues and largely arises from a reduced intrinsic temperature sensitivity of the *H. gigas* protein. Additional experiments on *H. gigas* β/δ82Asn→Lys mutant Hb further reveal this exchange renders Steller's sea cow Hb unresponsive to the potent intraerythrocytic allosteric effector 2,3-diphosphoglycerate, a radical modification that is the first documented example of this phenotype among mammals. Notably, β/δ82Lys→Asn moreover underlies the secondary evolution of a reduced blood–O_2 affinity phenotype that would have promoted heightened tissue and maternal/fetal O_2 delivery. This conclusion is bolstered by analyses of two Steller's sea cow prenatal Hb proteins (Hb Gower I; $\zeta_2\varepsilon_2$ and HbF; $\alpha_2\gamma_2$) that suggest an exclusive embryonic stage expression pattern, and reveal uncommon replacements in *H. gigas* HbF (γ38Thr→Ile and γ101Glu→Asp) that increased Hb–O_2 affinity relative to dugong HbF. Finally, the β/δ82Lys→Asn replacement of the adult/fetal protein is shown to increase protein solubility, which may have elevated red blood cell Hb content within both the adult and fetal circulations and contributed to meeting the elevated metabolic (thermoregulatory) requirements and fetal growth rates associated with this species cold adaptation.

Main Text

Introduction

The underwater foraging time of mammals is dictated by onboard oxygen stores and the efficiency of their use. Thus, evolutionary increases in oxygen stores, in the form of increased hemoglobin (Hb) and myoglobin—located within erythrocytes and skeletal/cardiac muscle, respectively—are nearly ubiquitous among mammalian divers (Ponganis, 2011). Notable exceptions to this rule are extant sirenians (sea cows), a group of strictly aquatic, (sub)tropical herbivores encompassing only four members; three species of manatee (family Trichechidae) and

the dugong, *Dugong dugon* (family Dugongidae). While sirenians are proficient divers, they do not exhibit the greatly elevated body O_2 stores or an enhanced dive reflex common to other lineages of marine mammals (Blessing, 1972; Scholander and Irving, 1941). Rather, previous work revealed that the sirenian's secondary transition to aquatic life coincided with a rapid evolution of their Hb encoding genes due, in part, to gene conversion events with a neighboring globin pseudogene (Signore et al., 2019). The resulting high blood–O_2 affinity phenotype presumably allows extant sea cows to maximize O_2 extraction from the lungs during submergence at the cost of somewhat reduced O_2 offloading, thus lowering overall metabolic intensity and fostering a prolonged breath-hold capacity (Signore et al., 2019).

While the relatively limited thermoregulatory capacity of extant sea cows confine them to (sub)tropical waters (Gallivan et al., 1986; Marsh et al., 2011), fossil evidence and first-hand accounts of the sub-Arctic Steller's sea cow (*Hydrodamalis gigas*) provide insights into the biological and morphological adaptations of this titanic sirenian to the harsh coastal conditions of the North Pacific, where they persisted from the Miocene (5 to 8 million years ago) until their demise in 1768 (Domning, 1976; Heritage and Seiffert, 2022; Stejneger, 1887; Steller, 1751). The retrieval of ancient genetic material from museum specimens has since been instrumental in clarifying the phylogenetic affinities and population history of this species, while providing additional details regarding the evolution of key morphological and physiological attributes (Gaudry et al., 2017; Le Duc et al., 2022; Mirceta et al., 2013; Sharko et al., 2019; Sharko et al., 2021; Signore et al., 2019; Springer et al., 2015) (Fig. 1A). For example, pilot experiments on "resurrected" Steller's sea cow recombinant Hb demonstrated that the Hb–O_2 affinity of this lineage secondarily decreased following their divergence from dugongs between the mid Oligocene and early Miocene (Signore et al., 2019). While sirenians do not possess the capacity for non-shivering thermogenesis due to pseudogenization of the *UCP1* gene (Gaudry et al., 2017), the reduced Hb–O_2 affinity shift in Steller's sea cow Hb was speculated to have promoted increased O_2 offloading to fuel increased thermogenesis to help cope with exposure to cold sub-Arctic waters. Although the Hb of this species accumulated 11 amino acid replacements since its divergence from the dugong (Figs. S1, S2), this decrease in Hb–O_2 affinity was hypothesized to arise from a highly unusual 82Lys→Asn exchange in the chimeric β-type (β/δ) chain (Signore et al., 2019). Data mined from more recent ancient DNA studies (Le Duc et al., 2022; Sharko et al., 2021) confirms that this substitution was fixed in the last remaining Steller's sea cow population (Fig. 1B), though the specific functional effect(s) of this substitution have not been characterized. This replacement is intriguing not only because β82Lys is invariant among characterized mammalian Hbs, but because human variants with substitutions at this position display profound alterations in both structural and functional properties (Abraham et al., 2011; Bonaventura et al., 1976; Ikkala et al., 1976; Sugihara et al., 1985). For example, the human Hb Providence (β82Lys→Asn) variant exhibits a decreased inherent Hb–O_2 affinity and

markedly reduced sensitivity to the allosteric effectors 2,3-diphosphoglycerate (DPG), Cl⁻, and H⁺

(Abraham et al., 2011; Bardakjian et al., 1985; Bonaventura et al., 1976; Charache et al., 1977;

Weickert et al., 1999), all of which preferentially bind and stabilize the (low O_2 affinity) deoxy-state

conformation of the protein. However, opposite to what was suggested for Steller's sea cows

(Signore et al., 2019), this exchange causes the whole blood O_2 affinity of Hb Providence carriers

to be noticeably higher than that of the general population (Bardakjian et al., 1985; Weber and

Campbell, 2011). It is thus unclear if and how the Steller's sea cow β/δ82Lys→Asn replacement

underlies the lower Hb–O_2 affinity of this extinct species relative to other sirenians, or whether this

attribute arises from one (or more) of the other 10 residue replacements that evolved in this lineage.

The β/δ82Lys→Asn exchange also raises other evolutionary significant questions, as it is

predicted to have altered multiple aspects of Hb function that may lead to antagonistic pleiotropic

effects. Notably, this exchange may detrimentally increase the effect of temperature on Hb–O_2

binding and release (Signore et al., 2019). The formation of the weak covalent bond between O_2

and the heme iron requires free energy, thus dictating an inverse relationship between Hb–O_2

affinity and temperature (Weber and Campbell, 2011). In temperate and Arctic endotherms this

inherent attribute of Hb potentially impedes O_2 delivery to the limbs and flukes, which are

maintained at substantively lower temperatures to minimize heat loss and hence energy

requirements (Campbell and Hofreiter, 2015). Accordingly, heterothermic mammals generally

possess Hbs whose O_2 binding properties are less sensitive to temperature than the Hbs of non-

cold-adapted species, thereby maintaining sufficient O_2 offloading in the face of decreasing tissue

temperatures. This reduction in thermal sensitivity (quantified as the overall enthalpy of

oxygenation, ΔH'), appears to predominantly arise from an increased interaction between allosteric

effectors and the Hb moiety (an exothermic process), which releases additional heat to assist with

deoxygenation (Weber and Campbell, 2011). Hence, the *H. gigas* β/δ82Lys→Asn replacement,

which deletes integral binding sites for the heterotropic ligands Cl⁻ and DPG (Bonaventura et al.,

1976), is puzzling in that it is expected to maladaptively increase the effect of temperature on O_2

uptake and release in the blood of the sub-Arctic Steller's sea cow.

Taken together, it remains unknown whether the *H. gigas* β/δ82Lys→Asn residue

exchange contributed adaptively to the species biology or is instead linked to small population sizes

(e.g., genetic drift) over the past half million years (Le Duc et al., 2022; Sharko et al., 2021). To

unravel the combined effects of evolved amino acid replacements on hemoglobin function in

relation to the extreme thermal biology of the extinct Steller's sea cow, we synthesized recombinant

Hb proteins of this extinct species together with those of the extant dugong (*Dugong dugon*) and

Florida manatee (*Trichechus manatus latirostris*), and measured their O_2 binding properties,

relative solubility, responses to allosteric effectors, and thermal sensitivities. We also synthesized

a *H. gigas* β/δ82Asn→Lys Hb mutant to assess the specific effects of this exchange, together with

the Hb of the last common ancestor ('ancestral dugongid') shared between the dugong and Steller's

sea cow (Fig. 1A) in order to assess the directionality of evolved physicochemical changes in Hb

function.

Results and Discussion

O_2 Affinity of Sirenian Hbs. Measured O_2-equilibrium curves of the five examined Hbs revealed

marked differences in intrinsic O_2 affinity (Figs. 2A, S3, S4 and Table S1). In the absence of

allosteric effectors (pH 7.2, 37°C), the P_{50} (the O_2 tension resulting in 50% Hb–O_2 saturation) of

Steller's sea cow Hb (P_{50} = 8.8 mm Hg) is ~2 times higher than that of dugong (3.5 mm Hg),

ancestral dugongid (4.3 mm Hg), and manatee (5.4 mm Hg) Hbs under the same conditions (Figs.

2A, S3 and Table S1). Site directed mutagenesis experiments reveal that the increased intrinsic

P_{50} of Steller's sea cow Hb predominantly arises from the β/δ82Asn substitution, as the

β/δ82Asn→Lys mutant exhibits an intrinsic P_{50} similar to that of the ancestral dugongid (Fig. 2A,

S4). Of note, the O_2 affinity of dugong, ancestral dugongid, and manatee Hbs was reduced in the

presence of Cl^- and DPG (P_{50} = 10.2, 9.9, and 10.9 mm Hg, respectively) by a similar degree to

that of Asian elephant Hb (Campbell et al., 2010b). This finding extends previous studies conducted

on sirenian Hbs (Farmer et al., 1979; McCabe et al., 1978; Signore et al., 2019), and reveals that

the high O_2 affinity of dugong and manatee blood is not attributable to decreased allosteric effector

sensitivity. Conversely, Steller's sea cow Hb was shown to be markedly less responsive to allosteric

effectors, as only a moderate reduction in O_2–affinity was observed in the presence of Cl^- and DPG

(P_{50} = 14.3 mm Hg). When the effects of these allosteric effectors were measured individually,

Steller's sea cow Hb exhibits lower DPG, Cl^-, and H^+ (Bohr) effects relative to those of the ancestral

dugongid and β/δ82Asn→Lys mutant (Fig. 2C-E; Table S1). These data confirm that a high intrinsic

(i.e. in the absence of allosteric effectors) Hb–O_2 affinity is an ancient sirenian trait that likely aided

the transition of the group to the aquatic environment, and that Hb–O_2 (and hence whole blood)

affinity was secondarily reduced in the Steller's sea cow lineage (Signore et al., 2019). This latter

finding contrasts with allometric expectations for mammals—whereby blood O_2 affinity and body

mass are inversely correlated (Schmidt-Neilsen and Larimer, 1958)—and thus further suggests this

modification served an adaptive function in this extinct species.

The single most distinct feature of *H. gigas* Hb is the lack of a discernable effect of DPG

on P_{50} ($\Delta logP_{50}^{(DPG-stripped)}$ = 0.02 at 37°C and pH 7.2; Fig. 2C and Table S1), relative to the Hbs of

the ancestral dugongid (0.27) and the extant manatee and dugong (0.14 and 0.30, respectively).

This intracellular effector generally occurs in equimolar concentrations to Hb (Bunn, 1980) and

strongly decreases the O_2 affinity of most mammalian Hbs via direct electrostatic interactions with

β2His and β82Lys (Fig. 3A), together with potential water-mediated interactions with β143His and

the α-NH$_2$ group of 1Val of the β$_2$ chain (Richard et al., 1993). However, unlike the other (ionizable)

residues whose ability to interact with DPG is highly pH dependent, β82Lys is strongly cationic and

thus is able to bind DPG across the entire physiological pH range. Presumably arising from this

indispensable role in DPG binding, this residue is uniformly conserved in mammalian Hbs (Fig. 3A),

with the exception of several heterozygous adult human HbA carriers with substitutions at this

position (Ikkala et al., 1976; Moo-Penn et al., 1976; Sugihara et al., 1985). Given that none of the

other six β/δ-chain replacements that evolved on the Steller's sea cow branch (Figs. 3B, S1 and

S2) are implicated in DPG binding, the deletion of the integral DPG binding site at β/δ82 in Steller's

sea cow Hb is fully consistent with its inability to bind DPG (Fig. 3B). This conclusion is further

supported by our measurements on the Steller's sea cow β/δ82Asn→Lys mutant, which show that

reversion to the ancestral state restores the DPG effect to the same level observed in ancestral

dugongid Hb (Fig. 2C). Notably, and despite possessing the identical DPG binding site residues as

Hb Providence, the *H. gigas* protein exhibits a distinctly lower DPG effect than this human variant

(0.08; 17,20). The lower DPG sensitivity of Steller's sea cow Hb thus implicates an epistatic

contribution from other amino acids in the vicinity of the DPG pocket. Importantly, the Lys→Asn

replacement in the DPG binding pocket causes the O$_2$ affinity of human HbA to increase in the

presence of allosteric cofactors (20,23), whereas results presented in Fig. 2A show Steller's sea

cow Hb–O$_2$ affinity is reduced relative to its ancestors carrying β/δ82Lys under all test conditions.

This observation highlights a growing body of research indicating that both the direction and overall

phenotypic effect of specific amino acid substitutions may be conditional on the genetic background

in which they occur (Natarajan et al., 2023; Storz, 2016).

The insensitivity of *H. gigas* Hb to DPG is also notable as it would have markedly reduced

their capacity to modulate blood–O$_2$ affinity *in vivo* (e.g., seasonally), and is the first demonstrated

example of a genuine DPG insensitive Hb phenotype among mammals. While eastern moles

(*Scalopus aquaticus*) are a possible exception (Campbell et al., 2010a), feliformid carnivores,

ruminants, and two species of lemurs have also traditionally been placed in the 'DPG insensitive'

category (Bunn, 1980) despite the fact their Hbs are moderately responsive to DPG in the absence

of Cl$^-$ (Bonaventura et al., 1976; Fronticelli et al., 1988; Janecka et al., 2015; Perutz et al., 1993).

Nonetheless, red blood cell DPG concentrations of species with suppressed DPG sensitivities are

markedly reduced relative to mammals whose Hb–O$_2$ affinity is regulated by DPG (<0.1-1.0 mM

vs. 4-10 mM, respectively; (Bunn, 1980)). While the potential benefits of a low DPG sensitivity

phenotype have been debated (Campbell et al., 2010a; Kay, 1977), approximately 20% of glucose

uptake by human erythrocytes is diverted to DPG synthesis via the Rapport-Luebering shunt (Duhm

et al., 1968), thereby bypassing production of both ATP molecules generated via the anaerobic

substrate level phosphorylation pathway (Bunn, 1980; Kauffman et al., 2002; Rapoport and

Luebering, 1950). Accordingly, since each molecule of DPG produced comes at the expense of

producing an ATP molecule, the probability that Steller's sea cow blood similarly contained low
levels of this organophosphate is high.

Hb Solubility. Ectopic expression of human Hb Providence β82Asp mutants in *E. coli* has been
shown to increase soluble protein production by 47-116% relative to the expression of human Hb
variants not carrying this substitution (Weickert et al., 1999). Consistent with this observation, we
found that Steller's sea cow Hb is more soluble than those of other sirenians and the engineered
H. gigas β/δ82Asn→Lys mutant (Fig. 2F and S5). While the precise mechanism underlying this
phenomenon is unknown, Hb Providence variants exhibit sharp reductions in irreversible oxidative
damage of nearby β93Cys that initiates Hb denaturation (Abraham et al., 2011; Jana et al., 2020;
Strader et al., 2017). These β82 replacements thereby presumably decrease the rate of Hb turnover
and increase the half-life of the protein (Strader et al., 2017), which may contribute to the mild
polycythemia in humans carrying this substitution (Bardakjian et al., 1985; Moo-Penn et al., 1976).
Blood with an elevated O_2 carrying capacity is typical of most mammalian divers, where it increases
onboard O_2 stores and extends dive times (Ponganis, 2011), but is not observed in extant sirenians
(Farmer et al., 1979; White et al., 1976; Wong et al., 2018). However, any solubility driven increases
in red blood cell Hb concentration resulting from the β82Lys→Asn exchange would have allowed
Steller's sea cows to maintain an elevated rate of tissue O_2 delivery to meet their metabolic
demands during extended underwater foraging intervals. Although this species was presumably
unable to completely submerge (Domning, 2022; Steller, 1751), this conjecture is corroborated by
Steller's account that, "*they keep their heads always under water* [foraging]*, without regard to life
and safety*" (Steller, 1751).

Thermal Sensitivity. The invariant energy change associated with forming the weak covalent bond
between O_2 and the heme iron (i.e. the enthalpy of heme oxygenation; ΔH^{O2}) is exothermic (-59 kJ
mol^{-1} O_2) (Atha and Ackers, 1974), and only moderately opposed by the endothermic solubilization
of O_2 (ΔH^{H2O}; 12.55 kJ mol^{-1} O_2), resulting in an inverse relationship between temperature and Hb–
O_2 affinity. However, the heat of the T→R conformational change ($\Delta H^{T \to R}$), and the oxygenation-
linked dissociation of H^+ (ΔH^{H+}), Cl^- (ΔH^{Cl-}), and DPG (ΔH^{DPG}) may offset this relationship, such that
the overall enthalpy of Hb oxygenation ($\Delta H'$) can become greatly minimized or even endothermic
(Weber and Campbell, 2011; Weber et al., 2010). By facilitating adequate oxygenation of cool
peripheral tissues, Hbs with numerically low $\Delta H'$ values are interpreted to be adaptive for cold-
tolerant, regionally heterothermic mammals. The evolution of this phenotype has predominantly
been attributed to the formation of additional heterotropic ligand binding sites on the protein moiety,
as has previously been demonstrated for the woolly mammoth, *Mammuthus primigenius* (Campbell
et al., 2010b; Weber and Campbell, 2011). Conversely, Steller's sea cow Hb lacks heterotropic

binding of DPG and displays lower Bohr (H^+) and Cl^- effects than all other sirenian Hbs measured
(Fig. 2; Table S1), yet exhibits a ΔH' value (-18.8 kJ mol^{-1} O_2; Fig. 2B) that is close to that of
mammoth Hb (-17.2 kJ mol^{-1} O_2) (Campbell et al., 2010b). This striking convergence largely arises
from the inherently low ΔH of stripped Steller's sea cow Hb (-34.2 kJ mol^{-1} O_2) relative to dugong,
ancestral dugongid, and manatee Hbs (range: -50.1 to -58.6 kJ mol^{-1} O_2) at pH 7.8—where
oxygenation-linked binding of protons is minimal—and indicates that structural differences
modifying the T→R transition largely underlie the low thermal sensitivity of *H. gigas* Hb. Recent
studies have shown that a large positive $ΔH^{T→R}$ may similarly contribute to the low ΔH' of deer
mouse, cow, shrew, and mole Hbs (Campbell et al., 2010a; Campbell et al., 2012; Jensen et al.,
2016; Signore et al., 2012; Weber et al., 2014) suggesting that this potential mechanism of
temperature adaptation may be more widespread than previously appreciated. Our experiments
with the Steller's sea cow β82Asn→Lys mutant further implicate substitutions at this position as a
key factor underlying the inherently low ΔH of the protein, as this modified protein displays a greatly
increased ΔH in the absence of allosteric effectors relative to the wild-type *Hydrodamalis* protein
(Fig. 2B). Interestingly, despite these inherent ΔH differences between the mutant and wild-type
Steller's sea cow Hbs, their ΔH' values are indistinguishable in the presence of allosteric effectors
(Fig. 2B). These data suggest that β82Asn uncouples thermal sensitivity from DPG concentration,
permanently conferring the *H. gigas* protein with a numerically low ΔH' by genetic assimilation while
simultaneously eliminating the energetic cost of DPG production within the red blood cells.

Given both the marked functional changes observed for *H. gigas* Hb and the
correspondingly large ecological and thermal shifts encountered by ancestral hydrodamalines
following their exploitation of the North Pacific in the Miocene (Heritage and Seiffert, 2022), it is
surprising that previous work failed to provide evidence for positive selection or an accelerated
amino acid substitution rate for any globin loci in the Steller's sea cow branch (Signore et al., 2019).
However, this result is not unique to hydrodamalines, as the Hb coding genes of woolly mammoths
and stem penguins also lack statistically significant signatures of positive selection accompanying
their niche transitions despite clear directional changes in their Hb properties (Campbell et al.,
2010b; Signore et al., 2021).

Paleophysiology of Steller's sea cows. The posthumously published behavioral and anatomical
accounts of the last remaining *H. gigas* population by naturalist Georg Wilhelm Steller while
stranded on Bering Island (55°N, 166°E) in 1741/1742 provide a rich tapestry to interpret the
paleophysiology of this colossal marine herbivore. For example, their protective thick bark-like hide
and extensive blubber layer give credence to the extreme nature of their shallow rocky and (during
winter) ice strewn habitat (Le Duc et al., 2022). Here, as Steller (Steller, 1751) remarked, they used
fingerless, bristle-covered forelimbs for support and to shear "*algae and seagrasses from the*

rocks", which they masticated "*not with teeth, which they lack altogether, but with*" large, ridged

keratinized pads located on the upper palate and lower mandible. Although they became visibly

thin during winter when "*their spinous processes can be seen*", Steller (Steller, 1751) noted that

"*(t)hese animals are very voracious, and eat incessantly*" such that their stupendous stomach ("*6*

feet [1.8 m] *long, 5 feet* [1.5 m] *wide*") and enormous intestines—which measured a remarkable

5,958 inches (~151.5 m) from esophagus to anus, equivalent to "*20½ times as long as the whole*

animal"—are constantly "*stuffed with food and seaweed*". The proportionally larger gut (Domning,

2022) is consistent with Steller's sea cow's higher energetic requirements relative to extant

manatees, which, owing to their low metabolic intensities become cold stressed and die if

chronically exposed to water temperatures below 15°C (O'Shea et al., 1985). The inferred reduction

in insulative blubber thickness of *H. gigas* during the winter months would likely have compounded

the rate of heat loss to sub-zero degree Celsius air and water, though may have been compensated

for by arteriovenous anastomoses that regulated blood flow to the skin, and by countercurrent rete

supplying the flippers and tail flukes, the latter of which are well developed in manatees and

presumably other sirenians (Marshall et al., 2022; Rommel and Caplan, 2003). These structures

conserve thermal energy by promoting profound cooling at the appendages and periphery (McCabe

et al., 1978), and presumably underlie the low thermal dependence of Steller's sea cow Hb relative

to those of extant sea cows.

Reductions in blood–O$_2$ affinity accompanying the *H. gigas* β/δ82Lys→Asn substitution is

expected to have further augmented tissue O$_2$ delivery without tangible effects on lung O$_2$ uptake,

thereby helping to fuel thermogenesis and maintain a stable core temperature. In the absence of

UCP1-dependent nonshivering thermogenesis (Gaudry et al., 2017), the latter was presumably

supplemented by a substantive heat increment arising from fermentation and other post-prandial

processes (Marshall et al., 2022). Although the attendant increase in the rate of O$_2$ consumption

would have mandated a reduction in breath-hold endurance—likely reflecting the relatively short

submergence times (4 to 5 minutes) observed by Steller (Steller, 1751)—our results suggest that

this may have been partially counteracted by an elevated blood–O$_2$ carrying capacity that was

potentially coupled to a greater lung volume (Domning, 2022). Underwater foraging times were

presumably further defended by key components of the dive reflex, namely bradycardia and

peripheral vasoconstriction. Indeed, Steller inadvertently was the first to (indirectly) describe this

phenomenon as he observed his crew hunting the animals with spears and knives, "*the blood from*

the wounded back spurted up like a fountain. As long as he kept his head under water the blood

did not flow out, but as soon as he raised his head to breathe the blood leaped forth anew".

A final compelling aspect of Steller's sea cow evolution was their immense size—up to

11,000 kg in mass and 10 m in length—relative to extant sirenians (Domning, 1976). While Steller

does not provide measurements of "*their tender little offspring*", Gerhard Friedrich Müller, who

edited Steller's manuscript prior to publication, noted calves "*weighed 1200 pounds* [544 kg] *and*

upwards" (Mueller, 1761). This value is ~10 to 50 times the mass of new born manatees and

dugongs (~10-50 kg) (Odell, 2009) and is suggestive of rapid prenatal growth during the ~1 year

gestational period indicated by Steller (Steller, 1751). While placental morphology and relative

blood flow are important factors affecting pre-natal growth rates, the efficiency of maternal/pre-natal

gas exchange is also influenced by differences in blood O_2-affinity between the two circulations

(Carter, 2015). During the early stages of mammalian development, O_2 diffusion is optimized via

the expression of embryonic Hb isoforms with high O_2-affinity (Weber et al., 1987). Briefly, the α

and β gene families of mammals possess multiple paralogs, with the 5'-3' linkage order and their

distance from the respective upstream locus control regions dictating the expression pattern of

each locus throughout development (Peterson and Stamatoyannopoulos, 1993). Thus, at two

weeks post-conception, developing human embryos begin expressing genes at the 5' end of the α

(HBZ) and β (HBE) clusters, which are translated into ζ- and ε-globin chains, respectively, to form

Hb Gower I (Fantoni et al., 1981). At week four, expression of the downstream HBA and HBG loci

add α- and γ-chains to the erythrocytes of the developing circulatory system to generate additional

Hb isoforms including HbF ($\alpha_2\gamma_2$) (Hecht et al., 1966). Notably, this pattern of gene expression

switching during development results in the temporal production of Hb isoforms with successively

lower O_2 affinities (i.e., each Hb isoform has a lower O_2 affinity than the protein it replaced), which

facilitates O_2 transfer from maternal to embryonic and fetal blood (Carter, 2015).

In all mammalian lineages examined to date, with the exception of bovid artiodactyls (e.g.,

goats, sheep, and cows) and simian primates, the expression of the above Hb isoforms is thought

to be limited to the embryonic stage of development (Carter, 2015); as such, most mammals

express the same Hb isoform (HbA) during both the fetal and post-natal stages of development.

However, observations suggest that sea cows and proboscideans (elephants) may also express

distinct fetal isoHbs. For example, blood from a 5-month old elephant fetus was shown to contain

two distinct Hb components, although only a single (adult) Hb component is present in 12-month

old fetal and adult blood (Riegel et al., 1967). Likewise, the blood of manatee calves contains a

second isoHb (comprising ~5% of total Hb; Farmer et al., 1979), which moreover appears to exhibit

O_2 binding properties distinct from that of maternal blood (White et al., 1976). It is thus conceivable

that the second Hb component in newborn manatee blood (and potentially other sirenians) arises

from the delayed expression of HBG (which expresses the γ-chain of HbF). Given that the timing

of globin gene expression is determined by its distance from the locus control region (Peterson and

Stamatoyannopoulos, 1993), the possible attenuation of HBG expression in sirenians HBG relative

to elephants is supported by synteny comparisons of the β-globin gene cluster, as the HBG locus

of sirenians is further downstream than the same locus is in the elephant cluster (see Fig. 1B of ref

4). If expression of the sirenian HBG locus is developmentally delayed to form a discrete isoHb in

fetal blood, it would be expected to display P_{50} and cooperativity (n_{50}) values that fall between
Gower I and HbA, and a response to pH that is similar to the latter.

To test this hypothesis and better understand the maternal/pre-natal gas exchange
strategy of sirenians, we thus expressed recombinant Hbs corresponding to Steller's sea cow
Gower I and HbF and dugong HbF (whose γ-chain differs from *H. gigas* γ at four positions (Signore
et al., 2019); Fig. S6), and measured their O_2 binding properties and response to allosteric effectors.
As expected, the P_{50} of *H. gigas* Gower I in the combined presence of Cl^- and DPG is markedly
less than adult Steller's sea cow Hb (3.1 vs. 15.2 mm Hg, respectively) (Fig. 4A). Similarly, the Hb–
O_2 affinity of Steller's sea cow and dugong HbF (P_{50} of 0.56 and 1.2 mm Hg, respectively) are
substantially higher than that of their respective adult counterparts and, unexpectedly, also higher
than that of Steller's sea cow Gower I (Fig. 4A). In line with the embryonic expressed Hb isoforms
of other mammals (Brittain, 2002; Weber et al., 1987), the Bohr and cooperativity coefficients of the
sirenian Gower I and HbF proteins were also substantially lower than that of the post-natal (HbA)
isoform (Fig. 4A, Table S2). Accordingly, their functional properties are consistent with the
embryonic (but not fetal) Hbs of other mammalian species. Although it remains possible that
expression of these isoforms lingers into late fetal development, the upstream HBG transcriptional
control motif ('CACCC') crucial for suppression of human HBB gene expression during the fetal
stage (Perez-Stable and Costantini, 1990) is mutated in both dugongs and Steller's sea cows (but
not manatees or elephants; Fig. S7). Consequently, the primary (if not sole) Hb isoform expressed
within the both the fetal and post-natal circulation of dugongids is almost certainly HbA. Intriguingly,
however, Steller's sea cow HbF exhibits a distinctly higher O_2 affinity but lower cooperativity than
dugong HbF, traits that are likely attributed to two exceedingly rare γ-chain amino acid
replacements positioned within the interior of the protein (γ38Thr→Ile and γ101Glu→Asp; Figs. 4B-
E and S6). Briefly, the central cavity γ101Glu→Asp replacement alters the highly conserved sliding
interface between the $\alpha_1\gamma_2$ dimer subunits by forming a hydrogen bond with γ104Arg (Fig. 4B) and
has been shown to increase the intrinsic affinity of human Hb Potomac (β101Glu→Asp) (Charache
et al., 1978; Shih et al., 1985). Residue γ38 is also potentially functionally relevant as it is located
along the $\alpha_2\gamma_1$ sliding interface and is in contact the distal heme (Fig. 4D) (Ropero et al., 2006). Of
note, however, the mRNA capping site of *H. gigas* HbF exhibits an A→G transversion mutation
(Fig. S7) that has been shown to lower transcript levels of human HBB by twofold (Meyers et al.,
1986). It thus remains unknown if *H. gigas* HbF exhibited a similar downregulation and hence to
what degree these γ-chain replacements may have altered O_2 transfer to the Steller's sea cow
embryo through the amniotic fluid prior to placental development.

Based on the above considerations, the apparently unique DPG insensitive phenotype of
H. gigas HbA is particularly noteworthy owing to its potential impact on maternal/fetal O_2 exchange.
Presumably to assist in this process, fetal blood cells expressing HbA contain only trace amounts

of DPG hence conferring fetal blood with an elevated O_2 affinity relative to that of the maternal
circulation (compare the dashed vs. solid red lines in Fig. 4A as an example) in the vast majority of
mammalian species (Bunn, 1980; Carter, 2015). By contrast, and owing to the inability of Steller's
sea cows Hb to respond to DPG in either the fetal or adult circulations, this species would represent
a rare example (feloids and eastern moles are others) in which fetal and maternal blood have the
same (albeit relatively low) O_2 affinity. However, placental O_2 delivery in these species will be
defended by the well-known double Bohr effect, whereby CO_2 transport from the fetal to maternal
circulation increases blood O_2 affinity in the former while lowering it in the latter (Carter, 2015).
More importantly, the evolved reduction in blood O_2 affinity of Steller's sea cows would have
stipulated that equilibrium between umbilical and uterine blood was reached at a higher PO_2, a
condition that is expected to substantially improve O_2 delivery to the fetal circulation (Carter, 2015).
The lower fetal blood O_2 affinity (relative to manatees and dugongs) and concomitant higher fetal
blood-to-tissue PO_2 gradients are further expected to have augmented O_2 delivery to the
developing tissues of this species. As such, these attributes, together with increases in Hb
solubility/reduced susceptibility to oxidative damage arising from β/δ82Lys→Asn that conceivably
also elevated the O_2 carrying capacity of fetal blood, may have been important contributors to the
enhanced fetal growth rate of these immense sirenians. The resulting increase in thermal inertia
and relatively low surface-area-to-volume ratio following birth, together with an adaptively reduced
Hb thermal sensitivity and thick 'bark-like' skin arising from inactivation of lipoxygenase genes (Le
Duc et al., 2022), were presumably central components of Steller's sea cows successful
exploitation of the harsh sub-Arctic marine environments of the North Pacific.

Materials and Methods

Sequence Collection and Analyses. The pre-natal (HBZ-T1, HBE, and HBG) and adult-
expressed Hb genes (HBA and HBB/HBD) of the Florida manatee, dugong, and Steller's sea cow,
and the most recent common ancestor shared by Steller's sea cow and the dugong ('ancestral
dugongid') have previously been determined (Signore et al., 2019). As the *H. gigas* β/δ82Lys→Asn
exchange is not known to occur in any living species, we mined recently deposited genomes for 13
additional Steller's sea cows (PRJNA484555, PRJEB43951) to test for the prevalence of this
replacement in the population. Briefly, we first searched SRA files of each specimen using the
megablast function against a previously determined *H. gigas HBB/HBD* gene sequence (GenBank
accession #: MK562081). All hits were then downloaded, trimmed of adapters and low-quality
regions using BBDuk (Joint Genome Institute), and assembled to *H. gigas HBB/HBD* using
Geneious Prime 2019 software (Biomatters Ltd, Auckland, New Zealand). Assemblies generated

using genome reads that were not pre-treated with uracil-DNA glycosylase and endonuclease VIII

to reduce C→T and G→A damage artifacts (Le Duc et al., 2022) were examined to ensure these

deamination artifacts did not affect the consensus sequences.

Construction of Recombinant Hb Expression Vectors. Coding sequences for Steller's sea cow

Gower I ($\zeta_2\varepsilon_2$), dugong and H. gigas HbF ($\alpha_2\gamma_2$), and the above four HbA ($\alpha_2\beta/\delta_2$) proteins were

optimized for expression in *E. coli* and synthesized *in vitro* by GenScript (Piscataway, NJ). The

resulting gene cassettes were digested with restriction enzymes and tandemly ligated into a custom

Hb expression vector (Natarajan et al., 2011) using a New England BioLabs Quick Ligation Kit as

recommended by the manufacturer. Chemically competent JM109 (DE3) *E. coli* (Promega) were

prepared using a Z-Competent *E. coli* Transformation Kit and Buffer Set (Zymo Research). We also

prepared a *H. gigas* β/δ82Asn→Lys Hb mutant via site-directed mutagenesis on the Steller's sea

cow Hb expression vector by whole plasmid amplification using mutagenic primers and Phusion

High-Fidelity DNA Polymerase (New England BioLabs), phosphorylation with T4 Polynucleotide

Kinase (New England BioLabs), and circularization with an NEB Quick Ligation Kit (New England

BioLabs). All site-directed mutagenesis steps were performed using the manufacture's

recommended protocol.

Hb expression vectors were co-transformed into JM109 (DE3) chemically competent *E.*

coli alongside a plasmid expressing methionine aminopeptidase (Natarajan et al., 2011), plated on

LB agar containing ampicillin (100 µg/ml) and kanamycin (50 µg/ml), and incubated for 16 hours at

37°C. A single colony from each transformation was cultured in 50 ml of 2xYT broth for 16 hours at

37°C while shaking at 200 rpm. Post incubation, 5 ml of the culture was pelleted by centrifugation

and plasmid DNA was isolated using a GeneJET Plasmid Miniprep Kit (Thermo Scientific). The

plasmid sequence was verified using BigDye 3.1 sequencing chemistry and an ABI3130 Genetic

Analyzer. The remainder of the culture was supplemented with glycerol to a final concentration of

10%, divided into 25 ml aliquots and stored at -80°C until needed for expression.

Expression and Purification of Recombinant Hb. 25 ml of starter culture (above) was added to

1250 ml of TB media containing ampicillin (100 µg/ul) and kanamycin (50 µg/ul) and distributed

evenly amongst five 1 L Erlenmeyer flasks. Cultures were grown at 37°C while shaking at 200 rpm

until the absorbance at 600 nm reached 0.6-0.8. Hb expression was induced by supplementing the

media with 0.2 mM isopropyl β-D-1-thiogalactopyranoside, 50 µg/ml of hemin and 20 g/L of glucose

and the culture was incubated at 28°C for 16 hours while shaking at 200 rpm. Once expression had

completed, dissolved O_2 was removed by adding sodium dithionite (1 mg/ml) to the culture, which

was promptly saturated with CO for 15 minutes. Bacterial cells were then pelleted by centrifugation

and Hb purified by ion exchange chromatography according to Natarajan et al. (Natarajan et al.,
2011)

It should be noted that the β82Asn residue of human Hb Providence is relatively uncommon
in that it slowly undergoes post-translational deamidation *in vivo* to form aspartic acid, with the latter
residue (β82Asp) comprising ~67-75% in mature mixed blood (Bardakjian et al., 1985; Perutz et
al., 1980). While it is unknown to what degree *H. gigas* β/δ82Asn was catalyzed into Asp in nature,
O_2 binding data (see below) of this species was collected from freshly purified recombinant samples
for which only one peak—presumably β/δ82Asn—was resolved during chromatography (data not
shown). Additionally, this reaction is dependent on the local protein environment (Robinson, 2002),
specifically two nearby residues β143His and β83Gly (Perutz et al., 1980). Importantly, the latter
residue was replaced by β/δ83Ser on the Steller's sea cow branch (Figs. 1B and 3B), which is
expected to slow (but not stop) the rate of deamidation (Robinson, 2002). Regardless, since the
two Hb Providence isoforms have similar O_2 affinities and functional properties (Bardakjian et al.,
1985; Bonaventura et al., 1976; Charache et al., 1977) it is unlikely that presence of β/δ82Asp in
Steller's sea cow blood would meaningfully alter the results and interpretations presented herein.

Functional Analyses of Hbs. O_2-equilibrium curves for HbA containing solutions (0.25–1.0 mM
heme in 0.1 M HEPES/0.0005 M EDTA buffers) were measured at 25 and 37°C using the thin film
technique (Weber, 1992), while curves for the three pre-natal Hb isoforms (0.25 mM heme in 0.1
M HEPES/0.0005 M EDTA) were measured at 37°C using a multi-cuvette tonometer cell described
by Lilly et al. (Lilly et al., 2013). Hb solutions varied in their pH (range: 6.8 to 7.9), chloride
concentration (0 or 0.1 M KCl), and organic phosphate concentration (0 or 2-fold molar excess of
DPG relative to tetrameric Hb concentrations) in order to test the influence of these cofactors on
Hb function. Each Hb solution was sequentially equilibrated with gas mixtures of three to five
different oxygen tensions (PO_2) that result in Hb–O_2 saturations between 30 to 70%. Hill plots
(log[fractional saturation/[1-fractional saturation]] vs logPO_2) constructed from these measurements
were used to determine the PO_2 (P_{50}) and the cooperativity coefficient (n_{50}) at half saturation, from
the χ-intercept and slope of these plots, respectively. By this method, the r^2 determination
coefficients for the fitted curves exceed 0.995 and the standard errors (SEM) are less than 3% of
the P_{50} and n_{50} values (Weber et al., 2014). A linear regression was fit to plots of logP_{50} vs. pH, and
the resulting equation was used to estimate P_{50}, Cl^- effect, and DPG effect values (± SE of the
regression estimate) at pH 7.20 for HbA samples, and pH 7.10 for Gower I and HbF samples (to
account for the lower pH of pre-natal blood). The slope of these plots (ΔlogP_{50}/ΔpH) represented
the Bohr effect. P_{50} values at 25 and 37°C were used to assess the thermal sensitivity of the Hbs
by calculating the apparent enthalpy of oxygenation using the van't Hoff isochore:

$$\Delta H = 2.303R \times \Delta \log P_{50} \times (1/T_1 - 1/T_2)^{-1}$$

where R is the universal gas constant and T_1 and T_2 are the absolute temperatures (°K) at which

the P_{50} values were measured. All ΔH values were corrected for the heat of O_2 solubilization (12.55

kJ mol^{-1} O_2).

Solubility assay. Ammonium sulfate was added to Hb solutions (0.074 ± 0.004 mM Hb_4) to

generate final concentrations that ranged from 0 to 3.5 M. These solutions were incubated for 60

minutes at 37°C and the remaining soluble Hb was measured via Drabkin's reagent, according to

the manufacturer's instructions (Sigma-Aldrich).

Homology modelling. To assess the structural effect of the *H. gigas* specific β/δ replacements on

the DPG binding site, homology models of ancestral dugongid and Steller's sea cow Hb were

constructed using the SWISS-MODEL server (62) using the three-dimensional human deoxy

structure with DPG bound (PDB: 1B86) as a template (30). The sequence conservation of amino

acid residues implicated in DPG binding were calculated by the ConSurf Server (63) from a

subsample of 51 mammalian beta-type hemoglobin chains downloaded from GenBank (Table S3).

Homology models were visualized with UCSF Chimera (64). To assess the structural differences

between *H. gigas* and *D. dugon* HbF ($\alpha_2\gamma_2$), homology models of these proteins were created as

above, but with human deoxy HbF used as template (PDB: 4MQJ).

Acknowledgments

This manuscript is dedicated to the memory of our dear friend Joseph (Joe) Bonaventura for his

pioneering work on the human Hb Providence Asn/Asp proteins. We thank Mike Gaudry, Diana

Hanna, and Elin Ellebæk Petersen for technical assistance, Chandrasekhar Natarajan for providing

us with a hemoglobin expression plasmid, and Jay Storz for constructive feedback on an earlier

version of this manuscript. Authorization to use paintings by Carl Buell was kindly provided by John

Gatesy. This study was supported by NSERC (Canada) Discovery and Accelerator Supplement

Grants (RGPIN/238838-2011, RGPIN/412336-2011, and RGPIN/06562-2016 to K.L.C;

RGPIN/261924-2013 and RGPIN/446005-2013 to C.J.B.), an NSERC Postgraduate Scholarship

(A.V.S.), the Faculty of Science and Technology, Aarhus University (R.E.W.), and the Independent

Research Fund Denmark (A.F.; Danmarks Frie Forskningsråd DFF-4181-00094).

529
530
531 **References**
532

Abraham, B., Hicks, W., Jia, Y., Baek, J. H., Miller, J. L. and Alayash, A. I. (2011). Isolated Hb
Providence β82Asn and β82Asp Fractions are more stable than native HbA0 under
oxidative stress conditions. *Biochemistry* 50, 9752–9766.

Atha, D. H. and Ackers, G. K. (1974). Calorimetric determination of the heat of oxygenation of
human hemoglobin as a function of pH and the extent of reaction. *Biochemistry* 13,
2376–2382.

Bardakjian, J., Leclerc, L., Blouquit, Y., Oules, O., Rafaillat, D., Arous, N., Bohn, B., Poyart, C.,
Rosa, J. and Galacteros, F. (1985). A new case of Hemoglobin Providence (α2 β2 82
(Ef6) Lys → Asn or Asp) discovered in a French caucasian family. Structural and
functional studies. *Hemoglobin* 9, 333–348.

Blessing, M. H. (1972). Studies on the concentration of myoglobin in the sea-cow and porpoise.
Comp. Biochem. Physiol. A 41, 475–480.

Bonaventura, J., Bonaventura, C., Sullivan, B., Ferruzzi, G., McCurdy, P. R., Fox, J. and Moo-
Penn, W. F. (1976). Hemoglobin providence. Functional consequences of two alterations
of the 2,3-diphosphoglycerate binding site at position beta 82. *J. Biol. Chem.* 251, 7563–
7571.

Brittain, T. (2002). Molecular aspects of embryonic hemoglobin function. *Mol. Aspects Med.* 23,
293–342.

Bunn, H. F. (1980). Regulation of Hemoglobin Function in Mammals. *Amer. Zool.* 20, 199–211.

Campbell, K. L. and Hofreiter, M. (2015). Resurrecting phenotypes from ancient DNA sequences:
promises and perspectives. *Can. J. Zool.*

Campbell, K. L., Storz, J. F., Signore, A. V., Moriyama, H., Catania, K. C., Payson, A. P.,
Bonaventura, J., Stetefeld, J. and Weber, R. E. (2010a). Molecular basis of a novel
adaptation to hypoxic-hypercapnia in a strictly fossorial mole. *BMC Evol. Biol.* 10, 214.

Campbell, K. L., Roberts, J. E. E., Watson, L. N., Stetefeld, J., Sloan, A. M., Signore, A. V.,
Howatt, J. W., Tame, J. R. H., Rohland, N., Shen, T.-J., et al. (2010b). Substitutions in
woolly mammoth hemoglobin confer biochemical properties adaptive for cold tolerance.
Nat. Genet. 42, 536–540.

Campbell, K. L., Signore, A. V., Harada, M. and Weber, R. E. (2012). Molecular and
physicochemical characterization of hemoglobin from the high-altitude Taiwanese brown-
toothed shrew (Episoriculus fumidus). *J. Comp. Physiol. B* 182, 821–829.

Carter, A. M. (2015). Placental Gas Exchange and the Oxygen Supply to the Fetus. *Compr.
Physiol.* 5, 1381–1403.

Charache, S., Fox, J., McCurdy, P., Kazazian, H., Winslow, R., Hathaway, P., Beneden, R. van
and Jessop, M. (1977). Postsynthetic deamidation of hemoglobin Providence (beta 82
Lys replaced by Asn, Asp) and its effect on oxygen transport. *J. Clin. Invest.* 59, 652–
658.

Charache, S., Jacobson, R., Brimhall, B., Murphy, E. A., Hathaway, P., Winslow, R., Jones, R.,
Rath, C. and Simkovich, J. (1978). Hb Potomac (101 Glu replaced by Asp): speculations
on placental oxygen transport in carriers of high-affinity hemoglobins. *Blood* 51, 331–338.

Domning, D. P. (1976). An ecological model for late tertiary Sirenian evolution in the North Pacific
Ocean. *Systemat. Biol.* 25, 352–362.

Domning, D. P. (2022). What Can We Infer About the Behavior of Extinct Sirenians? In *Ethology*
and Behavioral Ecology of Sirenia (ed. Marsh, H.), pp. 1–17. Cham: Springer
International Publishing.

Duhm, J., Deuticke, B. and Gerlach, E. (1968). Metabolism of 2,3-diphosphoglycerate and
glycolysis in human red blood cells under the influence of dipyridamole and inorganic
sulfur compounds. *Biochimica et Biophysica Acta* 170, 452–454.

Fantoni, A., Farace, M. G. and Gambari, R. (1981). Embryonic hemoglobins in man and other
mammals. *Blood* 57, 623–633.

Farmer, M., Weber, R. E., Bonaventura, J., Best, R. C. and Domning, D. (1979). Functional
properties of hemoglobin and whole blood in an aquatic mammal, the Amazonian
manatee (*Trichechus inunguis*). *Comp. Biochem. Physiol. A* 62, 231–238.

Fronticelli, C., Bucci, E. and Razynska, A. (1988). Modulation of oxygen affinity in hemoglobin by
solvent components: Interaction of bovine hemoglobin with 2,3-diphosphoglycerate and
monatomic anions. *J. Mol. Biol.* 202, 343–348.

Gallivan, G. J., Kanwisher, J. W. and Best, R. C. (1986). Heart rates and gas exchange in the
Amazonian manatee (*Trichechus inunguis*) in relation to diving. *J. Comp. Physiol. B* 156,
415–423.

Gaudry, M. J., Jastroch, M., Treberg, J. R., Hofreiter, M., Paijmans, J. L. A., Starrett, J., Wales,
N., Signore, A. V., Springer, M. S. and Campbell, K. L. (2017). Inactivation of
thermogenic UCP1 as a historical contingency in multiple placental mammal clades. *Sci.*
Adv. 3, e1602878.

Hecht, F., Motulsky, A. G., Lemire, R. J. and Shepard, T. E. (1966). Predominance of hemoglobin
Gower 1 in early human embryonic development. *Science*.

Heritage, S. and Seiffert, E. R. (2022). Total evidence time-scaled phylogenetic and
biogeographic models for the evolution of sea cows (Sirenia, Afrotheria). *PeerJ* 10,
e13886.

Ikkala, E., Koskela, J., Pikkarainen, P., Rahiala, E. L., El-Hazmi, M. A., Nagai, K., Lang, A. and
Lehmann, H. (1976). Hb Helsinki: A variant with a high oxygen affinity and a substitution
at a 2, 3-DPG binding site (β82 [EF6] Lys→ Met). *Acta. Haematol.* 56, 257–275.

Jana, S., Strader, M. B. and Alayash, A. I. (2020). The Providence Mutation (βK82D) in Human
Hemoglobin Substantially Reduces βCysteine 93 Oxidation and Oxidative Stress in
Endothelial Cells. *Int. J. Mol. Sci.* 21, 9453.

Janecka, J. E., Nielsen, S. S. E., Andersen, S. D., Hoffmann, F. G., Weber, R. E., Anderson, T.,
Storz, J. F. and Fago, A. (2015). Genetically based low oxygen affinities of felid
hemoglobins: lack of biochemical adaptation to high-altitude hypoxia in the snow leopard.
J. Exp. Biol. 218, 2402–2409.

Jensen, B., Storz, J. F. and Fago, A. (2016). Bohr effect and temperature sensitivity of
hemoglobins from highland and lowland deer mice. *Comp. Biochem. Physiol. A* 195, 10–
14.

Kauffman, K. J., Pajerowski, J. D., Jamshidi, N., Palsson, B. O. and Edwards, J. S. (2002).
Description and analysis of metabolic connectivity and dynamics in the human red blood
cell. *Biophys. J.* 83, 646–662.

Kay, F. R. (1977). 2,3-diphosphoglycerate, blood oxygen dissociation and the biology of
mammals. *Comp. Bio. Physiol. A* 57, 309–316.

Le Duc, D., Velluva, A., Cassatt-Johnstone, M., Olsen, R.-A., Baleka, S., Lin, C.-C., Lemke, J. R.,
Southon, J. R., Burdin, A., Wang, M.-S., et al. (2022). Genomic basis for skin phenotype
and cold adaptation in the extinct Steller's sea cow. *Sci. Adv.* 8, eabl6496.

Lilly, L. E., Blinebry, S. K., Viscardi, C. M., Perez, L., Bonaventura, J. and McMahon, T. J. (2013).
Parallel assay of oxygen equilibria of hemoglobin. *Anal. Biochem.* 441,
10.1016/j.ab.2013.06.010.

Marsh, H., O'Shea, T. J. and Reynolds III, J. E. (2011). *Ecology and conservation of the Sirenia:*
dugongs and manatees.

Marshall, C. D., Sarko, D. K. and Reep, R. L. (2022). Morphological and sensory innovations for
an aquatic lifestyle. In *Ethology and Behavioral Ecology of Sirenia* (ed. Marsh, H.), pp.
19–65. Springer International Publishing.

McCabe, M., Hamilton, R. and Marsh, H. (1978). Some studies on the oxygen affinity of
haemoglobin from the dugong. *Comp. Biochem. Physiol. A* 61,19–22

Meyers, R. M., Tilly, K. and Maniatis, T. (1986). Fine structure genetic analysis of a beta-globin
promoter. *Science* 232, 613–618.

Mirceta, S., Signore, A. V., Burns, J. M., Cossins, A. R., Campbell, K. L. and Berenbrink, M.
(2013). Evolution of mammalian diving capacity traced by myoglobin net surface charge.
Science 340, 1234192.

Moo-Penn, W. F., Jue, D. L., Bechtel, K. C., Johnson, M. H. and Schmidt, R. M. (1976).
Hemoglobin Providence. A human hemoglobin variant occurring in two forms in vivo. *J.*
Biol. Chem. 251, 7557–7562.

Mueller, G. F. (1761). *Voyages from Asia to America, for Completing the Discoveries of the North*
West Coast of America. T. Jefferys.

Natarajan, C., Jiang, X., Fago, A., Weber, R. E., Moriyama, H. and Storz, J. F. (2011). Expression
and purification of recombinant hemoglobin in *Escherichia coli*. *PLoS ONE* 6, e20176.

Natarajan, C., Signore, A. V., Bautista, N. M., Hoffmann, F. G., Tame, J. R. H., Fago, A. and
Storz, J. F. (2023). Evolution and molecular basis of a novel allosteric property of
crocodilian hemoglobin. *Curr. Biol.* 33, 98-108.e4.

Odell, D. K. (2009). Sirenian Life History. In *Encyclopedia of Marine Mammals (Second Edition)*
(ed. Perrin, W. F.), Würsig, B.), and Thewissen, J. G. M.), pp. 1019–1021. London:
Academic Press.

O'Shea, T. J., Beck, C. A., Bonde, R. K., Kochman, H. I. and Odell, D. K. (1985). An analysis of
manatee mortality patterns in Florida, 1976-81. *J. Wildl. Manag.* 49, 1–11.

Perez-Stable, C. and Costantini, F. (1990). Roles of fetal G gamma-globin promoter elements
and the adult beta-globin 3' enhancer in the stage-specific expression of globin genes.
Mol. Cell. Biol. 10, 1116–25.

Perutz, M. F., Fogg, J. H. and Fox, J. A. (1980). Mechanism of deamidation of haemoglobin
providence Asn. *J. Mol. Biol.* 138, 669–670.

Perutz, M. F., Fermi, G., Poyart, C., Pagnier, J. and Kister, J. (1993). A novel allosteric
mechanism in haemoglobin. Structure of bovine deoxyhaemoglobin, absence of specific
chloride-binding sites and origin of the chloride-linked Bohr effect in bovine and human
haemoglobin. *J. Mol. Biol.* 233, 536–545.

Peterson, K. R. and Stamatoyannopoulos, G. (1993). Role of gene order in developmental control
of human gamma-and beta-globin gene expression. *Mol. Cell.* 13, 4836–4843.

Ponganis, P. J. (2011). Diving mammals. *Compr. Physiol.* 1, 447–465.

Rapoport, S. and Luebering, J. (1950). *The formation of 2,3-diphosphoglycerate in rabbit*
erythrocytes: the existence of a diphosphoglycerate mutase. J. Biol. Chem. 183, 507-516

Richard, V., Dodson, G. G. and Mauguen, Y. (1993). Human deoxyhaemoglobin-2, 3-
diphosphoglycerate complex low-salt structure at 2· 5 Å resolution. *J. Mol. Biol.* 233,
270–274.

Riegel, K., Bartels, H., Buss, I. O. and Wright, P. G. (1967). Comparative studies of the
respiratory functions of mammalian blood IV. Fetal and adult African elephant blood.
Respir. Physiol. 2, 182-195

Robinson, N. E. (2002). Protein deamidation. *Proc. Nat. Acad. Sci.* 99, 5283–5288.

Rommel, S. A. and Caplan, H. (2003). Vascular adaptations for heat conservation in the tail of
Florida manatees (*Trichechus manatus latirostris*). *J. Anat.* 202, 343–353.

Ropero, P., Fernández-Lago, C., Villegas, A., Polo, M., Mateo, M., Mora, A. and González, F. A.
(2006). Hb LA Coruña [β38(C4)Thr→Ile]: A New Hemoglobin variant leading to familial
polycythemia. *LHEM* 30, 379–383.

Schmidt-Neilsen, K. and Larimer, J. L. (1958). Oxygen dissociation curves of mammalian blood in
relation to body size. *Am. J. Physiol.* 195, 424–428.

Scholander, P. F. and Irving, L. (1941). Experimental investigations on the respiration and diving
of the florida manatee. *J. Cell. Comp. Physiol.* 17, 169–191.

Sharko, F. S., Rastorguev, S. M., Boulygina, E. S., Tsygankova, S. V., Ibragimova, A. S.,
Tikhonov, A. N. and Nedoluzhko, A. V. (2019). Molecular phylogeny of the extinct
Steller's sea cow and other Sirenia species based on their complete mitochondrial
genomes. *Genomics* 111, 1543–1546.

Sharko, F. S., Boulygina, E. S., Tsygankova, S. V., Slobodova, N. V., Alekseev, D. A.,
Krasivskaya, A. A., Rastorguev, S. M., Tikhonov, A. N. and Nedoluzhko, A. V. (2021).
Steller's sea cow genome suggests this species began going extinct before the arrival of
Paleolithic humans. *Nat. Commun.* 12, 2215.

Shih, D. T., Jones, R. T., Imai, K. and Tyuma, I. (1985). Involvement of Glu G3(101)beta in the
691   function of hemoglobin. Comparative O2 equilibrium studies of human mutant
692   hemoglobins. *J. Biol. Chem.* 260, 5919–5924.

693   Signore, A. V., Stetefeld, J., Weber, R. E. and Campbell, K. L. (2012). Origin and mechanism of
694   thermal insensitivity in mole hemoglobins: a test of the 'additional' chloride binding site
695   hypothesis. *J. Exp. Biol.* 215, 518–525.

696   Signore, A. V., Paijmans, J. L. A., Hofreiter, M., Fago, A., Weber, R. E., Springer, M. S. and
697   Campbell, K. L. (2019). Emergence of a chimeric globin pseudogene and increased
698   hemoglobin oxygen affinity underlie the evolution of aquatic specializations in Sirenia.
699   *Mol. Biol. Evol.* 36, 1134–1147.

700   Signore, A. V., Tift, M. S., Hoffmann, F. G., Schmitt, Todd. L., Moriyama, H. and Storz, J. F.
701   (2021). Evolved increases in hemoglobin-oxygen affinity and the Bohr effect coincided
702   with the aquatic specialization of penguins. *Proc. Nat. Acad. Sci.* 118, e2023936118.

703   Springer, M. S., Signore, A. V., Paijmans, J. L. A., Velez-Juarbe, J., Domning, D. P., Bauer, C. E.,
704   He, K., Crerar, L., Campos, P. F., Murphy, W. J., et al. (2015). Interordinal gene capture,
705   the phylogenetic position of Steller's sea cow based on molecular and morphological
706   data, and the macroevolutionary history of Sirenia. *Mol. Phylogenet. Evol.* 91, 178–193.

707   Stejneger, L. (1887). How the great northern sea-cow (Rytina) became exterminated. *Am. Nat.*
708   21, 1047–1054.

709   Steller, G. W. (1751). *De bestiis marinis [The beasts of the sea]. Novi Commentarii Acad. Sci.*
710   *Imp. Petropoli.*

711   Storz, J. F. (2016). Causes of molecular convergence and parallelism in protein evolution. *Nat.*
712   *Rev.* 17, 239–250.

713   Strader, M. B., Bangle, R., Parker Siburt, C. J., Varnado, C. L., Soman, J., Benitez Cardenas, A.
714   S., Samuel, P. P., Singleton, E. W., Crumbliss, A. L., Olson, J. S., et al. (2017).
715   Engineering oxidative stability in human hemoglobin based on the Hb providence
716   (βK82D) mutation and genetic cross-linking. *Biochem. J.* 474, 4171–4192.

717   Sugihara, J., Imamura, T., Nagafuchi, S., Bonaventura, J., Bonaventura, C. and Cashon, R.
718   (1985). Hemoglobin Rahere, a human hemoglobin variant with amino acid substitution at
719   the 2,3-diphosphoglycerate binding site. Functional consequences of the alteration and
720   effects of bezafibrate on the oxygen bindings. *J. Clin. Invest.* 76, 1169–1173.

721   Weber, R. E. (1992). Use of ionic and zwitterionic (Tris/BisTris and HEPES) buffers in studies on
722   hemoglobin function. *J. Appl. Physiol.* 72, 1611–1615.

723   Weber, R. E. and Campbell, K. L. (2011). Temperature dependence of haemoglobin-oxygen
724   affinity in heterothermic vertebrates: mechanisms and biological significance. *Acta*
725   *Physiologica* 202, 549–562.

726   Weber, R. E., Kleinschmidt, T. and Braunitzer, G. (1987). Embryonic pig hemoglobins Gower I
727   (zeta 2 epsilon 2), Gower II (alpha 2 epsilon 2), Heide I (zeta 2 theta 2) and Heide II
728   (alpha 2 theta 2): oxygen-binding functions related to structure and embryonic oxygen
729   supply. *Respir. Physiol.* 69, 347–357.

Weber, R. E., Campbell, K. L., Fago, A., Malte, H. and Jensen, F. B. (2010). ATP-induced
temperature independence of hemoglobin-O_2 affinity in heterothermic billfish. *J. Exp. Biol.*
213, 1579–1585.

Weber, R. E., Fago, A. and Campbell, K. L. (2014). Enthalpic partitioning of the reduced
temperature sensitivity of O2 binding in bovine hemoglobin. *Comp. Biochem. Physiol. A*
176, 20–25.

Weickert, M. J., Pagratis, M., Glascock, C. B. and Blackmore, R. (1999). A mutation that
improves soluble recombinant hemoglobin accumulation in *Escherichia coli* in heme
excess. *App. Environ. Micro.* 65, 640–647.

White, J. R., Harkness, D. R., Isaacks, R. E. and Duffield, D. A. (1976). Some studies on blood of
the Florida manatee, *Trichechus manatus latirostris. Comp. Biochem. Physiol. A* 55, 413–
417.

Wong, A. W., Lanyon, J. M., Sneath, H. L., Leggatt, G. R. and Woolford, L. (2018). Comparison of
i-STAT® with Traditional Laboratory Analysers in the Measurement of Blood Analytes
from Field Captured Dugongs (Dugong dugon). *Aqua. Mamm.* 44, 19–31.

Figures and Tables
747

749
750

Figure 1. Evolution of notable morphological and genetic attributes within Sirenia. A) Phenotypic innovations contributing to the unique biology of Steller's sea cows are mapped along the sirenian phylogeny (red bars) with the underlying genetic causes shown in brackets where known. Note that the red bars do not represent the dating of these traits and that their placement order is arbitrary. Divergence dates in millions of years (Mya) are based on Springer et al. (2015) and Heritage and Seiffert (2022). The ancestral dugongid (Anc. dugongid) is represented by a late-Oligocene *Metaxytherium spp.* Images by Carl Buell and are used with permission. B) Partial nucleotide alignment of the sirenian β/δ-globin gene encoding the central region of the 2,3-diphosphoglycerate binding pocket of hemoglobin; corresponding amino acid residues (bolded) are provided below each sequence. The G→C nucleotide mutation underlying the otherwise invariant amino acid substitution (β/δ82Lys→Asn; K→N) of Steller's sea cow (*Hydrodamalis gigas*) hemoglobin is apparent in all 16 individuals for which sequence data is available. Values in brackets next to each *H. gigas* specimen represent the depth of sequence coverage for this nucleotide. Dots represent sequence identity with the Florida manatee (*Trichechus manatus*).

Figure 2. Biochemical properties of hemoglobins (Hbs) from manatee (*Trichechus manatus*), dugong (*Dugong dugon*), ancestral dugongid (Anc. dugongid), Steller's sea cow β/δ82Asn→Lys mutant (*H. gigas* β82K), and wild-type Steller's sea cow (*Hydrodamalis gigas*). All values were measured at 37°C and corrected to pH 7.2 (±SE of the regression estimate). A) Oxygen tensions at half O_2 saturation (P_{50}) in the absence (stripped) and presence of allosteric cofactors (2-fold molar excess of 2,3-diphosphoglycerate (DPG) and 0.1 M KCl). B) The enthalpy of oxygenation (ΔH) between 25 and 37°C in stripped Hb and in the presence of allosteric cofactors (2-fold molar excess DPG and 0.1 M KCl). C) The effect of DPG on sirenian Hbs determined from $logP_{50}^{(0.5\,mM\,DPG)}$ − $logP_{50}^{(stripped)}$. D) The effect of chloride on sirenian Hbs determined from $logP_{50}^{(0.1\,M\,KCl)}$ − $logP_{50}^{(stripped)}$. E) The Bohr effect of sirenian Hbs in the presence of allosteric cofactors (2-fold molar excess DPG and 0.1 M KCl), as calculated from $\Delta logP_{50}/\Delta pH$ over the pH range 6.9 and 7.8. F). The relative solubility of sirenian Hbs is denoted by the percentage of Hb protein precipitated by the addition of 3 M ammonium sulfate.

Variable ☐ Conserved

H. gigas derived ▦ Shared ancestral ▦

Figure 3. Homology models of the DPG binding site in ancestral dugongid and Steller's sea cow (*Hydrodamalis gigas*) hemoglobin. A) Model of the ancestral dugongid DPG binding site. Amino acids are colored according to the degree of sequence conservation. Notably, β/δ82Lys shows the highest level of sequence conservation, as it is able to bind to multiple sites on the DPG molecule (indicated by dashed pink lines), whereas β/δ2His is only able to directly interact with DPG in the protonated state. B) Model of Steller's sea cow hemoglobin (left) and a close-up of the DPG binding pocket (right). Dark blue colored residues represent the 11 *H. gigas* specific substitutions, while those in light blue denote the ancestral state. Homology modelling illustrates how the replacement of β/δ82Lys with neutral Asn inhibits DPG binding to the hemoglobin molecule.

832
833
Figure 4. Oxygen equilibrium curves of prenatal and adult sirenian hemoglobins. A) Oxygen
equilibrium curves for Steller's sea cow (*Hydrodamalis gigas*; blue) Hb Gower 1 ($\zeta_2\varepsilon_2$), HbF ($\alpha_2\gamma_2$),
and HbA ($\alpha_2\beta/\delta_2$) and dugong (*Dugong dugon*; red) HbF and HbA in the presence of allosteric
cofactors 2,3-diphosphoglycerate (2-fold molar excess) and KCl (100 mM) at pH 7.1 (prenatal Hbs)
or 7.2 (HbA). The dashed red line is for dugong HbA in the absence of DPG that illustrates relative
differences in O_2 affinity between the maternal (solid line) and fetal (dashed line) circulations.
Homology models of Steller's sea cow and dugong HbF denote structural alterations arising from
the *H. gigas* specific γ101 (B vs. C, respectively) and γ38 (D vs. E, respectively) replacements.
Solid pink lines denote predicted hydrogen bonds while the dashed pink lines represent predicted
Van der Waals interactions.