Evolution of an extreme hemoglobin phenotype contributed to the
sub-Arctic specialization of extinct Steller's sea cows.

Anthony V. Signore[1], Angela Fago[2], Roy E. Weber[2] and Kevin L. Campbell[1]*

[1]Department of Biological Sciences, University of Manitoba, Winnipeg, R3T 3N2, Canada

[2]Department of Biology, Zoophysiology, Aarhus University, DK-8000 Aarhus, Denmark.

*Corresponding author.

Email: Kevin.Campbell@umanitoba.ca

Author Contributions: AVS and KLC conceptualized the study; AVS performed research; AVS,
AF, REW and KLC analyzed data; AF, REW and KLC contributed reagents and analytical tools;
AVS and KLC drafted the manuscript; AVS, AF, REW and KLC edited the manuscript.

Competing Interest Statement: Authors declare that they have no competing interests.

Keywords: hemoglobin, thermal adaptation, protein biochemistry, paleophysiology, Sirenia

This PDF file includes:

Main Text
Figures 1 to 3

Abstract

The extinct Steller's sea cow (*Hydrodamalis gigas*; †1768) was a whale-sized marine mammal that

manifested profound morphological adjustments to exploit the harsh coastal climate of the North

Pacific. Yet despite first-hand accounts of their biology, little is known regarding the physiological

specializations underlying their evolution to this environment. Here, the adult-expressed

hemoglobin (Hb) of this species is shown to harbor a fixed amino acid replacement at an otherwise

invariant position (β/δ82Lys→Asn) that alters multiple aspects of Hb function. First, our functional

characterization of recombinant sirenian Hb proteins demonstrate that the Hb–O_2 affinity of this

sub-Arctic species was less affected by temperature than those of living (sub)tropical sea cows.

This phenotype presumably safeguarded O_2 delivery to cool peripheral tissues and largely arises

from a reduced intrinsic temperature sensitivity of the *H. gigas* protein owing to the β/δ82Lys→Asn

substitution. Additional experiments on *H. gigas* β/δ82Asn→Lys mutant Hb further reveal this

exchange renders Steller's sea cow Hb unresponsive to the potent intraerythrocytic allosteric

effector 2,3-diphosphoglycerate, a radical modification that is the first documented example of this

phenotype among mammals. Notably, β/δ82Lys→Asn moreover underlies the secondary evolution

of a reduced blood–O_2 affinity phenotype that would have promoted heightened tissue and

maternal/fetal O_2 delivery. Finally, this replacement is shown to increase protein solubility, which is

consistent with an increased O_2 carrying capacity within both the adult and pre-natal circulations

and may have further contributed to meeting the elevated metabolic (thermoregulatory)

requirements and fetal growth rates associated with this species cold adaptation.

Main Text

Introduction

During periods of submergence apnea when atmospheric gas exchange has ceased, the

underwater foraging time by mammals is dictated by onboard oxygen stores and the efficiency of

their use. Thus, evolutionary increases in oxygen stores, in the form of increased hemoglobin (Hb)

and myoglobin—located within erythrocytes and skeletal/cardiac muscle, respectively—are nearly

ubiquitous among aquatic species (1). Notable exceptions to this rule are extant sirenians (sea

cows), a group of strictly aquatic, (sub)tropical herbivores encompassing only four members, three

species of manatee (family Trichechidae) and the dugong, *Dugong dugon* (family Dugongidae).

While sirenians are proficient divers, they do not exhibit the strongly elevated body O_2 stores or an

enhanced dive reflex common to other lineages of marine mammals (2, 3). Rather, previous work

revealed that the sirenian's secondary transition to aquatic life coincided with a rapid evolution of their Hb encoding genes due, in part, to gene conversion events with a neighboring globin pseudogene (4). The resulting high blood–O_2 affinity phenotype presumably allows extant sea cows to maximize O_2 extraction from the lungs during submergence while curbing O_2 offloading, thus lowering overall metabolic intensity and fostering a prolonged breath-hold capacity (4).

While the relatively weak thermoregulatory capacity of extant sea cows confine them to (sub)tropical waters (5, 6), fossil evidence and first-hand accounts of the sub-Arctic Steller's sea cow (*Hydrodamalis gigas*) provide insights into the biological and morphological adaptations of this titanic sirenian to the harsh coastal conditions of the North Pacific, where they persisted from the Miocene until their demise in 1768 (7–9). The retrieval of ancient genetic material from museum specimens has since been instrumental in clarifying the phylogenetic affinities and population history of this species, while providing additional details regarding the evolution of key morphological and physiological attributes (4, 10–15) (Fig. 1A). For example, pilot experiments on "resurrected" Steller's sea cow recombinant Hb demonstrated that the Hb–O_2 affinity of this lineage secondarily decreased following their divergence from dugongs in the mid Oligocene (4). While sirenians do not possess the capacity for non-shivering thermogenesis due to pseudogenization of the *UCP1* gene (11), this affinity shift in Steller's sea cow Hb was speculated to have promoted increased O_2 offloading to fuel increased thermogenesis to help cope with exposure to cold sub-Arctic waters. Although the Hb of this species accumulated 11 amino acid replacements since its divergence from the dugong, this decrease in Hb–O_2 affinity was hypothesized to arise from a highly unusual 82Lys→Asn exchange in the chimeric β-type (β/δ) chain (4). Data mined from more recent ancient DNA studies (10, 14) confirms that this substitution was fixed in the last remaining Steller's sea cow population (Fig. 1B), though the specific functional effect(s) of this substitution have not been characterized. This replacement is intriguing not only because β82Lys is invariant among characterized mammalian Hbs, but because human variants with substitutions at this position display profound alterations in both structural and functional properties (16–19). For example, the human Hb Providence (β82Lys→Asn) variant exhibits a decreased inherent Hb–O_2 affinity and markedly reduced sensitivity to the allosteric effectors 2,3-diphosphoglycerate (DPG), Cl^-, and H^+ (16, 17, 20–23), which preferentially bind and stabilize the (low O_2 affinity) deoxy-state conformation of the protein. However, opposite to what was suggested for Steller's sea cows (4), this exchange causes the whole blood O_2 affinity of Hb Providence carriers to be noticeably higher than that of the general population (20,23). It is thus unclear if and how the Steller's sea cow β/δ82Lys→Asn replacement underlies the lower Hb–O_2 affinity of this extinct species relative to other sirenians, or whether this attribute arises from one (or more) of the other 10 residue replacements that evolved in this lineage.

The β/δ82Lys→Asn exchange also raises other evolutionary significant questions, as it is
predicted alter multiple aspects of Hb function that may lead to antagonistic pleotropic effects.
Notably, this exchange may detrimentally increase the effect of temperature on Hb–O_2 binding and
release (4). The formation of the weak covalent bond between O_2 and the heme iron requires free
energy, thus dictating an inverse relationship between Hb–O_2 affinity and temperature (23). In
temperate and Arctic endotherms this inherent attribute of Hb potentially impedes O_2 delivery to
the limbs and flukes, which are maintained at substantively lower temperatures to minimize heat
loss and hence energy requirements (24). Accordingly, heterothermic mammals generally possess
Hbs whose O_2 binding properties are less sensitive to temperature than the Hbs of non-cold-
adapted species, thereby maintaining sufficient O_2 offloading in the face of decreasing tissue
temperatures. This reduction in thermal sensitivity (the overall enthalpy of oxygenation, ΔH'),
appears to predominantly arise from an increased interaction between allosteric effectors and the
Hb moiety (an exothermic process), which releases additional heat to assist with deoxygenation
(23). Hence, the *H. gigas* β/δ82Lys→Asn replacement, which deletes integral binding sites for the
heterotropic ligands Cl^- and DPG (17), is puzzling in that it is expected to maladaptively increase
the effect of temperature on O_2 uptake and release in the blood of the sub-Arctic Steller's sea cow.

Taken together, it remains unknown whether the *H. gigas* β/δ82Lys→Asn residue
exchange contributed adaptively the species biology or is instead linked to small population sizes
(e.g., genetic drift) over the past half million years (10, 14). To unravel the combined effects of
evolved amino acid replacements on hemoglobin function in relation to the extreme thermal biology
of the extinct Steller's sea cow, we synthesized recombinant Hb proteins of this species together
with those of the dugong (*Dugong dugon*) and Florida manatee (*Trichechus manatus latirostris*),
and measured their O_2 binding properties, relative solubility, responses to allosteric effectors, and
thermal sensitivities. We also synthesized a *H. gigas* β/δ82Asn→Lys Hb mutant to assess the
specific effects of this exchange, together with the Hb of the last common ancestor ('ancestral
dugongid') shared between the dugong and Steller's sea cow (Fig. 1A) in order to assess the
directionality of evolved physicochemical changes in Hb function.

Results and Discussion

O_2 Affinity of Sirenian Hbs. Measured O_2-equilibrium curves of the five examined Hbs revealed
marked differences in intrinsic O_2 affinity (Figs. 2A, S1, S2 and Table S1). In the absence of
allosteric effectors (pH 7.2, 37°C), the P_{50} (the O_2 tension resulting in 50% Hb–O_2 saturation) of
Steller's sea cow Hb (P_{50} = 8.8 mmHg) is ~2 times higher than that of dugong (3.5 mmHg), ancestral
dugongid (4.3 mmHg), and manatee (5.4 mmHg) Hbs under the same conditions (Figs. 2A, S1 and
Table S1). Site directed mutagenesis experiments reveal that the increased intrinsic P_{50} of Steller's

sea cow Hb predominantly arises from the β/δ82Asn substitution, as the β/δ82Asn→Lys mutant

exhibits an intrinsic P_{50} similar to that of the ancestral dugongid (Fig. 2A, S2). Of note, the O_2 affinity

of dugong, ancestral dugongid, and manatee Hbs was reduced in the presence of Cl^- and DPG (P_{50}

= 10.2, 9.9, and 10.9 mmHg, respectively) by a similar degree to that of Asian elephant Hb (25).

This finding extends previous studies conducted on sirenian Hbs (4, 26, 27), and reveals that the

high O_2 affinity of dugong and manatee blood is not attributable to decreased allosteric effector

sensitivity. Conversely, Steller's sea cow Hb was shown to be markedly less responsive to allosteric

effectors, as only a moderate reduction in O_2–affinity was observed in the presence of Cl^- and DPG

(P_{50} = 14.3 mmHg). When the effects of these allosteric effectors were measured individually,

Steller's sea cow Hb exhibits markedly reduced DPG, Cl^-, and H^+ (Bohr) effects relative to those of

the ancestral dugongid and β/δ82Asn→Lys mutant (Fig. 2C-E; Table S1). These data confirm that

a high intrinsic (i.e. in the absence of allosteric effectors) Hb–O_2 affinity is an ancient sirenian trait

that likely aided the transition of the group to the aquatic environment, and that Hb–O_2 (and hence

whole blood) affinity was secondarily reduced in the Steller's sea cow lineage (4). This latter finding

contrasts with allometric expectations for mammals—whereby blood O_2 affinity and body mass are

inversely correlated (28)—and thus further suggests this modification serves an adaptive function.

The single most distinct feature of *H. gigas* Hb is the lack of a discernable effect of DPG

on P_{50} ($\Delta log P_{50}^{(0.2\ mM\ DPG\ –\ 0.0\ mM\ DPG)}$ = 0.02 at 37°C and pH 7.2; Fig. 2C and Table S1), relative to

the Hbs of the ancestral dugongid (0.27) and the extant manatee and dugong (0.14 and 0.30,

respectively). This intracellular effector generally occurs in equimolar concentrations to Hb (29) and

strongly decreases the O_2 affinity of most mammalian Hbs via direct electrostatic interactions with

β2His and β82Lys (Fig. 3A), together with potential water-mediated interactions with β143His and

the α-NH_2 group of 1Val of the $β_2$ chain (30). However, unlike the other (ionizable) residues whose

ability to interact with DPG is highly pH dependent, β82Lys is strongly cationic and thus is able to

bind DPG across the entire physiological pH range. Presumably arising from this indispensable role

in DPG binding, this residue is uniformly conserved in mammalian Hbs (Fig. 3A), with the exception

of several heterozygous adult human HbA carriers with substitutions at this position (18, 19, 31).

Given that none of the other six β/δ-chain replacements that evolved on the Steller's sea cow

branch (Figs. 3B and S3) are implicated in DPG binding, the deletion of the integral DPG binding

site at β/δ82 in Steller's sea cow Hb is consistent with its inability to bind DPG (Fig. 3B). This

conclusion is further supported by our measurements on the Steller's sea cow β/δ82Asn→Lys

mutant, which show that reversion to the ancestral state restores the DPG effect to the same level

observed in ancestral dugongid Hb (Fig. 2C). Notably, and despite possessing the identical DPG

binding site residues, the *H. gigas* protein exhibits a distinctly lower DPG effect than Hb Providence

(0.08; 17,20). The stronger reduction of DPG sensitivity in the former thus implicates an epistatic

contribution from other amino acids in the vicinity of the DPG pocket. Importantly, the Lys→Asn

replacement in the DPG binding pocket causes the O_2 affinity of human HbA to increase in the
presence of allosteric cofactors (20,23), whereas results presented in Fig. 2A show Steller's sea
cow Hb–O_2 affinity is reduced relative to its ancestors carrying β/δ82Lys under all test conditions.
This observation highlights a growing body of research indicating that both the direction and overall
phenotypic effect of specific amino acid substitutions may be conditional on the genetic background
in which they occur (32).
The insensitivity of *H. gigas* Hb to DPG it also notable as it would have markedly reduced
their capacity to modulate blood–O_2 affinity *in vivo* (e.g., seasonally), and is the first demonstrated
example of a genuine DPG insensitive Hb phenotype among mammals. While eastern moles
(*Scalopus aquaticus*) are a possible exception (33), feliformid carnivores, ruminants, and two
species of lemurs have also traditionally been placed in the 'DPG insensitive' category (29) despite
the fact their Hbs are moderately responsive to DPG in the absence of Cl^- (17, 34–36). Nonetheless,
red blood cell DPG concentrations of species with suppressed DPG sensitivities are markedly
reduced relative to mammals whose Hb–O_2 affinity is regulated by DPG (<0.1-1.0 mM vs. 4-10 mM,
respectively;(29)). While the potential benefits of a low DPG sensitivity phenotype have been
debated (33, 37) approximately 20% of glucose uptake by human erythrocytes is diverted to DPG
synthesis (38), thereby bypassing production of both ATP molecules generated via anaerobic
glycolysis (29, 39, 40). Accordingly, since each molecule of DPG produced comes at the expense
of producing an ATP molecule, the probability that Steller's sea cow blood similarly contained low
levels of this organophosphate is high.

Hb Solubility. Ectopic expression of human Hb Providence β82Asp mutants in *E. coli* has been
shown to improve soluble protein production by 47-116% relative to the expression of human Hb
variants not carrying this substitution (22). Consistent with this observation, we found that Steller's
sea cow Hb is more soluble than those of other sirenians and the engineered *H. gigas*
β/δ82Asn→Lys mutant (Fig. 2F and S4). While the precise mechanism underlying this
phenomenon is unknown, Hb Providence variants have been shown to strongly reduce irreversible
oxidative damage of nearby β93Cys that initiates Hb denaturation (16, 41, 42). These β82
replacements thereby presumably decrease the rate of Hb turnover and increase the half-life of the
protein (42), and may be the underlying cause of mild polycythemia in humans carrying this
substitution (20, 31). Blood with an elevated O_2 carrying capacity is typical of most mammalian
divers, where it increases onboard O_2 stores and extends dive times (1), but is not observed in
extant sirenians (27, 43, 44). However, solubility driven increases in blood Hb levels resulting from
the β82Lys→Asn exchange would have allowed Steller's sea cows to maintain an elevated rate of
tissue O_2 delivery to meet their metabolic demands during extended underwater foraging intervals.
Although this species was presumably unable to completely submerge (9, 45), this conjecture is

corroborated by Steller's account that, "*they keep their heads always under water* [foraging]*, without*

regard to life and safety" (9).

Thermal Sensitivity. The invariant energy change associated with forming the weak covalent bond

between O_2 and the heme iron (i.e. the enthalpy of heme oxygenation; ΔH^{O2}) is exothermic (-59 kJ

mol^{-1} O_2) (46), and only moderately opposed by the endothermic solubilization of O_2 (ΔH^{H2O}; 12.55

kJ mol^{-1} O_2), resulting in an inverse relationship between temperature and Hb–O_2 affinity. However,

the heat of the T→R conformational change ($\Delta H^{T\rightarrow R}$), and the oxygenation-linked dissociation of H^+

(ΔH^{H+}), Cl^- (ΔH^{Cl-}), and DPG (ΔH^{DPG}) may offset this relationship, such that the overall enthalpy of

Hb oxygenation ($\Delta H'$) can become greatly minimized or even endothermic (23, 47). By facilitating

adequate oxygenation of cool peripheral tissues, Hbs with numerically low $\Delta H'$ values are

interpreted to be adaptive for cold-tolerant, regionally heterothermic mammals. The evolution of

this phenotype has predominantly been attributed to the formation of additional heterotropic ligand

binding sites on the protein moiety, as has previously been demonstrated for the woolly mammoth,

Mammuthus primigenius (23, 25). Conversely, Steller's sea cow Hb lacks heterotropic binding of

DPG and displays lower Bohr (H^+) and Cl^- effects than all other sirenian Hbs measured (Fig. 2;

Table S1), yet exhibits a $\Delta H'$ value (-18.8 kJ mol^{-1} O_2; Fig. 2B) that is close to that of mammoth Hb

(-17.2 kJ mol^{-1} O_2) (25). This striking convergence largely arises from the inherently low ΔH of

stripped Steller's sea cow Hb (-34.2 kJ mol^{-1} O_2) relative to dugong, ancestral dugongid, and

manatee Hbs (range: -50.1 to -58.6 kJ mol^{-1} O_2) at pH 7.8—where oxygenation-linked binding of

protons is minimal—and indicates that structural differences modifying the T→R transition largely

underlie the low thermal sensitivity of *H. gigas* Hb. Recent studies have shown that a large positive

$\Delta H^{T\rightarrow R}$ may similarly contribute to the low $\Delta H'$ of deer mouse, cow, shrew, and mole Hbs (33, 48–

51) suggesting that this mechanism of temperature adaptation may be more widespread than

previously appreciated. Our experiments with the Steller's sea cow β82Asn→Lys mutant further

implicate substitutions at this position as a key factor underlying the inherently low ΔH of the protein,

as this modified protein displays a greatly increased ΔH in the absence of allosteric effectors

relative to the wild-type *Hydrodamalis* protein (Fig. 2B). Interestingly, despite these inherent ΔH

differences between the mutant and wild-type Steller's sea cow Hbs, their $\Delta H'$ values are

indistinguishable in the presence of allosteric effectors (Fig. 2B). These data suggest that β82Asn

uncouples thermal sensitivity from DPG concentration, permanently conferring the *H. gigas* protein

with a numerically low $\Delta H'$ by genetic assimilation while simultaneously eliminating the energetic

cost of DPG production within the red blood cells.

Paleophysiology of Steller's sea cows. The posthumously published behavioral and anatomical accounts of the last remaining *H. gigas* population by naturalist Georg Wilhelm Steller while stranded on Bering Island (55°N, 166°E) in 1741/1742 provide a rich tapestry to interpret the paleophysiology of this colossal marine herbivore. For example, their protective thick bark-like hide and extensive blubber layer give credence to the extreme nature of their shallow rocky and (during winter) ice strewn habitat (10). Here, as Steller (9) remarked, they used fingerless, bristle-covered forelimbs for support and to shear "*algae and seagrasses from the rocks*", which they masticated "*not with teeth, which they lack altogether, but with*" large, ridged keratinized pads located on the upper palate and lower mandible. Although they became visibly thin during winter when "*their spinous processes can be seen*", Steller (9) noted that "*(t)hese animals are very voracious, and eat incessantly*" such that their stupendous stomach ("*6 feet* [1.8 m] *long, 5 feet* [1.5 m] *wide*") and enormous intestines—which measured a remarkable 5,958 inches (~151.5 m) from esophagus to anus, equivalent to "*20½ times as long as the whole animal*"—are constantly "*stuffed with food and seaweed*". The proportionally larger gut (45) is consistent with Steller's sea cow's higher energetic requirements relative to extant manatees, which, owing to their low metabolic intensity become cold stressed and die if chronically exposed to water temperatures below 15°C (52). The reduction in insulative blubber thickness during the winter months would likely have compounded the rate of heat loss to sub-zero degree Celsius air and water, though may have been compensated for by arteriovenous anastomoses that regulated blood flow to the skin, and by countercurrent rete supplying the flippers and tail flukes, the latter of which are well developed in manatees and presumably other sirenians (53, 54). These structures conserve thermal energy by promoting profound cooling at the appendages and periphery (26), and presumably underlie the low thermal dependence of Steller's sea cow Hb relative to those of extant sea cows.

Reductions in blood–O_2 affinity accompanying the *H. gigas* β/δ82Lys→Asn substitution is expected to have further augmented tissue O_2 delivery without tangible effects on lung O_2 uptake, thereby helping to fuel thermogenesis and maintain a stable core temperature. The latter was presumably supplemented by a substantive heat increment arising from fermentation and other post-prandial processes (54). Although the attendant increase in the rate of O_2 consumption would have mandated a reduction in breath-hold endurance—likely reflecting the relatively short submergence times (4 to 5 minutes) observed by Steller (9)—our results suggests this may be been partially counteracted by an elevated blood–O_2 carrying capacity that was potentially coupled to a greater lung volume (45). Underwater foraging times were presumably further defended by key components of the dive reflex, namely bradycardia and peripheral vasoconstriction. Indeed, Steller inadvertently was the first to (indirectly) describe this phenomenon as he observed his crew hunting the animals with spears and knives, "*the blood from the wounded back spurted up like a fountain.*

As long as he kept his head under water the blood did not flow out, but as soon as he raised his

head to breathe the blood leaped forth anew".

A final compelling aspect of Steller's sea cow evolution was their immense increase in

size—up to 11,000 kg in mass and 10 m in length—relative to extant sirenians (7). While Steller

does not provide measurements of "*their tender little offspring*", Gerhard Friedrich Müller, who

edited Steller's manuscript prior to publication, noted calves "*weighed 1200 pounds* [544 kg] *and*

upwards" (55). This value is ~10 to 50 times the mass of new born manatees and dugongs (~10-

50 kg) (56) and is suggestive of rapid prenatal growth during the ~1 year gestational period

indicated by Steller (9). Given that all mammals except ruminants and anthropoid primates express

the same Hb isoform during both the fetal and post-natal stages of development (57), the apparently

unique DPG insensitive phenotype of *H. gigas* is particularly notable in this regard owing to its

potential impact on maternal/fetal O_2 exchange. Presumably to assist in this process, fetal blood

cells express only trace amounts of DPG and hence the O_2 affinity of fetal blood is elevated relative

to that of the maternal circulation in the vast majority of mammalian species (29, 57). By contrast,

and owing to the inability of Steller's sea cows Hb to respond to DPG in either the fetal or adult

circulations, this species would represent a rare example (feloids and eastern moles are others) in

which fetal and maternal blood have the same (relatively low) O_2 affinity. However, placental O_2

delivery in these species will still be defended by the well-known double Bohr effect, whereby CO_2

transport from the fetal to maternal circulation increases blood O_2 affinity in the former while

lowering it in the latter (57). More importantly, the evolved reduction in blood O_2 affinity of Steller's

sea cows would have stipulated that equilibrium between umbilical and uterine blood was reached

at a higher PO_2, a condition that is expected to substantially improve O_2 delivery to the fetal

circulation (57). The lower fetal blood O_2 affinity and concomitant higher fetal blood-to-tissue PO_2

gradients are further expected to have augmented O_2 delivery to the developing tissues of this

species. As such, these attributes, together with increases in Hb solubility/reduced susceptibility to

oxidative damage arising from β/δ82Lys→Asn that conceivably also elevated the O_2 carrying

capacity of fetal blood, may have been important contributors to the enhanced fetal growth rate of

these immense sirenians. The resulting increase in thermal inertia and relatively low surface-area-

to-volume ratio following birth, together with an adaptively reduced Hb thermal sensitivity and thick

a 'bark-like' skin arising from inactivation of lipoxygenase genes (10), were presumably central

components of Steller's sea cows successful exploitation of the harsh sub-Arctic marine

environments of the North Pacific.

Materials and Methods

Sequence Collection and Analyses. The adult-expressed Hb genes (HBA and HBB/HBD) of the Florida manatee, dugong, and Steller's sea cow, and the most recent common ancestor shared by Steller's sea cow and the dugong ('ancestral dugongid') have previously been determined (4). As the *H. gigas* β/δ82Lys→Asn exchange is not known to occur in any living species, we mined recently deposited genomes for 13 additional Steller's sea cows (PRJNA484555, PRJEB43951) to test for the prevalence of this replacement in the population. Briefly, we first searched SRA files of each specimen using the megablast function against a previously determined *H. gigas HBB/HBD* gene sequence (GenBank accession #: MK562081). All hits were then downloaded, trimmed of adapters and low-quality regions using BBDuk (Joint Genome Institute), and assembled to *H. gigas HBB/HBD* using Geneious Prime 2019 software (Biomatters Ltd, Auckland, New Zealand). Assemblies generated using genome reads that were not pre-treated with uracil-DNA glycosylase and endonuclease VIII to reduce C→T and G→A damage artifacts (10) were examined to ensure these deamination artifacts did not affect the consensus sequences.

Construction of Recombinant Hb Expression Vectors. Coding sequences for the above four Hb proteins were optimized for expression in *E. coli* and synthesized *in vitro* by GenScript (Piscataway, NJ). The resulting gene cassettes were digested with restriction enzymes and tandemly ligated into a custom Hb expression vector (58) using a New England BioLabs Quick Ligation Kit as recommended by the manufacturer. Chemically competent JM109 (DE3) *E. coli* (Promega) were prepared using a Z-Competent *E. coli* Transformation Kit and Buffer Set (Zymo Research). We also prepared a *H. gigas* β/δ82Asn→Lys Hb mutant via site-directed mutagenesis on the Steller's sea cow Hb expression vector by whole plasmid amplification using mutagenic primers and Phusion High-Fidelity DNA Polymerase (New England BioLabs), phosphorylation with T4 Polynucleotide Kinase (New England BioLabs), and circularization with an NEB Quick Ligation Kit (New England BioLabs). All site-directed mutagenesis steps were performed using the manufacture's recommended protocol.

Hb expression vectors were co-transformed into JM109 (DE3) chemically competent *E. coli* alongside a plasmid expressing methionine aminopeptidase (58), plated on LB agar containing ampicillin (100 µg/ml) and kanamycin (50 µg/ml), and incubated for 16 hours at 37°C. A single colony from each transformation was cultured in 50 ml of 2xYT broth for 16 hours at 37°C while shaking at 200 rpm. Post incubation, 5 ml of the culture was pelleted by centrifugation and plasmid DNA was isolated using a GeneJET Plasmid Miniprep Kit (Thermo Scientific). The plasmid sequence was verified using BigDye 3.1 sequencing chemistry and an ABI3130 Genetic Analyzer. The remainder of the culture was supplemented with glycerol to a final concentration of 10%, divided into 25 ml aliquots and stored at -80°C until needed for expression.

Expression and Purification of Recombinant Hb. 25 ml of starter culture (above) was added to 1250 ml of TB media containing ampicillin (100 µg/ul) and kanamycin (50 µg/ul) and distributed evenly amongst five 1 L Erlenmeyer flasks. Cultures were grown at 37°C while shaking at 200 rpm until the absorbance at 600 nm reached 0.6-0.8. Hb expression was induced by supplementing the media with 0.2 mM isopropyl β-D-1-thiogalactopyranoside, 50 µg/ml of hemin and 20 g/L of glucose and the culture was incubated at 28°C for 16 hours while shaking at 200 rpm. Once expression had completed, dissolved O_2 was removed by adding sodium dithionite (1 mg/ml) to the culture, which was promptly saturated with CO for 15 minutes. Bacterial cells were then pelleted by centrifugation and Hb purified by ion exchange chromatography according to Natarajan et al. [58]

It should be noted that the β82Asn residue of human Hb Providence is relatively uncommon in that it slowly undergoes post-translational deamidation *in vivo* to form aspartic acid, with the latter residue (β82Asp) comprising ~67-75% in mature mixed blood [20, 59]. While it is unknown to what degree *H. gigas* β/δ82Asn was catalyzed into Asp in nature, O_2 binding data (see below) of this species was collected from freshly purified recombinant samples for which only one peak— presumably β/δ82Asn—was resolved during chromatography (data not shown). Additionally, this reaction is dependent on the local protein environment [60], specifically two nearby residues β143His and β83Gly [59]. Importantly, the latter residue was replaced by β/δ83Ser on the Steller's sea cow branch (Figs. 1B and 3B), which is expected to slow (but not stop) the rate of deamidation [60]. Regardless, since the two Hb Providence isoforms have similar O_2 affinities and functional properties [17, 20, 21] it is unlikely that presence of β/δ82Asp in Steller's sea cow blood would meaningfully alter the results and interpretations presented herein.

Functional Analyses of Hbs. O_2-equillibrium curves for Hb solutions (0.25 mM heme in 0.1 M HEPES buffers) were measured at 25 and 37°C using the thin film technique [61]. Hb solutions varied in their pH (6.9, 7.4, and 7.9), chloride concentration (0 or 0.1 M KCl), and organic phosphate concentration (0 or 2-fold molar excess of DPG relative to tetrameric Hb concentrations) in order to test the influence of these cofactors on Hb function. Each Hb solution was sequentially equilibrated with three to five different oxygen tensions (PO_2) that result in Hb–O_2 saturations between 30 to 70%. Hill plots (log[fractional saturation/[1-fractional saturation]] vs logPO_2) constructed from these measurements were used to determine the PO_2 (P_{50}) and the cooperativity coefficient (n_{50}) at half saturation, from the χ-intercept and slope of these plots, respectively. By this method, the r^2 determination coefficients for the fitted curves exceed 0.995 and the standard errors (SEM) are less than 3% of the P_{50} and n_{50} values [51]. P_{50} values at 25 and 37°C were used to assess the thermal sensitivity of sirenian Hbs by calculating the apparent enthalpy of oxygenation using the van't Hoff isochore:

$\Delta H = 2.303R \times \Delta\log P_{50} \times (1/T_1 - 1/T_2)^{-1}$

where R is the universal gas constant and T_1 and T_2 are the absolute temperatures (°K) at which

the P_{50} values were measured. All ΔH values were corrected for the heat of O_2 solubilization (12.55

kJ mol^{-1} O_2).

Solubility assay. Ammonium sulfate was added to Hb solutions (0.074 ± 0.004 mM Hb_4) to

generate final concentrations that ranged from 0 to 3.5 M. These solutions were incubated for 60

minutes at 37°C and the remaining soluble Hb was measured via Drabkin's reagent, according to

the manufacturer's instructions (Sigma-Aldrich).

Homology modelling. To assess the structural effect of the *H. gigas* specific mutations on the

DPG binding site, homology models of ancestral dugongid and Steller's sea cow Hb were

constructed using the SWISS-MODEL server (62) using the three-dimensional human deoxy

structure with DPG bound (PDB: 1B86) as a template (30). The sequence conservation of amino

acid residues implicated in DPG binding were calculated by the ConSurf Server (63) from a

subsample of 51 mammalian beta-type hemoglobin chains downloaded from GenBank (Table S2).

Homology models were visualized with UCSF Chimera (64).

Acknowledgments

This manuscript is dedicated to the memory of our dear friend Joseph (Joe) Bonaventura for his

pioneering work on the human Hb Providence Asn/Asp proteins. We thank Mike Gaudry and Elin

Ellebæk Petersen for technical assistance, Chandrasekhar Natarajan for providing us with a

hemoglobin expression plasmid, and J. Storz for constructive feedback on an earlier version of this

manuscript. Authorization to use paintings by C. Buell was kindly provided by J. Gatesy. This study

was supported by NSERC (Canada) Discovery and Accelerator Supplement Grants (K.L.C.;

RGPIN/238838-2011, RGPIN/412336-2011, and RGPIN/06562-2016), an NSERC Postgraduate

Scholarship (A.V.S), the Faculty of Science and Technology, Aarhus University (REW), and the

Danish Council for Independent Research (AF; DFF - 4181-00094).

404
405
406 **References**
407

1.   P. J. Ponganis, Diving mammals. *Compr Physiol* **1**, 447–465 (2011).

2.   M. H. Blessing, Studies on the concentration of myoglobin in the sea-cow and porpoise.
Comp Biochem Physiol A **41**, 475–480 (1972).

3.   P. F. Scholander, L. Irving, Experimental investigations on the respiration and diving of the
florida manatee. *J Cell Comp Physiol* **17**, 169–191 (1941).

4.   A. V. Signore, *et al.*, Emergence of a chimeric globin pseudogene and increased hemoglobin
oxygen affinity underlie the evolution of aquatic specializations in Sirenia. *Mol Biol Evol* **36**,
1134–1147 (2019).

5.   G. J. Gallivan, J. W. Kanwisher, R. C. Best, Heart rates and gas exchange in the Amazonian
manatee (*Trichechus inunguis*) in relation to diving. *J Comp Physiol B* **156**, 415–423 (1986).

6.   H. Marsh, T. J. O'Shea, J. E. Reynolds III, *Ecology and conservation of the Sirenia: dugongs
and manatees* (2011).

7.   D. P. Domning, An ecological model for late tertiary Sirenian evolution in the North Pacific
Ocean. *Syst Biol* **25**, 352–362 (1976).

8.   L. Stejneger, How the great northern sea-cow (Rytina) became exterminated. *The American
Naturalist* **21**, 1047–1054 (1887).

9.   G. W. Steller, De bestiis marinis [The beasts of the sea]. *Novi Commentarii Acad Sci Imp
Petropoli* (1751).

10. D. Le Duc, *et al.*, Genomic basis for skin phenotype and cold adaptation in the extinct
Steller's sea cow. *Sci Adv* **8**, eabl6496 (2022).

11. M. J. Gaudry, *et al.*, Inactivation of thermogenic UCP1 as a historical contingency in multiple
placental mammal clades. *Sci Adv* **3**, e1602878 (2017).

12. S. Mirceta, *et al.*, Evolution of mammalian diving capacity traced by myoglobin net surface
charge. *Science* **340**, 1234192 (2013).

13. F. S. Sharko, *et al.*, Molecular phylogeny of the extinct Steller's sea cow and other Sirenia
species based on their complete mitochondrial genomes. *Genomics* **111**, 1543–1546
(2019).

14. F. S. Sharko, *et al.*, Steller's sea cow genome suggests this species began going extinct
before the arrival of Paleolithic humans. *Nat Commun* **12**, 2215 (2021).

15. M. S. Springer, *et al.*, Interordinal gene capture, the phylogenetic position of Steller's sea
cow based on molecular and morphological data, and the macroevolutionary history of
Sirenia. *Mol Phylogenet Evol* **91**, 178–193 (2015).

16. B. Abraham, *et al.*, Isolated Hb Providence β82Asn and β82Asp fractions are more stable
than native HbA0 under oxidative stress conditions. *Biochem* **50**, 9752–9766 (2011).

17. J. Bonaventura, *et al.*, Hemoglobin providence. Functional consequences of two alterations
of the 2,3-diphosphoglycerate binding site at position beta 82. *J Biol Chem* **251**, 7563–7571
(1976).

18. E. Ikkala, *et al.*, Hb Helsinki: A variant with a high oxygen affinity and a substitution at a 2, 3-
DPG binding site (β82 [EF6] Lys→ Met). *Acta Haematol* **56**, 257–275 (1976).

19. J. Sugihara, *et al.*, Hemoglobin Rahere, a human hemoglobin variant with amino acid
substitution at the 2,3-diphosphoglycerate binding site. Functional consequences of the
alteration and effects of bezafibrate on the oxygen bindings. *J Clin Invest* **76**, 1169–1173
(1985).

20. J. Bardakjian, *et al.*, A new case of Hemoglobin Providence (α2 β2 82 (Ef6) Lys → Asn or
Asp) discovered in a French Caucasian family. Structural and functional studies.
Hemoglobin **9**, 333–348 (1985).

21. S. Charache, *et al.*, Postsynthetic deamidation of hemoglobin Providence (beta 82 Lys
replaced by Asn, Asp) and its effect on oxygen transport. *J Clin Invest* **59**, 652–658 (1977).

22. M. J. Weickert, M. Pagratis, C. B. Glascock, R. Blackmore, A mutation that improves soluble
recombinant hemoglobin accumulation in *Escherichia coli* in heme excess. *App Enviro*
Micro **65**, 640–647 (1999).

23. R. E. Weber, K. L. Campbell, Temperature dependence of haemoglobin-oxygen affinity in
heterothermic vertebrates: mechanisms and biological significance. *Acta Physiol* **202**, 549–
562 (2011).

24. K. L. Campbell, M. Hofreiter, Resurrecting phenotypes from ancient DNA sequences:
promises and perspectives. *Can J Zool* **93***,* 701–710 (2015).

25. K. L. Campbell, *et al.*, Substitutions in woolly mammoth hemoglobin confer biochemical
properties adaptive for cold tolerance. *Nat Genet* **42**, 536–540 (2010).

26. M. McCabe, R. Hamilton, H. Marsh, Some studies on the oxygen affinity of haemoglobin from
the dugong. *Comp Biochem Physiol A* **61**,19–22 (1978).

27. M. Farmer, R. E. Weber, J. Bonaventura, R. C. Best, D. Domning, Functional properties of
hemoglobin and whole blood in an aquatic mammal, the Amazonian manatee (*Trichechus*
inunguis). *Comp Biochem Physiol A* **62**, 231–238 (1979).

28. K. Schmidt-Neilsen, J. L. Larimer, Oxygen dissociation curves of mammalian blood in relation
to body size. *Am J Physiol* **195**, 424–428 (1958).

29. H. F. Bunn, Regulation of Hemoglobin Function in Mammals. *Amer Zool* **20**, 199–211 (1980).

30. V. Richard, G. G. Dodson, Y. Mauguen, Human deoxyhaemoglobin-2, 3-diphosphoglycerate
complex low-salt structure at 2.5 Å resolution. *J Mol Biol* **233**, 270–274 (1993).

31. W. F. Moo-Penn, D. L. Jue, K. C. Bechtel, M. H. Johnson, R. M. Schmidt, Hemoglobin
Providence. A human hemoglobin variant occurring in two forms *in vivo*. *J Biol Chem* **251**,
7557–7562 (1976).

32. J. F. Storz, Causes of molecular convergence and parallelism in protein evolution. *Nat Rev*
Genet **17**, 239–250 (2016).

33. K. L. Campbell, *et al.*, Molecular basis of a novel adaptation to hypoxic-hypercapnia in a
strictly fossorial mole. *BMC Evol Biol* **10**, 214 (2010).

34. C. Fronticelli, E. Bucci, A. Razynska, Modulation of oxygen affinity in hemoglobin by solvent
components: Interaction of bovine hemoglobin with 2,3-diphosphoglycerate and monatomic
anions. *J Mol Biol* **202**, 343–348 (1988).

35. J. E. Janecka, *et al.*, Genetically based low oxygen affinities of felid hemoglobins: lack of
biochemical adaptation to high-altitude hypoxia in the snow leopard. *J Exp Biol* **218**, 2402–
2409 (2015).

36. M. F. Perutz, G. Fermi, C. Poyart, J. Pagnier, J. Kister, A novel allosteric mechanism in
haemoglobin. Structure of bovine deoxyhaemoglobin, absence of specific chloride-binding
sites and origin of the chloride-linked Bohr effect in bovine and human haemoglobin. *J Mol
Biol* **233**, 536–545 (1993).

37. F. R. Kay, 2,3-diphosphoglycerate, blood oxygen dissociation and the biology of mammals.
Comp Biochem Physiol A **57**, 309–316 (1977).

38. J. Duhm, B. Deuticke, E. Gerlach, Metabolism of 2,3-diphosphoglycerate and glycolysis in
human red blood cells under the influence of dipyridamole and inorganic sulfur compounds.
Biochimica et Biophysica Acta **170**, 452–454 (1968).

39. S. Rapoport, J. Luebering, The formation of 2, 3-diphosphoglycerate in rabbit erythrocytes:
the existence of a diphosphoglycerate mutase. *J Biol Chem* **183**, 507-516 (1950).

40. K. J. Kauffman, J. D. Pajerowski, N. Jamshidi, B. O. Palsson, J. S. Edwards, Description and
analysis of metabolic connectivity and dynamics in the human red blood cell. *Biophys J* **83**,
646–662 (2002).

41. S. Jana, M. B. Strader, A. I. Alayash, The Providence mutation (βK82D) in human
hemoglobin substantially reduces βCysteine 93 oxidation and oxidative stress in endothelial
cells. *Int J Mol Sci***21**, 9453 (2020).

42. M. B. Strader, *et al.*, Engineering oxidative stability in human hemoglobin based on the Hb
providence (βK82D) mutation and genetic cross-linking. *Biochem J* **474**, 4171–4192 (2017).

43. A. W. Wong, J. M. Lanyon, H. L. Sneath, G. R. Leggatt, L. Woolford, Comparison of i-STAT®
with traditional laboratory analysers in the measurement of blood analytes from field
captured dugongs (*Dugong dugon*). *Aquat Mam* **44**, 19–31 (2018).

44. J. R. White, D. R. Harkness, R. E. Isaacks, D. A. Duffield, Some studies on blood of the
Florida manatee, *Trichechus manatus latirostris*. *Comp Biochem Physiol A* **55**, 413–417
(1976).

45. D. P. Domning, "What can we infer about the behavior of extinct Sirenians?" in *Ethology and
Behavioral Ecology of Sirenia*, Ethology and Behavioral Ecology of Marine Mammals., H.
Marsh, Ed. (Springer International Publishing, 2022), pp. 1–17.

46. D. H. Atha, G. K. Ackers, Calorimetric determination of the heat of oxygenation of human
hemoglobin as a function of pH and the extent of reaction. *Biochem* **13**, 2376–2382 (1974).

47. R. E. Weber, K. L. Campbell, A. Fago, H. Malte, F. B. Jensen, ATP-induced temperature
independence of hemoglobin-O_2 affinity in heterothermic billfish. *J Exp Biol* **213**, 1579–1585
(2010).

48. K. L. Campbell, A. V. Signore, M. Harada, R. E. Weber, Molecular and physicochemical
characterization of hemoglobin from the high-altitude Taiwanese brown-toothed shrew
(*Episoriculus fumidus*). *J Comp Physiol B* **182**, 821–829 (2012).

49. B. Jensen, J. F. Storz, A. Fago, Bohr effect and temperature sensitivity of hemoglobins from
highland and lowland deer mice. *Comp Biochem Physiol A* **195**, 10–14 (2016).

50. A. V. Signore, J. Stetefeld, R. E. Weber, K. L. Campbell, Origin and mechanism of thermal
insensitivity in mole hemoglobins: a test of the 'additional' chloride binding site hypothesis. *J
Exp Biol* **215**, 518–525 (2012).

51. R. E. Weber, A. Fago, K. L. Campbell, Enthalpic partitioning of the reduced temperature
sensitivity of O_2 binding in bovine hemoglobin. *Comp Biochem Physiol A* **176**, 20–25 (2014).

52. T. J. O'Shea, C. A. Beck, R. K. Bonde, H. I. Kochman, D. K. Odell, An analysis of manatee
mortality patterns in Florida, 1976-81. *J Wildl Manag* **49**, 1–11 (1985).

53. S. A. Rommel, H. Caplan, Vascular adaptations for heat conservation in the tail of Florida
manatees (*Trichechus manatus latirostris*). *J Anat* **202**, 343–353 (2003).

54. C. D. Marshall, D. K. Sarko, R. L. Reep, "Morphological and sensory innovations for an
aquatic lifestyle" in *Ethology and Behavioral Ecology of Sirenia*, Ethology and Behavioral
Ecology of Marine Mammals., H. Marsh, Ed. (Springer International Publishing, 2022), pp.
19–65.

55. G. F. Mueller, Voyages from Asia to America, for Completing the Discoveries of the North
West Coast of America. London. (T. Jefferys, 1761).

56. D. K. Odell, "Sirenian life history" in *Encyclopedia of Marine Mammals (Second Edition)*, W.
F. Perrin, B. Würsig, J. G. M. Thewissen, Eds. (Academic Press, 2009), pp. 1019–1021.

57. A. M. Carter, Placental Gas Exchange and the Oxygen Supply to the Fetus. *Compr Physiol* **5**,
1381–1403 (2015).

58. C. Natarajan, *et al.*, Expression and purification of recombinant hemoglobin in *Escherichia
coli*. *PLoS ONE* **6**, e20176 (2011).

59. M. F. Perutz, J. H. Fogg, J. A. Fox, Mechanism of deamidation of Haemoglobin Providence
Asn. *J Mol Biol* **138**, 669–670 (1980).

60. N. E. Robinson, Protein deamidation. *Proc Nat Acad Sci* **99**, 5283–5288 (2002).

61. R. E. Weber, Use of ionic and zwitterionic (Tris/BisTris and HEPES) buffers in studies on
hemoglobin function. *J Appl Physiol* **72**, 1611–1615 (1992).

62. A. Waterhouse, *et al.*, SWISS-MODEL: homology modelling of protein structures and
complexes. *Nucleic Acid Res* **46**, W296–W303 (2018).

63. H. Ashkenazy, *et al.*, ConSurf 2016: an improved methodology to estimate and visualize
evolutionary conservation in macromolecules. *Nucleic Acid Res* **44**, W344–W350 (2016).

64. E.F. Pettersen, *et al.*, UCSF Chimera–a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605–1612 (2004).

65. S. Heritage, E. R. Seiffert, Total evidence time-scaled phylogenetic and biogeographic
models for the evolution of sea cows (Sirenia, Afrotheria). *PeerJ* **10**, e13886 (2022).

Figures and Tables
562

Figure 1. Evolution of notable morphological and genetic attributes within Sirenia. A) Phenotypic
innovations contributing to the unique biology of Steller's sea cows are mapped along the sirenian
phylogeny (red bars) with the underlying genetic causes shown in brackets where known. Note that
the red bars do not represent the dating of these traits and that their placement order is arbitrary.
Divergence dates in millions of years (Mya) are based on Heritage and Seiffert (65). The ancestral
dugongid (Anc. dugongid) is represented by a late-Oligocene *Metaxytherium spp*. Images by Carl
Buell, and are used with permission. B) Partial nucleotide alignment of the sirenian β/δ-globin gene
encoding the central region of the 2,3-diphosphoglycerate binding pocket of hemoglobin;
corresponding amino acid residues (bolded) are provided below each sequence. The G→C
nucleotide mutation underlying the otherwise invariant amino acid substitution (β/δ82Lys→Asn;
K→N) of Steller's sea cow hemoglobin is apparent in all 16 individuals for which sequence data is
available. Values in brackets next to each *H. gigas* specimen represent the depth of sequence
coverage for this nucleotide. Dots represent sequence identity with *T. manatus*.

580
581
Figure 2. Biochemical properties of hemoglobins (Hbs) from manatee (*T. manatus*), dugong (*D.*
dugon), ancestral dugongid (Anc. dugongid), Steller's sea cow β/δ82Asn→Lys mutant (*H. gigas*
β82K), and wild-type Steller's sea cow (*H. gigas*). All values were measured at 37°C and corrected
to pH 7.2, unless otherwise noted. A) Oxygen tensions at half O_2 saturation (P_{50}) in the absence
(stripped) and presence of allosteric cofactors (0.2 mM 2,3-diphosphoglycerate (DPG) and 0.1 M
KCl). B) The enthalpy of oxygenation (ΔH) between 25 and 37°C in stripped Hb and in the presence
of allosteric cofactors (0.2 mM DPG and 0.1 M KCl). C) The effect of DPG on sirenian Hbs
determined from $\log P_{50}^{(0.2 \text{ mM DPG})} - \log P_{50}^{(\text{stripped})}$. D) The effect of chloride on sirenian Hbs
determined from $\log P_{50}^{(0.1 \text{ M KCl})} - \log P_{50}^{(\text{stripped})}$. E) The Bohr effect of sirenian Hbs in the presence
of allosteric cofactors (0.2 mM DPG and 0.1 M KCl), as calculated from $\Delta \log P_{50}/\Delta pH$ over the pH
range 6.9 and 7.8. F). The relative solubility of sirenian Hbs is denoted by the percentage of Hb
protein precipitated by the addition of 3 M ammonium sulfate.
594

595
596

597
598
599

Figure 3. Homology models of the DPG binding site in ancestral dugongid and Steller's sea cow
(*H. gigas*) hemoglobin. A) Model of the ancestral dugongid DPG binding site. Amino acids are
colored according to the degree of sequence conservation. Notably, β/δ82Lys shows the highest
level of sequence conservation, as it is able to bind to multiple sites on the DPG molecule (indicated
by dashed pink lines), whereas β/δ2His is only able to directly interact with DPG in the protonated
state. B) Model of Steller's sea cow hemoglobin (left) and a close-up of the DPG binding pocket
(right). Dark blue colored residues represent the 11 *H. gigas* specific substitutions, while those in
light blue denote the ancestral state. Homology modelling illustrates how the replacement of
β/δ82Lys with neutral Asn inhibits DPG binding to the hemoglobin molecule.