Rodent trapping studies as an overlooked information source for understanding endemic and novel zoonotic spillover.

Rodent trapping to understand zoonotic spillover risk

David Simons[1, 2, 3]

Lauren A. Attfield[3, 5]

Kate E. Jones[3]

Deborah Watson-Jones[2, 4]

Richard Kock[1]

[1] Centre for Emerging, Endemic and Exotic Diseases, The Royal Veterinary College, London, UK

[2] Clinical Research Department, London School of Hygiene and Tropical Medicine, London, UK

[3] Centre for Biodiversity and Environment Research, University College London, London, UK

[4] Mwanza Intervention Trials Unit, National Institute for Medical Research, Mwanza, Tanzania

[5] Department of Life Sciences, Imperial College London, London, UK.

1. Abstract

Rodents a diverse, globally distributed and ecologically important order of mammals are nevertheless important reservoirs of known and novel zoonotic pathogens. Ongoing anthropogenic land use change is altering the composition of these species' abundance and distribution which among zoonotic host species may alter the risk of zoonoses spillover events. These changes demand that a better understanding of the current distribution of rodent species is vital for mitigating potential zoonotic disease hazard and managing risk. However, available species distribution and host-pathogen association datasets (e.g. IUCN, GBIF, CLOVER) are often taxonomically and spatially biased. Here, we synthesise data from West Africa from 127 rodent trapping studies, published between 1964-2022, as an additional source of information to characterise the range and presence of rodent species and identify the subgroup of species that are potential or known pathogen hosts. We identify that these rodent trapping studies, although biased towards human dominated landscapes across West Africa, can usefully complement current rodent species distribution datasets and we calculate the discrepancies between these datasets. For five regionally important zoonotic pathogens (Arenaviridae spp., Borrelia spp., *Lassa mammarenavirus*, Leptospira spp. and *Toxoplasma gondii*), we identify host-pathogen associations that have not been previously reported in host-association datasets. Finally, for these five pathogen groups, we find that the proportion of a rodent hosts range that has been sampled remains small with geographic clustering. A priority should be to sample rodent hosts across a greater geographic range to better characterise current and future risk of zoonotic spillover events. In the interim, studies of spatial pathogen risk informed by rodent distributions must incorporate a measure of the current sampling biases. The current synthesis of contextually rich rodent trapping data enriches available information from IUCN, GBIF and

38 CLOVER which can support a more complete understanding of the hazard of zoonotic spillover

39 events.

2. Author Summary

Emerging and endemic zoonotic diseases are projected to have increasing health impacts,

particularly under changing climate and land-use scenarios. Rodents, an ecologically vital order

of mammals carry a disproportionate number of zoonotic pathogens and are abundant across

West Africa. Prior modelling studies rely on large, consolidated data sources which do not

incorporate high resolution spatial and temporal data from rodent trapping studies. Here, we

synthesise these studies to quantify the bias in the sampling of rodent hosts and their

pathogens across West Africa. We find that rodent trapping studies are complementary to these

datasets and can provide additional, high-resolution data on the distribution of hosts and their

pathogens. Further, rodent trapping studies have identified additional potential host-pathogen

associations than are recorded in consolidated host-pathogen association datasets. This can

help to understand the risk of zoonotic diseases based on host distributions. Finally, we

quantify the current extent of known rodent presence and pathogen sampling within a species

range, highlighting that current knowledge is limited across much of the region. We hope that

this will support work to study rodent hosts and their pathogens in currently under sampled

regions to better understand the risk of emerging and endemic zoonoses in West Africa.

3. Introduction

There is increasing awareness of the global health and economic impacts of novel zoonotic pathogen spillover, driven by the ongoing SARS-CoV-2 pandemic and previous HIV/AIDs and Spanish Influenza pandemics [1]. The number of zoonotic disease spillover events and the frequency of the emergence of novel zoonotic pathogens from rodents are predicted to increase under intensifying anthropogenic pressure driven by increased human populations, urbanisation, intensification of agriculture, climate change leading to altered rodent species assemblages [2–5]. The impact of endemic zoonoses meanwhile remains underestimated [6]. Endemic zoonoses disproportionally affect those in the poorest sections of society, those living in close contact with their animals and those with limited access to healthcare [7–9].

Rodents along with bats contribute the greatest number of predicted novel zoonotic pathogens and known endemic zoonoses [10,11]. Of 2,220 extant rodent species, 244 (10.7%) are described as reservoirs of 85 zoonotic pathogens [10]. Most rodent species do not provide a direct risk to human health and all species provide important and beneficial ecosystem services including pest regulation and seed dispersal [12]. Increasing risks of zoonotic spillover events are driven by human actions rather than by rodents, for example, invasive rodent species being introduced to novel ranges through human transport routes. Rodents typically demonstrate "fast" life histories which allow them to exploit opportunities provided by anthropogenic disturbance [13]. Within rodents, species level traits such as early maturation and short gestation times are associated with increased probabilities of being zoonotic reservoirs [10,14]. Rodent species with these traits are able to thrive in human dominated landscapes, displacing species less likely to be reservoirs of zoonotic pathogens [15]. The widespread occurrence of reservoir species and their proximity to human activity make the description of rodent species

assemblages and host-pathogen associations vitally important to understanding the hazard of

zoonotic disease spillover and novel zoonotic pathogen emergence [16].

Despite the importance of understanding these complex systems, current evidence on host-

pathogen associations is considerably affected by taxonomic and geographical sampling biases

[11,17]. Curated biodiversity datasets such as the Global Biodiversity Information Facility

(GBIF) and resources produced by the International Union for Conservation of Nature (IUCN)

suffer from well described spatial and temporal sampling biases [18,19]. These data are

typically obtained from museum specimen collections and non-governmental organisation

surveys. These sampling biases can importantly distort produced species distribution models

that are used to infer risk of zoonotic disease risk [20]. Datasets on host-pathogen associations

(i.e. CLOVER) also can suffer from biases introduced from literature selection criteria and

taxonomic discrepancies resulting in differential likelihood of accurate host-pathogen

attribution by host species. These biases are important because identification of potential

geographic hotspots of zoonotic disease spillover and novel pathogen emergence are often

produced from these types of host species distributions and host-pathogen associations

[21,22]. For example, systematically increased sampling, over-representation of certain

habitats and clustering around areas of high human population could lead to an apparent

association between locations and hazard that is driven by these factors rather than underlying

host-pathogen associations [11,23,24]. Predictions of zoonotic disease spillover and novel

zoonotic pathogen emergence must account for these biases to understand the future hazard of

zoonotic diseases [22].

West Africa has been identified as a region at increased risk for rodent-borne zoonotic disease

spillover events, the probability of these events are predicted to increase under different

projected future land-use change scenarios [4,25]. Currently within West Africa, some rodent

species are known to be involved in the transmission of multiple endemic zoonoses with large

burdens on human health, these pathogens include Lassa fever, Schistosomiasis, Leptospirosis

and Toxoplasmosis [26,27]. The presence of other species within shared habitats may mitigate

the spread of these pathogens through the "dilution effect", ongoing loss of biodiversity may

further increase the risk to human populations [5]. Understanding of the distribution of these

zoonoses are limited by biases in consolidated datasets. Rodent trapping studies provide

contextually rich information on when, where and under what conditions rodents were

trapped, potentially enriching consolidated datasets [28]. Studies have been conducted in West

Africa to investigate the distribution of rodent species, their species assemblages, the

prevalence of endemic zoonoses within rodent hosts (e.g., Lassa fever, Schistosomiasis) and to

identify emerging and novel zoonotic pathogens [29–31]. However, individual level data from

these studies have not previously been synthesised for inclusion in assessments of zoonotic

disease spillover and novel zoonotic pathogen emergence.

Here, we synthesise rodent trapping studies conducted across West Africa published between

1964-2022. First, we use this dataset to investigate the geographic sampling biases in relation

to human population density and land use classification. Second, we compare this to curated

host datasets (IUCN and GBIF) to understand differences in reported host geographic

distributions. Third, we compare identified host-pathogen associations with a consolidated

dataset (CLOVER) to explore discrepancies in rodent host-pathogen associations and report the

proportion of positive assays for pathogens of interest. Finally, within our dataset we

investigate the spatial extent of current host-pathogen sampling to identify areas of sparse

sampling of pathogens within their host ranges. We expect that rodent trapping studies provide

an important additional source of high-resolution data that can be used to enrich available

consolidated datasets to better understand the hazard of zoonotic disease spillover and novel

zoonotic pathogen emergence across West Africa.

128 4. Methods

129 4.1. Data sources

130 4.1.1. Host and pathogen trapping data

To identify relevant literature, we conducted a search in Ovid MEDLINE, Web of Science (Core

collection and Zoological Record), JSTOR, BioOne, African Journals Online, Global Health and the

pre-print servers, BioRxiv and EcoEvoRxiv for the following terms as exploded keywords: (1)

Rodent OR Rodent trap* AND (2) West Africa, no date limits were set. We also searched other

resources including the UN Official Documents System, Open Grey, AGRIS FAO and Google

Scholar using combinations of the above terms. Searches were run on 2022-05-01, and

returned studies conducted between 1964-2021.

We included studies for further analysis if they met all of the following inclusion criteria; i)

reported findings from trapping studies where the target was a small mammal, ii) described the

type of trap used or the length of trapping activity or the location of the trapping activity, iii)

included trapping activity from at least one West African country, iv) recorded the genus or

species of trapped individuals, and v) were published in a peer-reviewed journal or as a pre-

print on a digital platform or as a report by a credible organisation. We excluded studies if they

met any of the following exclusion criteria: i) reported data that were duplicated from a

previously included study, ii) no full text available, iii) not available in English. One author

screened titles, abstracts and full texts against the inclusion and exclusion criteria. At each

stage; title screening, abstract screening and full text review, a random subset (10%) was

reviewed by a second author.

We extracted data from eligible studies using a standardised tool that was piloted on 5 studies

(Supplementary Table 1.). Data was abstracted into a Google Sheets document, which was

archived on completion of data extraction [32]. We identified the aims of included studies, for

example, whether it was conducted as a survey of small mammal species or specifically to

assess the risk of zoonotic disease spillover. we extracted data on study methodology, such as,

the number of trap nights, the type of traps used and whether the study attempted to estimate

abundance. For studies not reporting number of trap nights we used imputation based on the

number of trapped individuals, stratified by the habitat type from which they were obtained.

This was performed by multiplying the total number of trapped individuals within that study

site by the median trap success for study sites with the same reported habitat type.

Stratification was used as trap success varied importantly between traps placed in or around

buildings (13%, IQR 6-24%) compared with other habitats (3%, IQR 1-9%)

We also recorded how species were identified within a study and species identification was

assumed to be accurate. The number of individuals of these species or genera was extracted

with taxanomic names mapped to GBIF taxonomy [33]. We expanded species detection and

non-detection records by explicitly specifying non-detection at a trap site if a species was

recorded as detected at other trapping locations within the same study.

Geographic locations of trapping studies were extracted using GPS locations for the most

precise location presented. Missing locations were found using the National Geospatial-

Intelligence Agency GEOnet Names Server [34] based on placenames and maps presented in the

study. All locations were converted to decimal degrees. The year of rodent trapping was

extracted alongside the length of the trapping activity to understand seasonal

representativeness of trapping activity. The habitats of trapping sites were mapped to the IUCN

Habitat Classification Scheme (Version 3.1). For studies reporting multiple habitat types for a

single trap, trap-line or trapping grid, a higher order classification of habitat type was recorded.

For included studies with available data we extracted information on all microorganisms and

known zoonotic pathogens tested and the method used (e.g., molecular or serological

diagnosis). Where assays were able to identify the microorganism to species level this was

recorded, non-specific assays higher order attribution was used (e.g. to family level). A broad

definition of known zoonotic pathogen was used, a species of microorganism carried by an

animal that may transmit to humans and cause illness [35]. We do not include evolved

pathogens acquired originally through zoonotic pathways in our definition (i.e. HIV). The term

microorganism is used where either the microorganism is not identified to species level, in

which case it remains unclear whether it is a zoonotic pathogen (i.e. Arenaviridae), or the

species is not known to be a zoonotic pathogen (i.e. *Candidatus Ehrlichia senegalensis*). We

recorded the species of rodent host tested, the number of individuals tested and the number of

positive and negative results. For studies reporting summary results all testing data were

extracted, this may introduce double counting of individual rodents, for example, if a single

rodent was tested using both molecular and serological assays. Where studies reported

indeterminate results, these were also recorded.

4.1.2. Description of included studies

Out of 4,692 relevant citations, we identified 127 rodent trapping studies (Supplementary Table 2.). Of these, 55 (43%) were conducted to investigate rodent-borne zoonoses, with the remaining 77 (57%) conducted for ecological purposes (i.e. population dynamics, distribution) in rodents, including those known to be hosts of zoonotic pathogens. The earliest trapping studies were conducted in 1964, with a trend of increasing numbers of studies being performed annually since 2000. The median year of first trapping activity was 2007, with the median length of trapping activity being 1 year (IQR 0-2 years) (Supplementary Fig 1.). Studies were conducted in 14 West African countries, with no studies reported from The Gambia or Togo, at 1,611 trap sites (Fig 1A.).

Included studies explicitly reported on 601,184 trap nights, a further 341,445 trap nights were imputed from studies with no recording of trapping effort based on trap success, leading to an estimate of 942,629 trap nights (Fig 1B.). A minority of studies trapped at a single study site (30, 24%), with 46 (36%) trapping at between two and five sites, the remaining 51 studies (40%) trapped at between six and 93 study sites.

In total 76,275 small mammals were trapped with 65,628 (90%) identified to species level and 7,439 (10%) identified to genus, with the remaining classified to higher taxonomic level. The majority of the 132 identified species were Rodentia (102, 78%), of which Muridae (73, 72%) were the most common family. Soricomorpha were the second most identified order of small mammals (28, 21%). 57 studies tested for 32 microorganisms, defined to species or genus level that are known or potential pathogens. Most studies tested for a single microorganism (48, 84%). The most frequently assayed microorganisms were *Lassa mammarenavirus* or

Arenaviridae (21, 37%), *Borrelia sp.* (9, 16%), *Bartonella sp.* (4, 7%) and *Toxoplasma gondii* (4,

7%). Most studies used Polymerase Chain Reaction (PCR) to detect microorganisms (37, 65%),

with fewer studies using serology-based tests (11, 19%) or histological or direct visualisation

assays (11, 21%). From 32,920 individual rodent samples we produced 351 host-pathogen

pairs. With *Rattus rattus*, *Mus musculus*, *Mastomys erythroleucus*, *Mastomys natalensis* and

Arvicanthis niloticus being assayed for at least 18 microorganisms.

Fig 1: Rodent trapping sites across West Africa. A) The location of trapping sites in West

Africa. No sites were recorded from Togo or The Gambia. Heterogeneity is observed in the

coverage of each country by trap night (colour) and location of sites. For example, Senegal, Mali

and Sierra Leone have generally good coverage compared to Guinea and Burkina Faso. B)

Histogram of trap nights performed at each study site, a median of 248 trap nights (IQR 116-

500) was performed at each site. A labelled map of the study region is attached in

224   Supplementary Figure 5.

4.2. Analysis

4.2.1. What is the extent of spatial bias in the rodent trapping data?

To investigate the extent of spatial bias in the rodent trapping data, we calculated trap-night (TN) density within each West African level-2 administrative region. The `sf` package in the R statistical language (R version 4.1.2) was used to manipulate geographic data, administrative boundaries were obtained from GADM 4.0.4 [36–38]. Trap-night density ($TN_{density}$) was calculated by dividing the number of trap nights by the area of a level-2 administrative area (R_{area}). For studies not reporting trap nights, imputation was used as previously described. Human population density was obtained for the closest year (2005) to the median year of trapping (2007) from Socioeconomic Data and Applications Center (SEDAC) gridded population of the world v4 at ~ 1km resolution ($P_{density}$) [39]. Median population density was then calculated for each level-2 administrative region. Land cover classification was obtained from the Copernicus climate change service at ~300m resolution [40]. The proportion of cropland, shrubland, tree cover (ψ_{tree}) and urban land cover (ψ_{urban}) within a level-2 administrative region in 2005 was calculated.

We investigated the association between relative trapping effort, measured as TN density, and the proportion of urban, cropland, tree cover and human population density using Generalised Additive Models (GAM) incorporating a spatial interaction term (longitude and latitude) [41]. Spatial aggregation of relative trapping effort was modelled using an exponential dispersion distribution (Tweedie) [42]. The models were constructed in the `mgcv` package [43]. Selection of the most parsimonious model was based on Deviance explained and the Akaike Information Criterion for each model (Equations 1-5 below). Relative trapping effort was then predicted

247 across West Africa using these covariates. We performed two sensitivity analyses, first, by

248 removing sites with imputed trapping effort, second, by associating trap locations to ~1km

249 pixels rather than level-2 administrative areas.

$$\mathrm{TN_{density}} \sim \mathrm{Tweedie}(X * Y) \#(1)$$

$$\mathrm{TN_{density}} \sim \mathrm{Tweedie}\left(\mathrm{P_{density}} + (X * Y)\right) \#(2)$$

$$\mathrm{TN_{density}} \sim \mathrm{Tweedie}\left(\mathrm{P_{density}} + \mathrm{R_{area}} + (X * Y)\right) \#(3)$$

$$\mathrm{TN_{density}} \sim \mathrm{Tweedie}\left(\mathrm{P_{density}} + \psi_{\mathrm{tree}} + \psi_{\mathrm{urban}} + (X * Y)\right) \#(4)$$

$$\mathrm{TN_{density}} \sim \mathrm{Tweedie}\left(\mathrm{P_{density}} + \mathrm{R_{area}} + \psi_{\mathrm{urban}} + (X * Y)\right) \#(5)$$

4.2.2. What is the difference in rodent host distributions between curated

datasets and rodent trapping studies?

We assessed the concordance of curated rodent host distributions from IUCN and GBIF with

observed rodent detection and non-detection from rodent trapping studies for seven species

with the most trap locations (*M. natalensis*, *R. rattus*, *M. erythroleucus*, *M. musculus*, *A. niloticus*,

Praomys daltoni and *Cricetomys gambianus*). We obtained rodent species distribution maps as

shapefiles from the IUCN red list and translated these to a ~20km resolution raster [44].

Distributions were cropped to the study region for globally distributed rodent species. We

obtained rodent presence locations from GBIF as point data limited to the study region [45].

Presence locations were associated to cells of raster with a ~20km resolution produced for the
study region.

For each of the seven species, we first calculated the area of the IUCN expected range, and then
the percentage of this range covered by presence detections in GBIF, and from detections in the
rodent trapping data. We then calculated the area of both types of both GBIF and rodent
trapping detections outside of the IUCN expected range. For rodent trapping data, we
additionally calculated the area of non-detections within the IUCN expected area. Finally, we
calculated the combined area of detection from both GBIF and rodent trapping data.

4.2.3. Are rodent trapping derived host-pathogen associations present in a consolidated zoonoses dataset?

To examine the usefulness of rodent trapping studies as an additional source of data we
compared identified host-pathogen associations from trapping studies investigating zoonoses
with a consolidated zoonoses dataset (CLOVER) [11,46]. CLOVER is a synthesis of four host-
pathogen datasets (GMPD2, EID2, HP3 and Shaw, 2020) and was released in 2021, it contains
more than 25,000 host-pathogen associations for Bacteria, Viruses, Helminth, Protozoa and
Fungi. We compared the host-pathogen networks across the two datasets, where the CLOVER
data was subset for host species present in the rodent trapping data.

For host-pathogen pairs with assay results consistent with acute or prior infection, we
calculated the proportion positive and identify those absent from CLOVER. We expand the
analysis to host-pathogen pairs with pathogens identified to genus level in Supplementary Fig
4.

4.2.4. What is the spatial extent of pathogen testing within host ranges?

We use the sampled area of three pathogen groups and two pathogens (Arenaviridae, Borreliaceae, Leptospiraceae, *Lassa mammarenavirus* and *Toxoplasma gondii*) to quantify the bias of sampling within their hosts ranges. For each pathogen, we first describe the number of host species assayed, for the five most commonly tested species we associate the locations of sampled individuals to ~20km pixels and calculate the proportion of the IUCN range of the host in which sampling has occurred. We compare this figure to the total area in which the host has been detected to produce a measure of relative completeness of sampling within the included rodent trapping studies.

Data and code to reproduce all analyses are available in an archived Zenodo repository [32].

5. Results

5.1.   What is the extent of spatial bias in the rodent trapping data?

We found non-random, spatial clustering of rodent trapping locations across the study region, suggestive of underlying bias in the sampling or rodents across West Africa. Trap sites were situated in 256 of 1,450 (17.6%) level-2 administrative regions in 14 West African nations. The regions with the highest TN density included the capitals and large cities of Niger (Niamey), Nigeria (Ibadan), Ghana (Accra), Senegal (Dakar), and Benin (Cotonou). Outside of these cities, regions in, Northern Senegal, Southern Guinea, Edo and Ogun States in Nigeria and Eastern Sierra Leone had the greatest TN density (Fig 1A.).

The most parsimonious GAM model (adjusted R2 = 0.3, Deviance explained = 48.7%) reported

significant non-linear associations between relative trapping effort bias and human population

densities (Effective Degrees of Freedom (EDF) = 7.13, $p < 0.001$), proportion of urban landscape

(EDF = 1.92, $p < 0.002$) and region area (EDF = 3.63, $p < 0.001$), alongside significant spatial

associations (EDF = 27.3, $p < 0.001$) (Supplementary table 3.1). Greatest trapping effort bias

peaked at population densities between 5,000-7,500 individuals/km^2, proportion of urban

landscape >10% and region areas < 1,000km^2. Increased trapping effort was found in North

West Senegal, North and East Sierra Leone, Central Guinea and coastal regions of Nigeria, Benin

and Ghana; in contrast South East Nigeria, Northern Nigeria and Burkina Faso had an observed

bias towards a reduced trapping effort (Fig 2). In sensitivity analysis, excluding sites with

imputed trap nights, Mauritania, Northern Senegal and Sierra Leone remained as regions

trapped at higher rates, with Nigeria being trapped at lower than expected rates

(Supplementary Fig 3a.). In pixel-based sensitivity analysis spatial coverage was reduced with

similar patterns of bias observed to the primary analysis (Supplementary Fig 3b.).

Fig 2. **Relative trapping effort bias across West Africa**. Modelled relative trapping effort bias

316   adjusted for human population density, proportion urban land cover and area of the

317   administrative region. Brown regions represent areas with a bias towards increased trapping

318   effort (e.g., North West Senegal), Green regions represent areas with a bias towards reduced

319   trapping effort (e.g., Northern Nigeria).

5.2. What is the difference in rodent host distributions between curated datasets and rodent trapping studies?

We found that for six of the seven most detected rodent species (*M. natalensis*, *R. rattus*, *M. erythroleucus*, *M. musculus*, *A. niloticus* and *P. daltoni*), trapping studies provided more distinct locations of detection and non-detection than were available from GBIF. For the endemic rodent species (*M. natalensis*, *M. erythroleucus*, *A. niloticus*, *P. daltoni* and *C. gambianus*) IUCN ranges had good concordance to both trapping studies and GBIF, however, individuals of *A. niloticus* and *P. daltoni* were detected outside of IUCN ranges. In contrast, the non-native species *R. rattus* and *M. musculus* were detected across much greater ranges than were expected from IUCN distributions. Comparisons for *M. natalensis*, *R. rattus* and *M. musculus* are shown in Fig 3, the remaining species are shown in Supplementary Fig 4.

Mastomys natalensis

Rattus rattus

Mus musculus

Detection/Non-detection   ● Detection   ● Non-detection

Fig 3. Locations of detection and non-detection sites for rodent species in West Africa

Each row corresponds to a single rodent species. L) Presence recorded in GBIF (black points)

overlaid on IUCN species range (red-shaded area). R) Detection (purple) and non-detection

(orange) from rodent trapping studies overlaid on IUCN species ranges. M. musculus has no

IUCN West African range.

Comparison of the proportion of a species IUCN range in which detections and non-detections

occurred showed that sampling locations of these seven species within GBIF covered between

0.09-0.26% of expected ranges (Table 1.), compared to 0.03-0.24% for rodent trapping data.

Detections occurred outside IUCN ranges for all species in both the GBIF and rodent trapping

data, most noticeably for *A. niloticus* and *R. rattus*. Combining GBIF and rodent trapping data

increased the sampled area by a mean of 1.6 times compared to the GBIF area alone,

demonstrating limited overlap between the locations providing information to either dataset.

Non-detection of a species occurred across species ranges (mean = 0.11%, SD = 0.03%),

suggestive of spatial heterogeneity of presence within IUCN ranges.

    Table 1: Comparison of IUCN, GBIF and rodent trapping ranges for the 7 most detected

                                          rodent species.

Species	IUCN Range (1,000 km^2)	GBIF Area inside range (1,000 km^2) (% of IUCN range)	GBIF Area outside range (1,000 km^2)	Trapping studies Detection area inside range (1,000 km^2) (% of IUCN range)	Trapping studies Area outside range (1,000 km^2) (% of IUCN range)	Trapping studies Non-detection area inside range (1,000 km^2) (% of IUCN range)	Combined Detection area (1,000 km^2) (% of IUCN range)
Mastomys natalensis	3,257	6.83 (0.21%)	0.19	4.4 (0.14%)	0.17	3.12 (0.1%)	10.66 (0.33%)
Rattus rattus	1,019	2.61 (0.26%)	0.52	2.42 (0.24%)	1.21	1.3 (0.13%)	4.88 (0.48%)
Mastomys erythroleucus	3,735	4.48 (0.12%)	0.04	3.24 (0.09%)	0.12	4.35 (0.12%)	7.5 (0.2%)
Mus musculus			2.15		1.85		3.94
Arvicanthis niloticus	1,829	1.69 (0.09%)	2.41	1.98 (0.11%)	0.34	3.09 (0.17%)	3.59 (0.2%)
Praomys daltoni	2,658	4.03 (0.15%)	0.29	2.03 (0.08%)	0.15	2.78 (0.1%)	5.83 (0.22%)
Cricetomys gambianus	2,476	5 (0.2%)	0.17	0.75 (0.03%)	0.06	2.99 (0.12%)	5.69 (0.23%)

5.3.   Are rodent trapping derived host-pathogen associations present in a consolidated zoonoses dataset?

We found potentially important differences between the host-pathogen networks produced

from included rodent trapping studies and the consolidated CLOVER dataset. When limited to

taxonomic classification of both pathogen and host to species level we identified 25 host-

pathogen pairs among 14 rodent and 6 pathogen species (Fig 4. and Fig 5.). We identified

negative associations (non-detection through specific assays) for 45 host-pathogen pairs

among 35 rodent and 7 pathogen species. CLOVER contained 10 (40%) of our identified host-

pathogen associations, the remaining 15 (60%) were not found to be present in CLOVER,

additionally CLOVER recorded positive associations for 4 (9%) of the negative associations

produced from the rodent trapping data.

Fig 4. Host-Pathogen associations detected through acute infection. A) Identified species

level host-pathogen associations through detection of acute infection (i.e. PCR, culture).

Percentages and colour relate to the proportion of all assays that were positive, the number of

individuals tested for the pathogen is labelled N. Associations with a black border are present in

364   the CLOVER dataset.

Fig 5. Host-Pathogen associations detected through evidence of prior infection. B)

Identified species level host-pathogen associations through serological assays (i.e. ELISA).

Percentages and colour relate to the proportion of all assays that were positive, the number of

individuals tested for the pathogen is labelled N. Associations with a black border are present in

the CLOVER dataset.

CLOVER included an additional 492 host-pathogen associations we do not observe in rodent

trapping studies. The majority of these 392 (80%) pairs are from species with global

distributions (*M. musculus*, *R. rattus* and *R. norvegicus*), or from those with wide ranging

distributions in sub-Saharan Africa (38, 8%) (i.e., *A. niloticus*, *M. natalensis* and *Atelerix*

albiventris).

For pathogens not identified to species level (i.e. family or higher taxa only), we identified 148

host-pathogen pairs among 32 rodent species and 25 pathogen families (Supplementary Fig 4.),

with CLOVER containing 66 (45%) of these associations.

Rodent trapping studies identified additional rodent host species for six pathogens; *Lassa*

mammarenavirus (5), *Toxoplasma gondii* (4), Usutu virus (2), *Coxiella burnetii* (2), *Escherichia*

coli and *Klebsiella pneumoniae* (both 1), that were not present in this consolidated host-

pathogen association dataset.

5.4.   What is the spatial extent of microorganism testing within a

host's range?

The five most widely sampled microorganism species/families in included studies were

Arenaviridae, Borreliaceae, *Lassa mammarenavirus*, Leptospiraceae and *Toxoplasma gondii*

(Table 2.). Assays to identify Arenaviridae infection were performed in 44 rodent species with

evidence of viral infection in 15 species. Studies that reported Arenaviridae infection did not

identify the microorganism to species level and were distinct from those reporting *Lassa*

mammarenavirus infection. *Lassa mammarenavirus* was specifically tested for in 43 species

with 10 showing evidence of viral infection. The most commonly infected species for both

Arenaviridae, generally, and *Lassa mammarenavirus* specifically, were *M. natalensis* and *M.*

erythroleucus. These species were assayed across between 10-20% of their trapped area,

equating to ~0.02% of their IUCN range (Table 2.).

Infection with species of Borreliaceae was assessed in 42 species, with evidence of infection in

17 rodent species. The greatest rates of infection were among *A. niloticus* (16%), *Mastomys*

huberti (11%) and *M. erythroleucus* (9%). Testing was more widespread than for Arenviruses

with coverage between 15-34% of their trapped area, however, this remains a small area in

relation to their IUCN ranges (<0.05%). Leptospiraceae and *Toxoplasma gondii* was assessed in

8 species, with evidence of infection in 5 and 6 rodent species respectively. The spatial coverage

of testing for these microorganisms was more limited within IUCN host species ranges

(~0.01%).

Table 2: Comparison of microorganism sampling ranges for the 5 most widely sampled

microorganisms and the 5 most sampled rodent host species (* no IUCN range in West

African).

Microorganism	Host species	Tested (N)	Positive (N (%))	*Microorganism* testing area (1,000 km^2)	*Microorganism* testing area within trapped area (%)	*Microorganism* testing area within IUCN range (%)
Arenaviridae sp.						
	Mastomys natalensis	2,841	104 (4%)	0.61	13.45%	0.02%
	Praomys daltoni	854	6 (1%)	0.42	19.43%	0.02%
	Mastomys erythroleucus	398	20 (5%)	0.40	11.97%	0.01%

Microorganism	Host species	Tested (N)	Positive (N (%))	Microorganism testing area (1,000 km^2)	Microorganism testing area within trapped area (%)	Microorganism testing area within IUCN range (%)
Borrelia sp.	Rattus rattus	396	4 (1%)	0.38	10.5%	0.04%
	Praomys rostratus	310	5 (2%)	0.13	12.53%	0.02%
	Mastomys erythroleucus	1,586	140 (9%)	1.14	33.94%	0.03%
	Arvicanthis niloticus	1,551	253 (16%)	0.66	28.48%	0.03%
	Mastomys natalensis	733	54 (7%)	0.69	15.08%	0.02%
	Mastomys huberti	731	83 (11%)	0.23	29.83%	0.04%
	Mus musculus	686	26 (4%)	0.45	24.54%	*
Lassa mammarenavirus	Mastomys natalensis	3,199	580 (18%)	1.03	22.65%	0.03%
	Mastomys erythroleucus	352	14 (4%)	0.36	10.63%	0.01%
	Rattus rattus	177	2 (1%)	0.34	9.26%	0.03%
	Praomys rostratus	163	2 (1%)	0.27	27.02%	0.04%
	Mus musculus	147	0 (0%)	0.04	2.29%	*
Leptospira sp.	Rattus rattus	646	65 (10%)	0.40	11.1%	0.04%

Microorganism	Host species	Tested (N)	Positive (N (%))	Microorganism testing area (1,000 km²)	Microorganism testing area within trapped area (%)	Microorganism testing area within IUCN range (%)
Toxoplasma gondii	Arvicanthis niloticus	221	10 (5%)	0.02	0.9%	<0.01%
	Crocidura olivieri	141	14 (10%)	0.34	25.16%	*
	Mastomys natalensis	136	26 (19%)	0.36	7.91%	0.01%
	Rattus norvegicus	79	19 (24%)	0.21	40.08%	*
	Mus musculus	1,548	115 (7%)	0.62	33.64%	*
	Rattus rattus	428	8 (2%)	0.36	9.77%	0.03%
	Mastomys erythroleucus	292	13 (4%)	0.37	11.06%	0.01%
	Mastomys natalensis	107	2 (2%)	0.08	1.83%	<0.01%
	Cricetomys gambianus	47	13 (28%)	0.06	7.6%	<0.01%

6. Discussion

Endemic rodent zoonoses and novel pathogen emergence from rodent hosts are predicted to have an increasing burden in West Africa and globally [10]. Here we have synthesised data from 126 rodent trapping studies containing information on more than 72,000 rodents, from at least 132 species of small mammals, across 1,611 trap sites producing an estimated 942,669 trap nights across 14 West African countries. Locations studied are complementary to curated datasets (e.g. IUCN, GBIF), incorporation of our synthesised dataset when assessing zoonosis

risk based on host distributions could counteract some of the biases inherent to these curated

datasets [18]. Most assayed rodents were not found to be hosts of known zoonotic pathogens.

We identified 25 host-pathogen pairs reported from included studies, 15 of these were not

included in a consolidated host-pathogen dataset. Generally, the number of different species

tested for a microorganism and the spatial extent of these sampling locations were limited.

These findings highlight a number of sampling bias, supporting calls for further microorganism

sampling across diverse species in zoonotic hotspots [47].

We found that rodent trapping data, like biodiversity data, showed important spatial biases

[20]. Relative trapping effort bias was greater in Benin, Guinea, Senegal and Sierra Leone driven

by long-standing research collaborations investigating the invasion non-native rodent species

(*M. musculus* and *R. rattus*) and the hazard of endemic zoonosis outbreaks (e.g., *Lassa*

mammarenavirus). In addition to identifying point locations of prior rodent and pathogen

sampling (Figure 1.), additional information on the trapping effort (density of trap-nights),

human population density and land use type have been incorporated to produce a value of

relative effort that will assist researchers in identifying specific locations where predictions

based on these underlying data sources may suffer from increased effects of trapping effort

bias. This approach improves the ease of identifying under sampled locations, for example,

Figure 1. may suggest that South East Senegal, Southern Mali and Southern Niger are well

sampled based on locations of trapping sites. When the number of trap nights, human

population density and land use of these regions are taken into account (Figure 2.) and

compared with better sampled locations (i.e. Western Senegal, Eastern Sierra Leone) these

areas are found to be relatively under sampled and would benefit from further sampling effort.

This contrasts to North West Nigeria where no trapping has occurred (Figure 1.), the modelling

approach has perhaps highlighted this beyond a simple map of currently trapped locations as

an immediate priority for sampling of rodents and their pathogens given high human

population densities and a human dominated landscape.

Much of West Africaremains relatively under sampled, particularly Burkina Faso, Côte d'Ivoire,

Ghana and Nigeria, despite these countries facing many of the same challenges. For example,

annual outbreaks of Lassa fever are reported in Nigeria and there are potentially 60,000

unrecognised cases of Lassa fever every year in Côte d'Ivoire and Ghana [48]. Our estimates of

the proportion of a rodent species range that have been sampled along with pathogen testing

within their sampled range are sensitive to our choice of raster cell size, smaller area cells will

reduce the reported coverage while larger cells will have the opposite effect. Despite this, the

observed patterns are unlikely to importantly change, with the finding of sparse sampling of

both rodents and their pathogens remaining present across cell scales. Rodent sampling should

be targeted towards currently under sampled regions to reduce the potential impact of this bias

and improve our understanding of both the distribution of rodent hosts and the prevalence of

pathogens within their populations. This will allow for better estimation of risk from endemic

and novel zoonoses.

Rodent trapping studies provide geographic and temporally contextualised data on both

species detection and non-detection which are not available from curated datasets. Non-

detection data can improve models of species distributions, unfortunately, high levels of

missing data on trapping effort will continue to confound the allocations of non-detections as

true absences [49]. Models of host species occurrence and abundance, improved by

incorporating species absence, are important to assess the effect of land use and climate change

on endemic zoonosis spillover to human populations and direct limited public health resources

towards regions at greatest risk [50,51].

Currently available consolidated datasets on host-pathogen associations (e.g. CLOVER, EID2

and GMPD2) do not include spatial or temporal components [52]. The current synthesis of

rodent trapping studies has highlighted that pathogens have been sparsely sampled within a

host's range. Current zoonosis risk models dependent on these sources of data are therefore

not able to incorporate spatial heterogeneity in pathogen prevalence across the host range.

Additional uncertainty in current models of zoonotic disease risk arises from host-pathogen

associations that have not been reported in these consolidated datasets. For example,

Hylomyscus pamfi infected with *Lassa mammarenavirus* and *R. rattus* infected with *Coxiella*

burnetii, will not be included when solely based on consolidated host-pathogen datasets.

Further, detection of zoonotic pathogens in multiple co-occurring host species supports the

adoption of multi-species to better understand the potential range of endemic zoonoses [53].

Few studies stratified detection and non-detection of hosts or pathogen prevalence by time,

therefore limiting inference of changes in host and pathogen dynamics. This limitation prevents

calculation of incidence of infection and the abundance of infectious rodents which potentially

varies by both time and space [54]. Understanding of temporal changes in viral burden and

shedding for endemic zoonoses is required to accurately predict current and future risk of

pathogen spillover.

Finally, due to data sparsity, we were unable to account for temporal change over the six

decades of rodent trapping studies. Land use change and population density have changed

dramatically over this period in West Africa [55]. We attempted to mitigate against this by

using the median year of trapping to understand the spatial and land use biases in trapping

activity. It is possible that land use and population density at trapping sites varied importantly

between when rodent trapping was conducted and the conditions in 2005. Despite this

limitation, the finding that trapping is biased towards high density, human dominated

landscapes is unlikely to substantially change.

We have shown that synthesis of rodent trapping studies to supplement curated rodent

distributions can counteract some of the inherent biases in these data and that they can add

further contextual data to host-pathogen association data. Together this supports their

inclusion in efforts to model endemic zoonotic risk and novel pathogen emergence.

Contribution of rodent trapping studies as data sources can be improved by adopting reporting

standards and practices consistent with Open Science, namely sharing of disaggregated

datasets alongside publication [56].

Future rodent trapping studies should be targeted towards regions that are currently under-

studied. Further information on rodent presence and abundance across West Africa will aid the

modelling of changing endemic zoonosis risk and the potential for novel pathogen emergence.

Sharing of disaggregated data alongside research publications should be promoted with

adoption of data standards to support ongoing data synthesis. Specifically, inclusion of exact

locations of trapping sites, trapping effort and the dates at which trapping occurred would

support more detailed inference of the spatio-temporal dynamics of host populations and the

risk of endemic zoonosis spillover events. Despite these challenges we propose that rodent

trapping studies can provide an important source of data to supplement curated datasets on

rodent distributions to quantify the risk of endemic zoonosis spillover events and the hazard of

novel pathogen emergence.

7. Author contributions

DS – Conceptualisation, Data Curation, Formal Analysis, Writing

LAA - Data Curation, Validation, Writing – Review and Editing

KEJ – Conceptualisation, Supervision, Writing – Review and Editing

DWJ & RK – Supervision, Funding Acquisition, Writing – Review and Editing

8. Data availability

All data and code to reproduce this analysis is available in an archived Zenodo repository [32].

9. References

1.   Bernstein AS, Ando AW, Loch-Temzelides T, Vale MM, Li BV, Li H, et al. The costs and benefits of primary prevention of zoonotic pandemics. Science Advances. 2022;8: eabl4183. doi:10.1126/sciadv.abl4183

2.   Allen T, Murray KA, Zambrana-Torrelio C, Morse SS, Rondinini C, Di Marco M, et al. Global hotspots and correlates of emerging zoonotic diseases. Nature Communications. 2017;8: 1124. doi:10.1038/s41467-017-00923-8

3.   Hassell JM, Begon M, Ward MJ, Fèvre EM. Urbanization and disease emergence: Dynamics at the wildlife–livestock–human interface. Trends in Ecology & Evolution. 2017;32: 55–67. doi:10.1016/j.tree.2016.09.012

4.   García-Peña GE, Rubio AV, Mendoza H, Fernández M, Milholland MT, Aguirre AA, et al.

Land-use change and rodent-borne diseases: Hazards on the shared socioeconomic pathways.

Philosophical Transactions of the Royal Society B: Biological Sciences. 2021;376: 20200362.

doi:10.1098/rstb.2020.0362

5.   McMahon BJ, Morand S, Gray JS. Ecosystem change and zoonoses in the anthropocene.

Zoonoses and Public Health. 2018;65: 755–765. doi:10.1111/zph.12489

6.   Maudlin I, Eisler MC, Welburn SC. Neglected and endemic zoonoses. Philos Trans R Soc

Lond B Biol Sci. 2009;364: 2777–2787. doi:10.1098/rstb.2009.0067

7.   Halliday JEB, Allan KJ, Ekwem D, Cleaveland S, Kazwala RR, Crump JA. Endemic zoonoses

in the tropics: A public health problem hiding in plain sight. Vet Rec. 2015;176: 220–225.

doi:10.1136/vr.h798

8.   Molyneux D, Hallaj Z, Keusch GT, McManus DP, Ngowi H, Cleaveland S, et al. Zoonoses

and marginalised infectious diseases of poverty: Where do we stand? Parasites & Vectors.

2011;4: 106. doi:10.1186/1756-3305-4-106

9.   Judson SD, Rabinowitz PM. Zoonoses and global epidemics. Current Opinion in Infectious

Diseases. 2021;34: 385–392. doi:10.1097/QCO.0000000000000749

10.   Han BA, Schmidt JP, Bowden SE, Drake JM. Rodent reservoirs of future zoonotic diseases.

PNAS. 2015;112: 7039–7044. doi:10.1073/pnas.1501598112

11.   Gibb R, Albery GF, Mollentze N, Eskew EA, Brierley L, Ryan SJ, et al. Mammal virus

diversity estimates are unstable due to accelerating discovery effort. Biology Letters. 2021;18:

20210427. doi:10.1098/rsbl.2021.0427

12.   Fischer C, Gayer C, Kurucz K, Riesch F, Tscharntke T, Batáry P. Ecosystem services and

disservices provided by small rodents in arable fields: Effects of local and landscape

management. Journal of Applied Ecology. 2018;55: 548–558. doi:10.1111/1365-2664.13016

13.   Dobson FS, Oli MK. Fast and slow life histories of rodents. Rodent societies: an ecological

and evolutionary perspective. 2007; 99–105. Available:

https://www.jstor.org/stable/42902037

14.   Albery GF, Becker DJ. Fast-lived hosts and zoonotic risk. Trends in Parasitology.

2021;37: 117–129. doi:10.1016/j.pt.2020.10.012

15.   Gibb R, Redding DW, Chin KQ, Donnelly CA, Blackburn TM, Newbold T, et al. Zoonotic

host diversity increases in human-dominated ecosystems. Nature. 2020;584: 398–402.

doi:10.1038/s41586-020-2562-8

16.   Han BA, Kramer AM, Drake JM. Global patterns of zoonotic disease in mammals. Trends

in Parasitology. 2016;32: 565–577. doi:10.1016/j.pt.2016.04.007

17.   Gibb R, Franklinos LHV, Redding DW, Jones KE. Ecosystem perspectives are needed to

manage zoonotic risks in a changing climate. BMJ. 2020;371: m3389. doi:10.1136/bmj.m3389

18.   Boakes EH, McGowan PJK, Fuller RA, Chang-qing D, Clark NE, O'Connor K, et al. Distorted

views of biodiversity: Spatial and temporal bias in species occurrence data. PLOS Biology.

2010;8: e1000385. doi:10.1371/journal.pbio.1000385

19.   Bowler DE, Callaghan CT, Bhandari N, Henle K, Benjamin Barth M, Koppitz C, et al.

Temporal trends in the spatial bias of species occurrence records. Ecography. 2022;n/a:

e06219. doi:10.1111/ecog.06219

20.   Beck J, Böller M, Erhardt A, Schwanghart W. Spatial bias in the GBIF database and its

effect on modeling species' geographic distributions. Ecological Informatics. 2014;19: 10–15.

doi:10.1016/j.ecoinf.2013.11.002

21.   Plowright RK, Becker DJ, McCallum H, Manlove KR. Sampling to elucidate the dynamics

of infections in reservoir hosts. Philos Trans R Soc Lond B Biol Sci. 2019;374: 20180336.

doi:10.1098/rstb.2018.0336

22.   Carlson CJ, Farrell MJ, Grange Z, Han BA, Mollentze N, Phelan AL, et al. The future of

zoonotic risk prediction. Philosophical Transactions of the Royal Society B: Biological Sciences.

2021;376: 20200358. doi:10.1098/rstb.2020.0358

23.   Wille M, Geoghegan JL, Holmes EC. How accurately can we assess zoonotic risk? Dobson

AP, editor. PLoS Biol. 2021;19: e3001135. doi:10.1371/journal.pbio.3001135

24.   Redding DW, Lucas TCD, Blackburn TM, Jones KE. Evaluating bayesian spatial methods

for modelling species distributions with clumped and restricted occurrence data. PLOS ONE.

2017;12: e0187602. doi:10.1371/journal.pone.0187602

25.   Grace D, Mutua F, Ochungo P, Kruska R, Jones K, Brierley L, et al. Mapping of poverty and

likely zoonoses hotspots. 2012.

26.   Meerburg B, Singleton G, Kijlstra A. Rodent-borne diseases and their risks for public

health. Critical reviews in microbiology. 2009;35: 221–70. doi:10.1080/10408410902989837

27.   Galeh TM, Sarvi S, Montazeri M, Moosazadeh M, Nakhaei M, Shariatzadeh SA, et al. Global

status of toxoplasma gondii seroprevalence in rodents: A systematic review and meta-analysis.

Frontiers in Veterinary Science. 2020;7. Available:

https://www.frontiersin.org/article/10.3389/fvets.2020.00461

28.   Bovendorp RS, MCCleery RA, Galetti M. Optimising sampling methods for small mammal

communities in neotropical rainforests. Mammal Review. 2017;47: 148–158.

doi:10.1111/mam.12088

29.   Fichet-Calvet E, Lecompte E, Veyrunes F, Barrière P, Nicolas V, Koulémou K. Diversity

and dynamics in a community of small mammals in coastal guinea, west africa. Belgian Journal

of Zoology. 2009;139: 93–102. Available: http://www.vliz.be/nl/open-marien-

archief?module=ref&refid=204614

30.   Catalano S, Leger E, Fall CB, Borlase A, Diop SD, Berger D, et al. Multihost transmission of

schistosoma mansoni in senegal, 2015-2018. Emerging Infectious Diseases. 2020;26: 1234–

1242. doi:10.3201/eid2606.200107

31.   USAID. One health surveillance. PREDICT project. 2021 [cited 27 Sep 2021]. Available:

https://p2.predict.global/surveillance

32.   Simons D. DidDrog11/scoping_review: Repository to accompany submission. Zenodo;

2022. doi:10.5281/zenodo.7022637

33.   GBIF: The Global Biodiversity Information Facility. The global biodiversity information

facility. 2021 [cited 25 Apr 2021]. Available: https://www.gbif.org/

34.   National Geospatial-Intelligence Agency. NGA: GNS home. [cited 22 Mar 2021]. Available:

https://geonames.nga.mil/gns/html/

35.   CDC. Emerging and Zoonotic Infectious Disease Laws. 2022 [cited 31 Oct 2022].

Available: https://www.cdc.gov/phlp/publications/topic/zoonotic.html

36.   Pebesma E. Simple features for r: Standardized support for spatial vector data. The R

Journal. 2018;10: 439–446. Available: https://journal.r-project.org/archive/2018/RJ-2018-

009/index.html

37.   R Core Team. R: A language and environment for statistical computing. Vienna, Austria:

R Foundation for Statistical Computing; 2020. Available: https://www.R-project.org/

38.   Database of Global Administrative Areas. GADM. 2022 [cited 25 Apr 2021]. Available:

https://gadm.org/index.html

39.   Socioeconomic Data and Applications Center. Gridded population of the world (GPW), v4

SEDAC. 2021 [cited 4 Feb 2021]. Available:

https://sedac.ciesin.columbia.edu/data/collection/gpw-v4

40.   European Space Agency Climate Change Initiative. Land cover classification gridded

maps from 1992 to present derived from satellite observations. 2022 [cited 3 Mar 2022].

Available: https://cds.climate.copernicus.eu/cdsapp#!/dataset/satellite-land-

cover?tab=overview

41.   Pedersen EJ, Miller DL, Simpson GL, Ross N. Hierarchical generalized additive models in

ecology: An introduction with mgcv. PeerJ. 2019;7: e6876. doi:10.7717/peerj.6876

42. Kendal W. Spatial aggregation of the Colorado potato beetle described by an exponential

dispersion model. Ecological Modelling. 2002;151: 261-269. Doi: 10.1016/S0304-

3800(01)00494-X

43.   Wood SN. Generalized additive models: An introduction with r. 2nd ed. Chapman;

Hall/CRC; 2017.

44.   IUCN. The IUCN red list of threatened species. 2021. Available: www.iucnredlist.org

45.   GBIF: The Global Biodiversity Information Facility. Occurrence download. The Global

Biodiversity Information Facility; 2021. doi:10.15468/DL.S52MEH

46.   Gibb R, Carlson CJ, Farrell MJ. Viralemergence/clover: Preprint + zenodo. Zenodo; 2021.

doi:10.5281/zenodo.4435128

47.   Harvey E, Holmes EC. Diversity and evolution of the animal virome. Nat Rev Microbiol.

2022;20: 321–334. doi:10.1038/s41579-021-00665-x

48.   Basinski AJ, Fichet-Calvet E, Sjodin AR, Varrelman TJ, Remien CH, Layman NC, et al.

Bridging the gap: Using reservoir ecology and human serosurveys to estimate lassa virus

spillover in west africa. Wesolowski A, editor. PLoS Comput Biol. 2021;17: e1008811.

doi:10.1371/journal.pcbi.1008811

49.   Václavík T, Meentemeyer RK. Invasive species distribution modeling (iSDM): Are

absence data and dispersal constraints needed to predict actual distributions? Ecological

Modelling. 2009;220: 3248–3258. doi:10.1016/j.ecolmodel.2009.08.013

50.   Zeimes CB, Olsson GE, Ahlm C, Vanwambeke SO. Modelling zoonotic diseases in humans:

Comparison of methods for hantavirus in sweden. Int J Health Geogr. 2012;11: 39.

doi:10.1186/1476-072X-11-39

51.   Judson SD, LeBreton M, Fuller T, Hoffman RM, Njabo K, Brewer TF, et al. Translating

predictions of zoonotic viruses for policymakers. EcoHealth. 2018;15: 52–62.

doi:10.1007/s10393-017-1304-3

52.   Gibb R, Albery GF, Becker DJ, Brierley L, Connor R, Dallas TA, et al. Data proliferation,

reconciliation, and synthesis in viral ecology. BioScience. 2021;71: 1148–1156.

doi:10.1093/biosci/biab080

53.   Wilkinson DP, Golding N, Guillera-Arroita G, Tingley R, McCarthy MA. A comparison of

joint species distribution models for presence–absence data. Methods in Ecology and Evolution.

2019;10: 198–211. doi:10.1111/2041-210X.13106

54.   Fichet-Calvet E, Ölschläger S, Strecker T, Koivogui L, Becker-Ziaja B, Camara AB, et al.

Spatial and temporal evolution of lassa virus in the natural host population in upper guinea. Sci

Rep. 2016;6: 21977. doi:10.1038/srep21977

55.   Herrmann SM, Brandt M, Rasmussen K, Fensholt R. Accelerating land cover change in

west africa over four decades as population pressure increased. Commun Earth Environ.

2020;1: 1–10. doi:10.1038/s43247-020-00053-y

56.   Foster ED, Deardorff A. Open science framework (OSF). J Med Libr Assoc. 2017;105:

203–206. doi:10.5195/jmla.2017.88

A

Mastomys natalensis

Rattus rattus

Mus musculus

Detection/Non−detection ● Detection ● Non−detection

Acute infection

Serology