**Rodent trapping studies as an overlooked information source for**

**understanding endemic and novel zoonotic spillover.**

**Rodent trapping to understand zoonotic spillover risk**

David Simons[1, 2, 3]

Lauren A. Attfield[3]

Kate E. Jones[3]

Deborah Watson-Jones[2, 4]

Richard Kock[1]

[1] Centre for Emerging, Endemic and Exotic Diseases, The Royal Veterinary College, London, UK

[2] Clinical Research Department, London School of Hygiene and Tropical Medicine, London, UK

[3] Centre for Biodiversity and Environment Research, University College London, London, UK

[4] Mwanza Intervention Trials Unit, National Institute for Medical Research, Mwanza, Tanzania

## 1. Abstract

Rodents are important globally distributed reservoirs of known and novel zoonotic pathogens. Ongoing anthropogenic land use change is altering the composition of host species assemblages and modifying the risk of zoonoses spillover events. These changes mean that an understanding of the current distribution of rodent species is vital for accurately describing disease hazard and managing risk. However, available species distribution and host-pathogen association datasets (e.g. IUCN, GBIF, CLOVER) are often taxonomically and spatially biased. Here, we synthesise data from West Africa from 127 rodent trapping studies, published between 1964-2022, as an additional source of information to characterise the range and presence of important zoonotic pathogen host species in this region. We identify that these rodent trapping studies, although biased towards human dominated landscapes across West Africa, can usefully complement current rodent species distribution datasets and we calculate the discrepancies between these datasets. For five regionally important zoonotic pathogens (Arenaviridae spp., Borrelia spp., *Lassa mammarenavirus*, Leptospira spp. and *Toxoplasma gondii*), we identify host-pathogen associations that have not been previously reported in host-association datasets. These omissions have the potential for biasing estimates of current risk and drivers of zoonoses. Finally, for these five pathogen groups, we find that the proportion of a rodent hosts range that has been sampled remains small with geographic clustering. A priority of future rodent trapping studies should be to sample rodent hosts across a greater geographic range to better characterise current and future risk of zoonotic spillover events. In the interim, future studies of spatial pathogen risk informed by rodent distributions must incorporate a measure of the current sampling biases. The current synthesis of contextually rich rodent trapping data

36   enriches available information from IUCN, GBIF and CLOVER which can support a more

37   complete understanding of the hazard of zoonotic spillover events.

## 2. Author Summary

Emerging and endemic zoonotic diseases are projected to have increasing health impacts,

particularly under changing climate and land-use scenarios. Rodents carry a disproportionate

number of zoonotic pathogens and are abundant across West Africa, raising concerns about

increasing outbreaks of endemic disease and the emergence of pandemics. Prior modelling

studies rely on large, consolidated data sources which do not incorporate high resolution

spatial and temporal data from rodent trapping studies. Here, we synthesise these studies to

quantify the bias in the sampling of rodent hosts and their pathogens across West Africa. We

find that rodent trapping studies are complementary to these datasets and can provide

additional, high-resolution data on the distribution of hosts and their pathogens. Further,

rodent trapping studies have identified additional potential host-pathogen associations than

are recorded in consolidated host-pathogen association datasets. This can help to understand

the risk of zoonotic diseases based on host distributions. Finally, we quantify the current extent

of known rodent presence and pathogen sampling within a species range, highlighting that

current knowledge is limited across much of the region. We hope that this will support work to

study rodent hosts and their pathogens in currently under sampled regions to better

understand the risk of emerging and endemic zoonoses in West Africa.

## 3. Introduction

There is increasing awareness of the global health and economic impacts of novel zoonotic pathogen spillover, driven by the ongoing SARS-CoV-2 pandemic and previous HIV/AIDs and Spanish Influenza pandemics [1]. The number of zoonotic disease spillover events and the frequency of the emergence of novel zoonotic pathogens from rodents are predicted to increase under intensifying anthropogenic pressure driven by increased human populations, urbanisation, intensification of agriculture, climate change and wildlife defaunation [2–5]. The impact of endemic zoonoses meanwhile remains underestimated [6]. Endemic zoonoses disproportionally affect those in the poorest sections of society, those living in close contact with their animals and those with limited access to healthcare [7–9].

Rodents along with bats contribute the greatest number of predicted novel zoonotic pathogens and known endemic zoonoses [10,11]. Of 2,220 extant rodent species, 244 (10.7%) are described as reservoirs of 85 zoonotic pathogens [10], although many species provide important and beneficial ecosystem services including pest regulation and seed dispersal [12]. Rodents typically demonstrate "fast" life histories [13] with traits such as early maturation and short gestation times (<4 days) further associated with being zoonotic reservoirs [10,14]. Rodent species with "fast" life histories thrive in human dominated landscapes, displacing species less likely to be reservoirs of zoonotic pathogens [15]. The widespread occurrence of reservoir species and their proximity to human activity make the description of rodent species assemblages and host-pathogen associations vitally important to understanding the hazard of zoonotic disease spillover and novel zoonotic pathogen emergence [16].

Despite the importance of understanding these complex systems, current evidence on host-

pathogen associations is considerably affected by taxonomic and geographical sampling biases

[11,17]. Curated biodiversity datasets such as the Global Biodiversity Information Facility

(GBIF) and resources produced by the International Union for Conservation of Nature (IUCN)

suffer from well described spatial and temporal sampling biases [18,19]. These sampling biases

can importantly distort produced species distribution models [20]. Datasets on host-pathogen

associations also can suffer from biases introduced from literature selection criteria and

taxonomic discrepancies. These biases are important because identification of potential

geographic hotspots of zoonotic disease spillover and novel pathogen emergence are often

produced from these types of host species distributions and host-pathogen associations

[21,22]. For example, systematically increased sampling, over-representation of certain

habitats and clustering around areas of high human population could lead to an apparent

association between locations and hazard that is driven by these factors rather than underlying

host-pathogen associations [11,23,24]. Predictions of zoonotic disease spillover and novel

zoonotic pathogen emergence must account for these biases to understand the future hazard of

zoonotic diseases [22].

West Africa has been identified as a region at increased risk for rodent-borne zoonotic disease

spillover events, the probability of these events are predicted to increase under different

projected future land-use change scenarios [4,25]. Currently within West Africa, rodents are

involved in the transmission of multiple endemic zoonoses with large burdens on human

health, these pathogens include Lassa fever, Schistosomiasis, Leptospirosis and Toxoplasmosis

[26,27]. Understanding of the distribution of these zoonoses are limited by biases in

consolidated datasets. Rodent trapping studies provide contextually rich information on when,

where and under what conditions rodents were trapped, potentially enriching consolidated

datasets [28]. Studies have been conducted in West Africa to investigate the distribution of

rodent species, their species assemblages, the prevalence of endemic zoonoses within rodent

hosts (e.g., Lassa fever, Schistosomiasis) and to identify emerging and novel zoonotic pathogens

[29–31]. However, individual level data from these studies have not previously been

synthesised for inclusion in assessments of zoonotic disease spillover and novel zoonotic

pathogen emergence.

Here, we synthesise rodent trapping studies conducted across West Africa published between

1964-2022. First, we use this dataset to investigate the geographic sampling biases in relation

to human population density and land use classification. Second, we compare this to curated

host datasets (IUCN and GBIF) to understand differences in reported host geographic

distributions. Third, we compare identified host-pathogen associations with a consolidated

dataset (CLOVER) to explore discrepancies in rodent host-pathogen associations and report the

proportion of positive assays for pathogens of interest. Finally, within our dataset we

investigate the spatial extent of current host-pathogen sampling to identify areas of sparse

sampling of pathogens within their host ranges. We expect that rodent trapping studies provide

an important additional source of high-resolution data that can be used to enrich available

consolidated datasets to better understand the hazard of zoonotic disease spillover and novel

zoonotic pathogen emergence across West Africa.

## 4. Methods

### 4.1.    Data sources

#### 4.1.1. Host and pathogen trapping data

To identify relevant literature, we conducted a search in Ovid MEDLINE, Web of Science (Core collection and Zoological Record), JSTOR, BioOne, African Journals Online, Global Health and the pre-print servers, BioRxiv and EcoEvoRxiv for the following terms as exploded keywords: (1) Rodent OR Rodent trap* AND (2) West Africa, no date limits were set. We also searched other resources including the UN Official Documents System, Open Grey, AGRIS FAO and Google Scholar using combinations of the above terms. Searches were run on 2022-05-01, and returned studies conducted between 1964-2021.

We included studies for further analysis if they met all of the following inclusion criteria; i) reported findings from trapping studies where the target was a small mammal, ii) described the type of trap used or the length of trapping activity or the location of the trapping activity, iii) included trapping activity from at least one West African country, iv) recorded the genus or species of trapped individuals, and v) were published in a peer-reviewed journal or as a pre-print on a digital platform or as a report by a credible organisation. We excluded studies if they met any of the following exclusion criteria: i) reported data that were duplicated from a previously included study, ii) no full text available, iii) not available in English. One author screened titles, abstracts and full texts against the inclusion and exclusion criteria. At each stage; title screening, abstract screening and full text review, a random subset (10%) was reviewed by a second author.

We extracted data from eligible studies using a standardised tool that was piloted on 5 studies

(Supplementary Table 1.). Data was abstracted into a Google Sheets document, which was

archived on completion of data extraction [32]. We identified the aims of included studies, for

example, whether it was conducted as a survey of small mammal species or specifically to

assess the risk of zoonotic disease spillover. we extracted data on study methodology, such as,

the number of trap nights, the type of traps used and whether the study attempted to estimate

abundance. For studies not reporting number of trap nights we used imputation based on the

number of trapped individuals, stratified by the habitat type from which they were obtained.

This was performed by multiplying the total number of trapped individuals within that study

site by the median trap success for study sites with the same reported habitat type.

Stratification was used as trap success varied importantly between traps placed in or around

buildings (13%, IQR 6-24%) compared with other habitats (3%, IQR 1-9%)

We also recorded how species were identified within a study and species identification was

assumed to be accurate. The number of individuals of these species or genera was extracted

with taxanomic names mapped to GBIF taxonomy [33]. We expanded species detection and

non-detection records by explicitly specifying non-detection at a trap site if a species was

recorded as detected at other trapping locations within the same study.

Geographic locations of trapping studies were extracted using GPS locations for the most

precise location presented. Missing locations were found using the National Geospatial-

Intelligence Agency GEOnet Names Server [34] based on placenames and maps presented in the

study. All locations were converted to decimal degrees. The year of rodent trapping was

extracted alongside the length of the trapping activity to understand seasonal

representativeness of trapping activity. The habitats of trapping sites were mapped to the IUCN

Habitat Classification Scheme (Version 3.1). For studies reporting multiple habitat types for a

single trap, trap-line or trapping grid, a higher order classification of habitat type was recorded.

For relevant studies we extracted data on all microorganisms and zoonotic pathogens tested

and the method used (e.g., molecular or serological diagnosis). Where assays were able to

identify the microorganism to species level this was recorded, non-specific assays higher order

attribution was used (e.g. to family level). We recorded the species of rodent host tested, the

number of individuals tested and the number of positive and negative results. For studies

reporting summary results all testing data were extracted, this may introduce double counting

of individual rodents, for example, if a single rodent was tested using both molecular and

serological assays. Where studies reported indeterminate results, these were also recorded.

Out of 4,692 relevant citations, we identified 127 rodent trapping studies (Supplementary

Table 2.). The earliest trapping studies were conducted in 1964, with a trend of increasing

numbers of studies being performed annually since 2000. The median year of first trapping

activity was 2007, with the median length of trapping activity being 1 year (IQR 0-2 years)

(Supplementary Fig 1.). Studies were conducted in 14 West African countries, with no studies

reported from The Gambia or Togo, at 1,611 trap sites (Fig 1A.).

Included studies explicitly reported on 601,184 trap nights, a further 341,445 trap nights were

imputed from studies with no recording of trapping effort based on trap success, leading to an

estimate of 942,629 trap nights (Fig 1B.). A minority of studies trapped at a single study site

(30, 24%), with 46 (36%) trapping at between two and five sites, the remaining 51 studies

(40%) trapped at between six and 93 study sites.

In total 76,275 small mammals were trapped with 65,628 (90%) identified to species level and

7,439 (10%) identified to genus, with the remaining classified to higher taxonomic level. The

majority of the 132 identified species were Rodentia (102, 78%), of which Muridae (73, 72%)

were the most common family. Soricomorpha were the second most identified order of small

mammals (28, 21%). 57 studies tested for 32 microorganisms, defined to species or genus level

that are known or potential pathogens. Most studies tested for a single microorganism (48,

84%). The most frequently assayed microorganisms were *Lassa mammarenavirus* or

Arenaviridae (21, 37%), *Borrelia sp.* (9, 16%), *Bartonella sp.* (4, 7%) and *Toxoplasma gondii* (4,

7%). Most studies used Polymerase Chain Reaction (PCR) to detect microorganisms (37, 65%),

with fewer studies using serology-based tests (11, 19%) or histological or direct visualisation

assays (11, 21%). From 32,920 individual rodent samples we produced 351 host-pathogen

pairs. With *Rattus rattus*, *Mus musculus*, *Mastomys erythroleucus*, *Mastomys natalensis* and

*Arvicanthis niloticus* being assayed for at least 18 microorganisms.

**Fig 1: Rodent trapping sites across West Africa.** A) The location of trapping sites in West

Africa. No sites were recorded from Togo or The Gambia. Heterogeneity is observed in the

coverage of each country by trap night (colour) and location of sites. For example, Senegal, Mali

and Sierra Leone have generally good coverage compared to Guinea and Burkina Faso. B)

Histogram of trap nights performed at each study site, a median of 248 trap nights (IQR 116-

500) was performed at each site

## 4.2.     Analysis

### 4.2.1. What is the extent of spatial bias in the rodent trapping data?

To investigate the extent of spatial bias in the rodent trapping data, we calculated trap-night (TN) density within each West African level-2 administrative region. The `sf` package in the R statistical language (R version 4.1.2) was used to manipulate geographic data, administrative boundaries were obtained from GADM 4.0.4 [35–37]. Trap-night density ($TN_{density}$) was calculated by dividing the number of trap nights by the area of a level-2 administrative area ($R_{area}$). For studies not reporting trap nights, imputation was used as previously described. Human population density was obtained for the closest year (2005) to the median year of trapping (2007) from Socioeconomic Data and Applications Center (SEDAC) gridded population of the world v4 at ~ 1km resolution ($P_{density}$) [38]. Median population density was then calculated for each level-2 administrative region. Land cover classification was obtained from the Copernicus climate change service at ~300m resolution [39]. The proportion of cropland, shrubland, tree cover ($\psi_{tree}$) and urban land cover ($\psi_{urban}$) within a level-2 administrative region in 2005 was calculated.

We investigated the association between relative trapping effort, measured as TN density, and the proportion of urban, cropland, tree cover and human population density using Generalised Additive Models (GAM) incorporating a spatial interaction term (longitude and latitude) [40]. The models were constructed in the `mgcv` package [41]. Selection of the most parsimonious model was based on Deviance explained and the Akaike Information Criterion for each model (Equations 1-5 below). Relative trapping effort was then predicted across West Africa using these covariates. We performed two sensitivity analyses, first, by removing sites with imputed

trapping effort, second, by associating trap locations to ~1km pixels rather than level-2

administrative areas.

$$TN_{density} \sim Tweedie(X * Y) \#(1)$$

$$TN_{density} \sim Tweedie\big(P_{density} + (X * Y)\big) \#(2)$$

$$TN_{density} \sim Tweedie\big(P_{density} + R_{area} + (X * Y)\big) \#(3)$$

$$TN_{density} \sim Tweedie\big(P_{density} + \psi_{tree} + \psi_{urban} + (X * Y)\big) \#(4)$$

$$TN_{density} \sim Tweedie\big(P_{density} + R_{area} + \psi_{urban} + (X * Y)\big) \#(5)$$

**4.2.2. What is the difference in rodent host distributions between curated**

**datasets and rodent trapping studies?**

We assessed the concordance of curated rodent host distributions from IUCN and GBIF with

observed rodent detection and non-detection from rodent trapping studies for seven species

with the most trap locations (*M. natalensis, R. rattus, M. erythroleucus, M. musculus, A. niloticus,*

*Praomys daltoni* and *Cricetomys gambianus*). We obtained rodent species distribution maps as

shapefiles from the IUCN red list and translated these to a ~20km resolution raster [42].

Distributions were cropped to the study region for globally distributed rodent species. We

obtained rodent presence locations from GBIF as point data limited to the study region [43].

Presence locations were associated to cells of raster with a ~20km resolution produced for the

study region.

For each of the seven species, we first calculated the area of the IUCN expected range, and then

the percentage of this range covered by presence detections in GBIF, and from detections in the

rodent trapping data. We then calculated the area of both types of both GBIF and rodent

trapping detections outside of the IUCN expected range. For rodent trapping data, we

additionally calculated the area of non-detections within the IUCN expected area. Finally, we

calculated the combined area of detection from both GBIF and rodent trapping data.

**4.2.3. Are rodent trapping derived host-pathogen associations present in a**

**consolidated zoonoses dataset?**

To examine the usefulness of rodent trapping studies as an additional source of data we

compared identified host-pathogen associations from trapping studies investigating zoonoses

with a consolidated zoonoses dataset (CLOVER) [11,44]. CLOVER is a synthesis of four host-

pathogen datasets (GMPD2, EID2, HP3 and Shaw, 2020) and was released in 2021, it contains

more than 25,000 host-pathogen associations for Bacteria, Viruses, Helminth, Protozoa and

Fungi. We compared the host-pathogen networks across the two datasets, where the CLOVER

data was subset for host species present in the rodent trapping data.

For host-pathogen pairs with assay results consistent with acute or prior infection, we

calculated the proportion positive and identify those absent from CLOVER. We expand the

analysis to host-pathogen pairs with pathogens identified to genus level in Supplementary Fig

4.

### 4.2.4. What is the spatial extent of pathogen testing within host ranges?

We use the sampled area of three pathogen groups and two pathogens (Arenaviridae, Borreliaceae, Leptospiraceae, *Lassa mammarenavirus* and *Toxoplasma gondii*) to quantify the bias of sampling within their hosts ranges. For each pathogen, we first describe the number of host species assayed, for the five most commonly tested species we associate the locations of sampled individuals to ~20km pixels and calculate the proportion of the IUCN range of the host in which sampling has occurred. We compare this figure to the total area in which the host has been detected to produce a measure of relative completeness of sampling within the included rodent trapping studies.

Data and code to reproduce all analyses are available in an archived Zenodo repository [32].

## 5. Results

## 5.1.    What is the extent of spatial bias in the rodent trapping data?

We found non-random, spatial clustering of rodent trapping locations across the study region, suggestive of underlying bias in the sampling or rodents across West Africa. Trap sites were situated in 256 of 1,450 (17.6%) level-2 administrative regions in 14 West African nations. The regions with the highest TN density included the capitals and large cities of Niger (Niamey), Nigeria (Ibadan), Ghana (Accra), Senegal (Dakar), Ghana (Accra) and Benin (Cotonou). Outside of these cities, regions in, Northern Senegal, Southern Guinea, Edo and Ogun States in Nigeria and Eastern Sierra Leone had the greatest TN density (Fig 1A.).

The most parsimonious GAM model (adjusted R2 = 0.3, Deviance explained = 48.7%) reported

significant non-linear associations between relative trapping effort bias and human population

densities (Effective Degrees of Freedom (EDF) = 7.13, $p < 0.001$), proportion of urban landscape

(EDF = 1.92, $p < 0.002$) and region area (EDF = 3.63, $p < 0.001$), alongside significant spatial

associations (EDF = 27.3, $p < 0.001$) (Supplementary table 3.1). Greatest trapping effort bias

peaked at population densities between 5,000-7,500 individuals/km$^2$, proportion of urban

landscape >10% and region areas < 1,000km$^2$. Increased trapping effort was found in North

West Senegal, North and East Sierra Leone, Central Guinea and coastal regions of Nigeria, Benin

and Ghana; in contrast South East Nigeria, Northern Nigeria and Burkina Faso had an observed

bias towards a reduced trapping effort (Fig 2). In sensitivity analysis, excluding sites with

imputed trap nights, Mauritania, Northern Senegal and Sierra Leone remained as regions

trapped at higher rates, with Nigeria being trapped at lower than expected rates

(Supplementary Fig 3a.). In pixel-based sensitivity analysis spatial coverage was reduced with

similar patterns of bias observed to the primary analysis (Supplementary Fig 3b.).

**Fig 2. Relative trapping effort bias across West Africa.** Modelled relative trapping effort bias adjusted for human population density, proportion urban land cover and area of the administrative region. Brown regions represent areas with a bias towards increased trapping effort (e.g., North West Senegal), Green regions represent areas with a bias towards reduced trapping effort (e.g., Northern Nigeria).

## 5.2. What is the difference in rodent host distributions between curated datasets and rodent trapping studies?

We found that for six of the seven most detected rodent species (*M. natalensis*, *R. rattus*, *M. erythroleucus*, *M. musculus*, *A. niloticus* and *P. daltoni*), trapping studies provided more distinct locations of detection and non-detection than were available from GBIF. For the endemic rodent species (*M. natalensis*, *M. erythroleucus*, *A. niloticus*, *P. daltoni* and *C. gambianus*) IUCN ranges had good concordance to both trapping studies and GBIF, however, individuals of *A. niloticus* and *P. daltoni* were detected outside of IUCN ranges. In contrast, the non-native species *R. rattus* and *M. musculus* were detected across much greater ranges than were expected from IUCN distributions. Comparisons for *M. natalensis*, *R. rattus* and *M. musculus* are shown in Fig 3, the remaining species are shown in Supplementary Fig 4.

**Mastomys natalensis**

**Rattus rattus**

**Mus musculus**

Detection/Non-detection   ● Detection   ● Non-detection

**Fig 3. Locations of detection and non-detection sites for rodent species in West Africa**

Each row corresponds to a single rodent species. L) Presence recorded in GBIF (black points)

overlaid on IUCN species range (red-shaded area). R) Detection (purple) and non-detection

(orange) from rodent trapping studies overlaid on IUCN species ranges. M. musculus has no

IUCN West African range.

Comparison of the proportion of a species IUCN range in which detections and non-detections

occurred showed that sampling locations of these seven species within GBIF covered between

0.09-0.26% of expected ranges (Table 1.), compared to 0.03-0.24% for rodent trapping data.

Detections occurred outside IUCN ranges for all species in both the GBIF and rodent trapping

data, most noticeably for *A. niloticus* and *R. rattus*. Combining GBIF and rodent trapping data

increased the sampled area by a mean of 1.6 times compared to the GBIF area alone,

demonstrating limited overlap between the locations providing information to either dataset.

Non-detection of a species occurred across species ranges (mean = 0.11%, SD = 0.03%),

suggestive of spatial heterogeneity of presence within IUCN ranges.

**Table 1: Comparison of IUCN, GBIF and rodent trapping ranges for the 7 most detected**

329                                **rodent species.**

| Species | IUCN Range (1,000 km²) | GBIF Area inside range (1,000 km²) (% of IUCN) | GBIF Area outside range (1,000 km²) | Trapping studies Detection area inside range (1,000 km²) (% of IUCN) | Trapping studies Area outside range (1,000 km²) (% of IUCN) | Trapping studies Non-detection area inside range (1,000 km²) (% of IUCN) | Combined Detection area (1,000 km²) (% of IUCN) |
|---|---|---|---|---|---|---|---|
| *Mastomys natalensis* | 3,257 | 6.83 (0.21%) | 0.19 | 4.4 (0.14%) | 0.17 | 3.12 (0.1%) | 10.66 (0.33%) |
| *Rattus rattus* | 1,019 | 2.61 (0.26%) | 0.52 | 2.42 (0.24%) | 1.21 | 1.3 (0.13%) | 4.88 (0.48%) |
| *Mastomys erythroleucus* | 3,735 | 4.48 (0.12%) | 0.04 | 3.24 (0.09%) | 0.12 | 4.35 (0.12%) | 7.5 (0.2%) |
| *Mus musculus* | | | 2.15 | | 1.85 | | 3.94 |
| *Arvicanthis niloticus* | 1,829 | 1.69 (0.09%) | 2.41 | 1.98 (0.11%) | 0.34 | 3.09 (0.17%) | 3.59 (0.2%) |
| *Praomys daltoni* | 2,658 | 4.03 (0.15%) | 0.29 | 2.03 (0.08%) | 0.15 | 2.78 (0.1%) | 5.83 (0.22%) |
| *Cricetomys gambianus* | 2,476 | 5 (0.2%) | 0.17 | 0.75 (0.03%) | 0.06 | 2.99 (0.12%) | 5.69 (0.23%) |

## 5.3.     Are rodent trapping derived host-pathogen associations present in a consolidated zoonoses dataset?

We found potentially important differences between the host-pathogen networks produced from included rodent trapping studies and the consolidated CLOVER dataset. When limited to taxonomic classification of both pathogen and host to species level we identified 25 host-pathogen pairs among 14 rodent and 6 pathogen species (Fig 4. and Fig 5.). We identified

negative associations (non-detection through specific assays) for 45 host-pathogen pairs

among 35 rodent and 7 pathogen species. CLOVER contained 10 (40%) of our identified host-

pathogen associations, the remaining 15 (60%) were not found to be present in CLOVER,

additionally CLOVER recorded positive associations for 4 (9%) of the negative associations

produced from the rodent trapping data.

**Fig 4. Host-Pathogen associations detected through acute infection.** A) Identified species

level host-pathogen associations through detection of acute infection (i.e. PCR, culture).

Percentages and colour relate to the proportion of all assays that were positive, the number of

individuals tested for the pathogen is labelled N. Associations with a black border are present in

the CLOVER dataset.

**Fig 5. Host-Pathogen associations detected through evidence of prior infection.** B) Identified species level host-pathogen associations through serological assays (i.e. ELISA). Percentages and colour relate to the proportion of all assays that were positive, the number of individuals tested for the pathogen is labelled N. Associations with a black border are present in the CLOVER dataset.

CLOVER included an additional 492 host-pathogen associations we do not observe in rodent

trapping studies. The majority of these 392 (80%) pairs are from species with global

distributions (*M. musculus*, *R. rattus* and *R. norvegicus*), or from those with wide ranging

distributions in sub-Saharan Africa (38, 8%) (i.e., *A. niloticus*, *M. natalensis* and *Atelerix*

*albiventris*).

For pathogens not identified to species level (i.e. family or higher taxa only), we identified 148

host-pathogen pairs among 32 rodent species and 25 pathogen families (Supplementary Fig 4.),

with CLOVER containing 66 (45%) of these associations.

Rodent trapping studies identified additional rodent host species for six pathogens; *Lassa*

*mammarenavirus* (5), *Toxoplasma gondii* (4), Usutu virus (2), *Coxiella burnetii* (2), *Escherichia*

*coli* and *Klebsiella pneumoniae* (both 1), that were not present in this consolidated host-

pathogen association dataset.

## 5.4.    What is the spatial extent of pathogen testing within a host's

## range?

The five most widely sampled pathogen species/families in included studies were Arenaviridae,

Borreliaceae, *Lassa mammarenavirus*, Leptospiraceae and *Toxoplasma gondii* (Table 2.). Assays

to identify Arenaviridae infection were performed in 44 rodent species with evidence of viral

infection in 15 species. *Lassa mammarenavirus* was specifically tested for in 43 species with 10

showing evidence of viral infection. The most commonly infected species for both Arenaviridae,

generally, and *Lassa mammarenavirus* specifically, were *M. natalensis* and *M. erythroleucus*.

These species were assayed across between 10-20% of their trapped area, equating to ∼0.02%

of their IUCN range (Table 2.).

Infection with species of Borreliaceae was assessed in 42 species, with evidence of infection in

17 rodent species. The greatest rates of infection were among *A. niloticus* (16%), *Mastomys*

*huberti* (11%) and *M. erythroleucus* (9%). Testing was more widespread than for Arenviruses

with coverage between 15-34% of their trapped area, however, this remains a small area in

relation to their IUCN ranges (<0.05%). Leptospiraceae and *Toxoplasma gondii* was assessed in

8 species, with evidence of infection in 5 and 6 rodent species respectively. The spatial coverage

of testing for these pathogens was more limited within IUCN host species ranges (∼0.01%).

**Table 2: Comparison of pathogen sampling ranges for the 5 most widely sampled**

**pathogens and the 5 most sampled rodent host species (\* no IUCN range in West African).**

| Pathogen | Species | Tested | Positive | Pathogen testing area (1,000 km$^2$) | Pathogen testing area within trapped area (%) | Pathogen testing area within IUCN range (%) |
|---|---|---|---|---|---|---|
| Arenaviridae sp. | | | | | | |
| | *Mastomys natalensis* | 2,841 | 104 (4%) | 0.61 | 13.45% | 0.02% |
| | *Praomys daltoni* | 854 | 6 (1%) | 0.42 | 19.43% | 0.02% |
| | *Mastomys erythroleucus* | 398 | 20 (5%) | 0.40 | 11.97% | 0.01% |
| | *Rattus rattus* | 396 | 4 (1%) | 0.38 | 10.5% | 0.04% |
| | *Praomys rostratus* | 310 | 5 (2%) | 0.13 | 12.53% | 0.02% |
| Borrelia sp. | | | | | | |
| | *Mastomys erythroleucus* | 1,586 | 140 (9%) | 1.14 | 33.94% | 0.03% |

| Pathogen | Species | Tested | Positive | Pathogen testing area (1,000 km$^2$) | Pathogen testing area within trapped area (%) | Pathogen testing area within IUCN range (%) |
|---|---|---|---|---|---|---|
| Lassa mammarenavirus | *Arvicanthis niloticus* | 1,551 | 253 (16%) | 0.66 | 28.48% | 0.03% |
| | *Mastomys natalensis* | 733 | 54 (7%) | 0.69 | 15.08% | 0.02% |
| | *Mastomys huberti* | 731 | 83 (11%) | 0.23 | 29.83% | 0.04% |
| | *Mus musculus* | 686 | 26 (4%) | 0.45 | 24.54% | * |
| | *Mastomys natalensis* | 3,199 | 580 (18%) | 1.03 | 22.65% | 0.03% |
| | *Mastomys erythroleucus* | 352 | 14 (4%) | 0.36 | 10.63% | 0.01% |
| | *Rattus rattus* | 177 | 2 (1%) | 0.34 | 9.26% | 0.03% |
| | *Praomys rostratus* | 163 | 2 (1%) | 0.27 | 27.02% | 0.04% |
| | *Mus musculus* | 147 | 0 (0%) | 0.04 | 2.29% | * |
| Leptospira sp. | *Rattus rattus* | 646 | 65 (10%) | 0.40 | 11.1% | 0.04% |
| | *Arvicanthis niloticus* | 221 | 10 (5%) | 0.02 | 0.9% | <0.01% |
| | *Crocidura olivieri* | 141 | 14 (10%) | 0.34 | 25.16% | * |
| | *Mastomys natalensis* | 136 | 26 (19%) | 0.36 | 7.91% | 0.01% |
| | *Rattus norvegicus* | 79 | 19 (24%) | 0.21 | 40.08% | * |
| Toxoplasma gondii | | | | | | |

| Pathogen | Species | Tested | Positive | Pathogen testing area (1,000 km$^2$) | Pathogen testing area within trapped area (%) | Pathogen testing area within IUCN range (%) |
|---|---|---|---|---|---|---|
| | *Mus musculus* | 1,548 | 115 (7%) | 0.62 | 33.64% | * |
| | *Rattus rattus* | 428 | 8 (2%) | 0.36 | 9.77% | 0.03% |
| | *Mastomys erythroleucus* | 292 | 13 (4%) | 0.37 | 11.06% | 0.01% |
| | *Mastomys natalensis* | 107 | 2 (2%) | 0.08 | 1.83% | <0.01% |
| | *Cricetomys gambianus* | 47 | 13 (28%) | 0.06 | 7.6% | <0.01% |

## 6. Discussion

Endemic rodent zoonoses and novel pathogen emergence from rodent hosts are predicted to have an increasing burden in West Africa and globally [10]. Here we have synthesised data from 126 rodent trapping studies containing information on more than 72,000 rodents across 1,611 trap sites producing an estimated 942,669 trap nights across 14 West African countries. Locations studied are complementary to curated datasets (e.g. IUCN, GBIF), incorporation of our synthesised dataset when assessing zoonosis risk based on host distributions could counteract some of the biases inherent to these curated datasets [18]. We identified 25 host-pathogen pairs reported from included studies, 15 of these were not included in a consolidated host-pathogen dataset. Generally, the number of different species tested for a pathogen and the spatial extent of these sampling locations were limited. These findings highlight a number of sampling bias, supporting calls for further pathogen sampling across diverse species in zoonotic hotspots [45].

We found that rodent trapping data, like biodiversity data, showed important spatial biases

[20]. Relative trapping effort bias was greater in Benin, Guinea, Senegal and Sierra Leone driven

by long-standing research collaborations investigating the invasion non-native rodent species

(*M. musculus* and *R. rattus*) and the hazard of endemic zoonosis outbreaks (e.g., *Lassa*

*mammarenavirus*). Much of the region remains relatively under sampled, particularly Burkina

Faso, Côte d'Ivoire, Ghana and Nigeria, despite these countries facing many of the same

challenges. For example, annual outbreaks of Lassa fever are reported in Nigeria and there are

potentially 60,000 unrecognised cases of Lassa fever every year in Côte d'Ivoire and Ghana

[46]. Rodent sampling should be targeted towards currently under sampled regions to reduce

the potential impact of this bias and improve our understanding of both the distribution of

rodent hosts and the prevalence of pathogens within their populations. This will allow for

better estimation of risk from endemic and novel zoonoses.

Rodent trapping studies provide geographic and temporally contextualised data on both

species detection and non-detection which are not available from curated datasets. Non-

detection data can improve models of species distributions, unfortunately, high levels of

missing data on trapping effort will continue to confound the allocations of non-detections as

true absences [47]. Models of host species occurrence and abundance, improved by

incorporating species absence, are important to assess the effect of land use and climate change

on endemic zoonosis spillover to human populations and direct limited public health resources

towards regions at greatest risk [48,49].

Currently available consolidated datasets on host-pathogen associations (e.g. CLOVER, EID2

and GMPD2) do not include spatial or temporal components [50]. The current synthesis of

rodent trapping studies has highlighted that pathogens have been sparsely sampled within a

host's range. Current zoonosis risk models dependent on these sources of data are therefore

not able to incorporate spatial heterogeneity in pathogen prevalence across the host range.

Additional uncertainty in current models of zoonotic disease risk arises from host-pathogen

associations that have not been reported in these consolidated datasets. For example,

*Hylomyscus pamfi* infected with *Lassa mammarenavirus* and *R. rattus* infected with *Coxiella*

*burnetii*, will not be included when solely based on consolidated host-pathogen datasets.

Further, detection of zoonotic pathogens in multiple co-occurring host species supports the

adoption of multi-species to better understand the potential range of endemic zoonoses [51].

Few studies stratified detection and non-detection of hosts or pathogen prevalence by time,

therefore limiting inference of changes in host and pathogen dynamics. This limitation prevents

calculation of incidence of infection and the abundance of infectious rodents which potentially

varies by both time and space [52]. Understanding of temporal changes in viral burden and

shedding for endemic zoonoses is required to accurately predict current and future risk of

pathogen spillover.

Finally, due to data sparsity, we were unable to account for temporal change over the six

decades of rodent trapping studies. Land use change and population density have changed

dramatically over this period in West Africa [53]. We attempted to mitigate against this by

using the median year of trapping to understand the spatial and land use biases in trapping

activity. It is possible that land use and population density at trapping sites varied importantly

between when rodent trapping was conducted and the conditions in 2005. Despite this

limitation, the finding that trapping is biased towards high density, human dominated

landscapes is unlikely to substantially change.

We have shown that synthesis of rodent trapping studies to supplement curated rodent

distributions can counteract some of the inherent biases in these data and that they can add

further contextual data to host-pathogen association data. Together this supports their

inclusion in efforts to model endemic zoonotic risk and novel pathogen emergence.

Contribution of rodent trapping studies as data sources can be improved by adopting reporting

standards and practices consistent with Open Science, namely sharing of disaggregated

datasets alongside publication [54].

Future rodent trapping studies should be targeted towards regions that are currently under-

studied. Further information on rodent presence and abundance across West Africa will aid the

modelling of changing endemic zoonosis risk and the potential for novel pathogen emergence.

Sharing of disaggregated data alongside research publications should be promoted with

adoption of data standards to support ongoing data synthesis. Specifically, inclusion of exact

locations of trapping sites, trapping effort and the dates at which trapping occurred would

support more detailed inference of the spatio-temporal dynamics of host populations and the

risk of endemic zoonosis spillover events. Despite these challenges we propose that rodent

trapping studies can provide an important source of data to supplement curated datasets on

rodent distributions to quantify the risk of endemic zoonosis spillover events and the hazard of

novel pathogen emergence.

## 7. Author contributions

461 DS – Conceptualisation, Data Curation, Formal Analysis, Writing

462 LAA - Data Curation, Validation, Writing – Review and Editing

463  KEJ – Conceptualisation, Supervision, Writing – Review and Editing

464  DWJ &  RK – Supervision, Funding Acquisition, Writing – Review and Editing

## 8. Data availability

466  All data and code to reproduce this analysis is available in an archived Zenodo repository [32].

## 9. References

1.      Bernstein AS, Ando AW, Loch-Temzelides T, Vale MM, Li BV, Li H, et al. The costs and

benefits of primary prevention of zoonotic pandemics. Science Advances. 2022;8: eabl4183.

doi:10.1126/sciadv.abl4183

2.      Allen T, Murray KA, Zambrana-Torrelio C, Morse SS, Rondinini C, Di Marco M, et al.

Global hotspots and correlates of emerging zoonotic diseases. Nature Communications. 2017;8:

1124. doi:10.1038/s41467-017-00923-8

3.      Hassell JM, Begon M, Ward MJ, Fèvre EM. Urbanization and disease emergence:

Dynamics at the wildlife–livestock–human interface. Trends in Ecology & Evolution. 2017;32:

55–67. doi:10.1016/j.tree.2016.09.012

4.      García-Peña GE, Rubio AV, Mendoza H, Fernández M, Milholland MT, Aguirre AA, et al.

Land-use change and rodent-borne diseases: Hazards on the shared socioeconomic pathways.

Philosophical Transactions of the Royal Society B: Biological Sciences. 2021;376: 20200362.

doi:10.1098/rstb.2020.0362

5.    McMahon BJ, Morand S, Gray JS. Ecosystem change and zoonoses in the anthropocene.

Zoonoses and Public Health. 2018;65: 755–765. doi:10.1111/zph.12489

6.    Maudlin I, Eisler MC, Welburn SC. Neglected and endemic zoonoses. Philos Trans R Soc

Lond B Biol Sci. 2009;364: 2777–2787. doi:10.1098/rstb.2009.0067

7.    Halliday JEB, Allan KJ, Ekwem D, Cleaveland S, Kazwala RR, Crump JA. Endemic zoonoses

in the tropics: A public health problem hiding in plain sight. Vet Rec. 2015;176: 220–225.

doi:10.1136/vr.h798

8.    Molyneux D, Hallaj Z, Keusch GT, McManus DP, Ngowi H, Cleaveland S, et al. Zoonoses

and marginalised infectious diseases of poverty: Where do we stand? Parasites & Vectors.

2011;4: 106. doi:10.1186/1756-3305-4-106

9.    Judson SD, Rabinowitz PM. Zoonoses and global epidemics. Current Opinion in Infectious

Diseases. 2021;34: 385–392. doi:10.1097/QCO.0000000000000749

10.    Han BA, Schmidt JP, Bowden SE, Drake JM. Rodent reservoirs of future zoonotic diseases.

PNAS. 2015;112: 7039–7044. doi:10.1073/pnas.1501598112

11.    Gibb R, Albery GF, Mollentze N, Eskew EA, Brierley L, Ryan SJ, et al. Mammal virus

diversity estimates are unstable due to accelerating discovery effort. Biology Letters. 2021;18:

20210427. doi:10.1098/rsbl.2021.0427

12.    Fischer C, Gayer C, Kurucz K, Riesch F, Tscharntke T, Batáry P. Ecosystem services and

disservices provided by small rodents in arable fields: Effects of local and landscape

management. Journal of Applied Ecology. 2018;55: 548–558. doi:10.1111/1365-2664.13016

13.    Dobson FS, Oli MK. Fast and slow life histories of rodents. Rodent societies: an ecological

and evolutionary perspective. 2007; 99–105. Available:

https://www.jstor.org/stable/42902037

14.    Albery GF, Becker DJ. Fast-lived hosts and zoonotic risk. Trends in Parasitology.

2021;37: 117–129. doi:10.1016/j.pt.2020.10.012

15.    Gibb R, Redding DW, Chin KQ, Donnelly CA, Blackburn TM, Newbold T, et al. Zoonotic

host diversity increases in human-dominated ecosystems. Nature. 2020;584: 398–402.

doi:10.1038/s41586-020-2562-8

16.    Han BA, Kramer AM, Drake JM. Global patterns of zoonotic disease in mammals. Trends

in Parasitology. 2016;32: 565–577. doi:10.1016/j.pt.2016.04.007

17.    Gibb R, Franklinos LHV, Redding DW, Jones KE. Ecosystem perspectives are needed to

manage zoonotic risks in a changing climate. BMJ. 2020;371: m3389. doi:10.1136/bmj.m3389

18.    Boakes EH, McGowan PJK, Fuller RA, Chang-qing D, Clark NE, O'Connor K, et al. Distorted

views of biodiversity: Spatial and temporal bias in species occurrence data. PLOS Biology.

2010;8: e1000385. doi:10.1371/journal.pbio.1000385

19.    Bowler DE, Callaghan CT, Bhandari N, Henle K, Benjamin Barth M, Koppitz C, et al.

Temporal trends in the spatial bias of species occurrence records. Ecography. 2022;n/a:

e06219. doi:10.1111/ecog.06219

20.    Beck J, Böller M, Erhardt A, Schwanghart W. Spatial bias in the GBIF database and its

effect on modeling species' geographic distributions. Ecological Informatics. 2014;19: 10–15.

doi:10.1016/j.ecoinf.2013.11.002

21.    Plowright RK, Becker DJ, McCallum H, Manlove KR. Sampling to elucidate the dynamics

of infections in reservoir hosts. Philos Trans R Soc Lond B Biol Sci. 2019;374: 20180336.

doi:10.1098/rstb.2018.0336

22.    Carlson CJ, Farrell MJ, Grange Z, Han BA, Mollentze N, Phelan AL, et al. The future of

zoonotic risk prediction. Philosophical Transactions of the Royal Society B: Biological Sciences.

2021;376: 20200358. doi:10.1098/rstb.2020.0358

23.    Wille M, Geoghegan JL, Holmes EC. How accurately can we assess zoonotic risk? Dobson

AP, editor. PLoS Biol. 2021;19: e3001135. doi:10.1371/journal.pbio.3001135

24.    Redding DW, Lucas TCD, Blackburn TM, Jones KE. Evaluating bayesian spatial methods

for modelling species distributions with clumped and restricted occurrence data. PLOS ONE.

2017;12: e0187602. doi:10.1371/journal.pone.0187602

25.    Grace D, Mutua F, Ochungo P, Kruska R, Jones K, Brierley L, et al. Mapping of poverty and

likely zoonoses hotspots. 2012.

26.    Meerburg B, Singleton G, Kijlstra A. Rodent-borne diseases and their risks for public

health. Critical reviews in microbiology. 2009;35: 221–70. doi:10.1080/10408410902989837

27.    Galeh TM, Sarvi S, Montazeri M, Moosazadeh M, Nakhaei M, Shariatzadeh SA, et al. Global

status of toxoplasma gondii seroprevalence in rodents: A systematic review and meta-analysis.

Frontiers in Veterinary Science. 2020;7. Available:

https://www.frontiersin.org/article/10.3389/fvets.2020.00461

28.    Bovendorp RS, MCCleery RA, Galetti M. Optimising sampling methods for small mammal

communities in neotropical rainforests. Mammal Review. 2017;47: 148–158.

doi:10.1111/mam.12088

29.    Fichet-Calvet E, Lecompte E, Veyrunes F, Barrière P, Nicolas V, Koulémou K. Diversity

and dynamics in a community of small mammals in coastal guinea, west africa. Belgian Journal

of Zoology. 2009;139: 93–102. Available: http://www.vliz.be/nl/open-marien-

archief?module=ref&refid=204614

30.    Catalano S, Leger E, Fall CB, Borlase A, Diop SD, Berger D, et al. Multihost transmission of

schistosoma mansoni in senegal, 2015-2018. Emerging Infectious Diseases. 2020;26: 1234–

1242. doi:10.3201/eid2606.200107

31.    USAID. One health surveillance. PREDICT project. 2021 [cited 27 Sep 2021]. Available:

https://p2.predict.global/surveillance

32.    Simons D. DidDrog11/scoping_review: Repository to accompany submission. Zenodo;

2022. doi:10.5281/zenodo.7022637

33.    GBIF: The Global Biodiversity Information Facility. The global biodiversity information

facility. 2021 [cited 25 Apr 2021]. Available: https://www.gbif.org/

34.    National Geospatial-Intelligence Agency. NGA: GNS home. [cited 22 Mar 2021]. Available:

https://geonames.nga.mil/gns/html/

35.    Pebesma E. Simple features for r: Standardized support for spatial vector data. The R

Journal. 2018;10: 439–446. Available: https://journal.r-project.org/archive/2018/RJ-2018-

009/index.html

36.    R Core Team. R: A language and environment for statistical computing. Vienna, Austria:

R Foundation for Statistical Computing; 2020. Available: https://www.R-project.org/

37.    Database of Global Administrative Areas. GADM. 2022 [cited 25 Apr 2021]. Available:

https://gadm.org/index.html

38.    Socioeconomic Data and Applications Center. Gridded population of the world (GPW), v4

SEDAC. 2021 [cited 4 Feb 2021]. Available:

https://sedac.ciesin.columbia.edu/data/collection/gpw-v4

39.    European Space Agency Climate Change Initiative. Land cover classification gridded

maps from 1992 to present derived from satellite observations. 2022 [cited 3 Mar 2022].

Available: https://cds.climate.copernicus.eu/cdsapp#!/dataset/satellite-land-

cover?tab=overview

40.    Pedersen EJ, Miller DL, Simpson GL, Ross N. Hierarchical generalized additive models in

ecology: An introduction with mgcv. PeerJ. 2019;7: e6876. doi:10.7717/peerj.6876

41.    Wood SN. Generalized additive models: An introduction with r. 2nd ed. Chapman;

Hall/CRC; 2017.

42.    IUCN. The IUCN red list of threatened species. 2021. Available: www.iucnredlist.org

43.    GBIF: The Global Biodiversity Information Facility. Occurrence download. The Global

Biodiversity Information Facility; 2021. doi:10.15468/DL.S52MEH

44.    Gibb R, Carlson CJ, Farrell MJ. Viralemergence/clover: Preprint + zenodo. Zenodo; 2021.

doi:10.5281/zenodo.4435128

45.    Harvey E, Holmes EC. Diversity and evolution of the animal virome. Nat Rev Microbiol.

2022;20: 321–334. doi:10.1038/s41579-021-00665-x

46.    Basinski AJ, Fichet-Calvet E, Sjodin AR, Varrelman TJ, Remien CH, Layman NC, et al.

Bridging the gap: Using reservoir ecology and human serosurveys to estimate lassa virus

spillover in west africa. Wesolowski A, editor. PLoS Comput Biol. 2021;17: e1008811.

doi:10.1371/journal.pcbi.1008811

47.    Václavík T, Meentemeyer RK. Invasive species distribution modeling (iSDM): Are

absence data and dispersal constraints needed to predict actual distributions? Ecological

Modelling. 2009;220: 3248–3258. doi:10.1016/j.ecolmodel.2009.08.013

48.    Zeimes CB, Olsson GE, Ahlm C, Vanwambeke SO. Modelling zoonotic diseases in humans:

Comparison of methods for hantavirus in sweden. Int J Health Geogr. 2012;11: 39.

doi:10.1186/1476-072X-11-39

49.    Judson SD, LeBreton M, Fuller T, Hoffman RM, Njabo K, Brewer TF, et al. Translating

predictions of zoonotic viruses for policymakers. EcoHealth. 2018;15: 52–62.

doi:10.1007/s10393-017-1304-3

50.    Gibb R, Albery GF, Becker DJ, Brierley L, Connor R, Dallas TA, et al. Data proliferation,

reconciliation, and synthesis in viral ecology. BioScience. 2021;71: 1148–1156.

doi:10.1093/biosci/biab080

51.    Wilkinson DP, Golding N, Guillera-Arroita G, Tingley R, McCarthy MA. A comparison of

joint species distribution models for presence–absence data. Methods in Ecology and Evolution.

2019;10: 198–211. doi:10.1111/2041-210X.13106

52.    Fichet-Calvet E, Ölschläger S, Strecker T, Koivogui L, Becker-Ziaja B, Camara AB, et al.

Spatial and temporal evolution of lassa virus in the natural host population in upper guinea. Sci

Rep. 2016;6: 21977. doi:10.1038/srep21977

53.    Herrmann SM, Brandt M, Rasmussen K, Fensholt R. Accelerating land cover change in

west africa over four decades as population pressure increased. Commun Earth Environ.

2020;1: 1–10. doi:10.1038/s43247-020-00053-y

54.    Foster ED, Deardorff A. Open science framework (OSF). J Med Libr Assoc. 2017;105:

203–206. doi:10.5195/jmla.2017.88

Figure 1

Figure 2

**Mastomys natalensis**

**Rattus rattus**

**Mus musculus**

Detection/Non-detection ● Detection ● Non-detection

Figure 3

Figure 4

Figure 5