Ecological differences among hydrothermal vent symbioses may drive contrasting patterns

of symbiont population differentiation

Corinna Breusing[1*], Yao Xiao[2,3], Shelbi L. Russell[4], Russell B. Corbett-Detig[4], Sixuan Li[1], Jin

Sun[5], Chong Chen[6], Yi Lan[2,3], Pei-Yuan Qian[2,3], Roxanne A. Beinart[1*]

[1]Graduate School of Oceanography, University of Rhode Island, Narragansett, RI 02882, USA

[2]Department of Ocean Science, Division of Life Science and Hong Kong Branch of the Southern

Marine Science and Engineering Guangdong Laboratory (Guangzhou), The Hong Kong

University of Science and Technology, Hong Kong, China

[3]The Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou), Nansha,

Guangzhou, China

[4]Department of Biomolecular Engineering, University of California Santa Cruz, Santa Cruz, CA
95064, USA

[5]Institute of Evolution & Marine Biodiversity, Ocean University of China, Qingdao 266003,

China

[6]X-STAR, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokosuka,

Kanagawa Prefecture, Japan

*Corinna Breusing, Roxanne A. Beinart

Email: corinnabreusing@gmail.com, rbeinart@uri.edu

Keywords: Chemosynthetic symbiosis, hydrothermal vents, symbiont strain composition, niche

partitioning, homologous recombination, local adaptation, *Bathymodiolus septemdierum*

Competing interests: The authors declare no competing interests.

Abstract

The intra-host composition of horizontally transmitted microbial symbionts can vary across host

populations due to interactive effects of host genetics, environmental and geographic factors.

While adaptation to local habitat conditions can drive geographic subdivision of symbiont

strains, it is unknown whether these patterns are universal and how differences in ecological

characteristics among host-symbiont associations influence the genomic structure of symbiont

populations. To address this question, we sequenced metagenomes of different populations of the

deep-sea mussel *Bathymodiolus septemdierum*, which are common at Western Pacific deep-sea

hydrothermal vents and show characteristic patterns of niche partitioning with sympatric

gastropod symbioses. *Bathymodiolus septemdierum* lives in close symbiotic relationship with

sulfur-oxidizing chemosynthetic bacteria but supplements its symbiotrophic diet through filter-

feeding, enabling it to occupy ecological niches with less exposure to geochemical reductants.

Our analyses indicate that symbiont populations associated with *B. septemdierum* show

structuring by geographic location, but that the dominant symbiont strain is uncorrelated with

vent site. These patterns are in contrast to co-occurring *Alviniconcha* and *Ifremeria* symbioses

that exhibit greater symbiont nutritional dependence and occupy habitats with higher spatial

variability in environmental conditions. Our results suggest that relative habitat homogeneity

combined with sufficient symbiont dispersal and genomic mixing might promote persistence of

similar symbiont strains across geographic locations, while mixotrophy might decrease selective

pressures on the host to affiliate with locally adapted symbiont strains. Overall, these data

contribute to our understanding of the potential mechanisms influencing symbiont population

structure across a spectrum of marine microbial symbioses that vary in ecological niche and

relative host dependency.

Introduction

In the ocean, mutualistic symbioses between animals and bacteria are predominantly established through symbiont acquisition from the environment [1]. The composition of environmentally acquired symbionts within and across host populations is typically influenced by a combination of host genetic makeup, environmental selection, physical barriers to symbiont dispersal, symbiont competition and horizontal gene transfer, including homologous recombination (i.e., the exchange of homologous genetic material between bacterial strains or species) [2–4]. For instance, while host specificity, ecological partitioning or dispersal limitations may increase genetic divergence among symbiont populations, frequent recombination may promote genetic cohesiveness within a symbiont species. However, in most cases, the relative importance of these factors in shaping the genetic structure of symbiont populations remains unclear. Similarly, it is unknown how differences in ecological characteristics among symbiotic partnerships may impact patterns of symbiont population subdivision. Further work is necessary to untangle the effects of neutral and adaptive processes, as well as the dynamics of host-symbiont interaction, on horizontally transmitted symbiont populations.

Hydrothermal vents are dominated by invertebrates that obligately rely on the chemosynthesis of their bacterial symbionts for nutrition in the otherwise food-limited deep sea. Most symbiotic vent animals transmit their symbionts horizontally [5], such that at some early developmental stage, host animals must acquire their bacterial symbiont from the environment to survive. In these systems, where aposymbiotic larvae disperse between geographically distant and environmentally variable island-like vent fields, the acquisition of a local symbiont strain after settlement is thought to be common [6, 7]. Indeed, studies show that vent species typically

host distinct symbiont strains that vary by geographic site [7–13]. Our previous work on

Alviniconcha and *Ifremeria* symbioses from the Lau Basin indicated that site-specific variation in

symbiont populations is driven by environmental selection [7], though it is currently unknown

whether these processes apply to other hydrothermal vent symbioses that differ in their

ecological characteristics.

Owing to their ecological and evolutionary success, bathymodioline mussels belong to

the dominant fauna at deep-sea hydrothermal vents worldwide [14], with the genus

Bathymodiolus representing the most prominent member of the group. In the majority of cases,

Bathymodiolus species host thiotrophic gammaproteobacterial symbionts that use reduced sulfur

compounds as energy sources for chemosynthesis [15]. *Bathymodiolus septemdierum* is

widespread in the Indo-Pacific Ocean and although various names have been proposed for

different morphotypes, including *B. septemdierum*, *B. brevior*, and *B. marisindicus*, both

morphometric and genetic data have revealed that they represent populations of the same species,

leading to the other names being synonymized with *B. septemdierum* [16–18]. In the Western

Pacific, this mussel forms expansive beds that are typically arranged in concentric rings around

patches of sympatric gastropods of the genera *Alviniconcha* and *Ifremeria* [19]. Fine-scale

habitat partitioning among these holobionts is likely driven by contrasting thermal tolerances and

sulfide requirements for chemosynthesis, resulting in differential exposure of hosts and

symbionts to available geochemical reductants: *Alviniconcha* holobionts usually encounter the

highest and most variable sulfur concentrations, while *Bathymodiolus* holobionts experience the

lowest and most stable levels of sulfur compounds [19, 20]. The ability of *Bathymodiolus*

symbioses to thrive under diminished sulfide availability is hypothesized to be partly related to

the hosts' mixotrophic feeding mode [19, 21], which would decrease dependence on optimal

chemosynthetic production rates. Indeed, stable isotope and gill structure analyses suggest that *B.*

septemdierum is unlikely to fulfill its dietary requirements solely by assimilating symbiont-

derived organic carbon, especially given that symbiont densities appear to be too low to provide

the host with sufficient nutrition [22].

In this study we sequenced metagenomes of *B. septemdierum* populations from four

different vent sites in the Lau Basin to assess the population genetic structure and strain-level

functional variation of its thiotrophic symbiont, *Candidatus* Thiodubiliella endoseptemdiera [4],

in relation to host genotype, environmental factors and geographic location. To examine if

ecological differences and homologous recombination might play a role in symbiont geographic

partitioning, we subsequently compared the results to patterns in co-occurring *Alviniconcha* and

Ifremeria symbioses [7] that show strong contrasts in their niche preferences.

Materials and Methods

Sample collection, DNA extraction and sequencing

Mussel samples were collected from the Tu'i Malila (n = 40), ABE (n = 34), Tahi Moana (n =

11) and Tow Cam (n = 20) vent sites with the remotely operated vehicles (ROV) *ROPOS* and

Jason during research cruises to the Eastern Lau Spreading Center (ELSC), Southwest Pacific, in

2009 and 2016 (Fig. 1A; Table S1). The sites sampled are representative of the range of

geochemical habitat conditions *B. septemdierum* holobionts experience within the Lau Basin.

Upon ROV recovery, symbiont-bearing gill tissue of each individual was dissected on board ship

and frozen or stored in RNALater™ (Thermo Fisher Scientific, Inc., Waltham, MA, USA) at –

80°C until further analysis. DNA of 80 specimens was extracted from gill tissue snips with the

E.Z.N.A Mollusc DNA Kit (Omega Bio-tek, Norcross, GA) at the University of Rhode Island

and sent for plexWell384 library preparation and NovaSeq6000 2x150 bp sequencing at

Psomagen, Inc. (Rockville, MD, USA) (Table S1). A set of an additional 25 specimens was

extracted with the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) at the Hong Kong

University of Science and Technology and sent for metagenomic library preparation and

NovaSeq6000 2x150 bp sequencing at Novogene, Co. (Beijing, China) (Table S1), resulting in a

total of 105 samples for analysis. PERMANOVAs suggested that methodological differences had

negligible effects on patterns of symbiont genetic variation relative to biological factors (6.14%

versus 23.72 % explained variation), confirming the validity of combining the two metagenomic

datasets (Table S2). Raw reads for each individual were adaptor-clipped and quality-trimmed

with TRIMMOMATIC [23] before analysis. Host species identities were determined

morphologically and through assembly of the mitochondrial genome of each sample (see below).

Host transcriptome generation and population genomic analyses

To determine host population genomic structure, we *de novo* assembled the gill transcriptome of

B. septemdierum and used the resulting transcripts as reference for variant identification. Total

RNA of one RNALater™-preserved sample each from Tu'i Malila, ABE and Tahi Moana was

extracted with the Zymo Direct-zol RNA Miniprep Kit (Zymo Research, Inc., Irvine, CA, USA).

PolyA-enriched, 150-bp paired-end RNA-seq libraries were prepared with the TruSeq Stranded

mRNA Kit (Illumina, Inc., San Diego, CA, USA) and sequenced on a NovaSeq 6000 instrument

to a depth of 100 M reads per library by Psomagen, Inc. (Rockville, MD, USA). After initial

quality checks with FASTQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/), raw

reads from all three samples were trimmed with TRIMMOMATIC, error-corrected with

RCORRECTOR [24], filtered with TRANSCRIPTOMEASSEMBLYTOOLS

(https://github.com/harvardinformatics/TranscriptomeAssemblyTools), and co-assembled with

TRINITY in PASAFLY mode [25]. Transcript annotation and further curation, including removal of

transcript redundancies, followed the procedure in Breusing et al. [7]. Transcriptome

completeness was assessed with BUSCO [26] based on mollusk-specific conserved orthologous

genes. Single nucleotide polymorphisms (SNPs) among the 105 sequenced individuals were

subsequently estimated by mapping the metagenomic reads against the reference transcriptome

and assessing genotype likelihoods with ANGSD [27] based on a model correcting for deviations

from Hardy-Weinberg equilibrium (see ref. [7] for details). To evaluate host population genomic

differentiation between vent sites we determined pairwise fixation indices (F_{ST}s) using the

method described in Bhatia et al. [28] and performed principal component analyses of host

covariance matrices with the STATS and GGPLOT2 packages in R [29, 30].

Symbiont pangenome reconstruction and population genomic analyses

To assess symbiont population genomic structure, we reconstructed a pangenome for *Ca.*

Thiodubiliella endoseptemdiera out of individual metagenome-assembled genomes (MAGs) that

we used as scaffold for read mapping and variant calling. Using a pangenome instead of a single

reference genome allowed us to address limitations of individual assemblies, such as the risk of

missing accessory genes. Filtered reads from metagenomic sequencing were assembled with

METASPADES [31] using kmers from 21 to 121. Binning of MAGs was done following the

METAWRAP [32] pipeline and other binning procedures outlined in Breusing et al. [7]. MAGs

were taxonomically assigned with GTDB-TK [33] and evaluated for species-level genomic

similarity with FASTANI [34]. Ninety-one *Ca.* T. endoseptemdiera MAGs with CHECKM [35]

completeness scores > 85% and contamination scores < 10% were considered further for

pangenome reconstruction with PANAROO [36]. Genetic variants between symbiont populations

were identified by comparing metagenomic reads of all 105 samples against the *Ca.*

Thiodubiliella pangenome with FREEBAYES [37] as in Breusing et al. [7]. The identified variants

were subsequently used to infer symbiont population structure among vent sites through

calculation of pairwise F_{ST}s with SCIKIT-ALLEL (https://github.com/cggh/scikit-allel) and

principal coordinate analyses with the APE package in R [38]. F_{ST} values between symbiont

populations of individual mussel hosts and intra-host nucleotide diversities were calculated after

Romero Picazo et al. [39]. Statistical differences between genome-wide intra-host nucleotide

diversities among vent sites were determined via pairwise t-tests in R after confirming

assumptions of normality. The false discovery rate procedure was applied to correct for multiple

testing. To account for differences in variant detection probability within the pangenome, we

further calculated the pangenome fixation index (P_{ST}) based on gene diversities between vent

sites and mussel hosts after Romero Picazo et al. [40]. Associations of symbiont haplotypes with

host genotypes, geography and environment were assessed through redundancy analyses and

Mantel tests with the VEGAN and NCF packages [41, 42]. Geochemical data for these analyses

were taken from ref. [7]. Pairwise gene content differences between symbiont populations from

different vent sites were evaluated with PANPHLAN [43] based on coverage profiles of mapped

metagenomic reads against the *Ca.* Thiodubiliella pangenome. Genes that were present/absent in

~90% of samples from a given location were included in further analyses.

Symbiont strain decomposition and analysis

To quantify the abundance and number of distinct symbiont strains within each mussel host, we

inferred strain haplotypes on co-assembly graphs with STRONG [44]. All samples were

normalized to the same number of reads before analysis (~4.1 M reads/sample). STRONG was

run by evaluating multiple kmers for assembly (33, 55, 77) and using METABAT2 [45] for

binning. MAGs with qualities > 0.85 were considered for strain analysis. The BAYESPATHS

algorithm for strain resolution was run in five iterations, using five rounds of gene filtering and

allowing up to 15 strains to be detected. For comparative purposes, we also ran DESMAN [46]

on the metagenomic data within the STRONG pipeline (parameters: --nb_haplotypes 12 --

nb_repeat 5 --min_cov 1). Phylogenetic relationships among strains were assessed with

FASTTREE's Approximately-Maximum Likelihood method [47]. To evaluate if strain haplotypes

represented discrete or cohesive units, we calculated allele frequency spectra (AFS) for each

sample from variant calls performed for individual host-associated populations with FREEBAYES,

using the same parameters as above except that no threshold on the minimum alternate allele

fraction (-F) was applied and minimal post-filtering was performed to prevent bias in the shape

of the AFS. Strain abundance heatmaps and phylogenetic trees were produced with the

COMPLEXHEATMAP [48] and GGTREE [49] packages in R.

To confirm that strain detection across samples was not markedly influenced by potential

sample cross-contamination during processing or sequencing, we evaluated the mitochondrial

heterogeneity in each sample. Due to bottleneck effects during vertical transmission, intra-

individual mitochondrial diversity should be naturally low, though elevated levels are possible in

bivalves in the case of doubly uniparental inheritance where two divergent lineages of female (F)

and male (M) mitogenomes are present [50]. Individual host mitochondrial genomes were

assembled with MITOBIM [51] using a previously assembled *B. septemdierum* (AP014562)

mitogenome as bait for mitochondrial reconstruction. The mitogenome of sample R1931-6_463

could not be resolved through this approach and was reassembled with MITOBIM using the

sample's *COI* sequence as seed. Low frequency variants were then called with LOFREQ [52],

using default filters for coverage and strand bias, a minimum mapping quality of 30 and a

minimum base quality of 20.

Homologous recombination among symbiont genomes

To assess how homologous recombination might affect patterns of population genetic structure

in *Ca.* T. endoseptemdiera in comparison to co-occurring symbionts of *Alviniconcha* and

Ifremeria snails, we used three commonly applied methods: CLONALFRAMEML [53], MCORR

[54] and RHOMETA [55]. CLONALFRAMEML uses a maximum likelihood approach to calculate

recombination rates from MAGs, while the other two programs determine recombination rates

directly from metagenomic reads, thereby alleviating potential impacts of strain heterogeneity in

MAG-based recombination inference.

For the CLONALFRAMEML analyses we followed the approach by Osvatic et al. [56].

Alviniconcha and *Ifremeria* symbiont MAGs for this analysis were retrieved from Breusing et al.

[7]. We chose the eight most complete MAGs across the geographic range of each symbiont

species as this represented the lowest number of genomes available for a single symbiont. MAGs

of each symbiont species were first aligned in PROGRESSIVEMAUVE [57] with default parameters.

Core locally colinear blocks (LCBs) $\geqq$ 500 bp were then extracted with *stripSubsetLCBs* [57],

concatenated, and realigned with MAFFT [58] in auto mode. The core LCB alignment was

trimmed with TRIMAL [59] (parameters: -resoverlap 0.75 -seqoverlap 0.80) before phylogenetic

reconstruction in RAxML [60] (parameters: -f a -m GTRGAMMA -p 12345 -x 12345 -# 100).

The best RAxML tree and the trimmed core LCB alignment were subsequently used in

CLONALFRAMEML to infer homologous recombination events among symbiont MAGs. All

CLONALFRAMEML analyses were run with 100 pseudo-bootstrap replicates and applying the

transition/transversion rate ratios from RAxML as estimates for the kappa parameter. To

evaluate the algorithm's performance, we calculated the compound parameter Rδ [53]. Rδ values

greater than 1 typically imply that the relative rate of recombination to mutation R/θ is

underestimated due to repeated recombination events at any given genomic position that are not

accounted for by the program [53]. R values for the calculation of Rδ were inferred by

multiplying R/θ with an estimate of Watterson's θ according to the following formula: θ =

L$\times$G/a_n, where L is the sum of branch lengths in the recombination-corrected phylogeny, G is the

alignment length, and a_n is the (n-1)th harmonic number with n being the number of MAGs.

243   For calculations of recombination and mutation rates from metagenomic reads with

MCORR and RHOMETA we used the core and pangenomes of each symbiont species as reference

for mapping, respectively. Metagenomic datasets and symbiont pangenomes for *Alviniconcha*

and *Ifremeria* for this analysis were retrieved from ref. [7]. For each symbiont species, analyses

were run both across and within vent sites to get global and local estimates of recombination to

mutation rate ratios. Data were normalized to the same genome coverage across species and the

same number of samples per vent site to avoid bias in representation of symbiont strains. For all

local analyses, four samples per vent site were used, except in the case of ABE for the

Alviniconcha Gamma1 symbiont where only three samples were available. For all global

analyses, a total of 12 samples were used. All MCORR analyses were run with default settings.

However, as the resulting Φ_{pool} and γ/μ values appeared to be unrealistically high, likely due to a

poor fit of MCORR's coalescent model to our data, these results were not used for further

evaluation. For the RHOMETA analyses, the mean θ values estimated by the theta_est.nf pipeline

were used as input for the creation of lookup tables and inference of the recombination rate

parameter ρ. Lookup tables were generated for read depths of 3 to 85 for rho values from 1 to

100 by default. Rho values were divided by the recombination tract length (1000 bp) to get an

estimate of recombination rate per site. Recombination to mutation effect ratios were determined

according to Krishnan et al. [55] by multiplying the per site ρ/θ ratios with the recombination

tract lengths and substitution probabilities. Substitution probabilities were inferred by dividing

the number of filtered SNP sites by the pangenome length of each symbiont species.

Results

Host populations are undifferentiated across geographic localities

Co-assembly of representative mussel gill samples from Tu'i Malila, ABE and Tahi Moana

resulted in 35,060 transcripts that provided a reference for SNP estimation among host

populations (Table S3). The curated *B. septemdierum* gill transcriptome assembly comprised an

overall size of 57.71 Mb and was 88.90% complete. Assessment of variant sites based on

mappings of an average of 50,253,507 cleaned metagenomic reads per sample against the

reference transcriptome resulted in 279 SNPs for analysis of host population genomic

subdivision. Lowering the minimum sample calling threshold from 75% to 66% recovered 5,639

SNPs that produced equivalent results in terms of population structure. Both principal

component analyses based on host covariance matrices and pairwise F_{ST}s between vent localities

($\leqq 0.0139$) indicated an absence of geographic structuring in *B. septemdierum* and a more

individual-based partitioning of genomic diversity (Fig. 1B; Table 1). These patterns were

confirmed by Mantel tests that indicated no correlation between host identity-by-state and

geographic distance matrices (Table 2).

Symbiont populations show genetic structure across vent locations

Ninety-one species-level (ANI > 95%), moderate- to high-quality MAGs were combined to

reconstruct a ~3.26 Mb pangenome for *Ca.* T. endoseptemdiera, which consisted of 1040 core

and 4352 accessory genes (Table S4, S5). The MAGs underlying this pangenome ranged from

1.75 to 3.50 Mbp in size, had 374–2257 contigs and N50 values of 1836–9202 bp (Table S4).

Read mapping against the pangenome reference yielded 2,254 variant sites (SNPs and indels) for

symbiont population genomic analyses. Principal coordinate analyses and F_{ST} calculations based

on these variant sites indicated moderate partitioning of symbiont populations among vent

localities, while P_{ST} indices based on gene diversities suggested weaker differentiation (Fig. 1C;

Table 1: $F_{ST} = 0.0677–0.2949$, $P_{ST} = 0.0614–0.1465$). Maximum divergence based on pairwise

F_{ST}s was observed between Tow Cam and Tu'i Malila in accordance with the large geographic

distance between these sites, whereas P_{ST} indices did not show an obvious association with

geography (Table 1). Overall the degree of population differentiation in *Ca.* Thiodubiliella

appeared to be similar to that of the *Ifremeria* SOX symbiont ($F_{ST} = 0.0359– 0.3034$), but was

notably lower than in the co-occurring *Alviniconcha* Epsilon ($F_{ST} = 0.1012– 0.8682$) and

Gamma1 ($F_{ST} = 0.2892– 0.5589$) symbionts within the ELSC [7].

Intra-host nucleotide diversities were comparable among vent sites, except in the case of

Tow Cam which showed significantly lower π values than Tu'i Malila and ABE (Pairwise t-test:

p < 0.000002; Fig. S1), resulting in a higher distinctiveness of this site compared to others.

Similar to patterns in the sympatric *Ifremeria* and *Alviniconcha* symbionts, the association of *Ca.*

Thiodubiliella nucleotide and strain composition with vent locality was significant based on

traditional and partial Mantel tests (Table 2; p ≦ 0.002; r ≧ 0.6279), while no correlation

between symbiont and host genetic distance could be found. In contrast to the vent snail

symbioses, however, the dominant symbiont haplotype appeared to be unrelated to geography

(Table 2). Compared to levels of geographic subdivision, inter-sample symbiont population

differentiation was notably higher (Fig. 1D; Table S6: F_{ST} = 0.1708–0.7323, P_{ST} = 0.0231–

0.4035), suggesting that symbiont genetic diversity was predominantly partitioned among

individual hosts.

Some symbiont strains are associated with habitat geochemistry or geography

To get more detailed insights into strain composition within and among individual mussel hosts,

we decomposed strain haplotypes from the metagenomic dataset with BAYESPATHS. The

underlying *Ca*. Thiodubiliella MAG that was binned from the co-assembly and from which

strains were decomposed had a total coverage of 12929.84 (40.3–204.4 depth per sample).

Fourteen different strains (ANI: 98.88–99.70%) were resolved across samples via this method

and the settings chosen, with all strains being detectable in individual hosts – though with

drastically varying coverages (Fig. 2; Table S7). Although we cannot fully exclude that the

presence of some lowly abundant strains in individual samples might be due to sample cross-

contamination during processing or sequencing, analyses of mitochondrial heterogeneity

suggested that intra-sample mitochondrial diversity was in general low (0.00–1.32 variants/Kbp)

and relatively characteristic of vertically transmitted symbionts and organelles [61, 62]. One

mussel individual from ABE exhibited slightly elevated mitochondrial variant densities (3.54

variants/Kbp). The diversity in this sample might reflect natural variability, such as intra-host

mutations of the mitochondrial population, possibly due to increased oxidative stress in gill

tissue, or – in the case of doubly uniparental inheritance – potential presence of lowly abundant

M genomes that leaked into somatic tissue during transmission [50]. Overall, these results

indicate that potential sample cross-contamination is likely not an important factor influencing

the observed patterns (Table S8). Comparative strain quantifications based on DESMAN were

lower than those by BAYESPATHS, recovering only six strains (Fig. S2). Given that the detection

of symbiont strains tends to increase with both read depth and number of samples [45], the

amount of recovered strains presented here likely provides a snapshot of the true strain diversity

within *B. septemdierum*. In addition, due to the high amount of recombination within *Ca.*

Thiodubiliella (see below), the detected strains should be regarded as a continuum of possible

haplotypes rather than discrete ones, as also confirmed by the distribution of allele frequencies

within symbiont populations (Fig. S3). Of the 14 strains identified by BAYESPATHS, one was

dominant in almost all individuals regardless of vent site, while others showed a more distinctive

distribution according to geochemical habitat and correlated factors ($r^2_{adj} = 0.4617$; p = 0.001;

Fig. 2A, B). For example, based on redundancy analyses and assessments of intra-host strain

abundances, strains 4 and 13 were often relatively frequent in host individuals from Tow Cam,

whereas strains 2, 5 and 9 were typically abundant in host individuals from Tu'i Malila (Fig. 2).

The association of different strains with geographic location seemed not to be directly linked to

their phylogenetic relationships as more closely related strains were not necessarily abundant in

the same habitats (Fig. 2B). Given the dominance of a single symbiont strain at all vent sites, no

evidence for local adaptation could be found, as correlations between consensus variants

(representative of the most abundant strain) and environment were insignificant.

Symbiont populations from different vent sites exhibit variation in gene content

We profiled the metagenomic samples using pangenome-based phylogenomic analyses to

determine potential gene content variation among symbiont populations from distinct vent sites.

At a gene presence/absence threshold of ~90% between symbiont populations, the main

differences were observed for genes with unknown function, followed by genes involved in

DNA metabolism and antiviral defense. A small subset of differentially conserved genes was

associated with other metabolic categories (Fig. 3; Table S9). For example, *Ca.* T.

endoseptemdiera strains from Tow Cam, Tahi Moana and ABE contained a couple of

hydrogenase maturation and assembly factors and a heme synthase that were rare in populations

from Tu'i Malila (Table S9). By contrast, strains from Tu'i Malila typically encoded a

filamentous hemagglutinin, a copper metabolism related gene and an RNA polymerase that

decreased in frequency at ABE and Tahi Moana and were almost absent at Tow Cam (Table S9).

Homologous recombination contributes to cohesiveness of Ca. *T. endoseptemdiera populations*

Intra-specific homologous recombination plays a critical role in maintaining genetic connectivity

of bacterial populations [63]. To evaluate if homologous recombination might have a

homogenizing effect on *Ca.* T. endoseptemdiera population structure compared to co-occurring

Alviniconcha and *Ifremeria* symbiont species [7], we calculated the relative rates and importance

of recombination to mutation events from core genome alignments and metagenomic read

datasets of each symbiont group. R/θ ratios in core LCB alignments (1,111,427–2,480,742 bp) as

assessed by CLONALFRAMEML were at least four times higher in *Ca.* T. endoseptemdiera (0.44)

than in thiotrophic symbionts associated with *Alviniconcha* (0.04–0.10) and *Ifremeria* (0.10),

while r/m ratios were at least 2.5 times larger (2.38 compared to 0.28–0.94) (Table 3), suggesting

pervasive genome-wide recombination within *Ca.* Thiodubiliella (Fig. S4). Although all $R\delta$

values were markedly larger than 1 (Table 3) and CLONALFRAMEML is therefore likely to

underestimate R/θ within the inferred parameter ranges [53], the relative ratios among symbiont

species should be comparable. Equivalent inferences of recombination parameters from

metagenomic reads with RHOMETA suggested relatively high recombination to mutation rate and

effect ratios in all analyzed symbionts (Table 3; Fig. 4). Global ρ/θ and r/m values were largest in

the Gamma1 symbiont, followed by the two other gammaproteobacterial symbionts (*Ca.*

Thiodubiliella and Ifr-SOX) (Table 3; Fig. 4). Local estimates of ρ/θ and r/m showed notable

variation between sites for each symbiont species. Median per-site ρ/θ values were slightly

increased in the *Ifremeria* SOX symbiont compared to *Ca.* Thiodubiliella and the *Alviniconcha*

symbionts, while local recombination effect ratios showed the inverse pattern (Table 3, S10; Fig.

4). Similar to patterns of intra-host nucleotide diversities, the Tow Cam population of *Ca.*

Thiodubiliella had lower ρ/θ and r/m values than the other populations, further supporting a

higher genetic distinctiveness of this population (Table S10).

Discussion

Symbiont strain composition in horizontally transmitted symbioses is determined by a variety of

interacting factors, including host genetics, environment and geography [2]. At deep-sea

hydrothermal vent sites, local adaptation to geographically variable fluid geochemistry has been

shown to play a significant role in shaping strain-level variation of chemosynthetic

endosymbionts [7]. It is currently unknown how these patterns are affected by niche differences

among symbiotic associations that could influence the degree of selective pressures on both host

and symbiont. To address this question, we assessed symbiont strain composition in populations

of the deep-sea mussel *Bathymodiolus septemdierum* and compared our results to previous

findings in co-occurring vent snail symbioses [7] that segregate into contrasting ecological

niches [19].

Symbiont strain composition in *B. septemdierum* was unrelated to host genetics but

varied to some degree with vent habitat and geography. Although there are clear similarities to

the population genomic structures of sympatric *Alviniconcha* and *Ifremeria* symbionts given that

strain proportions were distinct among vent sites, the dominant *Ca.* Thiodubiliella

endoseptemdiera strain was contrastingly shared across localities without any evidence for local

adaptation. While we cannot directly determine the cause of the difference among the snail and

mussel symbionts given phylogenetic constraints and interlinked ecological variables that differ

among these taxa, there are notable habitat and feeding differences between the snails and

mussel species that could play a role. For example, the lower thermal and geochemical variation

in the microhabitat patches that *B. septemdierum* symbioses occupy [19, 20] might favor one

main *Ca.* T. endoseptemdiera strain across the symbiont's distributional range, thus allowing

host individuals to affiliate with the same symbiont genotype at different vent locations. In

addition, the increased mixotrophic capacity of *Bathymodiolus* relative to other chemosymbiotic

vent invertebrates, including *Alviniconcha* and *Ifremeria*, could reduce dependence on optimal

symbiont function and, thus, selection for a locally adapted strain, similar to what has been

suggested for facultative host-microbe associations [64]. *Bathymodiolus* species have a

functional albeit reduced digestive tract and can complement their symbiont-derived nutrition

with particulate and dissolved organic matter from their surroundings [65, 66], which is thought

to facilitate segregation into local habitat patches with barely detectable chemosynthetic

reductants [19, 20]. In fact, with sufficient food, bathymodioline mussels have been observed to

survive without their symbionts for months to possibly years, at least in a laboratory setting [67].

By contrast, there is no evidence that *Alviniconcha* and *Ifremeria* snails rely significantly on

heterotrophic feeding, though they still retain reduced and functional digestive systems [68–70].

Despite absence of habitat affiliation for the dominant *Ca.* T. endoseptemdiera strain,

symbiont populations exhibited differences in gene content and abundances of minority strains,

indicating that environmental conditions might at least partially shape the population genetic

structure of *Ca.* T. endoseptemdiera. For example, the strains from Tu'i Malila typically missed

several hydrogenase maturation factors that were present in the northern populations, possibly in

adaptation to the low hydrogen concentrations at this site. In addition, we observed some gene

content differences related to anti-viral defense. In line with the strong host and habitat

restriction of viruses at hydrothermal vents [71], the composition of viral communities might be

an important and possibly common mediator of symbiont genetic differentiation in the deep sea.

Similar impacts have been reported for symbiont populations associated with siboglinid

tubeworms [11, 72] and vent-associated snails [7].

We found that up to 14 distinct strains co-occurred within a single mussel individual,

although the actual number of strains is likely higher, given that strain detection depends notably

on read coverage, number of samples, sequencing method and bioinformatic algorithm

assumptions. In general, however, these estimates are within the range of variation reported for

thiotrophic symbionts of other *Bathymodiolus* species, including *B. brooksi* from cold seeps in

the Gulf of Mexico (up to 9 strains) [39] and *B. puteoserpentis* from hydrothermal vents on the

Mid-Atlantic Ridge (up to 16 strains) [73]. Symbiont diversity in bathymodioline mussels has

previously been shown to be maintained by functional heterogeneity among symbiont

subpopulations and is thought to confer metabolic flexibility of the symbiosis to changing

environmental conditions [73, 74]. Our data strongly support this hypothesis, suggesting that *B.*

septemdierum hosts retain one, broadly well-performing symbiont strain across their

distributional range and complement this strain by local, habitat-specific symbiont genotypes

with a variety of functions.

The widespread occurrence of one *Ca.* Thiodubiliella strain across vent localities

indicates that dispersal barriers are likely not the main drivers of symbiont population structure,

although overall population differentiation tended to increase with geographic distance (or

correlated factors). These patterns are in contrast to those in other *Bathymodiolus* symbionts

where geography is thought to be a major factor underlying population subdivision [12, 13], at

least over much larger spatial scales (1000s of km) than assessed here. It is possible that the

comparatively small distance among vent sites in the Lau Basin (<200 km), as well as contrasts

in oceanographic conditions or symbiont properties, contribute to these differences. Recent

biophysical models for the Lau Basin have implied that dispersal barriers within this region are

mostly absent [75], while evidence from a variety of biogeographic studies suggests that vent-

associated microbial communities are not significantly shaped by dispersal limitations [76].

However, the dispersal mechanisms of bacterial symbionts are not well understood, making it

difficult to assess how well dispersal patterns are represented by model simulations. In addition,

several population genetic studies on the symbionts of vent-associated animals, including those

of *Bathymodiolus* species from mid-ocean ridge systems and other mollusk taxa from the Lau

Basin, have highlighted the effect of topological barriers and geographic distance on symbiont

biogeography [7, 12, 13]. Despite contrasting evidence from these systems, it is possible that *Ca.*

T. endoseptemdiera faces less dispersal restrictions than co-occurring symbionts by employing

different dispersal strategies. For example, the bacterial symbionts of lucinid clams have been

hypothesized to be transported over long distances with seagrass fragments [56], whereas some

algal symbionts of gorgonian octocorals are known to hitchhike with the corals' larvae [77] – a

strategy that has also been suggested for the symbionts of bathymodioline mussels [78].

Considering the broad distribution of *B. septemdierum* in the Western Pacific and Indian Ocean

[16], vectored transport, if it can be confirmed, would greatly increase the dispersal ability of *Ca.*

T. endoseptemdiera, thereby reducing geographic partitioning among disparate populations. It is

currently unknown how widely distributed *Ca.* T. endoseptemdiera is, as sequence information

for bacterial symbionts of other *B. septemdierum* populations is scarce. Based on comparisons of

publicly available full-length 16S rRNA sequences, *Ca.* T. endoseptemdiera from the Lau Basin

is >97% identical to other *B. septemdierum* symbionts from the Izu-Ogasawara Arc vents in the

Northwest Pacific. In addition, our sequence searches within the Integrated Microbial NGS

platform (https://www.imngs.org) [79] at an identity cutoff of 100% suggest that *Ca.* T.

endoseptemdiera might occur at various locations in the Northeast Pacific Ocean and around the

Solomon Islands. Although further genomic analyses will be needed to confirm the species-level

designations of the recovered sequences, these results support a possible widespread distribution

of *Ca.* T. endoseptemdiera.

Compared to the genetic subdivision across geographic locations, the degree of symbiont

population divergence between individual mussels was approximately 2.5 times higher (based on

pairwise F_{ST}). The detected levels of inter-host symbiont population differentiation agree with

the patterns observed in other sulfur-oxidizing *Bathymodiolus* symbionts and were similarly

unrelated to host genotype [13, 39], suggesting that variation in symbiont composition between

host individuals is driven by neutral evolutionary processes [39]. Although *Bathymodiolus*

mussels have been shown to remain competent for symbiont uptake throughout their lifetime

[80], Romero Picazo et al. [39] suggested that strong genetic isolation among individual

symbiont populations can occur if environmental symbiont acquisition is restricted in later host

developmental stages and re-colonization happens increasingly through self-infection. This

model of symbiont transmission might explain how divergence of symbiont populations is more

strongly maintained on an individual than on a geographic level. Founder events combined with

possible priority effects [81] during symbiont infection may increase inter-host genetic

differentiation of symbiont populations. By contrast, occasional reshuffling of symbiont strains

between host and environment combined with adequate symbiont dispersal may promote

homologous recombination and mixing of the free-living symbiont pools, thereby diminishing

geographic and ecological isolation. *Candidatus* T. endoseptemdiera has previously been shown

to have some of the highest effective recombination rates observed for bacteria [4]. Our data

indicate that the relative rates of recombination to mutation in all analyzed mollusk symbionts

were elevated and surpassed an estimated threshold beyond which recombination is theoretically

sufficient to prevent population differentiation [82]. In *Ca*. T. endoseptemdiera high

recombination rates paired with diminished effects of environmental selection or dispersal

barriers might contribute to the lower geographic subdivision of this species compared to

sympatric *Alviniconcha* and *Ifremeria* symbionts. Similar observations have been made in

chemosynthetic lucinid clam symbioses, where increased levels of homologous recombination

are hypothesized to maintain genetic cohesiveness within a cosmopolitan symbiont species [56].

Conclusions

Our data provide evidence that symbiont populations associated with *Bathymodiolus*

septemdierum are structured among vent locations in the Lau Basin, but show no signs for

habitat affiliation of the dominant symbiont strain, in sharp contrast to thiotrophic symbionts of

co-occurring *Alviniconcha* and *Ifremeria* snails. While the lower dependence of *B. septemdierum*

on symbiont-derived nutrition might contribute to these patterns, our current data do not allow

disentangling these effects from impacts of correlated ecological partitioning. For example,

related to its mixotrophic feeding mode, *B. septemdierum* occupies niches with less variable fluid

flows, which might promote affiliations with the same symbiont strain across geographically

distinct, but ecologically similar habitats. Our findings further suggest that symbiont-related

factors, such as high recombination rates combined with the possibility of long-distance

dispersal, might decrease population differentiation among vent locations. However, the relative

importance of these processes for symbiont geographic structure and adaptation remains to be

determined. Future investigations of environmental versus intra-host strain abundances at

different vent sites as well as extracellular presence of symbionts on host larvae will help to shed

light on this question.

Acknowledgements

We thank the captains and crews of the R/Vs *Falkor* and *Thomas G. Thompson* as well as the

pilots of the ROVs *Ropos* and *Jason* for their invaluable support of the sample collections. The

technical staff at Novogene, Co. and Psomagen, Inc. are gratefully acknowledged for preparing

and sequencing the Illumina metagenomic and RNA-seq libraries. We further thank the

Kingdom of Tonga for providing access to their national waters and Abi Goodman for her help

in verifying the host species identities of mussel samples used in this study. This work was

funded by the Key Special Project for Introduced Talents Team of Southern Marine Science and

Engineering Guangdong Laboratory (Guangzhou) (GML2019ZD0409), the General Research

Fund of Hong Kong SAR (16101219, 16101822), the Schmidt Ocean Institute, the U.S. National

Science Foundation (grant numbers OCE-1536331, 1819530 and 1736932 to R.A.B. and

EPSCoR Cooperative Agreement OIA-#1655221) and the National Institutes of Health (grant

numbers 5K99GM135583-02 to S.L.R. and 5R35GM128932-03 to R.C.D.).

Competing Interests

536   The authors declare no competing interests.

Data Availability Statement

Raw metagenomic and RNA-seq reads are available in the National Center for Biotechnology

Information under BioProject numbers PRJNA855930 and PRJNA849542. The *Bathymodiolus*

septemdierum transcriptome has been deposited in the Transcriptome Shotgun Assembly

database under accession number GJZH00000000. Host mitochondrial genomes and *Ca.*

Thiodubiliella endoseptemdiera MAGs have been submitted to GenBank under accession

numbers listed in Supplementary Tables S1 and S4, respectively.

References

1.   Russell SL. Transmission mode is associated with environment type and taxa across

bacteria-eukaryote symbioses: a systematic review and meta-analysis. *FEMS Microbiol Lett*

2019; **366**: fnz013.

2.   LaJeunesse TC, Pettay DT, Sampayo EM, Phongsuwan N, Brown B, Obura DO, et al.

Long-standing environmental conditions, geographic isolation and host-symbiont

specificity influence the relative ecological dominance and genetic diversification of coral

endosymbionts in the genus *Symbiodinium*. *J Biogeogr* 2010; **37**: 785–800.

3.　McIlroy SE, Cunning R, Baker AC, Coffroth MA. Competition and succession among coral

endosymbionts. *Ecol Evol* 2019; **9**: 12767–12778.

4.　Russell SL, Pepper-Tunick E, Svedberg J, Byrne A, Ruelas Castillo J, Vollmers C, et al.

Horizontal transmission and recombination maintain forever young bacterial symbiont

genomes. *PLoS Genet* 2020; **16**: e1008935.

5.　Vrijenhoek RC. Genetic diversity and connectivity of deep-sea hydrothermal vent

metapopulations. *Mol Ecol* 2010; **19(20)**: 4391–4411.

6.　Vrijenhoek RC. Genetics and evolution of deep-sea chemosynthetic bacteria and their

invertebrate hosts. In: *The Vent and Seep Biota: Aspects from Microbes to Ecosystems*, S.

Kiel (ed.). 2010. Springer, Dordrecht, Netherlands, pp. 15–49.

7.　Breusing C, Genetti M, Russell SL, Corbett-Detig RB, Beinart RA. Horizontal transmission

enables flexible associations with locally adapted symbiont strains in deep-sea

hydrothermal vent symbioses. *Proc Natl Acad Sci USA* 2022; **119**: e2115608119.

8.　Di Meo CA, Wilbur AE, Holben WE, Feldman RA, Vrijenhoek RC, Cary SC. Genetic

variation among endosymbionts of widely distributed vestimentiferan tubeworms. *Appl*

Environ Microbiol 2000; **66(2)**: 651–658.

9.　Vrijenhoek RC, Duhaime M, Jones WJ. Subtype variation among bacterial endosymbionts

of tubeworms (Annelida: Siboglinidae) from the Gulf of California. *Biol Bull* 2007; **212(3)**:

180–184.

10.　Won Y, Jones WJ, Vrijenhoek RC. Absence of cospeciation between deep-sea mytilids and

their thiotrophic endosymbionts. *J Shellfish Res* 2008; **27(1)**: 129–138.

11. Perez M, Juniper SK. Insights into symbiont population structure among three

vestimentiferan tubeworm host species at Eastern Pacific spreading centers. *Appl Environ*

Microbiol 2016; **82**: 5197–5205.

12. Ho P-T, Park E, Hong SG, Kim E-H, Kim K, Jang S-J, et al. Geographical structure of

endosymbiotic bacteria hosted by *Bathymodiolus* mussels at eastern Pacific hydrothermal

vents. *BMC Evol Biol* 2017; **17**: 121.

13. Ücker M, Ansorge R, Sato Y, Sayavedra L, Breusing C, Dubilier N. Deep-sea mussels from

a hybrid zone on the Mid-Atlantic Ridge host genetically indistinguishable symbionts.

ISME J 2021; **15**: 3076–3083.

14. Roeselers G, Newton ILG. On the evolutionary ecology of symbioses between

chemosynthetic bacteria and bivalves. *Appl Microbiol Biotechnol* 2012; **94**: 1–10.

15. Duperron S, Lorion J, Samadi S, Gros O, Gaill F. Symbioses between deep-sea mussels

(Mytilidae: Bathymodiolinae) and chemosynthetic bacteria: diversity, function and

evolution. *C R Biol* 2009; **332**: 298–310.

16. Breusing C, Johnson SB, Tunnicliffe V, Vrijenhoek RC. Population structure and

connectivity in Indo-Pacific deep-sea mussels of the *Bathymodiolus septemdierum*

complex. *Conserv Genet* 2015; **16**: 1415–1430.

17. Thubaut J, Puillandre N, Faure B, Cruaud C, Samadi S. The contrasted evolutionary fates of

deep-sea chemosynthetic mussels (Bivalvia, Bathymodiolinae). *Ecol Evol* 2013; **3(14)**:

4748–4766.

18. Tunnicliffe V, Breusing C. Redescription of *Bathymodiolus septemdierum* Hashimoto and

Okutani, 1994 (Bivalvia, Mytilida, Mytilidae), a mussel broadly distributed across

hydrothermal vent locations in the western Pacific and Indian Oceans. *Zootaxa* 2022;

5214(3): 337–364.

19. Podowski E, Ma S, Luther G, Wardrop D, Fisher C. Biotic and abiotic factors affecting

distributions of megafauna in diffuse flow on andesite and basalt along the Eastern Lau

Spreading Center, Tonga. *Mar Ecol Prog Ser* 2010; **418**: 25–45.

20. Podowski EL, Moore TS, Zelnio KA, Luther GW, Fisher CR. Distribution of diffuse flow

megafauna in two sites on the Eastern Lau Spreading Center, Tonga. *Deep Sea Res Pt I*

2009; **56**: 2041–2056.

21. Fisher CR, Childress JJ, Arp AJ, Brooks JM, Distel D, Favuzzi JA, et al. Microhabitat

variation in the hydrothermal vent mussel, *Bathymodiolus thermophilus*, at the Rose Garden

vent on the Galapagos Rift. *Deep Sea Res Pt A* 1988; **35**: 1769–1791.

22. Dubilier N, Windoffer R, Giere O. Ultrastructure and stable carbon isotope composition of

the hydrothermal vent mussels *Bathymodiolus brevior* and *B.* sp. affinis *brevior* from the

North Fiji Basin, western Pacific. *Mar Ecol Prog Ser* 1998; **165**: 187–193.

23. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* 2014; **30**: 2114–2120.

24. Song L, Florea L. Rcorrector: efficient and accurate error correction for Illumina RNA-seq

reads. *GigaSci* 2015; **4**: 48.

25. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length

transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*

2011; **29**: 644–652.

26. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel and

streamlined workflows along with broader and deeper phylogenetic coverage for scoring of

eukaryotic, prokaryotic, and viral genomes. *Mol Biol Evol* 2021; **38(10)**: 4647–4654.

27. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics* 2014; **15**: 356.

28. Bhatia G, Patterson N, Sankararaman S, Price AL. Estimating and interpreting FST: The

impact of rare variants. *Genome Res* 2013; **23**: 1514–1521.

29. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016. Springer New York.

30. R Core Team. R: A language and environment for statistical computing. 2021. R

Foundation for Statistical Computing, Vienna, Austria.

31. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res* 2017; **27**: 824–834.

32. Uritskiy GV, DiRuggiero J, Taylor J. MetaWRAP—a flexible pipeline for genome-resolved

metagenomic data analysis. *Microbiome* 2018; **6**: 158.

33. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics* 2019; **36**: 1925–1927.

34. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI

analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018;

9: 5114.

35. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome Res 2015; **25**: 1043–1055.

36. Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, et al. Producing

polished prokaryotic pangenomes with the Panaroo pipeline. *Genome Biol* 2020; **21**: 180.

37. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing.

arXiv:12073907 [q-bio] 2012.

38. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics* 2019; **35**: 526–528.

39. Romero Picazo D, Dagan T, Ansorge R, Petersen JM, Dubilier N, Kupczok A. Horizontally

transmitted symbiont populations in deep-sea mussels are genetically isolated. *ISME J*

2019; **13**: 2954–2968.

40. Romero Picazo D, Werner A, Dagan T, Kupczok A. Pangenome evolution in

environmentally transmitted symbionts of deep-sea mussels is governed by vertical

inheritance. *Genome Biol Evol* 2022; **14(7)**: evac098.

41. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:

Community Ecology Package. 2020. https://CRAN.R-project.org/package=vegan

42. Bjornstad ON. ncf: Spatial Covariance Functions. 2020. https://CRAN.R-

project.org/package=ncf

43. Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, Maharjan S, et al.

Integrating taxonomic, functional, and strain-level profiling of diverse microbial

communities with bioBakery 3. *eLife* 2021; **10**: e65088.

44. Quince C, Nurk S, Raguideau S, James R, Soyer OS, Summers JK, et al. STRONG:

metagenomics strain resolution on assembly graphs. *Genome Biol* 2021; **22**: 214.

45.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies.

PeerJ 2019; **7**: e7359.

46.   Quince C, Delmont TO, Raguideau S, Alneberg J, Darling AE, et al. DESMAN: a new tool

for de novo extraction of strains from metagenomes. *Genome Biol.* 2017; **18(1)**: 181.

47.   Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for

large alignments. *PLoS ONE* 2010; **5(3)**: e9490.

48.   Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics* 2016; **32**: 2847–2849.

49.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT. ggtree: an R package for visualization and

annotation of phylogenetic trees with their covariates and other associated data. *Methods*

Ecol Evol 2017; **8**: 28–36.

50.   Breton S, Beaupré HD, Stewart DT, Hoeh WR, Blier PU. The unusual system of doubly

uniparental inheritance of mtDNA: isn't one enough? *Trends Genet* 2007; **23(9)**: 465–474.

51.   Hahn C, Bachmann L, Chevreux B. Reconstructing mitochondrial genomes directly from

genomic next-generation sequencing reads--a baiting and iterative mapping approach.

Nucleic Acids Res 2013; **41(13)**: e129.

52.   Wilm A, Aw PP, Bertrand D, Yeo GH, Ong SH, Wong CH, et al LoFreq: a sequence-

quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity

from high-throughput sequencing datasets. *Nucleic Acids Res* 2012; **40(22)**: 11189–11201.

53.   Didelot X, Wilson DJ. ClonalFrameML: efficient inference of recombination in whole

bacterial genomes. *PLoS Comput Biol* 2015; **11(2)**: e1004041.

54. Lin M, Kussell E. Inferring bacterial recombination rates from large-scale sequencing

datasets. *Nat Methods* 2019; **16(2)**: 199–204.

55. Krishnan S, DeMaere MZ, Beck D, Ostrowski M, Seymour JR, Darling AE. Rhometa:

Population recombination rate estimation from metagenomic read datasets. *bioRxiv* 2022;

doi: 10.1101/2022.08.04.502887.

56. Osvatic JT, Wilkins LGE, Leibrecht L, Leray M, Zauner S, Polzin J, et al. Global

biogeography of chemosynthetic symbionts reveals both localized and globally distributed

symbiont groups. *Proc Natl Acad Sci USA* 2021; **118**: e2104378118.

57. Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene

gain, loss and rearrangement. *PLoS ONE* 2010; **5(6)**: e11147.

58. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 2002; **30(14)**:

3059–3066.

59. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009; **25(15)**:

1972–1973.

60. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* 2014; **30(9)**: 1312–1313.

61. Russell SL, Cavanaugh CM. Intrahost genetic diversity of bacterial symbionts exhibits

evidence of mixed infections and recombinant haplotypes. *Mol Biol Evol* 2017; **34(11)**:

2747–2761.

62. Perez M, Breusing C, Angers B, Beinart RA, Won YJ, Young CR. Divergent paths in the

evolutionary history of maternally transmitted clam symbionts. *Proc Biol Sci* 2022;

289(1970): 20212137.

63. González-Torres P, Rodríguez-Mateos F, Antón J, Gabaldón T. Impact of homologous

recombination on the evolution of prokaryotic core genomes. *mBio* 2019; **10(1)**: e02494–

18.

64. Leydet KP, Hellberg ME. Discordant coral–symbiont structuring: factors shaping

geographical variation of *Symbiodinium* communities in a facultative zooxanthellate coral

genus, *Oculina*. *Coral Reefs* 2016; **35**: 583–595.

65. Page HM, Fiala-Medioni A, Fisher CR, Childress JJ. Experimental evidence for filter-

feeding by the hydrothermal vent mussel, *Bathymodiolus thermophilus*. *Deep Sea Res Pt A*

1991; **38**: 1455–1461.

66. Le Pennec M, Donval A, Herry A. Nutritional strategies of the hydrothermal ecosystem

bivalves. *Prog Oceanogr* 1990; **24**: 71–80.

67. Colaço A, Bettencourt R, Costa V, Lino S, Lopes H, Martins I, et al. LabHorta: a controlled

aquarium system for monitoring physiological characteristics of the hydrothermal vent

mussel *Bathymodiolus azoricus*. *ICES J Mar Sci* 2011; **68**: 349–356.

68. Yang Y, Sun J, Chen C, Zhou Y, Van Dover CL, Wang C, et al. Metagenomic and

metatranscriptomic analyses reveal minor-yet-crucial roles of gut microbiome in deep-sea

hydrothermal vent snail. *Anim Microbiome* 2022; **4(1)**: 3.

69. Laming SR, Hourdez S, Cambon-Bonavita MA, Pradillon F. Classical and computed

tomographic anatomical analyses in a not-so-cryptic *Alviniconcha* species complex from

hydrothermal vents in the SW Pacific. *Front Zool* 2020; **17**: 12.

70.   Chen C, Watanabe HK, Gena K, Johnson SB. Anatomical shifts linked with unusual diets in

728   deep-sea snails. *Ecology* 2022; doi: 10.1002/ecy.3847.

729   71.   Thomas E, Anderson RE, Li V, Rogan LJ, Huber JA. Diverse viruses in deep-sea

730   hydrothermal vent fluids have restricted dispersal across ocean basins. *mSystems* 2021;

731   **6(3)**: e0006821.

732   72.   Perez M, Angers B, Young CR, Juniper SK. Shining light on a deep-sea bacterial symbiont

733   population structure with CRISPR. *Microbial Genomics* 2021; **7**: 000625.

734   73.   Ansorge R, Romano S, Sayavedra L, Porras MÁG, Kupczok A, Tegetmeyer HE, et al.

735   Functional diversity enables multiple symbiont strains to coexist in deep-sea mussels. *Nat*

736   *Microbiol* 2019; **4**: 2487–2497.

737   74.   Ikuta T, Takaki Y, Nagai Y, Shimamura S, Tsuda M, Kawagucci S, et al. Heterogeneous

738   composition of key metabolic gene clusters in a vent mussel symbiont population. *ISME J*

739   2016; **10**: 990–1001.

740   75.   Breusing C, Johnson SB, Mitarai S, Beinart RA, Tunnicliffe V. Differential patterns of

741   connectivity in Western Pacific hydrothermal vent metapopulations: A comparison of

742   biophysical and genetic models. *Evol Appl* 2021; doi: 10.1111/eva.13326.

743   76.   Dick GJ. The microbiomes of deep-sea hydrothermal vents: distributed globally, shaped

744   locally. *Nat Rev Microbiol* 2019; **17**: 271–283.

745   77.   Wirshing HH, Feldheim KA, Baker AC. Vectored dispersal of *Symbiodinium* by larvae of a

746   Caribbean gorgonian octocoral. *Mol Ecol* 2013; **22**: 4413–4432.

747   78.   Won Y-J, Hallam SJ, O'Mullan GD, Pan IL, Buck KR, Vrijenhoek RC. Environmental

748   acquisition of thiotrophic endosymbionts by deep-sea mussels of the genus *Bathymodiolus*.

749   *Appl Environ Microbiol* 2003; **69**: 6785–6792.

79.   Lagkouvardos I, Joseph D, Kapfhammer M, Giritli S, Horn M, Haller D, et al. IMNGS: A

comprehensive open resource of processed 16S rRNA microbial profiles for ecology and

diversity studies. *Sci Rep* 2016; **6**: 33721.

80.   Wentrup C, Wendeberg A, Schimak M, Borowski C, Dubilier N. Forever competent:

deep-sea bivalves are colonized by their chemosynthetic symbionts throughout their

lifetime. *Environ Microbiol* 2014; **16**: 3699–3713.

81.   Debray R, Herbert RA, Jaffe AL, Crits-Christoph A, Power ME, Koskella B. Priority

effects in microbiome assembly. *Nat Rev Microbiol* 2022; 20: 109–121.

82.   Fraser C, Hanage WP, Spratt BG. Recombination and the nature of bacterial speciation.

Science 2007; **315**: 476–480.

Tables

Table 1 F_{ST} values between host (above diagonal) and symbiont (below diagonal) populations from different vent sites based on genomic variants. For the symbiont, the pangenome fixation index (P_{ST}) based on gene diversities is given in brackets.

	Tow Cam	Tahi Moana	ABE	Tu'i Malila
Tow Cam	*	0.0139	0.0049	0.0063
Tahi Moana	0.1818 (0.1172)	*	–0.0004	0.0021
ABE	0.1888 (0.0753)	0.0677 (0.1012)	*	–0.0002
Tu'i Malila	0.2949 (0.1250)	0.1896 (0.1465)	0.1678 (0.0614)	*

Table 2 Mantel tests between host, symbiont and geographic distance matrices. Significant tests are indicated in bold.

	Symbiont nucleotide counts		Dominant symbiont haplotype		Symbiont strain abundance		
	Host-Symbiont	*Symbiont-Geography*	*Host-Symbiont*	*Symbiont-Geography*	*Host-Symbiont*	*Symbiont-Geography*	*Host-Geography*
r statistic	–0.0163	**0.6655**	0.0552	–0.0158	–0.0162	**0.6279**	–0.0215
p-value	0.3623	**0.0002**	0.0742	0.2409	0.3473	**0.0002**	0.1727

Table 3 Rate and effect ratios of homologous recombination to mutation events in dominant chemosynthetic symbionts of *Bathymodiolus septemdierum*, *Alviniconcha* spp. and *Ifremeria nautilei* from the Lau Basin and Tonga Volcanic Arc. Estimates are given for inferences with i) CLONALFRAMEML based on MAGs and ii) RHOMETA based on metagenomic read alignments. For the CLONALFRAMEML analyses a 95% confidence interval is given for each ratio and alignment lengths for each symbiont are reported, with the percentage of the pangenome lengths provided in brackets. For the RHOMETA analyses, both global (across sites within species) and local (within sites within species) estimates of recombination are shown. R/θ, ρ/θ = recombination to mutation rate ratio inferred by CLONALFRAMEML and RHOMETA, respectively. r/m = relative effects of recombination to mutation. Rδ = key determinant for algorithm performance in CLONALFRAMEML.

	Ca. **Thiodubiliella** (*B. septemdierum*)	**Epsilon** (*A. boucheti*)	**Gamma1** (*A. kojimai, A. strummeri*)	**Ifr-SOX** (*I. nautilei*)
CLONALFRAMEML				
# MAGs	8	8	8	8
Alignment length [bp]	1,111,427 (34.11%)	1,152,639 (65.32%)	2,090,501 (51.74%)	2,480,742 (51.40%)

LCBs	550	672	895	998
R/θ [95% CI]	0.44 (0.42–0.45)	0.10 (0.10–0.11)	0.04 (0.03–0.05)	0.10 (0.09–0.11)
r/m [95% CI]	2.38 (2.23–2.53)	0.28 (0.25–0.30)	0.58 (0.48–0.70)	0.94 (0.81–1.09)
Rδ	85247.70	14109.17	7892.98	7713.08
RHOMETA				
Global				
# samples	12	12	12	12
Pangenome size [bp]	3,258,339	1,764,546	4,040,567	4,826,737
ρ/θ	51.14	35.19	61.81	49.47
r/m	1601.42	1518.53	2319.64	1636.89
Local				
# samples per site	4	4	3–4	4
Pangenome size [bp]	3,258,339	1,764,546	4,040,567	4,826,737
Median ρ/θ [range]	47.80 (29.86–59.10)	48.88 (38.84–74.85)	43.13 (31.90–60.99)	59.31 (45.82–71.17)
Median r/m [range]	912.94 (648.19–1095.06)	1105.59 (704.38–1827.99)	1004.44 (770.89–1036.48)	736.52 (655.45–1233.57)

Figure Legends

Fig. 1 Sampling locations of *Bathymodiolus septemdierum* in the Eastern Lau Spreading Center (A) and genetic structure of host (B) and symbiont (C) populations across vent sites. Vent locations are between 9 and 191 km apart. Host populations are genetically undifferentiated, while moderate structuring by geography is observed in the symbiont. (D) Pairwise F_{ST} values (lower diagonal) and P_{ST} values (upper diagonal) among symbiont populations of single mussel hosts. Compared to patterns observed on a geographic level, differentiation on a per-sample level is notably higher, indicating that symbiont genetic diversity is strongly driven by variation among host individuals. Map data ©2023 Google.

Fig. 2 Environmental associations (A) and intra-host abundances (B) of decomposed *Ca.* Thiodubiliella endoseptemdiera strains. Although one symbiont strain is clearly dominant irrespective of sampling locality, some strains are associated with fluid geochemistry or geography of the different vent sites, with strains 4 and 13 being relatively frequent in host individuals from Tow Cam, and strains 2, 5 and 9 being abundant in host individuals from Tu'i Malila. The environmental associations among the distinct symbiont strains are not necessarily related to phylogenetic proximity. Phylogenetic relationships are depicted as a mid-point rooted cladogram. Node values show local bootstrap support.

Fig. 3 Gene content differences among *Ca.* Thiodubiliella endoseptemdiera populations from vent locations in the Eastern Lau Spreading Center. Samples are ordered based on similarity. Symbiont populations appear to differ mostly in genes associated with unknown functions, DNA metabolism and anti-viral defense, with smaller variation being observed in other metabolic

802   categories. Overall, gene content is more similar among the northern populations at Tow Cam,

803   Tahi Moana and ABE compared to the southernmost population at Tu'i Malila.

805   **Fig. 4** Local (within vent site) recombination rate (A) and effect (B) ratios for chemosynthetic

806   symbionts of *Bathymodiolus*, *Alviniconcha* and *Ifremeria* species from the Lau Basin. Global

807   estimates (across vent sites) are superimposed on the boxplots as black dots.