Natural and pathological aging distinctively impact the vomeronasal detection system and social behavior

Adrián Portalés[1], Pablo Chamero[2], Sandra Jurado[1]*

[1]Institute of Neuroscience CSIC-UMH, San Juan de Alicante, 03550, Alicante, Spain

[2] Laboratoire de Physiologie de la Reproduction et des Comportements, CNRS, IFCE, INRAE, University of Tours, 37380, Nouzilly, France

*Correspondence: Dr. Sandra Jurado, Institute of Neuroscience CSIC-UMH, San Juan de Alicante, 03550, Alicante, Spain; sjurado@umh.es

KEYWORDS: Aging, pheromone, vomeronasal system, social behavior, odor discrimination, neurodegeneration

ABSTRACT

Normal aging and many age-related disorders such as Alzheimer´s disease cause deficits in olfaction, however it is currently unknown how natural and pathological aging impact the detection of social odors which might contribute to the impoverishment of social behavior at old age further worsening overall health. Here, we investigated the effect of aging in the recognition of social cues and the display of social behavior. Our findings indicate that aging distinctively disrupts the processing of social olfactory cues decreasing social odor exploration, discrimination and habituation in both wild type senescent (2-year-old) mice and in 1-year-old double mutant model of Alzheimer´s disease (APP/PS1[Het]). Furthermore, social novelty was diminished in 1-year-old APP/PS1[Het] mice, indicating that alterations in the processing of social cues are accelerated during pathological aging. Analysis of the vomeronasal organ, the main gateway to pheromone-encoded information, indicated that natural and pathological aging distinctively reduce the neurogenic ability of the vomeronasal sensory epithelium. Cell proliferation remained majorly preserved in 1-year model of Alzheimer´s disease (APP/PS1[Het]), whereas naturally aged animals exhibited significant deficiencies in the number of mature, proliferative and progenitor cells. This study reveals fundamental differences in the cellular processes by which natural and pathological aging disrupt the exploration of social cues and social behavior.

BACKGROUND

Social recognition is essential for survival allowing to appropriately adapt behavior across a variety of contexts (Adolphs, 2001, Brennan and Kendrick, 2006). As aging progresses, social odors have been shown to elicit attenuated responses (Hlinák and Krejcí, 1991, Garrat et al., 2011), a phenomenon interpreted as a natural consequence of age-related decline in sensory perception (Doty et al., 1984, Murphy et al., 2002, Doty and Kamath, 2014). In humans, social impoverishment has been identified as a major aggravating factor for decreased life expectancy, and an indicator of the appearance of dementia and neurodegenerative disorders such as Alzheimer´s disease (AD) (Heinrichs et al., 2003, Wilson et al., 2007, Donovan et al.,

2016). Despite the central role of social interaction in maintaining overall well-being, the mechanisms by which aging, natural or pathological, alter social information processing remain unclear.

Mouse social communication strongly depends on chemosignals comprised by volatile and non-volatile molecules (pheromones) (Beynon and Hurst, 2004, Yoon et al., 2005, Noack et al., 2010, Chamero et al., 2012, Li and Dulac, 2018). Volatile odors are primarily detected at the main olfactory epithelium (MOE) which projects to the main olfactory bulb (MOB), whereas pheromones are mainly detected by the vomeronasal sensory epithelium (VSE) (Jacobson, 1813, Halpern and Martínez-Marcos, 2003, Munger et al., 2009) which connects to the accessory olfactory bulb (AOB) (Ennis et al., 2015). Despite the segregated connectivity, both systems have overlapping functions (Brennan and Zufall, 2006, Zufall and Leinders -Zufall, 2007, Baum and Kelliher, 2009).

A remarkable property of the VSE is that contains proliferative niches capable of generating functional neurons during adulthood (Graziadei and Monti Graziadei, 1983, Weiler et al., 1999, Whitman and Greer, 2009, Brann and Firestein, 2010, Oboti et al., 2015). The neurogenic properties of olfactory structures are known to decline with age, a phenomenon proposed to underlie the olfactory deficits associated to both natural aging and neurodegenerative disorders (Doty et al., 1984, Murphy et al., 2002, Ahlenius et al., 2009, Doty and Kamath, 2014, Mobley et al., 2014, Devanand, 2016). However, how natural and pathological aging affect the regenerative capacity of the VSE has been scarcely studied despite being central to socio-sexual cue detection. An elegant study by Brann and Firestein (2010) showed that the proliferative capacity of the marginal region of the VSE of 2-year-old animals was attenuated in comparison to young animals. More recently, Mechin et al. (2021) reported a significant reduction in mature Olfactory Marker Protein (OMP)-expressing cells (OMP+), indicating structural modifications of the aged VNO. However, it is currently unclear whether these cell alterations have any impact in odor detection and social behavior, particularly in animal models of neurodegeneration.

To gain insight into these questions, we investigated the impact of natural and pathological

aging in social odor-evoked sniffing behavior, social-odor habituation/dishabituation and social

behavior during the aging process of wild type C57/BL6 mice and amyloid precursor protein

(APP) and presinilin-1 (PS1) double transgenic mice (APP/PS1^{Het} mice, therein), a widely used

animal model of AD (Jankowsky et al., 2001, Jankowsky et al., 2004). Analysis of the number

of progenitors, proliferative and mature neurons revealed that naturally aged animals show

reduced neuronal proliferation and decreased levels of sensory mature OMP$^+$ neurons. In

contrast, the VSE of 1-year-old APP/PS1^{Het} mice exhibited normal levels of mature neurons

and cell regeneration, suggesting that natural and pathological aging distinctively impact the

neurogenic capacity of the VSE.

We explored the functional consequences of these cell alterations observing reduced

exploration time of social odors and decreased performance in a social odor habituation-

dishabituation test in both senescent and middle age APP/PS1^{Het} mice, suggesting that

despite the overall normal VSE structure and cell composition, APP/PS1^{Het} animals exhibit an

accelerated decay of social cue decay. Moreover, 1-year-old APP/PS^{Het} mice exhibited a

significant reduction of social novelty which was no apparent in naturally aged animals,

exposing fundamental differences in how natural and pathological aging impact the regulation

of social behavior.

METHODS

Animals. Experiments were performed in either control C57/BL6 mice (young: 2-4 months-

old; adult: 6-8 months-old; middle age: 12-14 months-old; old (senescent): 20-24 months-old)

or APP/PS1 mice on a C57/BL6 genetic background (young: 2-4 months-old; middle-age: 12-

14 months-old) (Jackson Labs, Stock No. 004462, MMRRC Stock No. 34829). APP/PS1^{Het}

mice express a chimeric mouse/human amyloid precursor protein (Mo/HuAPP695swe) and a

mutant human presenilin 1 (PS1-dE9). These animals develop beta-amyloid deposits at ~ 6

months of age and exhibit early-onset cognitive impairments (Jankowsky et al., 2001, 2004,

Reiserer et al., 2007). Animals were kept group-housed and experimentally-naïve during all

their lifetime under pathogen-free conditions. Animals were housed in ventilated cages with

free access to food and water on a 12 h light/dark cycle.

All experiments were performed according to Spanish and European Union regulations

regarding animal research (2010/63/EU), and the experimental procedures were approved by

the Bioethical Committee at the Instituto de Neurociencias and the Consejo Superior de

Investigaciones Científicas (CSIC).

VNO dissection. Mice were perfused transcardially with PBS, pH 7.4, followed by 4%

paraformaldehyde in 0.1 M phosphate buffer (PB), pH 7.4. A fixed mouse head was placed

under a scope and the lower jaw was removed to get a palate view. To facilitate VNO

extraction, the palate and the nasal septum were removed and the bilateral VNOs were split

in two parts with a gentle twisting motion. Finally, the cartilaginous tissue that covers each

portion was carefully removed to extract the VNOs. Tissue was incubated in a 30 % sucrose

solution for cryoprotection and kept at -4 ºC until sectioning. Samples were embedded in OCT

and frozen at -80 ºC for cryosectioning.

Immunohistochemistry. VNOs embedded in OCT-medium were sliced in 16 μm thick

sections in a cryostat apparatus (Leica CM 3050S). Slices were incubated in blocking solution

containing 0.5 % Triton X-100 and 5 % normal horse serum in 0.1 M TBS for 2 h at room

temperature (RT). Primary antibody incubation was performed overnight (o/n) at 4 ºC against

OMP (Wako goat polyclonal; 1:2000), PCNA (Sigma, rabbit monoclonal; 1:2000); Sox2 (R&D

Systems, goat polyclonal; 1:300). For PCNA immunostaining, slices were incubated in 10 mM

citrate buffer (100 ºC; pH, 6.0) for 5 minutes prior staining. Sections were incubated with Alexa

Fluor 488- or 594- conjugated secondary antibodies (Jackson Laboratories, 1:500) for 2 h at

RT. Hoechst (Sigma, 1:10000) was added during 5 minutes after the secondary antibody

incubation for nuclei visualization. Imaging was performed using a vertical confocal

Microscope Leica SPEII. Final images were assembled in Adobe Illustrator.

Sox2 fluorescence intensity at the VNO supporting cell layer (SCL). For determining the Sox2 fluorescent labeling area in the VNO SCL where single-cell quantification was limited by the densely packed disposition of the cells, we calculated the corrected total fluorescence (CTF) of the area of interest employing the Freehand ROI tool of Image J. CTF was calculated by subtracting the background fluorescence from a minimum of 3 sections, 16 μm thick, from at least 4 animals per condition.

Stereology, cell quantification and AOB volume estimation

The total number of olfactory mature neurons (OMP$^+$ neurons) in the entire VSE was estimated using stereology (optical fractionator method, Wong et al., 2018) employing Stereo Investigator (MBF Bioscience). Two different tools are combined in this unbiased quantification method: a 3D optical dissector for cell counting and a fractionator, which is based on a systematic, uniform and random sampling over a known area of the tissue (West and Gundersen, 1990, Parrish-Aungst et al., 2007, Dorph-Petersen and Lewis, 2011). The number of neurons was estimated as:

$$N = (1/h \times 1/ssf \times 1/asf) \times \sum Q - t$$

In which $\sum Q$ is the total cell number in the region of interest acquired using the optical dissector method; **t** stands for the mean mounted section thickness; **h** is the optical dissector height; **asf** is the area sampling fraction and **ssf** is the frequency of sampling (section sampling fraction). For the VSE stereological analysis, sampling was performed with a 20x objective (Leica, NA 0.6), the counting frame area was 2500 μm^2 and the sampling grid area was 22500 μm^2. H for VSE stereology was 8 μm with 1 μm as upper and lower guard zones and t was set at 10 μm. Quantifications were performed for marginal regions of the VSE dividing the area in segments of equal length as previously described (Giacobini et al., 2000, Brann and Firestein, 2010). For the cell quantification in the anterior-posterior regions, the VSE was distributed in 10 sections per animal. Results were divided by the area to obtain the number of cells/mm^2 in each subdivision. Quantification of the cell number in the marginal area was performed in three

slices of each animal delineated with a 50x50 dissector employing Stereo Investigator (MBF

Bioscience).

VSE volume was calculated by multiplying the sampled area by the slice thickness (16 μm)

and the number of series (10 slices per animal). Ten sections per series were analyzed to

obtain an estimation of the VSE area and volume in young (4 months-old), aged (24 months-

old), middle aged (12-months old) APP/PS1^{WT} and middle aged APP/PS1^{Het} mice.

The volume of the AOB was estimated by measuring the AOB area and multiplying the

sampled area by the section thickness (50 μm) and the number of analyzed sections (5 series

per animal).

Odorants

Social and non-social scents were employed to discern the impact of aging in social odor

perception. As non-social scents, we used **i)** the banana-like odor isoamyl acetate (IA) (Sigma)

shown to be primarily detected at the MOE level (Xu et al., 2005) with neutral valence within

a broad dilution range (Root et al., 2014, Fortés-Marco et al., 2015, Saraiva et al., 2016), and

ii) food pellets to perform a food finding test (FFT, see experimental details below). As a social

scent, we used mouse urine as urine from conspecifics has been shown to elicit robust VNO

activity (Chamero et al., 2017). Urine samples were collected according to standard

procedures (Kurien et al., 2004). For habituation-dishabituation tests, frozen urine samples

from different young subjects and cage mates were used to test fine odor discrimination. For

sensitivity tests, urine samples were combined in a stock sample used for each round of odor

presentations until completing the experiments. For female urine, samples obtained at

different points of the estrous cycle were pooled to generate a uniform stock.

Behavioral assessment

All the behavioral experiments using social and non-social stimuli were performed in a

dedicated room with continuous air reposition under dim indirect light (20 lx). Odor dilutions

were prepared in a room outside the animal house. Odor presentation was performed in a

homemade methacrylate box with removable walls for cleaning. The chamber dimensions were 15 cm width, 15 cm length and 30 cm high. A small hole (1cm diameter) was performed in the middle of one of the sides located 5 cm above from the box ground to fit standard cotton sticks impregnated with 1 μL of the odorant (**Fig. 3A**) allowing direct contact with the stick. Habituation to the testing conditions was performed before odorant presentations consisting of: handling (5 min), free exploratory activity in the box and habituation to the cotton stick movement (5 min). Odors were presented embedded in a cotton stick in order to preserve non-volatile pheromones and mimic the natural detection method of physical contact between conspecifics (Leypold et al., 2002, Stowers et al., 2002, Kimchi et al., 2007). To avoid variability due to the intensity of the volatile components of the urine, only direct nose contact with the tip of the cotton stick were considered as explorative behavior (sniffing/exploration time). Experiments were monitored by a video recording camera fixed 15 cm above the box. Videos were collected and analyzed off-line using SMART video-tracking software (PanLab S.L.).

a) ***Odor exploration test***: After a period of habituation (10 min), mice were exposed to two control trials (mineral oil for isoamyl acetate assays or water for urine tests) during 1 min separated by intervals of 1 min to avoid lack of novelty and odor habituation (Breton-Provencher et al., 2009, Sanderson and Bannerman, 2011). Subsequent presentations consisted on serial dilutions of either urine samples (diluted in water: 1:1.000, 1:500, 1:250, 1:100, 1:50, 1:10 and non-diluted, nd) or isoamyl acetate were tested in ascending order (diluted in mineral oil: $1:5\times10^5$, 1:100.000, 1:10.000; 1:1.000, 1:100). Animals were considered to detect the olfactory stimulus when spent more time investigating the odors than the vehicles (water or mineral oil).

b) ***Food finding test (FFT)***: Food finding test (FFT) was performed following standard procedures (Deacon et al., 2009). The mice were food deprived for 24 h before testing. Five food pellets (~ 35 g) were placed a corner and covered by 5 cm of litter bedding. Animals were

considered to detect the food pellet when spent digging, touching and holding the food pellet

for more than 5 s.

c) *Social odor habituation-dishabituation test*: The effect of aging in social odor

discrimination and habituation was explored using an adapted habituation-dishabituation test

(Gheusi et al., 2000). After a period of cage habituation (10 min), two control trials were

performed employing cotton sticks soaked with 1µl of water (vehicle). Three successive

presentations of urine from an animal of the opposite sex (S1a-c) were followed by three

presentations of urine from a different subject of the opposite-sex (S2a-c) to evaluate fine

chemo-olfactory discrimination and habituation. Urine samples came from young animals of

the opposite sex to maximize approaching behavior (Xu et al., 2005, Garratt et al., 2011). Each

sample presentation lasted 1 min separated by 1 min intervals. Habituation was estimated as

a decrease in the exploration time (sniffing) over the cotton stick tip after the first exposure of

urine from the same animal (S1a). Dishabituation (discrimination) occurs in response to a new

odor presentation (S2a) as a measurable increase in the exploration time. The test allows to

explore two consecutive phases of discrimination (H_2O^B-S1a and S1c-S2a) and habituation

(S1a-S1b and S2a-S2b). Trend lines between H_2O^B-S1a, S1c-S2a, S1a-S1b and S2a-S2b

were fitted to obtain the slope values indicating the amplitude of the discrimination and

habituation effect in each tested condition. Higher positive values indicated more pronounced

social discrimination and higher negative values corresponded to more robust habituation

(**Fig. 6**).

d) *Long-term social habituation-dishabituation test*: The aforementioned social odor

habituation task was adapted to assay social cue memory by presenting the urine sample

employed in S1a, 24 hours after the first test was performed. Intact social odor memory

manifested as a reduction in the sniffing time during the presentation of S1a during the second

trial, an effect which was clearly apparent in young animals (**Fig. 5F, G**).

e) *Three chamber test*: Social testing was performed in a cage (60 × 40 ×22 cm) following

standard procedures (Nadler et al., 2004). Dividing walls were made from clear Plexiglas, with

openings allowing access into each chamber. The test mouse was first placed in the middle chamber and allowed to explore for 10 minutes. After the habituation period, an unfamiliar subject of the same sex (mouse 1, M1), was placed in one of the side chambers. The stranger mouse was enclosed in a small, round wire cage, which allowed nose contact between the bars. In this first session (sociability), the test mouse had a choice of spending time in either the empty chamber (E) or the chamber occupied by M1. At the end of the sociability session, each mouse was tested in a second 10-minute session to evaluate social preference for a new subject. A second, unfamiliar mouse (mouse 2, M2) of the same sex was placed in the chamber that had been empty during the first session. This second stranger was enclosed in an identical wire cage than M1. The test mouse had a choice between the first, already-investigated mouse (M1), and the novel unfamiliar mouse (M2) which indicates their social preference or social novelty (Nadler et al., 2004). Continuous video recordings were collected and analyzed off-line using BORIS (Friard and Gamba, 2016) and SMART video-tracking software (PanLab S.L.). Measures of time spent sniffing E, M1, and M2 were quantified for each session.

Data analysis. All data were tested for statistical significance using GraphPad Prism 8. Shapiro-Wilk test was used to determine data normality. One-way ANOVA with Tukey´s test for multiple comparisons with a single variable was implemented for cell quantifications and anatomical data. For the behavioral analysis, a two-way ANOVA with Tukey´s test was used to test multiple comparisons with more than one variable. $P \leq 0.05$ was considered statistically significant. **Additional files 2, 3, 5, 6** summarize the P and N values corresponding to all the statistical comparisons between different ages, genotypes and dilutions of the social odor sensitivity tests.

RESULTS

Structural modifications of the mouse VSE during natural and pathological aging

We sought to investigate the structure of the VSE during natural and pathological aging, as the main gateway of pheromone (non-volatile)-encoded information (Sharrow et al., 2002, Cheetham et al., 2007, Ferrero et al., 2013). Stereological analysis revealed significantly smaller VSE volumes in 2 year-old (senescent) mice (**Fig. 1A**: VSE volume mm^3: 4 month-old: 0.190 ± 0.008; 1-year-old APP/PS1^{WT}: 0.170 ± 0.007; 24 month-old: 0.160 ± 0.007; 1-year-old APP/PS1^{Het}: 0.160 ± 0.008). This observation was supported by the reduced number of olfactory marker protein positive cells (OMP$^+$) along the anterior and medial axis of the marginal VSE (Halpern and Martínez-Marcos, 2003) (**Fig. 1C, E, Additional file 1**: number of OMP$^+$ cells x10^3 /μm^2: **Anterior**: 4 month-old: 10.69 ± 0.56; 12 month-old APP/PS1^{WT}: 9.50 ± 0.45; 24 month-old: 7.80 ± 0.53; 12 month-old APP/PS1^{Het}: 8.60 ± 0.53; **Medial**: 4 month-old: 9.92 ± 0.23; 12 month-old APP/PS1^{WT}: 9.15 ± 0.78; 24 month-old: 7.70 ± 0.70; 12 month-old APP/PS1^{Het}: 8.40 ± 0.40; **Posterior**: 4 month-old: 9.15 ± 0.70; 12 month-old APP/PS1^{WT}: 9.53 ± 0.77; 24 month-old: 7.90 ± 0.75; 12 month-old APP/PS1^{Het}: 10.00 ± 0.84).

We hypothesized that the drop in the number of OMP$^+$ cells could translate into a reduced axonal projection to the AOB, the main VNO target area, resulting in smaller AOB volumes. As such, we observed reduced AOB size in 2 year-old mice (**Fig. 1B:** AOB volume mm^3: 4 month-old: 0.48 ± 0.03; 12 month-old APP/PS1^{WT}: 0.48 ± 0.04; 24 month-old: 0.41 ± 0.03; 12 month-old APP/PS1^{Het}: 0.44 ± 0.03), indicating that aging may alter basic structural features of the VNO-AOB axis.

We expanded our analysis to the SRY-box transcription factor 2 (Sox2)-expressing cells (Sox2$^+$) in the supporting cell layer (SCL), a neuronal stem cell-marker which also stain mature differentiated sustentacular cells (Guo et al., 2010, Taroc et al., 2020, Katreddi et al., 2021). The estimation of the labeling intensity of Sox2 in the SCL showed a drastic reduction in senescent but not in APP/PS1^{Het} mice (**Fig. 1D, F**: Sox2 fluorescence intensity / area (μm^2): 4

month-old: 33.24 ± 2.43; 12 month-old: 26.23 ± 3.95; 24 month-old: 10.74 ± 1.32; 12 month-old APP/PS1^{Het}: 27.37 ± 0.96), indicating natural aging may disrupt VSE structure by reducing the number of sensory neurons and sustentacular cells (OMP$^+$ and Sox2$^+$ cells, respectively).

Natural and pathological aging differentially impact VSE cell proliferation

The observed decrease in the number of mature OMP$^+$ and Sox2$^+$ cells in the VSE of senescent mice could be due to the reduction of the VSE´s regenerative ability. Given that there is currently scarce data on the properties of the VSE neurogenic niche in neurodegenerative animal models, we analyzed the number of proliferative cells identified by the expression of proliferative cell nuclear antigen (PCNA) in the anterior, medial and posterior VNO in senescent and APP/PS1^{Het} mice. Consistent with previous reports cell proliferation in young animals was abundant in the marginal zone of the anterior and medial VSE (Giacobini et al., 2000, Brann and Firestein 2010, **Fig. 2A, B**). According to previous reports (Brann and Firestein, 2010), senescent mice exhibited a significant reduction in PCNA$^+$ cells in the marginal VSE (**Fig. 2A, B**, number of PCNA$^+$ cells x 10^3/mm^3: **Anterior**: 4 month-old: 2.26 ± 0.80; 12 month-old APP/PS1^{WT}: 0.73 ± 0.3; 24 month-old: 0.75 ± 0.35; 12 month-old APP/PS1^{Het}: 3.42 ± 0.82; **Medial**: 4 month-old: 2.15 ± 0.52; 12 month-old APP/PS1^{WT}: 1.61 ± 0.32; 24 month-old: 0.23 ± 0.13; 12 month-old APP/PS1^{Het}: 1.57 ± 0.20; **Posterior**: 4 month-old: 3.04 ± 0.90; 12 month-old APP^{WT}: 2.42 ± 0.60; 24 month-old: 0.17± 0.08; 12 month-old APP/PS1^{Het}: 1.93 ± 0.57), indicating reduced proliferation. Surprisingly, the number of PCNA$^+$ cells in APP/PS1^{Het} mice was significantly higher that of aged-matched controls in the anterior VSE (**Fig. 2A, B**), suggesting a region-specific increase in cell generation in these animals. Importantly, no significant overlap between OMP and PCNA staining was observed indicating that these two markers recognize cell populations at different maturation stages (**Fig. 2A,** right panel).

We then sought to explore whether the reduced number of proliferative PCNA$^+$ cells in senescent mice could be due to a decrease of stem cell generation by analyzing the number of Sox2$^+$ neural precursor cells in the VSE (Tucker et al., 2010, Panaliappan et al., 2018). We

observed a significant reduction in the number of Sox2[+] cells in 2 year-old mice but not in APP/PS1[Het] animals, indicating a reduced ability to progenitor generation in the aged VSE (**Fig. 2C**: number of Sox2[+] cells in the VSE, number of cells x 10[4] cells/μm[2]: 4 month-old: 3.08 ± 0.40; 12 month-old APP/PS1[WT]: 2.92 ± 0.38; 24 month-old: 2.37 ± 0.36; 12 month-old APP/PS1[Het]: 2.15 ± 0.24). Altogether, these findings revealed a fundamental difference in how natural and diseased aging impact the VSE structure and proliferative capacity.

Late onset of social exploration deficits during natural aging

Our data indicated that natural aging induce significant alterations in the structure, VSE proliferative capacity and cellular composition, thus we investigated whether these adaptations might translate into impairments in the processing of socio-sexual information. Cumulative evidence points to olfactory decline as a common symptom of natural and pathological aging (Doty et al., 1984, Murphy et al., 2002, Rawson et al., 2012, Doty and Kamath, 2014, Roberts et al., 2016). However, most of these studies and diagnostic tests employed synthetic odors with reduced emotional valence, therefore, the temporal course and severity of the age-related involution of the recognition of social cues, largely processed by the VNO-AOB system, remains unclear.

To explore the impact of natural aging on social odor detection, we conducted an odor-evoked sniffing test in which serial dilutions of urine from young conspecifics of the opposite sex were presented to either male or female subjects over a range of different ages: young (2-4 mo.), adult (6-8 mo.), middle-aged (12-14 mo.) and old/senescent mice (20-24 mo.) (**Fig. 3A**, see Material and Methods for details on urine sample preparation). Our results indicated a significant reduction in the exploration time throughout the various urine dilutions in comparison to adult wild type mice (**Fig. 3B, C** sniffing time normalized to water control 2, **undiluted sample**: aged: 3.83 ± 0.71; adult: 6.06 ± 0.88; **1:10 dilution**: aged: 3.04 ± 0.50; adult: 4.90 ± 1.10; **1:50 dilution**: aged: 1.90 ± 0.33; adult: 3.14 ± 0.55; **1:100 dilution**: aged: 1.56 ± 0.37; adult: 2.95 ± 0.43; **1:250 dilution**: aged: 1.37 ± 0.36; adult: 2.36 ± 0.70; **1:500 dilution**: aged: 0.86 ± 0.20; adult: 2.50 ± 0.35; **1:1000 dilution**: aged: 0.60 ± 0.20; adult: 2.80

± 0.43). Raw sniffing time data showed a similar decrease in the exploration of social odors in

senescent animals across different dilutions (see details in **Additional files 2, 3**).

Furthermore, senescent mice progressively increased the exploration time in parallel with odor

concentration, whereas young, adult and middle-age animals showed a habituation phase at

intermediate dilutions (1:500, 1:250, 1:100 and 1:50), indicating effective odor detection and

recognition capabilities (Yang and Crawley, 2009). Lastly, data analysis disaggregated by sex

showed no sex-differences in the reduction of urine exploration time (**Additional file 4**; but

see Martínez et al., 2017 for an alternative view).

Reduction in the exploration of social information is accelerated in an animal model of

neurodegeneration

Next, we investigated whether pathological aging would alter the exploration of social olfactory

cues despite no obvious effects on VSE structure or proliferative capacity. Social odor

sensitivity tests in 1 year-old APP/PS1^{Het} mice revealed reduced sniffing times when compared

to age-matched APP/PS1^{WT} control mice for low urine dilutions, suggesting that pathological

aging accelerates the decline of exploration time of social odors (**Fig. 3D, E** sniffing time

normalized to water control 2, **undiluted sample**: 1 year-old APP/PS1^{Het}: 2.58 ± 0.43; 1 year-

old APP/PS1^{WT}: 5.65 ± 1.51; **1:10 dilution**: 1 year-old APP/PS1^{Het}: 1.82 ± 0.27; 1 year-old

APP/PS1^{WT}: 4.94 ± 1.21; **1:50 dilution**: 1 year-old APP/PS1^{Het}: 1.43 ± 0.46; 1 year-old

APP/PS1^{WT}: 1.40 ± 0.26; **1:100 dilution**: 1 year-old APP/PS1^{Het}: 1.37 ± 0.43; 1 year-old

APP/PS1^{WT}: 2.03 ± 0.55; **1:250 dilution**: 1 year-old APP/PS1^{Het}: 0.90 ± 0.17; 1 year-old

APP/PS1^{WT}: 0.70 ± 0.14; **1:500 dilution**: 1 year-old APP/PS1^{Het}: 0.52 ± 0.10; 1 year-old

APP/PS1^{WT}: 0.97 ± 0.20; **1:1000 dilution**: 1 year-old APP/PS1^{Het}: 1.00 ± 0.20; 1 year-old

APP/PS1^{WT}: 0.84 ± 0.31). Similar to senescent mice, these results were reproduced when raw

sniffing time data were compared (**Additional file 5, 6**).

Aging and neurodegeneration impact social odor recognition more severely than other odor modalities

Next, we asked whether non-social odor modalities were also affected in senescent and APP/PS1^{Het} animals. We investigated this question by analyzing the exploration time to both food and synthetic neutral odors. First, we exposed naturally aged and APP/PS1^{Het} mice to serial dilutions of IA, a synthetic banana like odor of neutral valence when used at high dilutions (Fortes-Marco, 2015, Saraiva et al., 2016). Consistent with previous studies, mice of all conditions showed significantly reduced responses to the neutral odorant than to urine (**Fig. 4A-D**) according to the higher emotional valence of urine in comparison to a synthetic odor (Kobayakawa et al. 2007, Mandairon et al. 2009, Saraiva et al. 2016, Jagetia et al. 2018). Our results showed no significant differences on the exploration times of IA in senescent mice and a modest but significant increase in the sniffing time of the 1:10^4 and 1:100 IA dilutions in APP/PS1^{Het} animals (**Figures 4A-D,** normalized sniffing time, **1:100 dilution**: 1 year-old APP/PS1^{Het}: 1.40 ± 0.12; 1 year-old APP/PS1^{WT}: 0.93 ± 0.16; young: 1.76 ± 0.26; aged: 1.24 ± 0.20; 1:1000 dilution: 1 year-old APP/PS1^{Het}: 1.40 ± 0.22; 1 year-old APP/PS1^{WT}: 1.42 ± 0.30; young: 1.90 ± 0.35; aged: 1.56 ± 0.50; **1:10^4 dilution**: 1 year-old APP/PS1^{Het}: 1.50 ± 0.25; 1 year-old APP/PS1^{WT}: 0.91 ± 0.13; young: 1.44 ± 0.32; aged: 1.20 ± 0.26; **1:10^5 dilution**: 1 year-old APP/PS1^{Het}: 0.75 ± 0.11; 1 year-old APP/PS1^{WT}: 0.75 ± 0.15; young: 1.41 ± 0.50; aged: 0.90 ± 0.20; **1:5x10^5 dilution**: 1 year-old APP/PS1^{Het}: 0.85 ± 0.18; 1 year-old APP/PS1^{WT}: 0.82 ± 0.11; young: 1.65 ± 0.42; aged: 1.23 ± 0.35). Similarly, the FFT showed no significant differences in the latency to find food pellets after 24 hours of food deprivation in either naturally aged or middle age APP/PS1^{Het} mice (**Fig. 4E** latency of finding food pellet (min), young: 5.22 ± 1.08; aged: 5.15 ± 0.95; 1-year-old APP/PS1^{Het}: 4.27 ± 0.58; 1-year-old APP/PS1^{WT}: 3.87 ± 0.32). These results suggest that deficits in social exploration time at advanced stages of natural aging and in an animal model of AD are more pronounced than other odor modalities.

Social odor discrimination and habituation are reduced in naturally aged and APP/PS1^{Het} mice

To further investigate the impact of aging and neurodegeneration in the detection of social information, we performed a habituation – dishabituation test, which relies on the animal's ability to discriminate novel smells (Yang and Crawley, 2009). For these experiments, young (2-4 mo.) and aged (20-24 mo.) wild type animals were presented three consecutive replicates of urine samples from two different animals (S1a-c and S2a-c) (**Fig. 5A**). Aged animals were able to discriminate between urine sources, but showed reduced sniffing times during both the first and second discrimination and habituation phases (**Fig. 5B, C**). Similarly, middle age APP/PS1^{Het} mice also exhibited reduced sniffing times in comparison to aged-matched APP/PS1^{WT} controls (**Fig. 5D, E** sniffing time in seconds, **water control 1**: aged: 0.56 ± 0.09; young: 0.42 ± 0.05; APP/PS1^{WT}: 0.72 ± 0.08; APP/PS1^{Het}: 0.57 ± 0.06; **water control 2**: aged: 0.56 ± 0.11; young: 0.67 ± 0.10; APP/PS1^{WT}: 0.50 ± 0.07; APP/PS1^{Het}: 0.71 ± 0.09; **S1a**: aged: 1.36 ± 0.27; young: 2.45 ± 0.40; APP/PS1^{WT}: 2.81 ± 0.40; APP/PS1^{Het}: 1.75 ± 0.22; **S1b**: aged: 0.70 ± 0.14; young: 0.87 ± 0.20; APP/PS1^{WT}: 0.95 ± 0.10; APP/PS1^{Het}: 0.68 ± 0.10; **S1c**: aged: 0.70 ± 0.21; young: 0.51 ± 0.10; APP/PS1^{WT}: 0.84 ± 0.22; APP/PS1^{Het}: 0.71 ± 0.13; **S2a**: aged: 1.51 ± 0.27; young: 2.44 ± 0.43; APP/PS1^{WT}: 1.85 ± 0.42; APP/PS1^{Het}: 1.00 ± 0.17; **S2b**: aged: 0.82 ± 0.14; young: 0.81 ± 0.15; APP/PS1^{WT}: 0.80 ± 0.22; APP/PS1^{Het}: 0.62 ± 0.08; **S2c**: aged: 0.96 ± 0.18; young: 0.55 ± 0.11); APP/PS1^{WT}: 0.60 ± 0.13; APP/PS1^{Het}: 0.50 ± 0.09). Analysis of the slope values of the first and second discrimination and habituation phases revealed that senescent mice exhibited significant deficits during both rounds of social habituation – dishabituation, and APP/PS1^{Het} mice only during the second phase of discrimination (**Fig. 6A, B**).

Then, we adapted the habituation-dishabituation test to assay potential changes in long-term social odor memory (**Fig. 5F**). To this aim, animals were exposed to the same S1a urine sample 24 hours after the first presentation. A reduction of the sniffing time during the first discrimination phase was interpreted as an indicator of memory. This reduction was clearly

observed in young animals, but absent in naturally aged and 1-year-old APP/PS1[Het] mice (**Fig. 5G:** sniffing time of S1a after 24 hours in seconds, young: 1.20 ± 0.20; aged: 1.12 ± 0.20; APP/PS1[WT]: 1.40 ± 0.50; APP/PS1[Het]: 1.20 ± 0.33), suggesting long-term social odor memory impairments during both natural and pathological aging.

Age-related deficits in social discrimination and habituation are not influenced by previous experience

Next, we compared animals exposed to either familiar (littermate urine, L) or novel social odors (novel urine, N) in the habituation-dishabituation test (**Fig. 7A**). Aged animals performed poorly in the L-N dishabituation task (no significant increase in sniffing time), suggesting that the reduction in social odor discrimination and habituation is independent of previous experience.

Since impairments in social odor discrimination were more significant in naturally aged mice, we asked whether these deficits also extended to the recognition of animal's own odors, as the loss of self-awareness is a disrupting symptom of common occurrence in senescent subjects (La Joie et al., 2016, Valech et al., 2018). To this aim, animals were presented to samples of their own urine (O) during the second discrimination phase of the habituation-dishabituation test (**Fig. 7C, D**). Our data indicated that animal´s own urine was effective to elicit a typical discrimination (dishabituation) response, (**Fig. 7C, D**: sniffing time in seconds, **water control 1**: aged: 0.70 ± 0.10; young: 0.81 ± 0.10; **water control 2**: aged: 0.31 ± 0.09; young: 0.48 ± 0.11; **La**: aged: 2.07 ± 0.43; young: 3.06 ± 0.35; **Lb**: aged: 0.64 ± 0.16; young: 0.74 ± 0.17; **Lc**: aged: 0.48 ± 0.10; young: 0.45 ± 0.10; **Oa**: aged: 1.23 ± 0.18; young: 1.22 ± 0.33; **Ob**: aged: 0.48 ± 0.13; young: 0.84 ± 0.13; **Oc**: aged: 0.49 ± 0.11; young: 0.55 ± 0.15), suggesting that self-recognition is overall preserved in senescent mice.

Social novelty is disrupted during pathological aging

Last, we explored whether the observed impairments in social odor exploration and discrimination may negatively impact social interaction in naturally aged and APP/PS1[Het] mice. To this aim, we performed a three-chamber test (Nadler et al., 2004, Yang et al., 2011; **Fig.**

8A) to assess general sociability and social novelty in senescent and APP/PS1[Het] animals. Our data indicated a reduction in social novelty in middle age APP1/PS1[Het] mice, which was not detected in 2 year-old animals (**Fig. 8B-E**, **Sociability**, sniffing time in seconds, **E**: aged: 14.50 ± 1.90; young: 25.06 ± 2.96; APP/PS1[WT]: 39.15 ± 3.63; APP/PS1[Het]: 41.00 ± 4.07; **M1[A]**: aged: 47.24 ± 4.60; young: 64.35 ± 5.23; APP/PS1[WT]: 69.40 ± 6.05; APP/PS1[Het]: 69.73 ± 7.10; **Social novelty**, sniffing time in seconds, **M1[B]**: aged: 49.15 ± 4.20; young: 62.37 ± 5.01; APP/PS1[WT]: 71.30 ± 5.75; APP/PS1[Het]: 65.12 ± 6.70; **M2**: aged: 63.70 ± 5.30; young: 99.93 ± 9.83; APP/PS1[WT]: 93.07 ± 11.70; APP/PS1[Het]: 77.02 ± 8.43). Although senescent mice did not show significant impairments in either sociability or social novelty, they exhibited a mild increase in the latency to approach M1 during the sociability phase (**Additional file 7**), consistent with the overall decrease in the exploration time of social odors **(Additional files 2, 3)**. These findings exposed exacerbated deficits in an animal model of AD, suggesting a distinct impact of natural and pathological aging on social behavior.

DISCUSSION

The quality of social life has been proposed as a predictive factor for developing dementia or mental illness (Cohen et al., 1985, Stockdale et al., 2007, Wilson et al., 2007). However, the impairment of social functions with age is poorly understood with some authors suggesting that social deficits might be the consequence of generalized brain impairments (Johansson et al., 2018, Moran et al., 2012, Cavallini et al., 2013) or a symptom which might be developed independently of cognitive deficits (Spalletta et al., 2004, Mohs et al., 2000).

Our results revealed that both natural and pathological aging affect several key aspects of social information processing in mice including the exploration of social odors, social odor discrimination and habituation. To gain insight into the mechanisms underlying these deficits, we explored the age-related adaptions of the VSE, a central gateway for pheromone-encoded information in mammals (Jacobson, 1813, Halpern and Martínez-Marcos, 2003, Munger et al., 2009) and part of the accessory olfactory system whose aging process has been largely overlooked in contrast to other aspects of olfaction (Murphy, 2019).

Our data showed VSE alterations during natural aging but not in APP/PS1[Het] mice, a

standardized animal model of AD. Whereas the VSE of APP/PS1[Het] mice maintained fairly

stable neurogenic capabilities, senescent wild type animals showed a reduction in proliferative

and stem cells (PCNA[+] and Sox2[+] cells) in the marginal VSE (**Fig. 2**) which in turn may explain

the reduction of mature OMP[+] neurons, sustentacular Sox2[+] cells and the organ volume (**Fig.**

1). Our findings are consistent with two previous studies (Brann and Firestein, 2010, Mechin

et al., 2021), which reported an overall thinning of the vomeronasal sensory epithelium

(Mechin et al., 2021) and reduced VSE neurogenesis in the marginal zone of 2-year-old

animals (Brann and Firestein, 2010). Here, we expand the current knowledge with novel

information from a commonly used AD animal model (APP/PS1[Het]), revealing an unexpected

preservation of the VSE proliferative capabilities in middle age APP/PS1[Het] mice. These results

contrast with the reduction of SVZ neurogenesis reported in similar animal models (Verret et

al., 2007, Zhang et al., 2007, Zeng et al., 2016, Scopa et al., 2020), suggesting that VSE

neurogenesis might be less susceptible to lesions (Brann and Firestein, 2010) and

pathological conditions than other proliferative areas. However, we cannot rule out the

possibility that the VSE of APP/PS1[Het] animals may develop potential alterations at a later age

(> 1 year-old). An aspect that we were prevented to investigate due to the short life expectancy

(~ 14-16 months-old) of this mouse model.

A relevant aspect of our work is that most previous studies addressing olfactory decline

employed synthetic or neutral odors, thus remaining uncertain the effect of healthy and

diseased aging in the recognition of social cues. To gain insight into this question, we

investigated the exploration time and the habituation-dishabituation response to social odors

(urine) (**Figs. 3, 5, 6**). Our findings revealed that despite the distinctive effects of natural and

pathological aging on the VSE, both processes impaired the exploration of socio-sexual cues,

social discrimination-habituation, and social behavior, suggesting fundamental differences in

the mechanisms by which healthy and diseased aging impact social information processing.

Further experiments are needed to establish causality, but our results suggest that the

observed VSE alterations in senescent mice underlie deficits in urine exploration time (**Fig. 3,**

Additional files 2, 3) which could impair the processing of social information. VSE volume

data from middle age APP/PS1^{WT} control animals suggest that VSE structural changes might

appear around 1 year-old of age (**Fig. 1A**: VES volume mm^3: 4 month-old: 0.190 ± 0.008; 1-

year-old APP/PS1^{WT}: 0.170 ± 0.007; $p = 0.03$; two-way ANOVA with Tuckey´s test), although

their functional consequences might not become apparent until advanced stages of aging (2-

year-old). This scenario suggests a parsimonious VNO decay that matches the aging of other

olfactory areas like the OB. Several studies have shown that despite the reduced regeneration

rate of olfactory sensory neurons (OSNs), OB alterations involving a reduced number of

synaptic contacts (Richard et al., 2010), expression of odorant receptor genes (Lee et al.,

2009; Khan et al., 2013) and changes in the OSNs dynamic range (Kass et al., 2018) just

become apparent in 2-year-old mice. This evidence suggests that although age-related

changes may start to appear earlier (Mobley et al., 2014), functional deficits might exhibit a

late onset, suggesting compensatory mechanisms to preserve the processing of olfactory

cues relevant for the survival of aged animals (Tobiansky et al., 2012, Tikhonova et al., 2015).

In contrast to natural steady decline, pathological conditions may accelerate functional deficits

even in the absence of detectable modifications of peripheral organs, suggesting impairments

in the central processing of social information. As such, defects in the exploration of social

odors were found exacerbated in middle age APP/PS1^{Het} mice (**Fig. 3D, E**), a condition likely

to aggravate frailty and reduce life expectancy in these animals (as observed for

APP/PSEN1^{Het} in our experimental conditions).

Furthermore, a detailed analysis of the first discrimination phase during the modified social

habituation-dishabituation test (presentation of S1a urine sample after 24 hours; **Fig. 5F, G**)

revealed significant impairments in both 2-year old wild type animals and middle age

APP/PS1^{Het} mice, suggesting that in parallel to peripheral sensory decline, the downstream

pathways involved in social cue recognition may be affected during both natural and

pathological aging. To control for a potential contribution of novelty in the social habituation-dishabituation test, we presented urine samples from littermate and strange animals finding similar deficits in discrimination (**Fig. 7**). Interestingly, although senescent animals showed severe discrimination deficits they preserved the ability to differentiate between urine from a novel subject and their own, preventing to explore further the underlying mechanisms of subjective perception loss, a disrupting symptom of senescence and dementia which currently lacks appropriate animal models for preclinical studies (Jessen et al., 2020).

Age-related decline of olfactory detection is likely to occur due to alterations of both the main olfactory epithelium and the VNO (Apfelbach et al., 1991, Lee et al., 2009, Ueha et al., 2018), thus potentially affecting other odor modalities. Here, we sought to explore this question by performing odor-evoked sniffing tests using social and neutral synthetic odors (**Fig. 4**). Quantification of the exploration time across various IA dilutions exposed equivalent measurements in senescent and young mice, and even enhanced responses in APP/PS1[Het] mice. Furthermore, results from a FFT, revealed no significant differences in the latency time to find the hidden food pellets (**Fig. 4E**) indicating that defects in social odor detection related to natural and pathological aging may be more severe than other odor modalities. A plausible explanation for the specific impairment in the exploration of social odors may involve the reduction of reproductive activity in old animals, but not in foraging and feeding behaviors (Harb et al., 2014). Furthermore, the decline to detect specific odors may rise from alterations on the number of sensory neurons and synaptic connections specifically related to the discrimination of conspecifics (Bielsky et al., 2005, Wacker et al., 2010, Tobin et al., 2010) and heterospecific social cues (e.g. predator) (Kang et al., 2009, Wacker et al., 2010). In particular, impairments in social odor exploration and discrimination may rise from maladaptations of the VNO-AOB axis, consistent with the observed decrease in AOB volume (**Fig. 1B**). Future studies should deepen into the anatomical and functional properties of the VNO-AOB circuit in order to obtain a complete picture on how healthy and pathological aging affect social information processing.

Finally, we sought to determine how the age-related defects in social odor sensitivity, discrimination and memory shaped social behavior. Results from a three chamber test (**Fig. 8**) indicated that sociability was overall preserved in senescent and APP/PS1[Het] animals. In contrast, social novelty was severely impaired in APP/PS1[Het] mice consistently with previous studies (Cheng et al., 2014, Olesen et al., 2016, Shoji and Miyakawa, 2019, Locci et al., 2021). Deficits in social novelty in APP/PS1[Het] mice could be linked to the deficits in social discrimination observed in the habituation-dishabituation test (**Fig. 5D, E** and **Fig. 6D-F**) which could impair the recognition of M2 as a novel subject (Enwere et al., 2004, Patel and Larson, 2009, Moreno et al., 2014). Consistent with the overall decrease in the exploration time of social odors **(Fig. 3, Additional files 2, 3)**, aged animals exhibited a mild increase in the latency to approach the novel mouse (M1) during the first sociability phase (**Additional file 7**). These findings indicate that despite the reduction in social odor exploration and discrimination, overall sociability and novelty are majorly preserved in naturally aged mice. In contrast, the APP/PS1 neurodegenerative model shows a measurable impairment in social preference. This evidence opens the intriguing possibility that neuronal circuits underlying specific social functions (sociability vs social memory) may be particularly susceptible to pathological aging.

CONCLUSION

Olfactory deficits are a common symptom of natural and pathological aging. While multiple aged-related changes of the olfactory sensory epithelium are described, the aging of the pheromone detection system, a major gateway for social information, has been largely overlooked. This study reveals that whereas natural aging elicits a decline in VSE cell proliferation, likely to reduce the number of mature sensory neurons and the overall volume of the organ, a common animal model of AD exhibits normal proliferation capacities and no obvious morphological alterations. Despite their distinctive effects at the cellular level, both natural and pathological aging disrupt the detection of social odors in a more severe way than other odor modalities. Furthermore, social detection and social behavior impairments were

579 exacerbated in APP/PS1^{Het} mice, indicating pathological aging may impact the downstream

580 processing of social information even in the absence of obvious alterations of the peripheral

581 organs.

583 **DECLARATIONS**

584 **Ethics approval**

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by grants of the Spanish Ministry of Science and Innovation

SAF2017-82524-R and PID2020-113878RB-I00 (to S.J.), the "Severo Ochoa" Program for

Centres of Excellence in R&D (SEV-2013-0317 and SEV-2017-0723), the Generalitat

Valenciana Prometeo/2019/014 (to S.J.), and a FPI contract (BES-2017-081243) to A.P;

Agence National de la Recherche (ANR): ANR-20-CE92-0003 (to P.C.) and Region Centre

Val de Loire: 201900134883 (to P.C.).

Author contributions

The project was led by S.J. Brain sample preparation, data acquisition, and analysis was done

by A.P. Result interpretation, manuscript preparation, and editing was done by S.J., P.C. and

A.P. with help of members from the Jurado Lab at the Instituto de Neurociencias (CSIC-UMH).

Acknowledgements

We are grateful to the teams of the Imaging and Animal Facilities at the Institute of

Neuroscience, and all members of the Jurado and Chamero laboratories for their support

during the realization of this work.

Author information

[1]Institute of Neuroscience CSIC-UMH, San Juan de Alicante, 03550, Alicante, Spain.[2]

Laboratoire de Physiologie de la Reproduction et des Comportements, CNRS, IFCE, INRAE,

University of Tours, 37380, Nouzilly, France.

REFERENCES

Adolphs R. The neurobiology of social cognition. Curr. Opin. Neurobiol. 2001;11:231-9.

Ahlenius H, Visan V, Kokaia M, Lindvall O, Koaia Z. Neuronal stem and progenitor cells retain

their potential for proliferation and differentiation into functional neurons despite lower number

in aged brain. J. Neurosci. 2009;29:4408-19.

Apfelbach R, Russ D, Slotnick BM. Ontogenic changes in odor sensitivity, olfactory receptor

area and olfactroy receptor density in the rat. Chem. Senses. 1991;16:209-18.

Baum MJ, Kelliher KR. Complementary roles of the main and accessory olfactory systems in

mammalian mate recognition. Annu. Rev. Physiol. 2009;71:141-60.

Beynon RJ, Hurst JL. Urinary proteins and the modulation of chemical scents in mice and rats.

Peptides. 2004;25:1553-63.

Bielsky IF, Hu S-B, Ren X, Terwilliger EF, Young LJ. The V1a vasopressin receptor is necessary and sufficient for normal social recognition: a gene replacement study. Neuron. 2005;47:503-13.

Brann JH, Firestein SJ. Regeneration of new neurons is preserved in aged vomeronasal epithelia. J. Neurosci. 2010;30:15686-94.

Brennan PA, Kendrick KM. Mammalian social odours: attraction and individual recognition. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 2006;361:2061-78.

Brennan PA, Zufall F. Pheromonal communication in vertebrates. Nature. 2006;444: 308-15.

Breton-Provencher V, Lemasson M, Peralta MR 3rd, Saghatelyan A. Interneurons produced in adulthood are required for the normal functioning of the olfactory bulb network and for the execution of selected olfactory behaviors. J Neurosci. 2009;29:15245-257.

Cavallini E, Lecce S, Bottiroli S, Palladino P, Pagnin A. Beyond false belief: theory of mind in young, young-old and old-old adults. Int. J. Aging Hum. Dev. 2013;76:181-98.

Chamero P, Leinders-Zufall T, Zufall F. From genes to social communication: molecular sensing by the vomeronasal organ. Trends Neurosci. 2012;35:597-606.

Chamero P, Weiss J, Alonso MT, Rodríguez-Prados M, Hisatsune C, Mikoshiba K, Leinders-Zufall T, Zufall F. Type 3 inositol 1,4,5-trisphosphate receptor is dispensable for sensory activation of the mammalian vomeronasal organ. Sci. Rep. 2017;7:10260.

Cheetham SA, Thom MD, Jury F, Ollier WER, Beynon RJ, Hurst JL. The genetic basis of individual-recognition signals in the mouse. Curr. Biol. 2007;17:177-77.

Cheng D, Spiro AS, Jenner AM, Garner B, Karl T. Long-term cannabidiol treatment prevents the development of social recognition memory deficits in Alzheimer's disease transgenic mice. J. Alzheimers Dis. 2014;42:1383-96.

Cohen S, Wills TA. Stress, social Support, and the buffering hypothesis. Psychological Bulletin. 1985;98:310-57.

Deacon RMJ, Koros E, Bornemann KD, Rawlins JNP. Aged Tg2576 mice are impaired on

social memory and open field habituation tests. Behav. Brain Res. 2009;197:466-68.

Devanand DP. Olfactory identification deficits, cognitive decline and dementia in older adults.

Am. J. Geriatr. Psychiatry. 2016;24:1151-57.

Donovan NJ, Okereke OI, Vannini P, Amariglio RE, Rentz DM, Marshall GA, Johnson KA,

Sperling RA. Association of higher cortical amyloid burden with loneliness in cognitively normal

older adults. JAMA Psychiatry. 2016;73:1230-37.

Dorph-Petersen KA, Lewis DA. Stereological approaches to identifying neuropathology in

psychosis. Biol. Psychiatry. 2011;69:113-26.

Doty RL, Kamath V. The influences of age on olfaction: a review. Front. Psychol. 2014;5.20

doi: 10.3389/fpsyg.2014.00020.

Doty RL, Shaman P, Applebaum SL, Giberson R, Siksorski L, Rosenberg L. Smell

identification ability: changes with age. Science. 1984;226:1441-43.

Ennis M, Puche AC, Holy T, Shipley MT. The olfactory system. In: the rat nervous system:

Fourth edition, Elsevier Inc; 2015. p. 761-803.

Enwere E, Shingo T, Gregg C, Fujikawa H, Ohta S, Weiss S. Aging results in reduced

epidermal growth factor receptor signaling, diminished olfactory neurogenesis, and deficits in

fine olfactory discrimination. J. Neurosci. 2004;24:8354-65.

Ferrero DM, Moeller LM, Osakada T, Horio N, Li Q, Roy DS, Cichy A, Spehr M, Touhara K,

Liberles SD. A juvenile mouse pheromone inhibits sexual behaviour through the vomeronasal

system. Nature. 2013;502:368-71.

Fortes-Marco L, Lanuza E, Martinez-Garcia F, Agustín-Pavón C. Avoidance and contextual

learning induced by a kairomone, a pheromone and a common odorant in female CD1 mice.

Front. Neurosci. 2015;9:336.

Friard O, Gamba M. BORIS: A free, versatile open-source event-logging software for

video/audio coding and live observations. Methods Ecol. Evol. 2016;7:1325-30.

Garratt M, Stockley P, Armstrong SD, Beynon RJ, Hurst JL. The scent of senescence: sexual

signaling and female preference in house mice. J. Evol. Biol. 2011;11:2398-2409.

Gheusi G, Cremer H, McLean H, Chazal G, Vincent JD, Lledo PM. Importance of newly

generated neurons in the adult OB for odor discrimination. Proc. Natl. Acad. Sci. U S A.

2000;97:1823-28.

Giacobini P, Benedetto A, Tirindelli R, Fasolo A. Proliferation and migration of receptor

neurons in the vomeronasal organ of the adult mouse. Brain Res. Dev. Brain Res.

2000;123:33-40.

Graziadei PP, Monti-Graziadei AG. Regeneration in the olfactory system of vertebrates. Am.

J. Otolaryngol. 1983;4:228-33.

Guo Z, Packard A, Krolewski RC, Harris MT, Manglapus GL, Schwob JE. Expression of pax6

and sox2 in adult olfactory epithelium. J. Comp. Neurol. 2010;18:4395-418.

Halpern M, Martínez-Marcos A. Structure and function of the vomeronasal system: an update.

Prog. Neurobiol. 2003;70:245-318.

Harb MR, Sousa N, Zihl J, Almeida OFX. Reward components of feeding behavior are

preserved during mouse aging. Frontiers in Aging Neurosci. 2014;6:242.

Heinrichs M, Baumgartner T, Kirschbaum C, Ehlert U. Social support and oxytocin interact to

suppress cortisol and subjective responses to psychosocial stress. Biol. Psychiatry.

2013;54:1389-98.

Hlinák Z, Krejcí I. Social recognition in male rats: age differences and modulation by MIF-I and

alaptide. Physiol. Res. 1991;40:59-67.

Jacobson L. Anatomisk beskrivelse over et nyt organ I huusdyrenes naese. Veter Salesk Skr.

1813;2:209-46.

Jagetia S, Milton AJ, Stetzik LA, Liu S, Pai K, Arakawa K, Mandairon N, Wesson DW. Inter-

and intra-mouse variability in odor preferences revealed in an olfactory multiple-choice test.

Behav. Neurosci. 2018;132:88-98.

Jankowsky JL, Fadale DJ, Anderson J, Xu GM, Gonzales V, Jenkins NA, Copeland NG, Lee

MK, Younkin LH, Wagner SL, Younkin SG, Borchelt DR. Mutant presenilins specifically

elevate the levels of the 42 residue beta-amyloid peptide in vivo: evidence for augmentation

of a 42-specific gamma secretase. Hum. Mol. Genet. 2004;13:159-170.

Jankowsky JL, Slunt HH, Ratovitski T, Jenkins NA, Copeland NG, Borchelt DR. Co-expression

of multiple transgenes in mouse CNS: a comparison of strategies. Biomol. Eng. 2001;17:157-

65.

Jessen F, Amariglio RE, Buckley RF, van der Flier WM, Han Y, Molinuevo JL, Rabin, L, Rentz

DM, Rodríguez-Gómez O, Saykin AJ, Sikkes SAM, Smart CM, Wolfsgruber S, Wagner M. The

characterisation of subjective cognitive decline. Lancet Neurol. 2020;19:271-78.

Johansson NE, Murray K, Happé F, Charlton RA. Cognitive and affective associations with an

ecologically valid test of theory of mind across the lifespan. Neuropsychology. 2018;32:754-

63.

Kang N, Baum MJ, Cherry JA. A direct main olfactory bulb projection to the 'vomeronasal'

amygdala in female mice selectively responds to volatile pheromones from males. Eur. J.

Neurosci. 2009;29:624-34.

Kass MD, Czarnecki LA, McGann JP. Stable olfactory sensory neuron in vivo physiology

during normal aging. Neurobiol. Aging. 2018;69:33-37.

Katreddi RR, Forni PE. Mechanisms underlying pre- and postnatal development of the

vomeronasal organ. Cell Mol. Life Sci. 20121;78:5069-82.

Khan M, Vaes E, Mombaerts P. Temporal patterns of odorant receptor gene expression in
adult and aged mice. Mol. Cell Neurosci. 2013; 57:120-29.

Kobayakawa K, Kobayakawa R, Matsumoto H, Oka Y, Imai T, Ikawa M, Okabe M, Ikeda T,
Itohara S, Kikusui T, Mori K, Sakano H. Innate versus learned odour processing in the mouse
olfactory bulb. Nature. 2007;450: 503-8.

Kurien BT, Everds NE, Scofield RH. Experimental animal urine collection: a review. Laboratory
Animals. 2004;38:333-61.

La Joie R, Perrotin A, Egret S, Pasquier F, Tomadesso C, Mézenge F, Desgranges B, de La
Sayette V, Chételat G. Qualitative and quantitative assessment of self-reported cognitive
difficulties in nondemented elders: Association with medical help seeking, cognitive deficits,
and β-amyloid imaging. Alzheimers Dement. 2016;5:23-34.

Lee AC, Tian H, Grosmaitre X, Ma M. Expression patterns of odorant receptors and response
properties of olfactory sensory neurons in aged mice. Chem. Senses. 2009;34:695-703.

Li Y, Dulac C. Neural coding of sex-specific social information in the mouse brain. Curr. Opin.
Neurobiol. 2018;53:120-30.

Locci A, Orellana H, Rodríguez G, Gottliebson M, McClarty B, Domínguez S, Keszycki R,
Dong H. Comparison of memory, affective behavior, and neuropathology in APPNLGF knock-
in mice to 5xFAD and APP/PS1 mice. Behavioural Brain Research. 2021;404:113192.

Mandairon N, Poncelet J, Bensafi M, Didier A. Humans and mice express similar olfactory
preferences. PLoS One. 2009;4:e4209.

Martínez B, Karunanayaka P, Wang J, Tobia MJ, Vasavada M, Eslinger PJ, Yang QX.
Different patterns of age-related central olfactory decline in men and women as quantified by
olfactory fMRI. Oncotarget. 2017;8:79212-222.

Mechin V, Pageat P, Teruel E, Asproni P. Histological and immunohistochemical
characterization of vomeronasal organ aging in mice. Animals. 2021;11:1211.

Mobley AS, Rodríguez-Gil DJ, Imamura F, Greer CA. Aging in the olfactory system. Trends

Neurosci. 2014;37:77-84.

Mohs RC, Schmeidler J, Aryan M. Longitudinal studies of cognitive, functional and behavioural

changes in patients with Alzheimer´s disease. Sta. Med. 2000;19:1401-09.

Moran JM, Jolly E, Mitchell JP. (2012) Social-cognitive deficits in normal aging. J. Neurosci.

2012;32:5553-61.

Moreno M, Richard M, Landrein B, Sacquet J, Didier A, Mandairon N. Alteration of olfactory

perceptual learning and its cellular basis in aged mice. Neurobiol. Aging. 2014;35:680-91.

Munger SD, Leinders-Zufall T, Zufall F. Subsystem organization of the mammalian sense of

smell. Annu. Rev. Physiol. 2009;71:115-40.

Murphy C. Olfactory and other sensory impairments in Alzheimer disease. Nat. Rev. Neurol.

2019;15:11-24.

Murphy C, Schubert CR, Cruickshanks KJ, Klein B.E.K., Klein R., Nondahl DM. Prevalence

of olfactory impairment in older adults. JAMA. 2002;288:2307-12.

Nadler JJ, Moy SS, Dold G, Trang D, Simmons N, Pérez A, Young NB, Barbaro RP, Piven J,

Magnuson TR, Crawley JN. Automated apparatus for quantitation of social approach

behaviors in mice. Genes Brain Behav. 2004;3:303-14.

Noack J, Richter K, Laube G, Haghgoo HA, Veh RW, Engelmann M. Different importance of

the volatile and non-volatile fractions of an olfactory signature for individual social recognition

in rats versus mice and short-term versus long-term memory. Neurobiol. Learn. Mem.

2010;94:568-75.

Olesen LØ, Bouzinova EV, Severino M, Sivasaravanaparan M, Hasselstrøm JB, Finsen B,

Wiborg O. Behavioural phenotyping of APPswe/PS1δE9 mice: Age related changes and effect

of long-term paroxetine treatment. PLoS One. 2016;11:e0165144,

10.1371/journal.pone.0165144 PMID: 27814403.

Oboti L, Ibarra-Soria X, Pérez-Gómez A, Schimd A, Pyrski M, Paschek N, Kircher S, Logan
DW, Leinders-Zufall T, Zufall F, Chamero P. Pregnancy and estrogen enhance neural
progenitor-cell proliferation in the vomeronasal sensory epithelium. BMC Biol. 2015;13:104
doi: 10.1186/s12915-015-0211-8.

Panaliappan TK, Wittmann W, Jidigam VK, Mercurio S, Bertolini JA, Sghari S, Bose R, Patthey
C, Nicolis SK, Gunhaga L. Sox2 is required for olfactory pit formation and olfactory
neurogenesis through BMP restriction and Hes5 upregulation. Development. 2018;145:
dev153791.

Parrish-Aungst S, Shipley MT, Erdelyi F, Szabo G, Puche AC. Quantitative analysis of
neuronal diversity in the mouse olfactory bulb. J. Comp. Neurol. 2007;501:825-36.

Patel RC, Larson J. Impaired olfactory discrimination learning and decreased olfactory
sensitivity in aged C57Bl/6 mice. Neurobiol. Aging. 2009;30:829-37.

Rawson NE, Gomez G, Cowart BJ, Kriete A, Pribitkin E, Restrepo D. Age-associated loss of
selectivity in human olfactory sensory neurons. Neurobiol. Aging. 2012;33:1913-19.

Reiserer RS, Harrison FE, Syverud DC, McDonald MP. Impaired spatial learning in the
APPSwe + PSEN1DeltaE9 bigenic mouse model of Alzheimer´s disease. Genes Brain Behav.
2007;6:54-65.

Richard MB, Taylor SR, Greer CA. Age-induced disruption of selective olfactory bulb synaptic
circuits. Proc. Natl. Acad. Sci. U S A. 2010;107:15613-618.

Roberts RO, Christianson TJ, Kremers WK, Mielke MM, Machulda MM, Vassilaki M, Alhurani
RE, Geda YE, Knopman DS, Petersen RC. Association between olfactory dysfunction and
amnestic mild cognitive impairment and Alzheimer disease dementia. JAMA Neurol.
2016;73:93-101.

Root CM, Denny CA, Hen R, Axel R. The participation of cortical amygdala in innate, odour-
driven behaviour. Nature. 2014;515:269-73.

Sanderson DJ, Bannerman DM. Competitive short-term and long-term memory processes in

spatial habituation. J. Exp. Psychol. Anim. Behav. Process. 2011;37:189-99.

Saraiva LR, Kondoh K, Ye X, Yoon KH, Hernandez M, Buck LB. Combinatorial effects of

odorants on mouse behavior. Proc. Natl. Acad. Sci. U S A. 2016;113: E3300-E3306.

Scopa C, Marrocco F, Latina V, Ruggeri F, Corvaglia V, La Regina F, Ammassari-Teule M,

Middei S, Amadoro G, Meli G, Scardigli R, Cattaneo A. Impaired adult neurogenesis is an

early event in Alzheimer's disease neurodegeneration, mediated by intracellular Aβ oligomers.

Cell Death Difer. 2020;27:934-48.

Spalletta G, Baldinetti, Buccione I, Fadda L, Perri R, Scalmana S, Serra L, Caltagirone C.

Cognition and behaviour are independente and hetreogenous dimensions in Alzheimer´s

disease. J. Neurol. 2004;251:68-95.

Stockdale SE, Wells KB, Tang L, Belin TR, Zhang L, Sherbourne CD. The importance of social

context: Neighborhood stressors, stress-buffering mechanisms, and alcohol, drug, and mental

health disorders. Social Science & Medicine. 2007;65:1867-81.

Taroc EZM, Katreddi RR, Forni PE. Identifying Isl1 genetic lineage in the developing olfactory

system and in GnRH-1 neurons. Front. Physiol. 2020;11:601923.

Tikhonova MA, Romaschenko AV, Akulov AE, Ho YJ, Kolosova NG, Moshkin MP,

Amstislavskaya TG. Comparative study of perception and processing of socially or sexually

significant odor information in male rats with normal or accelerated senescence using fMRI.

Behav. Brain Res. 2015;294:89-94.

Tobiansky DJ, Hattori T, Scott JM, Nutsch VL, Roma PG, Dominguez JM. Mating-relevant

olfactory stimuli activate the rat brain in an age-dependent manner. Neuroreport.

2012;23:1077-83.

Tobin VA, Hashimoto H, Wacker DW, Takayanagi Y, Langnaese K, Caquineau C, Noack J,

Landgrave R, Onaka T, Leng G, Meddle SL, Engelmann M, Ludwig M. An intrinsic vasopressin

system in the olfactory bulb is involved in social recognition. Nature. 2010;464:413-17.

Tucker ES, Lehtinen MK, Maynard T, Zirlinger M, Dulac C, Rawson N, Pevny L, La-Mantia

AS. Proliferative and transcriptional identity of distinct classes of neural precursors in the

mammalian olfactory epithelium. Development. 2010;137:2471-81.

Ueha R, Shichino S, Ueha S, Kondo K, Kikuta S, Nishijima H, Matsushima K, Yamasoba T.

Reduction of proliferating olfactory cells and low expression of extracellular matrix genes are

hallmarks of the aged olfactory mucosa. Front. Aging Neurosci. 2018;10:86.

Valech N, Tort-Merino A, Coll-Padrós N, Olives J, León M, Rami L, Molinuevo JL. Executive

and language subjective cognitive decline complaints discriminate preclinical Alzheimer's

disease from normal aging. J. Alzheimers Dis. 2018;61:689-703.

Verret L, Jankowsky JL, Xu GM, Borchelt DR, Rampon C. Alzheimer's-type amyloidosis in

transgenic mice impairs survival of newborn neurons derived from adult hippocampal

neurogenesis. J. Neurosci. 2007;27:6771-80.

Wacker DW, Tobin VA, Noack J, Bishop VR, Duszkiewicz AJ, Engelmann M, Meddle SL,

Ludwig M. Expression of early growth response protein 1 in vasopressin neurones of the rat

anterior olfactory nucleus following social odour exposure. J. Physiol. 2010;588:4705-17.

Weiler E, McCulloch MA, Farbman AI. Proliferation in the vomeronasal organ of the rat during

postnatal development. Eur. J. Neurosci. 1999;11:700-11.

West MJ, Gundersen HJ. Unbiased stereological estimation of the number of neurons in the

human hippocampus. J. Comp. Neurol. 1990;296:1-22.

Whitman MC, Greer CA. Adult neurogenesis and the olfactory system. Prog. Neurobiol.

2009;89:162-75.

Wilson RS, Krueger KR, Arnold, SE, Schneider JA, Kelly JF, Barnes LL, Tang Y, Bennett DA.

Loneliness and risk of Alzheimer disease. Arch. Gen. Psychiatry. 2007;64:234-40.

Wong FK, Bercsenyi K, Sreenivasan V, Portalés A, Fernández-Otero M, Marín O. Pyramidal

cell regulation of interneuron survival sculpts cortical networks. Nature. 2018;557:668-73.

Xu F, Schaefer M, Kida I, Schafer J, Liu N, Rothman DL, Hyder F, Restrepo D, Shepherd GM.

Simultaneous activation of mouse main and accessory olfactory bulbs by odors or

pheromones. J. Comp. Neurol. 2005;489:491-500.

Yang M, Crawley JN. Simple behavioral assessment of mouse olfaction. Curr. Protoc.

Neurosci. 2009;8:doi: 10.1002/0471142301.ns0824s48.

Yang M, Silverman JL, Crawley JN. Automated three-chambered social approach task for

mice. Curr. Protoc. Neurosci. 2011;8:8.26. doi: 10.1002/0471142301.ns0826s56.

Yoon H, Enquist LW, Dulac C. Olfactory inputs to hypothalamic neurons controlling

reproduction and fertility. Cell. 2005;123: 669-82.

Zeng Q, Zheng M, Zhang T, He G. Hippocampal neurogenesis in the APP/PS1/nestin-GFP

triple transgenic mouse model of Alzheimer´s disease. Neuroscience. 2016;314:64-74.

Zhang C, McNeil E, Dressler L, Siman R. Long-lasting impairment in hippocampal

neurogenesis associated with amyloid deposition in a knock-in mouse model of familial

Alzheimer's disease. Exp. Neurol. 2007;204:77-87.

Zufall F, Leinders-Zufall T. Mammalian pheromone sensing. Curr. Opin. Neurobiol.

2007;17:483-89.

FIGURE LEGENDS

Figure 1. Structural modifications of the mouse VSE during natural and pathological

aging. A) Dispersion plot of the VSE volume indicates a significant reduction during natural

but not pathological aging. **B)** Dispersion plot of AOB volume. Thick lines indicate the mean ±

SEM. Data were obtained from 4 animals per condition. **C)** Representative images of OMP

staining in the VSE during natural and pathological aging. Scale bar indicates 100 μm. **D)**

Representative images of Sox2 staining in the VSE and the supporting cell layer. Scale bar

indicates 100 μm. VL: vomeronasal lumen; SCL: supporting cell layer; NSE: non-sensory

epithelium; M: marginal zone; I: intermediate zone; C: central zone. **E, F)** Dispersion plots

represent the number of OMP$^+$ cells in the VSE (number of cells x $10^3/\mu m^3$) and the intensity

of Sox2 fluorescence in the SCL (Sox2 CTF $/\mu m^2$). Thick lines indicate the mean ± SEM. Data

were obtained from 10-16 slices from at least three animals per condition. Statistical

comparisons were calculated by two-way ANOVA with Tukey´s test. P ≤ 0.05 was considered

statistically significant.

Figure 2. Natural and pathological aging differentially impact cell proliferation in the

VSE. A) Representative confocal images of PCNA staining in the VSE proliferative niche

during natural and pathological aging. Panels on the right show confocal images of OMP and

PCNA staining in the VNO. To note there is no overlapping signal between the two markers

indicating the identification of cells at different maturation stages. Circles indicate the

proliferative marginal region of the VNO. Scale bar represents 20 μm. **B, C)** Dispersion plots

represent the number of PCNA$^+$ cells (number of cells x $10^3/mm^3$) and the number of Sox2$^+$

cells (number of cell x $10^4/\mu m^2$) in the marginal VNO. Thick lines indicate the mean ± SEM.

Data were obtained from ten slices from animal of each condition. Statistical analysis was

calculated by two-way ANOVA with Tukey´s test for multiple comparisons. P ≤ 0.05 was

considered statistically significant.

Figure 3. Natural and pathological aging reduce exploration to social odors. A)

Schematics of the olfactory test used in this study in which urine dilutions are presented as a

social signal. **B)** Average of the exploration (sniffing) time of urine serial dilutions normalized

to the exploration time of the vehicle (water) of adult and aged wild type mice. A typical

habituation was observed in adult mice at intermediate dilutions (1:500, 1:250, 1:100; 1:50).

C) Dispersion plot of the normalized sniffing time of each urine dilution (nd, non-diluted; 1:10;

1:50; 1:100; 1:250; 1:500; 1:1000) for adult and aged wild type mice. **D)** Average of the sniffing

time of urine dilutions normalized to the exploration time of the vehicle (water) of middle age

APP/PS1 controls (APP^{WT}) and middle age APP/PS1 heterozygous mice (APP/PS1^{Het}). **E)**

Dispersion plots of normalized sniffing time of middle age APP^{WT} control and APP/PS1^{Het} mice.

Grey lines in the dispersion plots indicate mean ± SEM. Data were analyzed by a one-way

ANOVA with Tukey´s test to test multiple comparisons with more than one variable. P ≤ 0.05

was considered statistically significant.

Figure 4. Aging and neurodegeneration impact social odor recognition more severely

than other odor modalities. A) Average of the sniffing time of young and aged mice in

response to a neutral synthetic odor isoamyl acetate (IA) normalized to the vehicle (mineral

oil). **B)** Dispersion plots of the normalized sniffing time of young and aged mice in response to

various dilutions of IA. **C)** Average of the sniffing time of middle age APP/PS1 controls (APP^{WT})

and middle age APP/PS1 heterozygous mice (APP/PS1^{Het}) in response to IA normalized to

the vehicle (mineral oil). **D)** Dispersion plots of normalized sniffing time of APP/PS1^{WT} and

APP/PS1^{Het} mice in response to various dilutions of IA. Note that APP/PS1^{Het} mice exhibit

significantly higher exploration times for the 1:10^4 and 1:100 IA dilutions. **E)** Food deprived

animals performed a food seeking test (schematics on the right) which revealed equivalent

latencies to the find hidden food pellets as shown in the dispersion data plot (left panel). Thick

lines in dispersion plots indicate mean ± SEM. Data were analyzed by a one-way ANOVA with

Tukey´s test to test multiple comparisons with more than one variable. Number of mice:

Neutral odor sensitivity: young: n = 19; aged: n = 27; middle age control: n = 10; middle age

APP/PS1^{Het}: n = 13. *Food finding test*: young: n = 21; aged: n = 27; APP^{WT}: n = 20; APP^{Het}: n

= 16. P ≤ 0.05 was considered statistically significant.

Figure 5. Social odor discrimination and habituation are reduced in naturally aged and

APP/PS1^{Het} mice. A) Schematics of the social habituation-dishabituation test used in this

study: after habituation to the experimental cage, the animal is exposed to the same urine

sample three times (S1a-c) which induces a typical increase in exploration time (1st

discrimination) to be followed by a reduction in the sniffing time (1st habituation). Dishabituation

induced by a urine sample from a new subject (S2a) elicits a second round of discrimination -

925 habituation. **B)** Average sniffing time of the social habituation-dishabituation test performed by

926 young and aged wild type mice. **C)** Dispersion plot of the sniffing time of young and aged

927 animals during the social habituation-dishabituation test. **D)** Average sniffing time of the social

928 habituation-dishabituation test performed by middle age APP/PS1^{WT} control and APP/PS1^{Het}

929 mice. **E)** Dispersion plot of the sniffing time of middle age APP/PS1^{WT} control and APP/PS1^{Het}

930 mice during the social habituation-dishabituation test. **F**) The habituation-dishabituation test

931 was modified to assess potential deficits in long-term discrimination due to natural or

932 pathological aging by presenting the same S1a sample 24 hours after. **G**) Paired data

933 corresponding to the sniffing time during the first and second S1a presentation (24 hours later)

934 is plotted for each condition. Thick lines in dispersion plots and black dots in G indicate mean

935 ± SEM. Data were analyzed by a one-way ANOVA with Tukey´s test to test multiple

936 comparisons with more than one variable. $P \leq 0.05$ was considered statistically significant.

937 **Figure 6. Social discrimination and habituation are more severely altered during natural**

938 **aging. A)** Paired data of the sniffing time young and aged wild type animals corresponding to

939 the first discrimination phase (H_2O-S1a). **B**) Paired data of the sniffing time of young and aged

940 wild type animals corresponding to the second discrimination phase (S1c-S2a). **C)** Dispersion

941 plots of the slope values corresponding to the first and second discrimination phases of young

942 and aged mice. **D)** Paired data of the sniffing time of APP^{WT} and APP^{Het} mice corresponding

943 to the first discrimination phase (H_2O-S1a). **E**) Paired data of the sniffing time of APP^{WT} and

944 APP^{Het} mice corresponding to the second discrimination phase (S1c-S2a). **F**) Dispersion plots

945 of APP^{WT} and APP^{Het} mice corresponding to the slope values of the first and second

946 discrimination phases. **G)** Paired data of the sniffing time of young and aged wild type animals

947 corresponding to the first habituation phase (S1a-S1b). **H)** Paired data of the sniffing time of

948 the young and aged wild type animals corresponding to the second habituation phase (S2a-

949 S2b). **I**) Dispersion plots of the slope values of young and aged mice corresponding to the first

950 and second habituation phases. **J)** Paired data of the sniffing time of APP^{WT} and APP^{Het} mice

951 corresponding to the first habituation phase (S1a-S1b). **K)** Paired data of the sniffing time of

952 APP^{WT} and APP^{Het} mice corresponding to the second habituation phase (S2a-S2b). **L)**

Dispersion plots of the slope values of APP^{WT} and APP^{Het} mice corresponding to the first and

second habituation phases. Black dots in A, B, D, E, G, H, J, K and thick lines in C, F, I, L

indicate mean ± SEM. Data were analyzed by one-way ANOVA with Tukey´s test. P ≤ 0.05

was considered statistically significant.

Figure 7. Age-related deficits in social discrimination and habituation are not influenced

by previous experience. A) Average sniffing time of social habituation-dishabituation test in

response to odors from novel (N) or littermate (L) subjects. **B)** Dispersion plot of the

exploration time of aged and young animals in response to novel or littermate urine samples.

C) Average sniffing time of social habituation-dishabituation test in response to littermate and

the animal´s own urine (O). **D)** Dispersion plot of exploration time of aged and young animals

in response to their own´s or littermate urine samples. Thick lines in dispersion plots indicate

mean ± SEM. Number of mice: young: n = 21; aged: n = 23. Data were analyzed by a one-

way ANOVA with Tukey´s test to test multiple comparisons with more than one variable. P ≤

0.05 was considered statistically significant.

Figure 8. Social novelty is disrupted during pathological aging. A) Schematics of the

three chamber test used in this study to test sociability and social novelty. In the sociability

phase the sniffing time of E and M1^A are compared. Social novelty is estimated by quantifying

the sniffing time of exploring M1^B versus M2 (see Methods). **B)** Paired data of the sniffing

times of young wild type mice during the sociability (E-M1^A) and social novelty (M1^B-M2)

phases. **C)** Paired data of the sniffing times of aged wild type mice during sociability and social

novelty. **D)** Paired data of the sniffing times of middle age APP^{WT} mice during sociability) and

social novelty. **E)** Paired data of the sniffing times of middle age APP^{Het} mice during sociability

and social novelty. Black dots in the paired plots indicate mean ± SEM. Number of mice:

young: n = 10; aged: n = 14; middle age control: n = 10; middle age APP/PS1^{Het}: n = 12. Data

were analyzed by a one-way ANOVA with Tukey´s test to test multiple comparisons with more

than one variable. P ≤ 0.05 was considered statistically significant.

Additional file 1. Structural modifications of the mouse VSE and AOB during natural

and pathological aging. A) Schematics depicting the VNO divisions into marginal (M),

intermediate (I) and central (C). **B**) Dispersion plots of the number of OMP⁺ cells in the

marginal zone of the VSE combining anterior, medial and posterior sections. Thick lines

indicate the mean ± SEM. Data were obtained from 10-16 slices from at least three animals

per condition. Statistical comparisons were calculated by two-way ANOVA with Tukey´s test.

$P \leq 0.05$ was considered statistically significant.

Additional file 2. Social odor exploration time during natural aging. Dispersion plots

represent the raw sniffing time of 2, 6, 12 and 24 months-old animals for the various serial

dilutions tested in **Fig. 3**. Thick grey lines indicate mean ± SEM. Data were analyzed by a one-

way ANOVA with Tukey´s test to test multiple comparisons with more than one variable. P

and n values are indicated in the tables. $P \leq 0.05$ was considered statistically significant.

Additional file 3. Statistical comparison between different urine dilutions for each age

tested (2, 6, 12 and 24 months-old). Dispersion plots represent the sniffing time and trend

curves of 2, 6, 12 and 24 months-old animals. Thick dots indicate mean ± SEM. P values for

each statistical comparison are indicated in the tables on the right. Data were analyzed by a

one-way ANOVA with Tukey´s test to test multiple comparisons with more than one variable.

$P \leq 0.05$ was considered statistically significant.

Additional files 4. Aged-related deficits in social odor exploration exhibit no significant

sex differences. A) Average sniffing time of urine serial dilutions of adult and aged female

wild type mice. **B**) Average exploration (sniffing) time of urine serial dilutions of adult and aged

male wild type mice. **C**) Dispersion plots of the sniffing time of urine dilutions that show

significant impairments in female mice (nd, non-diluted; 1:10; 1:500; 1:1000). **D**) Dispersion

plots of the sniffing time of the urine dilutions that show significant changes in male mice (nd,

non-diluted; 1:250; 1:500; 1:1000). Plots indicate mean ± SEM. Number of mice: adult

females: n = 13; adult males: n = 11; aged females: n = 14; aged males: n = 11. Data were

analyzed by a one-way ANOVA with Tukey´s test to test multiple comparisons with more than

one variable. $P \leq 0.05$ was considered statistically significant.

Additional file 5. Social odor exploration of APP^WT and APP^Het mice. Dispersion plots

represent the raw sniffing time of middle age APP^WT and APP^Het mice for the various serial

dilutions. Thick grey lines indicate mean ± SEM. Number of mice for each condition is indicated

in the table below. Data were analyzed by a one-way ANOVA with Tukey´s test to test multiple

comparisons with more than one variable. $P \leq 0.05$ was considered statistically significant.

Additional file 6. Statistical comparison between different urine dilutions of APP/PS1^{WT}

and APP/PS1^{Het} mice. Dispersion plots represent the sniffing time and trend curves of middle

age APP^{WT} and APP^{Het} mice for the various serial dilutions. Thick dots indicate mean ± SEM.

P values for each statistical comparison are indicated in the tables on the right. Data were

analyzed by a one-way ANOVA with Tukey´s test to test multiple comparisons with more than

one variable. $P \leq 0.05$ was considered statistically significant.

Additional file 7. Latency to approach the first stranger (M1^{A}) in the three chamber

sociability test is increased in naturally aged animals. A) The sociability phase of the three

chamber test involves exposing the test animal to a stranger mouse (M1^{A}) for the first time,

which elicits an increase in the exploration time in that side of the chamber. **B**) Dispersion

plots of the latency to approach M1 during the first encounter of the sociability phase. Thick

lines indicate mean ± SEM. Data were analyzed by one way-ANOVA with Tukey´s test. $P \leq$

0.05 was considered statistically significant.

ABBREVIATIONS

AD: Alzheimer´s disease

AOB: Anterior olfactory bulb

APP: Amyloid precursor protein

C: VNO central zone

CTF: Corrected total fluorescence

FFT: Food finding test

I: VNO intermediate zone

IA: Isoamyl acetate

M: VNO marginal zone

MOB: Main olfactory bulb

MOE: Main olfactory epithelium

NSE: Non-sensory epithelium

OCT: Optimal cutting temperature

OMP: Olfactory marker protein

PCNA: Proliferative cell nuclear antigen

PS1: Presinilin-1

SCL: Supporting cell layer

Sox2: SRY-box transcription factor 2

VL: Vomeronasal lumen

VNO: Vomeronasal organ

VSE: Vomeronasal epithelium

Figure 1

Figure 2

Figure 3

NEUTRAL ODOR EXPLORATION

A NATURAL AGING

B

C PATHOLOGICAL AGING

D

FOOD FINDING TEST

E

Figure 4

SOCIAL ODOR HABITUATION DISHABITUATION

A

Young 2-4 mo
APP^WT / **APP^HET** 12-14 mo
Aged 20-24 mo

URINE SAMPLES

1 min exposure — H_2O^A — H_2O^B

Sample 1 (S1): S1a, S1b, S1c
Sample 2 (S2): S2a, S2b, S2c

1 min interval

1 min trials

B NATURAL AGING

- Young
- Aged

discrimination / habituation
< 0.001
< 0.01
< 0.001
< 0.05

sniffing time (s) vs Samples (H_2O^A, H_2O^B, S1a, S1b, S1c, S2a, S2b, S2c)

C YOUNG   AGED

< 0.05   < 0.05
< 0.001, < 0.001, < 0.001, < 0.001
< 0.01, < 0.05, < 0.05, < 0.05

sniffing time (s) vs Samples

D PATHOLOGICAL AGING

- APP^WT
- APP^Het

discrimination / habituation
< 0.001
< 0.001
< 0.05
> 0.05

sniffing time (s) vs Samples

E APP^WT   APP^Het

< 0.05   < 0.05
< 0.001, < 0.001, < 0.05, < 0.05
< 0.001, < 0.001, > 0.05, < 0.05

sniffing time (s) vs Samples

F LONG-TERM SOCIAL DISCRIMINATION

Day 0 — 1st discrimination — 24 h → Day 1 — 1st discrimination

S1a time (s) vs S1a

G
- Young
- Aged
- APP^WT
- APP^Het

< 0.01   > 0.05   < 0.05   > 0.05

S1a sniffing time (s) — Day 0 / Day 1

Figure 5

SOCIAL DISCRIMINATION

SOCIAL HABITUATION

Figure 6

Figure 7

A   THREE CHAMBER SOCIABILITY TEST

Figure 8