Stress History Modulates CRF Neurons to Establish Resilience

Authors:

Sherod E Haynes[1,2,3], Anthony Lacagnina[4*], Hyun Seong Seo[1*], Muhammad Afzal[5], Carole Morel[3], Aurelie Menigoz[1], Kanaka Rajan[5,6], Roger L Clem[4], Barbara Juarez[7], Helen S Mayberg[5], Donald G. Rainnie[1], Larry J. Young[1], Ming-Hu Han[3,4,8†]

Affiliations:

[1]Center for Translational Social Neuroscience, Yerkes National Primate Research Center, Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, GA 30322, USA

[2]Neuroscience Graduate Program, Emory University, Atlanta, GA 30322, USA

[3]Department of Pharmacological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[4]Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[5]Nash Family Center for Advanced Circuit Therapeutics, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[6]Department of Neurobiology, Harvard Medical School, Kempner Institutes for the Study of Natural and Artificial Intelligence, Boston, MA 02115

[7]Department of Anatomy and Neurobiology, University of Maryland, Baltimore School of Medicine, Baltimore, MD, 21201 USA

[8]Department of Mental Health and Public Health, Faculty of Life and Health Sciences, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, Guangdong, China

35   *These authors contributed equally

36   †Corresponding electronic address: ming-hu.han@mssm.edu.

37
38

Abstract

Introduction: Cumulative stress is a major risk factor for developing major depressive disorder (MDD), yet not everyone experiencing chronic stress develops MDD. In those who do not, it is unclear at what point, or by what mechanism, a trajectory of stable resiliency emerges.

Methods: Utilizing a 10-day repeated social defeat stress model (RSDS) for MDD, we observed that a critical period between 7 and 10 daily defeats marks the phenotypical divergence of resilient from susceptible mice. Using cell-type selective electrophysiology, chemogenetics, optogenetics, fiber photometry and RNA quantification was employed to investigate the nature of stress effects on neuroadaptation in the oval nucleus of the bed nucleus of the stria terminalis (BNSTov) required to determine resilience.

Results: In response to ongoing stress, corticotropin-releasing factor (CRF^+, but not CRF^-) neurons of the (BNSTov) displayed a sustained increased firing rate in resilient, but not susceptible mice. This neurophysiological adaptation was self-sustaining, but only after 7 critical stress exposures, indicating that the process of developing resilience is dependent on stress history.

Conclusion: Our study reveals a novel process by which individuals might persist in the face of adversity by way of stress-provoked activation, not inhibition of a key CRF limbic region that establishes a pathway to resilience.

Introduction:

Major depressive disorder (MDD) is a crippling heterogeneous neuropsychiatric condition with

high morbidity and lifetime prevalence.[1,2] Major life stressors are key precipitants in the onset of

an episode[3,4]. While most people will report at least one major life stressor at some point in

their lives, not everyone will go on to develop MDD. It is also known that depression can

emerge after significant repeated stressor exposure, suggesting that cumulative exposure to

stress is associated with increased depression risk in vulnerable individuals[5–8]. Cumulative

stress causes numerous psychological insults that predispose neuropsychiatric conditions[5,9,10].

Many studies have explored the individual differences in stress susceptibility or resiliency, but

the process critical to driving the divergence in phenotypes remain elusive. With the odds of

treatment failure increasing with subsequent depressive episodes,[2,12,13] and increased

depression risk of cumulative stress, a potentially more effective strategy would be to target the

mechanisms mediating resilience, enhancing the ability to cope with cumulative stress.

Resiliency is defined as the "process of adapting well in the face of adversity… [to] threats, or

significant sources of stress."[14-16] Resiliency is unlikely to emerge from the absence of a

pathological stress response but instead is an active process involving a myriad of molecular,

cellular and circuit-level changes in the brain. Repeated social defeat stress (RSDS) induces

robust depression-like behavioral phenotypes in roughly 2/3 of mice[6,10]. The standard 10-day

RSDS protocol has been employed widely to identify and study neurobiological features of

susceptible and resilient subpopulations[6,7,17–23]. Studies on stress vulnerability often test

hypotheses after establishing vulnerable subgroups, making the processes that led to such

phenotypic diversification unclear.

The Bed Nucleus of the Stria Terminalis (BNST) is an important node for integrating sensory

cues, interoception, cognition, and motivational states to enact adaptive responses[24,25]. The

BNST is well-positioned in the social salience network to integrate external cues with internal

states to influence the outcome and context of social interactions[24–27]. The oval nucleus of the

BNST (BNSTov) is a stress-sensitive subregion that is a key candidate region for encoding

stress modulation of social behavior through its projections to areas such as the ventral

tegmental area (VTA)[28–31] and dorsal raphe[32,33]. Corticotrophin-releasing factor (CRF) neurons

of the BNSTov (BNSTov[CRF]) are a significant output of this region and can influence affective

states[34–37]. Since BNSTov[CRF] neurons are thought to promote arousal and adaptive responding

according to stress-related changes in internal state[24,25,27,38–44], we hypothesized they might play

a role in establishing resiliency to repeated social stress.

Using the 10-day RSDS paradigm, we used cell-type selective *ex vivo* electrophysiology,

chemogenetics, optogenetics, and *in vivo* fiber photometry to interrogate the BNSTov[CRF] system

to explore its role in the divergence of susceptible and resilient phenotypes. We observed that

BNSTov[CRF] neurons encode repetitive social stress and undergo adaptation that coincides with

the divergence of resilient and susceptible phenotypes. Unexpectedly, we observed that

resiliency entails cumulative stress-history dependent neuroadaptive changes, whereas

susceptibility emerges in the absence of similar adaptations. Activation of BNSTov[CRF] neurons

using Designer-Receptors-Exclusively-Activated-by-Designer-Drugs (DREADDs) during social

defeat led to a resilient phenotype while inhibition caused susceptibility. Finally, using

RNAscope and optogenetics, we provide intriguing evidence for a potential role of *Crfr1* in

BNSTov[CRF] neurons in mediating resilience.

48
49   **Methods & Materials**

50
Mice
The study used wild-type, Crf-ires-Cre (Jackson labs: 011087), ai14 (Cre-responsive tdTomato

reporter mouse; Jackson labs: 007915), ai32 (Cre-responsive channelrhodopsin-2/fused with

eYFP; Jackson labs: 012569) mice on C57BL/6J background that were bred at Icahn School of

Medicine at Mount Sinai and used were between 6-7 weeks at the start of experimental

manipulations. All experiments were approved by the Institutional Animal Care and Use

Committee and comply with institutional guidelines for the Animal Care and Use Committee set

forth by Icahn School of Medicine at Mount Sinai.

Repeated social defeat stress paradigm

The repeated social defeat stress paradigm was performed according to published

protocols[6,21,23,82,90–92]. Briefly, CD1 aggressors were singly housed in cages (26.7 cm width x

48.3 cm depth x 15.2 cm height; Allentown Inc) at least 24 hours before the start of the

experiment on one side with a clear perforated Plexiglas divider. Social stress consists of

physical and sensory stress components.

Viral constructs

For DREADD experiments in CRF neuronal populations, CRF-ires-Cre animals were injected

with AAV5-hSyn-DIO-hM4D(Gi)-mCherry (Addgene: 44362-AAV5) (≥7x10^12), AAV5-hSyn-

DIO-hM3D(Gq)-mCherry (Addgene: 44361-AAV5) (≥7x10^12), and AAV5-hSyn-DIO-mCherry

(Addgene: 50459). For fiber photometric recordings, CRF-ires-Cre animals were injected with

AAV9-syn-FLEX-jGCaMP7f-WPRE (addgene: 104492-AAV9) (>1x10^12). All viruses were

purchased from Addgene.

CNO-drinking water construct

Clozapine-N-Oxide (CNO) was obtained from Hello Bio (catalog no HB6149). The dry chemical

was dissolved in drinking water obtained from the vivarium and diluted such that each mouse

received 5 mg/kg/day based on previous studies. CNO was made fresh each day for the three

days it was administered. CNO solutions were protected from light throughout the experimental

procedure. On average, mice consumed ~4-5 mL of water per day. Water bottles and mice

were weighed daily.

Please see supplementary methods for further details.

Results:

Susceptible vs. resilience phenotypes emerge between 7 and 10 daily episodes of social

defeat stress.

RSDS produces divergent and enduring resilient/susceptible phenotypes following 10 episodes

of social defeat stress[6,21] and enables the investigation of the consequences of stress

accumulation[5,7]. To examine the temporal divergences of resilient and susceptible phenotypes

in mice, a modified RSDS stress protocol was employed in which mice were subjected to

discrete numbers of social defeat episodes (SDEs) interspersed with social interaction (SI), and

sucrose preference (SP) tests administered after 1, 4, 7, and 10 SDEs (Fig. 1A-D). Social

Interaction (SI) Ratio is a behavioral score of the SI test and measures the time spent in the

area proximal to the enclosure of a novel social target (social interaction zone). To generate the

SI ratio, interaction zone time is calculated both in the absence and presence of the social

target, such that SI ratio $\geqq$ 1 is resilient and <1 is susceptible[21]. Surprisingly, we found that the

susceptible phenotype emerged discretely between 7 and 10 SDEs (Fig. 1E, Supplemental

Data Fig 1A) (SI ratio: 2.062 +/- 0.226 after 7 SDEs to 0.559 +/- 0.093, p<0.0001, n = 11 for

susceptible and 1.967 +/- 0.188 (7 SDEs) to 1.554 +/- 0.163 (10 SDEs), p=0.2929, n=11 for

resilient). There were no significant differences in SI ratios after 7 SDEs between mice that went

on to become susceptible vs resilient after 10 SDEs (mean: 2.062 +/- 0.226 vs 1.967 +/- 0.188,

two-tailed t-test, p=0.7483, n= 11/group) (Supplemental Data Fig. 1B-C). This effect was not

due to repeated SI tests (Supplemental Data Fig. 2D-F). Resilient mice had an

indistinguishable SI ratio from all mice after 7 SDEs, while susceptible (10 SDEs) mice had an

SI ratio significantly lower than both groups (Fig. 1F, G). In the sucrose preference (SP) test,

resilient mice showed a similar preference for sucrose to all 7 SDE and stress-naïve control

mice. Previous studies reported that RSDS produces susceptible and resilient phenotypes after

10 SDEs in a bimodal distribution[21]. Here, we observed a unimodal distribution of social

interaction toward a novel conspecific in 7 SDE mice but a bimodal distribution in 10 SDE mice,

consistent with the emergence of distinct resilience and susceptible phenotypes (Kolmogorov-

Smirnov, p<0.0001; Fig. 1 (H,I,J). The time spent interacting with a novel conspecific was

significantly less in susceptible 10 SDE mice compared to the 7th SDE and 10th SDE resilient

mice, which were indistinguishable (Fig.1K). The phenotypic divergence was dependent upon

the number of SDEs rather than the passage of time and dependent upon the anterior dorsal

BNST(Supplemental Data Fig. 1G, 2A-E).

Divergence in BNSTov^{CRF} neuronal firing rates tracks the emergence of resilient and

susceptible phenotypes.

CRF neurons of the BNST are a major output source of the oval BNST that are sensitive to

chronic stressors[27,45−47]. Thus, we hypothesized that chronic stress would alter BNSTov^{CRF}

neuronal activity, coinciding with the divergence in resilient and susceptible behavioral

phenotypes. To test this hypothesis, *Crf-ires-Cre;*ai14 (tdTomato) mice[30,48−51] were subjected to

either 7 or 10 daily SDEs, and cell-attached *ex-vivo* electrophysiological recordings were

conducted in the BNSTov (Fig. 2A,B). CRF$^+$, but not CRF$^-$, BNSTov neurons of 7 SDE mice

had significantly increased firing rates compared to stress-naïve control mice. By contrast,

CRF[+] firing in susceptible 10 SDE mice was indistinguishable from that of controls and lower

than 7 SDE and resilient mice. CRF[-] neuronal firing did not differ between groups (Fig. 2C,D).

Moreover, there was a strong correlation between firing rate and social interaction ratio in CRF[+]

but not CRF[-] neurons in 10 SDE mice (CRF[+], R^2=0.5725, *p=0.0113; CRF- R^2=0.06096,

p=0.5219, Fig. 2E). CRF neurons display both burst and non-burst firing patterns[40,52,53] (Fig. 2F).

Burst firing patterns were prominent in resilient and 7 SDE mice compared to susceptible and

control mice (Fig. 2H). Additionally, percentage spikes within burst were higher in 7 SDE and

resilient compared to susceptible 10 SDE mice (***p=0.0004, one-way ANOVA, Fig. 2I), but not

susceptible or control mice. The number of spikes per burst and number of bursts per cell was

not significantly different amongst the groups (Fig. 2J,K). These data and correlation analysis

suggest the possibility that there is a causal link between the neuronal activity of BNSTov[CRF]

neurons and the divergence of behavioral phenotypes.

BNSTov[CRF] neurons bidirectionally modulate the emergence of resiliency.

To test the hypothesis that BNSTov[CRF] neurons regulate and maintain resilience over the last 3

episodes of RSDS, we injected *Crf*-ires-*Cre* mice with AAVs encoding Cre-dependent excitatory

(DIO-hM3Dq), inhibitory (DIO-hM4Di) DREADDs, or mCherry construct (control) into the

BNSTov and administered CNO via drinking water[54−56] (Fig. 3A,B). We presaged that this

experimental design afforded the opportunity to modulate neurons over a longer time span than

intraperitoneal (IP) injections would allow with minimal invasiveness, particularly because it was

not clear over when neuroadaptation occurs within the 3-day window. mCherry control mice

exhibited both susceptible and resilient phenotypes (SI ratio <1.0 and ≧1.0, respectively) in

roughly a 60/40 ratio as expected[6,21] (Fig. 3C,F). Interestingly, mice injected with inhibitory DIO-

hM4Di displayed a robust susceptible phenotype, while DIO-hM3Dq mice displayed resilient

phenotypes following CNO drinking water administration (Fig. 3C). Moreover, none of the DIO-

hM4Di + CNO mice went on to develop resilience (0/10, SI >1.0) while 89% (8/9, SI ≧1.0) of the

DIO-hM3Dq were resilient (Fig 3C,F). Notably, the social defeat experience was not affected by

the DREADDs manipulations (Supplemental Data Fig. 3A, B). The SP test — a test of hedonic

behavior — revealed differences between mCherry susceptible (mCherry (S)) and resilient

(mCherry (R)) mice that were mirrored in DIO-hM4Di and hM3Dq mice, respectively. mCherry

(S) and hM4Di mice displayed a significant decrease in SP relative to mCherry (R) and hM3Dq-

injected mice (Fig. 3D). There were no significant differences in locomotion (Fig. 3E). The

chemogenetic manipulation also produced bidirectional effects on anxiety-like behavior in

elevated plus maze (EPM) and open-field tests (Supplemental Data Fig. 4). Surprisingly, mice

injected with hM3Dq-DREADDs went on to become resilient (Fig 3C) although activation of *Crf*

neurons of the BNST have previously been shown to produce depressive- and anxiogenic-like

responses.[27,34,39,46,49,57] Interestingly, resilience was established if BNSTov[CRF] chemogenetic

activation occurred between 7 and 10 SDEs; excitatory hM3Dq-DREADDs activation during

episodes 4-7 or 10-13 failed to display resiliency (Fig. 3G-I), suggesting stress history plays a

critical role in the behavioral outcomes of CRF modulation. Modulating *Crf* neurons with

inhibitory hM4Di- or excitatory hM3Dq- DREADDs between 8 and 10 SDEs led to enduring

susceptible or resilient phenotypes, respectively, up to 6 weeks after CNO manipulation

(Supplemental Data Fig. 6). These observations strongly support that the behavioral outcomes

induced by the activation of BNSTov[CRF] neurons are stress-history dependent.

Calcium dynamics underlying stress adaptation

While we showed that we could bidirectionally drive the establishment of resilience, this

manipulation offers limited insight into the *in vivo* neural dynamics that reflect this divergence

point. To draw a link between neural changes and behavior *in vivo*, we combined fiber

photometry (gCAMP7f) with excitatory or inhibitory DREADDs (Cre-dependent -hM4Di, -hM3Dq

DREADDs or -mCherry viral vectors) to mimic the neuroadaptive changes in BNSTov^{CRF} and

bidirectionally drive the development of resilience/susceptibility (Fig 4A, B). Clozapine-N-oxide

(CNO) was administered via drinking water over the last 3 episodes of RSDS.

Given that mice with shared stress-history of 7 SDEs appeared to have unimodal distribution on

a variety of measures, we hypothesized that calcium-encoded neural activity may diverge in

susceptible/resilient mice concomitant with the display of their respective behaviors on SI test.

When comparing changes in calcium-encoded neural dynamics between SI Test 1 and 2,

mCherry (S) mice showed no difference, whereas mCherry (R) mice developed an increase

concomitant with the display of resilience (Fig. 4C-E), not observed in mice enduring only 7

SDEs. These findings suggest that the persistence of neural activity across SDEs #8-10 is

associated with resiliency.

During SI Test 1, similar to control mice, DREADDs injected mice experienced a decreased in

calcium-related neural activity upon initiating social interaction with a novel conspecific

(Supplemental Data Fig 7A), whose neural pattern was observed in both no target and target

trials (Supplemental Data Fig 7B). In hM4Di-injected mice, there were no significant differences

in neural activity during SI Test 1 and 2 trials, similar to control mCherry (S) mice (Fig. 4F-H;

Supplemental Data Fig. 7A-C). As was observed in the resilient (mCherry (R)) mice, the

hM3Dq-injected group showed an increase in calcium-encoded neural activity (Fig 4F-H). Mice

injected with cre-dependent hM4Di showed a significant decrease in calcium-related neural

activity. In contrast, social interaction initiation led to increased activity in DIO-hM3Dq injected

mice (Fig. 4F-H). There were no significant differences in SI of mice subjected to 7 SDEs, yet

differences emerge following 3 additional SDEs with a subset of mice becoming susceptible and

resilient (mCherry(R)/(S)) or activating/inhibiting DREADD promoting resiliency and

susceptibility in the direction predicted (Fig. 4I). We observed a strong correlation between SI

ratio and calcium activity only after 10 SDEs (Fig. 4J). There were no significant differences

observed in calcium-based neuronal activity upon social interaction with a novel conspecific in

mice subjected to 7 SDEs (Supplemental Data Fig. 7A,B), or in the absence of social interaction

(Supplemental Data Fig. 7C). In contrast, after 10 SDEs, mCherry (R) and DIO-hM3Dq injected

mice displayed significantly greater neuronal activation upon social contact relative to mCherry

(S) and DIO-hM4Di injected mice (Supplemental Data Fig. 7D). Despite the difference in time

spent interacting with a novel conspecific, the number of interaction zone entries were not

significantly different, nor differences in distance traveled (Fig. 4K, Supplemental Data Fig. 7E).

These data suggest that BNSTov^{CRF} neuronal dynamics are differentially altered by stress

modulation in accordance with phenotypical display resiliency/susceptibility.

***Crfr1* expression in BNSTov^{CRF} neurons mirrors the behavioral emergence of resilience.**

We observed a stress-induced enhancement in firing rates of BNSTov^{CRF} neurons in resilient

mice. To explore the effect of this BNSTov^{CRF} stress modulation on CRF receptor transmission,

we used RNAScope *in situ* hybridization to quantify *Crfr1* and *Crfr2* in accordance with stress

history. Mice were subjected to either 7- or 10-daily episodes of social defeat, and the BNST

and *Crfr1* and *Crfr2* genes were assessed (Fig. 5A-C) due to their reported role in mediating

stress responses[30,58–60]. The percentage of CRFR1-expressing neurons among CRF-

expressing neurons was higher in mice subjected to 7 SDEs than in susceptible mice but not

significantly different than resilient mice (Fig. 5D). In contrast, there were no significant

differences in BNSTov neurons co-expressing *Crfr2 and Crf mRNA* across groups of mice (Fig.

5E). The overlap between *Crfr1* and *Crf* in the BNST was significantly greater in the oval

nucleus than in anterolateral, anteromedial, and ventral subregions of the anterior dorsal BNST

(Fig. 5C,F).

To explore the role of firing rate changes on gene expression and the development of resiliency,

we optogenetically stimulated BNSTov^{CRF} neurons using transgenic *Crf-Cre*::ai32 mice, which

express channelrhodopsin-2 (ChR2) in *Crf*-containing neurons [61] (Fig. 5G). For control

experiments, transgenic mice of a similar background were used, except instead of ai32, mice

expressed the tdTomato fluorophore in CRF neurons (*Crf-tdTomato*). 5 Hz stimulation

frequency was used in order to mirror the average firing rate observed in the spontaneous firing

rate of resilient mice (Fig. 2D). Mice received 15-minute 5 Hz photo-stimulation of BNSTov^{CRF}

neurons following physical stress on SDEs #8-10. Mice were placed in the adjacent

compartment separating the aggressor, preventing physical contact. The compartment divides

the aggressor cage in half by a clear plexiglass that allows for continuous sensory cues (Fig.

5H,I). Photo-stimulation of *Crf*::ChR2 mice led to a significantly higher SI ratio and greater

percentage resilient than *Crf::tdTomato* mice (89% vs. 33%) when stimulated during SDEs #8-

10 (Fig. 5J). Surprisingly, *Crf::ChR2* mice that received photo-stimulation but were not

subjected to SDEs #8-10 showed a significant decrease in SI ratio and 100% of the mice,

becoming susceptible (0/6, SI ≧ 1.0) (Fig. 5J). Photo-stimulation paired with SDEs #8-10

increased the percentage of cells co-expressing *Crfr1* mRNA in CRF+ neurons relative to mice

that experienced photo-stimulation in the absence of additional stress (Fig. 5K, L). Additionally,

we obtained slice preparation from the optogenetically-induced resilient mice, and performed

cell-attached single unit recordings from BNSTov ChR2-expressing CRF neurons

(Supplemental Data Fig. 8A). Our recording data showed that optogenetic stimulation reliably

induced 2.5 Hz, 5 Hz and 10 Hz spike responses as expected (Supplemental Data Fig. 8B), and

interestingly triggered burst firing after the 5 Hz optical stimulation (Supplemental Data Fig.

8C,D). Importantly, Bath applied CRFR1-selective antagonist NBI 27914 significantly decreased

the firing rate of tested BNSTov ChR2-expressing CRF neurons (Supplemental Data Fig. 8E,F).

In summary, these data show that maintenance of resiliency requires social defeat stress

BNSTov^{CRF} activation and is correlated with the upregulation of *Crfr1* expression in a stress-

history dependent manner.

Discussion

The RSDS paradigm was modified to observe the effects of cumulative stress on neuroplasticity

in regions critical for mood regulation. Indeed, our work has uncovered a discrete window of

neuronal and behavioral plasticity between 7 and 10 SDEs, during which susceptible and

resilient phenotypes are established. By capturing behavioral, electrophysiological, and *in vivo*

fiber photometric measures during the intra-social defeat stress period, we uncover the

mechanisms underlying the establishment of resilience.

The oval nucleus of the BNST has been investigated in stress responses, we hypothesized that

the region would be instrumental in processing contexts associated with social stress. Indeed,

we observed that individual differences of stress effects on social behavior are encoded by

BNSTov^{CRF} neurons. Prior studies have implicated the overactivation of CRF neurons in the

BNST as being pro-depressive and anxiogenic[31,39,46,47,49], therefore, we were surprised to

observe that BNSTov^{CRF} neurons exhibited increased activity in resilient mice. When these

same cells were active, they yielded resilient mice. Indeed, we observed that Cre-dependent

hM3Dq action produced resilient mice only after a certain "dose" of daily stressors (7 SDEs),

suggesting that the BNSTov may be tightly modulated based on stress history. Our findings

support a view that resilience is a stress history-dependent state, which does not exist before

stress. This is highly consistent with previous demonstrations that resilience is a status

achieved by active regulation of genes and ion channel functions in this group more than in

susceptible animals[20-21,62–67].

CRF neurons have been observed to influence the salience of stressful contexts according to

stress exposure[24,27,60,64,68]. One feasible mechanism by which this occurs is through CRF

receptor dynamics. We hypothesized that the development of resilience occurs via CRF-CRFR

interactions in the BNST. Surprisingly, we observed *Crfr1* mRNA expression in CRF (and not

neighboring CRF⁻) neurons, suggesting that stress works as a pro-resilient agent based on

stress-history. This is in contrary to what has been observed, namely that CRFR1 has been

found largely on non-CRF neurons in the BNST.[68,69]

CRFR1 is selectively activated in the context of ongoing stress, serving as a coincidence-

detector [70]. Chronic stress has been shown to shift the connectivity of local CRF⁺ neurons from

CRF⁺-CRF⁻ to a larger percentage of CRF⁺-CRF⁺ cells[73]. Studies using prolonged

overactivation (over weeks to months) of CRF activity have yielded antidepressant and

anxiolytic results[29,34,35]. Here, we identify a more discrete timeline in the span of days to capture

the transition of when BNSTov^{CRF} activation becomes pro-resilient. By optogenetically

activating BNSTov^{CRF} neurons, we observed an increase in CRFR1 expression. Though

correlative, this exquisite regulation of CRFR1 according to stress-history and at times

demonstrating an opposing effect on depressive-like behavior, may underlie why clinical trials of

CRFR1 antagonists for MDD have been met with variable success[59,74−76]. Although beyond the

scope of the current study, future experiments using siRNA or CRISPR knockdown approaches

should be performed to determine whether upregulation of CRFR1 is a critical facet of

neuroadaptation leading to resiliency.

Stress-sensitive regions such as the BNST have been found to be of critical importance in

stress coping and reactivity[77−79]. Stress resilience has long been considered a response

separate from or in the absence of stimuli that gives rise to stress susceptibility, mediated by

parallel circuits or cell-types in a particular brain region[14,15,80]. Here, we observe that activity

dynamics of CRF neurons can shape and influence resiliency to stress, potentially through

(auto)regulation of *Crfr1* mRNA. Other substrates may be used to classify these neurons either

on the basis of dual-neuropeptide or circuit-specific identities and should be the object of future

exploration.

Previous work has shown that resiliency is influenced by dopaminergic VTA neurons in the

nucleus accumbens (NAc)[29], in part, through the actions of brain-derived-neurotrophic-factor

(BDNF)[21,81,82]. CRF peptide has been important for BDNF release in the NAc as a stress-

coincidence sensor[20], yet the sources of CRF important for altering stress effect on social and

hedonic behavior has not been extensively characterized. While long-range GABAergic BNST

neurons projecting to the VTA have been shown to influence reward and anxiety-like

behavior[48,83–86], it is unclear to what degree these cells comprise the oval nuclear BNST

population. The BNST also sends projections to the dorsal raphe, lateral and paraventricular

hypothalamus, and ventrolateral periaqueductal gray[27,31–33,42,77,87–89], among others, whose

modulation have been linked to stress on affect and social motivation. In this way, the BNST

acts as a node for integrating information regarding stress history and determining socio-

affective outcomes according, possibly due to CRFR1 receptor dynamics occurring on CRF

neurons of the oval nucleus, thereby shaping the long-lasting outcome of resiliency.

Our study highlights a previously unknown mechanism by which the BNST encodes cumulative

social stress and effectuates susceptible or resilient outcomes. Importantly, there are currently

no Food and Drug Administration-approved drugs aimed at preventing a depressive episode

from occurring. By targeting mechanisms involved in establishing resiliency, the possibility may

exist to therapeutically leverage windows of plasticity to effectuate resilience and evade the

development of MDD.

Statistical analysis

Animals were randomly assigned to control and experimental groups and all experimenters

were blinded. Mice were excluded if viral infection was off-target. No data was excluded for

other reasons. Student's two-tailed t-tests were used for comparisons of two experimental

groups. For parametric data sets, comparisons among three or more groups were performed

using one or two-way ANOVA tests followed by Tukey's or Bonferroni posthoc tests. For all

tests, $p<0.05$ was determined to be significant. Statistical analyses were performed using

Graph Pad Prism 9 .3.1 software (La Jolla, CA, USA). For data not normally distributed, non-

parametric analyses were performed.

Data & Code availability

All source data for this manuscript and MATLAB code used to analyze photometry data is

available upon request.

348
349 **Acknowledgements**
350
We would like to thank S.M. for generous sharing for input on the manuscript. We would like to

thank S.R. and L.L. for lending viral constructs for behavioral validation. We would like to thank

the animal care staff at ISMMS for the animal care staff and technical support. This research

was funded by the National Institutes of Health grants R01MH072908 to DGR and LJY,

R01MH1206387 to MHH and SJR, R21MH112081 to MHH and SJR, F31MH114624 to SEH,

and P51 OD011132 to YNPRC, by the National Key R&D Program of 1034 China No.

2021ZD0202900 to MHH.

Financial Disclosures

H.S.M. receives consulting and IP licensing fees from Abbott Neuromodulation.

All other authors report no biomedical financial interests or potential conflicts of interest.

References

363   1.   Veer, I. M. *et al.* Psycho-social factors associated with mental resilience in the Corona

364   lockdown. *Transl. Psychiatry* **11**, 67 (2021).

365   2.   Post, R. M. Transduction of psychosocial stress into the neurobiology of recurrent

366   affective disorder. *Am. J. Psychiatry* **149**, 999–1010 (1992).

367   3.   Gold, P. W. Evidence that stress per se has a role in the precipitation and natural history

368   of depressive illness. *Mol. Psychiatry* **18**, 954–956 (2013).

369   4.   Kendler, K. S., Karkowski, L. M. & Prescott, C. A. Causal relationship between stressful life

370   events and the onset of major depression. *Am. J. Psychiatry* **156**, 837–841 (1999).

371   5.   Barthas, F. *et al.* Cumulative Effects of Social Stress on Reward-Guided Actions and

372   Prefrontal Cortical Activity. *Biol. Psychiatry* **88**, 541–553 (2020).

373   6.   Golden, S. A., Covington, H. E., Berton, O. & Russo, S. J. A standardized protocol for

374   repeated social defeat stress in mice. *Nat. Protoc.* **6**, 1183–1191 (2011).

375   7.   Martinez, M., Phillips, P. J. & Herbert, J. Adaptation in patterns of *c-fos* expression in the

376   brain associated with exposure to either single or repeated social stress in male rats.

377   *European Journal of Neuroscience* **10**, 20–33 (1998).

378   8.   Zheng, Z. *et al.* Hypothalamus-habenula potentiation encodes chronic stress experience

379   and drives depression onset. *Neuron* (2022) doi:10.1016/j.neuron.2022.01.011.

9.   Kudryavtseva, N. N., Bakshtanovskaya, I. V. & Koryakina, L. A. Social model of depression

in mice of C57BL/6J strain. *Pharmacol. Biochem. Behav.* **38**, 315–320 (1991).

10.   Laine, M. A. *et al.* Brain activation induced by chronic psychosocial stress in mice. *Sci.*

Rep. **7**, 15061 (2017).

11.   Hultman, R. *et al.* Brain-wide Electrical Spatiotemporal Dynamics Encode Depression

Vulnerability. *Cell* **173**, 166-180.e14 (2018).

12.   Drysdale, A. T. *et al.* Resting-state connectivity biomarkers define neurophysiological

subtypes of depression. *Nat. Med.* **23**, 28–38 (2017).

13.   Samuels, B. A. *et al.* Modeling treatment-resistant depression. *Neuropharmacology* **61**,

408–413 (2011).

14.   Franklin, T. B., Saab, B. J. & Mansuy, I. M. Neural mechanisms of stress resilience and

vulnerability. *Neuron* **75**, 747–761 (2012).

15.   Russo, S. J., Murrough, J. W., Han, M.-H., Charney, D. S. & Nestler, E. J. Neurobiology of

resilience. *Nat. Neurosci.* **15**, 1475–1484 (2012).

16.   Karatsoreos, I. N. & McEwen, B. S. Psychobiological allostasis: resistance, resilience and

vulnerability. *Trends Cogn Sci (Regul Ed)* **15**, 576–584 (2011).

17.   Laine, M. A. *et al.* Genetic Control of Myelin Plasticity after Chronic Psychosocial Stress.

eNeuro **5**, (2018).

18.   Markham, C. M., Norvelle, A. & Huhman, K. L. Role of the bed nucleus of the stria

terminalis in the acquisition and expression of conditioned defeat in Syrian hamsters.

Behav. Brain Res. **198**, 69–73 (2009).

19.   Gururajan, A., Bastiaanssen, T. F. S., Ventura Silva, A. P., Moloney, G. M. & Cryan, J. F.

The impact of psychosocial defeat stress on the BNST transcriptome in adult male mice.

Eur. J. Neurosci. (2021) doi:10.1111/ejn.15567.

20.   Walsh, J. J. *et al.* Stress and CRF gate neural activation of BDNF in the mesolimbic reward

pathway. *Nat. Neurosci.* **17**, 27–29 (2014).

21.   Krishnan, V. *et al.* Molecular adaptations underlying susceptibility and resistance to

social defeat in brain reward regions. *Cell* **131**, 391–404 (2007).

22.   Matsuda, S. *et al.* Persistent c-fos expression in the brains of mice with chronic social

stress. *Neurosci. Res.* **26**, 157–170 (1996).

23.   Berton, O. *et al.* Essential role of BDNF in the mesolimbic dopamine pathway in social

defeat stress. *Science* **311**, 864–868 (2006).

24.   Waraczynski, M. Toward a systems-oriented approach to the role of the extended

amygdala in adaptive responding. *Neurosci. Biobehav. Rev.* **68**, 177–194 (2016).

25.   Flanigan, M. E. & Kash, T. L. Coordination of social behaviors by the bed nucleus of the

stria terminalis. *Eur. J. Neurosci.* (2020) doi:10.1111/ejn.14991.

26.   Lebow, M. A. & Chen, A. Overshadowed by the amygdala: the bed nucleus of the stria

terminalis emerges as key to psychiatric disorders. *Mol. Psychiatry* **21**, 450–463 (2016).

27.   Daniel, S. E. & Rainnie, D. G. Stress modulation of opposing circuits in the bed nucleus of

the stria terminalis. *Neuropsychopharmacology* **41**, 103–125 (2016).

28.   Jennings, J. H. *et al.* Distinct extended amygdala circuits for divergent motivational

states. *Nature* **496**, 224–228 (2013).

29.   Dedic, N. *et al.* Chronic CRH depletion from GABAergic, long-range projection neurons in

the extended amygdala reduces dopamine release and increases anxiety. *Nat. Neurosci.*

21, 803–807 (2018).

30.   Chen, Y., Molet, J., Gunn, B. G., Ressler, K. & Baram, T. Z. Diversity of Reporter

Expression Patterns in Transgenic Mouse Lines Targeting Corticotropin-Releasing

Hormone-Expressing Neurons. *Endocrinology* **156**, 4769–4780 (2015).

31.   Dabrowska, J., Martinon, D., Moaddab, M. & Rainnie, D. G. Targeting Corticotropin-

Releasing Factor Projections from the Oval Nucleus of the Bed Nucleus of the Stria

Terminalis Using Cell-Type Specific Neuronal Tracing Studies in Mouse and Rat Brain. *J.*

Neuroendocrinol. **28**, (2016).

32.   Matthews, G. A. *et al.* Dorsal raphe dopamine neurons represent the experience of

social isolation. *Cell* **164**, 617–631 (2016).

33.   Marcinkiewcz, C. A. *et al.* Serotonin engages an anxiety and fear-promoting circuit in the

extended amygdala. *Nature* **537**, 97–101 (2016).

34.   Regev, L. *et al.* Prolonged and site-specific over-expression of corticotropin-releasing

factor reveals differential roles for extended amygdala nuclei in emotional regulation.

Mol. Psychiatry **16**, 714–728 (2011).

35.   Sink, K. S. *et al.* Effects of continuously enhanced corticotropin releasing factor

expression within the bed nucleus of the stria terminalis on conditioned and

unconditioned anxiety. *Mol. Psychiatry* **18**, 308–319 (2013).

36.   Konishi, S., Kasagi, Y., Katsumata, H., Minami, S. & Imaki, T. Regulation of corticotropin-

releasing factor (CRF) type-1 receptor gene expression by CRF in the hypothalamus.

Endocr. J. **50**, 21–36 (2003).

37.   Jasnow, A. M., Banks, M. C., Owens, E. C. & Huhman, K. L. Differential effects of two

corticotropin-releasing factor antagonists on conditioned defeat in male Syrian hamsters

(Mesocricetus auratus). *Brain Res.* **846**, 122–128 (1999).

38.   Avery, S. N., Clauss, J. A. & Blackford, J. U. The human BNST: functional role in anxiety

and addiction. *Neuropsychopharmacology* **41**, 126–141 (2016).

39.   Kim, S.-Y. *et al.* Diverging neural pathways assemble a behavioural state from separable

features in anxiety. *Nature* **496**, 219–223 (2013).

40.   Rodríguez-Sierra, O. E., Goswami, S., Turesson, H. K. & Pare, D. Altered responsiveness of

453   BNST and amygdala neurons in trauma-induced anxiety. *Transl. Psychiatry* **6**, e857

454   (2016).

455   41.   Ventura-Silva, A. P. *et al.* Stress shifts the response of the bed nucleus of the stria

456   terminalis to an anxiogenic mode. *Eur. J. Neurosci.* **36**, 3396–3406 (2012).

457   42.   Giardino, W. J. *et al.* Parallel circuits from the bed nuclei of stria terminalis to the lateral

458   hypothalamus drive opposing emotional states. *Nat. Neurosci.* **21**, 1084–1095 (2018).

459   43.   Duvarci, S., Bauer, E. P. & Paré, D. The bed nucleus of the stria terminalis mediates inter-

460   individual variations in anxiety and fear. *J. Neurosci.* **29**, 10357–10361 (2009).

461   44.   Duque-Wilckens, N. *et al.* Extrahypothalamic oxytocin neurons drive stress-induced

462   social vigilance and avoidance. *Proc Natl Acad Sci USA* **117**, 26406–26413 (2020).

463   45.   Dabrowska, J. *et al.* Striatal-enriched protein tyrosine phosphatase-STEPs toward

464   understanding chronic stress-induced activation of corticotrophin releasing factor

465   neurons in the rat bed nucleus of the stria terminalis. *Biol. Psychiatry* **74**, 817–826

466   (2013).

467   46.   Hu, P. *et al.* Chronic Stress Induces Maladaptive Behaviors by Activating Corticotropin-

468   Releasing Hormone Signaling in the Mouse Oval Bed Nucleus of the Stria Terminalis. *J.*

469   *Neurosci.* **40**, 2519–2537 (2020).

47.   Daniel, S. E. *et al.* Chronic stress induces cell type-selective transcriptomic and

electrophysiological changes in the bed nucleus of the stria terminalis.

Neuropharmacology **150**, 80–90 (2019).

48.   Wanat, M. J., Bonci, A. & Phillips, P. E. M. CRF acts in the midbrain to attenuate

accumbens dopamine release to rewards but not their predictors. *Nat. Neurosci.* **16**,

383–385 (2013).

49.   Salimando, G. J., Hyun, M., Boyt, K. M. & Winder, D. G. BNST GluN2D-Containing NMDA

Receptors Influence Anxiety- and Depressive-like Behaviors and ModulateCell-Specific

Excitatory/Inhibitory Synaptic Balance. *J. Neurosci.* **40**, 3949–3968 (2020).

50.   Hartley, N. D. *et al.* Dynamic remodeling of a basolateral-to-central amygdala

glutamatergic circuit across fear states. *Nat. Neurosci.* **22**, 2000–2012 (2019).

51.   Sanford, C. A. *et al.* A Central Amygdala CRF Circuit Facilitates Learning about Weak

Threats. *Neuron* **93**, 164–178 (2017).

52.   Rodríguez-Sierra, O. E., Turesson, H. K. & Pare, D. Contrasting distribution of

physiological cell types in different regions of the bed nucleus of the stria terminalis. *J.*

Neurophysiol. **110**, 2037–2049 (2013).

53.   Gungor, N. Z. & Paré, D. Functional heterogeneity in the bed nucleus of the stria

terminalis. *J. Neurosci.* **36**, 8038–8049 (2016).

54.   Zhan, J., Komal, R., Keenan, W. T., Hattar, S. & Fernandez, D. C. Non-invasive Strategies

for Chronic Manipulation of DREADD-controlled Neuronal Activity. *J. Vis. Exp.* (2019)

doi:10.3791/59439.

55.   Schalbetter, S. M. *et al.* Oral application of clozapine-N-oxide using the micropipette-

guided drug administration (MDA) method in mouse DREADD systems. *Lab Anim (NY)*

50, 69–75 (2021).

56.   Wess, J., Nakajima, K. & Jain, S. Novel designer receptors to probe GPCR signaling and

physiology. *Trends Pharmacol. Sci.* **34**, 385–392 (2013).

57.   Walker, D. L., Miles, L. A. & Davis, M. Selective participation of the bed nucleus of the

stria terminalis and CRF in sustained anxiety-like versus phasic fear-like responses. *Prog*

Neuropsychopharmacol Biol Psychiatry **33**, 1291–1308 (2009).

58.   Tran, L., Schulkin, J. & Greenwood-Van Meerveld, B. Importance of CRF receptor-

mediated mechanisms of the bed nucleus of the stria terminalis in the processing of

anxiety and pain. *Neuropsychopharmacology* **39**, 2633–2645 (2014).

59.   Lee, Y., Fitz, S., Johnson, P. L. & Shekhar, A. Repeated stimulation of CRF receptors in the

BNST of rats selectively induces social but not panic-like anxiety.

Neuropsychopharmacology **33**, 2586–2594 (2008).

60.   Dedic, N. *et al.* Deletion of CRH From GABAergic Forebrain Neurons Promotes Stress

Resilience and Dampens Stress-Induced Changes in Neuronal Activity. *Front. Neurosci.*

13, 986 (2019).

61.   Wang, Y. *et al.* Single-cell morphological characterization of CRH neurons throughout the

whole mouse brain. *BMC Biol.* **19**, 47 (2021).

62.   Chen, P. *et al.* Prefrontal Cortex Corticotropin-Releasing Factor Neurons Control

Behavioral Style Selection under Challenging Situations. *Neuron* **106**, 301-315.e7 (2020).

63.   Elliott, E., Ezra-Nevo, G., Regev, L., Neufeld-Cohen, A. & Chen, A. Resilience to social

stress coincides with functional DNA methylation of the Crf gene in adult mice. *Nat.*

Neurosci. **13**, 1351–1353 (2010).

64.   Lemos, J. C. *et al.* Severe stress switches CRF action in the nucleus accumbens from

appetitive to aversive. *Nature* **490**, 402–406 (2012).

65.   Zhang, H. *et al.* α1- and β3-Adrenergic Receptor-Mediated Mesolimbic Homeostatic

Plasticity Confers Resilience to Social Stress in Susceptible Mice. *Biol. Psychiatry* **85**, 226–

236 (2019).

66.   Friedman, A. K. *et al.* Enhancing depression mechanisms in midbrain dopamine neurons

achieves homeostatic resilience. *Science* **344**, 313–319 (2014).

67.   Bagot, R. C. *et al.* Circuit-wide Transcriptional Profiling Reveals Brain Region-Specific

Gene Networks Regulating Depression Susceptibility. *Neuron* **90**, 969–983 (2016).

68.   Dabrowska, J., Hazra, R., Guo, J.-D., Dewitt, S. & Rainnie, D. G. Central CRF neurons are

525   not created equal: phenotypic differences in CRF-containing neurons of the rat

526   paraventricular hypothalamus and the bed nucleus of the stria terminalis. *Front.*

527   *Neurosci.* **7**, 156 (2013).

528   69.   Justice, N. J., Yuan, Z. F., Sawchenko, P. E. & Vale, W. Type 1 corticotropin-releasing

529   factor receptor expression reported in BAC transgenic mice: implications for reconciling

530   ligand-receptor mismatch in the central corticotropin-releasing factor system. *J. Comp.*

531   *Neurol.* **511**, 479–496 (2008).

532   70.   Ramot, A. *et al.* Hypothalamic CRFR1 is essential for HPA axis regulation following

533   chronic stress. *Nat. Neurosci.* **20**, 385–388 (2017).

534   71.   McClard, C. K. *et al.* POU6f1 Mediates Neuropeptide-Dependent Plasticity in the Adult

535   Brain. *J. Neurosci.* **38**, 1443–1461 (2018).

536   72.   Hillhouse, E. W. & Grammatopoulos, D. K. The molecular mechanisms underlying the

537   regulation of the biological activity of corticotropin-releasing hormone receptors:

538   implications for physiology and pathophysiology. *Endocr. Rev.* **27**, 260–286 (2006).

539   73.   Partridge, J. G. *et al.* Stress increases GABAergic neurotransmission in CRF neurons of

540   the central amygdala and bed nucleus stria terminalis. *Neuropharmacology* **107**, 239–

541   250 (2016).

74. Spierling, S. R. & Zorrilla, E. P. Don't stress about CRF: assessing the translational failures

of CRF1antagonists. *Psychopharmacology (Berl)* **234**, 1467–1481 (2017).

75. Waters, R. P. *et al.* Evidence for the role of corticotropin-releasing factor in major

depressive disorder. *Neurosci. Biobehav. Rev.* **58**, 63–78 (2015).

76. Reul, J. Corticotropin-releasing factor receptors 1 and 2 in anxiety and depression. *Curr.*

Opin. Pharmacol. **2**, 23–33 (2002).

77. Johnson, S. B. *et al.* A basal forebrain site coordinates the modulation of endocrine and

behavioral stress responses via divergent neural pathways. *J. Neurosci.* **36**, 8687–8699

(2016).

78. Crestani, C. C., Alves, F. H. F., Correa, F. M. A., Guimarães, F. S. & Joca, S. R. L. Acute

reversible inactivation of the bed nucleus of stria terminalis induces antidepressant-like

effect in the rat forced swimming test. *Behav. Brain Funct.* **6**, 30 (2010).

79. Fiedler, D. *et al.* Brain-Derived Neurotrophic Factor/Tropomyosin Receptor Kinase B

Signaling Controls Excitability and Long-Term Depression in Oval Nucleus of the BNST. *J.*

Neurosci. **41**, 435–445 (2021).

80. Cathomas, F., Murrough, J. W., Nestler, E. J., Han, M.-H. & Russo, S. J. Neurobiology of

resilience: interface between mind and body. *Biol. Psychiatry* **86**, 410–420 (2019).

81. Koo, J. W., Chaudhury, D., Han, M.-H. & Nestler, E. J. Role of Mesolimbic Brain-Derived

Neurotrophic Factor in Depression. *Biol. Psychiatry* **86**, 738–748 (2019).

82. Wook Koo, J. *et al.* Essential Role of Mesolimbic Brain-Derived Neurotrophic Factor in

Chronic Social Stress-Induced Depressive Behaviors. *Biol. Psychiatry* **80**, 469–478 (2016).

83. Takahashi, D. *et al.* Tonic Suppression of the Mesolimbic Dopaminergic System by

Enhanced Corticotropin-Releasing Factor Signaling Within the Bed Nucleus of the Stria

Terminalis in Chronic Pain Model Rats. *J. Neurosci.* **39**, 8376–8385 (2019).

84. Caillé, S., Guillem, K., Cador, M., Manzoni, O. & Georges, F. Voluntary nicotine

consumption triggers in vivo potentiation of cortical excitatory drives to midbrain

dopaminergic neurons. *J. Neurosci.* **29**, 10410–10415 (2009).

85. Silberman, Y., Matthews, R. T. & Winder, D. G. A corticotropin releasing factor pathway

for ethanol regulation of the ventral tegmental area in the bed nucleus of the stria

terminalis. *J. Neurosci.* **33**, 950–960 (2013).

86. Rodaros, D., Caruana, D. A., Amir, S. & Stewart, J. Corticotropin-releasing factor

projections from limbic forebrain and paraventricular nucleus of the hypothalamus to

the region of the ventral tegmental area. *Neuroscience* **150**, 8–13 (2007).

87. Dong, H. W., Petrovich, G. D., Watts, A. G. & Swanson, L. W. Basic organization of

projections from the oval and fusiform nuclei of the bed nuclei of the stria terminalis in

adult rat brain. *J. Comp. Neurol.* **436**, 430–455 (2001).

88. Maita, I., Bazer, A., Blackford, J. U. & Samuels, B. A. Functional anatomy of the bed

nucleus of the stria terminalis-hypothalamus neural circuitry: Implications for valence

surveillance, addiction, feeding, and social behaviors. *Handb. Clin. Neurol.* **179**, 403–418

(2021).

89.   Kaouane, N., Ada, S., Hausleitner, M. & Haubensak, W. Dorsal Bed Nucleus of the Stria

583   Terminalis-Subcortical Output Circuits Encode Positive Bias in Pavlovian Fear and

584   Reward. *Front. Neural Circuits* **15**, 772512 (2021).

585   90.   Cao, J.-L. *et al.* Mesolimbic dopamine neurons in the brain reward circuit mediate

586   susceptibility to social defeat and antidepressant action. *J. Neurosci.* **30**, 16453–16458

587   (2010).

588   91.   Morel, C. *et al.* Midbrain projection to the basolateral amygdala encodes anxiety-like but

589   not depression-like behaviors. *Nat. Commun.* **13**, 1532 (2022).

590   92.   Kumar, S. *et al.* Prefrontal cortex reactivity underlies trait vulnerability to chronic social

591   defeat stress. *Nat. Commun.* **5**, 4537 (2014).

592   93.   Friedman, A. K. *et al.* KCNQ channel openers reverse depressive symptoms via an active

593   resilience mechanism. *Nat. Commun.* **7**, 11671 (2016).

594   94.   Chaudhury, D. *et al.* Rapid regulation of depression-related behaviours by control of

595   midbrain dopamine neurons. *Nature* **493**, 532–536 (2013).

596   95.   Xiao, Q. *et al.* A new GABAergic somatostatin projection from the BNST onto accumbal

597   parvalbumin neurons controls anxiety. *Mol. Psychiatry* **26**, 4719–4741 (2021).

598   96.   Cooper, M. A. & Huhman, K. L. Blocking corticotropin-releasing factor-2 receptors, but

599   not corticotropin-releasing factor-1 receptors or glucocorticoid feedback, disrupts the

600   development of conditioned defeat. *Physiol. Behav.* **101**, 527–532 (2010).

601   97.   Challis, C. *et al.* Raphe GABAergic neurons mediate the acquisition of avoidance after

602   social defeat. *J. Neurosci.* **33**, 13978–88, 13988a (2013).

603   98.   Challis, C., Beck, S. G. & Berton, O. Optogenetic modulation of descending

604   prefrontocortical inputs to the dorsal raphe bidirectionally bias socioaffective choices

605   after social defeat. *Front. Behav. Neurosci.* **8**, 43 (2014).

606   99.   Calipari, E. S. *et al.* Dopaminergic dynamics underlying sex-specific cocaine reward. *Nat.*

607   *Commun.* **8**, 13877 (2017).

Repeated Social Defeat Stress establishes onset of depression-related behaviour

Figure 1

Firing rate alterations in BNSTov^CRF neurons occur as adaptation to social stress, persisting in resilient, not susceptible mice.

Figure 2

Chemogenetic Modulation of BNSTov^CRF Neurons Recapitulates Behavioural Tipping Point

Figure 3

BNSTov^{CRF} calcium-dynamics encode stress effect on social interaction

Figure 4

BNSTov *Crfr1* is associated with the emergence of resiliency

Figure 5

Figure 1. Susceptible and resilient subgroups emerge between 7 and 10 daily episodes of social defeat stress.

A. Experimental design for repeated social defeat stress (RSDS). **B.** Experimental timeline of social defeat stress and behavioral tests: social interaction (SI) and sucrose preference (SP). **C.** Social interaction test schema involving target and no target trials. **D.** Sucrose preference test schematic. **E.** Effect of cumulative social defeat stress on social interaction measured as SI ratios, after 1, 4, 7, and 10 social defeat episodes (SDEs) (n=11-18 mice/group), two-way ANOVA interaction $F_{(6,134)}=5.783$, ****$P<0.0001$, row factor $F_{(3,134)}=12.11$, ****$P<0.0001$, $F_{(2,134)}=4.086$, column factor *$P<0.05$, Tukey's post-hoc test susceptible vs resilient ****$P<0.0001$, susceptible vs control **$P<0.01$, resilient vs control $P=0.2751$. SI test susceptible (SI test stressed x7 vs stressed x10) ****$P<0.0001$. **F.** Aggregated data on social interaction test across experiments. One-way ANOVA treatment $F_{(3,109)}=14.61$, ****$P<0.0001$. Tukey's post-hoc test control vs stressed x7 *$P<0.05$, control vs susceptible **$P<0.01$, stressed x7 vs susceptible ****$P<0.0001$, susceptible vs resilient ****$P<0.0001$ (n=25-37 mice). **G.** Sucrose preference test. One-way ANOVA treatment $F_{(3,93)}=24.06$, ****$P<0.0001$). Tukey's post-hoc test control vs stressed x7 $P=0.77$, control vs susceptible ****$P<0.0001$, control vs resilient $P=0.9387$, stressed x7 vs susceptible ****$P<0.0001$, stressed x7 vs resilient $P=0.984$, susceptible vs resilient ****$P<0.0001$ (n=23-25 mice). **H.** Distribution of stressed mice who underwent 7 SDEs. **I.** Distribution of mice who underwent 10 SDEs and sorted into susceptible and resilient mice. **J.** Cumulative distribution of all stressed mice. Kolmogorov-Smirnov (distance) 0.2054 Susceptible vs Resilient, ****$P<0.0001$. **K.** Time spent interaction socially with a novel conspecific. One-Way ANOVA

$F_{(2,201)}=76.63$, ****$P<0.0001$. Tukey's post-hoc susceptible vs resilient ***$P<0.001$, susceptible vs stressed x7 ****$P<0.0001$, stressed vs resilient $P=0.4984$ (n=72 susceptible, 64 resilient mice). All data represent means ± SEM. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P<0.0001$, ns = not significant.

Figure 2. Firing rate alterations in BNSTov^{CRF} neurons occur as adaptation to social stress, persisting in resilient, not susceptible mice.

A. Mouse genotype and timeline of cell-attached electrophysiology experiments. **B.** Fluorescence-guided cell-attached electrophysiology setup, brain slice of the BNST (CRF cells, tdTomato) and DIC image of CRF neurons, scale bar 0.63 mm. **C.** Representative trace of BNSTov^{CRF} positive and negative neurons of control, stressed (SDEs x 7), susceptible, and resilience mice. **D.** Firing rate of CRF$^+$ neurons (n=9-31 cells/group), ****$P<0.0001$, one-way ANOVA, Tukey's multiple comparison's test, control vs stressed (SDEs x 7) **$P<0.01$, susceptible vs resilient **$P<0.01$, susceptible vs stressed (SDEs x7), ****$P<0.0001$. Firing rate of CRF$^-$ neurons (n=7-15 cells per 4-6 mice/group), one-way NOVA, $F_{(3, 45)}=3.113$, *$P<0.05$, Tukey's multiple comparison's test, control vs stressed x7 $P=0.9192$, control vs susceptible $P=0.0769$, control vs resilient $P=0.1742$, stressed (SDEs x 7) vs susceptible $P=0.1370$, stressed x7 vs resilient $P=0.3219$, susceptible vs resilient $P=0.9673$. **E.** Correlation of firing rate with social interaction ratio: CRF$^+$, $R^2=0.5726$, *$P<0.05$; CRF$^-$, $R^2=0.0609$, $P=0.5219$. **F.** Representative sample of burst firing. **G.** Representative sample of tonic firing. **H.**

Percentage of bursting cells per animal group. **I.** Percentage of spikes within burst. **J.** Number of spikes per burst. **K.** Number of bursts per cell. (n=10-52 cells per 4-6 mice/group). All data represent means ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, N.S. = not significant.

Figure 3. Chemogenetic modulation of BNSTov^{CRF} neurons bidirectional recapitulates behavioural tipping point.

A. Viral targeting of BNSTov. **B.** Experimental design of chemogenetic manipulation of BNSTov neurons with CNO-drinking water construct. **C.** Social interaction test, two-way ANOVA $F_{(3,68)}$=18.01, row ****P<0.0001, row factor $F_{(3, 68)}$=9.965, P=0.2616 (time), row factor x time $F_{(1,68)}$=1.281, ****P<0.0001, Sidak's multiple comparisons test mCherry (s), **P=0.0017 (n=10 mice); mCherry (r), P=0.1322 (n=9 mice); hM4Di, ****P<0.0001 (n=10 mice); hM3Dq, **P=0.0018 (n=9 mice). **D.** Sucrose preference test. one-way ANOVA treatment $F_{(3,27)}$=8.310, ***P<0.001, Tukey's post-hoc testing susceptible vs hM4Di, P=9619; susceptible vs resilient, **P<0.01; susceptible vs hM3Dq, **P<0.01; hM4Di vs resilient, **P<0.01; hM4Di vs hM3Dq, *P<0.05; resilient vs hM3Dq P=0.9368 (7-9 mice per group). **E.** Distance traveled, one-way ANOVA $F_{(3,27)}$=1.031, P=0.3944. **F.** Percentage of susceptible or resilient mice: DIO-hM4Di, 100% susceptible; DIO-hM3Dq, 89% resilient and 11% susceptible; mCherry, 40% resilient and 60% susceptible. Chi-square test=15.66, df=2, ***P<0.001 (n=21 mCherry, 10 hMDi, 9 hM3Dq); Fisher's post-hoc test mCherry vs hM4Di, *P<0.05; mCherry vs hM3Dq, *P<0.05; hM4Di vs hM3Dq,

****$P<0.0001$. **G.** Repeated social defeat DREADDs manipulation (4-7 episodes of stress), 2-way ANOVA treatment $F_{(1,16)}=0.01749$, $P=0.8964$; SI Test 1 vs SI Test 2, $F_{(1,16)}=0.2377$, $P=0.6325$; interaction $F_{(1,16)}=0.07055$, $P=0.6218$ (n=5-6 mice/group). **H.** RSDS DREADDs manipulation (7-10 episodes of stress), 2-way ANOVA treatment, $F_{(1,22)}=8.675$, ****$P< 0.01$; SI Test 1 vs SI Test 2, $F_{(1,22)}=2.549$, $P=0.1247$; interaction $F_{(1,22)}=2.549$, $P=0.1247$. Sidak's post-hoc test SI Test 1 vs SI Test 2, control, $P=0.5533$; DIO-hM3Dq, *$P<0.05$ (n=6-7 mice/group). **I.** RSDS DREADDs manipulation (10-13 episodes of stress), 2-way ANOVA $F_{(1,30)}=1.192$, treatment, $P=0.2623$. SI Test 1 vs Si Test 2, $F_{(1,30)}=0.3132$, $P=0.2836$. Interaction $F_{(1,30)}=0.2623$, $P=0.5799$. All data represent means ± SEM. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, ns = not significant.

Figure 4. BNSTov^{CRF} calcium-dynamics encode stress effect on social interaction

A. Viral targeting of the BNSTov. **B.** Experimental design of multiplexed chemogenetics with drinking water-CNO delivery and fiber photometry. **C.F.** Representative calcium recordings of mCherry susceptible (S), mCherry resilient (R), hM3Dq, and hM4Di groups, respectively. Vertical scale bar is equal to z-score of 1, and horizontal scale bar is equal to 10 s. **D. G.** Representative averaged trace centered around interaction bout for the four groups stated above. **E.** Two-way RM ANOVA row factor $F_{(3,14)}=1.401$; pre vs post-CNO $F_{(1,14)}=0.8179$; subject $F_{(14,14)}=1.195$; Row x SI Test 1/SI Test 2, $F_{(3, 14)}=9.343$, **$P<0.01$. Sidak's post-hoc test, susceptible vs resilient, $P=0.137$ vs *$P=0.05$, respectively (n=4 mice/group). **H.** hM4Di vs hM3Dq, $P=0.3069$ vs *$P<0.05$, respectively

(n=4-5 mice/grp). **I.** SI Test 1 social interaction test. One-way ANOVA $F_{(3,16)}=0.291$, P=0.8308 (n=5 mice/group). SI Test 2 social interaction test, One-Way ANOVA $F_{(3,16)}=17.83$, ****P<0.0001; Tukey's post-hoc test, susceptible vs resilient, ****P<0.0001; susceptible vs hM3Dq, ***P<0.001; susceptible vs hM4Di, P=0.0762; resilient vs hM3Dq, P=0.8506; resilient vs HM4Di, **P<0.01; hM3Dq vs hM4Di, *P<0.05 (n=5 mice/group). **J.** Correlation of SI ratio and z-score delta F/F upon social entry, simple linear regression pre-CNO $F_{(1,15)}=0.7674$, P=0.3948; SI test 2 $F_{(1,17)}=7.268$, *P<0.05. Intersection of lines, $F_{(1,32)}=6.896$, *P<0.05 (n=18 mice). **K.** Number of social interaction bouts, one-way ANOVA $F_{(3,16)}=1.346$, P=0.2949 (n=5 mice/group). All data represent means ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, ns = not significant.

Figure 5. BNSTov *Crfr1* is associated with the emergence of resiliency

A. Experimental timeline of RNAScope *in situ* hybridization of mice that underwent social defeat stress. **B**. Representative images of control, stressed x7, susceptible, and resilient. 20x magnification, scale bar 0.64 mm. **C.** Schematic of anterior dorsal BNST. **D.** *Crfr1* mRNA colocalization, one-way ANOVA $F_{(3,18)}=20.91$, ****P<0.0001; Tukey's posthoc test, control vs stressed x7, P=0.0701; control vs susceptible, P=0.3336; control vs resilient, ****P<0.0001; stressed x7 vs susceptible, **P<0.01; stressed x7 vs resilient, *P<0.05; susceptible vs resilient, ****P<0.0001 (n=4-6 BNST brain samples). **E.** *Crfr2* mRNA colocalization, one-way ANOVA $F_{(3,10)}=1.790$, P=0.2166 (n=3 BNST brain samples). **F.** RNAscope BNST subregion comparison, two-way ANOVA

interaction $F_{(6,24)}$=1.057, P=0.4147; row factor $F_{(2,24)}$=0.6092, P=0.5520; column factor $F_{(3,24)}$=1.549, P=0.2275 (n=3 BNST brain samples). **G.** Viral injection site. **H.** Experimental timeline. **I.** Optogenetic and social behavioral setup. **J.** Optogenetics social interaction, two-way ANOVA interaction $F_{(1,28)}$=5.059, *P<0.05; row factor $F_{(1,28)}$=1.061e-005, P=0.9974; column factor $F_{(1,28)}$=1.419, P=0.2436. Sidak's post-hoc test, *Crf*::tdT, *P<0.05; *Crf*::ChR2, P=0.7233 (n=7-9 mice/group). Optogenetic manipulation unpaired t-test, *Crf*::tdT, t=2.556, df=40, *P<0.05 (n=8-9 mice); *Crf*::ChR2, t=0.7758, df=40, P=0.442457 (n=8-9 mice); *Crf*::ChR2, t=2.369, df=40, *P<0.05 (n=6 mice). Holm-Sidak method for multiple comparisons. **K.** Representative image, scale bar 0.64mm. **I.** Optogenetics-RNAScope experiment. One-Way ANOVA $F_{(3,9)}$=13.53, **P<0.01. Tukey's post-hoc test, *Crf*::ChR2 (stress + stim) vs *Crf*::tdT (resilient), P=0.7872; *Crf*::ChR2 (stress + stim) vs. *Crf*::tdTomato (s), **P<0.01; *Crf*::ChR2 (stress + stim) vs. *Crf*::chR2 (stim), **P<0.01; *Crf*::tdTomato (r) vs. *Crf*::tdTomato (s), *P<0.05; *Crf*::tdTomato (r) vs. *Crf*::ChR2 (stim), *P<0.05; *Crf*::tdTomato (s) vs. *Crf*::ChR2 (stim), P=0.9988. All data represent means ± SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, ns = not significant.